

REGISTRATION DIVISION
PESTICIDES EPA
5761

TRACO SUPER-FINE COPPER

GUARANTEED ANALYSIS:

ACTIVE INGREDIENT	
Copper (in Basic Copper Sulphate)	
Expressed as Metallic	50.0%
INERT INGREDIENTS	50.0%
TOTAL	100.0%

Particle Size - Sub-Micron

CAUTION KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED: If taken internally, induce vomiting by giving mustard and water and tickling the throat. Give large doses of magnesia or lime water, followed with doses of Epsom Salts. Highly toxic to fish. Do not contaminate any body of water by direct application. Clearing of impurities out of nasal cavities and sinuses.

NOTICE TO USER: Traylor Chemical & Supply Co. can accept no responsibility for injury to persons or animals or for loss or damage to crops, soil or other property resulting from failure to follow directions for use or to observe cautions printed on this package.

DIRECTIONS FOR USE: Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations.

CAUTION: DO NOT USE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

TOMATOES, PEPPERS & POTATOES - For the control of Early and Late Blight of potatoes and tomatoes, Septoria Leaf Spot of tomatoes, Frog-eye Spot of peppers, use 2 to 4 lbs. per 100 gallons of spray. Begin spraying when diseases appear. Repeat every 3 to 7 days as needed.

CUCURBITS - (Cucumbers, watermelons, cantaloupes, squash) - For the control of Downy Mildew use 2 to 4 lbs. per 100 gallons of spray every 5 to 10 days as needed. Begin spraying at first sign of disease.

CELERY - For control of Early Blight use 2 to 4 lbs. per 100 gallons of spray every 5 to 14 days as needed. Begin spraying at first sign of disease.

CITRUS - Use 1 1/2 to 2 lbs. per 100 gallons of spray solution in Petal Fall or Post Bloom application for the control of Scab and/or Melanose. For dormant application for the control of Scab use 2 to 3 lbs. per 100 gallons spray.

NOTE: Traco Super Fine Copper is compatible with oil sprays used on citrus. The use of Copper and oil combinations should be avoided after the fruit has attained a size of 3/4 inch in diameter and when the temperature is excessively high.

ACCEPTED
U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF PESTICIDE REGULATION
WASHINGTON, D.C. 20250

NET WEIGHT 50 LBS.

Manufactured For:

EPA Reg. No. 20004-1

Traylor Chemical & Supply Co., Inc.

Orlando
Florida

Royal Center
Indiana

Fresno
California