

RECOMMENDATIONS

Use of this material in a manner other than in accordance with this label may cause serious plant, crop, or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

Unless otherwise specified for specific tree fruits, dosage rates are given as pints of THAT® FLOWABLE SULFUR per 100 gallons water for use in a thorough coverage spray.

APPLE:

Scab: Pre-bloom through Calyx sprays	1½ to 3½ pints
Cover sprays	1 to 2 pints
Powdery Mildew:	
Pre-bloom through Calyx sprays	2½ to 3 pints
Cover sprays	2 pints

PEACH:

Powdery Mildew, Brown Rot, Scab:	
Pink and bloom sprays	1½ to 3½ pints
Petal fall, shuck & cover sprays	1 to 1½ pints

CITRUS: (Grapefruit, Lemons, Lime, Oranges, Tangelos and Tangerines)

Rust Mite	1 to 4 pints
-----------	--------------

Do not apply within 4 weeks of an oil application.

PEANUTS:

Rust: Apply 2 to 4 pints per acre in sufficient water for thorough coverage. Begin application approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks of an oil application.

Consult your State Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your local conditions.

That® FLOWABLE SULFUR


ACTIVE INGREDIENT:

Sulfur	58%
INERT INGREDIENTS:	42%
Total	100%

(Contains 6.0 pounds Sulfur per gallon)

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not apply to any area not specified on label. Do not re-use empty container. Wash thoroughly with water and detergent. Puncture, make unusable. Dispose of in a safe place.



MANUFACTURED BY:
STOLLER CHEMICAL COMPANY, INC.
2211 SOUTH WHITTIER AVENUE
SPRINGFIELD, ILLINOIS 62704

NET CONTENTS 5 GALLONS

CAUTION: Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

During periods of high temperature sulfur may burn foliage and fruit. Do not make sulfur applications at excessively high temperatures. Do not use THAT® FLOWABLE SULFUR with oil or within 4 weeks of an oil application.

DIRECTIONS FOR MIXING BEFORE USING, STIR UNTIL SMOOTH

Pour recommended amount into partially filled spray tank. Keep agitator running during filling and spraying operation. **DO NOT ALLOW MIXTURE TO STAND.** Failure to maintain agitation will cause THAT® FLOWABLE SULFUR to settle and may necessitate manual stirring to redisperse.

The strong adhesive properties of THAT® FLOWABLE SULFUR act as a sticker on the plant, and the sticking characteristic necessitates the flushing of equipment with water after each day's use. Sulfur in any form is corrosive material. **TO REDUCE THE EFFECT, EQUIPMENT SHOULD BE FLUSHED DAILY.**

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use; but neither this warranty or any other warranty of merchantability of fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions not reasonably foreseeable to the seller; and buyer assumes the risk of any such use.

EPA Reg. No. 22555-1

ACCEPTED

MAY 1 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR THE FEDERAL POISON REGISTER
ED UNDER NO. ... SUBJECT
TO ATTACHED COMMENTS