



CARBARYL BAIT

N-Methyl Carbamate Insecticide

For Control of Cutworms, Armyworms, Crickets,
Darkling Ground Beetles, Grasshoppers,
and Sow Bugs.

ACTIVE INGREDIENT:	BY WEIGHT
Carbaryl (1-naphthyl-methyl carbamate)	5.0%
INERT INGREDIENTS:	95.0%
TOTAL	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Below For Other Precautions
NET WEIGHT 50 LBS. (22.68 KG)
EPA REG. NO. 34704-483
EPA EST. NO. _____

IHT

05Y02

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid contact with skin.
Avoid contamination of food, feed, water supplies, streams ponds.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets with requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-228-5635, EXT. 136, OR CALL COLLECT, 612-851-8180, EXT. 136. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Contains an N-methyl carbamate that inhibits cholinesterase.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Subsequent planting of food and feed crops in treated soils must be made with crops on this or other carbaryl labels.

ACCEPTED
MAY 15 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 34704-483

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated area during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is: coveralls, waterproof gloves and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.
Keep pets off treated area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry well-ventilated area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

APPLICATION INSTRUCTIONS

Broadcast application may be made with either ground equipment or aircraft. Apply recommended dosage for control of cutworms, armyworms, crickets, darkling ground beetles, grasshoppers and sow bugs.

For band application, reduce rates proportionately.

TURFGRASS AND ORNAMENTAL PLANTS

SITE	INSECT	POUNDS OF BAIT PER 1000 SQ. FT.	SPECIFIC DIRECTIONS
Turfgrass	Armyworms, Crickets, Cutworms, Darkling ground beetle, Grasshoppers, Mole crickets, Sowbugs	.50 to 1	Treat entire area to obtain uniform coverage, giving particular attention to building perimeters, edges of sidewalks, and along porch and patio foundations. For best results, mow lawn, remove clippings, and thoroughly water area to be treated prior to application.
Ornamental plants	Armyworms, Crickets, Cutworms, Darkling ground beetle, Grasshoppers, Mole crickets, Sowbugs	.50 to 1	Apply uniformly to soil around plants, shrubs, and trees. For best results, thoroughly water area to be treated prior to application.

CROPS

For control of Ants, Centipedes, Crickets, Cutworms, Earwigs, Grasshoppers, Millipedes, Sowbugs, Springtails, and ticks, apply product evenly over the soil surface.

CROP	POUNDS OF BAIT PER ACRE	POUNDS OF BAIT PER 1000 SQ. FT.	PREHARVEST INTERVAL	SPECIFIC DIRECTIONS
Asparagus	40	0.90	1 Day	Repeat applications as necessary up to a total of 3 times prior to harvest or a total of 4 times per year but not more often than once every 7 days.
Beans, Peas, Lentils*	30	0.70	3 Days (Fresh beans or peas) 14 Days (Forage) 21 Days (Dried beans or peas, seed, or hay)	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
*Not registered for use in California				
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard greens, Kohlrabi, Chinese Cabbage	40	0.90	3 Days (Broccoli, Brussel Sprouts, Cabbage, Cauliflower, Kohlrabi) 14 Days (Chinese Cabbage, Collards, Kale, Mustard greens)	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Carrots, Garden Beets (roots), Horseradish, Parsnips, Potatoes, Radishes, Rutabagas, Salsify, Sweet Potatoes, Turnips (roots)	40	0.90	7 Days	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Celery, Dandelion, Endive, Garden Beets (tops), Hanover Salad, Lettuce (head and leaf), Parsley, Spinach, Swiss chard, Turnips (tops)	40	0.90	14 Days	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Corn (Field and Pop)	40	0.90	14 Days (Forage, silage) 48 Days (Grain, Fodder)	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 14 days.

CARBARYL BAIT
EPA REG. NO. 34704-483

3/3

CROP	POUNDS OF BAIT PER ACRE	POUNDS OF BAIT PER 1000 SQ.FT.	PREHARVEST INTERVAL	SPECIFIC DIRECTIONS
Sweet Corn	40	0.90	2 Days (Ears) 14 Days (Forage) 48 Days (Fodder)	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
Cucumber, Melon, Squash	20	0.45	3 Days	Repeat applications as necessary up to a total of 6 times per year but not more often than once every 7 days.
Eggplant, Peppers, Tomatoes	40	0.90	3 Days	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.
Grain Sorghum (not sweet)	40	0.90	14 Days (Grazing, forage) 21 Days (Grain, fodder)	Repeat applications as necessary up to a total of 3 times per year but not more often than once every 7 days.
Strawberries	40	0.90	7 Days	Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days. Carbaryl may injure Early Dawn and Sunrise varieties of strawberries.
Sugar Beets	30	0.70	28 Days (Roots, forage)	Repeat applications as necessary up to a total of 2 times per year but not more often than once every 14 days.
Wheat	30	0.70	21 Days (Grain, straw) 7 Days (Forage)	Repeat applications as necessary up to a total of 2 times per year but not more often than once every 14 days.

NOTICE

IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE, THE MANUFACTURER OR SELLER. IN NO CASE SHALL PLATTE, THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE, THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.