

JAN 23 1996

5493601

464  
18

Jeannie K. Smith  
Platte Chemical Co.  
P.O. Box 667  
Greeley, CO 80632-0667

Dear Ms. Smith:

Subject: Dibrom 8EC Naled Insecticide  
EPA Registration No. 34704-546  
Amendment dated September 1, 1995

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the following changes to your label:

(a) Change the PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS to read as follows: "DANGER. CORROSIVE. Causes irreversible eye and skin damage. May be fatal if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe vapor or spray mist."

(b) Change FIRST AID to read as follows:

*If in eyes:* Immediately hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention immediately.

*If inhaled:* Remove victim to fresh air. If not breathing, give artificial respiration. Get medical attention immediately.

*If swallowed:* Drink 1 or 2 glasses of water (or milk) and induce vomiting by touching the back of the throat with finger. If possible contact a physician, Poison Control Center or emergency center before inducing vomiting. Do not induce vomiting or give anything by mouth to an unconscious person.

*If on skin:* Wash with plenty of soap and water. Get medical attention immediately.

NOTE TO PHYSICIAN: Naled is a cholinesterase inhibitor. Measurement of blood cholinesterase activity may be useful in monitoring exposure. If signs of cholinesterase inhibition appear, atropine is antidotal. Mucosal damage may contraindicate the use of gastric lavage."

B Edwards

A/34701/546

1/23/96

245

(c) To the DIRECTIONS FOR USE for BEANS/GREEN BEANS, add the following sentences: "Make only one application per 7 day period. Do not apply within 1 day of harvest. Do not feed treated vines."

(d) To the DIRECTIONS FOR USE for COTTON, add the following sentences: "Do not apply after first bolls open. Do not graze livestock in treated fields."

(e) To the DIRECTIONS FOR USE for GRAPES, add the following sentence: "Do not apply within 4 days of harvest."

(f) To the DIRECTIONS FOR USE for WALNUTS add the following sentence: "Do not graze livestock in treated orchards."

2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records. If you have any questions about this letter, please contact Beth Edwards at (703) 305-5400 or (703) 305-6600.

Sincerely yours,

Robert A. Forrest  
Product Manager (14)  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

Enclosure

CONCURRENCES

SYMBOL								
SURNAME								
DATE								



# DIBROM<sup>®</sup> 8 EC NALED INSECTICIDE

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

JAN 23 1996

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, this pesticide  
registered under EPA Reg. No.

34704-546

### For Agricultural or Commercial Use Only

ACTIVE INGREDIENT:	BY WT.
Naled (1,2-dibromo-2,2-dichloroethyl dimethyl phosphate) .....	58.0%
INERT INGREDIENTS: .....	42.0%
TOTAL 100.0%	

Contains 8 pounds technical naled per gallon equivalent to 7.2 pounds naled.

\*This product contains Xylene Range Aromatic Solvents.

### USER SAFETY RECOMMENDATIONS

**Users should:**  
 Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.  
 Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.  
 Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## KEEP OUT OF REACH OF CHILDREN DANGER—PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See Below For Additional Precautionary Statements.

EPA REG. NO. 34704-546 EPA EST. NO. 39184-WA-1

NET CONTENTS 1 GAL. (3.78 L)

4778 DIBROM is a Registered T.M. of Chevron Chemical Co. EXP07Y95

### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Store in cool, dry place.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system. Do not spray on Nectarines, Ornamental Cherries, Liquidambar or Chrysanthemums as injury may occur. Do not store diluted spray. Use new dilution for each use. Wash equipment thoroughly after use. Keep children and pets off treated areas.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER:** May be harmful or fatal if swallowed. Concentrate causes eye and skin damage. Do not get on skin, eyes or clothing. Use waterproof gloves and face shield or goggles when handling concentrate. Avoid breathing spray mist.

#### FIRST AID

In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water. For eyes, get medical attention.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN:** Atropine is antidotal. Mucosal damage may contraindicate the use of gastric lavage.

#### Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category F on the EPA chemical resistance category selection chart.

**Applicators and other handlers must wear:** coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate, butyl rubber > 14 mils, nitrile rubber > 14 mils and viton > 14 mils, chemical-resistant footwear plus socks, protective eyewear, chemical-resistant headgear for overhead exposure, chemical-resistant apron when cleaning equipment, mixing, or loading, and a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approved number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approved number prefix TC-24G).

**Exception:** When applying to hot pipes in greenhouses or entering vapor-treated greenhouses during application or before ventilation is complete, the protective eyewear must be goggles, unless a full-face respirator is worn.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

**Entry Restrictions:**  
 Outdoors and Non-Vapor treatments in greenhouses: Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

**Vapor Treatment in Greenhouses:** Entry by any person -- other than a correctly trained and equipped handler who is performing a WPS-defined handling task -- is PROHIBITED in the entire enclosed area from the start of application until one of the WPS ventilation criteria has been met. After ventilation is completed, do not enter or allow worker entry into the entire enclosed area during the restricted-entry interval (REI) of 24 hours, except as allowed by the WPS.

1485

**CLEAN CROP  
DIBROM® 8 EC**  
EPA REG. NO. 34704-546

**PPE for Entry During the Restricted Period:**

PPE for early entry to treated areas (1) outdoors, (2) in non-vapor treated greenhouses, or (3) in vapor-treated greenhouses after ventilation is complete - that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, such as barrier laminate, butyl rubber > 14 mils, nitrile rubber > 14 mils and vinyl > 14 mils, chemical-resistant footwear plus socks, protective eyewear, and chemical-resistant headgear for overhead exposure.

PPE for entry by handlers into vapor-treated greenhouses before WPS Ventilation Criteria have been met is listed in the "Hazards to Humans and Domestic Animals" section of this labeling.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of the operating areas or vicinity where there may be danger of drift. Do not enter treated areas prior to 24 hours after application without protective clothing (Does not apply to uses for adult mosquito control, or fly control, or rangeland pest control).

**STORAGE AND DISPOSAL**

**PESTICIDE STORAGE:** Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: dike surrounding area and absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers. Glass: Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved State and local procedures. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**APPLICATION**

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

**GROUND APPLICATION:** Vegetable and Field Crops—30 to 200 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—60 to 600 gals. Citrus—400 to 3000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre.)

**Where Recommended for AIR APPLICATION:** Field Crops—3 to 10 gals. A short residual contact insecticide for kill of insect and mite adults and larvae present at the time of application.

Begin application at first sign of insects. Repeat as necessary. Do not apply within 4 days of harvest unless otherwise noted. Do not use with highly alkaline materials such as lime or bordeaux.

**MIXING DIRECTIONS:** Add to spray tank when 1/4 to 1/2 full with agitator running and maintain agitation until spraying is completed.

**ALFALFA:** Aphids, Loopers, Leafhoppers, Lygus Bugs, Alfalfa Webworms, Armyworms—1 pt. per acre. **ALFALFA (Seed):** Lygus, Mite Suppression (1 application). Mites—(2 applications at 7 day intervals). Thrips—1/2 pt. per acre. Combine with recommended chemicals for residual control. By Air—Use at least 10 gals. water per acre.

**BEANS (Dry or Field), SOYBEANS, LIMA BEANS, and PEAS for Processing:** Alfalfa Looper, Aphids, Spider Mites, Lygus, Leaf Miner; **GREEN BEANS:** Aphids, Spider Mites, Loopers, Nitidulids, Leafhoppers—1 pt. per acre in water to cover 100 gals. per acre is preferable. **Large Loopers, Stink Bugs—** 1 1/2 pts. per acre in water to cover.

**BROCCOLI, CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS, KALE, COLLARDS:** Imported Cabbageworm, Diamond-Back Caterpillars, Aphids—1 pt. per acre. **Looper—** 2 pts. per acre. May be used 1 day before harvest, except do not apply to kale and collards within 4 days of harvest.

**CELERY:** Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 3/4 grown. On large plants use 1 1/2 pts. in 100 to 150 gals. water per acre.

**COTTON:** Lygus—1/2 to 1 pt. per acre by air or ground. Leaf Perforator, Fleahoppers, Armyworms, Bollworms, Loopers and for Suppression of Mites—1 pt. per acre by air or ground. Apply at 7 to 10 days intervals while pests persist. **FOR HEAVY POPULATION OF LISTED INSECTS—** Use 1/2 pt. per acre by air or ground in control programs with other insecticides registered for use on cotton.

**CUCUMBERS, SUMMER SQUASH:** Cucumber Beetles, Aphids, Spider Mites, Loopers, Thrips, Armyworms—1 pt. per acre in water to cover. Pickleworms, Leaf Miners—1 1/2 pts. per acre in water to cover. **By Air—** Use minimum of 10 gals. water per acre. May be used up to harvest.

**EGGPLANT, PEPPERS:** Aphids, Spider Mites, Blister Beetles, Flea Beetles, Leaf Miners, Whitefly—1 pt. per acre. May be used up to 1 day before harvest.

**GRAPES:** Drosophila, Leafhoppers—1/2 to 3/4 pt. per 100 gals. water (100 to 130 gals. diluted spray per acre depending on size of vine). Use only 1/2 pt. per 100 gals. water on Muscat varieties. Do not use on Italia varieties as fruit injury may occur.

**HOPS:** Aphids, Spider Mites, Corn Earworms, Hop Loopers, Cutworms—1 pt. per acre.

**SPINACH, CHARD, TURNIP GREENS:** Aphids, Loopers, Armyworms, Salt Marsh Caterpillars, Leaf Miners—Use ground application only with fine mist spray—1 pt. in 50 to 75 gals. water per acre on plants less than 3/4 grown. On larger plants use 1 1/2 pts. in 100 to 150 gals. water per acre. May be used 1 day before harvest.

**MELONS (Cantaloupes, Honeydew, Muskmelons, Watermelons), PUMPKINS, WINTER SQUASH:** Loopers, Armyworms, Leafhoppers, Spider Mites, Leaf Miners—1 to 2 pts. per acre (200 gals. diluted spray per acre). May be used 1 day before harvest. Do not apply when temperature is over 90°F.

**ORANGES, LEMONS, GRAPEFRUIT, TANGERINES:** Aphids, Soft Brown Scale Crawlers, Mites, Citrus Cutworms, Leaf Roller (Tortrix)—1 pt. to 100 gals. water. Thrips (Bloom)—1/2 pt. per 100 gals. water. Use up to 7 days of harvest. (Calif. and Ariz.—Do not apply within 1 day of harvest.)

**PEACHES (Western States Only):** Lygus, Suppression of Mites, Peach Twig Borer, Oriental Fruit Moth (Calif. Only)—3/4 pt. to 100 gals. water. To avoid fruit spotting, do not use within 30 days of harvest. Do not apply when temperature is over 90°F.

**STRAWBERRIES:** Leaf Roller, Spider Mites, Aphids, Omnivorous Leaf Tier, Spittlebugs (Northwest)—1 pt. per acre. May be used 1 day before harvest.

**SUGAR BEETS:** Spider Mites, Leafhoppers—1 pt. per acre by air or ground. **By Air—** Use in 1 to 5 gals. water per acre. Do not apply within 5 days of harvest.

**TOBACCO:** Hornworm—1 pt. to 100 gals. water, using up to 200 gals. diluted spray per acre. Flea Beetles—1 pt. per acre in water to cover. Apply at first signs of insects. Repeat as necessary.

**TOMATOES: (Field):** Fruit Flies (Drosophila)—1 pt. in no less than 40 gals. water per acre with ground sprayer, 7 to 10 gals. by air. Make first application 5 to 7 days before first picking and continue at 5 to 7 day intervals until the day before harvest. (Field and Greenhouse): Tomato Fruit Worm, Hornworms, Leaf Miners, Flea Beetles, Mites—1 pt. in 100 gals. water. Spray plants thoroughly. **For Field—** Use up to 200 gals. diluted spray per acre. Do not apply when temperature is over 90°F. May be used 1 day before harvest.

**WALNUTS:** Aphids—1/2 to 1 pt. in 100 gals. water. Use up to 10 days of harvest.

**FOREST AND SHADE TREES: CONIFERS (Arbovitae, Douglas Fir, Fir, Hemlock, Juniper, Pine, Spruce):** Suppression of Two-Spot and Red Spider Mites, Aphids, Zimmerman Moths, Juniper 1/2 b worms, Tussock Moths, Spruce Budworms, Adult Sierra Fir and Western Hemlock Bark Borers.

5 of 5

**CLEAN CROP  
DIBROM® 8 EC  
EPA REG. NO. 34704-546**

**BROADLEAF TREES** (Ash, Birch, Black Walnut, Box Elder, Dogwood, Elm, Evergreen Pear, Flowering Plum and Quince, Locust, Magnolia, Maple, Oak, Sycamore, Willow): Suppression of Two-Spot and Red Spider Mites, Aphids, Leaf Miners, Thrips, Elm Leaf Beetles, California Oakworms (Calif. Only), Spring and Fall Cankerworms, Fall Webworms, Oak Webworm and Tussock Moths — 1 pt. in 100 gals. water. Use when infestation is evident, repeat as necessary.

**GREENHOUSE, VAPOR TREATMENT OF ROSES; OTHER ORNAMENTAL PLANTS, TOMATOES, CUCUMBERS:** Spider Mites, Adult Whiteflies, Aphids, Leafrollers, Mealybugs — 1 fluid ounce per 10,000 cu. ft. when plants are dry. Apply undiluted on heat pipes (or in open pans on hot plates). Apply continuously or to intermittent sections of pipe on each side and on one or more of the pipes through center of each range depending upon width. Application to cold pipes should be followed by immediately heating pipes to 160°F. In steam heated houses, close vents for at least 1 hour after pipes are hot. With hot water systems, close house for at least 3 hours. Houses may remain closed overnight following treatment. When application is made to hot pipes, wear goggles and approved respirator. Ventilate house before workers reenter. Wear respirator when entering treated houses to ventilate. Protect hives of bees in cucumber and other houses by covering with plastic or rubber sheet during treatment and until house is ventilated. Do not apply to tomatoes or cucumbers closer than 1 day before harvest. Spider Mites (resistant and non-resistant) — Make 3 to 4 treatments at 3 or 4 day intervals. Repeat as necessary. Adult Whiteflies — Make 2 applications at 7 day intervals. Repeat as necessary. Aphids — Apply when aphids appear. Repeat as necessary. **WARNING** — Avoid overtreatment and direct application to plants as injury may result. White Butterfly and Golden Rapture Roses, Pink Champagne Chrysanthemums, Green Wandering Jew, Poinsettias and Dutchman's Pipe may be injured by DIBROM vapor.

**FRUIT FLIES (Drosophila sp.)** in an around Food Processing plants, Loading Docks, Cull Piles and Refuse Areas, Cider Mills — Use 2 ounces in 2½ gals. water (5 pts. to 100 gals.) as a coarse spray to walls, floors, doorways, windows, refuse and cull piles where insects congregate. Apply every 5 to 7 days as necessary. Foods should be removed or covered during treatment. All food processing surfaces should be covered before using. When using the product in these areas, apply only when the facility is not in operation.

**ADULT MOSQUITO, GNAT, STABLEFLY (Dog Flies), and HOUSEFLY CONTROL — RESIDENTIAL AREAS, CROPS LISTED ON LABEL, TIDAL MARSHES, SWAMPS, WOODLAND, LIVESTOCK PASTURES, FOOD LOTS, and PASTURES including DAIRY CATTLE:** Consult your State Fish and Game Agency before applying this product. It is not necessary to avoid farm buildings and dairy barns. Make applications during peak of infestation and repeat as necessary. See crop recommendations for use limitations near harvest. This rate of application will kill shrimp. Do not apply to tidal or marsh waters. Treat shrubbery and vegetation where mosquitoes may rest. Shrubby and vegetation around stagnant pools, marshy areas ponds and shore lines may be treated. Application of this product to any body of water is prohibited. **AIRCRAFT, East** — Apply 0.8 to 4 fluid ounces per acre diluted with water. Apply 2 to 8 quarts of diluted spray per acre. **MIST OR COLD FOG:** Use 3 to 5 quarts per 100 gals. water. Calibrate the equipment (rate of travel and output) to apply 0.1 to 0.25 pounds technical Dielder per acre.

**CORRALS, ADJACENT PASTURES, HOLDING PENS (Dairy and Beef Cattle, Hogs, Sheep, Horses):** Houseflies, Mosquitoes — **Airplane Application** — Use 1.6 to 4 fluid ounces per acre. Dilute 1 to 2½ pts. in 10 gals. water. Apply 1 gal. diluted spray per acre. Apply over areas with animals present.

**CLOVER MITES, ROACHES, EARWIGS and LEAFHOPPERS** Outside Dwellings and in Lawns: Use 1 fluid ounce in 3 gals. water. Apply to turf and to soil surfaces around flowers, shrubs and trees for general pest cleanup.

**NOTICE**

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER EXCEPT AS EXPRESSLY PROVIDED HEREIN. PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND PLATTE'S TOTAL LIABILITY, SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

FORMULATED FOR  
**PLATTE CHEMICAL CO.**

150 SO. MAIN STREET      REMONT, NEBRASKA 68025-5637