



LINDANE

EMULSIFIABLE CONCENTRATE

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

As A Livestock Dip

For the control of Sarcoptic and Psoroptic Mange Mites, Ticks, Hornflies and Lice on Animals except Dairy Animals, Chiggers and Ticks on Lawns. To be diluted with water.

ACTIVE INGREDIENTS:	
Lindane (Gamma isomer of benzene hexachloride)	11.14%
INERT INGREDIENTS:*	88.86%
TOTAL	100.00%

*This product contains xylene range aromatics.

KEEP OUT OF REACH OF CHILDREN CAUTION

See Below For Additional Precautionary Statements.

EPA REG. NO. 34704-591 EPA EST. NO. _____

NET CONTENTS _____ GALLONS

24859

EXP-1Y

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

HARMFUL IF SWALLOWED. May be absorbed through the skin. Avoid breathing spray mist and skin contact. In case of contact, wash immediately with soap and water. In case of contact with eyes, flush with water and get medical attention.

Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed or foodstuffs.

Applicators must wear the following protective clothing during the application process: long-sleeved shirt, long pants, waterproof gloves, full foot covering, and a head covering.

Mixers/loaders must wear goggles or face shield, waterproof apron, and waterproof gloves.

Do not allow children to handle or apply this product.

Children and pets should not be allowed in treated areas until sprayed surfaces are dry.

Do not treat sows within 3 weeks of farrowing nor for at least 3 weeks thereafter.

Do not use on dairy cattle or dairy goats.

Do not use on poultry.

Do not use a solution stronger than 1 pint mixed in 50 gallons of water on calves and other young livestock.

Do not use on household pets or humans.

Do not dip livestock within 60 days of slaughter.

Do not treat any animal under 3 months of age.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. If possible vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAN: Solvent presents aspiration hazard. Gastric lavage is indicated if material was taken internally.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Birds and other wildlife feeding in treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting vegetation blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, STORE OR SPILL NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not use or store near heat or open flame. Do not store below 32°F. Keep product in original container, tightly closed.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

FOLLOW DIRECTIONS CLOSELY, MEASURE ACCURATELY, AND STIR EMULSION THOROUGHLY JUST BEFORE USING.

DO NOT TREAT ANIMALS IN COLD, STORMY WEATHER. DO NOT TREAT OVERHEATED OR SICK ANIMALS. DO NOT TREAT ANIMALS YOUNGER THAN 3 MONTHS OF AGE. DO NOT USE ON DAIRY CATTLE. DO NOT DIP LIVESTOCK WITHIN 60 DAYS OF SLAUGHTER. DO NOT APPLY BY AIRCRAFT.

LIVESTOCK DIP ONLY:

HORNFLIES ON BEEF CATTLE, HORSES AND HOGS: Treat animals thoroughly with Lindane Solution prepared as follows: 1 pint to 45 gallons water. Repeat every 3 to 4 days or as often as necessary.

LICE AND TICKS ON BEEF CATTLE, SHEEP AND GOATS: Treat animals with 0.03% Lindane prepared as follows: 1 pint to 45 gallons water. Skin and hair of sheep and goats must be thoroughly wet. Repeat as necessary.

LICE ON HOGS: Treat hogs with a solution not to exceed 0.06% Lindane prepared as follows: 1 quart to 45 gallons water. Repeat as necessary.

SARCOPTIC AND PSOROPTIC MANGE ON BEEF CATTLE AND HOGS: Use 0.06% Lindane prepared as follows: 1 quart to 45 gallons water. Wet animals thoroughly including inside of legs. Pens and shelters should be sprayed to prevent re-infection. Do not use on Dairy Cattle.

Used dip solutions must be disposed of in accordance with the Resource Conservation and Recovery Act (RCRA). If the applicator generates more than 1000 kg (2200 lbs.) of used dip solution per month or more than 1000 kg (2200 lbs.) used dip solution in combination with other hazardous waste, the material must be treated as a hazardous waste subject to subpart C of RCRA. Any user who wishes to treat, store or dispose of hazardous waste must obtain a permit to serve as a hazardous waste facility pursuant to RCRA.

FLIES ON MANURE PILES, REFUSE HEAPS: Spray with 0.125% Lindane prepared as follows: Mix 1 pint LINDANE EMULSIFIABLE CONCENTRATE to 10 gallons water. Wet thoroughly every 3 to 4 weeks or as manure is added. Do not apply treated manure to land to be planted to

(continued)

282

LINDANE EMULSIFIABLE CONCENTRATE

EPA REG. NO. 34704-591

(continued)

root crops.

CHIGGERS, TICKS ON LAWNS: Spray thoroughly 0.06% Lindane prepared as follows: Mix 1 pint LINDANE EMULSIFIABLE CONCENTRATE to 20 gallons water. One-half gallon Lindane will be sufficient to cover approximately one acre (1/2 lb. Lindane per acre).

NOTICE OF WARRANTY

Manufacturer and Seller make no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

550

FORMULATED FOR
PLATTE CHEMICAL CO.
180 SO. MAIN STREET FREMONT, NEBRASKA 68025