



34704-883

08/08/2005

1/31

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	<p>EPA Reg. Number: 34704-883</p>	<p>Date of Issuance: AUG - 8 2005</p>
	<p>Term of Issuance: Conditional</p>	
	<p>Name of Pesticide Product: Mirage Plus</p>	
<p>Name and Address of Registrant (include ZIP Code):</p> <p>Loveland Products, Inc. P. O. Box 1286 Greeley, CO <del>34704</del>-1286 80632</p>		
<p><b>Note:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> <li>1. Submit/cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.</li> <li>2. Make the labeling changes listed below before you release the product for shipment. <ol style="list-style-type: none"> <li>a. Add the phrase "EPA Registration No. 34704-883."</li> <li>b. On page 2, in the Non Agricultural Use Requirements box, delete the phrase "to prevent transfer of this product onto desirable vegetation" from the last sentence.</li> <li>c. On page 2, under Storage and Disposal add the subheadings "Pesticide Storage", "Pesticide Disposal" and "Container Disposal"</li> <li>d. On page 2, under General Information, delete the second paragraph under "No Soil Activity", and the paragraphs entitled "Volatility" and "Toxicology Testing".</li> <li>e. On page 2, under General Information, Annual Maximum Use Rates, revise the last two sentences to read "The maximum use rates stated throughout Mirage Plus's labeling apply to Mirage Plus</li> </ol> </li> </ol>		
<p>Signature of Approving Official</p> 	<p>Date: 8-8-05</p>	

combined with the use of all other herbicides containing **glyphosate** or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other **glyphosate** or sulfosate containing products does not exceed stated maximum use rate.”

f. Move the last two sections of the first column on page 4 “Aerial Equipment” and “Ensure uniform application” to page 3, immediately under the chart for “Aerial Application In California” and before the section entitled “Aerial Applications in California–Industrial Turf and Ornamental Uses”.

g. On page 8, delete the section entitled “For New Cotton Lines With the Mirage Plus Tolerant Gene In-Crop Applications” and move to page 17 under “Cotton with the Mirage Plus Tolerant Gene”.

3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of labeling is enclosed for your records.

Please note that due to a recent ruling by EPA`s Office of General Council (OGC), in situations with 100% repacks, the formulators exemption may be accepted for the basic formulation (CSF). All alternate formulations (CSFs) must comply with Data Compensation.



# MIRAGE PLUS

**AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, (EXCEPT AS SPECIFIED FOR INDIVIDUAL GLYPHOSATE READY CROPS) DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

**ACTIVE INGREDIENT:**  
 \*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt ..... 41.0%

**OTHER INGREDIENTS:** ..... 59.0%

**TOTAL** ..... **100.0%**

\*Contains 480 grams per litre or 4 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

No license granted under any non-U.S. patent(s).

THIS IS AN END-USE PRODUCT. LOVELAND PRODUCTS, INC. DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

## KEEP OUT OF REACH OF CHILDREN WARNING—AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA REG. NO 34704-  
EPA EST. NO. \_\_\_\_\_  
NET CONTENTS \_\_\_\_\_  
IHT \_\_\_\_\_ 03/05

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. HARMFUL IF SWALLOWED OR INHALED.**  
Do not get in eyes or on clothing.  
Avoid breathing vapor or spray mist.

**Personal Protective Equipment (PPE)**  
Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**  
Users should:  
Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.  
Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ACCEPTED**  
with COMMENTS  
In EPA Letter Dated  
**AUG - 8 2005**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
**34704-883**

### FIRST AID

<b>FIRST AID:</b> Call a poison control center or doctor for treatment advice.	
<b>If in eyes:</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses if present after the first 5 minutes then continue rinsing eye.
<b>If inhaled:</b>	• Remove individual to fresh air. • If not breathing, give artificial respiration, preferably mouth-to-mouth. • Get medical attention.
<b>If swallowed:</b>	• This product will cause gastrointestinal tract irritation. • Immediately dilute by swallowing water or milk. • Get medical attention. • Never give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-800-301-7976.**

**Domestic Animals:** This product is considered to be relatively nontoxic to dogs and other domestic animals: however, ingestion of this product or large amounts of freshly sprayed vegetations may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters.

### PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

**AGRICULTURAL USE REQUIREMENTS**  
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.  
Do not enter or allow worker entry to treated areas during the restricted entry interval (REI) of 12 hours.  
PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves (EPA Chemical Resistance Category A) 8 mils in thickness or greater composed of materials such as butyl rubber, natural rubber, neoprene rubber, or nitrile rubber, shoes plus socks and protective eyewear.

# MIRAGE PLUS EPA REG. NO. 34704-

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are Not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

## STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Keep container closed to prevent spills and contamination. Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state, or local procedures. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned, or destroyed.

**(For Refillable Portable Containers)**  
Do not reuse this container except for refill in accordance with a valid Loveland Products, Inc. Repackaging or Toll Repackaging Agreement. If not refilled or returned to the authorized repackaging facility, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**(For Bulk Containers)**  
Triple rinse emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

**(For Plastic 1-Way Containers & Bottles)**  
Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**(For Drums)**  
Do not reuse container. Return container per the Loveland Products, Inc. container return program. If not returnable, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning, if burned, stay out of smoke.

**Container Disposal (Bulk and Minibulk)**  
**Instructions for User**  
When the container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase, or to an alternate location designated by the registrant at the time of purchase of this product. If not returned to the point of purchase or to a designated location, triple rinse or pressure rinse the empty container and offer for recycling if available.

**Instructions for Users and Refillers**  
The container must only be refilled with this pesticide product. Do not Reuse the Container for Any other Purpose. Do not transport if this container is damaged or leaking if the container is damaged, leaking, or obsolete, or to obtain information about recycling refillable containers, contact Loveland Products, Inc. at 1-970-356-4400. Cleaning is not necessary prior to refilling with the same product. Clean container before final disposal. Disposal of this container must be in compliance with state and local regulations.

**Instructions for Refillers**  
Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. If the container cannot be refilled, triple rinse or pressure rinse the empty container and offer for recycling if available.

## General Information (How This Product Works)

**Product Description:** Mirage Plus is a post-emergent, systemic herbicide with no soil residual activity. It is generally non-selective and gives broad-spectrum control of many annual weeds, perennial weeds, woody brush and trees. It is formulated as a water-soluble liquid. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water or other carriers according to label instructions.

**Time to Symptoms:** Mirage Plus move through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow activity of Mirage Plus and delay development of visual symptoms. Visible effects are a gradual wilting and yellowing of the plant, which advances to complete browning of aboveground growth and deterioration of underground plant parts.

**Stage of Weeds:** Annual weeds are easiest to control when they are small. Best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity. Refer to the "Annual Weeds, Perennial Weeds And Woody Brush Rate Tables" for recommendations for specific weeds.

Always use the higher rate of Mirage Plus per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

**Cultural Considerations:** Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed, or cut, and have not been allowed to re-grow to the recommended stage for treatment.

**Rainfastness:** Heavy rainfall soon after application may wash Mirage Plus off of the foliage and a repeat application may be required for adequate control.

**Spray Coverage:** For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

**Mode of Action:** The active ingredient in Mirage Plus inhibits an enzyme found only in plants and microorganisms that is essential to formation of specific amino acids.

**No Soil Activity:** Weeds must be emerged at the time of application to be controlled by Mirage Plus. Weeds germinating from seed after application will not be controlled. Unemerged plants arising from unattached underground rhizomes or rootstocks of perennials will not be affected by the herbicide and will continue to grow.

When this product comes in contact with soil, it is bound to soil particles. Under recommended use situations, once this product is bound to soil particles, it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treated area or if the soil is transported off-site. The strong affinity of this product to soil particles prevents this product from leaching out of the soil profile and entering ground water.

**Biological Degradation:** Degradation of Mirage Plus is primarily a biological process carried out by soil microbes.

**Volatility:** This product is non-volatile. Therefore, it cannot move as a vapor after application to affect nearby vegetation.

**Toxicology Testing:** Exposure to workers and other applicators generally is expected to pose minimal risks based on results of short-term toxicity studies. Glyphosate has been thoroughly tested and determined not to cause cancer or other adverse long-term health effects.

**Tank Mixing:** Mirage Plus does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of Mirage Plus with herbicides or other materials that are not expressly recommended in this labeling. Mixing Mirage Plus with herbicides or other materials not recommended on this label may result in reduced performance.

When this label recommends a tank mixture with a generic active ingredient such as diuron, atrazine, 2,4-D, or dicamba, the user is responsible for ensuring that the mixture product's label allows the specific application.

**Annual Maximum Use Rate:** Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 8 quarts of Mirage Plus per acre per year. For applications in non-crop sites or in tree, vine, or shrub crops, the combined total of all treatments must not exceed 10.6 quarts of Mirage Plus per acre per year. The maximum use rates stated throughout Mirage Plus's labeling apply to Mirage Plus combined with the use of all other herbicides containing Mirage Plus or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other Mirage Plus or sulfosate containing products does not exceed stated maximum use rate.

**NOTE:** Use of Mirage Plus in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

## MIXING

Clean sprayer parts immediately after using Mirage Plus by thoroughly flushing with water.

# MIRAGE PLUS

## EPA REG. NO. 34704-

5/31

**NOTE:** REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS VISIBLY MUDDY WATER OR WATER FROM PONDS AND DITCHES THAT IS NOT CLEAR.

### Mixing with Water

Mirage Plus mixes readily with water. Mix spray solutions of Mirage Plus as follows: Fill the mixing or spray tank with the required amount of water. Add the recommended amount of Mirage Plus near the end of the filling process and mix well. Use caution to avoid siphoning back into the carrier source. Use approved anti-back-siphoning devices where required by state or local regulations. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

### Tank Mixing Procedure

Mix labeled tank mixtures of Mirage Plus with water as follows:

1. Place a 20- to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the spray tank one-half full with water and start agitation.
3. If ammonium sulfate is used, add it slowly through the screen into the tank. Continue agitation. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding other products.
4. If a wettable powder is used, make a slurry with the water carrier, and add it slowly through the screen into the tank. Continue agitation.
5. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture slowly through the screen into the tank. Continue agitation.
6. If an emulsifiable concentrate formulation is used, pre-mix one part emulsifiable concentrate with two parts water. Add diluted mixture slowly through screen into tank. Continue agitation.
7. Continue filling the spray tank with water and add the required amount of Mirage Plus near the end of the filling process.
8. If a non-ionic surfactant is used, add it to the spray tank before completing the filling process.
9. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water-soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to re-suspend the mixture before spraying is resumed.

Keep by-pass line on or near the bottom of the tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh.

Always predetermine the compatibility of labeled tank mixtures of Mirage Plus with water carrier by mixing small proportional quantities in advance.

Refer to the "Tank Mixing" section of "General Information" for additional precautions.

### Mixing for Hand-Held Sprayers

Prepare the desired volume of spray solution by mixing the amount of Mirage Plus in water as shown in the following table:

Desired Volume	Spray Solution					
	Amount of Mirage Plus					
	0.5%	1%	1.5%	2%	5%	10%
1 gal	0.7 oz	1.3 oz	2 oz	2.7 oz	6.5 oz	3 oz
25 gal	1 pt	1 qt	1.5 qt	2 qt	5 qt	10 qt
100 gal	2 qt	1 gal	1.5 gal	2 gal	5 gal	10 gal

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of Mirage Plus be mixed with water in a larger container. Fill sprayer with the mixed solution.

### Surfactants

Nonionic surfactants (NIS) or wetting agents that have at least 70 percent active ingredient and are labeled for use with herbicides may be added to the spray solution. Do not reduce rates of this herbicide when adding surfactants. Read and carefully observe cautionary statements and other information appearing on the additives label.

### Ammonium Sulfate

The addition of 1 to 2 percent dry ammonium sulfate by weight or 8.5 to 17 pounds per 100 gallons of water may increase the performance of Mirage Plus, particularly under hard water conditions, drought conditions or when tank mixed with certain residual herbicides, on annual and perennial weeds. The equivalent rate of ammonium sulfate in a liquid formulation may also be used. Ensure that dry ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactants. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

**NOTE:** When using ammonium sulfate, apply Mirage Plus at rates recommended in this label. Lower rates will result in reduced performance. The use of ammonium sulfate as an additive does not preclude the need for additional surfactant.

### Colorants or Dyes

Agriculturally approved colorants or marking dyes may be added to Mirage Plus. Colorants or dyes used in spray solutions of Mirage Plus may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's recommendations.

### Drift Control Additives

Drift control additives may be used with all equipment types, except wiper applicators, sponge bars and Controlled Droplet Applicator (CDA) equipment. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label. The use of drift control additives can affect spray coverage which may result in reduce performance.

## APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply Mirage Plus through any type of irrigation system.

Mirage Plus may be applied with the following application equipment:

**Aerial** - Fixed Wing and Helicopter

**Ground Broadcast Spray** - Boom or boomless systems, pull-type sprayer, floaters, pick-up sprayers, spray coupes and other ground broadcast equipment.

**Hand-Held or High-Volume Spray Equipment** - Knapsack and backpack sprayers, pump-up pressure sprayers, handguns, handwands, mistblowers\*, lances and other hand-held and motorized spray equipment used to direct the spray onto weed foliage.

\*Mirage Plus is not registered in California or Arizona for use in mistblowers.

**Selective Equipment** - Recirculating sprayers, shielded and hooded sprayers, wiper applicators and sponge bars.

**Injection Systems** - Aerial or ground injection sprayers.

**Controlled Droplet Applicator (CDA)** - Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

## SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING MIRAGE PLUS TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of Mirage Plus can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

## FOR AERIAL APPLICATION IN CALIFORNIA ONLY

### Use Directions

Aerial applications of Mirage Plus are allowed in the following situations:

1. In fallow and reduced tillage systems prior to the emergence or transplanting of labeled crops.
2. In alfalfa and pasture renovation applications.
3. Over-the-top applications in glyphosate tolerant corn and cotton. Refer to the booklet and/or supplemental labels for Mirage Plus for specific application instructions for over-the-top applications in these crops.
4. Preharvest in alfalfa, corn, cotton, wheat, glyphosate tolerant corn and glyphosate tolerant cotton. Refer to the booklet and/or supplemental labels of Mirage Plus for specific preharvest application instructions for Mirage Plus for each individual crop.

Do not plant subsequent crops other than those listed in the label booklet for 30 days following application.

When applied as recommended under the conditions described, Mirage Plus controls annual and perennial weeds listed in the label booklet.

When tank mixing Mirage Plus with 2,4-D for aerial applications, only 2,4-D amine formulations may be used. This tank mixture may be used for fallow and reduced tillage systems and alfalfa and pasture renovation applications only.

DO NOT EXCEED THE FOLLOWING MAXIMUM RATES WHEN MAKING APPLICATIONS BY AIR:

1 quart per acre	2 quarts per acre
	Alfalfa
Corn	
Mirage Plus tolerant corn	
	Cotton
	Mirage Plus tolerant cotton
	Fallow
	Reduced Tillage Systems
	Pastures
Wheat	

## AERIAL APPLICATIONS IN CALIFORNIA - INDUSTRIAL, TURF, AND ORNAMENTAL USES

### Use Directions

Aerial applications of this product are allowed in the following situations:

1. Prior to the emergence or transplanting of labeled crops
2. Aid to burning for establishment and maintenance of fuel breaks
3. Establishing fire perimeters and black lines

**MIRAGE PLUS**  
**EPA REG. NO. 34704-**

4. Aid to prescribed burning
5. Along fire roads
6. Range conversion
7. Habitat restoration and management
8. Wildlife food plots

Apply 1 to 5 quarts of this product in 5 to 15 gallons of water per acre using aerial (helicopter only) applications.

To broaden the spectrum of control, Garlon™4 may be tank mixed with this product at the rate of 0.5 to 2 quarts per acre. The rate of Garlon should not exceed one-half of the rate of this product (e.g. 1 quart of Garlon to 2 quarts of this product) for best results.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

**AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of any desirable vegetation or crop(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within a minimum of 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desired vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.
5. **APPLY BY AIR ONLY TO NONRESIDENTIAL AREAS.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

**Ensure uniform application** - To avoid streaking, uneven, or overlapped application, use appropriate marking devices.

**PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear is most susceptible.

**Aerial Equipment**

Use the recommended rates of Mirage Plus in 3 to 15 gallons of water per acre. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

**AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

Use the following guidelines when aerial applications are made near crops or desirable perennial vegetation after bud break and before total leaf drop, and/or near other desirable vegetation or annual crops.

1. Do not apply within 100 feet of all desirable vegetation or crops(s).
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crop(s), do not apply within 500 feet of the desirable vegetation or crop(s).
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crop(s) may require buffer zones in excess of 500 feet.
4. Do not apply when winds are in excess of 10 miles per hour or when inversion conditions exist.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configuration that dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

**Ensure uniform application** - To avoid streaking, uneven, or over-lapped application, use appropriate marking devices. Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of Mirage Plus accumulated during spraying or from spills. **PROLONGED EXPOSURE OF MIRAGE PLUS TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413, may prevent corrosion.

**FOR AERIAL APPLICATION IN FRESNO COUNTY,  
CALIFORNIA ONLY (FROM FEBRUARY 15  
THROUGH MARCH 31 ONLY)**

6/31

**Applicable Area**

This only applies to the area contained inside the following boundaries within Fresno County, California.

North: Fresno County line  
South: Fresno County line  
East: State Highway 99  
West: Fresno County line

**General Information**

Always read and follow the label directions and precautionary statements for all products used in the aerial application.

Observe the following directions to minimize off-site movement during aerial application of Mirage Plus. Minimization of off-site movement is the responsibility of the grower, Pest Control Advisor and aerial applicator.

**Written Recommendations**

A written recommendation **MUST** be submitted by or on behalf of the applicator to the Fresno County Agricultural Commissioner 24 hours prior to the application. This written recommendation **MUST** state the proximity of surrounding crops, and that conditions of each manufacturer's product label and this label have been satisfied.

**Aerial Applicator Training and Equipment**

Aerial application of Mirage Plus is limited to pilots who have successfully completed a Fresno County Agricultural Commissioner and California Department of Pesticide Regulation approved training program for aerial application of herbicides. All aircraft must be inspected, critiqued in flight and certified at a Fresno County Agricultural Commissioner approved fly-in. Test and calibrate spray equipment at intervals sufficient to insure that proper rates of herbicides and adjuvants are being applied during commercial use. Applicator must document such calibrations and testing. Demonstration of performance at Fresno County Agricultural Commissioner approved fly-ins constitutes such documentation, or other written records showing calculations and measurements of flight and spray parameters acceptable to the Fresno County Agricultural Commissioner.

To report known or suspected misuse of Mirage Plus, call 1-800-332-3111.

For additional information on the proper aerial application of Mirage Plus, call 916-784-1718.

**NOTE:** For aerial application from April 1 through February 14, refer to the "For Aerial Application in California Only" section of the label.

**FOR AERIAL APPLICATION IN ARKANSAS ONLY**

**Use Directions**

**AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

Use the recommended rate of Mirage Plus in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply Mirage Plus when winds are in excess of 10 miles per hour. Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 miles per hour.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

**Aerial Equipment**

**DO NOT APPLY MIRAGE PLUS USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.**

# MIRAGE PLUS

## EPA REG. NO. 34704-

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. Unless otherwise specified, do not exceed 1 quart per acre. Refer to the individual use area sections of this label for recommended volumes, application rates, and further instructions.

FOR AERIAL APPLICATION IN CALIFORNIA OR SPECIFIC COUNTIES THEREIN, OR ARKANSAS, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE OR COUNTY FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS.

Mirage Plus plus dicamba tank mixtures may not be applied by air in California.

Ensure uniform application - To avoid streaked, uneven or overlapped application, use appropriate marking devices.

### AERIAL SPRAY DRIFT MANAGEMENT

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outermost nozzles on the boom must not exceed  $\frac{3}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward, parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

#### Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see the "Wind", "Temperature and Humidity", and "Temperature Inversions" sections of this label).

#### Controlling Droplet Size

**Volume:** Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with the higher rated flows produce larger droplets.

**Pressure:** Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rate are needed, use higher flow rate nozzles instead of increasing pressure.

**Number of Nozzles:** Use the minimum number of nozzles that provide uniform coverage.

**Nozzle Orientation:** Orienting nozzles so that the spray is released backwards, parallel to the air stream, will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

**Nozzle Type:** Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger produce larger droplets than other nozzle types.

**Boom Length:** For some use patterns, reducing the effective boom length to less than  $\frac{3}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

**Application Height:** Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces the exposure of the droplets to evaporation and wind.

#### Swath Adjustment

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller droplets, etc.).

#### Wind

Drift potential is lowest between wind speeds of 2 to 10 miles per hour. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 miles per hour due to variable wind directions and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

#### Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

#### Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

#### Sensitive Areas

Mirage Plus should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endanger species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

Avoid direct application to any body of water.

#### Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of Mirage Plus accumulated during spraying or from spills. **PROLONGED EXPOSURE OF MIRAGE PLUS TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint), which meets aerospace specifications MIL-C-38413, may prevent corrosion.

### FOR DISTRIBUTION AND USE ONLY WITHIN SOUTH DAKOTA. FOR NON-SELECTIVE CONTROL OF LISTED ANNUAL WEEDS IN SMALL GRAIN CROPPING SYSTEMS

#### Use Directions

Refer to the label of Mirage Plus for rate recommendations and weeds controlled.

For ground applications, use 3 to 5 gallons of water per acre. For aerial applications, use 2 to 3 gallons of water per acre.

#### ATTENTION

**AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING MIRAGE PLUS TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS**

Do not allow the herbicide solution to mist, drip, or splash onto desirable vegetation since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of injury occurring from the use of Mirage Plus is greatest when winds are gusty or in excess of 5 mile per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) that are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

**NOTE:** To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Adjust boom height on ground equipment to prevent streaked, overlapped or uneven applications. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations that dispense spray as fine spray droplets.

In aerial applications, do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Ensure uniform application. Use appropriate marking devices when applying herbicides by air.

Avoid spraying when weeds are subject to moisture stress, when dust is on foliage, or when straw canopy covers the weeds.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residue of Mirage Plus accumulated during spraying or from spills. **PROLONGED EXPOSURE OF MIRAGE PLUS TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint), which meets aerospace specification MIL-C-38413 may prevent corrosion.

#### Ground Broadcast Equipment

Use the recommended rates of Mirage Plus in 3 to 40 gallons of water per acre as a broadcast spray unless otherwise specified. As density of weeds increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzles to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

#### Hand-Held or High-Volume Equipment

Apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use coarse sprays only.

For recommended rates and timing, refer to the "Annual Weeds - Hand-Held Or High-Volume Equipment" section of the label.

#### Selective Equipment

Mirage Plus may be applied through re-circulating spray systems, shielded applicators, hooded sprayers, wiper applicators or sponge bars, after dilution and thorough mixing with water, to listed weeds growing in any non-crop site specified on this label.

In cropping systems, hooded sprayers, shielded sprayers, and wipers may be used in row middles (in between rows of crop plants) where any dripping or leaking will not contact crop foliage, when listed under "Types Of Application" in the crop sections of the labeling. Such equipment must be capable of preventing all crop contact with herbicide solutions and operated without leakage of spray mists or dripping onto crop. Wipers over-the-top of crops may be used only when specifically recommended in Mirage's labeling.

**AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desirable vegetation should be adjusted so that the lowest spray stream or wiper contact point is at least 2 inches above the desirable vegetation. Droplets, mist, foam or splatter of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

7131

# MIRAGE PLUS

## EPA REG. NO. 34704-

Applications made above the crops should be made when the weeds are a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatment may be necessary.

### Recirculating Spray System

A recirculating spray system directs the spray solution onto weeds growing above desirable vegetation, while spray solution not intercepted by weeds is collected and returned to the spray tank for reuse.

### Shielded and Hooded Applicators

When applied under the conditions described in the following paragraphs for shielded and hooded applications, Mirage Plus at recommended rates will control those weeds listed in the "Annual Weeds Rate Table" and "Perennial Weeds Rate Table" sections of this label. A hooded sprayer is a type of shielded applicator where the spray pattern is fully enclosed including top, sides, front and back, thereby shielding the crop from spray solution. Keep shields on these sprayers adjusted to protect desirable vegetation. When applying to crops grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows. **EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE WITH DESIRABLE VEGETATION.**

This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

Use hoods designed to minimize excessive dripping or run-off down the insides of the hoods. A single, low pressure/low drift flat-fan nozzle with an 80 to 95 degree spray angle positioned at the top center of the hood is recommended. Spray volume should be 20-30 gallons per acre.

These procedures will reduce the potential for crop injury:

- The spray hoods must be operated on the ground or skimming across the ground.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 miles per hour to avoid bouncing of the spray hoods.
- Maximum wind speed: 10 miles per hour.
- Use low-drift nozzles that provide uniform coverage within the treated area.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply Mirage Plus when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

### Wiper Applicators

When applied under the conditions described in the following paragraphs, Mirage Plus controls many weeds, including volunteer corn, Texas panicum, common rye, shatter-cane, sicklepod, spanishneedles and bristly starbur; and SUPPRESSES many weeds including Florida beggarweed, Bermudagrass, hemp dogbane, dogfennel, guineagrass, johnsongrass, milkweed, silverleaf nightshade, redroot pigweed, giant ragweed, smutgrass, sunflower, Canada thistle, musk thistle, vaseygrass, velvetleaf.

Wiper applicators are devices that physically wipe appropriate amounts of Mirage Plus directly onto the weed.

Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Operate this equipment at ground speeds no greater than 5 miles per hour. Performance may be improved by reducing speed in areas of heavy infestations to ensure adequate wiper saturation. Better results may be obtained if two applications are made in opposite directions.

Avoid leakage or dripping onto desirable vegetation. Adjust height of applicator to ensure adequate contact with weeds. Keep wiping surfaces clean. Be aware that, on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying of the wicks on the upper end of a wiper applicator.

Do not use wiper equipment when weeds are wet.

Mix only the amount of solution to be used during a 1-day period, as reduced activity may result from use of leftover solutions. Clean wiper parts immediately after using Mirage Plus by thoroughly flushing with water.

Do not add surfactant to the herbicide solution.

**For Rope or Sponge Wick Applicators** - Solutions ranging from 33 to 75 percent of Mirage Plus in water may be used. Apply this solution to weeds listed in this section.

**For Panel Applicators** - Solutions ranging from 33 to 100 percent of Mirage Plus in water may be used in panel wiper applicators.

### Injection Systems

Mirage Plus may be used in aerial or ground injection spray systems. It may be used as a liquid concentrate or diluted prior to injecting into the spray stream. Do not mix Mirage Plus with the concentrate of other products when using injections systems.

### CDA Equipment

The rate of Mirage Plus applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 2 to 15 gallons of water per acre. 8/31

For the control of annual weeds with hand-held CDA units, apply a 20 percent solution of Mirage Plus at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 miles per hour (1 quart per acre). For the control of perennial weeds, apply a 20 to 40 percent solution of Mirage Plus at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 miles per hour (2 to 4 quarts per acre).

Controlled droplet application equipment produces a spray pattern that is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation, as damage or destruction may result.

### ANNUAL AND PERENNIAL CROPS (Alphabetical)

**NOTE:** THIS SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL LISTED CROPS WITHIN "ANNUAL AND PERENNIAL CROPS" GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, AND ADDITIONAL PRECAUTIONS AND RESTRICTIONS.

See the "Mirage Plus Tolerant Crops" section of this label for treating glyphosate tolerant crops.

### Types Of Applications:

Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, At-Planting, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Applications in Row-Middles, and Post-Harvest Treatments.

### General Use Instructions:

Apply Mirage Plus during fallow intervals preceding planting, prior to planting or transplanting, at-planting, or preemergent to annual and perennial crops listed in this label, except where specifically limited. For any crop not listed in this label, applications must be made at least 30 days prior to planting. Unless otherwise specified, weed control applications may be made according to the rates listed in the "Annual Weeds, Perennial Weeds, and Woody Brush Rate Tables" in this label. Repeat applications may be made up to a maximum of 8 quarts per acre per year.

Post-directed hooded sprayers and wiper equipment capable of preventing all crop contact with herbicide solutions may be used in mulched or unmulched row middles after crop establishment. Where specifically noted below, wipers may also be used above certain crops to control tall weeds. Refer to the "Selective Equipment" section of this label for essential precautions when using hooded sprayers or wipers to avoid crop injury caused by leakage of spray mists or dripping onto crops. Crop injury is possible with these applications and shall be the sole responsibility of the applicator.

The maximum use rates stated throughout Mirage Plus's labeling apply to Mirage Plus combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rate.

### General Precautions, Restrictions:

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death to emerged seedlings. Apply before seed germination in coarse sandy soils to further minimize the risk of injury. Unless otherwise specified in Mirage Plus's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "Application Equipment And Techniques" section of this label for additional information.

In crops where spot treatments are allowed, do not treat more than 10 percent of the total field to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason.

For broadcast postemergent treatments, do not harvest or feed treated vegetation for 8 weeks following application, unless otherwise specified.

### CEREAL AND GRAIN CROPS

**Labeled Crops:** Barley, Buckwheat, Millet (pearl, proso), Oats, Rice, Quinoa, Teff, Teosinte, Triticale, Wheat (all types), Wild rice.

**Precautions, Restrictions:** Do not treat rice fields or levees when field contains water.

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: Red Rice Control Prior to Planting Rice, Spot Treatment (except rice), Over-the Top Wiper Applicators (Feed Barley and Wheat Only), Preharvest (Feed Barley and Wheat Only).

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied before, during or after planting of cereal crops. Applications must be made prior to emergence of the crop.

### Red Rice Control Prior to Planting Rice

**Use Instructions:** Apply 1.5 quarts of Mirage Plus in 5 to 10 gallons of water per acre. Flush fields prior to application to obtain uniform germination and stand of red rice. Make



# MIRAGE PLUS

## EPA REG. NO. 34704-

9/31

application when the majority of the red rice plants are in the 2-leaf stage and no more than 4 inches tall. Red rice plants with less than 2 true leaves may be only partially controlled.

**Precautions, Restrictions:** Avoid spraying during low humidity conditions, as reduced control may result. Do not treat rice fields or levees when the field contains floodwater. Do not re-flood treated fields for 8 days following application.

### FOR CONTROLLING BARNYARDGRASS (*ECHINOCHLOA CRUS-GALLI*) IN RICE USING RENOVATION TREATMENTS IN CALIFORNIA ONLY

#### Use Directions

##### Renovation Treatment

**Use Instructions:** Mirage Plus may be applied as a renovation treatment in rice crops to control barnyardgrass infestations using ground broadcast spray or hand-held equipment. Renovation is defined as herbicide treatment that will produce crop and weed destruction in an entire field or contiguous area treated within a field. Follow the application methods and recommended treatment rates in this label.

**Precautions, Restrictions:** The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason. The rice straw and stubble from the treated area, including a 25-foot buffer zone on all sides, shall not be used for grazing, animal bedding or any feed purposes. No aerial applications are permitted for rice renovation using this label.

##### Spot Treatment (Except Rice)

**Use Instructions:** Mirage Plus may be applied as a spot treatment in cereal crops. Apply Mirage Plus before heading in small grains.

**Precautions, Restrictions:** Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

##### Over-the-Top Wiper Applicators (Feed Barley and Wheat Only)

**Use Instructions:** Wiper applications may be used in wheat and feed barley. To control common rye or cereal rye, apply after the weeds have headed and achieved maximum growth, and when the rye is at least 6 inches above the wheat crop.

**Precautions, Restrictions:** Allow at least 35 days between application and harvest. Do not use roller applicators.

##### Preharvest (Feed Barley and Wheat Only)

**Use Instructions:** Mirage Plus provides weed control when applied prior to harvest of wheat or feed barley. For wheat, apply after the hard-dough stage of grain (30 percent or less grain moisture). For feed barley, apply after the hard-dough stage and when the grain contains 20 percent moisture or less. Stubble may be grazed immediately after harvest.

Mirage Plus may be applied using either aerial or ground spray equipment. For ground applications, apply Mirage Plus in 10 to 20 gallons of water per acre. For aerial applications, apply Mirage Plus in 3 to 10 gallons of water per acre.

**Precautions, Restrictions:** Do not apply more than 1 quart of Mirage Plus per acre. Allow 7 days between application and harvest, feeding, or grazing. Preharvest application is not recommended for wheat or barley grown for seed, as a reduction in germination or vigor may occur.

##### Post-Harvest

**Use Instructions:** Mirage Plus may be applied after harvest of cereal crops. Higher rates may be required for control of large weeds, which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

**Precautions, Restrictions:** For any crop not listed on this label, applications must be made at least 30 days prior to planting the next crop. Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

## CORN

**Types Of Corn:** Field corn, Seed corn, Silage corn, Sweet corn and Popcorn.

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: Preharvest. For Mirage Plus tolerant corn, see the "Mirage Plus Tolerant Crops" section of this label.

##### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied alone or in a tank mixture before, during or after planting corn. Mirage Plus may be tankmixed with the following products provided the specific product listed below is registered for use on these sites. Applications must be made prior to emergence of the crop.

**Tank Mixtures:** Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

2,4-D	Distinct™	Lariat®
Atrazine	Dual Magnum™	Lasso®/Alachlor
Axiom™	Dual II Magnum™	Linex™/Lorox™
Balance™	Epic™	Marksman™
Banvel™/Clarity™	Frontier™/Outlook™	Micro-Tech®
Bicep Magnum™	Fultime™	Prowl™
Bicep II Magnum™	Guardsman™/Leadoff™	Pythron™
Bullet®	Harness®	Simazine
Degree®	Harness Xtra	Topnotch™
Degree Xtra®	Harness Xtra 5.6L	

For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Mirage Plus at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of Mirage Plus per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, use rate may need to be increased for acceptable weed control.

**Precautions, Restrictions:** Applications of 2,4-D or dicamba must be made at least 7 days prior to planting corn.

For Southern states, do not apply in nitrogen solutions to tough-to-control grasses such as barnyardgrass, fall panicum, broadleaf signalgrass, annual ryegrass and any perennial weeds. The area covered by this recommendation includes from Route 50 South in Illinois and Indiana and the following states: Alabama, Arkansas, Delaware, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, New Jersey, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia and West Virginia.

The tank-mix recommendations in this section are not registered in California.

##### Hooded Sprayers

**Use Instructions:** Mirage Plus may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "Application Equipment And Techniques" section of this label.

**Precautions, Restrictions:** Corn must be at least 12 inches tall, measured without extending leaves. Contact of Mirage Plus in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. Do not apply more than 1 quart of Mirage Plus per acre for each application and no more than 3 quarts per acre per year for hooded sprayer applications.

##### Spot Treatment

**Use Instructions:** For spot treatments, apply Mirage Plus prior to silking of corn.

**Precautions, Restrictions:** Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

##### Preharvest

**Use Instructions:** Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed). For ground applications, apply up to 3 quarts of Mirage Plus per acre. For aerial applications, apply up to 2 quarts of Mirage Plus per acre.

**Precautions, Restrictions:** Allow a minimum of 7 days between application and harvest, feeding or grazing. Preharvest application is not recommended for corn grown for seed, as a reduction in germination or vigor may occur.

##### Post-Harvest

**Use Instructions:** Mirage Plus may be applied after harvest of corn. Higher rates may be required for control of large weeds, which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

**Precautions, Restrictions:** Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

### TANK MIX WITH AIM® HERBICIDE FOR IMPROVED SYMPTOMS IN PRE-PLANT APPLICATIONS IN CORN, SOYBEANS OR WHEAT

#### Use Directions

This tank mixture will significantly enhance the speed of symptomatology appearance on certain weeds when applied as a pre-plant treatment prior to corn, glyphosate tolerant corn, soybeans, glyphosate tolerant soybeans, or wheat. This mixture may be applied prior to planting or emergence of labeled crops. This mixture will not provide residual control of weeds that are un-emerged at the time of treatment.

Combine Mirage Plus (32 fluid ounces of product per acre) and Aim (0.144 ounce per acre) in the spray tank in 10 to 20 gallons of water per acre. For best results, make applications to actively growing weeds. Ammonium sulfate at 1 to 2 percent by weight or 8.5 to 17 pounds per 100 gallons of water may be added. The addition of 2,4-D or dicamba is not recommended.

#### Specific Mixing Recommendations

##### Option 1

1. Fill the spray tank one-half full with water and start the agitation.
2. Add Aim at 0.144 ounce per acre and mix thoroughly.
3. If desired, add ammonium sulfate at 1 to 2 percent by weight or 8.5 to 17 pounds per 100 gallons of spray.
4. Add Mirage Plus at 32 fluid ounces per acre and finish filling the spray tank with water.

##### Option 2

1. Fill the spray tank one-half full with water and start the agitation.
2. If desired, add ammonium sulfate at 1 to 2 percent by weight or 8.5 to 17 pounds per 100 gallons of spray.
3. Add Mirage Plus at 32 fluid ounces per acre.
4. Prepare a slurry of Aim with water and add to the spray tank.

Read and follow the Aim label for additional application instructions and precautions.

### TANK MIX WITH RESOURCE® FOR IMPROVED SYMPTOMS IN PRE-PLANT APPLICATIONS IN CORN OR SOYBEANS

#### Use Directions

This tank mixture will significantly enhance the speed of symptomatology on certain weeds when applied as a pre-plant treatment prior to corn or soybeans. This mixture

# MIRAGE PLUS

## EPA REG. NO. 34704-

6/31

may be applied prior to planting or emergence of labeled crops. This mixture will not provide residual control of weeds that are un-emerged at the time of treatment.

Combine Mirage Plus (32 fluid ounces of product per acre) and Resource (2.08 fluid ounces per acre) in the spray tank in 10 to 20 gallons of water per acre. For best results, make applications to actively growing weeds. Ammonium sulfate at 1 to 2 percent by weight or 8.5 to 17 pounds per 100 gallons of water may be added. The addition of 2,4-D or dicamba is not recommended.

### Specific Mixing Recommendations

#### Option 1

1. Fill the spray tank one-half full with water and start the agitation.
2. Add Resource at 2.08 fluid ounces per acre and mix thoroughly.
3. If desired, add ammonium sulfate at 1 to 2 percent by weight or 8.5 to 17 pounds per 100 gallons of spray.
4. Add Mirage Plus at 32 fluid ounces per acre and finish filling the spray tank with water.

#### Option 2

1. Fill the spray tank one-half full with water and start the agitation.
2. If desired, add ammonium sulfate at 1 to 2 percent by weight or 8.5 to 17 pounds per 100 gallons of spray.
3. Add Mirage Plus at 32 fluid ounces per acre.
4. Add Resource at 2.08 fluid ounces per acre to the spray tank.

Read and follow the Resource label for additional application instructions and precautions.

## COTTON

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: Selective equipment, Spot Treatment, Preharvest.

### FOR NEW COTTON LINES WITH THE MIRAGE PLUS TOLERANT GENE - IN-CROP APPLICATIONS

#### General Information

ATTENTION: LOVELAND PRODUCTS, INC. RECOMMENDS MIRAGE PLUS FOR USE ONLY OVER-THE-TOP OF OR DIRECTED ONTO IMPROVED COTTON VARIETIES THAT ARE DESIGNATED AS COTTON WITH THE MIRAGE PLUS TOLERANT GENE. SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE MIRAGE PLUS TOLERANT GENE ARE SPRAYED WITH MIRAGE PLUS. AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, OR ANY DESIRABLE PLANTS AND TREES, OTHER THAN CROPS WITH THE MIRAGE PLUS TOLERANT GENE, SINCE SEVERE INJURY OR DESTRUCTION WILL RESULT.

MIRAGE PLUS TOLERANT COTTON VARIETIES MUST BE PURCHASED FROM AN AUTHORIZED LICENSED SEED SUPPLIER. THE DESIGNATION, "MIRAGE PLUS TOLERANT", INDICATES THE COTTON VARIETY CONTAINS A PATENTED PROPRIETARY TRAIT.

#### Use Directions

Mirage Plus will control many troublesome weeds with over-the-top, post-directed, hooded sprayer, or preharvest applications in glyphosate tolerant cotton. See the APPLICATION EQUIPMENT AND TECHNIQUES section of the label for more information.

#### Maximum Allowable Yearly Rates

Combined total per year for all applications	8 quarts per acre
Preplant, Preemergence applications	5 quarts per acre
Total over-the-top applications from cracking to layby	2.5 quarts per acre
Total precision post-direct or hooded application through layby	2 quarts per acre
Maximum preharvest application rate	2 quarts per acre

**NOTE:** Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16-48 fluid ounces per acre of Mirage Plus.

**Sprayer Preparation:** It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of Mirage Plus to glyphosate tolerant cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

**There are no rotational crop restrictions following applications of Mirage Plus.**

**Over-the-top applications:** Up to 2.5 quarts per sprayed acre of Mirage Plus may be applied by aerial or ground broadcast application equipment postemergence to Mirage Plus tolerant cotton from the ground cracking stage until layby. The "Annual and Perennial Weeds Rate Tables" in the label booklet for Mirage Plus should be used to determine application rate. Any single over-the-top application should not exceed 2 quarts per sprayed acre. Sequential applications of Mirage Plus must be at least 7 days apart.

**With ground broadcast equipment:** apply Mirage Plus in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat-fan nozzles. Check for even distribution of spray droplets.

For aerial applications, apply Mirage Plus in 3 to 15 gallons of water per acre. DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF MIRAGE PLUS WHEN MAKING APPLICATIONS BY AIR. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING MIRAGE PLUS TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS, WHICH DO NOT CONTAIN THE MIRAGE PLUS TOLERANT GENE. Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions that favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

**Post-directed or hooded applications:** In addition to the over-the-top applications, up to 2 quarts per sprayed acre may be applied as a post-directed hooded application to Mirage Plus tolerant cotton through layby. These application methods may be preferred when there is a need to direct the spray onto weeds that are growing under the crop canopy. Equipment should be used which directs the spray into the lower crop canopy so that weeds in the row are covered. For best results, make applications while weeds are small (less than 3 inches). Sequential in-crop applications must be at least 7 days apart from any other in-crop application of Mirage Plus.

**ATTENTION: USE OF MIRAGE PLUS IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF MIRAGE PLUS TOLERANT COTTON, HOWEVER, VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS MAKE IT IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH MIRAGE PLUS, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN UNANTICIPATED RESULTS INCLUDING YIELD LOSS.**

**Weeds controlled.** For specific rates of application and instructions for control of various annual and perennial weeds, refer to the "Annual and Perennial Weeds Rate Table" sections of the label. Mirage Plus, applied at 1 to 2 quarts per acre, will control or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common Bermudagrass, silverleaf nightshade, trumpet creeper, and redvine. Fall preharvest applications may be required for control of these perennial weeds.

Tank mixtures with other herbicides may result in reduced weed control, or may cause crop injury and are not recommended for applications where the spray contacts the cotton plant.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of Mirage Plus for control.

**Preharvest applications:** Mirage Plus may be applied for preharvest annual and perennial weed control as a broadcast treatment to glyphosate tolerant cotton any time after layby up to 7 days prior to harvest. Mirage Plus may be tank mixed with DEFT™ 6, Folex™, Ginstar, or Prep™ to enhance cotton leaf drop. Allow a minimum of 7 days between final application and harvest. No more than 2 quarts of Mirage Plus per acre may be applied preharvest (between layby and seven days prior to harvest). Do not apply more than 1 quart of Mirage Plus per acre by air. **NOTE:** Mirage Plus will not enhance the performance of harvest aids when applied to glyphosate tolerant cotton. Do not apply Mirage Plus preharvest to cotton grown for seed.

**Precautions, Restrictions:** The combined total application from crop emergence until harvest must not exceed 8 quarts per acre. Allow a minimum of 7 days between final application and harvest. Tank mixtures with other herbicides may result in reduced weed control, or may cause crop injury and are not recommended for applications where the spray contacts the cotton plant. Do not apply Mirage Plus preharvest to cotton grown for seed.

Also see the "Mirage Plus Tolerant Crops" section of this label.

#### Preplant, Preemergence, At-Planting

**Use instructions:** Mirage Plus may be applied alone or in tank-mixture, before, during or after planting cotton. Applications must be made prior to emergence of the crop.

**Tank Mixtures:** Apply these tank mixtures in 10 to 20 gallons of water per acre.

Caparol®	Dual II Magnum
Clarity	Karmex
Command	Meturon®
Cotoran®	Prowl
Cotton-Pro®	Staple®
Direx	Zonal®
Dual Magnum	2,4-D

**Precautions, Restrictions:** Refer to individual product labels for rates, restrictions, precautionary statements and preplant intervals.

#### Hooded Sprayer, Selective Equipment

**Use instructions:** Mirage Plus may be applied through hooded sprayers, recirculating sprayers, shielded applicators or wiper applicators in cotton. Allow at least 7 days between application and harvest.

**Precautions, Restrictions:** See the "Selective Equipment" part of the "Application Equipment And Techniques" section of this label for information on proper use and calibration of this equipment.

#### Spot Treatment

**Use instructions:** For spot treatments, apply Mirage Plus prior to boll opening of cotton.

**Precautions, Restrictions:** Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

# MIRAGE PLUS

## EPA REG. NO. 34704-

### Preharvest

**Use Instructions:** Mirage Plus provides weed control and cotton regrowth inhibition when applied prior to harvest of cotton. For weed control, apply at rates given in the "Annual, Perennial and Woody Brush Weed Control Table" sections of this label. For cotton regrowth inhibition, apply 1 pint to 2 quarts of Mirage Plus per acre.

Up to 2 quarts of Mirage Plus may be applied using either aerial or ground spray equipment. Apply after sufficient bolls have developed to produce the desired yield of cotton. Applications made prior to this time could affect maximum yield potential.

**Tank Mixtures:** Mirage Plus may be tank mixed with Def™ 6, Folex™, Ginstar, or Prep™ to provide additional enhancement of cotton leaf drop.

**Precautions, Restrictions:** Allow a minimum of 7 days between application and harvest of cotton. Preharvest application is not recommended for cotton grown for seed, as a reduction in germination or vigor may occur. THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF MIRAGE PLUS TO COTTON IS PROHIBITED.

### Fallow Systems

**Labeled Crops:** Mirage Plus may be applied during the fallow period prior to planting or emergence of any crop on this label.

**Types Of Applications:** Chemical fallow, Preplant fallow beds, Aid-to-tillage.

### Chemical Fallow

**Use Instructions:** Mirage Plus may be applied during the fallow period prior to planting or emergence of any crop listed on this label. Mirage Plus may be used as a substitute for tillage to control annual weeds in fallow fields. Also, broadcast or spot treatments will control or suppress many perennial weeds in fallow fields. Ground or aerial application equipment may be used. Tank mixtures with 2,4-D and dicamba may be used. Applications up to 2 quarts per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift to adjacent crops.

**Precautions, Restrictions:** For any crop not listed on this label, applications must be made at least 30 days prior to planting. Do not apply dicamba tank mixtures by air in California.

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba is applied within 45 days of planting.

### Preplant Fallow Beds

**Use Instructions:** Mirage Plus may be applied to fallow beds prior to planting or emergence of any crop listed on this label. For any crop not listed on this label, applications must be made at least 30 days prior to planting. Mirage Plus will control weeds listed in the "Annual, Perennial and Woody Brush Weed Control Table" sections of this label.

**Tank Mixtures:** In addition, 12 fluid ounces of Mirage Plus plus 2 to 3 fluid ounces of Goal™ 2XL per acre will control the following weeds with the maximum height or length indicated: 3" - common cheeseweed, chickweed, groundsel; 6" - London rocket, shepherd's-purse.

16 fluid ounces of Mirage Plus plus 2 to 3 ounces of Goal™ 2XL per acre will control the following weeds with the maximum height or length indicated: 6" - common cheeseweed, groundsel, marehail (*Coryza canadensis*), 12" - chickweed, London rocket, shepherd's-purse.

### Aid-to-Tillage

**Use Instructions:** Mirage Plus may be used in conjunction with tillage practices in fallow systems or preplant to labeled crops to control downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 12 fluid ounces of Mirage Plus in 3 to 10 gallons of water per acre. Make applications before weeds are 6 inches in height. Application must be followed by conventional tillage practices no later than 15 days after treatment and before regrowth occurs. Allow at least 1 day after application before tillage.

**Precautions, Restrictions:** Tank mixtures with residual herbicides may result in reduced performance.

## GRAIN SORGHUM (MILO)

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: Spot Treatment, Over-the-Top Wiper Applicators, Preharvest.

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied alone or in tank mixture before, during or after planting grain sorghum. Mirage Plus may be tankmixed with the following products provided the specific product listed below is registered for use on these sites. Applications must be made prior to emergence of the crop.

**Tank Mixtures:** Apply these tank mixtures in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre.

Atrazine	Lariat
Bicep II Magnum	Lasso
Bullet	Micro-Tech
Dual II Magnum	

1/31  
For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Mirage Plus at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of Mirage Plus per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall. When using nitrogen solutions as the carrier, the use rate may need to be increased for acceptable weed control.

### Spot Treatment and Over-the-Top Wiper Applications

**Use Instructions:** Mirage Plus may be applied as a spot treatment in grain sorghum. Make spot treatments before heading of milo. Mirage Plus may be applied with wiper applicators to control or suppress the weeds listed under "Wiper Applicators" in the "Selective Equipment" section of this label.

**Precautions, Restrictions:** For spot treatment, do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

For wiper applicators, allow at least 40 days between application and harvest. Do not use roller applicators. Do not feed or graze treated milo fodder. Do not ensile treated vegetation.

### Hooded Sprayers

**Use Instructions:** Mirage Plus may be used through hooded sprayers for weed control between the rows of milo. Only hooded sprayers that completely enclose the spray pattern may be used. See additional instruction for the use of hooded sprayers in the "Application Equipment And Techniques" section of this label.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply Mirage Plus when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

**Precautions, Restrictions:** Milo must be at least 12 inches tall, measured without extending leaves. Treat before milo sends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed. Contact of Mirage Plus in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator. Do not graze or feed milo forage or fodder following applications of Mirage Plus through hooded sprayers. Do not apply more than 1 quart of Mirage Plus per acre per application and no more than 3 quarts per acre per year for hooded sprayer applications.

### Preharvest

**Use Instructions:** Make applications at 30 percent grain moisture or less.

**Precautions, Restrictions:** Do not apply more than 2 quarts of Mirage Plus per acre. As with other herbicides that cause sudden plant death, avoid preharvest applications of Mirage Plus to milo infected with charcoal rot as lodging can occur. Allow a minimum of 7 days between application and harvest, feeding, or grazing of sorghum. Preharvest application is not recommended for sorghum grown for seed, as a reduction in germination or vigor may occur. The use of Mirage Plus for preharvest grain sorghum (milo) is not registered in California.

### Post-Harvest

**Use Instructions:** Mirage Plus may be applied after harvest of grain sorghum. Higher rates may be required for control of large weeds, which were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

Mirage Plus may be applied to grain sorghum (milo) stubble following harvest to suppress or control regrowth. Apply 1 quart of Mirage Plus per acre for control, or 1.5 pints of Mirage Plus per acre for suppression.

**Precautions, Restrictions:** Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

## HERBS AND SPICES

**Labeled Crops:** Allspice, Angelica, Star anise, Annatto (seed), Balm, Basil, Borage, Burnet, Chamomile, Caper buds, Caraway, Black caraway, Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chive, Chinese chive, Cinnamon, Clary, Clove buds, Coriander leaf (cilantro or Chinese parsley), Coriander seed (cilantro), Costmary, Culantro (leaf), Culantro (seed), Cumin, Curry (leaf), Dill (dillweed), Dill (seed), Epazote, Fennel seed (common and Florence), Fenugreek, White ginger flower, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf and seed), Mace, Marigold, Marjoram (including oregano), Mexican oregano, Mioga flower, Mustard (seed), Nasturtium, Nutmeg, Parsley (dried), Pennyroyal, Pepper (black and white), Pepper leaves, Peppermint, Perilla, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory (summer and winter), Spearmint, Stevia leaves, Sweet bay, Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: Over-the-Top Wiper Applications (Peppermint and Spearmint Only), Spot Treatments (Peppermint and Spearmint Only).

**Precautions, Restrictions:** When applying Mirage Plus prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of Mirage Plus, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5 -inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to ensure that the water flushes off the plastic mulch and does not enter the transplant holes. For some crops below, it is recommended to make applications 3 days before transplanting or planting.

# MIRAGE PLUS

## EPA REG. NO. 34704-

12/31

### Over-the-Top Wiper Applications or Spot Treatments (Peppermint and Spearmint Only)

**Use Instructions:** Mirage Plus may be used as a spot treatment or wiper application in spearmint and peppermint. Apply spot treatments on a spray-to-wet basis with hand-held equipment, such as backpack and knapsack sprayers, pump-up pressure sprayers, hand-guns hand-wands or any other hand-held or motorized spray equipment used to direct the spray solution to a limited area. In wiper applications, the applicator should be adjusted so that the wiper contact point is at least 2 inches above the crop. Weeds should be a minimum of 6 inches taller than the crop.

**Precautions, Restrictions:** Allow at least 7 days between application and harvest. Further applications may be made in the same area at 30-day intervals. In spot treatment applications, no more than 10 percent of the total field area to be harvested should be treated at one time. The crop receiving spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for this reason. In wiper applications, contact of the herbicide solution with the crop may result in damage or destruction.

### OIL SEED CROPS

**Labeled Crops:** Borage, Buffalo gourd (seed), Canola, Crambe, Flax, Jojoba, Lesquerella, Meadowfoam, Mustard (seed), Rape, Safflower, Sesame, Sunflower. For Mirage Plus tolerant canola, see the "Mirage Plus Tolerant Crops" section of this label.

**Types of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application."

**Use Instructions:** Mirage Plus may be applied before, during or after planting oil seed crops. Broadcast applications must be made prior to emergence of the listed oil seed crops. Wiper applicators or hooded sprayers may be used between the rows once the crop is established.

**Tank Mixtures:** For sunflowers, a tank mixture with Prowl may be applied before, during or after planting in conventional tillage systems, into a cover crop, established sod or in previous crop residue.

**Precautions, Restrictions:** Do not apply more than 2 quarts of Mirage Plus per acre on canola. Do not apply more than 1 quart of Mirage Plus per acre for sunflowers as a single preplant or preemergence application per year. Do not feed or graze sunflower forage following application of Mirage Plus.

### SOYBEANS

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: spot treatment, preharvest, selective equipment. For Mirage Plus tolerant soybeans, see the "Mirage Plus Tolerant Crops" section of this label.

#### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied alone or in a tank mixture before, during or after planting soybeans. Applications must be made prior to emergence of the crop.

**Tank Mixtures:** Apply these tank mixtures in 10 to 20 gallons of water per acre.

Aim™	Gauntlet™
Amplify™	Lasso
Assure II™	Linex™
Authority™	Lorox/Linuron
Boundary™	Lorox Plus™
Canopy™	Micro-Tech
Canopy XI™	Prowl
Command™	Pursuit™
Command Xtra™	Pursuit Plus
Domain™	Reflex™
Dual Magnum	Scepter™
Dual II Magnum	Sencor™/Lexone™
Firstrate™	Squadron™
Flexstar™	Steel™
Frontier™/Outlook™	Valor™
Fusion™	

Mirage Plus may be tank mixed with 2,4-D or 2,4-DB. See the 2,4-D label for intervals between application and planting.

For difficult-to-control annual weeds such as fall panicum, barnyardgrass, crabgrass, shattercane and broadleaf signalgrass up to 2 inches tall, and Pennsylvania smartweed up to 6 inches tall, apply Mirage Plus at 2 pints per acre in these tank mixtures. For other labeled annual weeds, apply 1.5 to 2 pints of Mirage Plus per acre when weeds are less than 6 inches tall, and 2 to 3 pints when weeds are over 6 inches tall.

**Precautions, Restrictions:** Tank mixtures with some of the above listed herbicides may result in reduced weed control due to antagonism. Read and carefully observe the cautionary statements and all other information appearing on the product labels or fact sheets published separately for all herbicides used. Use according to the most restrictive directions for each product in the mixture.

#### Spot Treatment

**Use Instructions:** For spot treatments, apply Mirage Plus prior to initial pod set in soybeans.

**Precautions, Restrictions:** Do not treat more than 10 percent of the total field area to be harvested. The crop receiving spray in treated area will be killed. Take care to avoid drift or spray outside target area for the same reason.

#### Preharvest

**Use Instructions:** Mirage Plus provides weed control when applied prior to harvest of soybeans.

Apply at rates given in the "Annual Weeds, Perennial Weeds And Woody Brush Rate Tables". Mirage Plus may be applied using either aerial or ground spray equipment. Apply after pods have set and lost all green color. Care should be taken to avoid excessive seed shatter loss due to ground application equipment.

**Precautions, Restrictions:** Do not apply more than 5 quarts per acre of Mirage Plus for preharvest applications. Do not apply more than 2 quarts per acre of Mirage Plus by air. Allow a minimum of 7 days between application and harvest of soybeans. Do not graze or harvest treated hay or fodder for livestock feed within 25 days of last preharvest application. (If the application rate is 1 quart per acre or lower, the grazing restriction is reduced to 14 days after last preharvest application.) Preharvest application is not recommended for soybeans grown for seed, as a reduction in germination or vigor may occur.

#### Selective Equipment

**Use Instructions:** Mirage Plus may be applied through recirculating sprayers, shielded applicators, hooded sprayers, over-the-top wiper applicators or sponge bars in soybeans. Allow at least 7 days between application and harvest.

**Precautions, Restrictions:** See the "Selective Equipment" part of the "Application Equipment And Techniques" section of this label for information on proper use and calibration of this equipment.

### SUGARCANE

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application."

#### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied in or around sugarcane fields or in fields prior to the emergence of plant cane.

**Precautions, Restrictions:** Do not apply to vegetation in or around ditches, canals or ponds containing water to be used for irrigation.

#### Spot Treatment

**Use Instructions:** Mirage Plus may be applied as a spot treatment in sugarcane. For control of volunteer or diseased sugarcane, make a 1 percent solution of Mirage Plus in water and spray-to-wet the foliage of vegetation to be controlled. Volunteer or diseased sugarcane should have at least 7 new leaves.

**Precautions, Restrictions:** Avoid spray contact with healthy cane plants since severe damage or destruction may result. Do not feed or graze treated sugarcane foliage following application.

#### Fallow Treatments

**Use Instructions:** Mirage Plus may be used as a replacement for tillage in fields that are lying fallow between sugarcane crops. Mirage Plus may also be used to remove the last stubble of ratoon cane. For removal of last stubble of ratoon cane, apply 4 to 5 quarts of Mirage Plus in 10 to 40 gallons of water per acre to new growth having at least 7 new leaves. Allow 7 or more days after application before tillage. Ground or aerial application equipment may be used. Applications up to 3 quarts per acre may be made by aerial application in fallow sites where there is sufficient buffer to prevent injury due to drift onto adjacent crops. Tank mixtures with 2,4-D and dicamba may be used.

#### Hooded Sprayers

**Use Instructions:** Mirage Plus may be used through hooded sprayers for weed control between the rows of sugarcane. See the "Application Equipment And Techniques" section of this label for additional use instructions.

**Precautions, Restrictions:** Do not allow treated weeds to come into contact with the crop. Droplets, mist, foam or splatter of the herbicide solution settling on the crop may result in discoloration, stunting or destruction. Such damage shall be the sole responsibility of the applicator.

### VEGETABLE CROPS

**NOTE:** THIS "VEGETABLE CROPS" SECTION GIVES GENERAL DIRECTIONS THAT APPLY TO ALL LISTED VEGETABLE CROPS GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

**Types Of Applications:** Chemical Fallow, Preplant Fallow Beds, Preplant, Preemergence, Prior to Transplanting Vegetables, At-Planting, Hooded Sprayers in Row-Middles, Shielded Sprayers in Row-Middles, Wiper Applications in Row-Middles, and Post-Harvest, Directed Applications (Non-bearing Ginseng), Over-the-Top Wiper Applications (Rutabagas Only).

**Precautions, Restrictions:** When applying Mirage Plus prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of Mirage Plus, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to ensure that the water flushes off the plastic mulch and does not enter the transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

# MIRAGE PLUS

## EPA REG. NO. 34704-

Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death to emerged seedlings. In crops with vines, hooded sprayer, shielded sprayer, and wiper application to row middles should be made prior to vine development, otherwise severe injury or destruction may result. Unless otherwise specified in Mirage Plus's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "Application Equipment And Techniques" section of this label for additional information.

### BRASSICA VEGETABLES

**Labeled Crops:** Broccoli, Chinese broccoli (gai lon), Broccoli raab (rapini), Brussels sprouts, Cabbage, Chinese cabbage (bok choy), Chinese cabbage (napa), Chinese mustard cabbage (gai choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens.

### BULB VEGETABLES

**Labeled Crops:** Garlic, Great-headed garlic, Leek, Onion (dry bulb and green), Welsh onion, Shallot.

### CUCURBIT VEGETABLES AND FRUITS

**Labeled Crops:** Chayote (fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cucumber, Gherkin, Edible gourd (includes hyotan, cucuzza, hechima, Chinese okra), Melons (all), *Momordica* spp. (includes balsam apple, balsam pear, bittermelon, Chinese cucumber), Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey ball melon, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon), Pumpkin, Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini), Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash), Watermelon.

**Precautions, Restrictions:** For Cantaloupe, Casaba melon, Crenshaw melon, Cucumber, Gherkin, Gourds, Honeydew melon, Honey ball melon, Mango melon, Melons (all), Muskmelon, Persian melon, Pumpkin, Squash (summer, winter), and Watermelon, allow at least 3 days between application and planting.

### LEAFY VEGETABLES

**Labeled Crops:** Amaranth (Chinese spinach), Arugula (rocket), Beet greens, Cardoon, Celery, Chinese celery, Celtuce, Chaya, Chervil, Edible-leaved chrysanthemum, Garland chrysanthemum, Corn salad, Cress (garden and upland), Dandelion, Dock (sorrel), Dokudami, Endive (escarole), Florence fennel, Gow kee, Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Rhubarb, Spinach, New Zealand spinach, Vine spinach, Swiss chard, Watercress (upland), Water spinach.

**Precautions, Restrictions:** For Watercress, avoid applications within 3 days prior to seeding and during the period between seeding and emergence to minimize the risk of injury.

### FRUITING VEGETABLES

**Labeled Crops:** Eggplant, Ground cherry (*Physalis* spp.), Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Tomato.

**Precautions, Restrictions:** For Eggplant, Ground cherry, Pepper (all), and Tomatillo, allow at least 3 days between application and planting. For Tomato, hooded or shielded sprayer applications in row-middles are not recommended.

### LEGUME VEGETABLES (Succulent or Dried)

**Labeled Crops:** Bean (*Lupinus*: includes grain lupin, sweet lupin, white lupin, and white sweet lupin), Bean (*Phaseolus*: includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean), Bean (*Vigna*: includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean), Broad bean (fava), Chickpea (garbanzo), Guar, Jackbean, Lablab bean, Lentil, Pea (*Pisum*: includes dwarf pea, edible-podded pea, English pea, field pea, garden pea, green pea, snowpea, sugar snap pea), Pigeon pea, Soybean (immature seed), Sword bean.

### ROOT and TUBER VEGETABLES

**Labeled Crops:** Arracacha, Arrowroot, Chinese artichoke, Jerusalem artichoke, Beet (garden), Burdock, Canna, Carrot, Cassava (bitter and sweet), Celeriac, Chayote (root), Chervil (turnip-rooted), Chicory, Chufa, Dasheen (taro), Galangal, Ginger, Ginseng, Horseradish, Leren, Kava (turnip-rooted), Parsley (turnip rooted), Parsnip, Potato, Radish, Oriental radish, Rutabaga, Salsify, Black salsify, Spanish salsify, Skirret, Sweet potato, Tanier, Turmeric, Turnip, Wasabi, Yacon, Yam bean, True yam.

#### Directed Applications (Non-bearing Ginseng Only)

**Use Instructions:** Mirage Plus may be used for general weed control in established non-bearing ginseng. Applications may be made with boom equipment, CDA, shielded sprayers, hand-held and high volume wands, lances, and orchard guns or with wiper application equipment.

**Precautions, Restrictions:** Direct applications so that there is no contact of Mirage Plus with the ginseng plant. Applications must be made at least on year prior to harvest.

#### Over-the-Top Wiper Applications (Rutabagas Only)

**Use Instructions:** Wiper applicators may be used over-the-top of rutabagas.

**Precautions, Restrictions:** Allow at least 14 days between application and harvest of rutabagas.

### MISCELLANEOUS CROPS

**Labeled Crops:** Aloe vera, Asparagus, Bamboo shoots, Globe artichoke, Okra, Peanut (ground nut), Pineapple, Strawberry, Sugar beet.

**Types Of Applications:** Those listed in "Annual and Perennial Crops (Alphabetical) Types of Application" plus the following: General weed control, Site preparations, Spot Treatment (Asparagus).

For glyphosate tolerant sugarbeets, see the "Mirage Plus Tolerant Crops" section of this label.

**Precautions, Restrictions:** Avoid contact of herbicide with foliage, green shoots or stems, bark, exposed roots (including those emerging from plastic mulch), or fruit of crops because severe injury or destruction may result. When making preemergence and at-planting applications, applications must be made before crop emergence to avoid severe crop injury. Broadcast applications made at emergence will result in injury or death to emerged seedlings. In crops with vines, hooded sprayer, shielded sprayer, and wiper application to row-middles should be made prior to vine development otherwise severe injury or destruction may result. Unless otherwise specified in Mirage Plus's labeling, treatments with selective equipment including wipers and hooded sprayers must be made at least 14 days prior to harvest. Post-harvest or fallow applications must be made at least 30 days prior to planting any non-labeled crop. See "Application Equipment And Techniques" section of this label for additional information.

#### General Weed Control, Site Preparation

**Use Instructions:** Mirage Plus may be applied for general weed control or for site preparation prior to planting or transplanting crops listed in this section.

**Precautions, Restrictions:** When applying Mirage Plus prior to transplanting or direct-seeding crops into plastic mulch, care must be taken to remove residues of Mirage Plus, which could cause crop injury, from the plastic prior to planting. Residues can be removed by a single 0.5-inch application of water, either by natural rainfall or via a sprinkler system. Care should be taken to ensure that the water flushes off the plastic mulch and does not enter the transplant holes. Applications made at emergence will result in injury or death to emerged seedlings.

Do not apply within a week before the first asparagus spear emerges. Do not feed or graze treated pineapple forage following application.

#### Spot Treatment (Asparagus)

**Use Instructions:** Mirage Plus may be applied immediately after cutting, but prior to emergence of new spears.

**Precautions, Restrictions:** Do not treat more than 10 percent of the total field area to be harvested. Do not harvest within 5 days of treatment.

#### Post-Harvest (Asparagus)

**Use Instructions:** Mirage Plus may be applied after the last harvest and all spears have been removed. If spears are allowed to regrow, delay application until ferns have developed. Delayed treatments should be applied as a directed or shielded spray in order to avoid contact of the spray ferns, stems or spears.

**Precautions, Restrictions:** Direct contact of spray with the asparagus may result in serious crop injury. Select and use recommended types of spray equipment for post-emergence post-harvest applications. A directed spray is any application where the spray pattern is aligned in such a way as to avoid direct contact of the spray with the crop. A shielded spray is any application where a physical barrier is positioned and maintained between the spray and the crop to prevent contact of spray with the crop.

### TREE, VINE, AND SHRUB CROPS (Alphabetical)

**NOTE:** THIS SECTION GIVE GENERAL DIRECTIONS THAT APPLY TO ALL LISTED TREE, VINE, AND SHRUB CROPS GROUPED ALPHABETICALLY BELOW. SEE THE INDIVIDUAL CROP CATEGORIES FOR SPECIFIC INSTRUCTIONS, PREHARVEST INTERVALS, PRECAUTIONS AND RESTRICTIONS.

**Types Of Applications:** Preplant (Site Preparation) Broadcast Sprays, General Weed Control, Middles (between rows of trees, vines, or bushes), Strips (within rows of trees, vines or bushes), Selective Equipment (Shielded Sprayers, Wiper Applications), Directed Sprays, Spot Treatments, Perennial Grass Suppression, Cut Stump.

Applications may be made with boom equipment, CDA equipment, shielded sprayers, hand-held and high-volume wands, lances, orchard guns or with wiper applicator equipment, except as directed.

#### General Use Instructions:

Mirage Plus may be applied in middles (between rows of trees or vines), strips (within rows of trees or vines), and for general weed control or perennial grass suppression in established tree fruit and nut groves, orchards, berries, and vineyards. It may also be used for site preparation prior to planting or transplanting these crops. Apply 1 pint to 5 quarts per acre according to the "Annual and Perennial Weeds Rate Tables" sections of this label. Utilize rates at the higher end of the recommended rate range when weeds are stressed, growing in dense populations or are greater than 12 inches tall. Repeat applications may be made up to a maximum of 10.6 quarts per acre per year.

The maximum use rates stated throughout Mirage Plus's labeling apply to Mirage Plus combined with the use of all other herbicides containing Mirage Plus or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other Mirage Plus or sulfosate containing products does not exceed stated maximum use rate.

# MIRAGE PLUS

## EPA REG. NO. 34704-

### General Precautions, Restrictions:

Extreme care must be exercised to avoid contact of herbicide solution, spray, drift or mist with foliage or green bark of trunk, branches, suckers, fruit or other parts of trees, canes and vines. Avoid applications when recent pruning wounds or other mechanical injury has occurred. Contact of Mirage Plus with other than matured brown bark can result in serious crop damage or destructions. Only shielded or directed sprayers may be used in crops with potential for crop contact, and then only where there is sufficient clearance. For applications in strips (within rows of trees), only selective equipment (directed sprays, hooded sprayers, shielded applicators, or wipers) should be used to minimize the potential for leakage or drift of herbicide sprays onto crop. For berry crops, hooded or shielded sprayers must be fully enclosed including top, sides, front and back. Only wipers or shielded applicators capable of preventing all contact with crop may be used. See "Application Equipment And Techniques" section of this label for additional directions and precautions.

Allow a minimum 3 days between application and transplanting.

### Middles (between rows)

**Use Instructions:** Mirage Plus will control or suppress annual and perennial weeds and ground covers growing between the rows of labeled tree and vine crops. If weeds are under drought stress, irrigate prior to application. Reduced control may result if weeds have been mowed prior to application.

**Tank Mixtures:** A tank mixture of Mirage Plus plus Goal 2XL may be used for annual weeds in middles between rows of citrus crops, tree fruits, tree nuts and vine crops. This mixture is recommended when weeds are stressed or growing in dense populations. 16 to 32 fluid ounces per acre of Mirage Plus plus 3 to 12 fluid ounces per acre of Goal 2XL will control common cheeseweed (*Malva*) or hairy fleabane (*Conyza bonariensis*) with a maximum height or diameter of 3 inches, and annual weeds with a maximum height or diameter of 6 inches, including crabgrass, common groundsel, junglerice, common lambsquarters, redroot pigweed, London rocket, common ryegrass, shepherd's-purse, annual sowthistle, filaree (suppression), horseweed/marestail (*Conyza canadensis*), stinging nettle and common purslane (suppression).

### Strips (in rows)

**Tank Mixtures:** Mirage Plus may be applied in rows of tree or vine crops in tank mixtures with the following products:

Devrno™ 50 DF	Prowl	Solicam™ DF
Direx™ 4L	Princep Caliber™ 90	Surflan™ AS
Goal 2XL	Simazine 4L	Surflan 75W
Karmex DF	Simazine 80W	
Krovat	Sim-Trol™ 4L	

Do not apply these tank mixtures in Puerto Rico.

Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

### Perennial Grass Suppression

Mirage Plus will suppress perennial grasses such as bahiagrass, Bermudagrass, tall fescue, orchardgrass, Kentucky bluegrass, and quackgrass that are grown as ground covers in tree and vine crops.

For suppression of tall fescue, fine fescue, orchardgrass and quackgrass, apply 8 fluid ounces of Mirage Plus in 10 to 20 gallons of water per acre.

For suppression of Kentucky bluegrass covers, apply 6 fluid ounces of Mirage Plus per acre. Do not add ammonium sulfate.

For best results, mow cool season grass covers in the spring to even their height and apply Mirage Plus 3 to 4 days after mowing.

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of Mirage Plus in 10 to 25 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of Mirage Plus per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

For burndown of Bermudagrass, apply 1 to 2 quarts of Mirage Plus in 3 to 20 gallons of water per acre. Use this treatment only if reduction of the Bermudagrass stand can be tolerated. When burndown is required prior to harvest, allow at least 21 days to ensure sufficient time for burndown to occur.

For suppression of Bermudagrass, apply 6 to 16 fluid ounces of Mirage Plus per acre East of the Rocky Mountains and 16 fluid ounces of Mirage Plus per acre West of the Rocky Mountains. Apply in a total spray volume of 3 to 20 gallons per acre, no sooner than 1 to 2 weeks after full greenup. If the Bermudagrass is mowed prior to applications, maintain a minimum of 3 inches in height. Sequential applications may be made when regrowth occurs and Bermudagrass injury and stand reduction can be tolerated. East of the Rocky Mountains, rates of 6 to 10 fluid ounces of Mirage Plus per acre should be used in shaded conditions or where a lesser degree of suppression is desired.

### Cut Stump (Tree Crops)

**Use Instructions:** Cut stump applications of Mirage Plus may be made during site preparation or site renovation, prior to transplanting tree crops. Mirage Plus will control regrowth of cut stumps and resprouts of many types of tree species, some of which are listed below.

**Citrus Trees:** Calamondin, Chironja, Citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, Tangelo, Tangor.

**Fruit Trees:** Apple, Apricot, Cherry (sweet, sour), Crabapple, Loquat, Mayhaw, Nectarine, Olive, Peach, Pear, Plum/Prune (all), Quince.

**Nut Trees:** Almond, Beechnut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia, Pecan, Pistachio, Walnut (black, English).

Apply Mirage Plus using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of Mirage Plus to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansion.

**Precautions, Restrictions:** DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF ADJACENT DESIRABLE TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. INJURY RESULTING FROM ROOT GRAFTING MAY OCCUR IN ADJACENT TREES. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

### BERRY CROPS

**Labeled Crops:** Blackberry (including bingleberry, black satin berry, boysenberry, Cherokee blackberry, chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, juneberry, lavacaberry, lowberry, lucretiaberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenaberry, rangeberry, ravenberry, rossberry, Shawnee blackberry, and youngberry), Blueberry, Cranberry, Currant, Elderberry, Gooseberry, Huckleberry, Loganberry, Rasperry (black, red), Salal.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application" plus Spot Treatment in Cranberry Production and Post Harvest Treatments in Cranberry Production.

**Precautions, Restrictions:** To avoid damage, herbicide sprays must not be allowed to contact desirable vegetation, including green shoots, canes, or foliage. Allow a minimum of 30 days between last application and harvest in cranberries. Allow a minimum of 14 days between last application and harvest in other berry crops. Do not make directed sprays within the cranberry bush areas prior to berry harvest.

### Spot Treatment in Cranberry Production

**Use Instructions:** Spot treatments may be used to control weeds growing in dry ditches (interior and perimeter) of cranberry production areas. Hand-held sprayers or other appropriate application equipment listed under "Application Equipment And Techniques" in this label may be used. Drop water level to remove standing water in ditches prior to application. In hand-held sprayers, use 1 to 2 percent solution of Mirage Plus. Spray to wet vegetation, not to run-off.

**Precautions, Restrictions:** For treatments after draw down of water in dry ditches, allow 2 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply Mirage Plus within 1 day after draw down to ensure application to actively growing weeds. Allow a minimum of 30 days between last application and harvest of cranberries. Do not apply this material through the irrigation system. Do not make applications by air. Do not apply directly to water. Use nozzles that emit medium- to large-sized droplet to minimize drift in order to avoid crop injury.

### Post-Harvest Treatments in Cranberry Production

**Use Instructions:** Application of Mirage Plus may be made after the harvest of cranberries to control weeds growing within the field. Best results will be obtained if applications are made to vines that appear dormant (after they have turned red). Hand-held sprayers, wipers, or other appropriate application equipment listed under "Application Equipment And Techniques" in this label may be used. If using hand-held sprayers, use a 0.5 to 1 percent solution of Mirage Plus. Spray to wet vegetation, not to run-off. If using hand-held boom sprayers, apply 2 to 4 quarts of Mirage Plus per acre.

**Precautions, Restrictions:** Make applications only after cranberries have been harvested. Do not treat more than 10 percent of the total bog. Allow a minimum of 6 months after last application and next harvest of cranberries. Do not apply Mirage Plus through the irrigation system. Do not make applications by air. Do not apply directly to water. Even though vines appear dormant, contact of herbicide solution with desirable vegetation may result in damage or severe plant injury. Cranberry plants that are directly sprayed may be killed.

### CITRUS

**Labeled Crops:** Calamondin, Chironja, Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (all), Pummelo, Satsuma Mandarin, Tangelo (ugli), Tangor.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

**Use Instructions:** (The recommendations below pertain to applications in Florida and Texas).

For burndown or control of the weeds listed below, apply the recommended rates of Mirage Plus in 3 to 30 gallons of water per acre. Where weed foliage is dense, use 10 to 30 gallons of water per acre.

For goatweed, apply 2 to 3 quarts of Mirage Plus per acre. Apply in 20 to 30 gallons of water per acre when plants are actively growing. Use 2 quarts per acre when plants are



# MIRAGE PLUS

## EPA REG. NO. 34704-

less than 8 inches tall and 3 quarts per acre when plants are greater than 8 inches tall. If goatweed is greater than 8 inches tall, the addition of Krovar™ I or Karmex™ may improve control. Refer to the individual product labels for specific crops, rates, geographic restrictions and precautionary statements.

### Perennial weeds:

S = Suppression  
B = Burndown

PC = Partial control  
C = Control

Weed Species	Mirage Plus Rate Per Acre			
	1 Qt	2 Qt	3 Qt	5 Qt
Bermudagrass	B	-	PC	C
Guineagrass	-	-	-	-
Texas and Florida Ridge	B	C	C	C
Florida Flatwoods	-	B	C	C
Paragrass	B	C	C	C
Torpedograss	S	-	PC	C

**Precautions, Restrictions:** Allow a minimum of 1 day between last application and harvest in citrus crops. For citron groves, apply as directed sprays only.

### MISCELLANEOUS TREE FOOD CROPS

**Labeled Crops:** Cactus (fruit and pads), Palm (heart, leaves), Palm (oil).

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

### NON-FOOD TREE CROPS

**Labeled Crops:** Pine, Poplar, Eucalyptus, Christmas Trees, Other non-food tree crops.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

### Directed Sprays, Spot Treatment, Wiper Applications

**Use Instructions:** Mirage Plus may be used as a post-directed spray and spot treatment around established poplar, eucalyptus, Christmas trees and other non-food tree crops.

**Precautions, Restrictions:** Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established Christmas trees and other pine trees. Desirable plants may be protected from the spray solution by using shields or coverings made of cardboard or other impermeable material. UNLESS OTHERWISE DIRECTED, MIRAGE PLUS IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN CHRISTMAS TREES AND OTHER PINE TREES.

### Site Preparation

**Use Instructions:** Mirage Plus may be used prior to planting Christmas trees.

**Precautions, Restrictions:** Precautions should be taken to protect nontarget plants during site preparation applications.

### POME FRUIT

**Labeled Crops:** Apple, Crabapple, Loquat, Mayhaw, Pear (including oriental pear), Quince.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

**Precautions, Restrictions:** Allow a minimum of 1 day between last application and harvest in pome crops.

### STONE FRUIT

**Labeled Crops:** Apricot, Cherry (sweet, tart), Nectarine, Olive, Peach, Plum/Prune (all types), Plumcot.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

**Precautions, Restrictions:** Allow a minimum of 17 days between last application and harvest in stone fruit crops. For olive groves, apply as directed sprays only.

### Restrictions on Application Equipment

For cherries, any application equipment listed in this section may be used in all states.

Any application equipment listed in this section may be used in apricots, nectarines, peaches and plums/prunes growing in Arizona, California, Colorado, Idaho, Kansas, Kentucky, New Jersey, North Dakota, Oklahoma, Oregon, Texas, Utah and Washington, except for peaches grown in states specified in the following paragraph. In all other states, use wiper equipment only.

For PEACHES grown in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, North Carolina, South Carolina and Tennessee only, apply with a shielded boom sprayer or shielded wiper applicator, which prevents any contact of Mirage Plus with the foliage or bark of trees. Apply no later than 90 days after first bloom. Applications made after this time may result in severe damage. Remove suckers and low-hanging limbs at least 10 days prior to application. Avoid applications near trees with recent pruning wounds or other mechanical injury. Apply only near trees that have been planted in the orchard for 2 or more years. EXTREME CARE MUST BE TAKEN TO ENSURE NO PART OF THE PEACH TREE IS CONTACTED.

### TREE NUTS

**Labeled Crops:** Almond, Beechnut, Betelnut, Brazil nut, butternut, Cashew, Chestnut, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia, Pecan, Pine nut, Pistachio, Walnut (black, English).

15/3

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

**Precautions, Restrictions:** Allow a minimum of 3 days between last application and harvest of tree nuts, except coconut. Allow 14 days between application and harvest in coconut.

### TROPICAL and SUBTROPICAL TREES and FRUITS

**Labeled Crops:** Ambarella, Atemoya, Avocado, Banana, Barbados cherry (acerola), Biriba, Blimbe, Breadfruit, Cacao (cocoa) bean, Canistel, Carambola (starfruit), Cherimoya, Coffee, Custard apple, Dates, Durian, Feijoa, Figs, Governor's plum, Guava, Ilima, Imbe, Imbu, Jaboticaba, Jackfruit, Longan, Lychee, Mamey apple, Mango, Mangosteen, Marmaladebox (genip), Mountain papaya, Papaya, Pawpaw, Plantain, Persimmon, Pomegranate, Pulasan, Rambutan, Rose apple, Sapodilla, Sapote (black, mamey, white), Spanish lime, Soursop, Star apple, Sugar apple, Sunnam cherry, Tamarind, Tea, Ti (roots and leaves), Wax jambu.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application" plus Bananacide (banana only).

**Precautions, Restrictions:** Allow a minimum of 1 day between last application and harvest in banana, guava, papaya, and plantain crops. Allow a minimum of 14 days between last application and harvest for any other tropical or subtropical tree fruit. Allow a minimum of 28 days between last application and harvest in coffee crops. In coffee and banana, delay applications 3 months after transplanting to allow the new coffee or banana plant to become established.

### Bananacide (Banana Only)

**Use Instructions:** Mirage Plus may be used to destroy banana plants infected with the Banana Bunchy Top Virus as well as non-infected banana plants to establish disease free buffers around plantations. Remove all fruit from the plants within the treatment area prior to treatment. Inject 0.04 fluid ounces (1 mL) of Mirage Plus's concentrate per 2 to 3 inches of pseudostem diameter. Make the injection at least one foot above the ground, except for very small plants, which should be injected vertically into the top. Any subsequent regrowth must also be destroyed. All plants and mats (or units) adjacent (within a 4-foot radius) to a treated mat shall be mechanically destroyed.

For control of the Banana Bunchy Top Virus, it is critical that the grower follow a strict control program involving monitoring for diseased plants, spraying to control the aphid vector, and destruction of all infected mats (or units). An infected plant may not show symptoms of the banana bunchy top virus for up to 125 days, therefore it is critical that the entire mat (or unit) containing the diseased plant be destroyed immediately.

**Precautions, Restrictions:** Do not apply more than 0.5 fluid ounce (15 mL) of Mirage Plus's concentrate per mat (or unit). Remove all fruit from plants and mats (or units) prior to treatment. Do not harvest any fruit or plant materials from treated mats (or units) following injection. Do not allow livestock to consume treated plant materials. Follow transplant of new banana plants into treated areas, allow plants to become established for 3 months before applying Mirage Plus for general weed control.

### VINE CROPS

**Labeled Crops:** Grapes (juice, raisin, table, wine), Hops, Kiwi, Passion fruit.

**Types Of Applications:** Those listed in "Tree, Vine and Shrub Crops (Alphabetical) Types of Application."

Applications should not be made when green shoots, canes or foliage are in the spray zone.

In the northeast and Great Lakes regions, applications must be made prior to the end of bloom stage of grapes to avoid injury, or make applications with shielded sprayers or wiper equipment.

**Precautions, Restrictions:** Allow a minimum of 14 days between last application and harvest in vine crops. Do not use selective equipment in kiwi.

### PASTURE GRASSES, FORAGE LEGUMES and RANGELANDS ALFALFA, CLOVER, AND OTHER FORAGE LEGUMES

**Labeled Crops:** Alfalfa, Clover, Kenaf, Kudzu, Lespedeza, Leucaena, Lupin, Sainfoin, Trefoil, Velvet bean, Vetch (all types).

**Types Of Applications:** Preplant, Preemergence, At-planting, Spot Treatment (Alfalfa and Clover Only), Over-the-Top Wiper Applicators (Alfalfa and Clover Only), Renovation, Preharvest (Alfalfa Only).

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied before, during or after planting crops listed in this section. Applications must be made prior to emergence of the crop.

**Precautions, Restrictions:** If a single application is made at rates of 2 quarts per acre or less, no waiting period between treatment and feeding or grazing is required. If application rates greater than 2 quarts per acre are made, remove domestic livestock before application and wait 8 weeks after application before grazing or harvest.

1/31

**PREHARVEST (ALFALFA ONLY)**

**Use Instructions:** Mirage Plus may be used in declining alfalfa stands or any stand of alfalfa where crop destruction is acceptable. This application will severely injure or destroy the stand of alfalfa. Mirage Plus will control annual and perennial weeds, including quackgrass, when applied prior to the harvest of alfalfa. The treated crop and weeds can be harvested and fed to livestock after 36 hours. Allow a minimum of 36 hours between application and harvest. Applications may be made at any time of the year. Make only one application to an existing stand of alfalfa per year. For control of quackgrass, apply in the spring, late summer or fall when quackgrass is actively growing. Treatments for quackgrass must be followed by deep tillage for complete control.

**Precautions, Restrictions:** Do not apply more than 2 quarts of Mirage Plus per acre as a preharvest treatment. Preharvest application is not recommended for alfalfa grown for seed, as a reduction in germination or vigor may occur.

**Spot Treatment, Over-the-Top Wiper Applications (Alfalfa and Clover Only)**

**Use Instructions:** Mirage Plus may be applied as a spot treatment in alfalfa or clover. Mirage Plus may be applied with wiper applicators to control or suppress the weeds listed under "Wiper Applicators" in the "Selective Equipment" section of this label. Applications may be made in the same area at 30-day intervals.

**Precautions, Restrictions:** For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total field area should be treated at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

**Renovation**

**Use Instructions:** Mirage Plus may be applied as a broadcast spray to renovate existing stands of alfalfa, clover, and other labeled forage legumes. Labeled crops may be planted into the treated area.

**Precautions, Restrictions:** Remove domestic livestock before application. If application rates of 2 quarts per acre or less are used wait 36 hours after application before grazing or harvesting. If application rates greater than 2 quarts per acre are used, wait 8 weeks between applications and grazing or harvesting.

**FOR USE IN DORMANT ALFALFA**

**Use Directions**

Mirage Plus will control or suppress many weeds, including quackgrass, downy brome and cheatgrass in dormant alfalfa.

Apply 8 to 12 fluid ounces per acre of Mirage Plus. Apply in the spring to alfalfa that is dormant. Applications should be made after spring temperatures have warmed enough to encourage resumption of weed growth, but prior to initiation of trifoliolate leaf expansion of the alfalfa. Applications made after expansion of the first trifoliolate leaf of the alfalfa will cause growth reduction and reduced crop yield.

Do not use ammonium sulfate when spraying dormant alfalfa with Mirage Plus.

Do not use Mirage Plus where a slight yield reduction in the first cutting of alfalfa cannot be tolerated.

Do not make more than one application per year.

Allow 36 hours after application before grazing livestock or harvesting.

Slight discoloration of the alfalfa may occur, but the alfalfa will regreen and regrow under moist conditions as effects of Mirage Plus wear off.

Application of Mirage Plus is limited to persons who have attended a Loveland Products, Inc. approved training program.

Application of Mirage Plus can cause crop injury. Any crop injury is the sole responsibility of the applicator.

**Conservation Reserve Program (CRP)**

**Types Of Applications:** Renovation (rotating out of CRP), Site preparation, Postemergence weed control in dormant CRP grasses, Wiper applications.

**Renovation (Rotating Out of CRP), Site Preparation**

**Use Instructions:** Mirage Plus may be used to prepare CRP land for crop production. Refer to Federal, state or local use guides for CRP renovation recommendations.

**Postemergence Weed Control in Dormant CRP Grasses, Wiper Applications**

**Use Instructions:** Mirage Plus may be used to suppress competitive growth and seed production of undesirable vegetation in CRP acres. Such applications may be made with wiper application equipment or as a broadcast or spot treatment to dormant CRP grasses. For selective applications with broadcast spray equipment, apply 12 to 16 fluid ounces of Mirage Plus per acre in early spring before desirable CRP grasses, such as crested and tall wheatgrass, break dormancy and initiate green growth. Late fall applications can be made after desirable perennial grasses have reached dormancy.

**Precautions, Restrictions:** Some stunting of CRP perennial grasses will occur if broadcast applications are made when plants are not dormant.

**NOTE:** MIRAGE PLUS MAY BE USED FOR CONTROL OF NON-MIRAGE PLUS TOLERANT ALFALFA IN PRODUCTION FIELDS OF ALFALFA CONTAINING THE MIRAGE PLUS TOLERANT GENE. SEVERE INJURY OR DEATH OF ALFALFA WILL RESULT IF ALFALFA VARIETIES THAT DO NOT CONTAIN THE MIRAGE PLUS TOLERANT GENE ARE SPRAYED WITH MIRAGE PLUS.

**Use Directions**

Mirage Plus will control non-glyphosate tolerant alfalfa in seed production fields of alfalfa containing the glyphosate tolerant gene. Apply up to 64 fluid ounces of Mirage Plus in 5 to 20 gallons of spray solution per acre as broadcast spray. Subsequent applications of up to 64 fluid ounces per acre each may be applied, if needed to control non-glyphosate tolerant alfalfa plants.

DO NOT EXCEED A MAXIMUM RATE OF 8 QUARTS OF MIRAGE PLUS PER ACRE PER SEASON.

**Application timing** - Mirage Plus can be applied to glyphosate tolerant alfalfa from emergence to harvest.

Treated alfalfa or the resulting seed may not be used for food or feed. Do not feed or graze treated alfalfa. Do not process treated alfalfa or resulting seed for food or feed.

**USE ONLY FOR SEED PRODUCTION OF LETTUCE WITH THE**  
**MIRAGE PLUS TOLERANT GENE**

**NOTE:** MIRAGE PLUS MAY BE USED FOR CONTROL OF NON-MIRAGE PLUS TOLERANT LETTUCE IN PRODUCTION FIELDS OF LETTUCE CONTAINING THE MIRAGE PLUS TOLERANT GENE. SEVERE INJURY OR DEATH OF LETTUCE WILL RESULT IF LETTUCE VARIETIES THAT DO NOT CONTAIN THE MIRAGE PLUS TOLERANT GENE ARE SPRAYED WITH MIRAGE PLUS.

**Use Directions**

Mirage Plus will control non-glyphosate tolerant lettuce in seed production fields of lettuce containing the glyphosate tolerant gene. Apply up to 64 fluid ounces of Mirage Plus in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 64 fluid ounces per acre may be applied, if needed to control non-glyphosate tolerant lettuce plants.

DO NOT EXCEED A MAXIMUM RATE OF 4 QUARTS OF MIRAGE PLUS PER ACRE PER SEASON.

**Application timing** - Mirage Plus can be applied to glyphosate tolerant lettuce from emergence to harvest.

Treated lettuce may not be used for food or feed. Do not feed or graze treated lettuce. Do not process treated lettuce for food or feed.

**USE ONLY FOR SEED PRODUCTION OF RICE WITH THE**  
**MIRAGE PLUS TOLERANT GENE**

**NOTE:** MIRAGE PLUS MAY BE USED FOR CONTROL OF NON-MIRAGE PLUS TOLERANT RICE IN PRODUCTION FIELDS OF RICE CONTAINING THE MIRAGE PLUS TOLERANT GENE. SEVERE INJURY OR DEATH WILL RESULT IF RICE VARIETIES THAT DO NOT CONTAIN THE MIRAGE PLUS TOLERANT GENE ARE SPRAYED WITH MIRAGE PLUS.

**Use Directions**

Mirage Plus will control non-glyphosate tolerant rice in seed production fields of rice containing the glyphosate tolerant gene. Apply up to 64 ounces of Mirage Plus in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 64 ounces per acre may be applied, if needed to control non-glyphosate tolerant rice plants.

DO NOT EXCEED A MAXIMUM RATE OF 4 QUARTS OF MIRAGE PLUS PER ACRE PER SEASON.

**Application timing** - Mirage Plus can be applied to glyphosate tolerant rice from emergence to harvest.

Treated rice may not be used for food or feed. Do not feed or graze treated rice. Do not process treated rice for food or feed.

**FOR USE ONLY FOR SEED PRODUCTION OF WHEAT WITH**  
**THE MIRAGE PLUS TOLERANT GENE**

**NOTE:** MIRAGE PLUS MAY BE USED FOR CONTROL OF NON-MIRAGE PLUS TOLERANT WHEAT IN PRODUCTION FIELDS OF WHEAT CONTAINING THE MIRAGE PLUS TOLERANT GENE. SEVERE INJURY OR DEATH WILL RESULT IF WHEAT VARIETIES THAT DO NOT CONTAIN THE GENE ARE SPRAYED WITH MIRAGE PLUS.

**Use Directions**

Mirage Plus will control non-glyphosate tolerant wheat in seed production fields of wheat containing the gene. Apply up to 32 fluid ounces of Mirage Plus in 5 to 20 gallons of spray solution per acre as a broadcast spray. A second application up to 32 fluid ounces per acre may be applied, if needed to control non-glyphosate tolerant wheat plants.

DO NOT EXCEED A MAXIMUM RATE OF 2 QUARTS OF MIRAGE PLUS PER ACRE PER SEASON.

**Application timing** - Mirage Plus can be applied to wheat from emergence to harvest.



# MIRAGE PLUS

## EPA REG. NO. 34704-

Treated wheat may not be used for food or feed. Do not feed or graze treated wheat. Do not process treated wheat for food or feed.

### GRASS SEED PRODUCTION

**Labeled Crops:** Any grass (*Gramineae* family) except corn, sorghum, sugarcane and those listed above under "Cereal Crops".

**Types Of Applications:** Preplant, Preemergence, Renovation, Site Preparation, Shielded Sprayers, Wiper Applications, Spot Treatments, Creating Rows in Annual Ryegrass.

#### Preplant, Preemergence, Renovation

**Use Instructions:** Mirage Plus may be applied before, during, or after planting or for renovation of turf or forage grass areas grown for seed production. Applications must be made prior to the emergence of the crop to avoid injury. For maximum control of existing vegetation, delay planting to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses, such as Bermudagrass, summer or fall applications provide best control.

**Precautions, Restrictions:** Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

Do not feed or graze treated areas for 8 weeks following application.

#### Shielded Sprayers

**Use Instructions:** Apply 1 to 3 quarts of Mirage Plus as a broadcast spray in 10 to 20 gallons of water per acre to control weeds in the rows. Uniform planting in straight rows aid in shielded sprayer applications. Best results are obtained when the grass seed crop is small enough to easily pass by or through the protective shields.

**Precautions, Restrictions:** Contact of Mirage Plus in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

#### Wiper Applications

**Precautions, Restrictions:** Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators should be adjusted so that the wiper contact point is at least 2 inches above the desirable vegetation. Weeds should be a minimum of 6 inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when height of weeds varies so that not all weeds are contacted. In these instances, repeat treatments may be necessary. Better results may be obtained if 2 applications are made in opposite directions.

#### Spot Treatments

**Use Instructions:** Use a 1 to 1.5 percent solution.

**Precautions, Restrictions:** Apply Mirage Plus prior to heading of grasses. Do not treat more than 10 percent of the total field area. The crop receiving the spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason.

#### Creating Rows in Annual Ryegrass

**Use Instructions:** Use 16 to 32 fluid ounces of Mirage Plus per acre. Use the higher rate when the ryegrass is greater than 6 inches tall. Best results are obtained when applications are made before the ryegrass reaches 6 inches in height.

**Precautions, Restrictions:** Set nozzle heights to allow the establishment of the desired row spacing while preventing spray droplets, spray fines, or drift to contact the ryegrass plants not treated. Use of low-pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended.

Grower assumes all responsibility for crop losses from misapplication.

### PASTURES

**Labeled Crops:** Any grass (*Gramineae* family) except corn, sorghum, sugarcane and those listed above under "Cereal Crops". Including Bahiagrass, Bermudagrass, Bluegrass, Bromegrass, Fescue, Guineagrass, Kikuyu grass, Orchardgrass, Pangola grass, Ryegrass, Timothy, Wheatgrass.

**Types of Applications:** Spot Treatment, Over-the-Top Wiper Application, Preplant, Preemergence, Pasture Renovation.

#### Spot Treatment, Over-the-Top Wiper Application

**Use Instructions:** Mirage Plus may be applied as a spot treatment or with wiper applications in pastures. Applications may be made in the same area at 30-day intervals.

**Precautions, Restrictions:** For spot treatment and wiper applications, apply in areas where the movement of domestic livestock can be controlled. No more than 10 percent of the total pasture area should be treated at one time. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

#### Preplant, Preemergence: Pasture Renovation

**Use Instructions:** Mirage Plus may be applied prior to planting or emergence of forage

grasses. In addition, Mirage Plus may be used to control perennial pasture species listed on this label prior to re-planting.

**Precautions, Preemergence:** Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

### FOR THE CONTROL OF ANNUAL WEEDS IN COASTAL BERMUDAGRASS PASTURES PRIOR TO SPRING GROWTH OR IMMEDIATELY AFTER FIRST CUTTING

#### Use Directions

Mirage Plus may be applied at 16 fluid ounces per acre to control the weeds listed below and most other winter annual grass and broadleaf weeds in established coastal Bermudagrass pastures.

Annual bluegrass	Oats
Cheat	Ryegrass, Italian
Crabgrass	Sandbur, field
Henbit	Sunflower
Johnsongrass, seedling	Wheat
Little barley	Wild mustard

#### Timing Of Application

**Applications prior to spring growth:** Apply Mirage Plus in the late winter or early spring but before new coastal Bermudagrass growth begins in the spring. Applications to new growth can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 60 days after making this application before grazing or harvesting the treated area.

**Applications following the first cutting:** Apply Mirage Plus after the first Bermudagrass cutting when the Bermudagrass has not yet begun to regrow. Applications made after regrowth has begun can damage the Bermudagrass.

Remove domestic livestock from the pasture before making the application. Wait 28 days after making this application before grazing or harvesting the treated area.

**NOTE:** ONLY ONE APPLICATION PER YEAR MAY BE MADE TO ANY ONE FIELD. A SPRING APPLICATION PRIOR TO GROWTH AND AN APPLICATION FOLLOWING THE FIRST CUTTING MAY NOT BE MADE ON THE FIELD DURING THE SAME YEAR.

#### Rangelands

**Types Of Applications:** Postemergence.

**Use Instructions:** Mirage Plus will control or suppress many annual weeds growing in perennial cool and warm-season grass rangelands.

Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds.

Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

#### Postemergence

Apply 12 to 16 fluid ounces of Mirage Plus to control or suppress many weeds, including downy brome, cheatgrass, cereal rye and jointed goatgrass in rangelands. Apply when most mature brome plants are in early flower and before the plants, including seedheads, turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve and encourages perennial grass conversion on weedy sites. Fall applications are possible, and recommended, where spring moisture is usually limited and fall germination allows for good weed control.

For medusahead, apply 16 fluid ounces of Mirage Plus at the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of Mirage Plus wear off.

**Precautions, Restrictions:** Do not use ammonium sulfate when spraying rangeland grasses with Mirage Plus. Do not make more than one application per year.

### TURF GRASS SOD PRODUCTION

**Types Of Applications:** Preplant, Preemergence, Renovation, Site Preparation, Spot Treatments.

#### Preplant, Preemergence, Renovation, Site Preparation

**Use Instructions:** Mirage Plus controls most existing vegetation prior to renovating turf grass areas or establishing turf grass grown for seed or sod. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. For warm-season grasses such as Bermudagrass, summer or fall applications provide the best control.

Desirable turf grasses may be planted following the above procedures.

# MIRAGE PLUS

## EPA REG. NO. 34704-

### Precautions, Restrictions:

Do not feed or graze turf grass grown for seed or sod production for 8 weeks following application.

### Spot Treatments

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turf grass.

### MIRAGE PLUS TOLERANT CROPS

The following instructions or those separately published on Loveland Products, Inc. Supplemental labeling include all applications which can be made onto the specified glyphosate tolerant crops during the complete cropping season. Do NOT combine these instructions with other recommendations made for crop varieties that do not contain the glyphosate tolerant gene, in the "Annual And Perennial Crops (Alphabetical)" section of this label.

LOVELAND PRODUCTS, INC. RECOMMENDS USE OF MIRAGE PLUS FOR POSTEMERGENCE APPLICATION ONLY ON CROP VARIETIES DESIGNATED AS CONTAINING THE MIRAGE PLUS TOLERANT GENE.

Applying Mirage Plus to crop varieties that are not designated as glyphosate tolerant will result in severe crop injury and yield loss. Avoid contact with foliage, green stems, or fruit of crops, or any desirable plants that do not contain the glyphosate tolerant gene, since severe injury or destruction will result.

The glyphosate tolerant designation indicates that the crop variety contains a patented gene that provides tolerance to Mirage Plus. Information on glyphosate tolerant crop varieties may be obtained from your seed supplier or Loveland Products, Inc. representative. Mirage Plus tolerant crop varieties must be purchased from an authorized licensed seed supplier.

**Note: Seed, and the method of selectively controlling weeds using Mirage Plus on a crop, are protected under several U.S. Patents, including 5,352,605 and 5,633,435. A license to use seed must be obtained prior to use.** Monsanto retains ownership of the gene and process technologies, and the Purchaser of the seed receives the right to use the licensed genes and technologies subject to the limited use license conditions. Seed containing the trait cannot be used for research and demonstration, reverse engineering or in connection with herbicide registration. Progeny seed containing the trait cannot be saved for replanting. Contact your Authorized Monsanto retailer for information on obtaining a limited use license.

For ground applications with broadcast equipment, apply Mirage Plus in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat spray nozzles. Check for even distribution of spray droplets.

For aerial applications, apply Mirage Plus in 3 to 15 gallons of water per acre. See the "Application Equipment And Techniques" section of this label for procedures to avoid spray drift that may cause injury to any vegetation not intended for treatment. Use of appropriate buffer zones will help prevent injury to adjacent vegetation.

For proper stewardship of aerial applications over-the-top of crops, Loveland Products recommends that growers and applicators read and follow all precautions and procedures contained in the use guide "A Guide to On-Target Aerial Application" available by calling 1-800-ROUNDUP (1-800-768-6387) or on the internet at [www.FARM-SOURCE.com](http://www.FARM-SOURCE.com).

**ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING MIRAGE PLUS TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS, WHICH DO NOT CONTAIN THE GENE.**

**See the "Mixing" and "Application Equipment And Techniques" sections of this label for additional directions and restrictions on the application of Mirage Plus.**

Tank mixtures with other herbicides, insecticides, fungicides, micronutrients or foliar fertilizers may result in reduced weed control or crop injury and are NOT recommended for over-the-top applications of Mirage Plus unless otherwise noted in the Mirage Plus label or fact sheets published separately by Loveland Products, Inc. Unless otherwise directed, nonionic surfactant may be added to the spray solution for applications to Mirage Plus tolerant crops. The addition of certain surfactants to Mirage Plus may result in some crop response including leaf necrosis, leaf chlorosis or leaf speckling due to the surfactant added to the spray mixture. Read and carefully observe cautionary statements and other information appearing on the surfactant label.

Ammonium sulfate may be mixed with Mirage Plus for applications to glyphosate tolerant crops. Refer to the "Mixing" section for use instructions for ammonium sulfate.

**Sprayer Preparation:** It is important that sprayer and mixing equipment be clean and free of pesticide residue before making applications of Mirage Plus. Follow the cleaning procedures specified on the label of the product(s) previously used. THOROUGHLY CLEAN THE SPRAY TANK AND ALL LINES AND FILTERS TO ELIMINATE POTENTIAL CONTAMINATION FROM OTHER HERBICIDES PRIOR TO MIXING AND APPLYING MIRAGE PLUS.

**NOTE:** The following recommendations are based on a clean start at planting by using a burndown application or tillage to control existing weeds before crop emergence. In no-till and stale seedbed systems, a preplant burndown treatment of Mirage Plus is recommended to control existing weeds prior to crop emergence. Some weeds, such as black nightshade, broadleaf signalgrass, sicklepod, Texas panicum, sandbur, annual morning

glory, woolly cupgrass, shattercane, wild proso millet, burcumber, and giant ragweed with multiple germination times or suppressed (stunted) weeds may require a second application of Mirage Plus for complete control. The second application should be made after some regrowth has occurred and at least 10 days after a previous application of Mirage Plus.

### CANOLA WITH THE MIRAGE PLUS TOLERANT GENE

**Types Of Applications:** Preplant, At-Planting, Preemergence, Postemergence

DO NOT USE MIRAGE PLUS ON CANOLA WITH THE MIRAGE PLUS TOLERANT GENE PLANTED IN THE FOLLOWING STATES: ALABAMA, DELAWARE, FLORIDA, GEORGIA, KENTUCKY, MARYLAND, NEW JERSEY, NORTH CAROLINA, SOUTH CAROLINA, TENNESSEE, VIRGINIA AND WEST VIRGINIA.

THE USE OF MIRAGE PLUS FOR IN-CROP APPLICATIONS OVER MIRAGE PLUS TOLERANT CANOLA MAY NOT BE PRACTICED IN CALIFORNIA UNLESS THE APPLICATOR HAS THE TIME OF APPLICATION A CALIFORNIA-APPROVED SUPPLEMENTAL LABEL SPECIFYING THE ACCEPTED DIRECTION FOR USE.

Maximum Allowable Combined Application Quantities Per Season	
Preplant, At-planting, Preemergence applications	2 quarts per acre
Total in-crop application from emergence to 6-leaf stage	2 quarts per acre

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied before, during or after planting canola.

### Postemergence

**Use Instructions:** Mirage Plus may be applied postemergence to canola from emergence through the 6-leaf stage of development. Applications made during bolting or flowering may result in crop injury and yield loss. To maximize yield potential, make applications early to eliminate competing weeds.

**Weeds Controlled:** For specific rates of application and instructions, refer to the "Annual Weeds" and "Perennial Weeds Rate Tables" in this booklet.

**Single Application -** Apply 16 to 32 fluid ounces per acre no later than the 6-leaf stage for the control of annual weeds. Avoid overlapping applications that may result in temporary yellowing, delayed flowering, and/or growth reduction. Similar injury may result when applications of more than 16 fluid ounces per acre are applied after the 4-leaf stage.

**Sequential Application -** Apply 32 fluid ounces per acre to 1- to 3-leaf canola followed by a sequential application at a minimum interval of 10 days, but no later than the 6-leaf stage. Sequential applications are recommended for early emerging annual weeds and perennial weeds such as Canada thistle and quackgrass or when controlling weeds with multiple application times.

**Precautions, Restrictions:** See the "Mirage Plus Tolerant Crops" section of this label for general precautionary instructions for use in crops. No more than two over-the-top broadcast applications may be made from crop emergence through the 6-leaf stage of development and the total in-crop application should not exceed 64 fluid ounces per acre. Allow a minimum of 60 days between last application and canola harvest.

### CORN WITH THE MIRAGE PLUS TOLERANT GENE

**Types Of Applications:** Preplant, At-Planting, Preemergence (In-Crop), Spot Treatment, Preharvest, Post-Harvest.

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence Applications	5 quarts per acre
Total in-crop applications from emergence through the V8 stage = or 30 inches	2 quarts per acre

Maximum preharvest application rate after maximum kernel fill is complete and the crop is physiologically mature (black layer formation) until 7 days before harvest 1 quart per acre

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied alone or in a tank mixture before, during or after planting corn.

**Tank Mixtures:** Mirage Plus may be tank mixed with Bullet, Degree, Degree Xtra, Harness, Harness Xtra, Harness Xtra 5.6L, Lariat, Lasso or Micro-Tech at 50 to 100 percent of labeled. Refer to the specific product label and observe all precautions and limitations on the label for any preemergence herbicide application, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines - the more restrictive requirements apply.

**NOTE:** For maximum weed control, a postemergence (in-crop) application of Mirage Plus should be applied following the use of less than labeled rates of the preemergence residual products listed above.

### Postemergence (In-Crop)

**Use Instructions:** Mirage Plus may be applied postemergence to glyphosate tolerant corn from emergence through the V8 stage (8 leaves with collars) or until corn height reaches 30 inches, whichever comes first.

When applied as directed, Mirage Plus controls labeled annual grass and broadleaf weeds in glyphosate tolerant corn. Many perennial grasses and broadleaf weeds will be controlled or suppressed with one or more application of Mirage Plus. The postemergent application of 24 to 32 fluid ounces per acre of Mirage Plus should be made before the

# MIRAGE PLUS

## EPA REG. NO. 34704-

weeds reach a height and/or density that the weeds become competitive with the crop, generally 4 inch tall weeds or less.

Mirage Plus may be applied alone as a postemergence in-crop application to provide control of emerged weeds listed on this label. If new flushes of weeds occur, a sequential application of Mirage Plus at 24 to 32 fluid ounces per acre will control the labeled grasses and broadleaf weeds.

**Tank Mixtures:** Mirage Plus may be applied in tank mixture with Bullet, Degree, Degree Xtra, Harness, Harness Xtra, Harness Xtra 5.6L, and Micro-Tech at 50 to 100 percent of labeled rate, provided the specific product listed below is registered for use on these sites. Mirage Plus may be applied in tank mixture with Permit and Atrazine at labeled rates. Refer to the specific product label and observe all precautions and limitations on the label for all products used in tank mixtures, including application timing restrictions, soil restrictions, minimum recropping interval and rotational guidelines - the more restrictive requirements apply.

Tank-mix Partner	Maximum Height Of Corn For Application
Degree	11 inches
Degree Xtra	
Harness	
Harness Xtra	
Harness Xtra 5.6	
Bullet*	5 inches
Micro-tech*	
Permit	30 inches
Atrazine	12 inches

\*Bullet and Micro-Tech are not registered for use as a postemergence application in Texas.

**Precautions, Restrictions:** See the "Mirage Plus Tolerant Crops" section of this label for general precautionary instructions for use in glyphosate tolerant crops. Single in-crop applications of Mirage Plus are not to exceed 1 quart per acre. Sequential in-crop applications of Mirage Plus from emergence through the V8 stage or 30 inches must not exceed 2 quarts per acre per growing season. Allow a minimum of 10 days between in-crop applications of Mirage Plus. Allow a minimum of 50 days between application of Mirage Plus and harvest of corn forage.

### Preharvest

**Use Instructions:** In corn, up to 1 quart per acre of Mirage Plus can be applied preharvest. Make applications at 35 percent grain moisture or less. Ensure that maximum kernel fill is complete and the corn is physiologically mature (black layer formed).

**Precautions, Restrictions:** Allow a minimum of 7 days between application and harvest, feeding, or grazing.

### Post-Harvest

**Use Instructions:** Mirage Plus may be applied after harvest of corn. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

**Precautions, Restrictions:** Allow a minimum of 7 days between treatment and harvest or feeding of treated vegetation.

## COTTON WITH THE MIRAGE PLUS TOLERANT GENE

**Types Of Applications:** Preplant, At-Planting, Preemergence, Over-the-Top, Selective Equipment, Preharvest

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Total in-crop applications from ground cracking to layby	4 quarts per acre
Maximum preharvest application rate	2 quarts per acre

**Precautions, Restrictions:** See the "Mirage Plus Tolerant Crops" section of this label for general precautionary instructions for use in crops. The combined total application of Mirage Plus from cotton emergence until harvest must not exceed 6 quarts per acre. NO MORE THAN TWO OVER-THE-TOP BROADCAST APPLICATIONS MAY BE MADE FROM CROP EMERGENCE THROUGH THE 4-LEAF (NODE) STAGE OF DEVELOPMENT. NO MORE THAN TWO APPLICATIONS SHOULD BE MADE FROM THE 5-LEAF STAGE THROUGH LAYBY. SEQUENTIAL IN-CROP OVER-THE-TOP OR POST-DIRECTED APPLICATIONS OF MIRAGE PLUS MUST BE AT LEAST 10 DAYS APART AND COTTON MUST HAVE AT LEAST TWO NODES OF INCREMENTAL GROWTH BETWEEN APPLICATIONS. ALLOW A MINIMUM OF 7 DAYS BETWEEN APPLICATION AND HARVEST.

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied before, during or after planting cotton.

### Over-the-Top

**Use Instructions:** Mirage Plus may be applied by aerial or ground application equipment at rates up to 1 quart per acre per application postemergence to glyphosate tolerant cotton from the ground cracking stage until the 4-leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the 4-leaf (node) stage of development may result in bill loss, delayed maturity and/or yield loss.

**Precautions, Restrictions:** The addition of surfactant to the spray solution may result in crop injury and reduced yield and is not recommended for over-the-top applications of Mirage Plus to glyphosate tolerant cotton.

17/31

**Salvage Treatment.** This treatment may be used after the 4-leaf stage of development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top application or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

**NOTE:** For specific rates of application and instructions, refer to the "Annual and Perennial Weeds Rate Tables" in this booklet.

**Precautions, Restrictions:** See the "Mirage Plus Tolerant Crops" section of this label for general precautionary instructions for use in glyphosate tolerant crops.

### Selective Equipment

**Use Instructions:** Mirage Plus may be applied using precision post-directed or hooded sprayers at rates up to 1 quart per acre per application to glyphosate tolerant cotton through layby. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 psi). For best results, make applications while weeds are small (less than 3 inches).

**Precautions, Restrictions:** See the "Selective Equipment" part of the "Application Equipment And Techniques" section of this label for information on proper use and calibration of this equipment.

### Preharvest

**Use Instructions:** Mirage Plus may be applied for preharvest annual and perennial weed control as a broadcast treatment to glyphosate tolerant cotton after 20 percent boll crack. Up to 2 quarts of Mirage Plus may be applied using either aerial or ground spray equipment. **NOTE:** Mirage Plus will not enhance the performance of harvest aids when applied to glyphosate tolerant cotton.

**Precautions, Restrictions:** Allow a minimum of 7 days between application and harvest of cotton. Do not apply Mirage Plus to cotton grown for seed, as a reduction in germination or vigor may occur. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF MIRAGE PLUS TO MIRAGE PLUS TOLERANT COTTON IS PROHIBITED.**

**ATTENTION: Use of Mirage Plus in accordance with label directions is expected to result in normal growth of glyphosate tolerant cotton, however, various environmental conditions, agronomic practices and other factors make it impossible to eliminate all risks associated with Mirage Plus, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.**

## SOYBEANS WITH THE MIRAGE PLUS TOLERANT GENE

**Types Of Applications:** Preplant, At-planting, Preemergence, Postemergence (In-Crop), Preharvest, Post-Harvest

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence Applications	5 quarts per acre
Total in-crop applications from cracking throughout flowering	3 quarts per acre
Maximum preharvest application	1 quart per acre

**Precautions, Restrictions:** See the "Mirage Plus Tolerant Crops" section of this label for general precautionary instructions for use in glyphosate tolerant crops.

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied before, during or after planting soybeans.

### Postemergence (In-Crop)

**Use Instructions:** When applied as directed, Mirage Plus will control labeled annual grasses and broadleaf weeds in Mirage Plus tolerant soybeans. Applications of Mirage Plus can be made in glyphosate tolerant soybeans from emergence (cracking) through-out flowering. Refer to the "Annual Weeds Rate Table" in this label for rate recommendation for specific annual weeds. In general, an initial application of 1 quart per acre on 2 to 8-inch tall weeds is recommended. Weeds will generally be 2 to 8 inches tall. 2 to 5 weeks after planting. If the initial application is delayed and weeds are larger, apply a higher rate of Mirage Plus. Mirage Plus may be used up to 2 quarts per acre in any single in-crop application for control of annual weeds and where heavy weed densities exist. A 1 to 2-quarts per acre rate (single or multiple applications) of Mirage Plus will control or suppress perennial weeds such as: Bermudagrass, Canada thistle, common milkweed, field bindweed, hemp dogbane, horsenettle, marestail (horseweed), nutsedge, quackgrass, rhizome johnson grass, redvine, trumpetcreeper, swamp smartweed and wirestem muhly. For best results, allow perennial weed species to achieve at least 6 inches of growth before spraying with Mirage Plus.

Under adverse growing conditions such as drought, hail, wind damage or poor soybean stand that slows or delays canopy closure, a sequential application of Mirage Plus may be necessary to control late flushes of weeds. IN THE SOUTHERN STATES, A SEQUENTIAL APPLICATION OF MIRAGE PLUS WILL BE REQUIRED TO CONTROL NEW FLUSHES OF WEEDS IN THE MIRAGE PLUS TOLERANT SOYBEAN CROP. To control giant ragweed, it is recommended that 1 quart per acre of Mirage Plus be

# MIRAGE PLUS

## EPA REG. NO. 34704-

applied when the weed is 8 to 12 inches tall to increase control and possibly avoid the need for a sequential application.

**NOTE:** The use of Mirage Plus for in-crop applications over glyphosate tolerant soybeans may not be practiced in California unless the applicator has, at the time of application, a California-approved Supplemental Label specifying the accepted Directions for Use.

**Precautions, Restrictions:** The combined total application from crop emergence through harvest must not exceed 3 quarts per acre. The maximum rate for any single in-crop application is 2 quarts per acre. The maximum combined total of Mirage Plus that can be applied during flowering is 2 quarts per acre.

### Preharvest

**Use Instructions:** Mirage Plus provides weed control when applied prior to harvest of soybeans. Up to 1 quart per acre of Mirage Plus can be applied by aerial or ground application.

**Precautions, Restrictions:** Care should be taken to avoid excessive seed shatter loss due to ground application equipment. Allow a minimum of 14 days between final application and harvest of soybean grain or feeding of soybean grain, forage or hay.

### Post-Harvest

**Use Instructions:** Mirage Plus may be applied after harvest of glyphosate tolerant soybeans. Higher rates may be required for control of large weeds that were growing in the crop at the time of harvest. Tank mixtures with 2,4-D or dicamba may be used.

## SUGAR BEETS WITH THE MIRAGE PLUS TOLERANT GENE

**Types Of Applications:** Preplant, At-Planting, Preemergence, Postemergence (In-Crop).

Maximum Allowable Combined Application Quantities Per Season	
Combined total per year for all applications	8 quarts per acre
Preplant, At-planting, Preemergence applications	5 quarts per acre
Emergence to 8-leaf stage	2.5 quarts per acre
Between 8-leaf stage and canopy closure	2 quarts per acre

**Precautions, Restrictions:** See the "Mirage Plus Tolerant Crops" section of this label for general precautionary instructions for use in glyphosate tolerant crops. The combined total application from crop emergence through harvest must not exceed 4.5 quarts per acre. The maximum rate for any single application from crop emergence until the 8-leaf stage is 1.5 quarts per acre. The maximum rate for any single application between the 8-leaf stage and canopy closure is 1 quart per acre. Allow a minimum of 30 days between last application and sugar beet harvest.

### Preplant, Preemergence, At-Planting

**Use Instructions:** Mirage Plus may be applied before, during or after planting of glyphosate tolerant sugar beets.

### Postemergence (In-Crop)

**Use Instructions:** Mirage Plus may be applied postemergent over-the-top to Mirage Plus tolerant sugar beets from emergence to 30 days prior to harvest. To maximize yield potential, spray sugar beets early to eliminate competing weeds. Up to 4 sequential applications of Mirage Plus may be made with at least 10 days between applications. Refer to the "Annual Weeds Rate Tables" in this label for rate recommendations for specific annual weeds. Mirage Plus will control or suppress most perennial weeds. For some perennial weeds, repeat applications may be required to eliminate crop competition throughout the growing season.

## NON-CROP USES AROUND THE FARMSTEAD

**Types Of Applications:** General Non-Selective Weed Control, Trim-and-Edge, Greenhouse/Shadehouse, Chemical Mowing, Cut Stumps, Habitat Management.

### General Weed Control, Trim-And-Edge

**Use Instructions:** Mirage Plus may be used to control annual weeds, perennial weeds and woody brush which are found in any part of the farmstead, including building foundations, along and in fences, in dry ditches and canals, along ditchbanks, farm roads, shelterbelts, prior to landscape planting and equipment storage areas.

**Tank Mixtures:** Mirage Plus may be tankmixed with the following products provided the specific product listed below is registered for use on these sites. Refer to these product labels for approved farmstead sites and application rates. For annual weeds, use 1 quart per acre of Mirage Plus when weeds are less than 6 inches tall, 1.5 quarts per acre when weeds are 6 to 12 inches tall and 2 quarts per acre when weeds are greater than 12 inches tall. For perennial weeds, apply 2 to 5 quarts per acre in these tank mixes. For tank mixtures with these products through backpack sprayers, handguns or other high-volume spray-to-wet applications, see the "Annual Weeds - Hand-Held Or High Volume Equipment" section of this label for recommended rates.

Arsenal	Krovar 1 DF	Ronstar 50 WP
Banvel/Clarity	Oust	Sahara
Barricade 65WG	Pendulum 3.3EC	Simazine
Diuron	Pendulum WDG	Surflan
Endurance	Plateau	Telar
Escort	Princep DF	Vanquish
Karmex DF	Princep Liquid	2,4-D

Mirage Plus plus dicamba tank mixtures may not be applied by air in California.

20/31

## GREENHOUSE/SHADEHOUSE

Mirage Plus may be used to control weeds in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

## CHEMICAL MOWING

**Use Instructions:** Mirage Plus will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 6 fluid ounces of Mirage Plus per acre when treating Kentucky bluegrass. Use 8 fluid ounces of Mirage Plus per acre when treating tall fescue, fine fescue, orchardgrass, bahiagrass or quackgrass covers. Use 16 fluid ounces of Mirage Plus per acre when treating Bermudagrass. Use 64 fluid ounces of Mirage Plus per acre when treating torpedograss or paragrass. Apply treatments in 10 to 20 gallons of spray solution per acre. Chemical mowing applications may be made along farm ditches and other parts of farmsteads.

**Precautions, Restrictions:** Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

## CUT STUMPS

**Types Of Applications:** Treating cut stumps in any non-crop site listed on this label.

**Use Instructions:** Mirage Plus will control regrowth of cut stumps and resprouts of many types of woody brush and tree species, some of which are listed below. Apply Mirage Plus using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of Mirage Plus to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, applications should be made during period of active growth and full leaf expansion.

Alder	Reed, giant
Eucalyptus	Saltcedar
Madrone	Sweetgum
Oak	Tan oak
Pepper, Brazilian	Willow
Pine, Austrian	

**Precautions, Restrictions:** Do not make cut stump applications when the roots of desirable woody brush or trees may be grafted to the roots of the cut stump. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

## HABITAT MANAGEMENT

**Types Of Uses:** Habitat restoration and maintenance, Wildlife food plots.

### Habitat Restoration and Maintenance

**Use Instructions:** Mirage Plus may be used to control exotic and other undesirable vegetation in habitat management areas. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot Treatments can be made to selectively remove unwanted plants for habitat maintenance and enhancement.

### Wildlife Food Plots

**Use Instructions:** Mirage Plus may be used as a site preparation treatment to control annual and perennial weeds prior to planting wildlife food plots. Any wildlife food species may be planted after applying Mirage Plus, or native species may be allowed to repopulate the area. If tillage is needed to prepare a seedbed, wait 7 days after application before tillage.

## ANNUAL WEEDS RATE TABLE (Alphabetically by Species)

WATER CARRIER VOLUMES OF 3 TO 10 GALLONS PER ACRE FOR GROUND APPLICATIONS AND 3 TO 5 GALLONS PER ACRE FOR AERIAL APPLICATIONS ARE RECOMMENDED.

Apply to actively growing annual weeds. Annual weeds are generally easiest to control when they are small.

Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

Do not tank mix with soil residual herbicides when using these rates unless otherwise specified.

For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Mirage Plus may be used up to 48 fluid ounces per acre where heavy weed densities exist.

**MIRAGE PLUS**  
EPA REG. NO. 34704-

21/131

Weed Species	Annual Weeds Rate Table			
	16	24	32	48
	Rate (fluid ounces per acre)			
	Maximum height/length (in inches)			
Ammannia, purple	3	6	12	18
Annoda, spurred	-	2	3	5
Barley	18	18+	-	-
Barryardgrass	-	3	6	7
Bassia, fivehook	-	-	6	-
Beggarweed, Florida	-	5	8	-
Bittercress	12	20	-	-
Bluegrass, annual	10	-	-	-
Bluegrass, bulbous	6	-	-	-
Brome, downy <sup>1,2</sup>	6	12	-	-
Brome, Japanese	6	12	24	-
Browntop panicum	6	8	12	24
Buckwheat, wild <sup>3</sup>	-	1	2	-
Burcucumber	-	6	12	18
Buttercup	12	20	-	-
Carolina geranium	-	-	4	9
Carpetweed	-	6	12	-
Cheat <sup>2</sup>	6	20	-	-
Chervil	20	-	-	-
Chickweed	-	12	18	-
Cocklebur	12	18	24	36
Copperleaf, hophornbeam	-	2	4	6
Copperleaf, Virginia	-	2	4	6
Coreopsis, plains	-	6	12	18
Corn, volunteer	6	12	20	-
Corn speedwell	12	-	-	-
Crabgrass	3	6	12	-
Crowfootgrass	-	-	6	12
Cutleaf evening primrose	-	-	3	6
Devilsclaw (unicorn plant)	-	3	6	-
Dwarf dandelion	12	-	-	-
Eastern manna grass	8	12	-	-
Eclipta	-	4	8	12
Fall panicum	4	-	6	12
Falsedandelion	-	20	-	-
Falseflax, small seed	12	-	-	-
Fiddleneck	-	6	12	-
Field pennycress	6	12	-	-
Filaree	-	-	6	12
Flabane, annual	6	20	-	-
Flabane, hairy ( <i>Coryza bonariensis</i> )	-	-	6	10
Flabane, rough	3	6	12	-
Florida pusley	-	-	4	6
Foxtail, giant, bristly, yellow	6	12	20	-
Foxtail, Carolina	10	-	-	-
Foxtail, green	12	-	-	-
Goatgrass, jointed	6	12	-	-
Goosegrass	-	3	6	12
Grain sorghum (milo)	6	12	20	-
Groundcherry	-	3	6	9
Groundsel, common	-	6	10	-
Hemp sesbania	-	2	4	6
Henbit	-	-	6	12
Horseweed/Marestail ( <i>Coryza canadensis</i> )	-	6	12	18
Itchgrass	6	8	12	18
Jimsonweed	-	-	12	18
Johnsongrass, seedling	6	12	18	24
Junglerice	-	3	6	7
Knotweed	-	-	6	12
Kochia <sup>4</sup>	-	3 to 6	12	-
Lambsquarters	-	6	12	20
Little barley	6	12	-	-
London rocket	6	-	24	-
Mayweed	-	2	6	12
Morningglory, annual ( <i>Ipomoea</i> spp.)	-	-	3	6
Mustard, blue	6	12	18	-
Mustard, tansy	6	12	18	-
Mustard, tumble	6	12	18	-
Mustard, wild	6	12	18	-
Nightshade, black	-	4	6	12
Nightshade, hairy	-	4	6	12
Oats	3	6	18	-
Pigweed species	-	12	18	24
Prickly lettuce	-	6	12	-
Purslane	-	-	3	6
Ragweed, common	-	6	12	18
Ragweed, giant	-	6	12	18
Red rice	-	-	4	-
Rye, volunteer/cereal <sup>2</sup>	6	18	18+	-
Ryegrass	-	-	6	12
Sandbur, field	6	12	-	-

Weed Species	Annual Weeds Rate Table			
	16	24	32	48
	Rate (fluid ounces per acre)			
	Maximum height/length (in inches)			
Sandbur, longspine	6	12	-	-
Shattercane	6	12	20	-
Shepherd's-purse	6	12	-	-
Sicklepod	-	2	4	8
Signalgrass, broadleaf	-	3	6	7
Smartweed, ladythumb	-	-	6	9
Smartweed, Pennsylvania	-	-	6	9
Sowthistle, annual	-	-	6	12
Spanishneedles	-	-	6	12
Speedwell, purslane	12	-	-	-
Sprangletop	6	12	20	-
Spurge, prostrate	-	6	12	-
Spurge, spotted	-	6	12	-
Spurry, umbrella	6	-	-	-
Stinkgrass	-	12	-	-
Sunflower	12	18	-	-
Swinecress	-	5	12	-
Teaweed/Prickly sida	-	2	4	6
Texas panicum	6	8	12	-
Thistle, Russian <sup>5</sup>	-	6	12	24
Velvetleaf	-	-	6	12
Virginia pepperweed	-	18	-	-
Waterhemp	-	-	6	12
Wheat <sup>2</sup>	6	12	18	-
Wheat, (overwintered)	-	6	12	18
Wild oats	3	6	18	-
Wild proso millet	-	6	12	18
Witchgrass	-	12	-	-
Woolly cupgrass	-	6	12	-
Yellow rocket	-	12	20	-

- <sup>1</sup> For control of downy brome in no-till systems, use 24 fluid ounces per acre.
- <sup>2</sup> Performance is better if application is made before this weed reaches the boot stage of growth.
- <sup>3</sup> Use 24 fluid ounces per acre of Mirage Plus to control wild buckwheat in the cotyledon to 2-leaf stage. Use 32 fluid ounces per acre to control 2 to 4-leaf wild buckwheat. For improved control of wild buckwheat over 2 inches in size, use sequential treatments of 32 fluid ounces followed by 32 fluid ounces of Mirage Plus per acre.
- <sup>4</sup> Do not treat kochia in the button stage.
- <sup>5</sup> Control of Russian thistle may vary based on environmental conditions and spray coverage. Whenever possible, tank mixture with 2, 4-D as described below may improve control.

**ANNUAL WEEDS - RATES FOR 10 TO 40 GALLONS PER ACRE**  
Apply 1 to 2 quarts of Mirage Plus per acre. Use 1 quart per acre if weeds are less than 6 inches tall, 1.5 quarts per acre if weeds are 6 to 12 inches tall and 2 quarts per acre if weeds are greater than 12 inches tall. These rates will provide control of weeds listed in the annual weed control tables when water carrier volumes are 10 to 40 gallons per acre for ground applications. Older, mature (hardened) annual weed species may require higher rates even if they meet the size requirements.

**ANNUAL WEEDS - TANK MIXTURES WITH 2,4-D, DICAMBA OR TORDON 22K**

12 to 16 fluid ounces of Mirage Plus plus 0.25 pound of dicamba or 0.5 pound of 2, 4-D or 1 to 2 fluid ounces of Tordon 22K per acre will control the following weeds with the maximum height or length indicated: 6 inches - prickly lettuce, mare's tail/horseweed, morning glory, kochia (dicamba only) wild buckwheat (Tordon 22K only): 12 inches - cocklebur, lambsquarters, pigweed, Russian thistle (2,4-D only).

16 fluid ounces of Mirage Plus plus 0.5 pound of 2,4-D per acre, will control the following weeds when they are at maximum height or length of 6 inches: common ragweed, giant ragweed, Pennsylvania smartweed, and velvetleaf.

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if dicamba or Tordon 22K is applied within 45 days of planting.

DO NOT APPLY DICAMBA TANK MIXTURES BY AIR IN CALIFORNIA

**ANNUAL WEEDS - HAND-HELD OR HIGH-VOLUME EQUIPMENT**

For control of weeds listed in the "Annual Weeds Rate Tables", apply a 0.5 percent solution of Mirage Plus to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution.

For best results, use a 2 percent solution on harder-to-control perennials, such as Bermudagrass, dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

When using application methods that result in less than complete coverage, use a 5 percent solution for annual and perennial weeds and a 5 to 10 percent solution for woody brush and trees.

**MIRAGE PLUS**  
EPA REG. NO. 34704-

**ANNUAL WEEDS - TANK MIXTURES WITH ATRAZINE FOR FALLOW AND REDUCED TILLAGE SYSTEMS**

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota and Washington. In Oregon and Washington, do not exceed 1 pound of atrazine per acre.

24 to 28 fluid ounces of Mirage Plus plus 1 to 2 pounds of atrazine per acre will control the following weeds: Barnyardgrass (requires 28 ounces for control), Downy brome, Green foxtail, Lambsquarters, Prickly lettuce, Tansy mustard, Pigweed, Field sandbur, Stinkgrass, Russian thistle, Volunteer wheat, Witchgrass and Kochia (add 1/8 pound of dicamba for control).

**PERENNIAL WEEDS RATE TABLE (Alphabetically by Species)**

Apply to actively growing perennial weeds.

**NOTE:** If weeds have been mowed or tilled, do not treat until plants have resumed active growth and have reached the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

Unless otherwise stated, allow 7 or more days after application before tillage.

Best results are obtained when soil moisture is adequate for active weed growth.

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Alfalfa	1-2	3-10	2%

Make applications after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but before soil freeze-up.

Alligatorweed	4	3-20	1.5%
---------------	---	------	------

Partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain control.

Anise (fennel)	-	-	1-2%
----------------	---	---	------

For hand-held, apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth.

Bahiagrass	3-5	3-20	2%
------------	-----	------	----

Apply when most plants have reached the early head stage.

Bentgrass	1.5	10-20	2%
-----------	-----	-------	----

For suppression in grass seed production areas. For ground applications only. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results.

Bermudagrass	3-5	3-20	2%
--------------	-----	------	----

For control, apply 5 quarts of Mirage Plus per acre. For partial control, apply 3 quarts per acre. Treat when Bermudagrass is actively growing and seedheads are present. Retreatment may be necessary to maintain control.

Bermudagrass, water (knotgrass)	1-1.5	5-10	2%
---------------------------------	-------	------	----

Apply 1.5 quarts of Mirage Plus in 5 to 10 gallons of water per acre. Apply when water Bermudagrass is 12 to 18 inches in length. Allow 7 more days before tilling, flushing or flooding the field.

Fall applications only: Apply 1 quart of Mirage Plus in 5 to 10 gallons of water per acre. Fallow fields should be tilled prior to application. Apply prior to frost on water Bermudagrass that is 12 to 18 inches in length.

Mirage Plus is not registered in California for use on water Bermudagrass.

Blindweed, field	0.5-5	3-20	2%
------------------	-------	------	----

Do not treat when weeds are under drought stress as good soil moisture is necessary for active growth.

For control, apply 4 to 5 quarts of Mirage Plus per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when the weeds are at or beyond full bloom. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Also for control, apply 2 quarts of Mirage Plus plus 0.5 pound of dicamba in 10 to 20 gallons of water per acre. Do not apply by air.

For suppression on irrigated agricultural land, apply 1 to 2 quarts of Mirage Plus plus 1 pound 2,4-D in 10 to 20 gallons of water per acre with ground equipment only. Applications should be made following harvest or in fall fallow ground when the bindweed is actively growing and the majority of runners are 12 inches or more in length. The use of at least one irrigation will promote active bindweed growth.

For suppression, apply 16 fluid ounces of Mirage Plus plus 0.5 pound of 2, 4-D in 3 to 10 gallons of water per acre for ground application and 3 to 5 gallons of water per acre for aerial applications. Apply by air in fallow and reduced tillage systems only.

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
--------------	-------------	--------------------	----------------------

Blindweed, field cont'd.:

Applications should be delayed until maximum emergence has occurred and when vines are between 6 to 18 inches in length.

**In California Only,** apply 1 to 5 quarts of Mirage Plus per acre. Actual rate needed for suppression or control will vary within this range depending on local conditions. For suppression on irrigated land where annual tillage is performed, apply 1 quart of Mirage Plus in 3 to 10 gallons of water per acre. Apply to bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Allow 3 or more days after application before tillage.

Bluegrass, Kentucky	1-2	3-40	2%
---------------------	-----	------	----

Apply 2 quarts of Mirage Plus in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Mirage Plus in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Blueweed, Texas	3-5	3-40	2%
-----------------	-----	------	----

Apply 4 to 5 quarts of Mirage Plus per acre west of the Mississippi River and 3 to 4 quarts east of the Mississippi River. Apply when plants are at or beyond full bloom. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost.

Brackenfern	3-4	3-40	1-1.5%
-------------	-----	------	--------

Apply to fully expanded fronds that are at least 18 inches long.

Bromegrass, smooth	1-2	3-40	2%
--------------------	-----	------	----

Apply 2 quarts of Mirage Plus in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Mirage Plus in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Bursage, woolly-leaf	-	3-20	2%
----------------------	---	------	----

For control, apply 2 quarts of Mirage Plus plus 1/2 pound of dicamba per acre. For partial control, apply 1 quart of Mirage Plus plus 1/2 pound of dicamba per acre. Apply when plants are producing new, active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering.

Canarygrass, reed	2-3	3-40	2%
-------------------	-----	------	----

For best results, apply when most plants have reached the boot-to-head stage of growth.

Cattail	3-5	3-40	2%
---------	-----	------	----

Apply when most plants have reached the early head stage.

Clover; red or white	3-5	3-20	2%
----------------------	-----	------	----

Apply when most plants have reached the early bud stage.

Also for control, apply 16 to 32 fluid ounces of Mirage Plus plus 0.5 to 1 pound of 2,4-D in 3 to 10 gallons of water per acre.

Cogongrass	3-5	10-40	2%
------------	-----	-------	----

Apply when cogongrass is at least 18 inches tall in late summer or fall. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Dallisgrass	3-5	3-20	2%
-------------	-----	------	----

Apply when most plants have reached the early head stage.

Dandelion	3-5	3-40	2%
-----------	-----	------	----

Apply when most plants have reached the early bud stage of growth.

Also for control, apply 16 fluid ounces of Mirage Plus plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre.

Dock, curly	3-5	3-40	2%
-------------	-----	------	----

Apply when most plants have reached the early bud stage of growth.

Also for control, apply 16 to 32 fluid ounces of Mirage Plus plus 0.5 to 1 pound of 2,4-D in 3 to 10 gallons of water per acre.

Dogbane, hemp	4	3-40	2%
---------------	---	------	----

Apply when most plants have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. For suppression, apply 16 fluid ounces of Mirage Plus plus 0.5 pound of 2, 4-D in 3 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Delay application until maximum emergence of dogbane has occurred.

Fescue (except tall)	3-5	3-20	2%
----------------------	-----	------	----

Apply when most plants have reached the early head stage.

Fescue, tall	1-3	3-40	2%
--------------	-----	------	----

Apply 3 quarts of Mirage Plus per acre when most plants have reached boot-to-early seedhead stage of development.

Fall applications only: Apply 1 quart of Mirage Plus in 3 to 10 gallons of water per acre. Apply to fescue in the fall when plants have 6 to 12 inches of new growth. A sequential application of 1 pint per acre of Mirage Plus will improve long-term control and control seedling germinating after fall treatments or the following spring.

22/31



**MIRAGE PLUS**  
EPA REG. NO. 34704-

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Guineagrass	2-3	3-40	1%

Apply when most plants have reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. In Texas and ridge of Florida, use 2 quarts for control. In the flatwoods region of Florida, 3 quarts is required for control.

**Horsenettle** 3-5 3-20 2%

Apply when most plants have reached the early bud stage.

**Horseradish** 4 3-40 2%

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

**Iceplant** 1.5-2%

Iceplant should be at or beyond the early bud stage of growth. Thorough coverage is necessary for best control.

**Jerusalem artichoke** 3-5 3-20 2%

Apply when most plants have reached the early bud stage.

**Johnsongrass** 0.5-3 3-40 1%

In annual cropping systems apply 1 to 2 quarts of Mirage Plus per acre. Apply 1 quart of Mirage Plus in 3 to 10 gallons of water per acre. Use 2 quarts of Mirage Plus when applying 10 to 40 gallons of water per acre. In non-crop, or areas where annual tillage is not practiced (no-till), apply 2 to 3 quarts of Mirage Plus in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using 1 quart of Mirage Plus per acre.

For burndown of Johnsongrass, apply 1 pint of Mirage Plus 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.

Spot treatment (partial control or suppression) - Apply a 1 percent solution of Mirage Plus when Johnsongrass is 12 to 18 inches in height. Coverage should be uniform and complete.

**Kikuyugrass** 2-3 3-40 2%

Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth). Allow 3 or more days after application before tillage.

**Knapweed** 4 3-40 2%

Apply when most plants have reached the late bud to flower stage of growth. For best results, apply in late summer or fall.

**Lantana** 1-1.25%

Apply at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.

**Lespedeza** 3-5 3-20 2%

Apply when most plants have reached the early bud stage.

**Milkweed, common** 3 3-40 2%

Apply when most plants have reached the late bud to flower stage of growth.

**Muhly, wirestem** 1-2 3-40 2%

Use 1 quart of Mirage Plus in 3 to 10 gallons of water per acre. Use 2 quarts of Mirage Plus when applying 10 to 40 gallons of water per acre or in pasture, sod, or non-crop areas. Spray when the wirestem muhly is 8 inches or more in height. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 more days after application before tillage.

**Mullein, common** 3-5 3-20 2%

Apply when most plants are in the early bud stage.

**Napiergrass** 3-5 3-20 2%

Apply when most plants are in the early head stage.

**Nightshade, Silverleaf** 2 3-10 2%

Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost.

**Nutsedge, purple or yellow** 0.5-3 3-40 1.2%

Apply 3 quarts of Mirage Plus per acre or apply a 1 to 2 percent solution for control of nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets that have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control of ungerminated tubers.

Sequential applications: 1 to 2 quarts of Mirage Plus in 3 to 10 gallons of water per acre will also provide control. Make applications when a majority of the plants are in the 3- to 5-leaf stage (less than 6 inches tall). Repeat this application, as necessary, when newly emerging plants reach the 3- to 5-leaf stage. Subsequent applications will be necessary for long-term control.

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
<i>Nutsedge, purple or yellow cont'd.</i>			
Orchardgrass	1-2	3-40	2%

For partial control of existing plants, apply 1 pint to 2 quarts of Mirage Plus in 3 to 40 gallons of water per acre. Treat when plants have 3 to 5 leaves and most are less than 6 inches tall. Repeat treatments will be required to control subsequent emerging plants or regrowth of existing plants.

**Orchardgrass** 1-2 3-40 2%

Apply 2 quarts of Mirage Plus in 10 to 40 gallons of water per acre when most plants have reached boot-to-early seedhead stage of development. For partial control in pasture or hay crop renovation, apply 1 to 1.5 quarts of Mirage Plus in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12 inches in height.

Orchardgrass sods going to no-till corn: Apply 1 to 1.5 quarts of Mirage Plus in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least 3 days following application before planting. A sequential application of atrazine will be necessary for optimum results.

**Pampasgrass** 1.5-2%

Pampasgrass should be at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

**Paragrass** 3-5 3-20 2%

Apply when most plants are in the early bud stage.

**Phragmites** 3-5 10-40 1-2%

For partial control and for best results, treat during late summer or fall when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stage of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

**Poison hemlock** 1-2%

For hand-held, apply as a spray-to-wet treatment. Optimum results are obtained when plants are treated at the bud to full-bloom stage of growth. Thorough coverage is necessary for best control.

**Pokeweed, common** 1 3-40 2%

Apply to actively growing plants up to 24 inches tall.

**Quackgrass** 1-3 3-40 2%

In annual cropping systems, or in pastures and sods followed by deep tillage: Apply 1 quart of Mirage Plus in 3 to 10 gallons of water per acre. For 10 to 40 gallons of water per acre, apply 2 quarts of Mirage Plus. Do not tank mix with residual herbicides when using the 1-quart rate. Spray when quackgrass is 6 to 8 inches in height. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or sods, use a moldboard plow for best results.

In pastures, sods or non-crop areas where deep tillage does not follow application: Apply 2 to 3 quarts of Mirage Plus in 10 to 40 gallons of water per acre when the quackgrass is greater than 8 inches tall.

**Redvine** 0.75-2 5-10 2%

For suppression, apply 24 fluid ounces of Mirage Plus per acre at each of two applications 7 to 14 days apart or a single application of 2 quarts per acre. Apply recommended rates in 5 to 10 gallons of water per acre. Apply in late September or early October to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

**Reed, giant** 2%

Best results are obtained when applications are made in late summer to fall.

**Ryegrass, perennial** 1-3 3-40 1%

In annual cropping systems apply 1 to 2 quarts of Mirage Plus per acre. Apply 1 quart of Mirage Plus in 3 to 10 gallons of water per acre. Use 2 quarts of Mirage Plus when applying 10 to 40 gallons of water per acre. In non-crop, or areas where annual tillage is not practiced (no-till), apply 2 to 3 quarts of Mirage Plus in 10 to 40 gallons of water per acre.

For best results, apply when most plants have reached the boot-to-head stage of growth or in the fall prior to frost. Do not tank-mix with residual herbicides when using 1 quart of Mirage Plus per acre.

**Smartweed, swamp** 3-5 3-40 2%

Apply when most plants have reached the early bud stage of growth. Also for control, apply 16 fluid ounces of Mirage Plus plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall.

**Sowthistle, perennial** 2-3 3-40 2%

Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of Mirage Plus. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

**Spurge, leafy** 3-10 2%

For suppression, apply 16 fluid ounces of Mirage Plus plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre in the late summer or fall. If mowing has occurred prior to treatment, apply when most of the plants are 12 inches tall.

28/31

**MIRAGE PLUS**  
EPA REG. NO. 34704-

Weed Species	Rate (QT/A)	Water Volume (GPA)	Hand-Held % Solution
Starthistle, yellow	2	10-40	2%
Best results are obtained when applications are made during the rosette, bolting and early flower stage.			
Sweet potato, wild			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, artichoke			2%
For partial control, apply to plants that are at or beyond the bloom stage of growth. Repeat applications may be required.			
Thistle, Canada	2-3	3-40	2%

Apply when most plants are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall, allow at least 4 weeks for initiation of active growth and rosette development prior to the application of Mirage Plus. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

For suppression in the spring, apply 1 quart of Mirage Plus, or 1 pint of Mirage Plus plus 0.5 pound of 2,4-D in 3 to 10 gallons of water per acre. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.

Timothy	2-3	3-40	2%
For best results, apply when most plants have reached the boot-to-head stage of growth.			
Torpedograss	4-5	3-40	2%

For partial control, apply when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost.

Trumpet creeper	2	5-10	2%
-----------------	---	------	----

For partial control, apply in late September or October, to plants that are at least 18 inches tall and have been growing 45 to 60 days since the last tillage operation. Make applications at least 1 week before a killing frost.

Vaseygrass	3-5	3-20	2%
------------	-----	------	----

Apply when most plants are in the early head stage.

Velvetgrass	3-5	3-20	2%
-------------	-----	------	----

Apply when most plants are in the early head stage.

Wheatgrass, western	2-3	3-40	2%
---------------------	-----	------	----

For best results, apply when most plants have reached the boot-to-head stage of growth.

**WOODY BRUSH AND TREES RATE TABLE**

Apply Mirage Plus after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Unless otherwise directed, apply broadcast treatments in 3 to 40 gallons of water per acre. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Weed Species	Rate (QT/A)	Hand-Held % Solution
*Partial Control		
Alder	3-4	1-1.5%
Ash*	2-5	1-2%
Aspen, quaking	2-3	1-1.5%
Bearmat (Bearclover)*	2-5	1-2%
Beech*	2-5	1-2%
Birch	2-3	1-1.5%
Blackberry	3-4	1-1.5%

Make applications after plants have reached full leaf maturity. Best results are obtained when applications are made in late summer or fall. Applications may also be made after leaf drop and until a killing frost or as long as stems are green. After berries have set or dropped in late fall, blackberry can be controlled by applying a 0.75 percent solution of Mirage Plus. For control of blackberries after leaf drop and until killing frost or as long as stems are green, apply 3 to 4 quarts of Mirage Plus in 10 to 40 gallons of water per acre.

Blackgum	2-5	1-2%
Bracken	2-5	1-2%
Broom; French, Scotch	2-5	1.5-2%

Weed Species	Rate (QT/A)	Hand-Held % Solution
*Partial Control		
Buckwheat, California*	2-4	1-2%
Thorough coverage of foliage is necessary for best results.		
Casara*	2-5	1-2%
Catsclaw*	-	1-1.5%
Ceanothus*	2-5	1-2%
Chamise	-	1%
Thorough coverage of foliage is necessary for best results.		
Cherry; bitter, black, pin	2-3	1-1.5%
Coyote brush	3-5	1.5-2%
Apply when at least 50 percent of the new leaves are fully developed.		
Dogwood*	2-5	1-2%
Elderberry	2-3	1-1.5%
Elm*	2-5	1-2%
Eucalyptus	-	2%
For control of eucalyptus resprouts, apply when resprouts are 6 to 12 feet tall. Ensure complete coverage. Avoid application to drought-stressed plants.		
Florida holly		
(Brazilian Peppertree)*	2-5	1-2%
Gorse*	2-5	1-2%
Hasardia*	2-4	1-2%
Thorough coverage of foliage is necessary for best results.		
Hawthorn	2-3	1-1.5%
Hazel	2-3	1-1.5%
Hickory*	2-5	1-2%
Honeysuckle	3-4	1-1.5%
Hornbeam American*	2-5	1-2%
Kudzu	4-5	2%
Repeat applications may be required to maintain control.		
Locust, black*	2-4	1-2%
Madrone resprouts*	-	2%
Apply to resprouts that are 3 to 6 feet tall. Best results are obtained with spring/early summer treatments.		
Manzanita*	2-5	1-2%
Maple, red	2-4	1-1.5%
Apply a 1 to 1.5 percent solution when at least 50 percent of the new leaves are fully developed. For partial control, apply 2 to 4 quarts of Mirage Plus per acre.		
Maple, sugar	-	1-1.5%
Apply when at least 50 percent of the new leaves are fully developed.		
Monkey flower*	2-4	1-2%
Thorough coverage of foliage is necessary for best results.		
Oak; black, white*	2-4	1-2%
Oak, post	3-4	1-1.5%
Oak; northern	-	1-1.5%
Apply when at least 50 percent of the new pin leaves are fully developed.		
Oak; southern Red	2-3	1-1.5%
Persimmon*	2-5	1-2%
Pine	2-5	1-2%
Poison ivy/Poison oak	4-5	2%
Repeat application may be required to maintain control. Fall treatments must be applied before leaves lose green color.		
Poplar, yellow*	2-5	1-2%
Reobud, eastern	2-5	1-2%
Rose, multiflora	2	1%
Treatments should be made prior to leaf deterioration by leaf-eating insects.		
Russian olive*	2-5	1-2%
Sage, black	2-4	1%
Thorough coverage of foliage is necessary for best results.		
Sage, white*	2-5	1-2%
Sagebrush, California	2-4	1%
Thorough coverage of foliage is necessary for best results.		
Salmonberry	2-3	1-1.5%
Saltcedar	2-5	1-2%
Sassafras*	2-5	1-2%
Sourwood*	2-5	1-2%
Sumac; poison, smooth, winged*	2-4	1-2%
Sweetgum	2-3	1-1.5%
Swordfern*	2-5	1-2%
Tallowtree, Chinese	-	1%
Thorough coverage of foliage is necessary for best results.		
Tan oak resprouts*	-	2%
Apply to resprouts that are less than 3 to 6 feet tall. Best results are obtained with fall applications.		
Thimbleberry	2-3	1-1.5%
Tobacco, tree*	2-4	1-2%
Trumpet creeper	2-3	1-1.5%
Vine maple*	2-5	1-2%
Virginia creeper	2-5	1-2%
Waxmyrtle, southern*	2-5	1-2%
Willow	3-4	1-1.5%

**INDUSTRIAL, TURF AND ORNAMENTAL USE RECOMMENDATIONS**

Detailed instructions follow alphabetically, by site.

Unless otherwise specified, applications may be made to control any weeds listed in the annual, perennial and woody brush tables. Refer also to the "Selective Equipment" section.

24/31



# MIRAGE PLUS

## EPA REG. NO. 34704-

### CUT STUMPS

Cut stump treatments may be made on any site listed on this label. Mirage Plus will control many types of woody brush and tree species. Apply Mirage Plus using suitable equipment to ensure coverage of the entire cambium. Cut trees or resprouts close to the soil surface. Apply a 50 to 100 percent solution of Mirage Plus to the freshly cut surface **immediately after cutting**. Delays in application may result in reduced performance. For best results, applications should be made during periods of active growth and full leaf expansions.

DO NOT MAKE CUT STUMP APPLICATIONS WHEN THE ROOTS OF DESIRABLE WOODY BRUSH OR TREES MAY BE GRAFTED TO THE ROOTS OF THE CUT STUMP. Some sprouts, stems, or trees may share the same root system. Adjacent trees having a similar age, height and spacing may signal shared roots. Whether grafted or shared, injury is likely to occur to non-treated stems/trees when one or more trees sharing common roots are treated.

### FORESTRY SITE PREPARATION

Mirage Plus is recommended for the control or partial control of woody brush, trees and herbaceous weeds in forestry. Mirage Plus is also recommended for use in preparing or establishing wildlife openings within these sites and maintaining logging roads.

Mirage Plus is recommended for use in site preparation prior to planting any tree species, including Christmas trees, eucalyptus, hybrid tree cultivars and silvicultural nursery sites.

For applications using different types of equipment, see "Application Rates" table in "Hand-Held Equipment" section of this label.

**Tank Mixtures:** Tank mixtures of Mirage Plus may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture.

**NOTE:** For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions.

Any recommended rate of Mirage Plus may be used in a tank mix with the following products for forestry site preparation.

Product	Broadcast Rate
Arsenal Applicators Concentrate	2 to 16 fluid ounces per acre
Escort™	0.5 to 3.5 ounces per acre
Chopper™	4 to 32 fluid ounces per acre
Garlon 4	1 to 4 quarts per acre
Oust™	1 to 4 ounces per acre

Product	Spray-To-Wet Rates
Arsenal Applicators Concentrate	0.03 to 0.5 percent by volume

Product	Low Volume Directed Spray Rates
Arsenal Applicators Concentrate	0.1 to 0.5 percent by volume

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

**Do not apply Mirage Plus as an over-the-top broadcast spray for forestry conifer or hardwood release.**

### GENERAL NON-CROP AREAS AND INDUSTRIAL SITES

Use in areas such as airports, apartment complexes, Christmas tree farms, ditch banks, dry ditches, dry canals, fencerows, golf courses, industrial sites, lumber yards, manufacturing sites, office complexes, ornamental nurseries, parks, parking areas, petroleum tank farms and pumping installations, railroads, recreational areas, residential areas, roadsides, sod or turf seed farms, schools, storage areas, substations, warehouse areas, other public areas, and similar industrial and non-crop sites.

#### General Weed Control, Trim-and-Edge, Bare Ground

Mirage Plus may be used in general non-crop areas. It may be applied with any application equipment described in this label. Mirage Plus may be used to trim-and-edge around objects in non-crop sites, for spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. Mirage Plus may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

Repeated applications of Mirage Plus may be used, as weeds emerge, to maintain bare ground.

**Tank Mixtures:** Mirage Plus may be tankmixed with the following products provided the specific product listed below is registered for use on these sites. Refer to these products' labels for approved non-crop sites and application rates.

Arsenal™	Karmex™ DF	Ronstar™ 50 WP
Clarity	Krovar™ 1 DF	Sahara™
Barricade™ 65WG	Mangø	Simazine
Diuron	Oust	Spike 80DF
Endurance™	Pendulum™ 3.3 EC	Surflan™
Escort™	Pendulum WDG	Telar™
Garlon™ 3A	Plateau™	Vanquish™
Garlon 4	Princep™ DF	2,4-D
Hyvar X	Princep™ Liquid	

Mirage Plus plus dicamba tank mixtures may not be applied by air in California.

25/31

### Brush Control Tank Mixtures

Tank mixtures of Mirage Plus may be used to increase the spectrum of control for herbaceous weeds, woody brush and trees. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of Mirage Plus may be used in a tank mix.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

**NOTE:** For side trimming treatments, it is recommended that Mirage Plus be used alone or in tank mixture with Garlon 4.

Product	Broadcast Rate
Arsenal 2WSL	6 to 32 fluid ounces per acre
Escort	1 to 2 ounces per acre
Garlon 3A*, Garlon 4	1 to 4 quarts per acre

Product	Spray-To-Wet Rates
Arsenal 2WSL	0.06 to 0.12% by volume
Escort	1 to 2 ounces per acre

Product	Low Volume Directed Spray Rates
Arsenal 2WSL	0.1 to 0.5% by volume
Escort	1 to 2 ounces per acre

\*Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding Mirage Plus. Have spray mixture agitating at the time Mirage Plus is added to avoid spray compatibility problems.

### Chemical Mowing - Perennials

Mirage Plus will suppress perennial grasses listed in this section to serve as a substitute for mowing. Use 8 fluid ounces of Mirage Plus per acre when treating tall fescue, fine fescue, orchardgrass, quackgrass or reed canarygrass covers. Use 6 fluid ounces of Mirage Plus per acre when treating Kentucky bluegrass. Apply treatments in 10 to 40 gallons of spray solution per acre.

Use only in areas where some temporary injury or discoloration of perennial grasses can be tolerated.

### Chemical Mowing - Annuals

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 4 to 5 fluid ounces of Mirage Plus in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments may cause injury to the desired grasses.

### Bromus Species and Medusahead in Pastures and Rangelands

**Bromus species.** Mirage Plus may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial, rangeland and pasture sites. Apply 8 to 16 fluid ounces of Mirage Plus per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses can become reestablished on the site.

**Medusahead.** To treat medusahead, apply 16 fluid ounces of Mirage Plus per acre as soon as plants are actively growing, and prior to the 4-leaf stage. Applications may be made in the fall or spring.

Applications to brome and medusahead may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre. When applied as directed in this label, there are no grazing restrictions.

### Dormant Turfgrass

Mirage Plus may be used to control or suppress many winter annual weeds and tall fescue for effective release of dormant Bermudagrass and bahiagrass turf. Treat only when turf is dormant and prior to spring greenup.

Apply 8 to 64 fluid ounces of Mirage Plus per acre. Apply the recommended rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated.

Treatments in excess of 16 fluid ounces of Mirage Plus per acre may result in injury or delayed greenup in highly maintained areas such as golf courses and lawns. DO NOT apply tank mixtures of Mirage Plus plus Oust or Outrider in highly maintained turfgrass areas. For further uses, refer to the "Roadsides" section of this label, which gives rates for dormant Bermudagrass and bahiagrass treatments.

### Actively Growing Bermudagrass

Mirage Plus may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. DO NOT apply more than 16 fluid ounces of Mirage Plus per acre in highly maintained turfgrass areas. DO NOT

# MIRAGE PLUS

## EPA REG. NO. 34704-

apply tank mixtures of Mirage Plus plus Oust in highly maintained turfgrass areas. For further uses, refer to the "Roadsides" section of this label, which gives rates for actively growing Bermudagrass treatments. Use only in areas where some temporary injury or discoloration can be tolerated.

### Turfgrass Renovation, Seed, or Sod Production

Mirage Plus controls most existing vegetation prior to renovating turfgrass areas or establishing turfgrass grown for seed or sod. For maximum control of existing vegetation, delay planting or sodding to determine if any regrowth from escaped underground plant parts occurs. Where repeat treatments are necessary, sufficient regrowth must be attained prior to application. For warm-season grasses such as Bermudagrass, summer or fall application provide the best control. Where existing vegetation is growing under mowed turfgrass management, apply Mirage Plus after omitting at least on regular mowing to allow sufficient growth for good interception of spray.

Do not disturb soil or underground plant parts before treatment. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow translocation into underground plant parts.

Desirable turfgrasses may be planted following the above procedures.

Hand-held equipment may be used for spot treatment of unwanted vegetation growing in existing turfgrass. Broadcast or hand-held equipment may be used to control sod remnants or other unwanted vegetation after sod is harvested.

Do not feed or graze turfgrass grown for seed or sod production for 8 weeks following application.

### FOR USE FOR SELECTIVE WEED CONTROL ON MIRAGE PLUS TOLERANT PURE GOLD® TALL FESCUE AND AURORA GOLD® FINE FESCUE SELECTIONS.

#### Use Directions

##### Mirage Plus Tolerant Tall Fescue Selections For Seed Production

Use Mirage Plus on Pure Gold tolerant tall and Aurora Gold fine fescue grown for seed production only.

Mirage Plus may be applied at rates of 4 to 16 fluid ounces per acre as a postemergence spray on glyphosate tolerant tall fescue selections. See the label booklet for application instructions, rate recommendations, weeds controlled and proper growth stage of weeds.

When applied postemergence, Mirage Plus will control or suppress the following weeds: annual bluegrass mustards, downy brome, cheatgrass, chickweed, penny-cress, fleabane, shepherd's-purse, sowthistle, wild oat, dandelion, quackgrass, and Canada thistle. See the Mirage Plus label booklet for a complete list of weeds controlled or suppressed.

**NOTE:** The recommended rate for this use will limit the level of control of certain species of weeds.

**NOTE:** Some crop discoloration and yellowing may occur at higher rates of application with glyphosate tolerant tall and fine fescue selections. Reduction in stand of these selections may occur under stress conditions.

#### Timing Of Applications

Applications can be made 6 weeks after germination and to established crops after growth resumes in the Fall until onset of dormancy and in the Spring after dormancy break until 60 days prior to harvest.

Avoid spraying during or within two weeks after periods when air temperatures fall below 25°F.

Remove domestic livestock from the seed production field prior to application. Wait 60 days after making this application before grazing or harvesting the treated area.

**NOTE:** Only two applications per crop growth cycle may be made to any one site. If two applications are required, only on fall and spring application may be made during on 12-month cycle.

### HABITAT MANAGEMENT

#### Habitat Restoration and Management

Mirage Plus may be used to control exotic and other undesirable vegetation in habitat management and natural areas, including rangeland and wildlife refuges. Applications can be made to allow recovery of native plant species, prior to planting desirable native species, and for similar broad-spectrum vegetation control requirements. Spot treatments can be made to selectively remove unwanted plants for habitat management and enhancement.

#### Wildlife Food Plots

Mirage Plus may be used as a site preparation treatment prior to planting wildlife food plots. Any wildlife food species may be planted after applying Mirage Plus, or native species may be allowed to repopulate the area. If tillage is needed to prepare seedbed, wait 7 days after application before tillage to allow translocation into underground plant parts.

#### Injection and Frill (Woody Brush and Trees)

Mirage Plus may be used to control woody brush and trees by injection or frill applications. Apply Mirage Plus using suitable equipment that must penetrate into the living tissue. Apply the equivalent of 0.04 fluid ounce (1 mL) of Mirage Plus per each 2 to 3 inch-

es of trunk diameter at breast height (DBH). This is best achieved by applying a 50 to 100 percent concentration of Mirage Plus either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frilled or cut areas in species that exude sap freely. In species such as this, make the frill or cuts at an oblique angle to produce a cupping effect and use a 100 percent concentration of Mirage Plus. For best results, application should be made during periods of active growth and after full leaf expansion.

### FOR GROUND AND AERIAL APPLICATIONS TO BRUSH AND CHAPARRAL IN CALIFORNIA ONLY.

#### Use Directions

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

Nonionic surfactants which are labeled for use with herbicides may be used to improve wetting of foliage. Do not reduce rates of Mirage Plus when adding surfactant. Read and carefully observe surfactant rates, cautionary statements, and other information appearing on the surfactant label.

**Timing Of Application:** Apply Mirage Plus as a broadcast spray when plants are actively growing for partial control of undesirable vegetation listed on this label. Best results are obtained when application is made in the spring to early summer when brush species are at a high moisture content and flowering.

Mirage Plus may be used as recommended for:

- Aid to burning treatment to establish and maintain fuel breaks
- Establishing fire perimeters and black lines
- Aid to prescribed burning
- Along fire roads and rights-of-way
- Range conversion
- Site preparation in forestry

**Application Recommendations:** Apply 2 quarts of Mirage Plus per acre for partial control of the following emerged brush and chaparral species:

<i>Ceanothus</i>	Salge
<i>Ceanothus</i> spp.	<i>Salvia</i> spp.
Chamise	Scrub oak
<i>Adenostoma fasciculatum</i>	<i>Quercus dumosa</i>

Ground applications should be applied in 3 to 40 gallons of total spray solution per acre.

Aerial applications (helicopter only) should be applied in 3 to 15 gallons of total spray solution per acre.

Avoid direct application to any body of water.

**AVOID DRIFT - DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION WHICH FAVORS DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

For aerial application of Mirage Plus, please see the label directions for aerial application in California.

### ORNAMENTALS, PLANT NURSERIES, AND CHRISTMAS TREES

#### Post-Directed, Trim-and-Edge

Mirage Plus may be used as a post-directed spray around established woody ornamental species such as arborvitae, azalea, boxwood, crabapple, eucalyptus, euonymus, fir, Douglas fir, jojoba, hollies, lilac, magnolia, maple, oak, poplar, privet, pine, spruce and yew. Mirage Plus may also be used to trim and edge around trees, buildings, sidewalks and roads, potted plants and other objects in a nursery setting.

Desirable plants may be protected from the spray solution by using shields or covering made of cardboard or other impermeable material. UNLESS OTHERWISE DIRECTED, MIRAGE PLUS IS NOT RECOMMENDED FOR USE AS AN OVER-THE-TOP BROADCAST SPRAY IN ORNAMENTALS AND CHRISTMAS TREES. Care must be exercised to avoid contact of spray, drift or mist with foliage or green bark of established ornamental species.

#### Site Preparation

Mirage Plus may be used prior to planting any ornamental, nursery or Christmas tree species.

#### Wiper Applications

Mirage Plus may be used through wick or other suitable wiper applicators to control or partially control undesirable vegetation around established eucalyptus or poplar trees. See the "Selective Equipment" section of this label for further information about the proper use of wiper applicators.

### MIRAGE PLUS HERBICIDE FOR CONIFER RELEASE

#### Use Directions

##### Aerial Application

Mirage Plus may be applied using aerial spray equipment for conifer release treatments. See the "Application Equipment and Techniques" part of the "Mixing, Additives and Application Instructions" section of the label for Mirage Plus herbicide for information on how to properly spray Mirage Plus by air.

# MIRAGE PLUS

## EPA REG. NO. 34704-

DO NOT APPLY MIRAGE PLUS BY AIR TO RIGHTS-OF-WAY SITES IN THE STATE OF CALIFORNIA.

### Conifer Release

For release, apply only where conifers have been established for more than one year. Vegetation should not be disturbed prior to treatment or until visual symptoms appear after treatment. Symptoms of treatment are slow to appear, especially in woody species treated in late Fall. Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied or when applications are made during periods of active conifer growth. **Do not use additional surfactant with conifer release applications.**

Applications must be made after formation of final conifer resting buds in the Fall or prior to initial bud swelling in spring. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Use the following rates for conifer release to control or partially control the weeds listed in the "Weeds Controlled" section of the label.

### For release of the following conifer species:

Douglas Fir	<i>Pseudotsuga menziesii</i>
Fir	<i>Abies</i> spp.
Hemlock	<i>Tsuga</i> spp.
Pines*	<i>Pinus</i> spp.
Spruce	<i>Picea</i> spp.

\*Includes all species except eastern white pine, loblolly pine or slash pine.

Apply 1.5 to 2 quarts of Mirage Plus per acre except in Washington and Oregon, west of the crest of the Cascade Mountains. For spring treatments west of the crest of the Cascade Mountains, apply 1 quart of Mirage Plus per acre before conifer bud swell for control of annual weeds. For Fall treatments in Washington and Oregon, west of the crest of the Cascade Mountains, apply 1 to 1.5 quarts of Mirage Plus per acre before any major leaf drop of deciduous species.

For release of western hemlock apply 1 quart of Mirage Plus per acre.

### For release of the following conifer species:

Loblolly pine	<i>Pinus taeda</i>
Eastern white pine	<i>Pinus strobes</i>
Slash pine	<i>Pinus elliotii</i>

**Late Season Application** - Apply 1.5 to 2 quarts of Mirage Plus in a minimum of 5 gallons of spray solution per acre during early autumn. Applications made prior to September 1 or when conditions are conducive to rapid growth of conifers will result in potential for increased injury in the form of tip and/or needle burn. Injury may decrease with larger application. Some autumn colors are acceptable at time of application. Apply prior to frost or leaf drop of undesirable plants. Applications made according to label direction will release loblolly pine, eastern white pine and slash pine by reducing competition from the following species:

Ash	<i>Fraxinus</i> spp.
Cherry, black	<i>Prunus serotina</i>
Cherry, pin	<i>Prunus pensylvanica</i>
Elm	<i>Ulmus</i> spp.
Hawthorn	<i>Crataegus</i> spp.
Locust, black	<i>Robinia pseudoacacia</i>
Maple, red	<i>Acer rubra</i>
Oak, black	<i>Quercus velutina</i>
Oak, post	<i>Quercus stellata</i>
Oak, southern red	<i>Quercus falcata</i>
Oak, white	<i>Quercus alba</i>
Persimmon	<i>Diospyros</i> spp.
Poplar, yellow	<i>Liriodendron tulipifera</i>
Sassafras	<i>Sassafras albidum</i>
Sourwood	<i>Oxydendrum arboretum</i>
Sumac, poison	<i>Rhus vernix</i>
Sumac, smooth	<i>Rhus glabra</i>
Sumac, winged	<i>Rhus copallina</i>
Sweetgum	<i>Liquidambar styraciflua</i>

Apply only to those sites where wood brush and trees listed in this label constitute the majority of the undesirable species.

### Mirage Plus Plus Oust™ Tank Mixtures For Conifer Release From Herbaceous Weeds

To release loblolly pines from herbaceous weeds, tank mixtures of Mirage Plus with Oust will provide control of annual weeds listed in the "Weeds Controlled" section of the label booklet for Mirage Plus and the Oust label, and partial control of the perennial weeds listed below.

Apply 16 to 24 fluid ounces of Mirage Plus with 2 to 4 ounces of Oust in 10 to 30 gallons of spray solution per acre. Make application to actively growing weeds as a broadcast spray over the top to the young loblolly pines.

Mirage Plus plus Oust tank mixtures may not be applied by air in California.

This tank mixture may be applied using aerial equipment. When applying by air, use the recommended rate in 5 to 15 gallons of spray solution per acre.

For control of annual weeds below 12 inches in height (or runner length on annual vines), use the low rates of both products. Use the higher rates of both products when annual weeds are in more advanced stages of growth and approaching flower or seed formation.

29/31

Use the higher rates of both products for partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass	<i>Paspalum notatum</i>
Broomsedge	<i>Andropogon virginicus</i>
Dock, curly	<i>Rumex crispus</i>
Dogfennel	<i>Eupatorium capilliflorum</i>
Fescue, tall	<i>Festuca arundinacea</i>
Johnsongrass**	<i>Sorghum halepense</i>
Poorjoe**	<i>Diodia teres</i>
Trumpetcreeper*	<i>Campsis radicans</i>
Vaseygrass	<i>Paspalum urvillei</i>
Vervain, blue	<i>Verbena hastata</i>

\*Suppression at the higher rates only.

\*\*Control at the higher rates.

Pine damage may occur or can be accentuated if treatment takes place when young trees are under stress from drought, flood, water, insects or disease.

## BROADCAST APPLICATIONS FOR WEED CONTROL IN CHRISTMAS TREE PLANTATIONS IN THE STATES OF WASHINGTON AND OREGON ONLY

### Use Directions

**NOTE: IF IMPROPERLY APPLIED, MIRAGE PLUS HAS THE POTENTIAL TO CAUSE SEVERE CHRISTMAS TREE INJURY. FOLLOW ALL LABELED DIRECTIONS.**

Mirage Plus may be applied as a broadcast spray over established Christmas trees. Ensure that adequate buffers are maintained to prevent drift onto nearby desirable crops or vegetation. Read the entire "Application Equipment And Techniques" section of the label for additional application precautions.

### This application is approved for the following Christmas tree species:

Douglas fir	( <i>Pseudotsuga menziesii</i> )
Fir species	( <i>Abies</i> spp.)
Spruce species	( <i>Picea</i> spp.)

Applications may be made only after trees have completed at least a full growing season since planting or transplanting. Applications should not be made within 1 full year prior to tree harvest.

Applications may only be made in the fall after the formation of final conifer resting buds. Final resting buds must be fully hardened and in the dormant stage. Applications made at any other time may result in unacceptable Christmas tree injury.

Avoid spray pattern overlap, as injury may occur.

Apply 1 quart of Mirage Plus per acre in 5 to 30 gallons of water per acre.

**NOTE: DO NOT ADD SURFACTANTS, ADDITIVES CONTAINING SURFACTANTS, OR ANY OTHER ADDITIVES TO MIRAGE PLUS OR SEVERE CHRISTMAS TREE INJURY MAY RESULT.**

Mirage Plus may be used at rates from 1 to 2 quarts per acre in some areas. Consult your local Loveland Products, Inc. representative or Mirage Plus supplier for specific recommendations if you require rates greater than 1 quart per acre.

Drift control additives may increase Christmas tree injury and their use is not recommended.

The use of other herbicides tank mixed with Mirage Plus is not recommended since severe Christmas tree injury may result.

### Greenhouse/Shadehouse

Mirage Plus may be used to control weeds growing in and around greenhouses and shadehouses. Desirable vegetation must not be present during application and air circulation fans must be turned off.

## PARKS, RECREATIONAL AND RESIDENTIAL AREAS

Mirage Plus may be used in parks, recreational and residential areas. It may be applied with any application equipment described in this label. Mirage Plus may be used to trim-and-edge around trees, fences, and paths, around buildings, sidewalks, and other objects in these areas. Mirage Plus may be used to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. Mirage Plus may be used prior to planting an area to ornamentals, flowers, turfgrass (sod or seed), or prior to laying asphalt or beginning construction projects.

All of the instructions in the "General Non-Crop Areas And Industrial Sites" section apply to park and recreational areas.

# MIRAGE PLUS

EPA REG. NO. 34704-

28/31

## RAILROADS

All of the instructions in the "General Non-Crop Areas And Industrial Sites" section apply to railroads.

### Bare ground, Ballast and Shoulders, Crossings, Spot Treatments

Mirage Plus may be used to maintain bare ground on railroad ballast and shoulders. Repeat applications of Mirage Plus may be used, as weeds emerge to maintain bare ground. Mirage Plus may be used to control tall growing weeds to improve line-of-sight at railroad crossings and reduce the need for mowing along rights-of-way. For crossing applications, up to 80 gallons of spray solution per acre may be used.

**Tank Mixtures:** Mirage Plus may be tank mixed with the following products for ballast, shoulder, spot, bare ground and crossing treatments provided the specific product listed below is registered for use on these sites:

Arsenal	Garlon 4	Spike™
Clarity	Hyvar™ X	Telar
Diuron	Krovar I DF	Vanquish
Escort	Oust	2,4-D
Garlon 3A	Sahara	

### Brush Control

Mirage Plus may be used to control woody brush and trees on railroad rights-of-way. Apply 4 to 10 quarts of Mirage Plus per acre as a broadcast spray, using boom-type or boomless nozzles. Up to 80 gallons of spray solution per acre may be used. Apply a 0.75 to 2 percent solution of Mirage Plus when using high-volume spray-to-wet applications. Apply a 5 to 10 percent solution of Mirage Plus when using low volume directed sprays for spot treatment.

**Tank Mixtures:** Mirage Plus may be mixed with the following products for enhanced control of woody brush and trees:

Arsenal	Garlon 4
Escort	Tordon™ K
Garlon 3A	

### Bermudagrass Release

Mirage Plus may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of Mirage Plus in up to 80 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Fescue, tall	Trumpet creeper
Bluestem, silver	Johnsongrass	Vaseygrass

**Tank Mixtures:** Mirage Plus may be tank mixed with Oust. If tank mixed, use no more than 1 to 3 pints of Mirage Plus with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Dewberry	Poorjoe
Blackberry	Dock, curly	Raspberry
Bluestem, silver	Dogfennel	Trumpet creeper
Broomsedge	Fescue, tall	Vaseygrass
Dallisgrass	Johnsongrass	Vervain, blue

Use only on well-established Bermudagrass. Bermudagrass injury may result from the treatment, but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may occur.

## ROADSIDES

All of the instructions in the "General Non-Crop Areas And Industrial Sites" section apply to roadsides.

**Tank Mixtures:** Mirage Plus may be tank mixed with the following products for shoulder, guardrail, spot and bare ground treatments provided the specific product listed below is registered for use on these sites:

Clarity	Pendulum 3.3EC	Simazine
Diuron	Pendulum WDG	Surflan
Endurance	Princep DF	Telar
Escort	Princep Liquid	Vanquish
Krovar I DF	Ronstar 50WP	2,4-D
Oust	Sahara	

See the "General Non-Crop Areas And Industrial Sites" section of this label for general instructions for tank mixing.

### Shoulder Treatments

Mirage Plus may be used on road shoulders. It may be applied with boom sprayers, shielded boom sprayers, high-volume off-center nozzles, hand-held equipment, and similar equipment.

### Guardrails and Other Obstacles to Mowing

Mirage Plus may be used to control weeds growing under guardrails and around signposts and other objects along the roadside.

### Spot Treatment

Mirage Plus may be used as a spot treatment to control unwanted vegetation growing along roadsides.

### Release of Bermudagrass or Bahiagrass

#### Dormant Applications

Mirage Plus may be used to control or partially control many winter annual weeds and tall fescue for effective release of dormant Bermudagrass or bahiagrass. Treat only when turf is dormant and prior to spring greenup. Mirage Plus may also be tank-mixed with Outrider or Oust for residual control. Tank mixtures of Mirage Plus with Oust may delay greenup.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is at or beyond the 4- to 6-leaf stage.

Apply 8 to 64 fluid ounces of Mirage Plus in a tank mixture with 0.75 to 1.3 ounces Outrider herbicide per acre. Read and follow all label directions for Outrider herbicide.

Apply 8 to 64 fluid ounces of Mirage Plus per acre alone or in a tank mixture with 0.25 to 1 ounce per acre of Oust. Apply the recommended rates in 10 to 40 gallons of water per acre. Use only in areas where Bermudagrass or bahiagrass are desirable ground covers and where some temporary injury or discoloration can be tolerated. To avoid delays in greenup and minimize injury, add no more than 1 ounce of Oust per acre on Bermudagrass and no more than 0.5 ounce of Oust per acre on bahiagrass and avoid treatments when these grasses are in a semi-dormant condition.

#### Actively Growing Bermudagrass

Mirage Plus may be used to control or partially control many annual and perennial weeds for effective release of actively growing Bermudagrass. Apply 1 to 3 pints of Mirage Plus in 10 to 40 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or runner length). Use the higher rate as weeds increase in size or as they approach flower or seedhead formation. These rates will also provide partial control of the following perennial species:

Bahiagrass	Johnsongrass
Bluestem, silver	Trumpet creeper
Fescue, tall	Vaseygrass

**Tank Mixtures:** Mirage Plus may be tank mixed with Outrider for control or partial control of Johnsongrass and other weeds listed in the Outrider label. Use 8 to 32 fluid ounces of Mirage Plus with 0.75 to 1.3 ounces of Outrider. Use the higher rates of both products for control of perennial weeds or annual weeds greater than 6 inches in height.

Mirage Plus may be tank mixed with Oust. If tank-mixed, use no more than 1 to 2 pints of Mirage Plus with 1 to 2 ounces of Oust per acre. Use the lower rates of each product to control annual weeds less than 6 inches in height (or runner length) that are listed in this label and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages. These rates will also provide partial control of the following perennial weeds:

Bahiagrass	Dock, curly	Poorjoe
Bluestem, silver	Dogfennel	Trumpet creeper
Broomsedge	Fescue, tall	Vaseygrass
Dallisgrass	Johnsongrass	Vervain, blue

Use only on well-established Bermudagrass. Bermudagrass injury may result from treatment, but regrowth will occur under moist conditions. Repeat applications of the tank mix in the same season are not recommended, since severe injury may occur.

#### Actively Growing Bahiagrass

For suppression of vegetative growth and seedhead inhibition of bahiagrass for approximately 45 days, apply 6 fluid ounces of Mirage Plus in 10 to 40 gallons of water per acre. Apply 1 to 2 weeks after full greenup or after mowing to a uniform height of 3 to 4 inches. This application must be made prior to seedhead emergence.

For suppression up to 120 days, apply 4 fluid ounces of Mirage Plus per acre, followed by an application of 2 to 4 fluid ounces per acre about 45 days later. Make no more than 2 applications per year.

**Tank Mixtures:** Mirage Plus may be used for control or partial control of Johnson grass and other weeds listed on the Outrider label in actively growing bahiagrass. Apply 1.5 to 5 fluid ounces of Mirage Plus with 0.75 to 1.3 ounces of Outrider per acre. Use the higher rates for control of perennial weeds or annual weeds greater than 6 inches in height. Use only on well established bahiagrass.

A tank mixture of Mirage Plus plus Oust may be used. Apply 6 fluid ounces of Mirage Plus plus 0.25 ounce of Oust per acre 1 to 2 weeks following an initial spring mowing. Make only one application per year.

## UTILITY SITES

In utilities, Mirage Plus is recommended for use along electrical power, pipeline and telephone rights-of-way, and in other sites associated with these rights-of-way, such as substations, roadsides, railroads or similar rights-of-way that run in conjunction with utilities.

Mirage Plus is also recommended for use in preparing or establishing wildlife openings within these sites, maintaining access roads and for side trimming along utility rights-of-way.

**Tank Mixtures:** Tank mixtures of Mirage Plus may be used to increase the spectrum of control for herbaceous weeds, woody brush and trees. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of all products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of Mirage Plus may be used in a tank mix.

# MIRAGE PLUS

## EPA REG. NO. 34704-

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and tress, use the higher recommended rates.

**NOTE:** For side trimming treatments, it is recommended that Mirage Plus be used alone or in tank mixture with Garlon 4.

Product	Broadcast Rate	Use Sites
Arsenal 2WSL	6 to 32 fluid ounces per acre	Utility Sites
Escort	1 to 2 ounces per acre	Utility Sites
Garlon 3A*, Garlon 4	1 to 4 quarts per acre	Utility Sites/Side Trimming
Oust	1 to 4 ounces per acre	Utility Sites

Product	Spray-To-Wet Rates	Use Sites
Arsenal 2WSL	0.06 to 0.1 percent by volume	Utility Sites
Escort	1 to 2 ounces per acre	Utility sites

Product	Low Volume Directed Spray Rates	Use Sites
Arsenal 2WSL	0.1 to 0.5 percent by volume	Utility Sites
Escort	1 to 2 ounces per acre	Utility sites

\*Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding Mirage Plus. Have spray mixture agitating at the time Mirage Plus is added to avoid spray compatibility problems.

### Bare Ground, Trim-and-Edge

Mirage Plus may be used in utility sites and substations for bare ground, trim-and-edge around objects, spot treatment of unwanted vegetation and to eliminate unwanted weeds growing in established shrub beds or ornamental plantings. Mirage Plus may be used prior to planting a utility site to ornamentals, flowers, turfgrass (sod or seed), or beginning construction projects.

Repeated applications of Mirage Plus may be used, as weeds emerge, to maintain bare ground.

**Tank Mixtures:** Mirage Plus may be tankmixed with the following products provided the specific product listed below is registered for use on these sites. Refer to these products labels for approved non-crop sites and application rates.

Arsenal	Escort	Ronstar™50WP
Banvel	Garlon 3A	Sahara™
Barricade™ 65WG	Plateau™	Simazine
Diuron	Princep™DF	Surflan™
Endurance™	Princep™Liquid	

## MIRAGE PLUS AND TANK MIXTURES FOR NON-CROP AREAS, INCLUDING RAILROAD RIGHTS-OF-WAY, SUBSTATIONS, AIRPORTS, INDUSTRIAL PLANTS, ROADSIDES, STORAGE AREAS AND SIMILAR SITES.

### Use Directions

Do not allow spray mixtures of this herbicide to mist, drip, drift or splash onto desirable vegetation since injury or destruction may occur. Do not apply when wind or other conditions favor drift.

See the "Weeds Controlled" section of the Mirage Plus label booklet for rate recommendations. For difficult to control species, where dense stands occur, or where conditions for control are not ideal, 5 to 10 quarts per acre of Mirage Plus may be used for improved results.

### Tank Mixtures

Mirage Plus provides control of the emerged weeds listed in the label booklet. When applied as a tank mixture, the following herbicides will provide preemergence and/or postemergence control of weeds listed in the individual product labels.

The following list of products may be tank mixed with Mirage Plus provided the specific product listed below is registered for use on these sites. Any recommended rate of Mirage Plus may be used in a tank mixture with these products.

Tank-mix Product	Rate per Acre
Arsenal™	0.5 to 4 pints
Banvel	1 to 4 pints
2,4-D	0.5 to 1 pound
Garlon™3A	1 to 6 pints
Garlon 4	1 to 6 pints
Diuron	4 to 8 pounds
Diuron + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Diuron + Garlon 3A	4 to 10 pounds + 1 to 2 pints
Diuron + Garlon 4	4 to 10 pounds + 1 to 2 pints
Hyvar™X	4 to 8 pounds
Hyvar X + 2,4-D	4 to 8 pounds + 0.5 to 1 pound
Hyvar X + Garlon 3A	4 to 8 pounds + 1 to 2 pints
Hyvar X + Garlon 4	4 to 8 pounds + 1 to 2 pints
Krovar™1DF	4 to 6 pounds

Tank-mix Product	Rate per Acre
Krovar 1DF + 2,4-D	4 to 6 pounds + 0.5 to 1 pound
Krovar 1DF + Garlon 3A	4 to 6 pounds + 1 to 2 pints
Krovar 1DF + Garlon 4	4 to 6 pounds + 1 to 2 pints
Oust™	2 to 6 ounces
Oust + 2,4-D	2 to 6 ounces + 0.5 to 1 pound
Oust + Garlon 3A	2 to 6 ounces + 1 to 2 pints
Oust + Garlon 4	2 to 6 ounces + 1 to 2 pints
Spike™80W	2 to 5 pounds
Spike 80W + 2,4-D	2 to 5 pounds + 0.5 to 1 pound
Spike 80W + Garlon 3A	2 to 5 pounds + 1 to 2 pints
Spike 80W + Garlon 4	2 to 5 pounds + 1 to 2 pints

Refer to the individual product labels for specific non-crop sites, rates, carrier volumes and precautionary statements.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Maintain good agitation at all times during the mixing process. Ensure that the tank-mix products are well mixed with the spray solution before adding Mirage Plus.

Mix only the quantity of spray solution that can be used during the same day. Tank mixtures allowed to stand overnight may result in reduced weed control.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying it resumed.

When used in combination as recommended by Loveland Products, Inc. the liability of Loveland Products, Inc. shall in no manner extend to any damage, loss or injury not solely and directly caused by the inclusion of the Loveland Products, Inc. product in such combination use.

## WEEDS CONTROLLED

Always use the higher rate of Mirage Plus per acre within the recommended range when weed growth is heavy or dense or weeds are growing in an undisturbed (non-cultivated) area.

Reduced results may occur when treating weeds heavily covered with dust. For weeds that have been mowed, grazed or cut, allow regrowth to occur prior to treatment.

Refer to the following label sections for recommended rates for the control of annual and perennial weeds and woody brush and trees. For difficult to control perennial weeds and woody brush and trees, where plants are growing under stressed conditions, or where infestations are dense, Mirage Plus may be used at 5 to 10 quarts per acre for enhanced results.

## ANNUAL WEEDS

Use 1 quart per acre if weeds are less than 6 inches in height or runner length and 1.5 quarts to 4 quarts per acre if weeds are over 6 inches in height or runner length or when weeds are growing under stressed conditions.

For spray-to-wet applications, apply a 0.5 percent solution of Mirage Plus to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaf weeds. For annual weeds over 6 inches tall, or for smaller weeds growing under stressed conditions, use a 1 to 2 percent solution. Use the higher rate for tough-to-control species or for weeds over 24 inches tall.

### Weed Species

Annoda, spurred	Fall panicum*	Medusahead*
Barley*	Falsedandelion*	Morningglory
Barryardgrass*	Falselilax, smallseed*	( <i>Ipomoea</i> spp.)
Bittercress*	Fiddleneck	Mustard, blue*
Black nightshade*	Field pennycress*	Mustard, tansy*
Bluegrass, annual*	Flaree	Mustard, tumble*
Bluegrass, bulbous*	Fleabane, annual*	Mustard, wild*
Bassia, fivehook	Fleabane, hairy	Oats
Brome, downy*	( <i>Coryza bonariensis</i> )*	Pigweed*
Brome, Japanese*	Fleabane, rough*	Plains/Tickseed
Browntop panicum*	Florida pusley	coreopsis*
Buttercup*	Foxtail*	Prickly lettuce*
Carolina foxtail*	Goatgrass, jointed*	Puncturevine
Carolina geranium	Goosegrass	Purslane, common
Castor bean	Grain sorghum (milo)*	Ragweed, common*
Cheatgrass*	Groundsel, common*	Ragweed, giant
Cheeseweed	Hemp sesbania	Red rice
( <i>Malva parviflora</i> )	Henbit	Russian thistle
Chervil*	Horseweed/Marestail	Rye*
Chickweed*	( <i>Coryza canadensis</i> )	Ryegrass*
Cocklebur*	Itchgrass*	Sandbur, field*
Copperleaf,	Johnsongrass, seedling	Shattercane*
hophornbeam	Junglerice	Shepherd's purse*
Corn*	Knotweed	Sicklepod
Corn speedwell*	Kochia	Signalgrass, broadleaf*
Crabgrass*	Lamb's-quarters*	Smartweed, ladythumb*
Dwarf dandelion*	Little barley*	Smartweed,
Eastern mannagrass*	London rocket*	Pennsylvania*
Eclipta*	Mayweed	Sowthistle, annual

29/31

# MIRAGE PLUS

## EPA REG. NO. 34704-

### Weed Species cont'd.:

Spanishneedles	Starthistle, yellow	Virginia pepperweed*
Speedwell, purslane*	Stinkgrass*	Wheat*
Sprangletop*	Sunflower*	Wild oats*
Spurge, annual	Teaweed/Prickly sida	Witchgrass*
Spurge, Prostrate*	Texas panicum*	Woolly cupgrass*
Spurge, spotted*	Velvetleaf	Yellow rocket
Spurry, umbrella*	Virginia coppertleaf	

\*When using field broadcast equipment (aerial applications or boom sprayers using flat-fan nozzles) these species will be controlled or partially controlled using 1 pint of Mirage Plus per acre. Applications must be made using 3 to 10 gallons of carrier volume per acre. Use nozzles that ensure thorough coverage of foliage and treat when weeds are in an early growth stage.

### PERENNIAL WEEDS

Best results are obtained when perennial weeds are treated after they reach the reproductive stage of growth (seedhead initiation in grasses and bud formation in broadleaves). For non-flowering plants, best results are obtained when the plants reach a mature stage of growth. In many situations, treatments are required prior to these growth stages. Under these conditions, use the higher application rate within the recommended range.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment of low volume directed spot treatments, apply a 5 to 10 percent solution of Mirage Plus.

Allow 7 or more days after application before tillage.

Weed Species	Rate (QT/A)	Hand-Held % Solution
Alfalfa*	1	2
Alligatorweed*	4	1.5
Anise (fennel)	2-4	1-2
Bahiagrass	3-5	2
Beachgrass, European ( <i>Ammophila arenaria</i> )	-	5
Bentgrass*	1.5	2
Bermudagrass	5	2
Bermudagrass, water (knotgrass)	1.5	2
Bindweed, field	4-5	2
Bluegrass, Kentucky	2	2
Bluaweed, Texas	4-5	2
Brackenfern	3-4	1-1.5
Bromegrass, smooth	2	2
Bursage, woolly-leaf	-	2
Canarygrass, reed	2-3	2
Cattail	3-5	2
Clover, red, white	3-5	2
Cogongrass	3-5	2
Dallisgrass	3-5	2
Dandelion	3-5	2
Dock curly	3-5	2
Dogbane, hemp	4	2
Fescue (except tall)	3-5	2
Fescue, tall	1-3	2
German ivy	2-4	1-2
Guineagrass	3	1
Horsenettle	3-5	2
Horseradish	4	2
Iceplant	2	1.5-2
Jerusalem artichoke	3-5	2
Johnsongrass	2-3	1
Kikuyugrass	2-3	2
Knapweed	4	2
Lantana	-	1-1.25
Lespedeza	3-5	2
Milkweed, common	3	2
Muhly, wirestem	2	2
Mullein, common	3-5	2
Napiergrass	3-5	2
Nightshade, silverleaf	2	2
Nutsedge, purple, yellow	3	1-2
Orchardgrass	2	2
Pampasgrass	3-5	1.5-2
Paragrass	3-5	2
Pepperweed, perennial	4	2
Phragmites*	3-5	1-2
Poison hemlock	2-4	1-2
Quackgrass	2-3	2
Redvine*	2	2
Reed, giant	4-5	2
Ryegrass, perennial	2-3	1
Smartweed, swamp	3-5	2
Spurge, leafy*	-	2
Sweet potato, wild*	-	2
Thistle, artichoke	2-3	1-2
Thistle, Canada	2-3	2

Weed Species	Rate (QT/A)	Hand-Held % Solution
Timothy	2-3	2
Torpedograss*	4-5	2
Thumpetcreeper*	2-3	2
Vaseygrass	3-5	2
Velvetgrass	3-5	2
Wheatgrass, western	2-3	2

\*Partial control

### WOODY BRUSH AND TREES

Apply Mirage Plus after full leaf expansion, unless otherwise directed. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when applications are made in the spring to early summer when brush species are at high moisture content and are flowering.

Ensure thorough coverage when using spray-to-wet treatments using hand-held equipment. When using hand-held equipment for low volume directed-spray spot treatments, apply a 5 to 10 percent solution of Mirage Plus.

Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

Weed Species	Broadcast Rate (QT/A)	Hand-Held Spray-to-Wet % Solution
Alder	3-4	1-1.5
Ash*	2-5	1-2
Aspen, quaking	2-3	1-1.5
Bearclover (Bearmat)*	2-5	1-2
Beech*	2-5	1-2
Birch	2	1
Blackberry	3-4	1-1.5
Blackgum	2-5	1-2
Bracken	2-5	1-2
Broom; French, Scotch	2-5	1.5-2
Buckwheat, California*	2-4	1-2
Cascara*	2-5	1-2
Catsclaw*	-	1-1.5
Ceanothus*	2-5	1-2
Chamise*	2-5	1
Cherry; bitter, black, pin	2-3	1-1.5
Coyote brush	3-4	1.5-2
Deerweed	2-5	1
Dogwood*	2-5	1-2
Elderberry	2	1
Elm*	2-5	1-2
Eucalyptus	-	2
Gorse*	2-5	1-2
Hasardia*	2-4	1-2
Hawthorn	2-3	1-1.5
Hazel	2	1
Hickory*	2-5	1-2
Honeysuckle	3-4	1-1.5
Hornbeam, American*	2-5	1-2
Kudzu	4	2
Locust, black*	2-4	1-2
Madrone resprouts*	-	2
Mananzita*	2-5	1-2
Maple, red	2-4	1-1.5
Maple, sugar	-	1-1.5
Monkey flower*	2-4	1-2
Oak; black, white*	2-4	1-2
Oak, post	3-4	1-1.5
Oak; northern, pin	2-4	1-1.5
Oak, Scrub*	2-4	1-1.5
Oak; southern red	2-3	1-1.5
Peppertree, Brazilian (Florida holly)*	2-5	1-2
Persimmon*	2-5	1-2
Pine	2-5	1-2
Poison ivy	4-5	2
Poison oak	4-5	2
Poplar, yellow*	2-5	1-2
Redbud, eastern	2-5	1-2
Rose, multiflora	2	1
Russian olive*	2-5	1-2
Sage, black	2-4	1
Sage, white*	2-4	1-2
Sagebrush, California	2-4	1
Salmonberry	2	1
Saltcedar	2-5	1-2
Sassafras*	2-5	1-2
Sourwood*	2-5	1-2
Sumac; laurel, poison, smooth, sugarbush, winged*	2-4	1-2

39/31

**MIRAGE PLUS**  
EPA REG. NO. 34704-

Devrinol is a trademark of United Phosphorus Inc.

31/31

Sim-Trol is trademark of Sipcam Agro USA, Inc.

Resource and Valor are trademarks of Valent USA Corporation.

Desirable Plants And Trees, Because Severe Injury Or Destruction May Result.

See "General Information" and "Mixing" sections of this label booklet for Mirage Plus for essential product performance information.

<b>Broadcast Weed Species</b>	<b>Rate (QT/A)</b>	<b>Hand-Held Spray-to-Wet % Solution</b>
Sweetgum	2-3	1-1.5
Swordfern*	2-5	1-2
Tallowtree, Chinese	-	1
Tan oak resprouts*	-	2
Thimbleberry	2	1
Tobacco, tree*	2-4	1-2
Toyon*	-	2
Trumpet creeper	2-3	1-1.5
Vine maple*	2-5	1-2
Virginia creeper	2-5	1-2
Waxmyrtle, southern*	2-5	1-2
Willow	.3	1
Yerbasanta*	-	2

\*Partial control

**MIRAGE PLUS TOLERANT AND GLYPHOSATE  
TOLERANT HAVE THE SAME MEANING  
THROUGHOUT THIS LABEL.**

**WARRANTY DISCLAIMER AND NOTICE**

THE DIRECTIONS FOR USE OF THIS PRODUCT ARE BELIEVED TO BE ADEQUATE AND SHOULD BE FOLLOWED CAREFULLY. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT DUE TO SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OR ABSENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER.

THE PRODUCTS SOLD TO YOU ARE FURNISHED "AS IS" BY LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER, AND ARE SUBJECT ONLY TO THE MANUFACTURER'S WARRANTIES, IF ANY, WHICH APPEAR ON THE LABELS TO THE PRODUCTS SOLD TO YOU. EXCEPT AS EXPRESSLY PROVIDED HEREIN, LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD OR USE OF THE PRODUCT, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. EXCEPT AS EXPRESSLY STATED HEREIN, LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER MAKES NO WARRANTY OF RESULTS TO BE OBTAINED BY USE OF THE PRODUCT. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND LOVELAND PRODUCTS, INC.'S, THE MANUFACTURER'S OR SELLER'S TOTAL LIABILITY, SHALL BE LIMITED TO DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. NO AGENT OR EMPLOYEE OF LOVELAND PRODUCTS, INC. OR SELLER IS AUTHORIZED TO AMEND THE TERMS OF THIS WARRANTY DISCLAIMER OR THE PRODUCT'S LABEL OR TO MAKE A REPRESENTATION OR RECOMMENDATION DIFFERENT FROM OR INCONSISTENT WITH THE LABEL OF THIS PRODUCT.

IN NO EVENT SHALL LOVELAND PRODUCTS, INC., THE MANUFACTURER OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES AND THE BUYER AND USER WAIVE ANY RIGHT THEY MAY HAVE TO SUCH DAMAGES.

Amplify, Bullet, Degree, Farmsource, Harness, Lariat, Lasso, Micro-Tech, Loveland Products and Vine symbol, and Permit are trademarks of Loveland Products Technology LLC.

Assure, Authority, Canopy, Escort, Hyvar, Krovar, Leadoff, Lexone, Oust, Staple and Telar are trademarks of E.I. duPont de Nemours & Co., Inc.

Coloran, Cotton-Pro, Direx, Karmex, Linex, LoroX and Meturon are trademarks of Griffin LLC.

Barricade, Bicep MAGNUM, Boundary, Caporal, Dual MAGNUM, Endurance, Flexstar, Fusion, Princep, Caliber, Reflex and Solicam. Vanquish and Zorial are trademarks of Syngenta CropProtection Inc.

Firstrate, Fulltime, Garlon, Goal, Python, Spike, Surflan, Topnotch and Tordon are trademarks of Dow AgroSciences LLC.

Arsenal, Banvel, Clarity, Distinct, Frontier, Guardsman, Marksman, Outlook, Pendulum, Plateau, Prowl, Pursuit, Pursuit Plus, Sahara, Scepter, Squadron and Steel are trademarks of BASF Corp.

Balance, Folex, Ginstar, Prep and Ronstar are trademarks of Aventis CropSciences.

Axiom, DEF, Domain, Epic, and Sencor are trademarks of Bayer Crop Protection.

Aim, Command and Gauntlet are trademarks of FMC Corporation.

FORMULATED FOR



P.O. BOX 1286, GREELEY, COLORADO 80632-1286