



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

34913-2

Date of Issuance:

8/24/21

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

SSI MAXIM TRANSFORMER 5G
GRANULAR HERBICIDE

Name and Address of Registrant (include ZIP Code):

Landis International, Inc.
3185 Madison Highway
P.O. Box 5126
Valdosta, GA 31603, USA

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A). You must comply with the following conditions:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product under FIFRA when the Agency requires all registrants of similar products to submit such data.

Signature of Approving Official:

Emily Schmid

Emily Schmid, Product Manager 25
Herbicide Branch, Registration Division (7505P)

Date:

8/24/21

2. You are required to comply with the data requirements described in the DCI Order identified below:

- a. Tebuthiuron GDCI-105501-1540

You must comply with all of the data requirements within the established deadlines. If you have questions about the Generic DCI Order listed above, you may contact the Chemical Review Manager in the Pesticide Reevaluation Division:

<http://iaspub.epa.gov/apex/pesticides/f?p=chemicalsearch:1>

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 34913-2.”
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you fail to satisfy these data requirements, EPA will consider appropriate regulatory action including, among other things, cancellation under FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:


- Basic CSF dated 7/19/2021
- Alternate CSF 1-3 dated 7/19/2021

If you have any questions, please contact Lydia Crawford by phone at 703-347-0622, or via email at Crawford.Lydia@epa.gov.

Enclosure

TEBUTHIURON	GROUP	7	HERBICIDE
IMAZAPYR	GROUP	2	HERBICIDE

SSI MAXIM



TRANSFORMER 5G

GRANULAR HERBICIDE

The Herbicide Designed for Utilities

EPA Reg. No. 34913-XX

EPA Est. No. 39578-TX-1

ACTIVE INGREDIENT:

Tebuthiuron4.0%
 Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1 H-imidazol-2-yl]-3-pyridinecarboxylic acid)1.0%

OTHER INGREDIENTS95.0%
TOTAL100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION


PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For non-emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 Monday through Friday, 8:00 am to 12:00 pm Pacific Time (NPIC Web site: www.npic.orst.edu) For emergencies, call the poison control center at 1-800-222-1222, 24 hours a day, 7 days a week.



Net contents: 40 pounds

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION!**

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear if appropriate. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are polyethylene, polyvinyl chloride, barrier laminate, and butyl, nitrile, neoprene, and natural rubber.

Mixers, loaders and applicators, and other handlers must wear:

- Long-sleeve shirt and long pants,
- Shoes plus socks,
- Chemical resistant gloves made of any waterproof material for all mixers and loaders, plus applicators using handheld equipment.

Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other materials that have been drenched or heavily contaminated with this product. Do not reuse them.

Pilots must use an enclosed cockpit that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to plants. Drift and run-off may be hazardous to plants in water adjacent to treated areas. In case of spills, collect, cover or incorporate granules spilled on the soil surfaces to prevent contamination to water. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater, or rinsate. See Directions for Use for additional precautions and requirements. Treatment of aquatic weeds may result in oxygen depletion or loss due to decomposition of dead plants.

GROUND WATER ADVISORY: This product contains a chemical that is known to leach through soil into ground water under certain conditions as a result of registered (rangeland and non-crop use) uses. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. A shallow water table is defined as depth to water table of 30 feet or less. Permeable soils include, but are not limited to sandy soils.

USE RESTRICTIONS FOR GROUND WATER PROTECTION: Vulnerable Sites: To minimize any movement of tebuthiuron to subsurface water, do not exceed the application rates specified below on treatment sites where soils have a sand or loamy sand texture throughout the soil profile and all of the following characteristics:

1. Rapid to very rapid permeability.
2. Absence of well-defined organic layers or a textural B-horizon (restricting layer of fine textured soil).
3. The water table of an underlying aquifer is shallow.



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781

An aquifer is defined as an "underground saturated, permeable geologic formation capable of producing significant quantities of water to a well or spring". It is the ability of the saturated zone, or portion of that zone, to yield water which makes it an aquifer (American Chemical Society, 1983). Local agricultural agencies can provide further information on the type of soil in your area and the location of shallow ground water aquifers.

Do not apply in areas where the water table is predominately shallow (5 feet or less), such as marshy or sub irrigated areas, or areas immediately adjacent to streams or lakes which are periodically flooded, unless such use is allowed under a state-approved pesticide management program.

Note: Also on such areas, woody plants rooted directly in a shallow water table are minimally affected by applications of tebuthiuron, and poor woody plant control will result.

Do not apply within areas identified by state or local authorities as protected ground water recharge zones.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

PHYSICAL AND CHEMICAL HAZARDS

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE must be mixed and stored only in lined paper bags, stainless steel, fiberglass, plastic and plastic lined steel containers. DO NOT mix or store SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE in unlined steel (except stainless steel) containers.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

NON-CROP WEED CONTROL INDUSTRIAL USE ONLY.

PRODUCT INFORMATION

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE is applied for control of most annual and perennial grasses, broadleaf weeds, brush, and trees. The herbicide is readily absorbed through the roots and is translocated rapidly throughout the plant, with accumulation in the meristematic regions. Treated plants stop growing soon after application. Chlorosis appears first in the newest leaves, and necrosis spreads from this point. In perennials, the herbicide is translocated into and kills underground storage organs, thus preventing regrowth. Chlorosis and necrosis may not be apparent in some plant species until two weeks after application. Complete kill of plants may not occur for several weeks.

Effects are often slow to appear and will not become apparent until sufficient moisture has carried the active ingredients into the root zone and they have been absorbed by the roots and into the plant. The time required to achieve control is dependent on factors including species, soil type, rainfall, and timing of application. Plants are most easily controlled by SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE when applied before emergence or when they are small and actively growing. Drought stress can reduce effectiveness.

An arboriculturist (tree expert) should be consulted to help you to determine if the area of proposed application is free of all roots of desirable vegetation. The effect of this product on desirable vegetation may be irreversible, and its presence in the soil may prevent growth of other desirable vegetation for some years after application. Read the entire label before using this product to determine if this product is suitable for the desired purpose.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE contains both a Group 7 (tebuthiuron) and a Group 2 (imazapyr) herbicide. Any weed population may contain plants naturally resistant to Group 7 and/or Group 2 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE or other Group 7 or Group 2 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is expected, prevent weed seed production in the affected area by an alternate herbicide from a different group or by a mechanical method including hoeing or tillage.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist for additional pesticide resistance-management and/or integrated weed management recommendations for specific weed biotypes.

Contact your local sales representative, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. Do not assume that each listed weed is being controlled by multiple mechanisms of action. Co-formulated active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredient in this product.

For vegetation control by broadcast and spot applications a preemergent or postemergent application of SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may be used to control most annual grasses and broadleaf weeds; as well as most vines, brambles, brush, trees, and other woody species.

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may be used for vegetation control on utility, railroad, pipeline and highway rights-of-way, petroleum tank farms, pumping installations, fence rows, storage areas, and communication tower sites. Utility applications include those made around wooden utility poles, guy wire anchors, switch poles and grates, and under steel transmission towers, It may also be used for fire protection and groundline moisture control around wooden poles. Applications may also be made around fire plugs, utility cabinets, pad mounted transformers, and pipeline markers. Broadcast applications of SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may also be made to utility rights-of-way to eliminate or suppress vegetation and to maintain accessibility.

The maximum labeled rate and frequency of application is 50 lb. of SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE per acre once every three years for vulnerable areas (where soils are sandy and depth of water table is shallow). For all other areas, SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may be applied one time in a three year period at a rate of up to 100 lb. per acre; however, no more than a total of 150 lb. per acre may be applied in two consecutive treatments in any 6 year period.

Spot treatments (hand-held equipment): SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may be applied at rates up to 150 lb./A, (6 lb tebuthiuron/1.5 lb imazapyr) or 5.5 ounces/100 sq. ft. (0.22 oz tebuthiuron/0.055 oz imazapyr) when needed. One-half cup will treat about 100 sq. ft.

RESTRICTIONS

DO NOT enter or allow others to enter treated areas until dusts have settled.

DO NOT apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT use on food or feed crops. DO NOT apply on or near field crops.

DO NOT apply on ditches used to transport irrigation water.

DO NOT apply where runoff or irrigation water may flow onto agricultural land as injury to crops may result.



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781

The use of treated waters on irrigated crops within 120 days of treatment is prohibited.
DO NOT apply more than 150 lbs of product (6lbs tebuthiuron/1.5lbs imazapyr) per acre per application.

DO NOT apply more than 150 lbs of product (6lbs tebuthiuron/1.5lbs imazapyr) per acre per year.
Spot Treatments (Hand-held equipment): SSI Maxim Transformer 5G Granular Herbicide may be applied at rates up to the equivalency of 150 lbs (6lbs tebuthiuron/1.5lbs imazapyr) per acre when needed.
Use only 1 application per year for herbaceous treatments and 1-2 applications per 10 years, not to exceed 1 application per year, for brush treatment. Minimum retreatment interval is 1 year.
Keep from contact with fertilizers, insecticides, fungicides and seeds.
DO NOT apply in locations where the chemical may be washed by runoff or sheet erosion in contact with roots of desirable vegetation, as severe injury or death may occur. Roots of trees, shrubs, and other desirable vegetation may extend far beyond the drip line of the plant's branches.
DO NOT apply on any area into which the roots of desirable vegetation may extend.
DO NOT use on areas such as lawns, walks, driveways, streets, tennis courts, patios, swimming pools, cemeteries, or other landscaped areas, or under asphalt or concrete pavement where future landscaping is planned.

Ditchbank Usage: DO NOT apply to interior ditchbanks (areas which slope toward the drainage). Do not apply to ditches used to transport irrigation or potable water.

DO NOT use in California.

DO NOT APPLY TO THE ROOT ZONE OF DESIRABLE VEGETATION.

Thoroughly clean all traces of this product from application equipment after use. DO NOT empty residues cleaned from application equipment on areas where they may come in contact with the roots of desirable vegetation or the water source for such vegetation.

Herbaceous Perennials (nonwoody plants): Do not cut forage from treated areas for feeding livestock. May injure or suppress certain herbaceous vegetation in the treated area. DO NOT apply where such injury cannot be tolerated.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Except for the protected applicator, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- shoes plus socks
- chemical-resistant gloves made of any waterproof material
- protective eyewear

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Except for protected applicator, do not enter or allow others to enter the treated area until dusts have settled.



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781

APPLICATION INSTRUCTIONS

A preemergence or postemergent application of SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may be used to control most annual and perennial grasses and broadleaf weeds, vines, brush, and trees on noncropland areas including railroad, utility, pipeline and highway rights-of-way, petroleum tank farms, pumping installations, non-agricultural fence rows, and storage areas. SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE may be applied anytime that the ground is not frozen.

For vegetation control by broadcast (ground equipment) and banded applications: If the treated area is to be used for haying, do not apply more than 100 lbs (4 lbs tebuthiuron/1 lb imazapyr) per acre of product, and do not apply this product more than once per year. Grazing is allowed in areas receiving band or individual plant treatments of 100 lbs (4 lbs tebuthiuron/1 lb imazapyr) per acre or less of product.

The maximum label rate and frequency of application is 50 lbs (2 lbs tebuthiuron/0.5 lb imazapyr) per acre once every three years for vulnerable areas (where soils are sandy and depth to water table is shallow.) For all other areas, this product may be applied one time in a 3-year period at a rate of up to 100 lbs. (4 lbs tebuthiuron/1 lb imazapyr) per acre of product; however, no more than 150 lbs. (6 lbs tebuthiuron/1.5 lbs imazapyr) per acre may be applied in two consecutive treatments in any 6 year period.

Weeds Controlled by SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE		
GRASSES	BROADLEAF WEEDS	VINES AND BRAMBLES
ANNUAL BLUEGRASS (<i>Poa annual</i>) BAHIAGRASS (<i>Paspalum notatum</i>) BEARDGRASS (<i>Andropogon</i> spp) BERMUDAGRASS (<i>Cynodon dactylon</i>) BIG BLUESTEM (<i>Andropogon geradi</i>) BROADLEAF SIGNALGRASS (<i>Brachiaria platphyllia</i>) CANADA BLUEGRASS (<i>Poa compressa</i>) CATTAIL (<i>Typha</i> spp) CHEAT (<i>Bromus secalinas</i>) CRABGRASS (<i>Digitaria</i> spp) DALLISGRASS (<i>Paspalum dilatatum</i>) DOWNY BROME (<i>Bromus tectorum</i>) FALL PANICUM (<i>Panicum dichotomiflorum</i>) FEATHER TOP (<i>Penniselum villosum</i>) FESCUE (<i>Festuca</i> spp) FOXTAIL (<i>Setaria</i> spp) GOOSEGRASS (<i>Eleusine indica</i>) GUINEAGRASS (<i>Panicum maximum</i>) ITALIAN RYEGRASS (<i>Lolium multiflorum</i>) JOHNSONGRASS (<i>Sorghum halepense</i>) KENTUCKY BLUEGRASS (<i>Poa pratensis</i>) LOVEGRASS (<i>Eragrostis</i> spp) ORCHARDGRASS (<i>Dactylis glomerata</i>) PARAGRASS (<i>Brachiaria mutica</i>) PHRAGMITES (<i>Phragmites australis</i>) PRARIE CORDGRASS (<i>Spartina pectinata</i>) PRARIE THREEAWN (<i>Aristida oliganina</i>) QUACKGRASS (<i>Agropyron repens</i>) REED CANARY GRASS (<i>Phalaris arundinacea</i>) SALTGRASS (<i>Distichis stracta</i>) SAND DROPSEED (<i>Sporobolus cryptandrus</i>) SANDBUR (<i>Cenchrus</i> spp) SMOOTH BROME (<i>Bromus inermis</i>) TIMOTHY (<i>Phleum pretense</i>) TORPEDOGRASS (<i>Panicum repens</i>) VASEYGRASS (<i>Paspalum urvillei</i>) WILD BARLEY (<i>Hordeum</i> spp) WILD OATS (<i>Avena fatua</i>) WIRESTEM MUHLY (<i>Muhlenbergia Irondosa</i>) WITCHGRASS (<i>Panicum capillare</i>)	BULL THISTLE (<i>Cirsium volgare</i>) BURDOCK (<i>Arctium</i> spp) CAMPHORWEED (<i>Heterotheca subaxillaris</i>) CANADA THISTLE (<i>Cirsium arvense</i>) CAROLINA GERANIUM (<i>Geranium carolinianum</i>) CARPETWEED (<i>Mullugo verticillata</i>) CLOVER (<i>Trifolium</i> spp) COCKLEBUR (<i>Xanthium strumarium</i>) COMMON CHICKWEED (<i>Stellaria media</i>) DANDELION (<i>Taraxacum officinale</i>) DOCK (<i>Rumex</i> spp) DOGFENNEL (<i>Eupatorium capillifolium</i>) FILAREE (<i>Erodium</i> spp) FLEABANE (<i>Erigeron</i> spp) GOLDENROD (<i>Solidago</i> spp) HOARY VERVAIN (<i>Verbena stricta</i>) INDIAN MUSTARD (<i>Brassica juncea</i>) KOCHIA (<i>Kochia scoparia</i>) LAMBSQUARTERS (<i>Chenopodium album</i>) LESPEDEZA (<i>Lespedeza</i> spp) LITTLE MALLOW (<i>Malva parvillora</i>) MILKWEED (<i>Asclepias</i> spp) MINERS LETTUCE (<i>Montia perfoliata</i>) MULLEIN (<i>Verbascum</i> spp) NETTLELEAF GOOSEFOOT (<i>Chenopodium murale</i>) OXEYE DAISY (<i>Chrysanthemum leucanthemum</i>) PEPPERWEED (<i>Lepidium</i> spp) PIGWEED (<i>Amaranthus</i> spp) PLANTAIN (<i>Plantago</i> spp) POKEWEEED (<i>Phytolacca americana</i>) PRIMROSE (<i>Oenothera kunthiana</i>) PURSLANE (<i>Portulaca</i> spp) SILVERLEAF NIGHTSHADE (<i>Solanum elaeagnifolium</i>) SMARTWEED (<i>Polygonum</i> spp) SORREL (<i>Rumex</i> spp) SOWTHISTLE (<i>Sonchus</i> spp) SUNFLOWER (<i>Helianthus</i> spp) SWEET CLOVER (<i>Melilotus</i> spp) TANSYMUSTARD (<i>Descurainia pinnata</i>) TEXAS THISTLE (<i>Cirsium texanum</i>) WILD CARROT (<i>Daucus carota</i>) WILD LETTUCE (<i>Lactuca</i> spp) WILD PARSNIP (<i>Pastinaca sativa</i>) WILDTURNIP (<i>Brassica campestris</i>) WOOLLYLEAF BURSAGE (<i>Ambrosia grayi</i>) YELLOW STARHISTLE (<i>Centaurea solstitialis</i>) YELLOW WOODSORREL (<i>Oxalis stricta</i>)	BLACKBERRY (<i>Rubus</i> spp)* DEWBERRY (<i>Rubus</i> spp)* FIELD BINDWEED (<i>Convolvulus arvensis</i>) GREENBRIAR (<i>Smilax</i> spp) HEDGE BINDWEED (<i>Calystegia sequium</i>) HONEYSUCKLE (<i>Lonicera</i> spp) KUDZU (<i>Pueraria lobata</i>) MORNINGGLORY (<i>Ipomoea</i> spp) POISON IVY (<i>Rhus radicans</i>) REDVINE (<i>Brunnichia cirrhosa</i>) TRUMPETCREEPER (<i>Campsis radicans</i>) VIRGINIA CREEPER (<i>Parthenocissus quinquefolia</i>) WILD BUCKWHEAT (<i>Polygonum convolvulus</i>) WILD GRAPE (<i>Vitis</i> spp) WILD ROSE (<i>Rosa</i> spp) Including: MULTIFLORA ROSE (<i>Rosa multiflora</i>) MACARTNEY ROSE (<i>Rosa bractreata</i>) * The degree of control may be species dependent. Some <i>Rubus</i> species may not be controlled.
		WOODY BRUSH AND TREES ASH (<i>Fraxinus</i> spp) BLACKGUM (<i>Nyssa sylvatica</i>) BOXELDER (<i>Acer negundo</i>) CHERRY (<i>Prunus</i> spp) CHINESE TALLOW-TREE (<i>Sapium sebiterum</i>) DOGWOOD (<i>Cornus</i> spp) HAWTHORN (<i>Crataegus</i> spp) HICKORY (<i>Carya</i> spp) MAPLE (<i>Acer</i> spp) MULBERRY (<i>Morus</i> spp) ** OAK (<i>Quercus</i> spp) PERSIMMON (<i>Diospyros virginiana</i>) POPLAR (<i>Populus</i> spp) PRIVET (<i>Ligustrum vulgare</i>) RED MAPLE (<i>Acer rubrum</i>) SASSAFRAS (<i>Sassafras albidum</i>) SUMAC (<i>Rhus</i> spp) SWEETGUM (<i>Liquidambar styracitula</i>)



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781

INDIVIDUAL (SPOT) WOODY PLANT APPLICATION Apply SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE by hand tossing 2.5 ounces (0.1 oz tebuthiuron (0.625 lb)/0.025 oz imazapyr (0.0016 lb)) of the granules around the base of the unwanted individual woody stem. Wear protective gloves when applying granules by hand. Repeat this process for each unwanted woody plant in the area. Caution: Do not use SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE in this manner in any area where desirable species are in the vicinity of the plants to be eliminated. SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE will injure or control other herbaceous vegetation in the treated area. Do not apply where such injury cannot be tolerated.

WOODY PLANT CONTROL – APPLICATION RATES (BROADCAST)

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE applied at the rate of 50lbs/A (2lb tebuthiuron/0.5lb imazapyr) will control the following species:

Ailanthus altissima (Tree-of-heaven)	Nyssa sylvatica (Black gum)
Aloysia lycioides (Whitebrush)	Quercus nigra (Water oak)
Artemisia vulgaris (Mugort)	Ulmus Americana (American elm)
Liriodendrom tulipifera (Tuliptree)	

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE applied at the rate of 75lbs/A (3lb tebuthiuron/0.75lb imazapyr) will control the following species:

Carya texana (Black hickory)	Symphoricarpos orbiculatus (Buckbrush)
Quercus marilandica (Blackjack Oak)	Ulmus alata (Winged elm)
Quercus stellata (Post oak)	

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE applied at the rate of 100lbs/A (4lb tebuthiuron/1lb imazapyr) will control the following species:

Acacia berlandieri (Guajillo)	Larrea tridentata (Creosotebush)
Acacia rigidula (Blackbrush acacia)	Myrica cerifera (Southern waxmyrtle)
Acer negundo (Boxelder)	Rhus copallina (Shining sumac)
Campsis radicans (Trumpet creeper)	Rosa multiflora (Multiflora Rose)
Celtis occidentalis (Western hackberry)	Rubus allegheniensis (Allegheny blackberry)
Celtis pallida (Grajeno)	

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE applied at the rate of 125lbs/A (5lb tebuthiuron/1.25lb imazapyr) will control the following species:

Acer macrophyllum (Bigleaf maple)	Pinus virginiana (Virginia pine)
Alnus rubra (Red alder)	Prunus emarginata (Bitter cherry)
Cornus florida (Flowering dogwood)	Pseudotsuga menziesii (Douglas fir)
Gaultheria shallon (Salal)	Quercus alba (White oak)
Juniperus virginiana (Eastern redcedar)	Rubus laciniatus (Evergreen blackberry)
Pinus echinata (Shortleaf pine)	Rubus occidentalis (Black raspberry)

SSI MAXIM TRANSFORMER 5G GRANULAR HERBICIDE applied at the rate of 150lbs/A (6lb tebuthiuron/1.5lb imazapyr) will control the following species:

Acer circinatum (Vine maple)	Prunus serotina (Black Cherry)
Cornus spp. (Dogwood)	Rhus glabra (Smooth sumac)
Crataegus spp. (Hawthorn)	Rhus typhina (Staghorn sumac)
Fraxinus americana (White ash)	Salix spp. (Willow)
Haplopappus tenuisectus (Burweed)	Smilax rotundifolia (Common greenbriar)
Liquidambar styraciflua (Sweetgum)	Ulmus parvifolia (Chinese Elm)
Prosopis juliflora (Mesquite)	Ulmus rubra (Slippery elm)



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** DO NOT store below 10°F. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of by using on sites according to label directions or at an approved waste disposal facility.

CONTAINER HANDLING FOR BAG: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONTAINER HANDLING FOR PAIL OR JUG: Nonrefillable container. Do not reuse or refill this container. Completely empty pail into application equipment. Triple rinse (or equivalent). Then offer for recycling or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The use directions are believed to be reliable and must be followed. To the extent permitted by applicable law, it is impossible to eliminate all risks inherently associated with use of this product. To the extent permitted by applicable law, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use or application of the product contrary to label instructions, all of which are beyond the control of SSI Maxim Company. All such risks shall be assumed by the user. SSI Maxim Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions of use, subject to the risks referred to above. To the extent permitted by applicable law, all such risks shall be assumed by the user. To the extent permitted by applicable law, SSI Maxim Company makes no other express or implied warranty, including other express or implied warranty of FITNESS or of MERCHANTABILITY.



SSI MAXIM COMPANY, INC.

P.O. Box 1954 • Kilgore, TX 75663 • 8003464781