RIGHT SIDE PANEL CONTINUED

		after spraying, irrigate treated area with \$ to 1 inch of water to wash the insecticide into the underlying soil
Bluegrass billbug	1½ - 3 fl. oz. per 100 sq. ft.	Spray early in the season when adults first appear. Retreat as needed.

^{* 3/4} fl. oz. per 1000 sq. ft. = 3 3/4 fl. oz. per 5000 sq. ft. = \frac{1}{2} pint per 10,700 sq. ft. = 1 quart per acre

Perimeter Treatment around buildings: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes and sowbugs (pillbuge), treat a band of soil 6 to 10 feet wide adjacent to the building. Also treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Use 2½ fl. oz. of ATLANTIC TC 4 per 100 gal. of water. Apply as a coarse spray to wet the band thoroughly and uniformly.

WARRANTY STATESENT

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTARILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably forseeable to seller, and buyer assumes the risk of any such use.

		VELP		
U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS	EPA REGISTRATION NO.	°N5v° 350° 1582 '		
REGI'TRATION DIVISION (WH-567) WASHINGTO G.C. 27467	TERM OF ISSUANCE			
NOTICE OF PESTICIDE: REGISTRATION	NAME OF PESTICIDE PRODUCT	· · · · · · · · · · · · · · · · · · ·		
Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended:				
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)		1 84578		
r	$\neg \frac{16}{1}$	87 1		
· 				
L	T			
NOTE: Changes in labeling formula differing in substance submitted to and accepted by the Registration Division product always refer to the above U.S. EPA registration nur	or to use of the label in commerce other.	. In any correspondence on this		
On the basis of information furnished by the registrant, the the Federal Insecticide, Fungicide, and Rodenticide Act.	above named pesticide is hereby	Registered/Reregistered under		
A copy of the labeling accepted in connection with this Re	gistration/Reregistration is retur	ned herewith.		
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.				
	_			
	4 .			
	•			
ATTACHMENT IS APPLICABLE				
SIGNATURE OF APPROVING OFFICIAL		DAYE		

FRONT PANEL

ACCEPTED
with COMMENTS
in EPA Letter Dated:



NOV 3 0 1982

1.415 ~

ATLANTIC TC 4

Under the Federal Interfede.
Fungacide, and Redentions According DURSBAN* and VAPONA**
By accorded, for the polytools Plus Wetting Agent
registered under EPA By. No.

Only for sale to, use and storage by Service Fersons such as 'Commercial Lawn Spraymen, Pest Control Operators, and other persons responsible for insect control in ornamental turf areas such as golf courses, playing fields, and parks.

For control of Chinch Bugs, Sod Webworms, Ants, Earwigs, Crickets, Ticks, Fleas, Armyworms, and Grasshoppers.

Active Ingredients:

Tris product contains lbs. of chlorpyrifes and lbs. of dichlorves per gallon.

- *DURSBALD Trademark of the Dow Chemical Company U.S. Patent No. 3,244,586
- **VAFONA® Trademark of the Shell Chemical Company

KEEL OUT OF REACH OF CHILDREN

DANGER - POISON



DIATEMENT OF PRACTICAL TREATMENT

If SWALLOWED: All a physician or Poison Control Center immediately.

If possible, vom. ting should be induced under medical supervision. Drais one or two glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN! Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAL: Chlorpyrifos is a cholinesterase inhibitor.
Treat symptomatically. Atropine only by injection is an antidote.

See live paner for recautionary Statements

manufactured by

- ATLANTIC FERTILIZER & CHEMICAL CO. Homestead, Florida 33030

EFA Est. No.35455-FL-01

CHA Reg. No. 35455- .

NET CONTENTO

FRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PCISON



May be fatal is swallowed - May be absorbed through skin - May be injurious to eyes and skin.

Do not take internally. Do not get in eyes or on ski .. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Avoid spraying food crops. Avoid breathing vapors and spray mist. Handle concentrate in ventilated area. Keep children and pets off treated areas until spray has dried.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the turf, weeds, or other vegetation blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

ATIANTIC TC 4 is toxic to fish, birds and other wildlife. Keep out of lates, streams, ponds, tidal marshes and estuaries. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply when these are important resources. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

THYSICAL OR CHEMICAL MAZARDO

- Combustible -

Do not use or store near heat or open flame. Do not cut or weld container. - Do not ship or store with food, feeds, or clothing.

PRECAUTION AL USUARIO: Si usted no lee ingles, no ase este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: "To the User: If you cannot read anglish, do not use this inoduct until the laber has been fully explained to you."

LEFT SIDE PANEL CONTINUED

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISFOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep away from food, feedstuffs and domestic water supplies. Keep container closed.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or Local proceedures under the Resource Conservation and Recovery Act.

of in a sanitary landfill or by other approved State and Local proceedures. Metal containers may be offered for recycling or reconditioning.

. . .

RIGHT SIDE PANEL

General Information: Use ATLANTIC TC 4 for control of certain pests infesting lawns and ornamental turf, and for area control of fleas, ticks and chiggers. Consult the following directions for dosage and application details to control the pest listed.

Pests of Lawns and other Ornamental and Recreational Turfgrass:
Mix ATLANTIC TC 4 with water according to directions
given in the following table. Apply as a coarse, low
pressure spray. For best results, turf should be
moist at the time of treatment and care should be
taken to obtain complete and uniform coverage of pest
infested areas. Spray when pests first appear and
re-treat as needed.

Dosages and Directions for use on Turf:

ATLANTIC TO 4

Dilute with 30 gal. of water per 1000 sq. ft.

PEST'	LOSAGE	REMARKS	
Ants Armyworms Chiggers Chinch bugs Clover mite Crickets Cutworms Earwigs Fleas Grasshoppers Sod webworms Ticks	3/4 fl. oz.* per 1000 sq. ft.	Spray when pests first appear, re- treat when needed. For sod webworms delay watering or mowing treated area for 12 to 24 hours after treatment.	
Turfgrass (Hyperodes) weevil	1; fl. oz. per 100 sq. ft.	Openay suspected problem areas in mid-April and again in mid-may or as recommended by your local agricultural extension Service.	
Tapanese beetle larvae only)	15 - 3 °1. oz. per 100 sq. it.	for white grubs, pray when grubs are young and active, feeding near the soil surface, usually during late July and active, or as your formal sericultural extension Dervice recommends. Immediate	