



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

DEC 16 1997

H.R. McLane
Howard Fertilizer Corporation
P.O. Box 13800
Orlando, FL 32859

Dear Mr. McLane:

SUBJECT: Turf Pride With 75% Ronstar
EPA Registration No. 35512-26
Your Application Dated October 24, 1997

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable. A stamped copy is enclosed for your records. Submit one copy of your final printed labeling before you release the product for shipment.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

son. Cooke
12/16/97

2014

ACCEPTED
with COMMENTS
In EPA Letter Dated

DEC 16 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

35512-26

2014



TURF PRIDE® with .75% RONSTAR®

50 lb. Bag Treats up to 11,000 Sq.Ft.

Ask for a Copy of the Supplemental Label Available For The Use Of This Product On Ornamental Shrubs, Vines & Trees FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY. FOR COMMERCIAL USE ON PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS.

A PRE-EMERGENT WEED CONTROL PLUS FERTILIZER FOR PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS, ST. AUGUSTINEGRASS, BERMUDAGRASS, TALL FESCUE AND ZOYSIAGRASS

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed & Other Weeds Listed.

ACTIVE INGREDIENT:

Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)7H-1,3,4-oxadiazolin-5-one] - 0.75%

INERT INGREDIENTS: 99.25%
Total 100.00%

Contains RONSTAR® Herbicide
a Registered Trade Mark of Rhone Poulenc Inc.

FOR LAWNS AND ORNAMENTAL TURF NOT FOR USE ON HOME LAWNS. FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF & LANDSCAPE PERSONNEL ONLY

When applied according to Directions this product can be used safely on the following established turfgrass varieties: BERMUDAGRASS, PERENNIAL BLUEGRASS, PERENNIAL RYEGRASS VARIETIES, ST. AUGUSTINE GRASS VARIETIES, TALL FESCUE, & ZOYSIAGRASS lawns & ornamental turf. It may be used for Apartment and Condominium complexes, Parks, Golf Courses, Athletic Fields, Cemeteries and all other ornamental and recreational turfgrass areas.

CONTROLS GRASSY WEEDS & BROADLEAF WEEDS
Annual Bluegrass (Poa Annua), Barnyard Grass, Crabgrass, Fall Panicum, Florida Pusley, Goose grass, Green Foxtail, Oxalis, Stinging Nettle, Carpet grass, Pigweed.

HOW TO APPLY

For best weed control rake the area free of leaves and thatch and mow if necessary so that granules will reach the soil surface. Measure the area to be treated and uniformly apply the required amount using a fertilizer spreader (see Spreader Settings chart). Do not apply to turf when it is wet. Do not over-lap spreader applications, on St. Augustinegrass since discoloration "streaking" may temporarily occur from "double rate". After applications if rain is not expected within 24 hours sprinkle the treated turf with 1/4 to 1/2 inch of water to get maximum weed control.

HOW MUCH TO USE

FOR CRABGRASS: Apply 50 lbs. / 8,150 sq.ft. when turf has been in a yearly maintenance program (equal to 2 lbs. A.I. herbicide per acre.) Apply 50 lbs. to treat 5,500 sq.ft. of lawn if it has been heavily infested with Crabgrass; equal to 3 lb. A.I. herbicide per acre. Apply in early spring 1 to 2 weeks prior to weed seed germination. This product is a pre-emergence herbicide. It will not control emerged and growing weeds; only germinating weed seeds. Late applications will control only those weed seeds that germinate after the application. See "When To Apply" Chart.

GOOSEGRASS and PROSTRATE SPURGE: These weeds do not usually germinate until late spring or early summer. Apply 50 lbs. to treat 4,100 sq.ft. (4 lbs. A.I./acre) and apply 1 to 2 weeks prior to seed germination. See Chart for When To Apply. Observe maximum rate for St. Augustine grass.

FOR ANNUAL BLUEGRASS (POA ANNUA): Apply 50 lbs. per 4,100 to 5,500 sq.ft. of Lawn; equal to 3 to 4 lbs. active herbicide per acre. Apply in late summer to fall or in early spring 1 to 2 weeks before expected weed seed germination. Will not control annual bluegrass present at time of application; therefore only germinating seeds. Refer to chart for proper timing in your locale. Annual Barnyardgrass and Prostrate Spurge may also be controlled by this same rate. Observe maximum rate for St. Augustine grass, which is 3 # active ingredient / acre.

POA ANNUA "ANNUAL BLUEGRASS" CONTROL IN BERMUDAGRASS FAIRWAYS OVERSEEDED WITH PERENNIAL RYEGRASS ONLY Apply 50 lbs. to treat 8,200 sq.ft. (2 lbs. a.i./Acre) Apply in late summer to early fall prior to Poa Annua spp. annua seed germination. Established plants will not be controlled. This product should be applied no closer than 60 days prior to overseeding. **NOTE:** Do not use where Poa trivialis is present in overseed mix. **IMPORTANT:** In many areas annual bluegrass exists as at least two biotypes or subspecies of Poa Annua. The true annual biotype, "Poa Annua spp. annua" which germinates from seed every year is controlled, while the biotype, "Poa Annua spp. raptans" which behaves like a perennial and survives through the summer, as an established plant, cannot be controlled. Consult the extension service or University weed specialist in your area for more information concerning your locale.

FOR PIGWEED, FLORIDA PUSLEY, OXALIS, STINGING NETTLE AND CARPETGRASS: Where these weeds are a problem apply 50 lbs. to treat 4,100 to 4,600 sq.ft. of Lawn; (equal to 3.5 to 4 lbs. Active Herbicide per acre) in early spring or 1 to 2 weeks before expected weed seed germination. Will not control of emerged and growing weeds; only effective against germinating weed seeds. Observe maximum rate for St. Augustine grass. (50#/5,500)

SPREADER SETTINGS FOR SQUAHE FOOT-COVERAGE

| SPREADER PRODUCT | AMT. OF FOR 8,150 Sq.Ft. | 5,500 sq.Ft. | 4,100 Sq.Ft. |
|------------------|--------------------------|--------------|--------------|
| CYCLONE B-1 | 50 LBS. 4 1/2" | 5 1/2" | 5 3/4" |
| SCOTTS #35 | 50 LBS. 5" | 6" | 7" |

NOTE: Spreader rate varies with walking speed. Check spreader over known area.

WHEN TO APPLY

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE

| JAN. | FEB. | MAR. | APR | MAY | JUNE | JULY | AUG. | SEPT. | OCT. | NOV. | DEC. |
|------|------|------|-----|-----|------|------|------|-------|------|------|------|
| | | 1. | | | | | | 3. | | | |
| | | | | | | | 3a. | | | | |
| | | | | | | | | | 4. | | |

1. For CRABGRASS & other Grasses & Broadleaf Weeds germinating in the Spring in Northern turf.
2. For GOOSEGRASS & other Late Germinating Annual Grasses & Broadleaf Weeds such as PROSTRATE SPURGE. 2a. GOOSEGRASS in Southern Turf. PRECAUTIONS IN USING
3. For POA ANNUA in Northern Turf. 3a. For POA ANNUA in Southern Turf.
4. For Fall-Winter-Spring control of ANNUAL WEEDS in Southern Turf.

PRECAUTIONS IN USING

Do not use on desirable Red Fescue, Bentgrass varieties, Dichondra, or Centipede grass since damage will occur. Do not use on Putting greens or "Tees" due to varying cultural practices used. Use only the rate of 50 lbs. to treat 5,500 sq.ft. on St. Augustinegrass since temporary discoloration will occur from heavier rates. Avoid contact with flowers and shrubs except as recommended on the Ornamental Use Supplemental Labeling available for this product. For newly seeded lawns do not apply this product until after the second or third cutting. For best lawn care, place new grass seed in the fall of the year. Spring pre-emergence weed control can then be applied.

RESEEDING TREATED AREAS

Delay new seeding until 4 months after treatment.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (28WPSOK PM5 / WPS/CERTIFICATION/10-25-93 \ EPA acpt'd 03-14-94amd for Poa10-22-97))

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks; protective eyewear; chemical resistant headgear for overhead exposure; chemical resistant apron when cleaning equipment, mixing or loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short sleeved shirt and short pants; waterproof gloves, chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS

| | |
|---|------------|
| Total Nitrogen (N) | variable % |
| Nitrate Nitrogen | 0% |
| Ammoniacal Nitrogen | 0% |
| Water Soluble Nitrogen and/or Urea Nitrogen | 0% |
| Water Insoluble Nitrogen | 0% |
| Available Phosphoric Acid (P ₂ O ₅) | variable % |
| Soluble Potash (K ₂ O) | variable % |
| Derived from Ammonium Sulfate, Ureaform, Muriate of Potash; Manganese as Mn .5%; Iron as Fe 1.0%; Chlorine <15% | |

NET WEIGHT 50 POUNDS (25; 40;)

EPA Reg. No. 35512-26 EPA EST. NO. 35512-FL-1

HOWARD FERTILIZER COMPANY
P.O. BOX 13800 - ORLANDO, FL 32859

404



TURF PRIDE® with .75% RONSTAR

SUPPLEMENTAL LABELING FOR WOODY ORNAMENTAL SHRUBS, VINES, TREES and CONTAINER GROWN ORNAMENTALS

50 lb. Bag Treats up to 8,150 Sq.Ft.

A Commercial Weed & Feed "Preemergent + Fertilizer" FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY

SEE BASIC TURF PRIDE® PRODUCT LABEL EPA REG NO. 35512-24 FOR THE COMMERCIAL USE OF THIS PRODUCT ON PARKS, GOLF COURSES, ATHLETIC FIELDS, AND OTHER RECREATIONAL AND ORNAMENTAL TURFGRASS AREAS

Controls: Crabgrass, Annual Bluegrass, Goosegrass, Florida Pusley, Oxalis, Stinging Nettle, Carpetgrass, Pigweed and Many Other Weeds as Listed

ACTIVE INGREDIENT:

Oxadiazon; [2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl) N-1,3,4-oxadiazolin-5-one] 0.75%

INERT INGREDIENTS: 99.25%
Total 100.00%

Contains RONSTAR® Herbicide a Registered Trade Mark of Rhone Poulenc Inc.

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye and skin irritation. Avoid contact with skin, eyes or clothing. Do not allow domestic animals to graze treated areas. Avoid contamination of feed and foodstuffs.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical resistant apron when cleaning equipment, mixing and loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the Restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over short-sleeved shirt and short pants; waterproof gloves; chemical resistant footwear plus socks, protective eyewear, and chemical resistant headgear for overhead exposure.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store product in its original bag, in a cool, dry, locked place out of reach of children. In case of spill, sweep up, and use according to directions or dispose of as follows. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GUARANTEED FERTILIZER ANALYSIS

| | |
|---|------------|
| Total Nitrogen (N) | variable % |
| Nitrate Nitrogen | 0% |
| Ammoniacal Nitrogen | 12% |
| Water Soluble Nitrogen and/or Urea Nitrogen | 0% |
| Water Insoluble Nitrogen | 3% |
| Available Phosphoric Acid (P ₂ O ₅) | variable % |
| Soluble Potash (K ₂ O) | variable % |
| Derived from Ammonium Sulfate, Ureaform, Muriate of Potash, Manganese as Mn .5% Iron as Fe 1.0% Chlorine <15% | |

WEED CONTROL AROUND WOODY ORNAMENTAL SHRUBS, VINES & TREES. CONTROLS the FOLLOWING GRASSY WEEDS and BROADLEAF WEEDS

Annual Bluegrass, Barnyardgrass, Bittercress, Carpetweed, Carpetgrass, Common Groundsel, Common Purslane, Crabgrass, *Evening Primrose, Green Foxtail, Fall Panicum, Florida Pusley, *Galinsoga, Golden Ragwort, *Goosegrass, Lambsquarter, Oxalis, Pennsylvania Smartweed, Petty Spurge, *Prostrate Spurge, Redroot Pigweed, Spotted Catsear, Stinging Nettle, *Swinecress, Yellow Woodsorrel.

Can be applied to both newly transplanted and established ornamentals plants where both weed control and fertilization is desired. Rainfall or overhead irrigation will improve weed control. Do not disturb or cultivate soil after treatment. Do not apply to wet foliage or under conditions in which granules will collect on leaves.

USE PRECAUTIONS

Avoid contact with flowers and shrubs except as recommended on this label.

HOW MUCH TO USE

For susceptible weeds apply 50 lbs. to treat 8,150 to 5,500 sq.ft. of Ornamental planted area. See Chart WHEN TO APPLY. (50 lbs. / 5,500 sq.ft. is preferred rate.)

*For hard to kill weeds as indicated use the rate of 50 lbs. for 5,500 sq.ft. or 50 lbs. for 4,100 sq.ft. (3 to 4 lbs. A.I./Acre) = 9 lbs. to 12.2 lbs. actual product per 1000 sq. ft. of plant bed.

FOR AREAS WHERE WEEDS ARE GROWING

If treatment is desired in non-turf areas where weeds are presently growing first remove existing weeds. Rake and smooth the soil by "breaking-up" clumps. Apply at rate as directed for specific weeds. (50 lbs. to treat either 5,500 or 8,150 sq.ft.) This treatment will only control weeds germinating from seeds. It will not control weeds sprouting from roots! Irrigate the area after treatment with 1/4 to 1/2 inch water to get maximum weed control.

WEED AND FEED FOR CONTAINER GROWN ORNAMENTAL PLANTS

This product is recommended on a wide variety of Container and Field Grown Woody Ornamental Shrubs, Vines and Trees. It may be used on the following list of Container Grown Plants. Use at the rate of 50 lbs. per 4,100 sq.ft. of pot soil surface area. Plants in 14 inch containers set out in 12 foot wide rows X 100 feet long will require 10 3/4 lbs. product per row (Equivalent to 4 lbs. A.I. per acre). Apply with a "shoulder strap held" grass-seeder or granule applicator that will direct the granules to each pot soil surface.

| | | | | |
|---------------------------|-------------|-------------------|----------------|--------------|
| Agapanthus | Cedar | Heath Heather | Ornithogalum | Rose |
| Andromeda | Cotoneaster | Hedera | Petunias | Star Jasmine |
| Arborvita | Cypress | Holly | Phlox | Syzygium |
| Archostaphylos | Dogwood | Honeysuckle | Pine-Varieties | Viburnum |
| Azalea | Douglas Fir | Jasmine | Platanus | Xylocopa |
| Barberry | Eucalyptus | Juniper-Varieties | Podocarpus | |
| Boxwood | Euonymus | Jusica | Privet | |
| Common-Japanese evergreen | Leucodermis | Leucodermis | Pyracantha | |
| Callistemon | Forsythia | Magnolia | Quercus | |
| Camellia | Gardenia | Malvaceae | Raphanistrum | |
| Ceanothus | Ginkgo | Myrtle | Rhododendron | |

FOR FIELD GROWN ORNAMENTALS

*Apply at the rate of 50 lbs. per 5,000 or 8,150 sq.ft. of area.

This product may be used around the following Established Plants.

| | | | |
|------------------|-------------|---------------|----------------|
| Arborvita | Euonymus | Pittosporum | Ground Covers: |
| Ash | Forsythia | Privet | *Carobrotus |
| Azalea | Holly | Rhododendron | Osteospermum |
| Barberry | Honeysuckle | Rose | *Sedum |
| Japanese Juniper | Juniper | Russian Olive | Algerian Ivy |
| Birch | Lilac | Spruce | Vinca Major |
| Camellia | Maple | Taxus | English Ivy |
| Cotoneaster | Oak | Viburnum | Pachasandra |
| Crabapple | Periwinkle | Xylocopa | Ajuga |
| Dogwood | Pine | | Euonymus |

*Use higher rate (50 lbs./5,500 sq.ft.) for hard to kill weeds. May be used around the following Newly Transplanted Ornamentals.

| | | | |
|-----------|------------|---------------|--------|
| Camellia | Holly | Pine | Spruce |
| Dogwood | Juniper | Privet | Taxus |
| Forsythia | Oak | Rhododendron | Teuga |
| Hedera | Periwinkle | Russian Olive | |

To obtain continued season long weed control on the following list of plants additional applications may be made 60 days after the previous application in the following situations.

APPLY 2 REPEAT TREATMENTS

50 lbs / 5,500 SQ.FT ON FOLLOWING PLANTS

Established - Container Grown Plants - Abelia, Common Boxwood, English Ivy, Holly, Juniper, Pine, Rhododendron. Newly Transplanted - Container Grown Plants: Common Boxwood, English Ivy, Juniper, Pine, Rhododendron.

APPLY 3 REPEAT TREATMENTS

50 lbs / 5,500 SQ.FT. ON FOLLOWING PLANTS

Established - Container Grown Plants - Abelia, Holly, Rhododendron.

Forest Nurseries - For Pines - Loblolly, Slash, Eastern White.

NOTICE: Buyer and user assume all risk and liability of use, storage and/or handling of this product not in accordance with the terms of this label. (26WPSPOK.PMS / WPS CERT/ EPA acc'd 03-14-04)

EPA Reg. No. 35512-26 EPA EST. NO. 35512-FL-1

NET WEIGHT 25 (40)(50) POUNDS NET

HOWARD FERTILIZER COMPANY

P.O. BOX 13800 - ORLANDO, FL 32859