



**ADDITIONAL INSTRUCTIONS:** Determine the amount of Fouaphene E-6 to be used in the spray by multiplying the number of acres to be treated by the quantity of Fouaphene E-6 recommended per acre. Add this amount of Fouaphene E-6 to one-half the water required for coverage. Mix well the total remaining water. Continue mixing until spray mixture is uniform. By Air Use 10 to 15 gallons water per acre. BY GROUND Use 20 to 30 gallons of water per acre.

**ORNAMENTALS**

**CANERWORM and FALL WEAVER** (on Elm and Maple) **CATALEPA WORM** (on Catalpa) **TENT CATERpillARS** (on Oak, Willow, Poplar) Use 2 quarts Fouaphene E-6 in 100 gallons of water. **MAY BEETLES** (on nursery stock of Elm, Maple, Oak) **BARK WORM** (on evergreens) Use 1 1/2 to 2 quarts per 100 gallons of water. Use full coverage spray.

**LIVESTOCK SPRAYING**

**DO NOT USE FOR DIPPING.** Do not use on animals within 30 days of slaughter.

**CATTLE:** For control of Horn Flies, Ticks and Lice. Dilute 1 pint in 15 gallons of water. Apply thoroughly and apply as spray only. Use driving spray to wet skin. 2 quarts of solution per adult animal. Repeat as needed. Do not use on dairy cattle or calves under 6 months of age.

**SHEEP and Goats:** For control of Fleecce Worms and Ticks (Linnite) and Winter Ticks. Dilute 1 pint in 15 gallons of water. Keep spray agitated. spray animal thoroughly. For lice and Sheep Ticks use pint in 10 gallons of water. Do not apply to dairy goats.

Avoid treating animals during cold weather, overheated or sick animals.

**COTTON**

Do not feed treated cotton trash to dairy animals or animals being raised for slaughter. Do not contaminate lakes or streams. The dosage depends on size of plants. Repeat application as necessary.

INSECT	AMOUNT E-6 PER ACRE	TIME FOR TREATMENT
Boll Weevil	2-2 3 pts	From first square until maturity
Cotton Fleahopper	1 qt	When cotton begins to square
Cotton Leafworm	2-2 3 pts	As soon as young worms hatch
Cutworms	2-2 3 pts	Soon after seeding cotton is up
Fall Armyworm	2-2 3 pts	When young worms appear
Grass-hoppers	2-2 3 pts	Usually early in growing season
Lygus Bugs	2-2 3 pts	When cotton begins to square
Thrips	1 pt	2 to 4 leaf stage of cotton

CROP	INSECT	AMOUNT	TIME FOR TREATMENT	REMARKS
Corn, Sorghum	Yellow Stippled Armyworm, Fall Armyworm, Corn Earworm, Cutworm, Caterpillar, Flea Hopper, Lygus Bug, Green Clover Worm	2-2 3 pts	When insects first appear	Do not apply to corn 30 days before harvest. Do not apply to sorghum 15 days before harvest.
Corn, Soybeans, Wheat, Oats, Rice, Barley, Rye	Yellow Stippled Armyworm, Green Clover Worm, Climbing Cutworm, Fall Armyworm	2-2 3 pts	When insects first appear	Do not feed treated forage to cattle or animals being raised for slaughter. Do not use on forage to be shipped to market.
Grain, Alfalfa	Yellow Stippled Armyworm, Webworm, Climbing Cutworm, Green Clover Worm	2-2 3 pts	When insects first appear	Do not apply to GRAIN, DAIRY ANIMALS, or ANIMALS BEING RAISED FOR SLAUGHTER.
Beans	Armyworms, Grasshoppers, Lygus Bugs	2-2 3 pts	When insects first appear	Do not apply to BEANS or FEED. Do not use on beans for animal feed.
Tomatoes	Armyworms, Corn Earworms	2-2 3 pts	When insects first appear	Do not apply to TOMATOES. Do not use on tomatoes for animal feed.
Peanuts	Fall Armyworms, Velvet Beans, Cutworm	2-2 3 pts	When insects first appear	Do not apply to PEANUTS. Do not use on peanuts for animal feed.
Soybeans	Yellow Bean Caterpillar, Armyworm, Corn Earworm, Bean Leaf Beetle, Armyworms, Grasshoppers, Beetle, Green Clover Worm	2-2 3 pts	When insects first appear	Do not apply to soybeans 30 days before harvest. Do not use on soybeans for animal feed.
Cabbages	Cutworm, Cabbage Looper, Imported Cabbageworm	2-2 3 pts	When insects first appear	Do not apply to cabbages 30 days before harvest. Do not use on cabbages for animal feed.
Cole Crops, Kale	Cutworm, Cabbage Looper, Imported Cabbageworm	2-2 3 pts	When insects first appear	Apply 3 times at 7, 14, and 21 days after planting. Do not use on cole crops for animal feed.
Corn, Beans, Soybeans, Sorghum	Yellow Stippled Armyworm, Green Clover Worm, Fall Armyworm, Cutworm, Caterpillar, Flea Hopper, Lygus Bug, Green Clover Worm	2-2 3 pts	When insects first appear	Do not apply to LEAF LET. Do not use on leaf let for animal feed.

Fouaphene E-6 is a registered trademark of the Dow Chemical Company.