36638-29





COTTON



PINK BOLLWORM




NolMate® PBW Spiral is a pheromone communication disruptant for control of the pink bollworm (*Pectinophora gossypiella*)

Active Ingredients:

(Z,E)-7,11 Hexadecadien-1-yl Acetate 5.37%
(Z,Z)-7,11 Hexadecadien-1-yl Acetate 5.37%
Inert Ingredients
Total
*165ma of pheromone per dispenser.

KEEP OUT OF REACH OF CHILDREN CAUTION



PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid prolonged contact with skin. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.



610 Central Avenue Billings, Montana 59102 (406) 248-5856 800-735-5323 www.scentry.com

EPA REG. No. 36638-29 EPA EST. No. 36638-MT-01 Net contents: (500 spirals)

SCENTRY BIOLOGICALS

loMate[®]

PBW SPIRAL

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool place until used. Prolonged storage should be at or less than 72° F (22°C).

Pesticide Disposal: After use, used spirals should be retrieved if practical. Discard any retrieved spirals in trash.

Container Disposal: Do not reuse empty pouch. Discard pouch in trash.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NoMate PBW Spiral is intended to prevent damage to cotton by the pink bollworm. The product functions by disrupting the mating communication between the sexes of the adult moths. By interfering with the natural mating process, fertile egg laying and subsequent larval infestation can be suppressed, thus reducing the need for conventional insecticides. Reduction in the use of conventional insecticides permits an increase in the populations of parasites and predators which assist in the biological control of other pests.

RATE OF APPLICATION

The rate of application is 100-400 spirals per acre (250-1000 spirals per hectare). Use the spirals in a single application per year.

METHOD OF APPLICATION

NoMate PBW Spiral is applied by wrapping the spiral around stems or branches. Open the container just prior to application. Reseal unused product and refrigerate.

Apply the product as high in the crop canopy as possible to avoid covering spirals during cultivation. A uniform field distribution is necessary for maximum effectiveness.

TIMING OF APPLICATION

Monitor pink bollworm populations using the Scentry® "Delta" Style traps and Scentry® PBW pheromone lures. Place traps in fields soon after planting and continue monitoring throughout the season to assess treatment effectiveness. NoMate PBW Spiral is to be applied when the following conditions exist in the field:

- 1. Trap counts average 1 moth per trap per night.
- Susceptible pinsquares or cotton bolls are present.
- 3. Cotton stems are strong enough to support the spirals.

If plants are not of sufficient size to support the weight of the spirals, an application of NoMate PBW Fiber or MEC may be needed until plants reach the appropriate stage.

WARRANTY AND CONDITIONS OF SALE Scentry Biologicals warrants that this product conforms to the description of this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use.

SCENTRY BIOLOGICALS MAKES NO WAR-RANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

If this product is defective, Buyer's exclusive remedy shall be the replacement of the product, or if replacement is impracticable refund of the purchase price. In no case will Scentry Biologicals be liable for incidental, consequential or special damages resulting from the handling, storage or use of this product.

NoMate and Scentry are trademarks of Scentry Biologicals, Inc. Ver. 3/01 - Copyright @2001 Scentry Biologicals, Inc.