

[37060-1]

NET WT. POUNDS

TRI-EXCEL[®] DS

A general purpose Insecticide and Miticide for use on
Roses, Flowers, Vegetable crops, Fruit Trees, Shrubs, Evergreens and other ornamentals

ACTIVE INGREDIENTS:

Pyrethrins	0.100%	w/w
Rotenone	0.750%	
Other cube resins	1.500%	
*Ryanodine	0.055%	
INERT INGREDIENTS:	97.595%	
	100.000%	

*From ground stems of *Ryania Speciosa*.

CAUTION

KEEP OUT OF REACH OF CHILDREN.
Avoid inhalation of dust.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of waste. Apply this product only as specified on this label.

DIRECTIONS FOR USE

Tri-Excel DS can be used undiluted as a dust or as a spray diluted with water at the rate of 8 tablespoonfuls per gallon.

AS A DUST Tri-Excel DS can be used in any type of hand or power duster. Apply a light film of dust to all parts of plants, especially the underside of leaves, before insects cause damage.

AS A SPRAY Tri-Excel DS can be used in most type hand or power sprayers. Apply the spray in such a manner as to thoroughly wet all parts of plants, especially the underside of leaves. Tri-Excel DS whether applied as a dust or a spray should be used at 7 to 14 day intervals and after heavy rains during the growing season.

ORNAMENTAL TREES AND SHRUBS such as Poplar, Elm, Dogwood, Birch, Buxus, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Laurel, Rose and Lilac.

For the control of Aphids, Leafhoppers, Oblique-banded Leaf Roller, Sawfly larvae, Boxelder Bug, Japanese Beetle, Elm Leaf Beetle, Cucumber Beetles, Rose Chater and Mites.

VEGETABLE PLANTS such as Pea, Bean, Beet, Cucumber, Squash, Muskmelon, Potato, Tomato, Cabbage, Spinach, Lettuce, Celery, Asparagus, Pepper, Eggplant, Onion, Carrot, and Cor.

CAUTION: Do not apply Tri-Excel DS within 1 day of harvest.

For the control of Aphids, Thrips, exposed, Leafhoppers, Cabbage worms, Bear Beetles, Beet Webworm, Jod Webworm, Asparagus Beetles, Japanese Beetles, Lygus Bugs, White Fly, Garden Fleaopper, Harlequin Bug, Oblique-banded Leaf Roller, Stink Bugs, Celery Leafminer, Melon worm, Tortoise Beetles, Garden Webworm, Alfalfa Webworm, Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bug, nymphs, and Squash Vine Borer. For European Corn Borer, 3 to 4 applications or more as eggs start to hatch and repeated at weekly intervals.

FLOWERING PLANTS such as Zinnia, Aster, Petunia, Geranium, Dahlia, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation, and Chrysanthemum.

For the control of Aphids, Thrips, exposed, Leafhoppers, Lygus Bugs, Lace Bugs, Greenhouse White Fly, Stink Bugs, Garden Fleahoppers, Japanese Beetle, Oblique-banded Leaf Roller, Greenhouse Leafminer, Cucumber Beetles, Aster Beetles, Flea Beetles, Ants, and Mites.

~~**FRUITS, BERRIES** such as Apple, Pear, Peach, Plum, Strawberry, Raspberry, and Blueberry.~~

CAUTION: Do not apply Tri-Excel DS within 1 day of harvest.

For the control of Aphids, Thrips, exposed, Leafhoppers, Imported Currant Worm, Gooseberry Fruit Worm, Raspberry Root Worm, adult, Blueberry Maggot, Cherry Fruit Fly, Strawberry Root Worm, adult, Red Necked Borer, Japanese Beetle, Strawberry Leaf roller, Rose Chater and Mites.

For Codling Moth Control: make first application at petal fall when 75% of blossom petals have fallen followed by three cover sprays at 10 to 14 day intervals. Spray applications for second and third treatments should be timed according to seasonal development. Information concerning the best control may be obtained from your State Agricultural Experiment Station.

USDA Reg. No. 8723-1

ACCEPTED
MAY 27 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
UNDER NO. 8723-1

Fred A. Voth
3505 Madison Avenue
Cincinnati, Ohio

water by cleaning of equipment,
or disposal of waste. Apply
this product only as specified
on this label.

DIRECTIONS FOR USE

Tri-Excel DS can be used undiluted as a dust or as a spray diluted with water at the rate of 8 tablespoonfuls per gallon.

AS A DUST: Tri-Excel DS can be used in any type of hand or power duster. Apply a light film of dust to all parts of plants, especially the underside of leaves, before insects cause damage.

AS A SPRAY: Tri-Excel DS can be used in most types of hand or power sprayers. Apply the spray in such a manner as to thoroughly wet all parts of plants, especially the underside of leaves. Tri-Excel DS whether applied as a dust or a spray should be used at 7 to 14 day intervals and after heavy rains during the growing season.

ORNAMENTAL TREES AND SHRUBS, such as Poplar, Elm, Dogwood, Willow, Birch, Boxelder, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Spirea, Rose and Lilac.

For the control of Aphids, Leafhoppers, Oblique-banded Leaf Roller, Sawfly larvae, Boxelder Bug, Japanese Beetle, Elm Leaf Beetle (larvae), Cucumber Beetles, Rose Chafer and Mites.

VEGETABLE PLANTS, such as Pea, Bean, Beet, Cucumber, Squash, Muskmelon, Potato, Tomato, Cabbage, Spinach, Lettuce, Celery, Asparagus, Pepper, Eggplant, Onion, Carrot, and Corn.

CAUTION: Do not apply Tri-Excel DS within 1 day of harvest.

For the control of Aphids, Thrips (exposed), Leafhoppers, Cabbage worms, Bean Beetles, Beet Webworm, Sod Webworm, Asparagus Beetles, Japanese Beetles, Lygus Bugs, White Fly, Garden Fleahopper, Harlequin Bug, Oblique-banded Leaf Roller, Stink Bugs, Celery Leafhopper, Melon worm, Tortoise Beetles, Garden Webworm, Alfalfa Webworm, Cucumber Beetles, Flea Beetles, Vegetable Weevil, Squash Bug (nymphs), and Squash Vine Borer. For European Corn Borer, 3 to 4 applications or more as eggs start to hatch and repeated at weekly intervals.

FLOWERING PLANTS, such as Zinnia, Aster, Petunia, Geranium, Dahlia, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation, and Chrysanthemum.

For the control of Aphids, Thrips (exposed), Leafhoppers, Lygus Bugs, Lace Bugs, Greenhouse White Fly, Stink Bugs, Garden Fleahoppers, Japanese Beetle, Oblique-banded Leaf Roller, Greenhouse Leafhopper, Cucumber Beetles, Aster Beetles, Flea Beetles, Ants, and Mites.

~~FRUITS, BERRIES, such as Apple, Pear, Peach, Strawberry, Raspberry, Blueberry, and Cherry.~~

CAUTION: Do not apply Tri-Excel DS within 1 day of harvest.

For the control of Aphids, Thrips (exposed), Leafhoppers, Imported Currant Worm, Gooseberry Fruit Worm, Raspberry Root Worm (adult), Blueberry Maggot, Cherry Fruit Fly, Strawberry Root Worm (adult), Red-Necked Borer, Japanese Beetle, Strawberry Leaf-roller, Rose Chafer and Mites.

For Codling Moth Control: Make first application at petal fall (when 75% of blossom petals have fallen) followed by three cover sprays at 1 to 2 week intervals. Spray applications for second and third brood worms should be timed according to seasonal development. Information concerning the best schedule may be obtained from your State Agricultural Experiment Station.

USDA Reg. No. 8723-1

ACCEPTED
MAY 27 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 8723-1 SUBJECT
TO ATTACHED CONDITIONS

Fred A. Veith
3505 Mozart Avenue
Cincinnati, Ohio