r (l)	1 (GH 40-1-1
US ENVIRONMENTAL PHOTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	TERM OF ISSUANCE
NOTICE OF PESTICIDE: [] REGISTRATION	NAME OF PESTICIDE PRODUCT
(Under the Federal Insecticide, Fungicide, and Rodenticide Act. a: amended)	
NAME AND ADDRESS OF REGISTRANT (Include ZIP code)	
MIAR, M.	7 161 754 487
F.O. tox 01 175 Irving, TV 7:601	17/1
L	_ // //
	′
NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.	
On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.	
A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.	
Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.	
This product is conditionally registered in accommode with FIDT section $\beta(\mathbf{c})(7)(K)$ provided that yith	
 Submit/cits all data required for registration/reregistration of your product under FIFPN section 3(c)(5) when the Agency requires all redistrants 	
of similar products to submit such data. 2. Add the thruse "EPA Degretration No. 40300-2" to your label before you release the product for shipment.	
3. Submit iin (5) copies of your final printed labeling select you release the product for support. Postr to the L-74 sold one for a further	
description of final oristed is eling. If these comists is are not complied with, the registration will be	
subject to cuncellation in accordance with FIFAA section 5(6). Your release for shipment of the problem constitutes accordance of these conditions. A stanged coping for large is emplosed for your records.	
A dealer to the fact of the property of the pr	
	il uttur Smurt (nem - (17)
I r.	sections = Is of the limit transfer quit ation live into the party
Enclosores	
ATTACHMENT IS APPLICABLE	
SIGNATURE OF APPROVING OFFICIAL	DATE
PA Form 8570-6 (Rev. 5-76) PREVIOUS EDITION	MAY BE USED UNTIL SUPPLY IS EXMAUSTED

THE TERMINATOR

TRANSPARENT EMULSION ** DILUTABLE CONCENTRATE 2%

Contains 0.17 lb/qal of SBP-1382 (20 g/L)
DUAL ACTION INSECT KILLER
KILLS AS SPACE AND RESIDUAL SPRAY
KILLS ROACHES UP TO 4 WEEKS

CTIVE INCREDIGHTS:

*Cls/trans isomers ratio:max.30Z(2)cls and min.70Z(2)trans. †U.S.Patent Hos.3,465,007 and 3,542,928. *Penick-Bio UCLAF'S SEP-1382° brand of resmethrin insecticide. Penick-Bio UCLAF Corp..1050 Wall Street West,Lyndhurst,M.J.07971 **U.S.Patent Me. 3,683,078.

REEP OUT OF REACH OF CHILDREN CAUTION

Mave clean water supply at hand in case of an eye emergency when diluting this concentrate. See side panel for additional Precautionary Statements and Statement of Pra tital Treatment

CAN BE USED IN MON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS.

Net Contents: EFA EST. NO. 40300-TX-1 EFA REG. NO.

MAINTRO. INC. P.O. NOX 610025 • IRVING, TX 75061 • (214) 438-2096

WARRANTT STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or im combination with other materials.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

OCT 24 1989

Under the rederal insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 403 00 - 8

PRECAUTIONARY STATEMENTS WARNING

BAZARDS TO HUMANS AND DURESTIC ARTHALS

Harmful if swallowed. Avoid contact with skin or ciothing. Causes eye irritation. Do not get in eyes. Wash after handling tool breathing vapor. Do not spray on humans. I apray where food or drink is exposed or bailing consumed.

Remove pets, birds, and cover fish squariums before space spraying. Do not remise in treated areas. Ventilate the area when treatment is complete

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting by sticking finger in back of throat or by administration of syrup of Ipacac. Gat medical attention.

IF IM EYES: Flush with water for 15 minutes. If irritation develops or presists, get medical attention.

FOR SKID: When with plenty of soap and vater. If irritation develos, get medical attention.

ENVIRORMENTAL BAZARDS

This product is textz to fish. Keep out of lakes, streams or ponds. Be not contaminate water by cleaning of equipment or disposal of wastes. When used on ships do not wash or flush freshly sprayed surfaces into adjacent waters.

MAYSICAL OR CHEMICAL MAZARDS DO NOT USE OR STORE MEAR HEAT OR OPEN FLAME. Plant point minima $150^{\circ} F(65.5^{\circ}C)$

DIRECTIONS FOR USE

It is a violation of Federal law to use this product
in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, foor or feed by storage or

STORAGE: Do not store this water-base formulation at temperatures below 32°F(0°C), there may be precipatation. Check for crystalization. If evident, warm to 80°F(26.5°C) and thoroughly mix before using, DO NOT USE OPEN FLAME. For containers 1 gallon and under: Store product in original container in a locked storage area.

PESTICIDE DISPOSAL: For Containers Over 1 gallon: Westeresulting from the use of this product may be disposed of on site or at approved waste disposal facility. CONTAINER DISPOSAL: For Containers ! Gallon and Under: Do not reuse empty container. Wrap container and put in trash. For Containers Over ! Gallon: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and Local suthorities.

DIRECTIONS CONT'D ON NEXT PANEL

For use in non-food __as of food handling astablishments, in and around n. mises of commercial, industrial, office, residential, public, instituctional and storage buildings, hotels, motels, homes, preschools, nurseries, day camps, seed houses, packaging and processing plants, wilk handling areas of milk plants, milk storage rounce egg processing plants, supertarkets, restaurants, bottling plants, canneries, flour and feed mills; baseries, boats, ship cabins and hospitals. IN MOSPITALS, patients should be removed prior to treatment. Also for use in campers, aphile, bomes, buses, automobiles, trucks, boxcars and saip campo, holds. Also for use in dog kannels and house in stables. IN HORSE STABLES; not for application if horses will be used for food.

DO NOT USE ON ELECTRICAL EQUIPMENT OR ON SURFACES THAT CAN BE HARMED O'S STAINED BY MATER.

Do not use in edible product areas of tood processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. In the home, cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces.

GENERAL INFORMATION: Use this product to control the pests indicated in the areas listed below by application as a 0.145%, 0.25% or 0.35% emulsion spray. For Contact and Space Sprays use either the 0.145%, 0.25% or 0.35% emulsion sprays. For Residual Sprays use the 0.35% emulsion.


IMPORTANT! To avoid gelling and assure good mixture.
THE TERMINATOR should always be added to water. Use per the following:

SPRAY DILUTION CHART Repeat the Treatment as Yeaded.

Amount of Spray P1.Czs. *SBP-1382 TEDC IX to use Mixture Desired 0.145% Spray 0.25% Spray 0.35% Spray 0.35% Spray 8 fl oz(237 ml) .06 (18 ml) 1 fl oz(20 ml) 19 fl oz(44 ml) 16 fl oz(473 ml) 1.2 (45 ml) 2 fl oz(59 ml) 3 fl oz(39 ml) 32 fl oz(946 ml) 2.4 (72 ml) 4 fl oz(118 ml) 6 fl oz(177 ml) 1 gal (1.69 L) 4.7 (144 ml) 8 fl oz(237 ml) 111 fl oz(340 ml) 1 gal (3.79 L) 9.6 (288 ml) 16 fl oz(473 ml) 23 fl oz(680 ml)

*MOTE: I fluid ounce = 1 tablespoonfuls = 5 teaspoons= 30 milliliters

VIRECTIONS FOR USE CONT'D UN REVERGE PROCE



AS A SPACE SPRAY FOR USE IN AUTOMATIC NOZZLE STSTEMS: Barrel should be empty, fill barrel 1/2-2/3 with water, add 4 gallons THE TERMINATOR agitate to mix well flush out container of THE TERMINATOR, real well into barrel, finish filling barrel with water.

CRAWLING INSECTS: To kill fless, cockrosches, water-bugs, crickers, silverfish; earwigs, spiders, sowbugs, tentipedes, millipedes, grain mites and grain inhects fadult and accessible stages of confused flouts; electles, saw-toothed grain beetles, cigaretre beetles; whiny spider beetles, carpet beetles, grain weevils, languagois grain moths). Apply as a contact spray. Bix well before using, Fill spray tank with spray dilution. Keep spray notale about 8 to 12 inches (20 to 30 cm) from surface to be treated. Adjust pressure and nozale to deliver a coarse spray with corticle size larger than that if the average zero-aoi. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, storage areas and other insect harborage areas. Spray directly on insects where possible. Repeat as nec-

TO RILL SPIDERS: Spray directly on these peats and use repeated applications to the webbing and all cracks and crevices which harbor spiders. TO RILL ANTS: Spray loors, around window frames,

ant trails and hills.
TO KILL BEDBUGS AND LICE:Apply as a spot treatment to cracke and cravices stound baseboards, floor-boards and wells. First, thoroughly clean and air mattresses and springs. Them spot treat only areas that may harbor peets, concentrating on tufts, folds and seams, epeat is necessary. For persistent infestations, consult professional applicator.

TO KILL FIEAS AND TICKS ON DOGS.CATS & REMISES: Apply directly to animals along back, under forelage, and on other areas except scrotum and amus. Part bein to contact skin and insure good coverage. Bold morele about 1 fr. (0.0m) away while spraying.Apply mutil hair is thoroughly moist.Do not spray directive in eyes or mouth of reinel.Repeat within 24 hours of first treatment, if necessary. To reduce infestation, choroughly spray the animal's bedding and other resting areas until the reface in slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches (10 to 26 cm). Also apray entire inside surface of dog houses. Repeat as necessary, thoroughly apraying floors, cracks, crevices and bedding. Bit as many insects directly as possible.

TO KILL WASPS, HORNETS, TELLOW JACKETS, BEES: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Repeat as necessary.

RESIDUAL SERAT: Aids in the control of COCKRCACHES. CRICKETS, SILVERFISH, ANTS, MILLIFEDES, CENTIPEDES, EAR-WIGS, GRAIN MITES, SOMBUGS, SPIDERS: Kills for up to 4 weeks. Spray baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back of and under appliances, sinks and and cupboards in kitchens and bathrooms. In basements, apray miding places in cracks and cravices, lower walls and dark places. Apply as a coarse wet apray in a 4 to 6 inch (10 to 15 cm) bank until area is thoroughly wet but not to the point of run-off. Repeat as nec-

DIRECTIONS CONT'D ON NEXT PANEL

CHOOCES

SPACE SPRAY: Remove or cover plants. Do not allow the spray to contact plants because of possible injury. FLYING INSECTS-To kill house flies, weertlies, stable files, gnats, cluster flies, vinegar flies, mosquitoes, fruit flies, flying mothstmillers):

Apply as a space spray. Mix well before using. Fill spray tank. Adjust preshure and mozzle to deliver a file mack, spray Close all doors and windows. Point nozzle upward. direct the spray misc to all patts of the race. especially windows and other light sources which attract these insects. Keep the spray mozzle at all times at least 4 ft. (1.2m) from fabrics to avoid wetting or staining. Fill the room with mist at the doesn, of I fl.oz. per 1000 cu. ft. of space (30 ml per 28.3 cu.m.). Then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the foom when treatment is completed and before rewentry?

OUTDOORS

To control nuisance peets around buildings.petics and paved areas. Apoly as a contact spray using either 0.5% of 0.35% spray dilution. Spray directly on insects avoid spraying Turf of Plants. TO KILL HOUSE FLIES.HOSQUITOES.GNATS (Midees; DEER FLIES.STABLE FLIES.CLUSTER FLIES.FLING MOTHS (Millers).FRUIT FLIES.Spray resting areas to reduce annowance from these peets.Foliow precautions listed above.Hold container 18 to 74 inches (46 to 01 cm) from the areas to be treated. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space (0.76 cu.m) while moving away from the treated areas.Treat open porches and patios in a similar manner.Spray only when air is still. Allow spray to dissipate before occupying the treated area. Papeat as necessary.

MASPS.HORNETS.YELLOW JACKETS. BEES:Same and directed above for indoor use.Application should be made in the late evening when insects are at rest Aim spray at nest openings in cracks and crevices which may harbor peets.saturating nest openings and coutacting as many insects as possible.

ARTS AND FLEAS: Same as directed above for indoor use.

CRICKETS, SILVERFISH, EARWIGS, SPIDERS, SOWBUGS, CINTIPEDES, AND MILLIPEDES: Same as directed above for indoor use.

FIRE ARTS: Rills and temporarily reduces annovance from these ants and keep them from entering homes, spray patios, betbecue areas, driveways, door sills. Window vills, other entrance areas, toundations of buildings, and trails, mounds, around pools and areas where the ants are noticed. Spray until the area is wer contacting as many insects as possible. This treatment will not kill the colonies but will reduce the immediate population for up to four nours. Repeat as necessary. Avoid contacting plants, shrubbery and large areas of lawns.

