


4375-4
PM 12
1/2



ACCEPTED
NOV 14 1978
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 11556-74



WARNING

Read entire label before using. Use strictly according to directions and limitations.
WARNING: May be fatal if swallowed. Do not get into eyes, on skin or clothing. Do not breath spray mist. Wash thoroughly after using. Avoid contamination of food and foodstuffs. Do not use around poultry, dogs or fish. Do not apply in dairy barns or on dairy animals. Do not use on animals intended for food within 30 days of slaughter. Do not Use, Pour, Spill, or Store Near Heat or Open Flame.

NOTICE

CAUTION: As a general precaution in spraying and dipping animals, avoid treating during cold, stormy weather; avoid treating overheated or sick animals, water animals before dipping to prevent them from drinking the dip. If calves must be treated, spray a minimum amount of material. Do not use on calves under 3 months.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

FIRST AID

INTERNAL: If swallowed, induce vomiting immediately. Give a tablespoon of salt in a glass of warm water and repeat until vomit is clear. Have victim lie down and keep quiet. Call a physician immediately.
EXTERNAL: If on skin, wash immediately and thoroughly with soap and water.

DIRECTIONS FOR USE

Spray-Dip Machines

Dilute TOXAPHENE-61 as directed. Follow manufacturer's directions for operation of machine. Tests indicate that recycling TOXAPHENE-61 emulsions does not cause the strength to weaken. The addition of a suitable wetting agent will greatly increase the wetting properties of the spray, thereby improving the effectiveness of the treatment

Dipping Vats

Use TOXAPHENE-61 in dipping vats at the dilution given in the directions. Always stir well just before using. Water animals before dipping to prevent them from drinking dip solution. Methods used for determining the dipping vat concentration of toxaphene emulsions may be used with TOXAPHENE-61. Do not dip calves under 3 months of age.

TOXAPHENE-61
LIVESTOCK SPRAY AND DIP

For Use on Cattle, Sheep, Goats and Hogs.
Controls Scabies (Sarcoptic and Chorioptic Mange and Psoroptic Scab).
Kills Ticks, Lice, Hornflies, Keds and Cures Sheep Scab.

Approved as a Permitted Pesticide by Animal and Health Inspection Service (APHIS) of the U.S. Department of Agriculture for the control of Scabies and Ticks in Federal Eradication Programs when used according to the directions of APHIS Veterinary Services Regulations and/or Memoranda

BEST AVAILABLE COPY

WARNING
KEEP OUT OF REACH OF CHILDREN
See Antidote and other Warnings on Side Panel

ACTIVE INGREDIENTS	
Toxaphene (Technical Chlorinated camphene containing 67.69% Chlorine)	61.0%
*Petroleum Distillate	27.0%
INERT INGREDIENTS	
	12.0%
	100.0%

*also known as deobase

EPA Reg. No. 11556-74 EPA Est. No. 2737 KS 1
DO NOT RE-USE THIS CONTAINER FOR ANY PURPOSE.

NET CONTENTS 5 GALLONS

Manufactured For

BEST AVAILABLE COPY

ACCEPTED

V 14 1978

Federal Insecticide,
and Rodenticide Act,
for the pesticide
under

No. 11556-74



TOXAPHENE-61

LIVESTOCK SPRAY AND DIP

For Use on Cattle, Sheep, Goats and Hogs.
Controls Scabies (Sarcoptic and Chorioptic
Mange and Psoroptic Scab).

Kills Ticks, Lice, Hornflies, Keds and Cures Sheep Scab.

Approved as a "Permitted Pesticide" by Animal and Health Inspection Service (APHIS) of the U.S. Department of Agriculture for the control of Scabies and Ticks in Federal Eradication Programs when used according to the directions of APHIS Veterinary Services Regulations and or Memoranda

WARNING

KEEP OUT OF REACH OF CHILDREN

See Antidote and other Warnings on Side Panel

ACTIVE INGREDIENTS

Toxaphene Technical Chlorinated
Camphene containing 61%
Chlorine

*Petroleum Emulsifier

NET INGREDIENTS

NET CONTENTS 5 GALLONS

Beef Cattle

To control Scabies, (Sarcoptic and Chorioptic Mange, Psoroptic Scab), Hornflies, Cattle Lice, Ear Tick, Lone Star Tick, Gulf Coast Tick, Winter Tick, and Cattle Fever Tick, mix 1 gallon of TOXAPHENE-61 with 150 gallons of water and stir well to form a creamy emulsion. Spray or dip animals to penetrate to the skin. Be sure not to overdose young animals. Repeat treatment for mange may be necessary and should be given in 10-14 days, if needed.

Back Rubber

As an aid in the control of Hornflies and Lice on beef cattle; dilute 1½ gallons of TOXAPHENE-61 with 34 gallons of fuel oil or other suitable carrier and fill insecticide compartment of back rubber device or soak sack or cloth as the occasion requires. Recharge device as necessary. Do not spray animals with this oil solution. Do not allow calves under 6 months of age to use rubbing devices. Do not allow beef animals to use backrubber for 30 days prior to slaughter.

Sheep and Goats

To control Psoroptic Sheep Scab, Fleece Worms and Ticks, dilute 1 gallon of TOXAPHENE-61 with 150 gallons of water and spray or dip animals thoroughly so that the solution penetrates to the skin. A second dipping may be given in 2 weeks if necessary. For control of Sheep Lice, Goat Lice and Sheep Ticks (Keds) use 1 gallon of TOXAPHENE 61 with 500 gallons of water and dip animals thoroughly. When using a power sprayer, a dilution of 1 gallon of TOXAPHENE-61 to 200 gallons of water will give better results.

Hogs

To control Sarcoptic Mange and Hog Lice, dilute 1 gallon of TOXAPHENE-61 with 150 gallons of water, mix well, and spray or wash animals thoroughly. Be sure not to overdose young animals. Repeat treatment for mange may be necessary and should be given in 10-14 days, if needed. Do not treat sows within 2 weeks of farrowing or at least 3 weeks thereafter.

NOTE: DO NOT USE ON CATTLE, SHEEP, GOATS, OR HOGS WITHIN 30 DAYS OF SLAUGHTER.

LIMITED WARRANTY AND LIMITATION OF DAMAGES

We warrant that at the time of manufacture this product was true to label and reasonably fit for the purposes described in its labeling. However, there are numerous factors beyond our control, e.g. storage, diagnosis, dosage, differences among individual animals, etc., which make it impossible to eliminate all risks associated with the use of this product, even if the directions are followed. This statement is made in lieu of any other warranty, either expressed or implied, including any warranty of merchantability or fitness, and no representative or agent of this company is authorized to vary the terms of this warranty or the contents of the labeling of this product except by printed notice from the company's Shawnee, Kansas head office. In the event of damages arising from our breach of this warranty, damages shall be limited to direct damages and shall not include consequential commercial damages such as loss of profit or values, etc.