

CRABGRASS PREVENTER

25-3-3



ACCEPTED
NOV 12 1985
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. *4515-5*

HERBICIDE

ACTIVE INGREDIENTS:

N-butyl-n-ethyl-alpha, alpha, alpha-trifluoro-2, 6-Dinitro-p-Toluidine 1.02%

INERT INGREDIENTS 98.98%

TOTAL 100.00%

BY WEIGHT

KEEP OUT OF REACH OF CHILDREN
CAUTION
See Back Panel for Additional Precautions

EPA Est. No.

NET WT. 18 LBS. (8.2 kg) TREATS 5,000 SQ. FT.

- previous fall.
3. ESTABLISHES A WEED CONTROL ZONE—Nutro "2 in 1" establishes a weed control zone by killing certain annual weed seeds as they germinate.
 4. WEEDS CONTROLLED—Application prior to weed germination will prevent smooth and hairy crabgrass, annual bluegrass (poa annua), goosegrass (silver crabgrass or crowfoot), watergrass (barleygrass) and yellow and green foxtail.
 5. FEEDS YOUR LAWN—Nutro "2 in 1" feeds your lawn a balanced diet of nitrogen, phosphorus, and potash.

WHEN TO APPLY

FOR CRABGRASS CONTROL: Apply in the spring before crabgrass germinates. Reseeding should be delayed for 90 to 120 days after application. To insure satisfactory reseeding results after using a pre-emergence chemical, the amount of seed should be increased.

FOR POA ANNUA (ANNUAL BLUEGRASS) CONTROL: Apply late to early fall (mid-July to mid-September) prior to annual bluegrass (poa annua) germination. Chemicals that effectively control poa annua seeds may thin established poa annua and fine-leaved fescues. Reseeding should be delayed 12 to 16 weeks after application.

HOW TO APPLY

Apply Nutro "2 in 1" LAWN FOOD PLUS CRABGRASS PREVENTER evenly over the turf area without skips or misses. A more uniform application can be made by spreading half the required amount of this product over the area, and then applying the remaining half at right angles or crosswise. Both drop-type and rotary-type spreaders may be used to apply.

Listed below are suggested spreader settings. These settings are only approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. This bag should cover approximately 5,000 Sq. Ft. (example, 50 ft. x 100 ft. = 5000 sq. ft.). Calculate the size of area to be treated and do not over or under apply. Close spreader when filling, stopping, or turning. DO NOT HAND SPREAD.

SUGGESTED SPREADER SETTINGS

Model	Recommended Two-Step Settings (half rate)	One step Settings (not recommended)
Central	6-6½	7-7½
Central Broadcast	3½	3¾-4
Cyclone	3½-3¾	4-4¼
Greenview	6-6½	7-7½
Ortho Drop	3½-4	4½-5
Ortho Broadcast	Not Recommended	1-1½
Scotts Drop	4¾-5¼	5¾-6¼
Scotts Broadcast	C-D½	E½-G
Seers Drop	5¼-6	6½-7
Seers Broadcast	3½	3¾-4

STORAGE AND DISPOSAL

Storage: Keep container closed and away from food, feedstuff, or domestic water supplies. Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bag. Discard bag in trash.

"Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with any directions provided, with proper equipment in good working order and under normal conditions of use. Buyer assumes the risk of any contrary use. Seller makes no other express or implied warranty including the warranty of fitness for a particular purpose, and no agent of seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made."

4 IN 1

CRABGRASS PREVENTER 25-3-3

GUARANTEED ANALYSIS

Total Nitrogen (N) 2:
 1.2% Ammoniacal Nitrogen
 23.8% Urea Nitrogen
 Available Phosphoric Acid (P₂O₅) :
 Soluble Potash (K₂O) :
 Primary Plant Foods Derived from Ammonium Phosphate, Urea, and Muriate of Potash.
 Potential Acidity 620 lbs. Calcium Carbonate Equivalent per ton.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact flush with water. Do not contaminate foodstuffs, or feed.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by direct application, cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

A selective pre-emergence herbicide for the control of crabgrass, annual bluegrass (poa annua), goosegrass other annual weed grasses in established:

BLUEGRASS (perennial)	FESCUE	BERMUDAGRASS	ST. AUGUSTINEGRASS
RYEGRASS (perennial)	BAHIAGRASS	CENTIPEDEGRASS	ZOYSIAGRASS

GENERAL INFORMATION

Nutrico "2 In 1" LAWN FOOD PLUS CRABGRASS PREVENTER has the following advantages:

1. CONVENIENT & ECONOMICAL—*Nutrico* "2 In 1" is on a convenient carrier, and can be used effectively in a rotary spreader as well as a drop-type spreader.
2. GOOD SAFETY MARGIN—Applied as directed, will not injure perennial Bluegrass, perennial ryegrass, Fescue, Bermudagrass, Zoysia, St. Augustinegrass, Centipede and Bahiagrass. Do not apply to new lawns started previous fall.
3. ESTABLISHES A WEED CONTROL ZONE—*Nutrico* "2 In 1" establishes a weed control zone by killing certain air weed seeds as they germinate.
4. WEEDS CONTROLLED—Application prior to weed germination will prevent smooth and hairy crabgrass, air bluegrass (poa annua), goosegrass (sliver crabgrass or crowfoot), watergrass (barnyardgrass) and yellow green foxtail.
5. FEEDS YOUR LAWN—*Nutrico* "2 In 1" feeds your lawn a balanced diet of nitrogen, phosphorous, and potash.

WHEN TO APPLY

FOR CRABGRASS CONTROL: Apply in the spring before crabgrass germinates. Re seeding should be delayed 10 to 120 days after application. To insure satisfactory re seeding results after using a pre-emergence chemical, the amount of seed should be increased.

FOR POA ANNUA (ANNUAL BLUEGRASS) CONTROL: Apply late to early fall (mid-July to mid-September) prior to air bluegrass (poa annua) germination. Chemicals that effectively control poa annua seeds may thin established poa annua and fine-leaved fescues. Re seeding should be delayed 12 to 16 weeks after application.

HOW TO APPLY

Apply *Nutrico* "2 In 1" LAWN FOOD PLUS CRABGRASS PREVENTER evenly over the turf area without skips or rills. A more uniform application can be made by spreading half the required amount of this product over the area, and applying the remaining half at right angles or crosswise. Both drop-type and rotary-type spreaders may be used as directed.

Listed below are suggested spreader settings. These settings are only approximate. Differences may occur due to condition of spreader, speed at which applied, and roughness of area treated. This bag should cover approximately 6,000 Ft. (example, 50 ft. x 100 ft. = 6000 sq. ft.). Calculate the size of area to be treated and do not over or under apply. Close spreader when filling, stopping, or turning. DO NOT HAND SPREAD.

SUGGESTED SPREADER SETTINGS

Model	Recommended Two-Step Settings (half rate)	One step Settings (not recommended)
Central	6-8 1/2	7-7 1/2
Central Broadcast	3 1/2	3 3/4-4
Cyclone	3 1/2-6 1/4	4-4 1/4
Greenview	6-8 1/2	7-7 1/2
Ortho Drop	3 1/2-4	4 1/2-6
Ortho Broadcast	Not Recommended	1-1 1/2
Scotts Drop	4 3/4-6 1/4	6 3/4-8 1/4
Scotts Broadcast	C-0 1/2	E 1/2-G
Sears Drop	5 1/4-6	6 1/2-7
Sears Broadcast	3 1/2	3 3/4-4

STORAGE AND DISPOSAL

Storage: Keep container closed and away from food, feedstuff, or domestic water supplies. Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bag. Discard bag in trash.

*Seller warrants that this material conforms to its chemical description and is accurately fit for the purposes stated on the label when used in a