

PM 25

44561-7

page 1 of 2

J.M. Magness
Nutrico, Inc.
70 Oak Ledge Lane
Wilton, CT 06897

Subject: Fertilizer With Team
EPA Reg. No. (s) 44561-7, -8, and -9
RE: Submission of labels bearing "directions for use".
Your submission dated February 12, 1993.

Dear Mr. Magness:

The above mentioned amendment submitted in connection with registration under FIFRA is accepted, however please refer to the comments listed below. A stamped copy is enclosed for your records.

1. The statement "contains 40% sulfur coated urea" must be removed from your label (44561-9) so that it will coincide with your previous label.
2. It will be assumed that the blank spaces left in front of the word "fertilizer" on each label will contain the appropriate number (guaranteed analysis) pertaining to that particular label.
3. Update the "Environmental Hazard" statements to read; "Do not contaminate water when disposing of equipment washwaters."

Sincerely,

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C						
SURNAME	HICKS						
DATE	2/22/93						

20-5-10 FERTILIZER WITH

A GRANULAR FERTILIZER PLUS SELECTIVE PREEMERGENCE HERBICIDE FOR THE CONTROL OF CRABGRASS AND MOST OTHER ANNUAL WEED GRASSES ON LAWNS AND GOLF COURSES IN ESTABLISHED:

BLUEGRASS (PERENNIAL) BERMUDAGRASS
 BENTGRASS FAIRWAYS RYEGRASS (PERENNIAL)
 ST. AUGUSTINEGRASS ZOYSIAGRASS
 FESCUE CENTIPEDEGRASS
 BAHIAGRASS

ACTIVE INGREDIENTS

N-butyl-N-ethyl-p,p'-dithioure-2,6-dinitro-p-toluidine	0.813%
s,s'-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.306%
INERT INGREDIENTS	99.081%
TOTALS	100.000%



TEAM and the TEAM logo are trademarks of DowElanco

Total Nitrogen (N)	20.0%
1.9% ammoniacal nitrogen	
12.0% water soluble nitrogen	
6.0% coated slow release nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	5.0%
Soluble Potash (K ₂ O)	10.0%
Derived from urea, sulfur coated urea, ammoniated phosphates and muriate of potash.	

WARNING
 KEEP OUT OF THE REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: CAUSES SUBSTANTIAL BUT TEMPORARY EYE IRRITATION. DO NOT GET IN EYES OR ON CLOTHING. WEAR GOGGLES, FACE SHIELD, OR SAFETY GLASSES. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING. REMOVE CONTAMINATED CLOTHING AND WASH BEFORE REUSE. AVOID BREATHING DUST AND CONTACT WITH SKIN. HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: CALL A PHYSICIAN OR POISON CONTROL CENTER. HOLD EYELIDS OPEN AND FLUSH WITH A STEADY, GENTLE STREAM OF WATER FOR 15 MINUTES.
 IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER. PROMPTLY DRINK A LARGE QUANTITY OF MILK, EGG WHITE, OR GELATIN SOLUTION, OR IF THESE ARE NOT AVAILABLE, LARGE QUANTITIES OF WATER. AVOID ALCOHOL.
 IF ON SKIN: WASH WITH PLENTY OF SOAP AND WATER. GET MEDICAL ATTENTION IF IRRITATION PERSISTS.
 IF INHALED: IF BREATHING DISCOMFORT OCCURS MOVE INDIVIDUAL TO FRESH AIR. GET MEDICAL ATTENTION IF BREATHING DIFFICULTY OCCURS.
 ENVIRONMENTAL HAZARDS: DO NOT CONTAMINATE ANY BODY OF WATER BY DIRECT APPLICATION, CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

STORAGE AND DISPOSAL

STORAGE: STORE IN ORIGINAL CONTAINER ONLY. IN CASE OF SPILL CONTAIN MATERIAL AND DISPOSE AS WASTE.
 PESTICIDE DISPOSAL: DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE AND DISPOSAL. WASTES RESULTING FROM USE OF THIS PRODUCT MAY BE DISPOSED OF ON SITE OR AT AN APPROVED WASTE DISPOSAL FACILITY.

CONTAINER DISPOSAL: COMPLETELY EMPTY BAG INTO APPLICATION EQUIPMENT. THEN DISPOSE OF EMPTY BAG IN A SANITARY LANDFILL OR BY INCINERATION OR, IF ALLOWED BY STATE AND LOCAL AUTHORITIES, BY BURNING. IF BURNED STAY OUT OF SMOKE.

BEST AVAILABLE CONTROL

READ ALL DIRECTIONS BEFORE USING

FEDERAL LAW PROHIBITS USE OF THIS PRODUCT IN ANY MANNER INCONSISTANT WITH LABEL DIRECTIONS. FERTILIZER WITH TEAM CONTAINS A PREEMERGENCE HERBICIDE FOR THE CONTROL OF ANNUAL BLUEGRASS (POA ANNUA), GOOSEGRASS, BAHIAGRASS, SMOOTH AND Hairy CRABGRASS AND GREEN AND YELLOW FOXTAIL. IT CONTROLS THESE ANNUAL WEEDS BY KILLING THEIR SEEDS AS THEY GERMINATE AND MUST BE APPLIED BEFORE THESE SEEDS GERMINATE. THIS PRODUCT DOES NOT CONTROL ESTABLISHED WEEDS. THIS PRODUCT WILL NOT HARM ESTABLISHED ORNAMENTALS.

APPLICATION DIRECTIONS

TIMING OF APPLICATION IS CRITICAL FOR SUCCESSFUL PREEMERGENCE CONTROL OF ANNUAL GRASS WEEDS. APPLY FERTILIZER WITH TEAM APPROXIMATELY ONE TO TWO WEEKS BEFORE GERMINATION OF GRASS WEEDS. BEST CONTROL WILL BE ACHIEVED IF TREATED AREAS ARE IRRIGATED SOON AFTER APPLICATION. ADDITIONALLY, ANY SOIL DISRUPTION (VERTICUTTING, CORING OR AERIFICATIONS) SHOULD BE COMPLETED PRIOR TO APPLICATION OF THIS PRODUCT TO AVOID BREAKUP OF THE VAPOR BARRIER.

ANNUAL GRASSES - CRABGRASS - FOXTAIL AND BAHIAGRASS

NORTHERN (COOL SEASON) TURFGRASSES

APPLY 165 TO 215 LBS. PER ACRE (1.5 TO 2 LBS. AM OF FERTILIZER WITH TEAM ONE TO TWO WEEKS PRIOR TO THE GERMINATION OF ANNUAL GRASS SEEDS. THIS RANGE WILL RESULT IN .75 TO 1.0 LBS. OF NITROGEN PER 1000 SQ. FT.

IN GEOGRAPHIC AREAS WHERE GERMINATION OF ANNUAL GRASS WEEDS CONTINUES FOR EXTENDED PERIODS OF TIME A SECOND APPLICATION AT THE RECOMMENDED RATE MAY BE MADE 9 TO 10 WEEKS AFTER THE INITIAL APPLICATION FOR CONTINUED CONTROL.

LATE SPRING - EARLY SUMMER GERMINATING ANNUAL GRASSES
 GOOSEGRASS
 GOOSEGRASS NORMALLY GERMINATES 4 TO 6 WEEKS LATER THAN FOXTAIL, CRAB OR OTHER ANNUAL GRASS AND APPLICATIONS SHOULD THEREFORE BE TIMED ACCORDINGLY TO REFLECT THIS VARIATION.

NORTHERN (COOL SEASON) TURFGRASSES

APPLY 215 LBS. OF FERTILIZER WITH TEAM PER ACRE IN EARLY SPRING OR EARLY SUMMER ONE TO TWO WEEKS PRIOR TO GERMINATION OF GOOSEGRASS SEEDS.

NORTHERN - SOUTHERN TURFGRASSES

IN GEOGRAPHIC AREAS WHERE PREEMERGENCE CONTROL OF ANNUAL GRASSES (CRABGRASS, FOXTAIL, AND BAHIAGRASS) IS NEEDED, LATER GERMINATING GOOSEGRASS IS NECESSARY. STRONG APPLICATIONS OF THIS PRODUCT SEVERAL WEEKS APART TO PROVIDE FULL LENGTH SEASONAL CONTROL.

POA ANNUA (ANNUAL BLUEGRASS)

THE ELIMINATION OF HEAVILY INFESTED AREAS OF POA ANNUA IN TIGHTLY MAINTAINED TURFGRASS. PROLONGED IRRIGATION AND RESEEDING WILL BOTH HELP ERADICATE POA AS WELL AS HELP TO FILL IN THESE THINNED AREAS.

NORTHERN (COOL SEASON) TURFGRASSES

APPLY 215 LBS. OF FERTILIZER WITH TEAM PER ACRE IN EARLY FALL AND AGAIN IN THE SPRING ONE TO TWO WEEKS PRIOR TO GERMINATION OF POA ANNUA (ANNUAL BLUEGRASS) SEEDS.

ROTARY SPREADER SETTINGS

NOTE: THESE SETTINGS ARE APPROXIMATE AND VARY WITH THE AGE AND CONDITION OF THE EQUIPMENT. ALWAYS CALIBRATE SPREADER WITH EACH APPLICATION.

	4 LBS. PER 1000 SQ. FT.
SCOTT'S PRO BROADCAST	0 1/2
CYCLONE	5
LESCO	0 1/2
SPYER	5
EARTHWAY	15 1/2

RESEEDING

THE CHEMICALS USED TO FORMULATE TEAM ARE PREEMERGENCE CONTROL OF ANNUAL GRASS WEEDS. ALSO AFFECT NEW DESIRABLE TURFGRASS SEEDS. RESEEDING SHOULD BE DELAYED AT LEAST 8 WEEKS AFTER APPLICATION OF FERTILIZER WITH TEAM. IT IS IMPORTANT TO FOLLOW CULTIVATION AND IRRIGATION PRACTICES AFTER APPLICATION TO INCREASE THE NORMAL AMOUNT OF TURFGRASS THAT HAVE BEEN TREATED WITH PREEMERGENCE CONTROL IN THE SPRING TO TURFGRASS AREAS THAT WERE TREATED PREVIOUS FALL.

PRECAUTION

IN BERMUDAGRASS AREAS THAT HAVE BEEN OVERSEED WITH A SPRING APPLICATION OF PREEMERGENCE CONTROL, THE OVERSEEDING AREAS. DUE TO VARIATION IN PERSISTENCE OF PREEMERGENCE CONTROL ON CENTIPEDEGRASS UNLESS THESE AREAS ARE RESEEDED.

E.P.A. REG. NO. 44581-7 E.P.A. REG. NO. 44581-7

COVERS 10,000 SQUARE FEET

NET WEIGHT 50 POUNDS

MANUFACTURED BY:
 NUTRICO, INC. 92 HOWARD AVENUE BOSTON, MASSACHUSETTS 02118