

J.M. Magness
Nutrico, Inc.
70 Oak Ledge Lane
Wilton, CT 06897

Subject: Fertilizer With Team
EPA Reg. No.(s) 44561-7, ~~8~~, and -9
RE: Submission of labels bearing "directions for use".
Your submission dated February 12, 1993.

Dear Mr. Magness:

The above mentioned amendment submitted in connection with registration under FIFRA is accepted, however please refer to the comments listed below. A stamped copy is enclosed for your records.

1. The statement "contains 40% sulfur coated urea" must be removed from your label (44561-9) so that it will coincide with your previous label.
2. It will be assumed that the blank spaces left in front of the word "fertilizer" on each label will contain the appropriate number (guaranteed analysis) pertaining to that particular label.
3. Update the "Environmental Hazard" statements to read; "Do not contaminate water when disposing of equipment washwaters."

Sincerely,

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C						
SURNAME	HICKS						
DATE	2/22/93						

25-3-9 FERTILIZER WITH

CONTAINS 50% SULFUR COATED UREA

A granular fertilizer plus selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

- | | | | |
|-----------------------|----------------------|--------------|----------------|
| Bluegrass (Perennial) | Fescue | Bermudagrass | Centipedegrass |
| Bentgrass Fairways | Eyegrass (Perennial) | | Bahiagrass |
| St. Augustinegrass | Zoysiagrass | | |

GUARANTEED ANALYSIS

Total Nitrogen (N)..... 25.0%

1.17% Ammoniacal Nitrogen
11.33% Urea Nitrogen
12.50% Coated Slow Release Nitrogen

Available Phosphoric Acid (P₂O₅)..... 1.0%

Soluble Potash (K₂O) 3.0%

Derived from Sulfur Coated Urea, Urea, Ammonium Phosphate and Muriate of Potash.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-a,a,a-trifluoro-2,6-dinitro-p-toluidine... 0.766%

a,a,a-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine..... 0.383%

Inert Ingredients..... 98.851%

BEST AVAILABLE COPY

READ ALL DIRECTIONS BEFORE USING

Federal law prohibits use of this product in any matter inconsistent with label directions. With Team is a preemergence herbicide for the control of annual bluegrass (poa annual), goosegrass, barnyardgrass, smooth and hairy crabgrass and green and yellow foxtail. It controls these annual weeds by killing their seeds as they germinate and must be applied before these weeds germinate. This product does not control established weeds. This product will not harm established ornamentals.

APPLICATION DIRECTIONS

Timing of application is critical for successful preemergence control of annual grass weeds. Apply Fertilizer with Team approximately one to two weeks before germination of grass weeds. Best control will be achieved if treated areas are irrigated soon after application. Additionally, any soil disruption (verticutting, coring or aeration) should be completed prior to application of this product to avoid breakup of the vapor barrier.

Annual Grasses-Crabgrasses-Foxtail and Barnyardgrass

Northern (cool season) turfgrasses
Apply 174 pounds per acre (1.5 to 2 lbs. aia) of fertilizer with Team one to two weeks prior to the germination of annual grass seeds. This range will result in .75 to 1.8 lbs. of nitrogen per 1000 sq.ft.

In geographic areas where germination of annual grass weeds continues for extended periods of time a second application at the recommended rate may be made 8 to 10 weeks after the initial application for continued control.

NET WEIGHT 50 POUNDS

Covers 12,500 square feet.



*TEAM and trademarks of

late Spring-
Goosegrass
Goosegrass a
foxtail, cra
should there
variation...
Northern (co
Apply 174
Team per ac
to two week
Northern-Sou
In geographi
early annual
grass) and t
two "all sta
rate : to 10
control.

roo Annual (o
The eliminat
will result
ilization, a
ine turfgras
areas with
Northern (c
Apply 174
Team per ac
the spring
annual (annu
NOTE

Note: These
affected by
rent. Be t
with each

Scotts Pro
Cyclone
Lesco
Spykar

The chem
in the p
therefor
seedling
weeks af
Team. I
and irri
advisabl
used in
Do not
were see

PRECAUTI
In Berm
winter c
may res
Due to
bermuda
unless

E.P.A. 1
..

WARNING: KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing dust and contact with skin. Harmful if swallowed, inhaled, or absorbed through the skin.

STATEMENT OF PRACTICAL TREATMENT

In eyes: Call a physician or poison control center. Hold eyelids open and flush with a gentle stream of water for 15 minutes.

Swallowed: Call a physician or poison control center. Promptly drink a large quantity of milk, egg white, or gelatin solution, or if these are not available, large quantities of water. Avoid alcohol.

On skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Inhaled: If breathing discomfort occurs move individual to fresh air. Get medical attention if breathing difficulty occurs.

Environmental hazards: Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

STORAGE and DISPOSAL

Storage: Store in original container only. In case spill contain material and dispose as waste.
Toxicity Disposal: Do not contaminate water, food or feed by storage and disposal. Wastes resulting from use of this product may be disposed of on site at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning.

FEE 29

44561-8