

J.M. Magness
Nutrico, Inc.
70 Oak Ledge Lane
Wilton, CT 06897

Subject: Fertilizer With Team
EPA Reg. No. (s) 44561-7, -8, and ~~9~~
RE: Submission of labels bearing "directions for use".
Your submission dated February 12, 1993.

Dear Mr. Magness:

The above mentioned amendment submitted in connection with registration under FIFRA is accepted, however please refer to the comments listed below. A stamped copy is enclosed for your records.

1. The statement "contains 40% sulfur coated urea" must be removed from your label (44561-9) so that it will coincide with your previous label.
2. It will be assumed that the blank spaces left in front of the word "fertilizer" on each label will contain the appropriate number (guaranteed analysis) pertaining to that particular label.
3. Update the "Environmental Hazard" statements to read; "Do not contaminate water when disposing of equipment washwaters."

Sincerely,

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C							
SURNAME	HICKS							
DATE	2/22/93							

IN ESTABLISHED:

Bluegrass (Perennial)
Bentgrass Fairways
St. Augustinegrass

Poa
Bermudagrass
Ryegrass (Perennial)
Zoysiagrass

Centipedegrass
Bahingrass



GUARANTEED ANALYSIS

Total Nitrogen (%)..... 30.00
1.960 Ammoniacal Nitrogen
16.040 Urea Nitrogen
12.000 Coated Slow Release Nitrogen
Available Phosphoric Acid (P₂O₅)..... 5.00
Soluble Potash (K₂O)..... 7.00

Derived from Sulfur Coated Urea, Urea, Ammonium Phosphate, Salts of Potash.

WARNING: KEEP OUT OF REACH OF CHILDREN

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid breathing dust and contact with skin. Harmful if swallowed, inhaled, or absorbed through the skin.

STATEMENT OF PRACTICAL TREATMENT

In eyes: Call a physician or poison control center. Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes.
If swallowed: Call a physician or poison control center. Promptly drink a large quantity of milk, egg white, or gelatin solution, or if these are not available, large quantities of water. Avoid alcohol.
On skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
If inhaled: If breathing discomfort occurs move individual to fresh air. Get medical attention if breathing difficulty occurs.
Environmental hazards: Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

STORAGE and DISPOSAL

Storage: Store in original container only. In case of spill contain material and dispose as waste.
pesticide Disposal: Do not contaminate water, food or feed by storage and disposal. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. Do not burn if burned stay out of smoke.

CONTAINS 10% SULFUR COATED UREA...

ACTIVE INGREDIENTS

N-butyl-N-ethyl-s,s-trifluoro-2,6-dinitro-p-toluidine... 0.9190
s,s-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine..... 0.4590
Inert Ingredients..... 98.6220

BEST AVAILABLE COPY

READ ALL DIRECTIONS BEFORE USING

Federal law prohibits use of this product in any manner inconsistent with label directions. Team is a preemergence herbicide for the control of annual bluegrass (poa annual), goosegrass, barnyardgrass, smooth and hairy crabgrass and green and yellow foxtail. It controls these annual weeds by killing their seeds as they germinate and must be applied before these weeds germinate. This product does not control established weeds. This product will not harm established ornamentals.

APPLICATION DIRECTIONS

Timing of application is critical for successful preemergence control of annual grass weeds. Apply Fertilizer with Team approximately one to two weeks before germination of grass weeds. Best control will be achieved if treated areas are irrigated soon after application. Additionally, any soil disruption (verticutting, coring or aeration) should be completed prior to application of this product to avoid breakup of the vapor barrier.

Annual Grasses-Crabgrasses-Foxtail and Barnyardgrass

Northern (cool season) turfgrasses
Apply 100 to 145 pounds per acre (1.5 to 2 lbs. aia) of fertilizer with Team one to two weeks prior to the germination of annual grass seeds. This range will result in .75 to 1.0 lbs. of nitrogen per 1000 sq. ft.

In geographic areas where germination of annual grass weeds continues for extended periods of time a second application at the recommended rate may be made 8 to 10 weeks after the initial application for continued control.

NET WEIGHT 50 POUNDS

Covers 15,000 square feet
Spread at 3 lbs. per 1000 square feet

FFB 22 1002
Under
Fungus
ES
REG
44561-9

Bluegrass
foxtail, c
annual, the
varieties:
Northern (c
Apply 100
pounds per
to 145 lbs
Northern (c
In geographic
early (anne
grass) and
two full c
made 8 to
control.

Poa Annual
for elimin
will resul
il action,
in turfgr
areas with
Northern (c
Apply 100
Team per a
the spring
annual (ann

Note: Th
affected
ment. Be
with each

Scotts Pr
Cyclone
Lasco
Spiker
Do
The ch
in the
therefo
seedlin
weeks i
Team.
and ir
advised
used i
Do not
were s
PRECAU
In bet
winter
may re
Due to
bermud
unless
E.P.A.
E.P.A.
Read label