

Mr. Ralph P. Burton
ISK Biotech Corporation
P.O. Box 8000
Mentor, OH 44061-8000

Dear Mr. Burton:

Subject: Add Peanut Tank Mix Directions with Solubor and Surfix
Bravo 90DG
EPA Registration No. 50534-157
Your Submission Dated November 5, 1992

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Revise the precautionary statements to include the following statements:

Wear goggles, face shield or safety glasses.

b. In the note to user under the directions for use section include the requirement for goggles, face shield, or safety glasses. When you chose to repeat the personal protective clothing and equipment requirements, all required personal protective clothing and equipment must be listed. Goggles, face shield, or safety glasses must not only be worn when handling this product but also when applying it.

c. We note the way the tank mix directions are written they imply that Surfix must always be added to the tank mix. Based on our discussions we are not sure that is what you intended.

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

-2-

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

151

Cynthia Giles-Parker
Product Manger (22)
Fungicide Herbicide Branch
Registration Division (H7505C)

Enclosure

ACCEPTED
with COMMENTS
In EPA Letter Dated

Under the Federal Insect
Fungicide, and Rodenticide
Act as amended, for the pest
registered under EPA Reg.

Net Contents: 25 Pounds

ISK BIOTECH™

Bravo® 90 DC

Agricultural Fungicide
(90% Water Dispersible Granules)

Active Ingredient: Chlorothalonil (tetrachloroisophthalonitrile)*	90.0%
Inert Ingredients:	10.0%
Total:	100.0%

Keep Out of Reach of Children
DANGER-PELIGRO

See side panel for additional precautionary statements.

Precaucion al Usuario:

Si usted no lee ingles, no use este producto
que la etiqueta haya sido explicado ampliamente.

ISK Biotech Corporation
5968 Heisley Road
P.O. Box 8000
Mentor, Ohio 44061-8000

BRAVO is a registered trademark of ISK Biotech Corporation.

EPA Reg. No. 5
EPA Est. No. 37

3298-3038-1

Bravo 90DG

A Broad Spectrum Agricultural Fungicide

Read entire label carefully and use only as directed.

Precautionary Statements

Hazards to Humans and Domestic Animals

Danger

Corrosive, causes severe eye damage. May cause skin irritation. May be a potential skin sensitizer.

Do not get in eyes. Wear goggles or face shield when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Avoid prolonged contact with skin.

Avoid breathing spray mist.

Do not take internally.

Note to User: This product may produce mild bronchial irritation and temporary irritation of the skin characterized by redness or rash on exposed skin areas. Affected persons should consult a physician.

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

First Aid:

In case of contact with eyes, immediately flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes immediately.

Wash exposed areas of skin with soap and warm water after handling or using.

Note to Physician: Persons having temporary irritation symptoms may respond to treatment with antihistamines or steroid creams and/or systemic steroids.

Environmental Hazards

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

4/2/12

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Note to User: Wear goggles or face shield when handling this product. Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

Brave 900G is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases.

Brave 900G can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

Do not use on greenhouse-grown crops.

Do not combine Brave 900G in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. Do not combine Brave 900G with Dipel 4L, Foli, Triton AG-98, Triton B-1956, Latron B-1956 or Latron AG-98 as phytotoxicity may result from the combination when applied to some species on this label.

Dipel is a registered trademark of Abbott Laboratories. Foli is a registered trademark of Ecogen Corporation.

Triton AG-98, Triton B-1956, Latron B-1956 and Latron AG-98 are trademarks of Fohm and Haas Company.

Dosage rates on this label indicate pounds of Brave 900G per acre, unless otherwise stated. Under conditions favoring disease development the high rate specified and shortest application interval should be used.

The required amount of Brave 900G should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of Brave 900G in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

Reentry:

Note to User: Do not enter treated area to perform hand labor within 24 hours of application unless protective clothing is worn.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be

given if there is reason to believe that written warnings cannot be understood by workers. Oral and written warnings must include the following information: "DANGER. Area treated with Brave 900G on (date of application). Do not enter without appropriate protective clothing for 24 hours. In case of accidental exposure, wash exposed area with plenty of water and get medical attention. For further information see 'Precautionary Statements' on the label."

Field and Row Crops:

Apply Brave 900G in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See application and calibration instruction below.

5
2
12

Application and Calibration Techniques for Sprinkler Irrigation

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set and portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue

pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject BRAVO 900G into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Posting of areas to be chemigated is required when: 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads; or 2) when the chemigated area is open to the public.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in

(Continued on back of bag)

6412

and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2 - 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required,

can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of Bravo 900G for acreage to be covered with water so that the total mixture of Bravo 900G plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Agitation is recommended. Bravo 900G can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until Bravo 900G has been cleared from last sprinkler head.

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Bean (Snap)	Rust	1-1/8 to 2-1/2 lbs	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat at weekly intervals or as necessary to maintain control. Do not apply within 7 days of harvest. Do not graze treated areas or feed treated plant parts to livestock.
	Botrytis blight (gray mold)	2-1/2 lbs	
Beans (Dry)	Rust, Anthracnose, Downy mildew, Cercospora leaf spot (blackeye only)	1-1/8 to 1-3/4 lbs	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage and repeat at 7 to 10 day intervals. For use only on beans harvested dry with pods removed. Do not apply within 6 weeks before harvest. Do not allow livestock to graze in treated areas or feed treated plant parts to livestock.
Cabbage, Chinese Cabbage (tight-headed varieties only), Cauliflower, Broccoli, Chinese Broccoli, Brussels Sprouts	Alternaria leaf spot, Downy mildew	1-1/4 lbs	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control. Do not apply within 7 days of harvest to Chinese cabbage or Chinese broccoli.
	Ring spot (California only)	1-1/2 lbs	
Carrot	Cercospora (Early) blight, Alternaria (Late) blight	1-1/4 to 1-1/2 lbs	Use in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals or as necessary to maintain control. Bravo 900G may be applied through sprinkler irrigation equipment. See calibration directions preceding this section.
Celery	Cercospora (Early) blight, Septoria (Late) blight	7/8 to 1-1/4 lbs	Use 7/8 to 1-1/4 lb per acre on a 3 to 5 day spray schedule or 1-3/4 to 2-1/2 lb per acre on a 7 day schedule. Start applications when transplants are set in the field. Apply in sufficient water to obtain adequate coverage. Do not apply within 7 days of harvest. Bravo 900G may be applied through sprinkler irrigation equipment. See calibration directions preceding this section.
	Preal stalk rot (Septoria solani)	1-3/4 to 2-1/2 lbs	
	Early blight, Late blight	1-1/4 to 1-1/2 lbs per 100 gal	

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS												
Corn (sweet), Corn grown for seed	Helminthosporium leaf blights, Rust	5/8 to 1-1/2 lbs	Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at 4 to 7 day intervals or as required to maintain control. Under severe disease conditions, use 1-1/4 to 1-1/2 lb per acre. Do not apply within 14 days of harvest. Do not apply to sweet corn to be processed. Do not allow livestock to graze in treated fields. Do not ensile treated corn or use as livestock forage.												
Cranberry	Fruit rots, Lophodermium leaf/twig blight	3-1/2 to 5-3/4 lbs	Apply at early bloom and repeat at 10 to 14 day intervals. Under severe disease conditions use the 5-3/4 lbs/acre rate on a 10 day schedule. Do not apply Bravo 900G more than 3 times per season, or within 50 days before harvest. Do not apply to bogs when flooded or allow release of irrigation water from bogs for at least 3 days following application. Bravo 900G may be applied through sprinkler irrigation equipment. Use 300 gallons of water per acre through solid set systems only. See calibration directions preceding this section.												
Cucurbits: Cucumber, Cantaloupe, Muskmelon, Honeydew melon, Watermelon, Squash, Pumpkin	Anthracnose, Downy mildew, Target spot	1-1/4 to 1-1/2 lbs	Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe disease conditions, shorten spray interval. Bravo 900G may be applied through sprinkler irrigation equipment (solid set, portable wheel move or center pivot systems only). See calibration directions preceding this section.												
	Cercospora leaf spot, Gummy stem blight (black rot), Alternaria leaf blight, Scab, Powdery mildew (Sphaerotheca only)	1-1/2 to 2-1/2 lbs													
	Cucumber belly rot (<i>Rhizoctonia solani</i>)	7 lbs		Use Bravo 900G in sufficient water to obtain runoff to soil surface. Make a single application when vines begin to form. Bravo 900G may be applied through sprinkler irrigation equipment as directed above.											
Grasses Grown for Seed	Stern rust, Leaf rust, Stripe rust, Septoria leaf spot, Glume blotch, Bipolaris and Drechslera leaf spots	7/8 to 1-1/8 lbs	Use in sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat applications at 14 day intervals. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated areas or feed treated plant parts to livestock												
	Selenophoma (eyespot)	7/8 to 1-3/4 lbs													
Mint	Rust, Septoria leaf spot	1-1/8 lbs	Use in sufficient water to obtain adequate coverage. Begin applications when emerging plants are 4-8 inches high. Repeat applications at 7 to 10 day intervals or as necessary to maintain control. Do not apply more than 3 times per season. Do not apply within 60 days of harvest. Do not feed fresh or extracted mint hay from treated fields to livestock. Based on available residue data, use of Bravo 900G on mint is restricted to Indiana, Michigan and Wisconsin.												
Onion (dry bulb)	Botrytis leaf blight/ blast, Botrytis neck rot (suppression), Purple blotch		<p>Apply in sufficient water to obtain thorough coverage of tops. Bravo 900G is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows:</p> <table border="1"> <thead> <tr> <th></th> <th>Low Disease Hazard & Prior to Infection</th> <th>Low Disease Hazard & Some Disease Present</th> <th>High Disease Hazard</th> </tr> </thead> <tbody> <tr> <td>Rate per Acre:</td> <td>7/8 lb</td> <td>1-1/8 lbs</td> <td>1-3/4 lbs</td> </tr> <tr> <td>Frequency:</td> <td>10 days</td> <td>7 to 10 days</td> <td>7 days</td> </tr> </tbody> </table> <p>For suppression of neck rot (<i>Botrytis</i> spp.) during storage, a minimum of three weekly applications prior to lifting, using 1-1/8 to 1-3/4 lbs of Bravo 900G per acre, is recommended. Do not apply within 7 days of harvest.</p>		Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard	Rate per Acre:	7/8 lb	1-1/8 lbs	1-3/4 lbs	Frequency:	10 days	7 to 10 days	7 days
	Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard												
Rate per Acre:	7/8 lb	1-1/8 lbs	1-3/4 lbs												
Frequency:	10 days	7 to 10 days	7 days												
Onion (green bunching), Leek, Shallot, Garlic, Onion grown for seed	Botrytis leaf blight/ blast, Purple blotch, Downy mildew (suppression)	1-1/8 to 2-1/4 lbs	Use in sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease. Use the high rate and a 7 day schedule of applications when heavy dew or rain persists. Do not apply within 7 days of harvest on garlic. Do not apply more than 3 times per season or within 14 days of harvest on green bunching onions, leeks or shallots. If additional disease control is needed before harvest, use another registered fungicide												

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Papaya	Alemaria fruit spot, Anthracnose, Stem and rot	1-3/4 to 3-1/4 lbs	Apply with ground equipment only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development. Do not graze livestock in treated area or feed processing by-products to livestock.
Parsnip	Alemaria leaf spot, Downy mildew, Anthracnose, Botrytis blight (gray mold), Bottom rot (Rhizoctonia)	1-1/8 to 1-3/4 lbs	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule. Do not apply more than 4 times per season or within 10 days of harvest. Do not feed treated plant parts to livestock.
Passion Fruit (Hawaii only)	Alemaria fruit and leaf spot, (passion fruit brown spot)	1-1/2 lbs	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when fruit spots appear (April to July) and continue treatments at 14 day intervals until weather conditions no longer favor disease development. Do not graze in treated area or feed vines or processing by-products to livestock used for food.
Pearl	Cercospora (early) leafspot Cercospondium (late) leafspot	1 to 1-1/4 lbs	Apply in sufficient water for adequate coverage when leaf wetness first occurs and before disease symptoms appear, generally 30 to 40 days after planting; repeat at 10 to 14 day intervals. When late leafspot prevails or when rust or web blotch occur, apply 1-1/4 lbs per acre at 10 day intervals for the remainder of the season. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated areas. Do not feed hay or threshings from treated fields to livestock. Bravo 90DG may be applied through sprinkler irrigation equipment. See calibration directions preceding this section. TANK MIXTURES: Bravo 90DG may be tank mixed with the boron fertilizer Solubor® and/or with the surfactant Surfex®. Read and observe all label directions and limitations for these products. Solubor® is a registered trademark of U.S. Borax. Surfex® is a registered trademark of Helena Chemical Co.
	Rust, Web blotch	1-1/4 lbs	
Potato	Late blight, Early blight, Botrytis vine rot	5/8 lb — then —	Apply as a banded treatment directed over the foliage, beginning when vines are first exposed and leaf wetness occurs. Repeat applications at 7 to 10 day intervals until vines close, then increase rate as described below. In addition to the early-season applications above, as vines close between rows increase water carrier volume to cover the denser canopy and begin broadcast application patterns. Continue applications at 7 to 10 day intervals. Use the highest registered rate weekly when disease conditions are severe. Bravo 90DG may be applied through sprinkler irrigation equipment (solid set, portable wheel move, center pivot or motorized lateral move systems only). Do not exceed a 10 day interval between applications when using this technique. Follow calibration and application directions preceding this section.
		7/8 to 1-1/4 lbs	
Soybean Determinate (Southern) Varieties	Anthracnose, Diaporthe pod and stem blight, Frogeye leaf spot (Cercospora soenkei), Purple seed stain, Cercospora leaf blight (Cercospora leaf blight), Septoria brown spot	1-1/8 to 2 lbs	Two application program — Make the first application at early pod set (R3) stage, when majority of pods are 1/8 to 3/8 inch in length and the second at beginning seed formation (R5) which occurs about 14 days later.
		7/8 to 1-1/2 lbs	Three application program — Make the first application at the beginning of flowering (R1), the second at early pod set (R3) and the third at beginning of seed formation (R5). Do not apply Bravo 90DG within 6 weeks of harvest. Do not feed soybean hay or threshings from treated fields to livestock.
		7/8 lb	Apply in 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make one application at emergence of second trifoliate leaves (V2). If conditions favor continued infection, make additional applications at 10 to 14 day intervals.
	Stem canker (Diaporthe phaseolorum var. caulivora)	7/8 lb	

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Soybean Indeterminate (Northern) Vanebes	Anthracnose, Diaporthe pod and stem blight, Frogeye leaf spot (<i>Cercospora soynai</i>), Purple seed stain, Cercospora leaf blight (<i>Cercospora kikuchii</i>), Septoria brown spot		Apply in sufficient water to obtain complete coverage, using at least five gallons of water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. Brave 900G may be applied through sprayer irrigation equipment. Follow application and calibration directions preceding this section. Do not exceed total of 3 applications per season. Do not apply within 6 weeks of harvest. Do not feed soybean hay or threshings from treated fields to livestock.
		1-1/8 to 2 lbs	Two application program — Make the first application when largest pods are 1 to 1-1/2 inches in length and make the second application 14 days later. Brave 900G may be co-applied with Benlate® 50WP as a tank mix for disease control in indeterminate (northern) soybeans. Use 7/8 lb of Brave 900G plus 8 ounces of Benlate 50WP per acre. Make the first application when pods near the top of plants are 1/2 to 1 inch in length and a second application 14 days later.
		7/8 to 1-1/2 lbs	Three application program — Make the first application one week after first flowering and continue applications at 14 day intervals.
Tomato	FOLIAGE (apply every 7-10 days): Early blight, Late blight, Gray leaf spot, Gray leaf mold, Septoria leaf spot, Target spot	1-1/8 to 1-3/4 lbs	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Use the highest rate and shortest interval specified when disease conditions are severe. Brave 900G may be combined in the spray tank with a copper fungicide which is registered to control bacterial diseases. Check the copper manufacturer's label for specific instructions, precautions and limitations prior to mixing with Brave 900G. Do not use with Copper-Court™ N in concentrated spray suspensions.
	FRUIT (apply every 7-14 days beginning at fruit set): Anthracnose, Alternaria fruit rot (black mold), Rhizoctonia fruit rot, Botrytis gray mold, Late blight fruit rot	1-3/4 to 2-1/4 lbs	Brave 900G may be applied through sprayer irrigation equipment. See calibration directions preceding this section.

*Copper-Court is a registered trademark of Mineral Research & Development Corporation.

®Benlate is a registered trademark of E.I. DuPont de Nemours and Co., Inc.

Tree and Orchard Crops

Apply Brave 900G in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, Brave 900G may be applied with aircraft using at least 20 gallons of spray per acre. When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of Brave 900G listed may be used. Do not allow livestock to graze in treated areas. The following spray volumes are recommended as gallons of spray per acre:

CROP	SPRAY VOLUME (Gallons per Acre)	
Peach, Nectarine, Apricot, Tart Cherry, Plum, Prune	20 (concentrate) to 300 (full dilute)	
Sweet Cherry	20 (concentrate) to 400 (full dilute)	
Conifers	Dilute	Concentrate
Forest stands	Not used	10 to 20 (aircraft)
Christmas trees	100	10 to 50 (aircraft or ground equipment)
Nursery beds	100	5 to 10 (ground equipment only)

CROP	DISEASES	BRAVO 880G RATE PER		APPLICATION DIRECTIONS
		ACRE	100 GAL*	
Peach, Nectarine, Apricot, Cherry, Plum, Prune	Leaf curl, Coryneum blight (shot-hole)	2-1/2 to 3-1/2 lbs	7/8 to 1-1/8 lbs	For best control of both diseases apply at leaf fall in late autumn, using sufficient water and proper spray calibration to obtain uniform coverage. When conditions favor high disease levels use the high rate of application and apply once or twice more in mid to late winter before budswell. If the leaf fall application is not practical, application of Bravo 880G for control of leaf curl may be made at any time prior to budswell the following spring. Where Coryneum blight (shot-hole) occurs, also apply at budbreak to protect newly emerging leaves and at shuck-split to prevent fruit infections.
	Brown rot blossom blight Lacy (necrotic) scab (plum/prune)	2-1/2 to 4-1/2 lbs	7/8 to 1-1/8 lbs	Use 3-1/2 to 4-1/2 lb per acre on trees taller than 20 feet and 2-1/2 to 3-1/2 lb per acre on smaller trees. Make one application at popcorn (pink, red or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry leaf spot; Peach, Nectarine, Apricot scab	2-1/2 to 4-1/2 lbs	7/8 to 1-1/8 lbs	In addition to the bloom application listed above, make one application at shuck-split. Do not apply Bravo 880G after shuck-split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10-14 days later.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

CROP	DISEASES	BRAVO 880G RATE PER		APPLICATION DIRECTIONS
		ACRE	100 GAL*	
Conifers	Swiss needlecast	2-1/4 to 4-1/2 lbs	2-1/4 to 4-1/2 lbs	Single application technique: In Christmas plantations or forest stands make one application in the spring when new shoot growth is 1/2 to 2 inches in length.
	Sclerotinia canker (pines), Swiss needlecast	1-1/8 to 2-1/4 lbs	1-1/8 to 2-1/4 lbs	Make the first application in spring when new shoot growth is 1/2 to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
	Sirococcus tip blight	1-3/4 to 3 lbs	1-3/4 to 3 lbs	
	Rhizosphaera needlecast (spruces), Scirrhia brown spot (pines)	4-1/2 lbs	4-1/2 lbs	
	Cyclaneusma and Lophodermium needlecasts (pines)	2-1/4 to 4-1/2 lbs	2-1/4 to 4-1/2 lbs	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific Northwest). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

CROP	DISEASES	BRAVO 880G RATE PER		APPLICATION DIRECTIONS
		ACRE	100 GAL*	
Conifers	Rhabdocline needlecast (Douglas-fir)	1-1/8 to 2-1/4 lbs	1-1/8 to 2-1/4 lbs	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
	Botrytis seedling blight, Phoma twig blight	1-1/8 to 2-1/4 lbs	1-1/8 to 2-1/4 lbs	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Storage: Store in a dry place.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Warranty and Limitation of Damages

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of Fitness or of Merchantability, and no agent of Seller is authorized to do so except in writing with a specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.