

- 2) At the end of the "DIRECTIONS FOR USE (Cont.)" box at the top of the right panel, insert:

See back panel for remainder of DIRECTIONS FOR USE including directions for mixing bait, prebaiting, and bait application.

- c. In the first sentence of the second paragraph of the "USE RESTRICTIONS," change "Seiver" to "Sevier."
- d. To the third paragraph of the "USE RESTRICTIONS," add:

Do not apply strychnine baits on or above food or feed crops.

3. It is a condition of this registration that the following data be submitted by the due date noted below.

<u>Guideline Reference</u> <u>Number</u>	<u>Name</u>	<u>Due Date</u>
63-17	Storage Stability	October 15, 1990

The test is to be conducted on the 1.6% strychnine paste.

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



William H. Milier
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

DANGER

Convulsive Poison! Poisonous if swallowed. Do not breathe dust. Do not contaminate food and food-stuffs. Keep away from children, pets, and domestic animals. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Baits exposed on soil surface may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**ENDANGERED SPECIES
CONSIDERATIONS**

NOTICE: The killing of an endangered species during strychnine baiting operations may result in a fine and/or imprisonment under the Endangered Species Act. Before baiting, the user is advised to contact the Regional U.S. Fish and Wildlife Service (Endangered Species Specialist) or the local Fish and Game Office for specific information on endangered species. Strychnine baits should not be used in the geographic ranges of the following species except under programs and procedures approved by the USEPA: California Condor, San Joaquin Kit Fox, Aleutian Canada Goose, Morro Bay Kangaroo Rat, Gray Wolf, Grizzly Bear, and Salt Marsh Harvest Mouse, Utah Prairie Dog and Masked Bob White Quail.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicators certification.

**FOR DISTRIBUTION AND USE ONLY WITHIN ARIZONA,
IDAHO, NEVADA, NEW MEXICO, AND UTAH**

1.6% STRYCHNINE PASTE

ACTIVE INGREDIENT: Strychnine Alkaloid 1.6%
INERT INGREDIENTS: 98.4%
TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN
DANGER-POISON**



STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY!

If less than ten (10) minutes have passed since the poison was taken, give 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Have patient lie down in quiet, darkened room and keep him warm and quiet. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and water.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN

A. Administer 100% OXYGEN by positive pressure to provide as much pulmonary gas exchange as possible, despite seizures.

B. Administer ANTICONVULSANT DRUGS intravenously to control convulsions.

CAUTION: It may be difficult or impossible to stop the seizure activity without stopping respiration. Be prepared to maintain pulmonary ventilation mechanically. Tracheotomy may be necessary if seizures are prolonged.

**UNITED STATES DEPARTMENT OF AGRICULTURE
ANIMAL AND PLANT HEALTH INSPECTION SERVICE**

Animal Damage Control
Hyattsville, MD 20782
EPA Est. No. 56228-ID-01
EPA Reg. No. 56228

Net Contents: 1 Gallon

DIRECTIONS FOR USE (Cont.)

USE RESTRICTIONS

For use against the following rabbit and hare species in rangelands, pasture, and non-agricultural areas (around airports only):

- Desert cottontail rabbits (*Sylvilagus auduboni*)
- Eastern cottontail rabbits (*S. floridanus*)
- Black-tailed jackrabbits (*Lepus californicus*)
- White-tailed jackrabbits (*L. townsendii*)
- Antelope jackrabbits (*L. alleni*)

Do not use in areas occupied by the Utah Prairie Dog in Garfield, Iron, Kane, Piute, Sevier, and Wayne counties, Utah. Do not use in areas occupied by the Masked Bobwhite Quail in Pima and Santa Cruz counties, Arizona.

Do not apply this bait intentionally to kill any species not claimed on this label.

STORAGE AND DISPOSAL

PROHIBITION: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers.

Keep cool in shade, out of direct sunlight. Not for use or store in or around the home. Do not store next to feed or transport in or on vehicles containing foodstuffs or feeds.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled paste, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact the appropriate State agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Pesticide, paste, or rinsate that cannot be used or chemically reprocessed should be disposed of in accordance with the regulations of the appropriate State agency. Pick up and burn or bury any uneaten bait and dead animals daily.

CONTAINER DISPOSAL: Thoroughly rinse (or equivalent) offer for recycling or reconditioning, or dispose of in approved landfill and in accordance with the regulations of the appropriate State Agency.

5/89

JUL 17 1989

Under Federal Law
Pesticide Products
Registered for the Use
of Strychnine Alkaloid
5/22/89-27

DIRECTIONS FOR USE (Cont.)

MIXING DIRECTIONS:

**SHAKE STRYCHNINE PASTE WELL BEFORE USE.
REFRIGERATE STRYCHNINE PASTE UNTIL USED**

CARROTS: Cut carrots into 1/4 to 1/2 inch cubes or slices. Use untreated carrot cubes or slices for prebait. To prepare toxic bait, place 25 pounds of carrot cubes or slices into a tub. Heat a ONE GALLON can of Strychnine Paste until lukewarm. Shake well and sprinkle the warmed contents of the ONE GALLON container over the 25 pounds of bait material in the tub. Stir until carrots are evenly coated.

ALFALFA OR CLOVER TIPS: Use untreated tips for prebait. To prepare toxic bait, place 25 pounds of tips into a tub. Heat a ONE GALLON can of Strychnine Paste until lukewarm. Shake well and sprinkle the warmed contents of the ONE GALLON container over the 25 pounds of bait material in the tub. Stir until tips are evenly coated.

DRY ALFALFA OR CLOVER TIPS: Use untreated dry tips for prebait. To prepare toxic bait, place 16 pounds of tips into a tub. Heat a ONE GALLON can of Strychnine Paste until lukewarm. Shake well and sprinkle the warmed contents of the ONE GALLON container over the 16 pounds of bait material in the tub. Stir until tips are evenly coated.

PREBAITING DIRECTIONS:

Before exposing poisoned bait, prebait with untreated bait for 3 to 5 days to condition animals to feed in locations other than where they are doing damage. Prebait with the same plant material that you intend to use in toxic baits. Prebaiting is most effective when done in the late afternoon.

Place prebaits at 15-foot intervals adjacent to pasture, range area, or airport area where target animals are causing problems.

Place prebait along trails used by target rabbits or jackrabbits if such trails are evident.

CARROTS: Use 6 to 8 cubes or slices per bait placement.

ALFALFA OR CLOVER BAITS: Use one cup of bait per placement.

BAITING DIRECTIONS:

In the late afternoon, remove or cover all prebait and replace with equal amounts of poisoned bait made from the same plant material. Place bait in same locations as were used for prebaits.