

MA 15 RE-101-10 1/2

**ACCEPTED**

SEP 16 1987

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, this pesticide registered with EPA Reg. No. 8780-59



TM







THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOO FARMS

**Arthroban-S**

**LAWN** 

 **INSECT KILLER**

**CONTROLS THESE INSECTS IN TURF**

			
<b>EUROPEAN CHAFER AND JAPANESE BEETLE GRUB</b>	<b>BLUE GRASS BILL BUG</b>	<b>SOO WEBWORM</b>	<b>CRUNCH BUG</b>
<b>TREAT AT RATE OF 4 LBS. PER 1000 SQ. FT. (PACKAGE TREATS 3000 SQ. FT. 30' x 100')</b>		<b>TREAT AT RATE OF 3.1 LBS. PER 1000 SQ. FT. (PACKAGE TREATS 6,000 SQ. FT.)</b>	

**TURF LINE ARTHROBAN-S LAWN INSECT KILLER**

For control of White Grubs of European Chaffer and Japanese Beetles, Sod Webworms, Crunch Bugs, Billbugs, Hyperodes Weevil, and Ants, in home lawns, recreational and ornamental turfgrass areas.

**NET WEIGHT 20 LBS.**

**ACTIVE INGREDIENT:**  
0,0-dimethyl 0-(2-isopropyl 6-methyl-4-pyrimidinyl) ph aphorothoate 3.13%

**INERT INGREDIENTS** 96.87%

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured By **HIGH POINT, U.S.**  
A Division of HighPoint Chemicals, Inc.  
1225 Lehigh Station Road  
Hanover, New York 14877

EPA Establishment No. 8780-NY-1  
EPA Registration No. 8780-86

**PRODUCT CODE 8101**



TM



# Arthroban-S

LAWN



INSECT KILLER

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING**

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**  
Causes eye irritation. Do not get in eyes, on skin, or clothing. Wash contaminated clothing before reuse. Avoid if swallowed or absorbed through skin. Avoid breathing dust. Avoid contamination of food, feedstuffs, and domestic water supplies. Keep children and pets off treated areas until this product is washed into the soil and the grass is dry.

**STATEMENT OF PRACTICAL TREATMENT**  
**IF SWALLOWED:** Call a physician or poison control center immediately. Drink one or two glasses of water and induce vomiting by washing the back of the throat with a finger. DO NOT induce vomiting or give anything by mouth to an unconscious person.

**IF IN EYES:** Flush eyes with water for at least 25 minutes. Get medical attention.

**IF IRRITATED:** Show victim to fresh air. Apply compression if necessary.

**IF ON SKIN:** Wash thoroughly with soap and water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** If symptoms of cholinesterase inhibition occur, administer atropine therapeutically.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Keep out of streams, lakes, and ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**NOTE THIS PRODUCT IS FOR:** Controlling existing infestations of turf of grub (larvae) of the European Chaffer and Japanese

Beetle. Controlling surface infestations of Sod Webworm, and Check Bug in Turf. Controlling populations of Sheargrass Biting and Appropriate Weevil in Turfgrass Areas.

**WHERE TO USE:** Use only on the grass of lawn areas, or other turf areas not used for pasture or hay. Do not use on edible food crop areas. This product will not harm rose seedlings.

**WHEN TO USE:** Apply when surface insects are active beginning in May for surface insects, and lasting through August. For soil grubs, make applications in August (preferred) or in the early spring (April).

**HOW TO USE:** Spread evenly over the lawn with a good fertilizer spreader. Settings given in the table are intended to be a guide in setting your fertilizer spreader due to variations in each spreader, and the varying techniques of application; it is important to make certain you are applying the correct amount of material. The correct use rate for control of soil grubs is: 4 lbs./1,000 sq. ft. The correct use rate for control of surface insects is: 3.3 lbs./1,000 sq. ft. Always lightly after application using 20 gallons per 1,000 sq. ft. When treating for soil grubs water thoroughly after application using 1/2-1 inch of water.

Spreader Make	Spreader Setting Number	
Cycloco Model B	3.3 lbs./1,000	4 lbs./1,000
Scout PF-1	3.8	4.2
Scout Compact Rotary	5.4	5.75
	1	2

**STORAGE & DISPOSAL**

Store this product in a safe place, beyond the reach of children and pets. Avoid contaminating foods or other products to be used in food bins.

Do not contaminate water, food, or feed by the storage or disposal of this product.

Do not reuse empty container. Wrap container and put in the trash collection.

