



U.S. ENVIRONMENTAL PROTECTION AGENCY  
 Office of Pesticide Programs  
 Registration Division (7505P)  
 1200 Pennsylvania Ave., N.W.  
 Washington, D.C. 20460

**EPA Reg. Number:**  
 59639-256

**Date of Issuance:**  
 3/17/22

**NOTICE OF PESTICIDE:**

Registration  
 Reregistration  
 (under FIFRA, as amended)

**Term of Issuance:**  
 Unconditional

**Name of Pesticide Product:**  
 TetraSan EZ Miticide/Ovicide

**Name and Address of Registrant (include ZIP Code):**

Sue-Chi Shen  
 Sr. Regulatory Project Associate Director  
 Registration and Regulatory Affairs  
 Valent U.S.A. LLC  
 1600 Riviera Ave, Suite 200, P.O. Box 8025  
 Walnut Creek, CA 94596-8025

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.

**Signature of Approving Official:**

Venus Eagle, Product Manager 01  
 Invertebrate and Vertebrate Branch 3, Registration Division (7505P)  
 Office of Pesticide Programs

**Date:**

3/17/22

2. Make the following label changes before you release the product for shipment:
  - Revise the EPA Registration Number to read, “EPA Reg. No. 59639-256.”
3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Please also note that the record for this product currently contains the following CSFs:

- Basic CSF dated 3/16/21
- Alternate CSF 1 dated 3/17/21

If you have any questions, please contact Marianne Lewis via email at [lewis.marianne@epa.gov](mailto:lewis.marianne@epa.gov)

Enclosure: Stamped label



**ACCEPTED**  
 03/17/2022  
 Under the Federal Insecticide, Fungicide  
 and Rodenticide Act as amended, for the  
 pesticide registered under  
 EPA Reg. No. 59639-256

ETOXAZOLE	GROUP	<b>10B</b>	ACARICIDE
-----------	-------	------------	-----------

# TetraSan<sup>®</sup> EZ MITICIDE/OVICIDE

For Ornamental Plants, Christmas Trees, and Greenhouse Tomato Production

For control of Spider Mites on Ornamental Plants, Christmas trees and Greenhouse Tomatoes

Affects all life stages of spider mites.

Active Ingredient	By Wt
Etoxazole* .....	31.7%
Other Ingredients .....	<u>68.3%</u>
Total	100.0%

\*2-(2,6-difluorophenyl)-4-[4-(1,1-dimethylethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole

*TetraSan* EZ MITICIDE/OVICIDE is a soluble concentrate containing 2.88 lb active ingredient etoxazole per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

**SEE NEXT PAGE [BACK][PANEL][BOOKLET]FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

**EPA Reg. No. 59639-XXX**

**EPA Est. No.**

Lot No. :

**NET Content :**

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

#### CAUTION

##### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact **800-892-0099** for emergency medical treatment information.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, shoes, socks, and chemical-resistant gloves made out of: barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride ≥14 mils, or Viton ≥14 mils..

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS:

This pesticide is toxic to freshwater and marine/estuarine aquatic invertebrates, including oysters and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval.

Do not enter or allow workers entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants, socks plus shoes, and chemical-resistant gloves.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses.

Keep all unprotected persons out of operating areas, or vicinity where there may be drift.  
Do not enter treated areas without protective clothing until sprays have dried.

**Always mix product thoroughly before use.**

## RESISTANCE MANAGEMENT

For resistance management, TetraSan EZ MITICIDE/OVICIDE contains a Group 10B acaricide. Any mite population may contain individuals naturally resistant to TetraSan EZ MITICIDE/OVICIDE and other Group 10B acaricides. The resistant individuals may dominate the mite population if this group of acaricides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay acaricide resistance, take the following steps:

- Rotate the use of TetraSan EZ MITICIDE/OVICIDE or other Group 10B acaricides within a growing season, or among growing seasons, with different groups that control the same pests.
- Use tank mixtures with acaricides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests that the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Valent U.S.A. LLC at 1-800-6-VALENT (682-5368).

## TANK MIXES

**NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, to the extent allowed by applicable law.**

## INFORMATION

- **Intended for commercial use.**

*TetraSan* EZ MITICIDE/OVICIDE is a miticide that controls major tetranychid mite species in egg and nymph stages of growth. Foliar applications of *TetraSan* EZ MITICIDE/OVICIDE can be made for control of spider mites infesting indoor (greenhouses, lathhouses, shadehouses and interiorscapes) and outdoor ornamentals, including flowering and foliage crops, ground covers, shrubs and ornamental trees, non-bearing fruit and nut trees, Christmas trees and indoor tomato production (greenhouses, lathhouses and shadehouses). *TetraSan* EZ MITICIDE/OVICIDE may be used in ornamental production areas and in the landscape in residential and commercial areas. *TetraSan* EZ MITICIDE/OVICIDE exhibits strong translaminar movement within plant leaves. Spray contacting the upper surface of leaves will absorb into the leaves and locally translocate within the leaf. Thus mites feeding on the undersides of leaves will be controlled.

**IMPORTANT:** Applications of *TetraSan* EZ MITICIDE/OVICIDE may be made to a wide range of ornamental plants. To ensure that *TetraSan* EZ MITICIDE/OVICIDE is compatible with the variety or cultivar under your specific conditions, test the product on a limited scale and observe for phytotoxicity for two weeks before making large scale applications. The large number of existing ornamental varieties and cultivars coupled with the constant introduction of new varieties makes it impossible to field test *TetraSan* EZ MITICIDE/OVICIDE in every locale where sold or in all of the combinations created by differences in growing practices. Factors that can influence phytotoxicity to any applied pesticide include soil or media type, pH, moisture or fertility, environmental conditions such as temperature, lighting or degree days and horticultural practices including planting dates, control of other pests, and timing of various horticultural practices. *TetraSan* EZ MITICIDE/OVICIDE has not been tested in tank mix combinations, sequential pesticide treatments or with various adjuvants and surfactants. Do not apply to Poinsettia after bract formation.

### SPRAY EQUIPMENT

Thoroughly clean spray equipment prior to mixing and applying *TetraSan* EZ MITICIDE/OVICIDE. Thoroughly rinse the spray tank, spray boom and all nozzles before adding *TetraSan* EZ MITICIDE/OVICIDE to the tank. Apply *TetraSan* EZ MITICIDE/OVICIDE using equipment such as compressed air sprayers, hydraulic sprayers, or ground boom sprayers.

### TANK MIXING

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### COMPATIBILITY

Before mixing *TetraSan* EZ MITICIDE/OVICIDE with other insecticide products, test mix small portions of products in a small quantity to ensure compatibility.

## MANDATORY SPRAY DRIFT MANAGEMENT

### SPRAY DRIFT

#### Aerial Applications

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select a nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572.1).
- If the windspeed is 10 miles per hour or less, applicators must use  $\frac{1}{2}$  swath displacement upwind at the downwind edge of the field. When the windspeed is between 11-15 miles per hour, applicators must use  $\frac{3}{4}$  swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Do not apply during temperature inversions

#### Airblast Applications

- Sprays must be directed into the canopy.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- User must turn off outward pointing nozzles at low ends and when spraying outer row.
- Do not apply during temperature inversions.

#### Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to select a nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

#### Boom-less Ground Applications

- Applicators are required to use a medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

### SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.



### **Controlling Droplet Size – Ground Boom**

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### **Controlling Droplet Size – Aircraft**

- Adjust Nozzles - Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

### **BOOM HEIGHT – Ground Boom**

For ground equipment, the boom should remain level with the crop and have minimal bounce.

### **RELEASE HEIGHT - Aircraft**

Higher release heights increase the potential for spray drift.

### **SHIELDED SPRAYERS**

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### **TEMPERATURE AND HUMIDITY**

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

### **WIND**

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **Boom-less Ground Applications**

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

### **Handheld Technology Applications**

- Take precautions to minimize spray drift.

**DIRECTIONS FOR USE ON ORNAMENTAL PLANTS**

CROP	PESTS	PRODUCT RATE	SPECIAL INSTRUCTIONS
<p>Ornamental Plants Including: Bedding Plants Flowering Plants Foliage Plants Ground Covers Non-Bearing Fruit Trees Non-Bearing Nut Trees Ornamentals Ornamental Trees Shrubs</p>	<p>Spider Mites Including: Citrus Red Mite European Red Mite Lewis Spider Mite Pacific Spider Mite Southern Red Mite Spruce Spider Mite Twospotted Spider Mite</p>	<p>1.1 to 2.2 fl oz/100 gallons (0.025 to 0.05 lb ai/100 gallons)</p> <p>2.2 to 4.4 fl oz/A (0.05 to 0.1 lb ai/A)</p>	<p>Mix with sufficient water and apply as a foliar spray to obtain uniform coverage.</p> <p><i>TetraSan EZ</i> MITICIDE/OVICIDE kills mite eggs (ovicidal) and nymphs, but not adult mites. For best control, apply at first sign of infestation and before large numbers of adult mites are present. Make a second application if necessary, but no sooner than 14 days after the first application.</p> <p><i>TetraSan EZ</i> MITICIDE/OVICIDE is transovarial, meaning that treated adult mites will not produce viable eggs.</p> <p>Use higher rates within the rate range for moderate to heavy infestations, especially with dense plant canopies.</p> <p>If rapid control (&lt; 7 days) of adult mites is required, apply <i>TetraSan EZ</i> MITICIDE / OVICIDE in combination with a contact adulticide.</p> <p>Non-bearing: will not bear fruit/nuts within 365 days after application</p>
<p><b>IMPORTANT:</b> Since ornamental varieties are numerous, constantly changing, and may react differently to <i>TetraSan EZ</i> MITICIDE/OVICIDE and tank mixtures, test the product(s) on a small scale before making large-scale applications.</p> <p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>Do not apply <i>TetraSan EZ</i> MITICIDE/OVICIDE more than two times per cropping season or within a six month period.</li> <li>Do not apply this product through any type of irrigation or chemigation system.</li> </ul>			

**DIRECTIONS FOR USE ON CHRISTMAS TREES**

CROP	PESTS	PRODUCT RATE	SPECIAL INSTRUCTIONS
Christmas Trees	McDaniel Spider Mite Pacific Spider Mite Spruce Spider Mite Twospotted Spider Mite	3.9 to 5.6 fl oz/A (0.088 to 0.125 lb ai/ A )	<p>Apply by ground as full coverage spray or by air in a minimum of 10 gallons/A. Coverage is critical for effective control.</p> <p>Best results are achieved when mite populations are low. <i>TetraSan</i> EZ MITICIDE/OVICIDE is predominately an ovicide/larvicide. Apply <i>TetraSan</i> EZ MITICIDE/OVICIDE at or prior to threshold for your area but not greater than the maximum rate listed.</p> <p>Do not use rates below 3.9 fl oz/A (0.088 lb ai/A) since it may result in poor control and contribute to the development of resistance to etoxazole among mite populations.</p> <p>Use higher rates within the rate range for moderate to heavy infestations.</p>
<p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Do not make more than one (1) <i>TetraSan</i> EZ MITICIDE/OVICIDE application per calendar year.</li> <li>• Do not apply more than 5.6 fl oz (0.125 lb ai) <i>TetraSan</i> EZ MITICIDE/OVICIDE per acre calendar year.</li> <li>• Do not apply this product through any type of irrigation or chemigation system.</li> </ul>			

## DIRECTIONS FOR USE IN GREENHOUSE TOMATO PRODUCTION

CROP	PESTS	PRODUCT RATE	SPECIAL INSTRUCTIONS
Tomatoes	Spider Mites Including: Citrus Red Mite European Red Mite Lewis Spider Mite Pacific Spider Mite Southern Red Mite Spruce Spider Mite Twospotted Spider Mite	1.1 to 2.8 fl oz/100 gallons (0.025 to 0.063 lb ai/100 gallons)  2.2 to 5.6 fl oz/A (0.05 to 0.125 lb ai/A)	<p>Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. When applied as a high volume spray, 100 gallons of spray solution will treat approximately 20,000 sq ft.</p> <p><i>TetraSan</i> EZ MITICIDE/OVICIDE kills mite eggs (ovicidal) and nymphs, but not adult mites. For best control, apply at first sign of infestation and before large numbers of adult mites are present. Make a second application if necessary, but no sooner than 21 days after the first application.</p> <p><i>TetraSan</i> EZ MITICIDE/OVICIDE is transovarial, meaning that treated adult mites will not produce viable eggs.</p> <p>Use higher rates within the rate range for moderate to heavy infestations, especially with dense plant canopies.</p> <p>If rapid control (&lt; 7 days) of adult mites is required, apply <i>TetraSan</i> EZ MITICIDE/OVICIDE in combination with a contact adulticide.</p> <p>May also be used in lathhouses and shadehouses.</p>
<p><b>IMPORTANT:</b> Since tomato varieties are numerous, constantly changing and may react differently to <i>TetraSan</i> EZ MITICIDE/OVICIDE and tank mixtures, test the product(s) on a small scale before making large-scale applications.</p> <p><b>Restrictions</b></p> <ul style="list-style-type: none"> <li>• Do not apply within one (1) day of harvest.</li> <li>• Do not make more than two (2) <i>TetraSan</i> EZ MITICIDE/OVICIDE applications per cropping season. Maximum two crop seasons per calendar year.</li> <li>• Do not exceed 11.1 fl oz <i>TetraSan</i> EZ MITICIDE/OVICIDE (0.25 lb ai) per acre per cropping season.</li> <li>• Do not apply <i>TetraSan</i> EZ MITICIDE/OVICIDE more than two times per cropping season.</li> <li>• Do not apply more than 5.6 fl oz (0.125 lb ai) per acre per application.</li> <li>• Do not use with an adjuvant or surfactant.</li> <li>• Do not apply this product through any type of irrigation or chemigation system.</li> </ul>			

## **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

### **PESTICIDE STORAGE**

Keep pesticide in original container.

Do not put concentrate or dilute into food or drink containers.

Store in cool, dry place.

Do not store diluted spray.

For help with any spill, leak, fire, or exposure involving this material, call day or night **(800) 892-0099**.

### **PESTICIDE DISPOSAL**

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### **CONTAINER HANDLING**

Nonrefillable container {HDPE plastic, < 5 gallons}. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available; otherwise puncture and dispose of in a sanitary landfill.

### **RECYCLING**

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 877-952-2272 (toll free) or [www.acrecycle.org](http://www.acrecycle.org).

**RISKS OF USING THIS PRODUCT,  
LIMITED WARRANTY AND DISCLAIMER,  
AND LIMITATION OF LIABILITY**

**IMPORTANT: Read the entire Label including this section titled Risks of Using this Product, Limited Warranty and Disclaimer, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.**

**RISKS OF USING THIS PRODUCT**

The buyer and user (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND TO THE FULLEST EXTENT ALLOWED BY LAW, AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

The Directions for Use of this product must be followed carefully. Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

**LIMITED WARRANTY AND DISCLAIMER**

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. To the extent consistent with applicable law, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

**LIMITATION OF LIABILITY**

**To the fullest extent allowed by law, Valent or Seller is not liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. TO THE FULLEST EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

**PROMPT NOTICE OF CLAIM**

To the extent consistent with applicable law allowing such requirements, Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

**NO AMENDMENTS**

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing **Risks of Using This Product, Limited Warranty and Disclaimer, and Limitation of Liability**, which may not be modified by any oral or written agreement.

©2022 Valent U.S.A. LLC

*TetraSan* is a registered trademark of Valent U.S.A. LLC.

Manufactured for:

**Valent U. S. A. LLC**

P.O. Box 5075

San Ramon CA 94583

(800)-89-VALENT (898-2536)

Made in U.S.A.

EPA Reg. No. 59639-XXX

EPA Est. No.

059639-00XXX.20220316.TET\_EZ\_NPR.Clean