



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 18, 2020

Ms. Mary McKinney Hunt
Authorized Representative
OHP, Inc.
P. O. Box 746
Bluffton, S.C. 29910-0746

Subject: Label Amendment – Update WSP mixing and loading language
Product Name: **OHP 50 WP Fungicide**
EPA Registration Number: 59807-6
Application Date: October 3, 2017
Decision Number: 538453

Dear Ms. Hunt:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 59807-6
Decision No. 538453

with FIFRA section 6. If you have any questions, please contact Eleanor Thornton by phone at 703-305-6799, or via email at thornton.eleanor@epa.gov.

A handwritten signature in black ink that reads "Shaja B. Joyner". The signature is written in a cursive style with a large, looping initial "S".

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

THIOPHANATE-METHYL GROUP 1 FUNGICIDE

OHP 50 WP Fungicide

ABN: OHP 6672 50 WP

Systemic Turf and Ornamental Fungicide

For control of a broad spectrum of diseases of herbaceous, bedding, flowering and tropical foliage plants, shrubs, trees and flowers in the landscape, interiorscape, nursery and greenhouse, containerized woody shrubs and trees, turfgrass, and vegetable transplants.

ACTIVE INGREDIENT:

Thiophanate-methyl 50.0%

OTHER INGREDIENTS:..... 50.0%

TOTAL:..... 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**


| FIRST AID | |
|---|---|
| IF IN EYES: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye • Call a poison control center or doctor for treatment advice |
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person. |
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice |
| IF INHALED: | <ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible • Call a poison control center or doctor for further treatment advice |
| HOT LINE NUMBER | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-356-4647 for emergency medical treatment information. | |

See inside label booklet for additional Precautionary Statements

**EPA Reg. No. 59807- 6
EPA Est. No.**

Manufactured for:
OHP, Inc.
PO Box 746
Bluffton, SC 29910-0746
(800) 356-4647

Net Contents: Four 8-ounce water soluble bags



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq mils, polyvinyl chloride (PVC) \geq 14 mils or viton \geq 14 mils.

Mixers, loaders, applicators and other handlers supporting dip treatment must wear:

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves
- Chemical-resistant apron

All other mixers, loaders, applicators and handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves for all mixers and loaders and for applicators using hand held equipment

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement

Water soluble packets, when used correctly, qualify as a closed mixing/loading system under the Worker Protection Standard [40 CFR 170.607(d)]. Mixers and loaders handling this product while it is enclosed in intact water soluble packets may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes, socks, a chemical-resistant apron, and chemical-resistant gloves. When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Remove clothing/**PPE** immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove **PPE** immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for homeowner use. For use only by certified applicators or those under their immediate supervision. Not for use on turf being grown for sale or other commercial use as sod. Do not apply to home orchards/backyard fruit trees after fruit set. Do not apply with fixed wing or rotary aircraft.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exemption: The Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Coveralls over long sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposures

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter treated areas without protective clothing until sprays have dried.

PRODUCT INFORMATION

OHP 50 WP Fungicide provides broad spectrum disease control on woody, flowering, herbaceous, and tropical foliage ornamental plants and trees in containers, greenhouses and landscapes and also on turfgrass.

This canister/bag contains [four] 8-ounce OHP 50 WP Fungicide water soluble bags.

Instructions for Using Water Soluble Packages Directly into Spray tanks:

Water Soluble Packages (WSPs) are designed to dissolve in water. Agitation may be used, if necessary, to help dissolve the WSP. Failure to follow handling and mixing instructions can increase your exposure to the pesticide products in WSPs. WSPs, when used properly, qualify as a closed mixing/loading system under the Agricultural Worker Protection Standard [40 CFR 170.607(d)].

Handling Instructions

Follow these steps when handling pesticide products in WSPs.

1. Mix in spray tank only.

2. Handle WSP(s) in a manner that protects package from breakage and/or unintended release of contents. If package is broken, put on PPE required for clean-up and then continue with mixing instructions.
3. Keep the WSP(s) in outer packaging until just before use.
4. Keep the WSP dry prior to adding to the spray tank.
5. Handle with dry gloves and according to the label instructions for PPE.
6. Keep WSP intact. Do not cut or puncture WSP.
7. Reseal the WSP outer packaging to protect any unused WSP(s).

Mixing Instructions

Follow the steps below when mixing this product, including if tank mixed with other pesticide products. If being tank mixed, the mixing directions 1 through 9 below take precedence over the mixing directions of the other tank mix products. WSPs may, in some cases, be mixed with other pesticide products so long as the directions for use of all mixed products do not conflict. Do not tank mix this product with products that prohibit tank mixing or have conflicting mixing directions.

1. If a basket or strainer is present in the tank hatch, remove prior to adding the WSP to the tank.
2. Fill tank with water to approximately one-third to one-half of the desired final volume of spray.
3. Stop adding water and stop any agitation.
4. Place intact/unopened WSP(s) into the tank.
5. Do not spray water from a hose or fill pipe to break or dissolve the WSP(s).
6. Start mechanical and recirculation agitation from the bottom of tank without using any overhead recirculation, if possible. If overhead recirculation cannot be turned off, close the hatch before starting agitation.
7. Dissolving the WSP(s) may take up to 5 minutes or longer, depending on water temperature, water hardness and intensity of agitation.
8. Stop agitation before tank lid is opened.
9. Open the lid to the tank, exercising caution to avoid contact with dusts or spray mix, to verify that the WSPs have fully dissolved and the contents have been thoroughly mixed into the solution.
10. Do not add other allowed products or complete filling the tank until the bags have fully dissolved and pesticide is thoroughly mixed.
11. Once the WSP have fully dissolved and any other products have been added to the tank, resume filling the tank with water to the desired level, close the tank lid, and resume agitation.
12. Use the spray solution when mixing is complete.
13. Maintain agitation of the diluted pesticide mix during transport and application.
14. It is unlawful to use any registered pesticide, including WSPs, in a manner inconsistent with its label.

Maintain agitation to ensure even suspension of the product. Make fresh daily. Spray uniformly over the area to be treated with a boom type or other power sprayer.

Make applications of OHP 50 WP Fungicide with ground equipment, using sufficient spray volume to provide thorough coverage. Do not tank mix with copper containing materials or with highly alkaline pesticides such as Bordeaux mixture or lime sulfur. No claim of compatibility with other pesticides is implied. Under conditions of severe disease pressure or when application intervals are shorter than 14 days due to persistent rainfall, use the higher concentration or rates provided in this label. Contact your local State Extension Service specialist for application schedule recommendations.

Important: If recommended treatments of OHP are ineffective, a tolerant strain of fungus may be present. Consult your OHP, Inc. representative or distributor, your local State Agricultural Experiment Station or State Agricultural Extension Service for advice on prompt use of some other labeled fungicides.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

{note to reviewer-California restriction used until vegetable transplants approved for use in California. See EPA letter dated February 26, 2016}

[NOTE: This product is not for use on vegetable transplants in California.]

Resistance Management:

For resistance management, OHP 50 WP Fungicide contains a Group 1 fungicide. Any fungal population may contain individuals naturally resistant to OHP 50 WP Fungicide and other Group 1 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of OHP 50 WP Fungicide or other Group 1 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- OHP, Inc. does not recommend the use of products containing thiabendazole in combination or rotation with OHP 50 WP Fungicide. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Consult with your local OHP, Inc. representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. If, after using OHP 50 WP Fungicide as recommended, the treatment is not effective, a tolerant strain of fungi may be present.

CHEMIGATION INSTRUCTIONS

For Use on Ornamentals Only

Apply this product only through pressurized drench (flood), sprinkler, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system.

Do not connect chemigation system to a public water system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Should the need arise, a person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments.

Pressurized Drench (Flood) System

Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from back flow if water flow stops. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops,
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Sprinkler Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops,
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

Drip (Trickle) Chemigation

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back towards the injection pump.
- The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being drawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops,
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

HORTICULTURAL APPLICATIONS

GREENHOUSE, NURSERY, LANDSCAPE AND INTERIORSCAPE

OHP 50 WP Fungicide provides broad spectrum control of many foliar, stem, and below ground diseases on a wide range of horticultural plants grown or maintained under a variety of conditions. Apply 14-21 days prior to when a particular disease usually appears and at the very latest, upon first sign of disease. Spray intervals usually range from 7 to 14 days with 14 days being for preventative treatments and the 7 day interval for times when conditions are judged acceptable for disease development. For hard to wet foliage, an acceptable wetting agent may be added to the spray tank to increase product efficacy. Use of a spreader-sticker is recommended when excessive and repeated foliar wetting occurs. OHP 50 WP Fungicide may be used to control listed diseases on non-commercial fruit and nut trees.

Do not apply to home orchards or backyard fruit trees after fruit set.

Do not use fruit, nuts or sap from treated trees as food or feed.

Note: OHP 50 WP Fungicide has been determined to be safe for use on the plant types listed in these directions for use based on cumulative data derived from research product trials and historical field use. As all species and cultivars have not been tested, it is recommended that trial applications be performed on a similar use site listed on the label and for disease listed on the label. To conduct a trial application, apply at least two applications to at least 25 trial plants at the highest concentration, 7 days apart. Evaluate 7 days after the last application before initiating full-scale application.

Restriction: Do not use this product on the following plants: Swedish Ivy (*Plectranthus australis*), Boston Fern (*Nephrolepis exhalta*), and Easter Cactus (*Hatiora gaertneri*).

| Plant Type | Such as: |
|--------------------|--|
| Herbaceous Bedding | Ageratum, Begonia, Canna, Coleus, Dahlia, Dusty Miller, Foxglove, Fuchsia, Geranium, Impatiens, Lavender, Marigold, Pansy, Petunia, Pinks, Primrose, Salvia, Statice, Strawflower, Tickseed, Verbena |
| Flowering | Chrysanthemum, Hydrangea, Hollyhock, Iris, Lily, Poinsettia |
| Tropical Foliage | Dieffenbachia, Dracaena, English Ivy, Philodendron, Pothos |
| Woody Ornamentals | Azalea, Hibiscus, Holly, Ligustrum, Rhododendron, Rose, Pyracantha |

| | |
|------------------|---|
| Evergreen Trees | Douglas Fir, Fir, Larch, Pine, Spruce |
| Deciduous Trees* | Ash, London Plane, Maple, Oak, Sycamore, Walnut |
| Flowering Trees* | Cherry, Crabapple, Hawthorn, Mountain Ash, Pear |

*Do not use fruit, nuts or sap from treated trees as food or feed.

FOLIAR SPRAY APPLICATIONS

Hydraulic Application Mixing Instructions

Add the required amount of OHP 50 WP Fungicide to a partially filled spray tank agitated by mechanical or hydraulic means and then add the remaining recommended volume of water. Maintain continuous agitation to keep the material in suspension and apply with properly calibrated spray equipment.

Application Concentrations (Mechanical or Hand Held):

Use the specified amount of OHP 50 WP Fungicide per 100 gallons of water for the prevention and control of the diseases shown below.

FOLIAR DISEASES

| Diseases/Pathogens Controlled | Concentration of OHP 50 WP Fungicide oz/100 gal | Application Instructions |
|---|---|--|
| Anthraco­nose <i>Colletotrichum</i> | 12-16 oz/100 gallon | Apply as buds break or at first sign of disease. Repeat application at 7-14 day intervals during disease period. |
| Black Spot of Rose <i>Diplocarpon rosae</i> | 12-16 oz/100 gallon | Apply early summer or at first sign of disease. Repeat every 7-14 days during disease period. |
| Brown Rot and Blight <i>Monilinia, Sclerotini, Whetzellinia</i> | 12-16 oz/100 gallon | Apply late spring or at first sign of disease. Repeat every 7-14 days during disease period. |
| Fusicladium and Venturia Leaf Scabs on: Crabapple, Hawthorn, Pear, Mountain Ash, Pyracantha, etc. | 12-16 oz/100 gallon | Apply as buds break. Repeat every 7-14 days during disease period. Effective control requires coverage during leaf expansion. Rotations and/or tank mix combinations with chlorothalonil or propiconazole can be utilized. |
| Leaf Spots and Blights caused by: <i>Ascochyta, Blumeriella, Botrytis, Cercospora, Coccomyces, Corynespora, Curvularia, Didymellina, Entomosporium, Fabraea, Fusarium, Ramularia, Rhizoctonia, Marssonina, Mycosphaerella, Myrothecium, Phoma, Physalospora, Schizothyrium, Septoria, Sphaceloma</i> | 12-16 oz/100 gallon | Make applications when disease symptoms first appear. Repeat every 7-14 days during disease period. Rotations chlorothalonil may be used. |
| Ovulinia Blight | 8-16 oz/100 gallon | Apply as flowers open. Repeat every 7-14 days during disease period. |
| Powdery Mildews <i>Erysiphe, Microsphaera, Phyllactinia, Podosphaera, Oidium, Sphaerotheca</i> | 8-16 oz/100 gallon | Apply when disease first appears and repeat every 7-14. Rotations with other effective products are recommended. |
| Rust Diseases caused by: | 12-16 oz/100 | Apply late spring or when symptoms |

| Diseases/Pathogens Controlled | Concentration of OHP 50 WP Fungicide oz/100 gal | Application Instructions |
|---|---|---|
| <i>Puccinia, Gymnosporangium, Uromyces</i> | gallon | first appear. Repeat application every 7-14 days during disease period. Rotations with other effective products are recommended. |
| Tip Blight of Pine <i>Sphaeropsis sapinea, Diplodia pinea</i> | 12-16 oz/100 gallon | Begin applications in the spring when new growth starts. Make a second application just prior to needle emergence from the sheath and a third application 7 days later. Ensure thorough coverage. |
| Twig Blights, Cankers, and Diebacks <i>Diaporthe, Kabatina, Phoma, Phomopsis</i> | 16-32 oz/100 gallon | Apply when symptoms first appear. Repeat application every 7-14 days during disease period. |

Adjuvants: Where rainfall and/or overhead irrigation is the norm, use of a compatible spreader/sticker is suggested. Where wetting of foliage is difficult, use a compatible wetting agent. Follow the phytotoxicity precautions described in the "HORTICULTURAL APPLICATIONS" section of this label.

SOIL DRENCH APPLICATIONS

Mixing Instructions: Add the required amount of OHP 50 WP Fungicide to a partially filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation throughout application to keep the material in suspension.

Application Concentrations/Rates and Timing for Disease Control: Use 8 to 16 oz of OHP 50 WP Fungicide per 100 gallons of water. Apply as a drench or heavy spray at the rate of ½ to 2 pints per square foot (100 gallons per 400 to 1,600 square feet). For small pots and shallow flats up to 4 inches in size, use 8 oz/100 gallons applied at 1 pint per square foot. For containers and pots 4 inches or larger, refer to the following table for the volume to apply. Repeat applications may be made at 4 to 8 week intervals depending on disease presence and conditions for disease development.

| Container Type | Volume to Apply/Container | |
|----------------|---------------------------|-----------------|
| | 1 pt/sq ft Rate | 2 pt/sq ft Rate |
| 4 inch | 2 fl oz | |
| 5 inch | 2 ½ fl oz | |
| 6 inch | | 6 ½ fl oz |
| 7 inch | | 8 ½ fl oz |
| 8 inch | | 11 fl oz |
| 9 inch | | 14 fl oz |
| 10 inch | | 17 ½ fl oz |

For containers larger than 10 inches, a drench volume of 2 ½ to 3 pints per square foot of surface area may be required.

Plant Types: Containerized woody shrubs, trees, herbaceous/bedding, flowering, and tropical foliage plants and flowers and bedding plants in the landscape.

Restriction: Do not apply this product to plug trays or seedling flats at time of seeding.

Soil Diseases Controlled: *Botrytis, Cythrodcladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia and Thielaviopsis.*

Note: *Pythium, Phytophthora and Cythrodcladium spathiphylli are not controlled by OHP 50 WP Fungicide.*

PLANT DIP TREATMENT

Mixing Instructions: Mix as described in the “FOLIAR DISEASES” and “SOIL DRENCH APPLICATIONS” sections of this label. Maintain continuous agitation during application.

Application Concentration and Dipping Time

Plants or Cuttings: Use 12-16 oz of OHP 50 WP Fungicide per 100 gallons of water. Immerse plants or cuttings for 10 to 15 minutes, remove and allow to drain and dry. Wear protective clothing as described under the “PERSONAL PROTECTIVE EQUIPMENT” section of this label.

Note: Follow accepted hygiene practices to minimize the introduction and spread of water borne bacterial and water mold fungal diseases.

Bulbs, Corms, Tubers and Rhizomes: Use 16-32 oz of OHP 50 WP Fungicide per 100 gallons of water. Soak cleaned bulbs for 15 to 30 minutes in warm dip (80-85°F) within 48 hours after digging. Dry bulbs after treatment. If bulbs are for forcing, treat bulbs that have been heat-cured.

Note: Follow accepted hygiene practices to minimize the introduction and spread of water born bacterial and water mold fungal diseases.

Plant Types: Propagated units of woody, herbaceous, flowering and tropical foliage plants. Bulbs, corms, tubers, and rhizomes of plants such as but not limited to Caladium, Easter Lily, Tulip, Gladiolus, Daffodil, Iris.

Diseases Controlled: *Botrytis, Cythrodcladium, Fusarium, Gliocladium, Myrothecium, Penicillium, Ramularia, Rhizoctonia, Sclerotinia, Thielaviopsis*

INTERIORSCAPE USE

Spray Application: Mix well before spraying. Ensure complete coverage by spraying foliage to runoff, thoroughly wetting upper and lower leaf surfaces. Use concentrations under the “FOLIAR DISEASES” and “SOIL DRENCH APPLICATIONS” sections of this label for foliar, twig and drench application. Use a coarse, low pressure spray when applying in public areas. Apply during non-business hours or cordon off area during spraying and until spray has dried. Avoid spraying wall coverings, plastic (vinyl) surfaces, drapes, carpets, and upholstery.

TURF APPLICATIONS

OHP 50 WP Fungicide may be used against certain foliar and soil diseases for use on all turf types such as golf course greens, tees and fairways, athletic fields, cemeteries, parks, and commercial and home lawns. OHP 50 WP Fungicide may be used both preventatively and curatively and is not phytotoxic.

Restrictions: Do not use OHP 50 WP Fungicide on turf being grown for sale or other commercial uses as sod. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

Mixing Instructions: Add the required amount of OHP 50WP Fungicide to a partially filled tank agitated by mechanical or hydraulic means. Add the remaining required amount of water. Maintain continuous agitation to keep the material in suspension. For best results, use spray mixture the same day it is prepared.

Turf Types: All cool season and warm season grasses (such as but not limited to Bentgrasses, Bermudagrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustinegrasses, and Zoysiagrasses) or their mixtures.

| Disease(s) Controlled | Rate of OHP 50 WP Fungicide oz/1000 sq ft * | Application Instructions |
|---|---|--|
| Anthracnose <i>Colletotricum graminicola</i> | 2 – 4 (4 – 6)** | Apply when disease first appears. Make additional applications at 14 day intervals. Allow spray to dry on leaves with no “watering in.” |
| Dollar Spot <i>Sclerotinia homoeocarpa</i> (<i>Mollerodiscus</i> spp., <i>Lanzia</i> spp.), Copper Spot (<i>Gloeocerospora sorghi</i>), Brown Patch and Zoysia Patch <i>Rhizoctonia solani</i> , Red Thread <i>Laetisaria fusiformis</i> (<i>Corticium fuciforme</i>) | 2 – 4 | Apply when disease first appears. Make additional applications at 14 day intervals. Allow spray to dry on leaves with no “watering in.” |
| Pink Snow Mold <i>Microdochium nivale</i> (Only for those areas where snow cover is not present the entire winter) | 2 – 4 | Apply OHP 50 WP Fungicide in middle to late November before turf has stopped all growth activity. Lightly water this application into the root zone for best results. For best results, use a spreader-sticker. Second spray should dry on leaf surfaces with no “watering in”. Minimum Spray interval is 14 days. |
| Gray Leaf Spot <i>Pyricularia grisea</i> | 4 – 6 | Apply when conditions are favorable for disease development. Continue applications at 14 day intervals. Allow spray to dry on leaves with no “watering in.” |
| Summer Patch <i>Magnaportha poae</i> | 4 – 6 | For preventative treatment, make 3 applications at 21 day intervals beginning in early May. Water product into the root zone thoroughly after application. For suppression apply two applications at 14 day intervals beginning when disease first appears. |
| Fusarium Blight <i>Fusarium</i> spp. (<i>Fusarium roseum</i> and <i>tritricum</i>), Necrotic Ring Spot and Spring Dead Spot <i>Leptosphaeria korrea</i> (<i>Dreschlera</i> spp.) (<i>Helminthosporium</i> spp.) | 4 – 6 | Make two applications at 14 day intervals beginning when disease first appears. |
| Stripe Smut <i>Ustilago striiformis</i> | 4 – 6 | Make two applications at 14 day intervals when disease first appears. Water product into the root zone after application. For prevention, apply in the spring (just before grass begins growth), and in the fall. |

*Refer to the “Use Sites and Maximum Application Rates” table to determine allowable rates for each application.

**Use 4 – 6 oz rate for curative response to Basal Stem Anthracnose.

Turf Application Directions

Apply OHP 50 WP Fungicide uniformly over the area to be treated with a properly calibrated sprayer. Apply after mowing or avoid mowing 12 hours after application. Apply sufficient water to obtain thorough coverage, usually 1 ½ to 2 ½ gallons per 1,000 square feet of turf area. When treating golf greens, always treat aprons and approaches to golf greens. The minimum retreatment interval for turf applications is 14 days.

Use Sites and Maximum Application Rates

| SITE | MAXIMUM SINGLE APPLICATION RATE | | MAXIMUM YEARLY APPLICATION RATE oz/1000 sq ft | |
|---|---------------------------------|---------------------|---|---------------------|
| | oz/1000 sq ft | Pounds Product/Acre | Oz/1000 sq ft | Pounds Product/Acre |
| Golf Courses – Greens, Tees and Aprons | 6 | 16.33 | 16 | 43.6 |
| Golf Courses - Fairways (except Florida) only during overseeding | 4 | 10.8 | 4 | 10.88 |
| Golf Courses - Fairways (Florida only) | 2 | 5.44 | 2 | 5.44 |
| Residential and Public Areas (Home lawns, parks, athletic fields, schools and day care centers) | 2 | 5.44 | 8 | 21.76 |

Do not graze animals on treated turf. Do not feed clippings to livestock or poultry.

VEGETABLE TRANSPLANTS:

Not for use on field vegetable production

{note to reviewer-California restriction used until vegetable transplants approved for use in California}

[CALIFORNIA ONLY – This product is not for use on vegetable transplants.]

| Vegetable Transplants | Diseases Controlled | Rate of OHP 50 WP Fungicide | Application Instructions |
|--|---|--|---|
| Beans, dry and succulent including: Lima bean, Snap bean, Kidney bean, Mung bean, Navy bean, Pinto bean, Wax bean, Broad bean, Fava bean, Asparagus bean, Blackeyed pea, Cowpea, Sweet lupine, White lupine, White sweet lupine, Grain lupine, Chick pea, Garbanzo bean | Anthracnose <i>Colletotrichum</i> Gray Mold <i>Botrytis</i> White Mold <i>Sclerotinia</i> | 2.1-2.8 lbs/Acre ¹ or 1.4-2.1 lbs/Acre ² | For one application: Apply when 100% of plants have at least one open bloom or when conditions are favorable for disease development. or For multiple applications: Make the first application when 10% to 30% of plants have at least one open bloom and follow with sequential applications on a 4 to 7 day interval. Apply prior to the development of disease for best results. Do not apply more than 5.6 lbs. of product (2.8 lbs a.i./A/crop. |

| Vegetable Transplants | Diseases Controlled | Rate of OHP 50 WP Fungicide | Application Instructions |
|--|--|---------------------------------|--|
| | | | <p>[Pre-harvest interval: California only, 14 days for succulent beans, 28 days for dry beans and lima beans.]</p> <p>Pre-harvest interval: [all other states,] 14 days for succulent beans and lima beans, 28 days for dry beans.</p> |
| <p>CUCURBITS (Cantaloupe, Casaba, Cucumbers, Melons, Pumpkins, Summer and Winter Squash, and Watermelons)</p> | | | <p>Do not apply more than 4.2 lbs of product (2.1 lbs a.i.)/A/crop from any combination of application timings.</p> <p>Follow resistance management guidelines under Directions for Use.</p> |
| | <p>Anthracnose[*] <i>Colletotrichum</i> Gummy Stem Blight[*] <i>Didymella</i> Powdery Mildew[*] <i>Erysiphe,</i> <i>Sphaerotheca,</i> <i>Podosphaera</i> Target Spot[*] <i>Corynespora</i></p> | <p>0.7 lbs/Acre³</p> | <p>Begin applications when plants begin to run or when disease first appears, and repeat at 7 to 14 day intervals.</p> <p>For Target Spot use at 7 day intervals.</p> |
| | <p>Belly Rots[*] <i>Rhizoctonia,</i> <i>Fusarium</i></p> | <p>0.7 lbs/Acre³</p> | <p>Apply in sufficient volume to allow runoff to the soil. Will not control <i>Pythium</i> or <i>Phytophthora</i>.</p> |

[*Not for this use in California]

{note to reviewer-California restriction used until vegetable transplants approved for use in California}

1. Apply, for example in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 2.1 pounds per acre. If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 2.8 pounds per acre. For example, if 200 gallons of water are required, use 1.4 pounds per 100 gallons.
2. Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 1.4 pounds per acre. If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 2.1 pounds per acre. For example, if 200 gallons of water are required, use 1.05 pounds per 100 gallons.
3. Apply, for example, in 50-200 gallons of water per acre. In volumes of water below 50 gallons, use a minimum of 0.7 pounds per acre. If more than 200 gallons of water per acre are required for good plant coverage, apply a maximum rate of 0.7 pounds per acre. For example, if 200 gallons of water are required, use 0.35 pounds per 100 gallons.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container in a dry, temperature controlled, secure place.

PESTICIDE HANDLING: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

[Containers Less Than 5 Gallons]:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

CONTAINER DISPOSAL: Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

[Containers Greater Than 5 Gallons]:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying.

CONTAINER DISPOSAL: Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of OHP, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold OHP, Inc. and Seller harmless for any claims relating to such factors.

OHP, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or OHP, Inc., and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, OHP, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither OHP, Inc. or Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF OHP, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT,**

{Brackets indicate note to reviewer}

Label Amendment 8/12/2020

EPA Reg. No. 59807-6

SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF OHP, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

OHP, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of OHP, Inc.