

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Nova 40W
EPA Registration Number 62719-411
Your submission of November 17, 2006 adding peppers (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) and eggplant
OPPIN decision number 372358

The supplemental label for peppers (bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) and eggplant, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

2 8 3

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

< S >

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES							
SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

3 7 3

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Nova[®] 40W

EPA Reg. No. 62719-411

Control of Powdery Mildew on Peppers (Bell Pepper, Chili Pepper, Cooking Pepper, Pimento, Sweet Pepper) and Eggplant

(Not for Use in the States of Alaska, Arizona, California, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Nova[®] 40W fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Nova 40W according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Nova 40W.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Nova 40W (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Leveillula taurica</i>)	2.5 - 5	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. A lower rate and 14-day re-treatment interval may be used for small plants and under low disease pressure conditions. The highest rate and shorter re-treatment interval are recommended for large plants and severe or threatening disease conditions. If reliable predictive modeling (risk index) systems are available, these can help to indicate disease pressure conditions. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 5 oz (0.31 lb) of Nova 40W (0.125 lb ai) per acre per application or 20 oz (1.25 lb) of Nova 40W (0.5 lb ai) per acre per season. • Do not make more than 4 applications per season. • Pre-harvest Interval: None. • Minimum Re-treatment Interval: 10 to 14 days

Resistance Management: Nova 40W belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Nova 40W should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R866-006
EPA accepted ___/___/___
Initial printing

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

MAR 20 2008
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

62719-411

NEXT

LABEL

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D.
Regulatory Leader
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054

MAR 20 2008

Dear Dr. Brinkmeyer:

SUBJECT: Nova 40W
EPA Registration Number 62719-411
Your submission of November 17, 2006 adding okra
OPPIN decision number 372358

The supplemental label for okra, referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended is acceptable provided you make the following changes:

Move the "Attention" box so that it immediately follows the heading "Directions for Use."

Move the text beginning "Control of Powdery Mildew on . . ." so that it follows the new position of the "Attention" box.

This supplemental label must be incorporated into the main label within 18 months or at the next label printing, whichever comes first.

As a condition of registration you must provide the following ecotox data:

GRN 72-4c, Estuarine/marine life cycle (mysid), due July 2009.
GRN 162-3, Benthic Anaerobic Aquatic Metabolism, due July 2010.
GRN 162-4, Aerobic Aquatic Metabolism, due July 2010.

Please submit one copy of your final printed supplemental labeling before the product is released for shipment. A stamped copy of the label is enclosed for your records.

If you have any questions, please contact Lisa Jones on my team at (703) 308-9424 or jones.lisa@epa.gov.

Sincerely yours,

< S >

Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure: Stamped supplemental label

CONCURRENCES							
SYMBOL <	7505P						
SURNAME <	Lisa Jones						
DATE <	Mar 20, 2008						

Supplemental Labeling



Dow AgroSciences LLC 9330 Zionsville Road Indianapolis, IN 46268-1054 USA

Nova[®] 40W

EPA Reg. No. 62719-411

Control of Powdery Mildew on Okra

(Not for Use in the States of Alaska, Arizona, California, Hawaii, Idaho, Montana, Nevada, Oregon, Utah, Washington, Wyoming)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Nova[®] 40W fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Nova 40W according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Nova 40W.

Directions for Use

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Nova 40W (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Sphaerotheca fuliginea</i>)	2.5 - 5	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. A lower rate and 14-day re-treatment interval may be used for small plants and under low disease pressure conditions. The highest rate and shorter re-treatment interval are recommended for large plants and severe or threatening disease conditions. If reliable predictive modeling (risk index) systems are available, these can help to indicate disease pressure conditions. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 5 oz (0.31 lb) of Nova 40W (0.125 lb ai) per acre per application or 20 oz (1.25 lb) of Nova 40W (0.5 lb ai) per acre per season. • Do not make more than 4 applications per season. • Pre-harvest Interval: None. • Minimum Re-treatment Interval: 10 to 14 days

Resistance Management: Nova 40W belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Nova 40W should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R866-005
EPA accepted ___/___/___
Initial printing

ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAR 20 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

62719-411