1/2

ENVIRONMENTAL PROTECTION AGENCY

Raymond S. Brinkmeyer, Ph.D. Regulatory Leader Dow AgroSciences, LLC 9330 Zionsville Road Indianapolis, IN 46268-1054

JAN 0 3 2008

Dear Dr. Brinkmeyer:

SUBJECT:

Laredo EC

EPA Registration Number 62719-412 Your submission dated April 4, 2007

Supplemental label increasing application rate for cotton seed

treatment

OPPIN Decision Number 378128

The supplemental label referred to above, submitted in connection with registration under section (3) of the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA) is acceptable provided that the supplemental labeling must be incorporated into the Section 3 master label within 18 months or at the next label printing, whichever comes first.

Please submit one copy of your final printed labeling before you release the product for shipment. A stamped copy of the label is enclosed for your records. If you have any questions, please contact Lisa Jones on my team by phone at (703) 308-9424 or by e-mail at jones.lisa@epa.gov.

Sincerely yours,

Mary L. Waller Product Manager (21) Fungicide Branch Registration Division (7505C)

Enclosure:

Stamped supplemental label

CONCURRENCES					
SYMBOL <	7505P				
SURNAME <	Lisa Jones				
DATE <	Dec 20, 2007				

EPA Form 1320-1 (12-70)

OFFICIAL FILE COPY

Supplemental Labeling



Dow AgroSciences LLC:

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Laredo® EC

EPA Reg. No. 62719-412

Cotton Seed Treatment

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of pesticide application.
- Read the label affixed to the container for Laredo® EC fungicide before applying. Carefully follow all
 precautionary statements and applicable use directions.
- Use of Laredo EC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Laredo EC is recommended as a seed treatment to protect against sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*) which impairs good cotton seed germination and seedling development. Laredo EC can be tank mixed with other properly labeled cotton seed protectants such as Apron TL fungicide for Pythium control. Laredo EC is a liquid formulation that can be pumped or poured directly into the cotton seed treater or diluted with water, provided the recommended amount of product is applied.

Commercial Cotton Seed Treatment

Label Treated Seed "This seed is treated with Laredo EC which contains the active ingredient myclobutanil. Do not use for food, feed, oil production or any other purpose except planting. Do not graze cotton plants grown from treated seed. Store away from feed and foodstuffs."

In California, this product must be applied through a closed system such as WECO-RED (Returnable Evacuation Device).

Use 1.25 to 4 fl oz of Laredo EC (0.019 to 0.062 lb active ingredient) per 100 lb of cotton seed to reduce seedling damage caused by sore shin (*Rhizoctonia solani*) and black root rot (*Thielaviopsis basicola*).

For cotton seed treatment, always use Laredo EC with an EPA approved dye or colorant that imparts an unnatural color to the seed.

On Farm Cotton Seed Treatment

For mechanical treaters or hopper box treatments, dilute 1.25 to 4 oz of Laredo EC (0.019 to 0.062 lb active ingredient) with 8 to 10 fl oz of water and apply to 100 lb of cotton seed. Other cotton seed hopper box products as DeltaCoat AD or DeltaCoat ND may be applied with Laredo EC. Always follow the label recommendations on the respective products.

Chemigation: Do not apply this product through any type of irrigation system.

Trademark of Dow AgroSciences Ll	_C
R153-085 Accepted:// Initial printing.	

ACCEPTED

JAN 0 3 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

C2719-417

Page 1 of 1