

Until Reregistration

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

MAXFORCE® Professional Insect
Control™ Roach Gel

PM13
6 4248-5

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

The Clorox Company
7200 Johnson Drive
Pleasanton, CA 94588

~~1166/5918796~~
18/1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) since you have agreed in writing to the conditions outlined in our letter dated March 20, 1992.

Submit five copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (H7505C)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

CSA 7/23/92

2 of 10

**MAXFORCE® Professional Insect Control™
Roach Gel**

Carton for Triple-Pack of Plastic Dispensers
(front)

MAXFORCE® Professional Insect Control™ Roach Gel

MAXFORCE® Gel may be used as a spot or crack and crevice treatment for the indoor and outdoor control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments including hospitals, laboratories, computer facilities, sewers, buses, boats/ships, trains, pet shops and zoos.

ACTIVE INGREDIENT: Hydramethylnon [tetrahydro-5,5-dimethyl-2(1H)-pyrimidinone {3-[4-(trifluoromethyl)phenyl]-1-{2-[4-(trifluoromethyl)phenyl]ethenyl}-2-propenylidene}hydrazonone.....	2.15%
INERT INGREDIENTS.....	97.85%
TOTAL.....	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See directions on back panel
Use only in areas not easily accessible to children and pets
Product will stain porous surfaces

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid contact with skin and clothing. Wear impervious gloves (plastic or rubber) during application. Keep exposed gel away from open food and food contact surfaces. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, drink two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. In case of emergency, call day or night, 800-426-6228.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wrap used dispensers and place in a trash can. Store in cool, dry area.

Contains 3 pre-filled plastic dispensers
Total Net Wt. 6.3 oz. (180 g)
©1991 MAXFORCE Insect Control Systems
1221 Broadway
Oakland, CA 94612

EPA Reg. No. 64243-5

ACCEPTED
with COMMENTS
in EPA Letter Dated



JUL 23 1992

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
64243-5

3 of 10

**MAXFORCE® Professional Insect Control™
Roach Gel**

**Carton for Triple-pack of Plastic Dispensers
(back)**

MAXFORCE® Professional Insect Control™ Roach Gel

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use MAXFORCE® Gel only in areas not easily accessible to children and pets. Wear impervious gloves (plastic or rubber) when applying product. Keep exposed gel away from open food and food contact surfaces.

Do not use in food/feed product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared, processed, stored or exposed. Do not use in serving areas while food/feed is exposed. (Serving areas are considered areas prepared foods are served such as dining rooms, but exclude areas where food may be prepared or held.)

NOTE: Do not place bait in locations that are routinely washed, as bait will be removed by washing. Do not place treatments in locations where routine cleansing operations may transfer diluted bait to food or food preparation surfaces. Do not treat food preparation surfaces.


Non-food areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Avoid contact with porous surfaces as bait may stain.

Not for use in USDA Meat and Poultry plants.

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

MAXFORCE® Gel is packaged in a plastic dispenser containing 2.1 oz. (60 g) of bait. Application rate is dependent on level of infestation and species to be controlled. Bait can be applied as spots or as a bead in cracks and crevices.

Carton for Triple Pack of Plastic Dispensers (beck cont.)

<u>Species</u>	<u>Application Rate (per sq. yd.)</u>	<u>Recommended Spot Size (4-1/2g)</u>
Small, such as German or brown-banded	1-2 spots, or 3-6 inch bead	
Large, such as American, smokey-brown or oriental	2-4 spots, or 6-12 inch bead	

The lower application rate should be used for low to moderate infestations or retreatments and the higher rate for more severe infestations or where the population occurs in inaccessible areas, such as in walls. Gel placements should be at or near harborages or aggregation areas, such as corners, areas of movement or cracks and crevices. Numerous smaller placements will provide faster control than few larger spots, especially for German cockroach control. For outdoor use place in areas adjacent to structure where roaches may rest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints.

Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushings.

To apply gel, remove the cap on the nozzle, touch the tip to surface, and depress the plunger. Recap the dispenser after treatment is completed. The bait will adhere to non-greasy or non-dusty surfaces and will remain pliable and palatable to cockroaches as long as it is visibly present. Reapply when bait is no longer visibly present, according to the remaining level of infestation. A visual inspection of bait placements is recommended at one month after the initial treatment. Reapplication should occur if the bait is visibly depleted.

Pest population reduction will be apparent within a week or less.

**MAXFORCE® Professional Insect Control™
Roach Gel**

**Carton for Single Plastic Dispenser
(back)**

MAXFORCE® Professional Insect Control™ Roach Gel


DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MAXFORCE® Gel may be used as a spot or crack and crevice treatment for the indoor and outdoor control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments including hospitals, laboratories, computer facilities, sewers, buses, boats/ships, trains, pet shops and zoos.

Use MAXFORCE® Roach Control Gel only in areas not easily accessible to children and pets. Wear impervious gloves when applying the product. Keep exposed gel away from open food and food contact surfaces.

MAXFORCE® Gel is packaged in a plastic dispenser containing 2.1 oz. (60 g) of bait. Application rate is dependent on level of infestation and species to be controlled. Bait can be applied as spots or as a bead in cracks and crevices. For outdoor use place bait in areas adjacent to structures where roaches may rest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints.

<u>Species</u>	<u>Application Rate (per sq. yd.)</u>	<u>Recommended Spot Size (1/4-1/2g)</u>
Small, such as German, or brown-banded	1-2 spots, or 3-6 inch bead	
Large, such as American, smokey-brown or oriental	2-4 spots, or 6-12 inch bead	

6 of 10

**MAXFORCE® Professional Insect Control™
Roach Gel**

**Carton for Single Plastic Dispenser
(Side)**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: May be harmful if swallowed. Avoid contact with skin and clothing. Wear impervious gloves (plastic or rubber) during application. Keep exposed gel away from open food and food contact surfaces. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, drink two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. In case of emergency, call day or night, 800-426-6228.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wrap used dispensers and place in a trash can. Store in cool, dry area.

MAXFORCE® Professional Insect Control™
Roach Gel

7-10

Carton for Single Plastic Dispenser
(Front)

MAXFORCE® Professional Insect Control™ Roach Gel

MAXFORCE® Gel may be used for as a spot or crack and crevice treatment for the indoor and outdoor control of cockroaches in residential areas and the non-food/feed areas of institutional, warehousing and commercial establishments. Read insert for full use directions.

ACTIVE INGREDIENT: Hydramethylnon [tetrahydro-5,5-dimethyl-2(1H)-pyrimidinone {3-[4-(trifluoromethyl)phenyl]-1-[2-[4-(trifluoromethyl)phenyl]ethenyl]-2-propenylidene}hydrazonone.....2.15%

INERT INGREDIENTS.....97.85%

TOTAL.....100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See precautions on side panel

Use only in areas not easily accessible to children and pets

Product will stain porous surfaces

Contains 1 pre-filled plastic dispenser.

Net Wt. 2.1 oz. (60 g)

©MAXFORCE Insect Control Systems
1221 Broadway
Oakland, CA 94612

EPA Reg. No. 64248-5

EPA Est. No. 35132-OH-001



8/10

**MAXFORCE® Professional Insect Control™
Roach Gel**

Plastic Dispenser Label

MAXFORCE® Professional Insect Control™ Roach Gel

ACTIVE INGREDIENT: Hydramethylnon [tetrahydro-5,5-dimethyl-2(1H)-pyrimidinone
{3-[4-(trifluoromethyl)phenyl]-1-[2-[4-(trifluoromethyl)phenyl]ethenyl]-2-
propenylidene]hydrazone..2.15%
INERT INGREDIENTS.....97.85%
TOTAL.....100.00%

KEEP OUT OF REACH OF CHILDREN

Use only in areas not easily accessible to children and pets

Product will stain porous surfaces

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with skin and clothing. Wear impervious gloves (plastic or rubber) during application. Keep exposed gel away from open food and food contact surfaces. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, drink two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. In case of emergency, call day or night, 800-426-6228.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wrap used dispensers and place in a trash can. Store in cool, dry place.

Net Wt. 2.1 oz. (60 g)

©1991 MAXFORCE Insect Control Systems
1221 Broadway
Oakland, CA 94612


EPA Reg. No. 64248-5

EPA Est. No. 35132-OH-001



Carton for Single Plastic Dispenser (insert cont.)

9 of 10

<u>Species</u>	<u>Application Rate (per sq. yd.)</u>	<u>Recommended Spot Size (4-1/2g)</u>
Small, such as German or brown-banded	1-2 spots, or 3-6 inch bead	
Large, such as American, smokey-brown or oriental	2-4 spots, or 6-12 inch bead	

The lower application rate should be used for low to moderate infestations or retreatments and the higher rate for more severe infestations or where the population occurs in inaccessible areas, such as in walls. Gel placements should be at or near harborage or aggregation areas, such as corners, areas of movement or cracks and crevices. Numerous smaller placements will provide faster control than few larger spots, especially for German cockroach control. For outdoor use, place bait in areas adjacent to structure where roaches may rest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints.

Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushing.

To apply gel, remove the cap on the nozzle, touch the tip to surface, and depress the plunger. Recap the dispenser after treatment is completed. The bait will adhere to non-greasy or non-dusty surfaces and will remain pliable and palatable to cockroaches as long as it is visibly present. Reapply when bait is no longer visibly present, according to the remaining level of infestation. A visual inspection of bait placements is recommended at one month after the initial treatment. Reapplication should occur if the bait is visibly depleted.

Pest population reduction will be apparent within a week or less.

10 of 10

**MAXFORCE® Professional Insect Control™
Roach Gel**

**Carton for Single Plastic Dispenser
(insert)**

MAXFORCE® Professional Insect Control™ Roach Gel

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use MAXFORCE® Gel only in areas not easily accessible to children and pets. Wear rubber gloves when applying product.

Do not use in the food/feed product areas of food/feed processing plants, restaurants or other areas where food/feed is commercially prepared, processed, stored or exposed. Do not use in serving areas while food/feed is exposed. (Serving areas are considered areas where prepared foods are served such as dining rooms, but exclude areas where food may be prepared or held.

NOTE: Do not place bait in locations that are routinely washed, as bait will be removed by washing. Do not place treatments in locations where routine cleansing operations may transfer diluted bait to food or food preparation surfaces. Do not treat food preparation surfaces.

Non-food areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Avoid contact with porous surfaces as bait may stain.

Not for use in USDA Meat and Poultry plants.

Do not use this product in or on electrical equipment due to the possibility of shock hazard.

MAXFORCE® Gel is packaged in a plastic dispenser containing 2.1 oz. (60 g) of bait. Application rate is dependent on level of infestation and species to be controlled. Bait can be applied as spots or as a bead in cracks and crevices.