

69361-37

9/13/2012

1/16



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Chemical Safety and Pollution Prevention
Office of Pesticide Programs
Registration Division (7504P)
1200 Pennsylvania Ave., N.W.
Washington, DC 20460

EPA Reg. Number:

69361-37

Date of Issuance:

SEP 13 2012

Term of Issuance:

Unconditional

Name of Pesticide Product:

**Tebuconazole 45DF
Fungicide**

NOTICE OF PESTICIDE:

- Registration
 - Reregistration
- Under FIFRA, as amended

Name and Address of Registrant (include ZIP Code):

**Repar Corporation
P.O. Box 4321
Silver Spring, MD 20914**

Mailed to:

**Bushan Mandava
Mandava Associates
1050 Connecticut Ave., Suite 1000
Washington, DC 20036-4549**

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

Signature of Approving Official:

Mary L. Waller

Mary L. Waller, Product Manager (21)
Fungicide Branch/Registration Division/OPP/OCSP (7504P)

Date:

9/13/2012

2/16

2. You must submit the following data before the due date of 4/1/2014:
 - a. Storage Stability (830.6317) and Corrosion Characteristics (830.6320) studies.
3. Make the following changes to the label:
 - a. Change the product registration number to "EPA Reg. No. 69361-37"
 - b. On page 1 immediately below the product name add (in smaller print): "Water Dispersible Granules"
 - c. In the **FIRST AID** box, move the entry for **If on skin or clothing** to below the entry for **If Inhaled**.
 - d. On page 13 in the notes for Plums (Post Harvest) , in the last line change "...do not rise." to "...do not rinse."
4. Submit one copy of the revised final printed label for the record before the product is released for shipment.

Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "Accepted with Comments" is enclosed for your records.



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7504P)

Enclosure:

Label stamped "Accepted with Comments"
Product Chemistry Review DP404231 dated August 17, 2012
Acute Toxicity Review DP404233 dated August 28, 2012

3/16



TEBUCONAZOLE 45DF FUNGICIDE

For control of specified diseases on fruit crops

ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....45%

OTHER INGREDIENTS:55%

TOTAL:100.0%

KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (TO THE USER: If you do not understand the label, find someone to explain it to you in detail.)

For **PRODUCT USE** information call 1-866-248-7426

FIRST AID

If in the eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing the eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

Note to Physician: No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

Have a product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergency Assistance, call the National Pesticide Information Center 1-800-858-7378

For chemical emergency: spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

Manufactured For:
 Repar Corporation
 P.O. Box 4321
 Silver Spring, MD 20914

ACCEPTED
 with **COMMENTS**
 In EPA Letter Dated:
 9/13/2012

Under the Federal Insecticide, Fungicide, and Rodenticide Act, EPA has established for the pesticide registered under EPA Reg. No.

EPA Reg. No.: 69361-

NET WEIGHT: LBS

69361-37

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Harmful if swallowed, inhaled or absorbed through the skin. Causes substantial but temporary eye injury. Prolonged or frequently repeated skin contact may cause allergic skin reactions in some individuals. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA Chemical Resistance Category Selection Chart.

Mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Shoes plus socks,
- Use goggles or face shield.

Remove and wash contaminated clothing before reuse. Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory: Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Surface Water Label Advisory: This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- Shoes plus socks,
- Protective eyewear.

PRODUCT INFORMATION

Spray Volume: Apply TEBUCONAZOLE 45DF FUNGICIDE in 50 gallons or more of spray solution per acre by ground spray equipment or in 15 gallons or more per acre by aircraft equipment except as noted under "REMARKS" for each crop. Increase spray volume as crop growth increases to ensure through coverage of foliage and fruit. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied.

Aerial application is not allowed in the State of New York.

Chemigation: Do not apply this product through any type of irrigation system.

Mixing: Add labeled amount of TEBUCONAZOLE 45DF FUNGICIDE into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the TEBUCONAZOLE 45DF FUNGICIDE should be thoroughly dispersed in the spray tank prior to the addition of any other materials.

Compatibility: The following procedure should be followed when determining the compatibility of TEBUCONAZOLE 45DF FUNGICIDE with other products: Pour the recommended proportions of the products into a suitable container of water. Mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information, contact your local Repair Corporation representative.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS; MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Spray Drift Management: The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Buffer Zone Requirements: For soil or foliar applications, do not apply by ground within 25 feet, or by air within 150 feet of lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries and commercial fish farm ponds.

Recommendations for Aerial Applications: The spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Importance of Droplet Size: An important factor influencing drift is droplet size. Small droplets (<150 - 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Wind Speed Restrictions: Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Avoiding applications when wind direction is toward the aquatic area can reduce risk of exposure to sensitive aquatic areas.

Restrictions During Temperature Inversions: Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF TEBUCONAZOLE 45DF FUNGICIDE
Grape	Black Rot (<i>Guignardia bidwellii</i>)	4 oz./A
	Powdery mildew (<i>Uncinula necator</i>)	
	Botrytis (Suppression)	

Powdery Mildew

Apply in a preventive spray schedule. Make the first application of TEBUCONAZOLE 45DF FUNGICIDE before bloom and continue applications using spray intervals of up to 21 days in low to moderate disease pressure. Use a 14-day schedule when disease pressure is severe.

Black Rot

Preventive Schedule: Make the first application at 1 to 3 inches of new shoot growth and continue at 7- to 14 day intervals through 5⁰ Brix stage or until veraison (berry coloring) is complete. Apply at 1 inch new growth and 7- to 10-day intervals on highly susceptible varieties or under severe disease conditions.

Post-Infection Schedule: A post-infection schedule may be followed from 1-inch new shoot growth through 5⁰ Brix stage. Apply within 72 hours after the beginning of an infection period. TEBUCONAZOLE 45DF FUNGICIDE applications should not be closer than 7 days apart. Continue TEBUCONAZOLE 45DF FUNGICIDE applications using the preventive schedule if the post-infection schedule is discontinued.

Botrytis

TEBUCONAZOLE 45DF FUNGICIDE, applied in a powdery mildew spray schedule, will enhance the activity of registered fungicides used for control of Botrytis bunch rot. Applications must be made on a 14-day schedule for suppression of Botrytis.

APPLICATION RESTRICTIONS: TEBUCONAZOLE 45DF FUNGICIDE may be applied up to 14 days before harvest. A maximum of 2 pounds of TEBUCONAZOLE 45DF FUNGICIDE may be applied per acre per crop season. **The minimum interval between applications is 7 days.**

Apply product in sufficient water for thorough coverage of vines and fruit. Increase spray volume as vine growth increases.

Restricted-entry interval (REI) = 12 hours

RESISTANCE MANAGEMENT STRATEGY: TEBUCONAZOLE 45DF FUNGICIDE is a sterol demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tank-mixing TEBUCONAZOLE 45DF FUNGICIDE with other DMI fungicides is not recommended.

Almond	Brown rot blossom blight (<i>Monilinia laxa</i> , <i>M. fructicola</i>)	8 oz./A
<p>Blossom blight: Begin application at pink bud. If the bloom period is extended, and/or severe disease conditions exist, make a second application at full bloom. If conditions remain favorable for disease, make another application at petal fall.</p> <p>APPLICATION RESTRICTIONS: Apply TEBUCONAZOLE 45DF FUNGICIDE in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after petal fall is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.</p> <p>A maximum of 32 ounces of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. TEBUCONAZOLE 45DF FUNGICIDE may be applied up to 35 days before harvest.</p> <p>Restricted-entry interval (REI) = 12 hours</p> <p>RESISTANCE MANAGEMENT STRATEGY: TEBUCONAZOLE 45DF FUNGICIDE is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing TEBUCONAZOLE 45DF FUNGICIDE with other DMI fungicides is not recommended.</p>		
Pistachio	Botryosphaeria panicle and shoot blight (<i>Botryosphaeria dothidea</i>)	8 oz./A
<p>Botryosphaeria: Begin application at 25 to 50% bloom and repeat again 10 to 14 days later to protect young flower clusters and fruit. Make two additional applications of REPAR 45DF FUNGICIDE 14 days apart beginning 49 days before harvest.*</p> <p>APPLICATION RESTRICTIONS: Apply TEBUCONAZOLE 45DF FUNGICIDE in a minimum spray volume of 15 gallons per acre by air or 50 gallons per acre by ground. Reduce the application interval for varieties that are highly susceptible to the indicated diseases or when severe disease conditions exist. The use of ground application after flowering is preferred because of difficulty in penetrating the canopy and obtaining thorough coverage of the foliage and fruit by air.</p> <p>A maximum of 32 ounces of product may be applied per acre per crop season. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. TEBUCONAZOLE 45DF FUNGICIDE may be applied up to 35 days before harvest.</p> <p>Control of insect vectors and a thorough pruning program to remove plant tissue infected by <i>Botryosphaeria</i> are critical for optimum control of this disease.</p> <p>Restricted-entry interval (REI) = 12 hours</p> <p>RESISTANCE MANAGEMENT STRATEGY: TEBUCONAZOLE 45DF FUNGICIDE is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Tankmixing TEBUCONAZOLE 45DF FUNGICIDE with other DMI fungicides is not recommended.</p>		

13/16

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCONAZOLE 45DF FUNGICIDE	
Pome Fruits: Apple Crabapple Loquat Mayhaw Pear Pear Oriental Quince	Scab (<i>Venturia</i> spp.) Cedar-apple rust (<i>Gymno-sporangium juniperi-virginianae</i>) Powdery mildew (<i>Podosphaera leucotricha</i>)	4 to 8 oz./acre*	2 oz./100 gal.**

Scab:

Preventive Schedule: Apply at 7- to 10-day intervals from green tip through the second cover spray.

Post-Infection Schedule: Apply within 72 hours after the beginning of a scab infection period. TEBUCONAZOLE 45DF FUNGICIDE applications should not be closer than 7 days apart. Continue TEBUCONAZOLE 45DF FUNGICIDE applications using the preventive schedule if the post-infection schedule is discontinued.

Cedar-apple rust: Apply at 7- to 10-day intervals from pink stage through the second cover spray.

Powdery mildew: Apply at 7- to 10-day intervals from tight cluster through the second cover spray.

APPLICATION RESTRICTIONS:

*The amount of TEBUCONAZOLE 45DF FUNGICIDE required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated.

**For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume.

TEBUCONAZOLE 45DF FUNGICIDE may be applied up to 75 days before harvest. A total of 3 pounds of TEBUCONAZOLE 45DF FUNGICIDE may be applied per acre per crop season.

Restricted-entry interval (REI) = 5 days

RESISTANCE MANAGEMENT STRATEGY: TEBUCONAZOLE 45DF FUNGICIDE is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Do not mix or alternate TEBUCONAZOLE 45DF FUNGICIDE with other DMI fungicides.

14/16

APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCONAZOLE 45DF FUNGICIDE	
Stone Fruits: Apricot Cherry Nectarine Peach Plum Plumcot Prune (fresh)	Brown rot (blossom blight, fruit rot) <i>(Monilinia laxa, M. fructicola)</i> Cherry Leaf Spot <i>(Blumeriella jaapii)</i> Powdery mildew <i>(Podosphaera spp., Sphaerotheca pannosa)</i>	4 to 8 oz./acre*	2 oz./100gal.**
Plum Plumcot Prune (fresh) Peach	Rust (<i>Tranzschelia discolor</i>)	5 to 8 oz./acre*	2 oz./100gal.**

Blossom blight: Begin application at white bud on cherry, pink bud on peach and nectarine, and green tip on plums and prunes. Apply again at 50% bloom and at petal fall if conditions continue to be favorable for disease development.

Fruit rot: Begin application at two to three weeks before harvest and continue at 7-day intervals through the day of harvest. The blossom and fruit stages must be protected for optimum control of brown rot. If TEBUCONAZOLE 45DF FUNGICIDE is applied during only one of these stages, another registered fungicide should be applied to the other stage to provide optimum protection. Additional cover sprays during the early post-bloom period are also important for preventing quiescent fruit infections on sweet cherry and peach.

Leaf spot: Begin application at petal fall or when first leaves unfold and continue applications at 7- to 14-day intervals. Applications should be made at 7-day intervals early in the growing season when terminal growth is rapid and/or under severe disease conditions. A post-harvest application may be made to maintain control and reduce overwintering inoculum.

Powdery mildew: Follow leaf spot schedule until terminal growth ceases.

Rust: Begin applications after canker emergence and continue applications at 14-day intervals under severe disease conditions.

APPLICATION RESTRICTIONS:

TEBUCONAZOLE 45DF FUNGICIDE may be applied up to and including the day of harvest (0% PHI). Do not apply more than 3 lb. of product per acre per crop season.

* The amount of TEBUCONAZOLE 45DF FUNGICIDE required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees. For smaller trees, multiply 2 ounces times the number of 100 gallons of spray solution required to thoroughly wet, to the point of runoff, one acre of the trees being treated. For concentrate sprays, apply the same amount of product per acre as would be applied in a dilute spray based on tree size and foliage volume, but not less than 4 ounces of TEBUCONAZOLE 45DF FUNGICIDE per acre. Apply 6 to 8 ounces per acre of TEBUCONAZOLE 45DF FUNGICIDE when severe disease conditions exist.

** When using the 2 oz./100 gal. rate, use at least 4 ounces of TEBUCONAZOLE 45DF FUNGICIDE per acre.

Restricted-entry interval (REI) = 5 days

RESISTANCE MANAGEMENT STRATEGY: TEBUCONAZOLE 45DF FUNGICIDE is a demethylation inhibitor (DMI) fungicide. It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Do not mix TEBUCONAZOLE 45DF FUNGICIDE with other DMI fungicides.

Cherry (Post-Harvest)	Brown rot (<i>Monilinia</i> spp.)	8.0 oz. per 25,000 lb. of fruit
	Grey Mold (<i>Botrytis cinerea</i>)	
	Rhizopus rot (<i>Rhizopus</i> spp.)	

Mix specified rate of TEBUCONAZOLE 45DF FUNGICIDE in 100 gallons of water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion). Apply through a spray or flood system on a commercial packing line after stem cutting, sizing, sorting or washing. For spray treatments, use conventional spray equipment such as T-Jet, flat-fan, or flood jet spray nozzles. Use enough fruit coating material to fully coat fruit. Workers handling treated fruit need to wear waterproof gloves. Do not apply more than once prior to boxing and do not rinse.

Plums (Post-Harvest) Including Chickasaw, Damson, Japanese, Plumcot, Prune(Fresh)	Brown rot (<i>Monilinia</i> spp.)	8.0 oz. per 200,000 lb. of fruit
	Grey Mold (<i>Botrytis cinerea</i>)	
	Rhizopus rot (<i>Rhizopus</i> spp.)	

Mix specified rate of TEBUCONAZOLE 45DF FUNGICIDE in 100 gallons of water, with or without wax. Maintain TEBUCONAZOLE 45DF FUNGICIDE suspension by direct or by-pass agitation sufficient to prevent settling of product in spray mixture. Apply as spray to fruit as they travel along packing line. After sizing, sorting or washing, TEBUCONAZOLE 45DF FUNGICIDE application may be incorporated into a waxing system using conventional equipment that provides for even and complete coverage of fruit. Do not apply more than once prior to boxing and do not rise.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry, secure place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed below. In spill or leak incidents, keep unauthorized people away.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved disposal facility.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

