



70126 -1

10/09/2003

1/7

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

OCT 9 2003

Organic Resources, Inc.
c/o Sharon M. Johnston, Esq.
509 Tower Valley Drive
Hillsboro, MO 63050

**Subject: Label Amendment Updating First Aid per PR Notice 2001-1 and Clarifying
Directions for Use to Comply with Other Agency Requirements
Organic Resources Multipurpose Insecticide
EPA Reg. No. 70126-1
Your Submission, Dated July 10, 2003**

Dear Ms. Johnston:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comments below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

1. Revise the product name on this master label to agree with the primary brand name of record: "Organic Resources Multipurpose Insecticide."
2. Move all of the statements in the second paragraph of the "Hazards to Humans & Domestic Animals" section ("Do not use on dogs or cats under twelve (12) weeks old. Consult a veterinarian ... If signs continue, consult a veterinarian immediately.") to the directions for use on dogs and cats. You must also change the heading "Dogs Cats and Pets" on the last page to "Dog and Cats", since the product may not be used on other pets.
3. You are encouraged to revise the disposal instructions for "Containers 1 Pound and Smaller" as recommended in PR Notice 2001-6 for residential use products in non-pressurized containers:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or [insert a toll free number which meets the criteria in paragraph II.E. of PR Notice 2001-6] for disposal instructions.
Never place unused product down any indoor or outdoor drain.

4. Correct the following spelling errors noted during our review:

- Leaftiers (2 places) under "Vegetable Crops on page 2
- Chervil under "Leaves of Root and Tuber Vegetables" on page 2
- Kohlrabi under "Brassica (Cole) Leafy Vegetables" on page 2
- Begonia and Petunia under "Ornamentals" on page 3
- then in the first line of the Container Disposal instruction on the last page (i.e., "... then offer for recycling ...").

If you have any questions regarding this action, please contact Susan Stanton of my team at (703) 305-5218.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505C)

Enclosure

ORGANIC RESOURCES

ALL-CROP

MULTIPURPOSE COMMERCIAL & AGRICULTURAL INSECTICIDE

CONTAINS DIATOMACEOUS EARTH, PYRETHRIN, AND PIPERONYL BUTOXIDE

For use indoors and outdoors including but not limited to: homes, animal quarters, on pets, grasses, lawns & turf (including golf courses), gardens, vegetable & growing crops (outdoors & greenhouses), ornamentals, specialty crops, field grain crops, fruit, nut, vine and oriental vegetable crops.

ACTIVE INGREDIENTS:

Pyrethrin.....	0.2%
Piperonyl Butoxide*	1.0%
Silicon Dioxide (from Diatomaceous Earth).....	82.9%
OTHER INGREDIENTS.....	15.9%
Total.....	100.0%

* Equivalent to 0.8 (butylcarbityl)(6-propylpiperonyl) ether and 0.2% related compounds.

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

READ ENTIRE LABEL BEFORE EACH USE

Pet Use Applies to Use Only on Dogs & Cats

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Use adequate ventilation when applying this product indoors.

Do not use on dogs or cats under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Personal Protective Equipment:

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes and socks and a NIOSH approved respirator with any N, R, P or HE filter. Follow manufacturer's instructions for cleaning and maintaining personal protective equipment. If no instructions for washing are given, use detergent and hot water. Keep and wash Personal Protective Equipment separately from other laundry.

User Safety Recommendations

User Should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove Personal Protective Equipment immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothes.

Environmental Hazards:

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from the treatment area. Do not contaminate water when cleaning equipment or when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

with COMMENTS

In EPA Letter Dated:

OCT 9 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

70126-1

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.	

See Side/Back Panel for Additional Precautions

WARRANTY STATEMENT

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and conditions of application equipment will vary, seller makes no claims other than those on this label.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are: Coveralls, waterproof gloves, shoes and socks. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nursery or greenhouses. Keep unprotected persons children and pets out of the treated area until sprays have dried

GENERAL INFORMATION: Carefully follow the directions for each type of use. This material may be used on edible crops growing outdoors or in greenhouses, up to and including the day of harvest. Use rates will vary with insect populations and amount of foliage.

Do not apply in classrooms when in use. Do not apply this product in hospital patient rooms while occupied or in any rooms occupied by the elderly or infirm.

USE RATES FOR VEGETABLES, FIELD CROPS AND MISCELLANEOUS CROPS: Use 2 to 6 pounds of product per acre depending upon the insect population and amount of foliage. The material can be used either:

AS A WETTABLE POWDER, apply in sufficient water to obtain maximum coverage. Dilution of the material should be at the maximum rate of one half pound in one half gallon of water or at a minimum of one pound to 25 gallons of water (4 Tbsp. per gallon). It is best to keep the material agitated during application. Apply the material uniformly to both the upper and lower leaf surfaces. Reapply only as necessary. Rates will vary with insect populations and amount of foliage, but adequate control can be obtained at 2 to 6 pounds product per acre.

AS A DUST, the best results are obtained when the plants are wet or dew still remains on the leaf surface. It is suggested that the plants are misted prior to application to obtain the maximum benefits and use the least amount of product for control.

VEGETABLE CROPS:

FOR THE CONTROL OF INSECTS SUCH AS: Ants, Aphids, Mexican Bean Beetle, 12 Spotted Cucumber Beetle, Army Worms, Beet Webworms, Blister Beetles, Cabbage Looper, Cabbage Worms, Caterpillars, Celery Leafliars, Colorado Potato Beetle, Corn Earworm, Cucumber Beetles, Diamond Black Moth Larva, European Corn Borer, Fire Ants, Flea Beetles, Garden Fleahoppers, Harlequin Bugs, Imported Cabbageworms, Japanese Beetles, Leafhoppers, Leaf Miners, Leafliars, Loopers, Lygus Bugs, Mites, Oblique-banded Leafrollers, Plant Bugs, Stink Bugs, Squash Vine Borers, Thrips, Vegetable Weevils, Webworms, and White Flies.

GROWING CROPS (Outdoor and Greenhouses):

Root and Tuber Vegetables: Including but not limited to Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Beets, Sugar Beets, Edible Burdock, Carrots, Cassava (Bitter & Sweet), Celeriac (Celery Root), Chervil (Turnip Root), Chicory, Chufa, Dasheen, Ginger, Ginseng, Horseradish, Laren, Parsely (Turnip Rooted), Parsnip, Potato, Radish, Japanese Radish, Rutabaga, Salsify, Black Salsify, Sweet Potato, Tanier, Tarrow Root, Tumeric, Turnip, Yam, Bean Yam.

Leaves of Root and Tuber Vegetables: Including but not limited to Beet, Sugar Beet, Edible Burdock, Carrot, Cassava (Bitter & Sweet), Celeriac, Cheryil, Chicory, Dasheen, Parsnip, Radish, Japanese Radish, Rutabaga, Black Salsify, Sweet Potato, Tanier, Turnip and Yam (True).

Bulb Vegetable: Including but not limited to Garlic, Leek, Onion (Bulb & Green), Shallot.

Leafy Vegetables: Including but not limited to Amaranth, Leafy Amaranth, Chinese Spinach, Tampala, Arrugula, Celery, Cettuce, Chervil, Cilantro, Corn Salad, Chrysanthemum (edible leaves), Chrysanthemum Garland, Cress (Garden), Upland Cress (Yellow Rocket Winter Cress), Dandelion, Dock, Endive, Fennel, Lettuce (Head and Leafy), Orach, Parsley, Purslane (Garden and Winter), Rhubarb, Spinach, Fine Spinach (Melabar Ceylon), Spinach (New Zealand), Swiss Chard.

Brassica (Cole) Leafy Vegetables: Including but not limited to Broccoli, Chinese Broccoli, Broccoli Raab, Brussel Sprouts Cabbage, Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Collards, Kale, Kohirabi, Mustard Greens, Rape Greens.

Legume Vegetables (Succulent or Dried): Including but not limited to Adzuki Beans, Field Beans, French Beans, Kidney Beans, Lima Beans, Moh Beans, Mung Beans, Navy Beans, Pinto Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black Eyed Peas, Catjang, Chinese Longbeans, Cowpeas, Crowder Peas, Southern Peas, Yard-Longbeans, Broad Beans (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jack Beans (Sword Beans), Lablab (Hyacinth Beans), Lentils, Peas (Garden Peas, Field Peas, Sugar Peas), Pigeon Peas, Soybeans.

Foliage of Legume Vegetables: Including but not limited to plant parts of any Legume vegetable included in the legume vegetable group that will be used as animal food including any variety of Beans, Field Peas, Soybeans.

Fruiting Vegetables: Including but not limited to Eggplant, Ground Cherry, Okra, Pepinos, Pepper (Bell Pepper, Chili Pepper, Cooking Pepper, Pimentos, Sweet Peppers), Tomatillo, Tomatoes.

Cucurbit Vegetables: Including but not limited to Balsam Pear (Bitter Melon), Chinese Waxgourd, Citron Melon, Cucumber, Gherkin, Edible Gourds, Melons including Hybrids (Cantaloupe, Casaba, Crenshaw, Honeydew Melons, Honey Balls, Mango Melon, Muskmelon, Persian Melon), Pumpkin, Squash (Summer & Winter), Watermelon including hybrids.

Oriental Vegetables: Including but not limited to Japanese Artichoke, Chinese Broccoli (Gai Lon), Chinese Cabbage (Bok Choy Napa), Chinese Mustard Cabbage (Gai Choy), Cilantro, Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Balsam Pear (Bitter, Melon), Japanese Radish (Diakon), Chinese Spinach, Chinese Waxgourd.

FIELD GRAIN CROPS

FOR THE CONTROL INSECTS SUCH AS: Ants, Aphids, Armyworms, Chinch Bugs, Boll Weevil, Bollworm, Budworm, Caterpillars, Corn Earworms, Fire Ants, Fleabeetles, Fleahoppers, Flies, Hornworms, Loopers, Lygus Bugs, Midges, Mites, Pear Curculia, Pink Boll Worms, Thrips, White Flies.

Cereal Grains: Including but not limited to Barley, Buckwheat, Corn, Millet (Proso & Pearl), Oats, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticale, Wheat, Wild Rice.

Grass Forage, Fodder and Hay: Including but not limited to Bermudagrass, Bluegrass, Bromegrass, Fescue and any type grass, Gramineae Family (Green and Cured), Barley, Buckwheat, Corn, Millet (Proso & Pearl), Oats, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticale, Wheat, Wild Rice, that will be fed to or grazed by livestock, all pasture, range grasses and grasses grown for hay or silage.

Non-Grass Animal Feeds: Including but not limited to Alfalfa, Velvet Bean, Clover, Kudzu, Lespedeza, Lupine, Sainfoin, Trefoil Vetch, Crown Vetch, Milk Vetch.

Ornamentals:

For the control of insects such as: Ants, Aphids, Armyworms, Chinch Bugs, Caterpillars, Fire Ants, Flea Beetles, Fleahoppers, Flies, Fruit Flies, Japanese Beetles, Leafhoppers, Leafminers, Leafrollers, Loopers, Lygus Bugs, Mealy Bugs, Mites, Plant Bugs, Thrips, White Flies.

Ornamentals: Such as African Violets, Aster, Azalea, Begonia, Calceolaria, Calendula, Calia, Camellia, Carnation, Cineraria, Chrysanthemum, Cypress, Daffodil, Dahlia, Dogwood, Elm, Eucalyptus, Fern, Ficus, Geranium, Gladiolus, Gypsophila, Holly, Juniper, Lily,

Marigold, Oak, Palm, Peony, Patunia, Philodendron, Pine, Roses, Snapdragons, Sweetpeas, Tulips, Viburnum, Wandering Jew, Yew, Zinnia.

HERBS, SPICES:

FOR THE CONTROL OF INSECTS SUCH AS: Ants, Aphids, Fire Ants, Loopers, Mites, Plant Bugs, Thrips, White Flies.

Herbs and Spices: Including but not limited to Anise, Balm, Basil, Burnet, Borage, Camomile, Caraway, Catnip, Chives, Clary, Coriander, Costmary, Cumin, Curry Leaf, Dill, Fennel (Italian & Sweet), Fenugreek, Horehound, Hyssop, Marigold, Marjoram (Sweet & Wild), Mint, Nasturtium, Oregano, Pennyroyal, Rosemary, Rue, Sage, Savory (Winter & Summer), Sweet Bay (Bay Leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

FRUIT, NUT, VINE AND SPECIALTY CROPS:

Use rate: Use 2 to 6 pounds per acre depending upon the insect population and amount of foliage.

As a **WETTABLE POWDER**, apply in sufficient water to obtain maximum coverage. Dilution of the material should be at the maximum rate of one half pound in one half gallon of water or at a minimum of one pound to 25 gallons of water (4 tsp. per gallon). It is best to keep the material agitated during application. Apply the material uniformly to both the upper and lower leaf surfaces. Reapply only as necessary. Rates will vary with insect populations and amount of foliage, but adequate control can be obtained at 2 to 6 pounds product per acre.

As a **DUST**, the best results are obtained when the plants are wet or dew still remains on the leaf surface. It is suggested that the plants are misted prior to application to obtain the maximum benefits and use the least amount of product for control.

FOR THE CONTROL OF SUCH INSECTS AS: Ants, Armyworms, Aphids, Blueberry Maggots, Cabbage Loopers, Caterpillars, Cherry Fruit Flies, Cutworms, Fire Ants, Fireworms, Fruit Flies, Fuller Rose Beetle, Gooseberry Fruit Worms, Imported Cabbage worms, Japanese Beetles, Leafhoppers, Mites, Peach Borers, Pecan Weevil, Red-necked Borers, Raspberry Fruit Worms, Rose Chafers, Stink Bugs, Strawberry Leaf Rollers, Thrips, Weevils, Webworms, White Flies.

CITRUS FRUITS: Including but not limited to Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (Sweet & Sour), Pummelo, Sats'ima Mandarin.

POME FRUITS: Including but not limited to Apple, Crabapple, Loquat, Pear, Oriental Pear, Quince.

STONE FRUITS: Including but not limited to Apricot, Cherry (Sweet & Sour), Nectarine, Peaches, Plums, Prunes, Chickasaw Plums, Damson Plum, Japanese Plum.

SMALL FRUITS AND BERRIES: Including but not limited to Blackberry, Blueberry, Boysenberry, Cranberry, Currants, Dewberry, Elderberry, Gooseberry, Grapes,

Huckleberry, Loganberry, Olallie Berry, Raspberry (Black & Red), Strawberry, Youngberry.

SUBTROPICAL FRUITS: Including but not limited to Avocado, Banana, Carob, Barbados Cherry, Cherimoya, Dates, Feijoa, Figs (Adriatic, Calimyma, Kadota, Black Mission, California Brown Turkey, Brunswick), Guava, Kiwifruit, Lychee, Mango, Papaya, Passion Fruit, Perisimmon, Pineapple, Pomegranate.

TREE NUTS: Including but not limited to Almond, Beech Nut, Brazil Nut, Butternut, Cashews, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Japanese Horsechestnut, Macadamia Nut (Bushnut), Pecan, Pistachio, Walnut (Black & English).

SPECIALTY CROPS: Such as Artichoke, Chayote, Asparagus, Coffee, Cotton, Hops, Jojoba, Ornamental, Turf Grass, Sesame, Sunflower (Leaves & Seed), Tea, Tobacco.

On harvested tomatoes and fruit including grapes: Fruit Flies, Vinegar Flies. Apply with a hand or power sprayer or other suitable means on tomatoes and fruits in baskets, on trucks or in plants, raw stock stacked in the yard. Apply liberally at the rate of 8 oz. per 100 sq. ft. of area.

AS A GRAIN PROTECTANT: To protect against grain storage insects, dilute one pound in 5 gallons of water and apply directly to the grain at the rate of 4 to 5 gallons per 1000 bushels as the grain is carried along a belt or as it enters the auger or elevator. Monthly inspection should be made if the top two or three inches are found to be reinfested. Retreat at the rate of 1 to 2 gallons of dilution per 1000 bushels of stored product.

IN MUSHROOM PRODUCTION AND PROCESSING: Mushroom Flies, Fungus Gnats. Apply dilution at a rate of 5 oz. per 1000 cu. ft. of space. Direct spray 8 to 10 feet from beds and slightly upwards over beds. Keep circulating fans running but turn off refrigerators during application. In processing plants, use at the rate of 5 oz. per 2000 cu. ft. of space.

MOSQUITO CONTROL: May be used on croplands as an adulticide to protect agricultural workers during harvesting. Use 2 to 12 ounces per acre when applied by fixing wing aircraft, helicopters or truck mounted equipment. Do not apply in a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

COMMERCIAL INDOOR AND OUTDOOR USE: For use in and around aircraft (cargo areas only), apartment buildings, bakeries, bottling facilities, breweries, buses, cafeterias, candy plants, canneries, dairy product processing plants, food manufacturing plants, food processing plants, food service establishments, granaries & grain mills, hospitals, hotels, homes, industrial buildings, installations, kitchens, laboratories, meat, poultry & egg processing & packaging plants, mobile and motor homes, motels, nursing homes, offices, railcars, restaurants, schools, ships & vessels, supermarkets, theaters, trailers, trucks, warehouses and wineries. This product is intended for application with hand or power duster, sprayer or other

suitable means, to hiding and running areas, and places where pests are found. Remove or cover food prior to use. Cover all food preparation surfaces and utensils during treatment, or thoroughly wash before use.

CONTROLS: Cockroaches, Ants, Fire Ants, Earwigs, Silverfish, Crickets, Millipedes, Box Elder Bugs, Scorpions and Other Similar Crawling Insects: Coat with a layer of dust, or spray as a wettable powder (mix 6 Tbsp. per gallon of water) in areas where these pests are found or may hide such as cracks and crevices, behind and beneath stoves, refrigerators, sinks, cabinets, garbage cans, garbage trucks, in allies, basements, driveways, sidewalks, patios around houses and buildings. Apply directly to insects when possible.

GRASSES, LAWNS & TURF (INCLUDING GOLF COURSES): Controls Armyworms, Caterpillars, Chinch Bugs, Cutworms, Fleas, Ants, Fire Ants, Ticks.

Use Rate: Use 16 oz. (1lb.) per 1000 sq. ft. of area. Apply either as a DUST or as a WETTABLE POWDER. Use hose-end sprayer for lawns, add a small amount of water to the jar and add the amount needed for the measured area, stir to make a slurry, then add the remaining water to the top of the jar. Empty entire contents of the jar on the premeasured area.

FIRE ANTS:

Mound Drench or Water Method: Use 0.6 oz. (4 Tbsp.) to one gallon of water. For best results, saturate the perimeter of the mound first working toward the center of the mound in a circular motion (you may want the mixture to puddle). An ant mound 12-14 inches in diameter requires approximately one gallon of the mix.

Hand Sprayer: Use 0.6 oz. (4 Tbsp.) to one gallon of water. It is best to keep the material agitated during application. Apply the contents to a premeasured area.

Hose end Sprayer: Use 16 oz. (1 lb.) per 1000 sq. ft. of area. Use a hose-end sprayer for lawns, add a small amount of water to the jar and add the amount needed for the measured area, stir to make a slurry; then add the remaining water to the top of the jar. Empty the entire contents of the jar on the premeasured area.

As a Dust: For best results, dust the perimeter of the mound first. With a stick, disturb the mound, then dust mound in a circular motion working toward the center until the entire mound is thoroughly dusted.

HOUSE FLIES: Apply to surface of window frames around light fixtures, on porches, in garages and other similar places where these insects light or congregate.

BEDBUGS: Treat liberally baseboards, moldings, cracks, bedstead, and both sides of mattress.

FLEAS: Thoroughly treat floor, bedding in and around pet's sleeping quarters. Treat surrounding cracks and crevices, baseboards, furniture, carpeting and wherever fleas are suspected. Dust rugs and carpets with thin coat of dust. Rake (not sweep) nap back and forth, causing dust to enter rug or carpet. Leave overnight. When cleaning do not

make special effort to remove dust in rug or carpet. It will kill live fleas, repel others and help deodorize carpet or rug.

ROACH CONTROL IN PUBLIC SEWERS: A power blower using 1 pound of product will coat the dry part of the sewer line and manhole for a maximum of 300 ft. each direction. It will also enter the laterals and dust may be seen emerging from the vent pipes on the homes on each side of the street.

FOOD/FEED HANDLING ESTABLISHMENTS: For control of all labeled pests, applications are permitted in food/feed and non-food/non-feed areas of food/feed handling and processing establishments as a spot or crack and crevice treatment.

Food/Feed Handling Establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas when food is exposed and facility is in operation are also considered food areas.

Non-Food/Non-Feed Areas: Examples of non-food/non-feed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after packaging, canning or bottling).

Crack and Crevice Treatments: Apply in small amounts directly into cracks and crevices using equipment capable of directing insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where cockroaches, ants, silverfish and crickets hide. Spot treatment may also be made to areas where any of the above pests hide or inhabit such as baseboards, storage areas, closets, around water pipes, doors, windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment. Care should be taken to avoid depositing the product onto exposed food contact surfaces or introducing the material into the air.

Spot Treatment: No individual spot will exceed two (2) square feet. Limit spot treatments to wall or floor surfaces where pests have been seen or where pests are suspected of hiding.

In the home, do not allow spray dust to contact food/feed surfaces. If it does, clean surfaces with soap and water before use.

DOGS, CATS AND PETS:

CONTROLS: Fleas, Flies, Lice and Ticks.

As a dust, lightly work the material into the hair of the animal, down to the skin. Pay strict attention to places where the animal cannot reach, such as the scalp. Dust the ears of the animal with care. Dust the animal's bedding and areas where the animal sleeps, as well as cracks and crevices adjacent to them. Repeat application weekly if

necessary. Individual ticks observed on the animal should be dusted thoroughly.

As a spray, mix 4 Tbsp. per gallon of water. Spray animal thoroughly. Repeat application weekly if necessary.

For wet application, mix 4 Tbsp. per gallon of water. Dip animals in solution paying attention to make sure animals are completely covered with material. Repeat application weekly if necessary.

ANIMAL QUARTERS (including and not limited to Barns, Dairies, Milkrooms, Pet Kennels, Poultry Houses, Runs and Stables).

CONTROLS: Ants, Gnats, Fire Ants, Fleas, Flies, Lice, Mites, Mosquitoes, Ticks.

USE RATE: Use at the rate of 2 oz. Per 400 sq. ft. of surface area. Apply with a hand or power duster or sprayer or other suitable means to the hiding places where pests are found.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container in a cool dry place, out of sunlight. Prevent cross-contamination with other pesticides and fertilizers.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) the offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities. Containers 1 Pound and Smaller: Do not reuse container. Flush container with water and use the water in application. Place empty flushed container in trash.

Organic Resources, Inc.
8745 Grissom Rd. #197
San Antonio, Texas 78251
800-862-7482

www.organicresources.com
EPA Reg. No. 70126-1

