

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460 EPA Reg. Number:

Date of Issuance:

70933-1

FEB 10 2000

NOTICE OF PESTICIDE:

x Registration Reregistration

Unconditional

Term of Issuance:

Name of Pesticide Product:

MLS Fly contROL Tub

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Midcontinent Livestock Supplements, Inc.

Route 2. Box 74B

Moberly, MO 65270

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5). Once a pesticide is registered, however, it is not regarded as permanently acceptable. Registration does not eliminate the need for continual reassessment of pesticides. If the Agency determines that, at any time, additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under FIFRA section (3)(c)(2)(B).

- Make the following label changes:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 70933-1".
 - b. Based upon the toxicity profile for this product, you must have a "Statement of Practical Treatment" of "First Aid" statement. It should be located under the Precautionary Statements section and state the following:

Signature of Approving Official:

Date

George T. LaRocca, Product Manager 13

EPA Form 8570-6

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

Also, add the following statement near the first aid statements: "Have the product container or label with you when calling a poison control center or doctor or going for treatment."

Add a NOTE TO PHYSICIAN section. Should be located directly below your "Statement of Practical Treatment: section and should contain the following information: 1) Technical information on symptomatology; 2) Use of supportive measures to maintain life functions; 3Mediciane that will counteract the specific physiological effects of the pesticide; 4) Company telephone number to specific medical personnel who can provide specialized medical advice.

EPA recommends that these statements be boxed and bulleted to present the information in a clear, easy-to-read fashion.

c. Change PRECAUTIONARY STATEMENT to read: "Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and handlers must wear:

- 1) Long-sleeved shirt and long pants,
- 2) Chemical-resistant gloves,
- 3) Shoes and socks

)

)

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- d. We are unsure what age of cattle you refer to with "mature and growing beef cattle". Provide more specific information in order to prevent the product from inappropriate use.
- e. Remove the term, "all natural" from the front panel. The Agency considers such terms, a false or misleading statement. Refer to 40 CFR 156.10(a)(5) for further information.
- f. Under DIRECTIONS FOR USE, paragraph 2, replace the # sign with "lb.".
- g. Per the tetrachlorvinphos reregistration eligibility document (RED), September 1995, add the following precautionary statement, "This product may cause skin sensitization reactions in certain individuals."
- Submit two (2) copies of your final printed label before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions.

If you have any questions regarding this action, please contact Linda Werrell of my team at (703) 308-8033. A stamped copy of the label is enclosed for your records.

Sincerely

George T. LaRocca

Product Manager (13)

Insecticide Branch

Registration Division (7505C)

Enclosure

}

1-800-406-4609

Route 2, Box 748, Moberly, MO 65270

This all natural protein, vitamin, mineral package with RABON® Oral Larvicide is formulated for mature and growing beef cattle.

INDICATIONS FOR USE: Prevents development of horn flies, house flies, face flies and slable flies in the manure of treated cattle. Provide plenty of drinking water and free choice salt.

ACTIVE INGREDIENT: BY WEIGHT

Total 100.00%

* Each pound contains 2.1 grams RABON
**Refers only to ingredients which are not Inrvicidal.

GUARANTEED ANALYSIS

Crude Protein, min.————— 4.0%
Crude Fat, min 5.0%
Crude Fiber, max.——— 1.0%
Calcium (Ca), max.———— 5.0%
Calcium (Ca), min. 4.0%
Phosphorus (P), min.——— 6.0%
Salt (NaCl), max 3.0%
Salt (NaCl), min.————— 2.5%
Magnesium (Mg), min.———— 4.0%
Potassium (K), min. 4.0%
lodine (I), min 50 PPM
Selenium (Se), min. 9 PPM
Vitamin A, min. IU/lb.——— 200,000
Vitamin D ₃ , min. IU/lb. 27,000
Vitamin E, min. IU/lb. 100

INGREDIENTS

Molasses Products, Monocalcium Phosphate, Animat fat, Salt, Dehydrated Aspergillus Oryzae Fermentation Extract, Magnesium Oxide, Vitamin A Supplement, D-activated Animal Sterol (source of vitamin D.), Vitamin E Supplement, Sodium Selenite, Calcium Pantothenate, Thiamine Hydrochloride, FyrioJoxine Hydrochloride, Zinc. Oxide, Copper Sulfate, Manganous Oxide, Riboflavin Supplement, Vitamin B₁₂ Supplement, Niacin, Choline Chloride, Biotin, Folic Acid, Ascorbic Acid, Potassium Iodide, Ethylenediamine Dihydriodide, and Sodium Molybdate.

CAUTION: Keep out of reach of children.

PRECAUTIONARY STATEMENTS CAUTION: HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with it's labeling.

Provide one 200# tub per 35 head of cattle. Cattle should consume an average of 1/3 lb. of the tub per 1000 lbs. of body weight per day. This will supply the recommended average intake of 70 mg. RABON Oral Larvicide per 100 lbs. of body weight. Animals should not be starved for minerals before initiation of the MLS contROL Tub program. Place tub near loafing areas.

If consumption of MLS contROL Tub is above 1/3 lb. per 1000 lbs. of body weight per day, reduce the number of feeding locations. If consumption is below 1/3 lb. per 1000 lbs. of body weight per day, increase the number of feeding locations.

Start feeding the MLS contROL Tub early in the Spring before flies begin to appear, and continue feeding throughout the Summer and Fall.

EPA Reg. #70933<u>-1</u> EPA Est. #<u>70933-MO-01</u> FEB 1 0 2000

NET WEIGHT:

125 / 200 Lbs 709 33-/ (56.7kg / 90.7kg)

#8 MLS Fly ContROL Tub

RABON® ORAL LARVACIDE

Midcontinent Livestock Supplements, Inc.



Route 2, Box 748, Moberly, MO 65270

This all natural protein, vitamin, mineral package with RABON® Oral Larvicide is formulated for mature and growing beel cuttle.

INDICATIONS FOR USE: Prevents development of horn flies, house flies, frace flies and stable flies in the manure of treated callle. Provide plenty of drinking water and free choice salt.

ACTIVE INGREDIENT: BY WEIGHT

Total 100.00%

* Each pound contains 2.1 grams RABON ** Refers only to ingredients which are not larvioidal

GUARANTEED ANALYSIS

Crude Protein, min.————	
Crude Fat, min.	- 5.0%
Crude Fiber, max.—————	1.0%
Calcium (Ca), max.	- 5.0%
Calcium (Ca), mln.	4.0%
Phosphorus (P), min.————	- 6.0%
Salt (NaCl), max	- 3.0%
Salt (NaCl), min.	- 2.5%
Magnesium (Mg), min	- 4.0%
Potassium (K), min.————	- 4.0%
lodine (I), min.————	50 PPM
Selenium (Se), min.	9 PPM
Vitamin A, min. IU/lb	200,000
Vitamin D ₃ , min. IU/lb.————	27,000
Vitamin E, min. IU/lb.	
•	

INGREDIENTS

Molasses Products, Monocalcium Phosphate, Animal fat, Salt, Dehydrated Aspergillus Oryzae Fermentalion Extract, Magnesium Oxide, Vitamin A Supplement, D-activated Animal Sterol (source of vitamin D₃), Vitamin E Supplement, Sodium Selenite, Calcium Pantothenate, Thiamine Hydrochloride, Pyriodoxine Hydrochloride, Zinc Oxide, Copper Sulfate, Manganous Oxide, Riboflavin Supplement, Vitamin B₁₂ Supplement, Niacin, Choline Chloride, Biotin, Folic Acid, Ascorbic Acid, Potassium Iodide, Ethylenediamine Dihydriodide, and Sodium Molybdate.

CAUTION: Keep out of reach of children.

PRECAUTIONARY STATEMENTS CAUTION: HAZARDS TO HUMANS

Harmful if swallowed. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water. If irritation persists, get medical attention.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment washwater.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with it's labeling.

Provide one 200# tub per 35 head of cattle. Cattle should consume an average of 1/3 lb. of the tub per 1000 lbs. of body weight per day. This will supply the recommended average intake of 70 mg. RABON Oral Larvicide per 100 lbs. of body weight. Animals should not be starved for minerals before initiation of the MLS contROL Tub program. Place tub near loafing areas.

If consumption of MLS contROL Tub is above 1/3 lb. per 1000 lbs. of body weight per day, reduce the number of feeding locations. If consumption is below 1/3 lb. per 1000 lbs. of body weight per day, increase the number of feeding locations.

Start feeding the MLS contROL Tub early in the Spring before flies begin to appear, and continue feeding throughout the Summer and Fall.

EPA Reg. #70933<u>-1</u> EPA Est. #<u>70933-MO-01</u>

NET WEIGHT:

125 / 200 Lbs. (56.7kg / 90.7kg)

#8 MLS Fly ContROL Tub

RABON® ORAL LARVACIDE

