

71368-23

8/25/2009

Jacket
1/36



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

71368-23

Date of Issuance:

AUG 25 2009

NOTICE OF PESTICIDE:

Registration
 Reregistration
(under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:

KAMBA 4SL

Name and Address of Registrant (include ZIP Code):

Nufarm Inc.
150 Harvester Drive, Suite 200
Burr Ridge, IL 60527

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.

Signature of Approving Official:

Joanne Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

AUG 25 2009

2) The signal word currently on the label “WARNING” must be revised to read “CAUTION” and the Spanish signal word “AVISO” must be deleted from the label.

3) Based on toxicity ranking per the acute toxicity review, the First Aid statements should be placed on the label in the following order:

“IF SWALLOWED:…
IF INHALED:…
IF ON SKIN OR CLOTHING:…
IF IN EYES:…”

4) The first sentence in the handler PPE section must be revised to read “Some materials that are chemical-resistant to this product are made of any waterproof material.”

Note: Eyewear is not required for this product but may remain in this section of the label.

5) The following text must be added to the Engineering Control text on the label:

“Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].”

6) Delete “For terrestrial use” from the Environmental Hazards section of the label.

7) The double notification text appearing in the Agricultural Use Requirements box (“Notify workers of the application by warning them orally and by posting warning signs at entrances to the treated area”) is no longer needed and should be deleted from the label.

8) To the Storage and Disposal section change “Do not contaminate water, foodstuffs, feed, or seed by storage or disposal” to “Do not contaminate water, food, or feed by storage or disposal”.

9) Per the revised RED label table, the following revisions are needed to the directions for use:

Sorghum:

Replace the text “Delay harvest until 30 days after treatment” with:

- “30-day PHI for sorghum grain.
- 0-day PHI for sorghum forage.
- 30-day PHI for sorghum fodder.”

Small Grains (Wheat, barley and oats):

Add:

- “ 7-day PHI for barley.
- 7-day PHI for oats (grain).

Page 3
EPA Reg. 71368-23

7-day PHI for wheat (grain).”

Sugarcane:

-The rates of up to 2 lbs ai per acre to treat perennial weeds in the table on Page 20 of the label exceed the maximum allowable application rate of 1.0 lb ai per acre per the RED. The label must be revised.

-Per the revised RED label table, a 87-day PHI for sugarcane (cane) must be added to the label.

Turf and Lawns:

The directions on Page 26 of the label to “apply 1/5 to 1/2 pint of this product per treated acre with 1/2 to 1 1/2 lbs acid equivalent of 2,4-D, MCPA, or MCPP-p” is not acceptable. Per the MCPP-p RED, the maximum individual application rate is 0.75 lbs ae MCPP-p per acre per application for broadcast application and 1.2 lbs ae MCPP-p ae per acre for spot treatment only. The label must be revised.

Preplant Directions (Post Harvest/Fallow/Crop Stubble/Set-A-Side):

-The text on Page 29 of the label “...do not exceed a total of 4 pints of this product per treated acre during any given fallow period” must be revised to read “Do not exceed 2.0 lbs ae Dicamba per acre per year.”

-The statements “apply 1 to 4 pints (2 lbs ae) of Kamba 4SL with,....” appearing above the table on Page 30 and the text “The following recommendations are based on this product use rates up to 4 pints (2 lbs ae) per treated acre” on Page 31 of the label are unacceptable because rates up to 4 pints exceed the maximum allowable individual application rate of 1.0 lbs ae Dicamba per acre. The label must be revised.

Wiper Application Uses:

An area to be treated must be added to the use directions for wiper application and the rate must comply with the rates specified in the RED.

10) Page 3, change “General Information” to “Product Information”. Page 7 and 22, delete “General” from “General Weed List” and Page 20 “General Farmstead”. Page 26, 31 and anywhere else it occurs in the label, change “recommended rates” to “specified rates”.

A stamped copy of the label is enclosed for your records. You must submit one copy of the final printed label before you release the product for shipment. Products shipped after 12 months from the date of this letter or the next printing of the label whichever occurs first, must bear the new revised label. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA. Your release for shipment of the product constitutes acceptance of these conditions. This label supercedes all other previously accepted labels. If you have any questions please call Erik Kraft at 703-308-9358 or email at Kraft.Erik@epa.gov.

4/3b

KAMBA 4SL

Herbicide

For weed control in corn, sorghum, small grains, pasture, hay rangeland, general farmstead (non-cropland), rights-of-way, public utility and industrial areas, fallow, sugarcane, asparagus, turf and grass seed crops.

ACTIVE INGREDIENT:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)*	49.77%
OTHER INGREDIENTS:	<u>50.23%</u>
TOTAL:	100.00%

*This product contains 41.35% 3,6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon (480 g/L).

KEEP OUT OF REACH OF CHILDREN

WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)


SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300
For Medical Emergencies Only, Call (877) 325-1840

ACCEPTED
with COMMENTS
In EPA Letter Dated:

AUG 25 2009
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

71368-23



EPA REG. NO. 71368-23
EPA EST. NO. 228-IL-1

NET CONTENTS: 2.5 GALLONS (9.46L)

071368-00023.20090629.Dicamba RED Pending

Manufactured For
NUFARM INC.
150 Harvester Drive
Burr Ridge, IL 60527



**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING / AVISO**

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid breathing spray mist. Wash thoroughly after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on the EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (except for applicator using groundboom equipment, pilots and flaggers) and
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statement:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [140 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Apply this product only as directed on label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls worn over short-sleeve shirt and short pants, chemical-resistant footwear plus socks, chemical-resistant gloves made of any waterproof material, chemical-resistant headgear for overhead exposure and protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

Before applying this product, read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury, or illegal residues.

GENERAL INFORMATION

The following directions apply to all uses of this product. Additional precautions and restrictions will be found in each specific use section.

Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

Do not apply this product through any type of irrigation system.

Do not apply more than 2 pints of this product per application, and no more than 2 applications per year.

MIXING AND APPLICATION

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF THIS PRODUCT. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

This product is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see Compatibility Test) should be made prior to tank mixing.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of a diluted spray per treated acre when using ground application equipment, or 1 to 10 gallons of a diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, this product should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

BEST STEWARDSHIP PRACTICES

This product provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

GROUND AND SURFACE WATERS PROTECTION

1. Point source contamination - To prevent point source contamination, do not mix, load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment, or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or antisiphoning devices must be used on all mixing equipment.

2. Movement by surface runoff or through soil - Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the general information section of this label.
3. Movement by water erosion of treated soil - Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

SENSITIVE CROP PRECAUTIONS

This product may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems, or foliage. These plants are most sensitive to this product during their development or growing stage. FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING THIS PRODUCT.

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of this product with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles such as D10, TK10 or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 gpa, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- Do not apply this product adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.
- To avoid injury to desirable plants, equipment used to apply this product should be thoroughly cleaned (see Procedure For Cleaning Spray Equipment) before reusing to apply any other chemicals.

All crop uses of this product are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. **Tank mix recommendations are for use only in states where the tank mix product and application site are registered.**

BAND TREATMENTS

This product may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE treated acre} = \text{Band RATE per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per treated acre} = \text{Band VOLUME per treated acre}$$

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

**Amount of Herbicide to Add to One Pint of Spray Carrier
(Assuming Volume is 25 Gallons per Acre)**

HERBICIDE FORMULATIONS	RATE PER ACRE	LEVEL TEASPOONS
Dry	1 lb.	1-1/2
Liquid	1 pt.	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Re-run the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of this product or tank mixes of this product or tank mixes of this product plus 2,4-D amine.

1. Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply this product as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. This product tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

5. Complete Step 1.
6. Fill tank with water while adding 2 pounds of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
7. Flush the detergent solution out of the spray tank through the boom.
8. Repeat step 1, and follow with steps 2, 3, and 4.

GENERAL WEED LIST

This is a general list of weeds which may be treated with this product in accordance with this label as recommended under the rates and timing sections of the Individual Use Headings. Proper usage of this product will give control or growth suppression of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species including:

ANNUALS

Amaranth, Spiny (Spiny Pigweed)	Ladysthumb	Radish, Wild
Aster, Slender	Lambsquarters, Common	Ragweed, Common
Bedstraw	Lambsquarters (triazine resistant)	Ragweed, Giant (Buffaloweed)
Beggarweed, Florida	Lettuce, Prickly	Ragweed Lance-Leaf
Broomweed, Common	Mallow, Common	Rubberweed, Bitter (Bitterweed)
Buckwheat, Wild	Mallow, Venice	Sesbania, Hemp
Buffalobur	Mayweed	Shepherdspurse
Burclover, California	Morningglory, Ivyleaf	Sicklepod
Burcucumber	Morningglory, Tall	Sida, Prickly (Teaweed)
Buttercup, Roughseed	Mustard, Tansy	Smartweed, Green
Carpetweed	Mustard, Wild	Smartweed, Pennsylvania
Catchfly, Nightflowering	Mustard (Yellowtops)	Sneezeweed, Bitter
Chamomile, Corn	Nightshade, Black	Sowthistle, Annual
Chickweed, Common	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)	Sowthistle, Spiny
Clovers (Annual)	Pepperweed, Virginia (Peppergrass)	Spikeweed, Common
Cockle, Corn	Pigweed, Prostrate	Spurge, Prostrate
Cockle, Cow	Pigweed, Redroot (Carelessweed)	Spurry, Corn
Cocklebur, Common	Pigweed, Rough	Starbur, Bristly
Croton, Tropic	Pigweed, Smooth	Sumpweed, Rough
Croton, Woolly	Pigweed (triazine resistant)	Sunflower, Common (Wild)
Daisy, English	Pigweed, Tumble	Sunflower, volunteer
Evening primrose, Cutleaf	Poorjoe	Thistle, Russian
Fleabane, Annual	Puncturevine	Velvetleaf
Goosefoot, Nettleleaf	Purslane, Common	Waterhemp
Henbit	Pusley, Florida	Waterprimrose, Winged
Jimsonweed		Wormwood, Annual
Knotweed		
Kochia		

BIENNIALS

- | | |
|----------------------------------|---------------------|
| Burdock, Common | Plantain, Bracted |
| Carrot, Wild (Queen Anne's Lace) | Ragwort, Tansy |
| Cockle, White | Starthistle, Yellow |
| Evening primrose, Common | Sweetclover |
| Geranium, Carolina | Teasel |
| Gromwell | Thistle, Bull |
| Knapweed, Diffuse | Thistle, Milk |
| Knapweed, Spotted | Thistle, Musk |
| Mallow, Dwarf | Thistle, Plumeless |

PERENNIALS

- | | | |
|---------------------------------|-------------------------------|-----------------------------|
| *Alfalfa | Fern, Bracken | Redvine |
| Artichoke, Jerusalem | Garlic, Wild | Sericia Lespedeza |
| Aster, Spiny | Goldenrod, Canada | Smartweed, Swamp |
| Waster, Whiteheath | Goldenrod, Missouri | Snakeweed, Broom |
| Beadstraw, Smooth | Goldenweed, Common | *Sorrel, Red (Sheep Sorrel) |
| Bindweed, Field | Hawkweed | Sowthistle |
| Bindweed, Hedge | Horsenettle, Carolina | Sowthistle, Perennial |
| Blueweed, Texas | Ironweed | Spurge, Leafy |
| *Bursage, | Knapweed, Black | Sundrop, Halfshrub |
| (Bur | Knapweed, Russian | (Eveningprimrose) |
| Ragweed)(Lakeweed)(Povertyweed) | Mare's Tail (Horseweed) | Thistle, Canada |
| Bursage, Woollyleaf (Lakeweed) | Milkweed, Climbing | Toadflex, Dalmatian |
| Buttercup, Tall | Milkweed, Common | Tropical Soda Apple |
| Campion, Bladder | Milkweed, Honeyvine | Trumpetcreeper (Buckvine) |
| Chickweed, Field | Milkweed, Western Whorled | Vetch |
| Chickweed (Mouseear, Canada) | Nettle, Stinging | Waterhemlock |
| Chicory | Nightshade, Silverleaf (White | Waterprimrose, Creeping |
| *Clover, Hop | Horsenettle) | *Woodsorrel, Creeping |
| *Dandelion, Common | Onion, Wild | Common Yellow |
| *Dock, Broadleaf (Bitterdock) | *Plantain, Broadleaf | Wormwood, Common |
| *Dock, Curly | Plantain, Buckhorn | Wormwood, Louisiana |
| Dogbane, Hemp | Pokeweed | *Yankeeweed |
| *Dogfennel (Cypressweed) | Ragweed, Western | Yarrow, Common |

* Noted perennials may be controlled using this product at rates lower than those recommended for other listed perennial weeds. (See Application Rates and Timings sections).

WOODY

Alder	* Dewberry	Maple	Sagebrush, Fringe
Ash	* Dogwood Elm	Mesquite	Sassafras
Aspen	Grape	Oak	Serviceberry
Basswood	*Hawthorn (Thornapple)	Oak, Poison	Spicebush
Beech	Hemlock	Olive, Russian	Spruce
Birch	Hickory	Persimmon, Eastern	Sumac
*Blackberry	Honeylocust	Pine	*Sweetgum
*Blackgum	Honeysuckle	*Plum, Sand (Wild	Sycamore
*Cedar	Hornbeam	Plum)	Tarbrush
Cherry	Huckleberry	Poplar	Willow
Chinquapin	Huisache	Rabbitbrush	Witchhazel
Cottonwood	Ivy, Poison	*Redcedar, Eastern	*Yaupon
*Creosotebush	Kudzu	*Rose, McCartney	*Yucca
Cucumbertree	Locust, Black	*Rose, Multiflora	

* Growth Suppression

FIELD, SEED, POPCORN AND SILAGE CORN

Observe all PRECAUTIONS, MIXING and APPLICATION instructions as well as the following:

*Do not apply this product to seed corn or popcorn without first verifying with your local seed corn company (supplier) this product selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

This product is not registered for use on sweet corn.

Direct contact of this product with corn seed must be avoided. If corn seeds are less than 1-1/2 inches below the surface, delay application until corn has emerged.

Up to 2 applications of this product may be made during a growing season. Do not exceed a total 1-1/2 pints of this product per treated acre per crop year. Allow two weeks or more between applications of this product. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.

Applications of this product to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Agriculturally approved surfactants or sprayable fertilizers (1/2 to 1 gallon per acre of 28%, 30%, 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate*) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum-based oils after crop emergence or crop injury may result.

Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

Several synthetic pyrethroid insecticides are labeled for tankmix applications with this product. Refer to their label for specific recommendations.

* Not for use in California

WEEDS CONTROLLED

This product will control many ANNUAL broadleaf weeds or give growth suppression of many PERENNIAL broadleaf weeds commonly found in corn. (Refer to the General Weeds List).

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

PREPLANT/PREEMERGENCE IN NO-TILLAGE CORN

Applications of this product may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply this product at 1 pint per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 1/2 pint per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply this product after 4 to 6 inches of regrowth has occurred.

PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN

This product may be applied after planting and prior to corn emergence. Application at 1 pint per treated acre may be made to medium or fine textured soils, which contain 2% or greater organic matter. DO NOT apply to coarse textured soils (sand, sandy loam and loamy sand) until after crop emergence (see Early Postemergence uses below).

Preemergence application of this product does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

**EARLY POSTEMERGENCE (ALL TILLAGE SYSTEMS)
(Spike Through 8 Inch Tall Corn)**

This product at 1 pint per treated acre may be applied during the period from corn emergence through the five leaf stage or 8 inches tall, whichever comes first. Reduce the rate to 1/2 pint per treated acre if corn is growing on coarse textured soils (sand, sandy loam, loamy sand). See Late Postemergence applications given below if the 6th true leaf is emerging from whorl or corn is greater than 8 inches tall.

**LATE POSTEMERGENCE (ALL TILLAGE SYSTEMS)
(8 to 36 Inch Tall Corn)**

Application of this product at 1/2 pint per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when: (1) corn leaves prevent proper spray coverage; (2) sensitive crops are growing nearby; (3) tank mixing with 2,4-D.

DO NOT apply this product when soybeans are growing nearby if any of these conditions exist:

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

OVERLAY (SEQUENTIAL) TREATMENTS

This product may be applied to ground previously treated with one or more of the following herbicides:

acetochlor (Surpass, Harness)	Guardman
alachlor (Lasso)	glyphosate (Roundup)
atrazine	halosulfuron (Permit)
Bicep II Magnum	Lariat
Bullet	Marksman
butylate (Sutan +)	metolachlor (Dual II Magnum)
Clarity	paraquat (Gramoxone)
dimethenamid (Frontier)	pendimethalin (Prowl)

EPTC (Eradicane)

simazine (Princep)
Surpass

Apply this product at 1/2 pint per treated acre to ground previously treated with full rates of Clarity or Marksman herbicides. Allow at least 2 weeks between applications.

READ AND FOLLOW LABEL DIRECTIONS FOR EACH OF THE ABOVE PRODUCTS.

TANK MIX TREATMENTS FOR CORN

This product may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, and other restrictions.

RATES AND TIMINGS

This Product	Preplant/ Preemergence (No Tillage Corn)	Preemergence (Conventional or Reduced Tillage Corn)	Early Post- Emergence (All Tillage Systems)	Late Postemergence (All Tillage Systems)	Additional Directions
ACCENT Nicosulfuron	--	--	1/2 - 1 ounce ai/A	1/2 - 1 ounce ai/A (To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.)	Application may be made to emerged weeds before corn is greater than 24 inches tall. Use non-ionic surfactant at .25% (V/V) with this tank mixture.
Atrazine	1-1/4 to 2 lbs. ai/A	1-1/4 to 2 lbs. ai/A	1-1/4 to 2 lbs. ai/A. Crop oil concentrates may be used with this mixture if corn is 5 inches or less in height.	1-1/4 to 2 lbs. ai/A. Do not apply if corn is greater than 12 inches.	Application may be made before grasses are 1-1/2 inches tall. Follow all state and Federal restrictions pertaining to atrazine applications.
BEACON Primisulfuron	--	--	0.31 to 0.62 ounce ai/A	0.31 to 0.62 ounce ai/A (To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.)	Application may be made to emerged weeds when corn is 4 to 24 inches tall. Use non-ionic surfactant at .25% (V/V) with this tank mixture.
DUAL II Magnum Metolachlor	0.8 to 1.6 lbs. ai/A	0.8 to 1.6 lbs. ai/A (Use only on fine or medium soils with 2-1/2% or greater organic matter.)	0.8 to 1.6 lbs. ai/A	--	Application may be made before grasses reach the 2-leaf state and before corn is greater than 3 inches tall.

RATES AND TIMINGS (Continued)

This Product	Preplant/ Preemergence (No Tillage Corn)	Preemergence (Conventional or Reduced Tillage Corn)	Early Post- Emergence (All Tillage Systems)	Late Postemergence (All Tillage Systems)	Additional Directions
FRONTIER Dimethenamid	13 to 25 fl. oz/A	13 to 25 fl. oz/A (Use only on fine or medium textured soils with 2.5% or greater organic matter.)	13 to 25 fl. oz/A	--	Application may be made up to 8- inch tall corn. This treatment must be combined with a herbicide that provides postemergence control of grass weeds if they are greater than 1 inch tall at the time of application.
GRAMOXONE Paraquat	1/4 to 1 lb. ai/A	1/4 to 1 lb. ai/A	--	--	Application may be made to emerged weeds but prior to corn emergence.
HARNESS or SURPASS acetochlor	1-1/2 to 3 lbs. ai/A	1-1/2 to 3 lbs. ai/A (Use only on fine or medium textured soils with 2.5% or greater organic matter.)	--	--	Application should be made prior to corn emergence.
LASSO alachlor	1-1/2 to 4 lbs. ai/A	1-1/2 to 4 lbs. ai/A (Use only on fine textured soils with greater than 2- 1/2% organic matter.)	1-1/2 to 4 lbs. ai/A	--	Application may be made before grasses reach the 2-leaf stage and before corn is greater than 3 inches tall. If microencapsulated forms of alachlor are used (Lasso MT, Partner), applications must be made prior to grass emergence.
PRINCEP simazine	2.0 to 3.0 lbs. ai/A	2.0 to 3.0 lbs. ai/A	--	--	Application may be made prior to corn or weed emergence.
PROWL pendimethalin	--	3/4 to 1-1/2 lbs. ai/A (Use only on fine or medium textured soils with 2-1/2% or greater organic matter.)	3/4 to 1-1/2 lbs. ai/A	--	Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2-leaf stage of growth.

RATES AND TIMINGS (Continued)

This Product	Preplant/ Preemergence (No Tillage Corn)	Preemergence (Conventional or Reduced Tillage Corn)	Early Post- Emergence (All Tillage Systems)	Late Postemergenc e (All Tillage Systems)	Additional Directions
ROUNDUP glyphosate	1.0 to 3.0 lbs. ai/A	1.0 to 3.0 lbs. ai/A	--	--	Application may be made to emerged weeds but prior to corn emergence.
2,4-D	1/4 to 1/2 lbs. ai/A	1/4 to 1/2 lbs. ai/A	Not recommended	1/8 lbs. ai/A	Drop pipes are to be used when corn height is 8 inches or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of the weed foliage.

SORGHUM (MILO)

Observe all precautions, including the reference to crops growing under stress.

Read and follow mixing and application instructions.

Applications of this product to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture or hay, refer to the Pasture Use section of this label. Do not apply this product to sorghum grown for seed production.

Make no more than one application per growing season.

WEEDS CONTROLLED

This product, when applied at the recommended rate for sorghum, will control many actively growing ANNUAL broadleaf weeds and will reduce competition from established PERENNIAL broadleaf weeds as well as control their seedlings. (Refer to GENERAL WEED LIST.)

RATES AND TIMINGS

This product may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of this product must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3 to 5 leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

BROADCAST RATE PER TREATED ACRE: 1/2 pint (1/4 lb. a.i.)

**TANK MIX TREATMENT
KAMBA 4LS PLUS ATRAZINE**

For improved control of emerged, actively growing broadleaf weeds including triazine-resistant species and added suppression of perennial broadleaf weeds, tank mix 1/2 pint this product with 0.5 to 1.25 lbs. a.i. atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix 1/2 pint this product with 2 lbs. a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3 to 8 inches tall and when broadleaf weeds are small (less than 6 inches tall). Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will

depend upon soil texture and length of residual weed control desired. Follow all State and Federal restrictions pertaining to atrazine applications.

KAMBA 4LS PLUS BUCTRIL

For improved control of broadleaf weeds, tank mix 1/2 pint of this product with 1 to 1-1/2 pints (0.25 to 0.375 lbs. ai/A) Buctril Herbicide per treated acre. Make application at 4-leaf to 15-inch-tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall.

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS AND OTHER RESTRICTIONS.

OVERLAY (SEQUENTIAL) TREATMENTS

This product may be applied to ground previously treated with one or more of the following herbicides:

Herbicide	Maximum Rate Per Treated Acre (lbs. a.i.)
alachlor (Lasso) (Screen treated seed)	4
atrazine ¹	2.5
metolachlor (Dual II Magnum) (Concep treated seed)	1.33

¹ Maximum use rate for atrazine is determined by soil type, tillage practices used, surface residue, and state or local restrictions. follow the more restrictive requirements when determining the maximum use rate for atrazine.

PREHARVEST USES

For Use Only in the States of Texas and Oklahoma.

This product may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial applications use at least 2 gallons of water-based carrier per treated acre.

Delay harvest until 30 days after treatment.

BROADCAST RATE PER TREATED ACRE: 1/2 pint (1/4 lbs. a.i.)

SMALL GRAINS (WHEAT, BARLEY AND OATS)

Not Underseeded to Legumes

Important: Observe all PRECAUTIONS. Read and follow CLEANING, MIXING and APPLICATION instructions. If small grains are used for pasture hay, the following restrictions apply:

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
- There is no waiting period between treatment and grazing for non-lactating dairy animals.
- Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
- Do not harvest hay from treated areas before 37 days after treatment.

NOTE: Observe all precautions and restrictions on the labels of products used in tank mix treatments.

WEEDS CONTROLLED

This product, or combinations with listed tank mix partners, will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds, it is recommended that this product be applied in a tank mix with other herbicides. Refer to specific crop tank mix options.

Alkanet ¹	Knawel (German Moss)	Pigweed, Tumble
Bedstraw, Catchweed ¹	Knotweed, Prostrate	Pineappleweed ¹
Bindweed, Field ²	Kochia	Plantain, Broadleaf ²
Buckwheat, Tartary	Ladysthumb	Poppy, Red Horned ¹
Buckwheat, Wild	Lambsquarters, Common	Puncturevine ¹
Carpetweed ¹	Lettuce, Miners ¹	Purslane, Common ¹
Chamomile, Corn	Lettuce, Prickly	Radish, Wild ¹
Chervil, Bur ¹	Mallow, Common	Ragweed, Common ¹
Chickweed, Common ¹	Mayweed, Chamomile	Ragweed, Giant (Buffaloweed) ¹
Cockle, Corn	(Dogfenne) ¹	Rocket, London ¹
Cockle, Cow	Mustard, Blue (Purple) ¹	Rocket, Yellow ¹
Cocklebur, Common	Mustard, Tansy	Salsify (Goatsbeard) ¹
Cornflower (Bachelorbutton) ¹	Mustard Treacle ¹	Sheperdspurse ¹
Dandelion, Common ²	Mustard, Tumble (Jim Hill) ¹	Smartweed, Green
Dock, Curly ²	Mustard, Wild ¹	Smartweed, Pennsylvania
Dragonhead, American ¹	Nightshade, Black	Sorrel, Red (Sheep Sorrel) ¹
Evening Primrose, Cutleaf ¹	Nightshade, Cutleaf ¹	Sowthistle, Annual
Falseflax, Smallseeded ¹	Nightshade, Silverleaf ² (White	Starthistle, Yellow ¹
Fiddleneck (Tarweed) ¹	Horsenettle)	Sunflower, Common (Wild)
Flixweed ¹	Pennycress, Field (Fanweed,	Thistle, Canada ²
Fumitory ¹	Frenchweed,	Thistle, Russian
Gromwell, Corn ¹	Stinkweed)	Yarrow, Common ²
Groundsel, Common ¹	Pepperweed, Peppergrass ¹	Velvetleaf
Hempnettle ¹	Pigweed, Redroot	Vetch ¹
Henbit	(Carelessweed)	
Jacobs Ladder ¹	Pigweed, Rough	

¹ These weeds will be controlled with this product's tank mixtures. Refer to tank mix label for specific weeds controlled.

² Kamba 4SL tank mixes will provide suppression of established broadleaf weeds and control of their seedlings.

RATES AND TIMINGS

Application of this product may be made before, during or after planting small grains. For best performance, make applications when weeds are in the 2- to 3-leaf stage and rosettes are less than 2 inches across. Application of this product to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use this product at 2 to 4 fluid ounces per treated acre in wheat, fall seeded barley, and oats, and at 2 to 3 fluid ounces per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia, Russian thistle and prickly lettuce or dense vegetative growth.

This product used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for this product rate and application timing.

For applications prior to the emergence of weeds or when sulfonylurea resistant weeds are present or suspected, use a minimum of 3 fluid ounces per treated acre of this product with a tank mix herbicide. Non-sulfonylurea herbicides such as 2,4-D or MCPA tank mixed with this product will offer more consistent control of sulfonylurea resistant weeds.

When tank mixing with sulfonylurea herbicides, such as Ally, Amber, Express, Finesse, Glean and Harmony Extra, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1 to 4 pints per 100 gallons of spray or not more than 0.25 to 0.5% by volume. Use the highest rate of

surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

FALL AND SPRING SEEDED WHEAT

THIS PRODUCT MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE. APPLICATIONS TO SPRING SEEDED WHEAT MUST BE MADE BEFORE WHEAT EXCEEDS THE 5-LEAF STAGE. Early developing wheat varieties such as TAM 107, MADISON, or WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

TANK MIX TREATMENTS

This product may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2 to 4 fluid ounces of Kamba 4SL with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8 to 12 fl oz. (0.25 to 0.375 lb ai/A) ¹
MCPA Amine or Ester	MCPA	4 lb/gal	8 to 12 fl oz. (0.25 to 0.375 lb ai/A) ¹
Ally	metsulfuron-methyl	60% DF	1/10 oz.
Amber	triasulfuron	75% DF	0.28 oz.
Express	thifensulfuron + tribenuron-methyl	75% DF	1/6 oz.
Finesse	chlorsulfuron + metsulfuron-methyl	75% DF	1/3 oz.
Glean	chlorsulfuron	75% DF	1/6 oz.
Harmony Extra	thifensulfuron + tribenuron-methyl	75% DF	1/3 oz.
Buctril	bromoxynil ²	2 lb/gal	1 to 1-1/2 pts.
Bronate	bromoxynil + MCPA	4 lb/gal	1 to 2 pts.
Curtail	clopyralid + 2,4-D	2.38 lb/gal	2 to 2-2/3 pts.
Stinger	clopyralid	3 lb/gal	1/4 to 1/3 pt.
Karmex ³	diuron ²	80% DF	1/2 to 1-1/2 lbs.
Sencor ³	metribuzin ²	75% DF	1 to 10 oz.
Tiller ⁴	fenoxaprop-ethyl + MCPA +	2.7 lb/gal	1 to 1.7 pts.

¹ When using formulations other than 4 lb/gal use pounds active per acre listed.

² Herbicides with the same active ingredient and/or different formulations may be used.

³ Tank mixtures for fall seeded wheat only.

⁴ Use 2 fluid ounces of Kamba 4SL only. Do not use if wild oats is the target weed. Do not use on Durum wheat.

SPECIAL USE TANK MIXES FOR SPRING AND FALL SEEDED WHEAT

(See Footnotes for Application Uses)

Apply 3 to 4¹ fluid ounces of Kamba 4SL with:

Product ²	Active Ingredient	Formulation	Amount of Product
2,4-D or MCPA Amine	2,4-D or MCPA	4 lb/gal	1 to 2 pts. ³ (1/2 to 1.0 lb ai/A) ⁴
2,4-D or MCPA Ester	2,4-D or MCPA	4 lb/gal	1 to 1-1/2 pts. ³ (1/2 to 3/4 lb ai/A) ⁴
Ally	metsulfuron-methyl	60% DF	1/20 to 1/10 oz.
Amber	triasulfuron	75% DF	0.14 to 0.28 oz.
Express	thifensulfuron + tribenuron-methyl	75% DF	1/12 to 1/6 oz.
Finesse	chlorsulfuron + metsulfuron-methyl	75% DF	1/6 to 1/3 oz.
Glean	chlorsulfuron	75% DF	1/6 oz.
Harmony Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6 to 1/3 oz
Ally + 2,4-D Amine or Ester ⁵	metsulfuron-methyl + 2,4-D	60% DF + 4 lb/gal	1/20 to 1/10 oz + 8 fl oz.
Amber + 2,4-D Amine or Ester ⁵	triasulfuron + 2,4-D	75% DF + 4 lb/gal	0.14 to 0.28 oz + 8 fl oz.
Express + 2,4-D Amine or Ester ⁵	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/12 to 1/6 oz + 8 fl oz.
Finesse + 2,4-D Amine or Ester ⁵	(chlorsulfuron + metsulfuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/6 to 1/3 oz + 8 fl oz.
Glean + 2,4-D or Ester ⁵	chlorsulfuron + 2,4-D	75% DF + 4 lb/gal	1/6 oz + 8 fl oz.
Harmony Extra + 2,4-D Amine or Ester ⁵	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/6 to 1/3 oz + 8 fl oz.
Roundup ⁶	glyphosate	3.0 lb/gal	12 to 16 fl oz.

¹ This product may be used at 6 fluid ounces on fall seeded wheat in Western Oregon as a spring application only. In CO, KS, NM, OK and TX up to 8 fluid ounces of this product may be applied on fall seeded wheat after it exceeds the 3-leaf stage for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. This product may be tank mixed with 2,4-D amine at 8 fluid ounces after wheat begins to tiller. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For Fall applications only, do not use if the potential for crop injury is not acceptable.

² Do not use low rates of sulfonylurea herbicides, such as Ally, Amber, Express, Finesse, Glean, and Harmony Extra on more mature weeds and/or on dense vegetative growth.

³ NOTE: For use on Fall Seeded Wheat Only. Do Not Use unless potential crop injury will be acceptable.

⁴ When using formulations other than 4 lb/gal use pounds active/acre listed.

⁵ Use for improved control of Russian thistle, flixweed, gromwell, mayweed and fiddleneck.

⁶ This product may be applied at 2 fluid ounces with Roundup as a preplant application to small grains with no waiting period prior to planting. Add 0.5% by volume of an agriculturally approved non-ionic surfactant.

FALL SEEDED BARLEY

THIS PRODUCT MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE.

NOTE: For spring barley varieties that are seeded during the winter months or later, follow the rates and timings given for spring seeded barley.

TANK MIX TREATMENTS

This product may be tank mixed with one or more, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2 to 4 fluid ounces of Kamba 4SL with:

Product ¹	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8 fl oz. (0.25 lb ai/A) ²
MCPA Amine or Ester	MCPA	4 lb/gal	8 to 12 fl oz. (0.25-0.375 lb ai/A)
Ally	metsulfuron-methyl	60% DF	1/20 to 1/10 oz.
Amber	triasulfuron	75% DF	0.14 to 0.28 oz.
Express	thifensulfuron + tribenuron-methyl	75% DF	1/12 to 1/6 oz.
Finesse	chlorsulfuron + metsulfuron-methyl	75% DF	1/6 to 1/3 oz.
Glean	Chlorsulfuron	75% DF	1/6 oz.
Harmony Extra	thinensulfuron + tribenuron-methyl	75% DF	1/6 to 1/3 oz.
Sencor	metibuzin ³	75% DF	1 to 10 oz.
Buctril	bromoxynil	2 lb/gal	1 to 1-1/2 pts.
Bronate	bromoxynil + MCPA	4 lb/gal	3/4 to 1-1/2 pts.

¹ Do not use low rates of sulfonylureas (Ally, Amber, Express, Finesse, Glean, and Harmony Extra) on more mature weeds and/or on dense vegetative growth.

² When using formulations other than 4 lb/gal use pounds active/acre listed.

³ Herbicides with the same active ingredient and/or different formulations may be used.

SPRING SEEDED BARLEY

THIS PRODUCT MUST BE APPLIED BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4-LEAF STAGE.

TANK MIX TREATMENTS

This product may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2 to 3 fluid ounces Kamba 4SL with:

Product ¹	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8 -12 fl. oz. (0.25 to 0.375 lb ai/A) ²
Ally	metsulfuron-methyl	60% DF	1/20 to 1/10 oz.
Amber	Triasulfuron	75% DF	0.14 to 0.28 oz.
Express	thifensulfuron + tribernuron methyl	75% DF	1/12 to 1/6 oz.
Finesse	chlorsulfuron + metsulfuron-methyl	75% DF	1/6 to 1/3 oz.
Glean	Chlorsulfuron	75% DF	1/6 oz.
Harmony Extra	thifensulfuron + tribernuron-methyl	75% DF	1/6 to 1/3 oz.
Sencor	metribuzin ³	75% DF	1 to 10 oz.
Buctril	bromoxynil	2 lb/gal	1 to 1-1/2 pts.
Bronate	bromoxynil + MCPA	4 lb/gal	3/4- 1-1/2 pts.

¹ Do not use low rates of sulfonyleureas (Ally, Amber, Express, Finesse, Glean, and Harmony Extra) on more mature weeds and/or on dense vegetative growth.

² When using formulations other than 4 lb/gal use pounds active/acre listed.

³ Herbicides with the same active ingredient and/or different formulations may be used.

FALL AND SPRING SEEDED OATS

THIS PRODUCT MUST BE APPLIED BEFORE SPRING SEEDED OATS EXCEED THE 5 LEAF STAGE. APPLICATIONS TO FALL SEEDED OATS MUST BE MADE PRIOR TO THE JOINTING STAGE.

TANK MIX TREATMENTS

This product may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2-4 fluid ounces of Kamba 4SL with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8 to 12 fluid oz. (0.25 to 0.375 lb ai/A) ¹

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

SUGARCANE

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions.

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

WEEDS CONTROLLED

This product, when applied at recommended rates, will control many ANNUAL, BIENNIAL and PERENNIAL broadleaf weeds commonly found in sugarcane. (Refer to GENERAL WEED LIST.)

RATES AND TIMINGS

Application of this product may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane. Application rates and timings of this product are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage and Type	Amount Product	Broadcast Rate Per Treated Acre lbs. a.i.
Annual		
Small, actively growing	1/2 to 1 pt	1/4 to 1/2
Established weed growth	1 to 1-1/2 pts	1/2 to 3/4
Biennial	1 to 2 pts	1/2 to 1
Perennial	2 pts	1 to 2 ¹

¹Application made over the top of actively growing sugarcane may result in crop injury.

When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

Retreatments may be made as needed, however, do not exceed a total of 4 pints (2 lbs. a.i.) of this product per treated acre during a growing season.

TANK MIX TREATMENTS

This product may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

Herbicide	Rates per treated acre
Ametryn (Evik)	2/5 to 8
Asulam (Asulox)	2 to 3-1/3
atrazine	2/5 to 4
2,4-D	1/2 to 3 ¹

¹ Application of this product plus 2,4-D tank mix at the higher listed rate range may result in crop injury.

PASTURE, HAY, RANGELAND AND GENERAL FARMSTEAD (NON-CROPLAND)

This product is recommended for use on pasture, hay, rangeland, general farmstead (non-cropland) (including fence rows and non-irrigation ditchbanks), for broadleaf weed and brush control. This product may also be applied to non-cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatment of roadsides and highways, utilities, railroad and pipeline rights-of-way. Noxious weeds must be recognized at the State level but programs may be administered at State, County or other levels.

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions.

This product uses described in this section also pertain to small grains (such as barley, forage, sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

NEWLY SEEDED AREAS, including small grains grown for pasture may be severely injured if rates of this product greater than 1 pint per acre are applied.

ESTABLISHED GRASS CROPS growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Bentgrass, carpetgrass, buffalograss and St. Augustine grass may be injured at rates exceeding 1 pint this product (1/2 lb. a.i.) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill alfalfa, clovers, lespedeza, wild winter peas, vetch and other legumes.

ANIMALS CANNOT BE REMOVED FROM TREATED AREA FOR SLAUGHTER PRIOR TO 30 DAYS AFTER LAST APPLICATION.

THERE IS NO WAITING PERIOD BETWEEN TREATMENT AND GRAZING FOR NON-LACTATING ANIMALS.

TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT

Kamba 4SL Rate per Treated Acre	Days Before Grazing	Days Before Hay Harvest
Up to 1 pint (1/2 lb. a.i.)	7 days	37 days
Up to 1 quart (1 lb. a.i.)	21 days	51 days

NOTE: Observe all precautions and restrictions on labels of products used in tank mixtures.

MIXING AND APPLICATION

This product can be applied using water, oil in water emulsions (including invert systems), or sprayable fluid fertilizer as a carrier. A COMPATIBILITY TEST should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

This product may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment apply 1 to 40 gallons of diluted spray per treated acre in a water-based carrier.

This product may be applied to individual clumps or small areas (SPOT TREATMENT) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to run-off) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

WEEDS CONTROLLED

This product, when applied at recommended rates, will give control of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species commonly found in pasture, hay, rangeland, and general farmstead (non-cropland) areas. (Refer to GENERAL WEED LIST.)

Noted (*) PERENNIAL weeds may be controlled with lower rates of either this product or this product plus 2,4-D. See RATES AND TIMINGS below.

RATES AND TIMINGS

Application rates and timing of this product are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage & Type	Amount Product	Broadcast Rate per Treated Acre (lbs. a.i.)
Annual		
Small, actively growing	1/2 to 1 pt.	1/4 to 1/2
Established weed growth	1 to 1-1/2 pts.	1/2 to 3/4
Biennial ¹		
Rosette diameter		
Less than 3 inches	1/2 to 1 pt.	1/4 to 1/2
3 inches or more	1 to 2 pts.	1/2 to 1
Bolting	2 pts	1
Perennial		
Suppression or top growth	1/2 to 1 qt.	1/2 to 1
Noted (*) Perennials	1 qt.	1
Other perennials	1 qt.	1
Woody Brush & Vines		
Top growth suppression	1/2 to 1 qt.	1/2 to 1
Top growth control ²	1 qt.	1
Stems and stem suppression	1 qt.	1

¹ For best performance, make application when BIENNIAL weeds are in the rosette stage.

² Species noted in GENERAL WEED LIST will require tank mixtures for adequate control.

Retreatments may be made as needed; however, do not exceed a total of 2 quarts (2 lbs. a.i.) of this product per treated acre during a growing season.

TANK MIX TREATMENTS

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND OTHER RESTRICTIONS. This product may be tank mixed with one or more, but not limited to, of the following herbicides for control of grasses, additional broadleaf weeds, and woody brush and vines.

Herbicide	Rates Per Treated Acre (lbs. a.i.)
Pasture, hay, rangeland and general farmstead (non-cropland) use:	
glyphosate (Roundup)	3/4 to 3-3/4
metsulfuron methyl (Ally)	0.0038 to 0.011
paraquat (Gramoxone)	1/2 to 1
picloram (Tordon)	1/8 to 3
triclopyr (Garlon)	3/4 to 8
2,4-D	1/4 to 4

Due to the variations that may occur on formulated products and specific use ingredients (e.g. water supplies), a COMPATIBILITY TEST is recommended prior to actual tank mixing.

CUT SURFACE TREE TREATMENTS

This product may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. A mix of 1 part this product with 1 to 3 parts water should be used in application. Use the lower dilution when treating difficult-to-control species.

FRILL OR GIRDLE TREATMENTS: Make a continuous cut or a series of overlapping cuts using an ax to girdle tree trunk. Spray or paint cut surface with the this product/water mix.

STUMP TREATMENTS: Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

NOTE: For more rapid foliar effects, 2,4-D may be added to this product/water mix.

DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE

This product can be applied when plants are dormant as an undiluted SPOT-CONCENTRATE directly to the soil or as a LO-OIL BASAL BARK treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of this product should be applied directly to the soil as close as possible to the root crown but within 6 to 8 inches of the crown. On sloping terrain, application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying this product directly to the soil. The use rate of this product is dependent on the canopy diameter of the multiflora rose. Examples: Use this product at 1/4, 1 or 2-1/4 fluid ounces of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 2 quarts of this product per acre per year.

LO-OIL BASAL BARK applications of this product should be applied to the basal stem region from the ground up to a height of 12 to 18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying this product to the ground line. Refer to Mixing and Applications above in this section for method of preparing oil-in-water emulsion. Example for making approximately 2 gallons of a Lo-Oil spray solution mixture: combine 1-1/2 gallons water plus 1 ounce emulsifier plus 1 pint this product plus 2-1/2 pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

CONSERVATION RESERVE PROGRAM (CRP) ACRE

This product is recommended for use on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs.

Observe all PRECAUTIONS, MIXING AND APPLICATION directions.

This product treatment will cause injury or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.

Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after grass emergence on newly seeded grasses.

NEWLY SEEDED AREAS

This product may be applied either preplant or postemergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat, or other grain species grown as a cover crop. Postemergence applications may be made after seedling grasses exceed the 3-leaf stage. Rates of this product greater than 1 pint per treated acre may severely injure newly seeded grasses. Preplant

applications - injury to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of this product per treated acre West of the Mississippi River or 20 days per pint East of the Mississippi River.

ESTABLISHED GRASS STANDS

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species: bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustine grass may be injured when treated with this product at rates exceeding 1 pint per treated acre.

WEEDS CONTROLLED

This product, when applied at recommended rates, will control many annual and biennial weeds and provide control and suppression of many perennial weeds. (Refer to GENERAL WEED LIST.)

RATES AND TIMINGS

Application rates and timing of this product treatment are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature.

Weed Type ¹ & Stage	Broadcast Rate Per Treated Acre	
	Amount of Formulated This product (pints)	Equivalent lbs. a.i.
Annuals Small actively growing Established weed growth	1/4 to 1 1	1/8 to 1/2 1/2
Biennials ² Rosette diameter a) less than 3 inches b) 3 inches or greater c) bolting	1/2 to 1 1 to 2 2	1/4 to 1/2 1/2 to 1 1
Perennials ² Suppression/Control	2	1

¹ For best results, treat biennial weeds with this product when they are in the rosette stage of growth. Retreatments may be made as needed; however, DO NOT EXCEED A TOTAL OF 2 QUARTS (2 lbs. a.i.) of this product per treated acre during the growing season.

² Biennial and perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

TANK MIX TREATMENTS

To control grasses and additional broadleaf weeds, this product may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2,4-D, glyphosate (Roundup), paraquat (Gramoxone), metsulfuron (Ally) and others.

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES, AND OTHER RESTRICTIONS.

28
36

ASPARAGUS

IMPORTANT

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions. If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

Do not harvest prior to 24 hours after treatment.

Do not use in the Coachella Valley of California.

Multiple applications may be made per growing season. Do not exceed a total of 1 pint of this product per treated acre per crop year.

RATES AND TIMINGS

Apply this product to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

Weeds	Rate per Treated Acre
Mustard, Black Pigweed, Redroot (Carelessweed) Sowthistle, Annual Thistle, Canada ¹ Thistle, Russian	1/2 to 1 pt. (1/4 to 1/2 lb. a.i.)
Bindweed, Field ¹ Chickweed, Common Goosefoot, Nettleleaf Radish, Wild Thistle, Milk	1pt. (1/2 lb. a.i.)

¹ This product may be applied in a tank mixture with either 2,4-D or Roundup Herbicide for improved control of Canada thistle and field bindweed.

READ AND FOLLOW 2,4-D AND ROUNDUP HERBICIDE PRODUCT LABELING FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

TURF AND LAWNS FOR USE IN GENERAL FARMSTEAD (Non-Cropland) AND SOD FARMS

IMPORTANT

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions.

To avoid injury to newly seeded grasses, application of this product should be delayed until after the second mowing. Furthermore, application rates in excess of 1 pint (1/2 lb. a.i.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass, carpetgrass, buffalograss, and St. Augustine grass.

In areas where roots of sensitive plants extend, do not apply in excess of 1/4 pint (1/8 lb. a.i.) of this product per treated acre on coarse textured (sandy-type) soils, or in excess of 1/2 pint (1/4 lb. a.i.) per treated acre on fine textured (clay-type) soils.

Do not make repeat applications in these areas for 30 days and until previous applications of this product have been activated in the soil by rain or irrigation.

WEEDS CONTROLLED

This product, when applied at recommended rates, will give control of many ANNUAL, BIENNIAL, and noted (*) PERENNIAL broadleaf weeds commonly found in turf. This product will also give growth suppression of many other listed PERENNIAL broadleaf weeds and WOODY brush and vine species. (Refer to GENERAL WEED LIST.)

MIXING AND APPLICATION

Apply 30 to 200 gallons of diluted spray per treated acre (3 quarts to 4-1/4 gallons per 1,000 square feet) depending on density or height of weeds treated and on the type of equipment used.

RATES AND TIMINGS

Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	KAMBA 4SL		
	Pints per treated acre	Lbs. a.i. per treated acre	Teaspoons per 1,000 sq. ft.
Annual			
Small, actively growing	1/2 to 1	1/4 to 1/2	1 to 2-1/4
Established weed growth	1 to 1-1/2	1/2 to 3/4	2-1/4 to 3-1/4
Biennial Rosette diameter			
Less than 3 inches	1/2 to 1	1/4 to 1/2	1 to 2-1/4
3 inches or more	1 to 2	1/2 to 1	2-1/4 to 4-1/2
Perennials and Woody			
Brush and Vines	1 to 2	1/2 to 1	2-1/4 to 4-1/2

For best performance, apply when weeds are emerged and actively growing.

Retreatments may be made as needed; however, do not exceed a total of 2 pints (1 lb. a.i.) this product per treated acre during the growing season.

TANK MIX TREATMENTS

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

Tank mix treatments of this product may be made with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product.

Apply 1/5 to 1/2 pint (1/10 to 1/4 lb. a.i.) of this product per treated acre with 1/2 to 1-1/2 lbs. acid equivalent of 2,4-D, MCPA, or MCPP, or with 3/8 to 1/2 lb. a.i. of bromoxynil. Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed; however, do not exceed 2 pints (1 lb. a.i.) of this product per treated acre during the growing season.

GRASS SEED CROPS:

GRASSES GROWN FOR SEED SUCH AS BERMUDA GRASS, BLUEGRASS, FESCUE AND RYEGRASS

IMPORTANT

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions.

Refer to PASTURE, HAY, RANGELAND, AND GENERAL FARMSTEAD (Non-Cropland) section for possible grazing and feeding restrictions.

Do not use on bentgrass unless possible crop injury can be tolerated.

WEEDS CONTROLLED

This product will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, it is recommended that this product be applied in a tank mix with other herbicides.

Alfalfa ¹	Dock, Curly
Bedstraw, Catchweed	Hemlock, Poison
Bindweed, Field	Knapweed, Russian ¹
Buttercup, Corn	Knawel
Buttercup, Creeping	Knotweed, Prostrate
Buttercup, Western Field	Kochia
Catchfly, Nightflowering	Ladysthumb
Chamomile, Corn	Lambsquarters, Common
Chickweed, Common	Lettuce, Prickly
Chickweed, Mouseear	Mayweed (Dogfennel)
Clover	Ragwort, Tansy
Cockle, White	Sorrel, Red (Sheep Sorrel)
Dock, Broadleaf	Sowthistle, Annual Starwort, Little Thistle, Canada ¹

¹ Top growth only.

RATES AND TIMINGS

Apply 1/2 to 1 pint of this product per treated acre on SEEDLING GRASS after the crop reaches the 3- to 5-leaf stage. Apply up to 2 pints of this product on well-established Perennial grass. DO NOT APPLY AFTER THE GRASS SEED CROP BEGINS TO JOINT. For best performance, make applications when weeds are in the 2- to 4-leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, this product may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 1/2 to 2 pints of Kamba 4SL with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	1 to 4 pts. (0.5 - 2.0 lb ai/A) ¹
MCPA Amine	MCPA	4 lb/gal	1 to 2 pts (0.5 - 1.0 lb ai/A) ¹
Buctril	bromoxynil ²	2 lb/gal	1 to 2 pts.
Curtail	clopyralid + 2,4-D	2.38 lb/gal	1-3/4 to 4 pts.
Karmex	diuron ²	80% DF	2 to 4 lbs.
Stinger	clopyralid	3 lb/gal	1/4 to 1 pt.

- ¹ When using formulations other than 4 lb/gal use pounds active/acre listed.
- ² Herbicides with the same common name and/or different formulations may be used.

ANNUAL GRASS CONTROL

For suppression of ANNUAL GRASS WEEDS such as: Brume, Downy (Cheatgrass), Brame, Ripgut, Fescue, Rattail and Windgrass, apply up to 4 pints of this product per treated acre in the fall or late summer after harvest and burning of established grass seed crops. Applications should be made immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

**PREPLANT DIRECTIONS
(POST HARVEST/FALLOW/CROP STUBBLE/SET-A-SIDE)
FOR BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS**

IMPORTANT

Observe all precautions. Read and follow mixing and application instructions.

WEEDS CONTROLLED

This product may be applied alone or in tank mix combinations with other herbicides registered for this use.

This product can be applied either POST HARVEST in the fall, spring or summer during the FALLOW period or to CROP STUBBLE/SET-A-SIDE acres. This product, when applied at the recommended rates, will control many ANNUAL broadleaf weeds, see the WEEDS CONTROLLED section under small grains. In addition, This product will control or suppress the following BIENNIAL and PERENNIAL broadleaf weeds:

Alfalfa ¹	Knapweed, Spotted
Artichoke, Jerusalem	Nightshade, Silver
Bindweed, Field	Redvine
Bindweed, Hedge	Smartweed, Swamp
Blueweed, Texas	Sowthistle, Perennial ¹
Bursage (Bur Ragweed, Povertyweed,	Spurge, Leafy
Lakeweed) ¹	Thistle, Bull
Dandelion, Common ¹	Thistle, Canada ²
Dock, Curly ¹	Thistle, Milk
Dogbane, Hemp	Thistle, Musk
Garlic, Wild ²	Thistle, Plumeless
Horsenettle, Carolina	Thistle, Scotch
Knapweed, Diffuse	Trumpet creeper (Buckvine)

¹ These perennials may be controlled using this product at rates lower than those recommended for other listed perennial weeds. See RATES AND TIMINGS under this heading).

² See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control program for these weeds.

RATES AND TIMINGS

Apply product as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or cropland or crop stubble the following spring or summer. Agriculturally approved spray additives, such as surfactants or oils, may be used to enhance spray coverage and the herbicides penetration of weed foliage. See CROPPING RESTRICTIONS for recommended interval between application and planting to prevent crop injury.

For best performance, make application when ANNUAL weeds are less than 6 inches tall, when BIENNIAL weeds are in the rosette stage, and to PERENNIAL weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright PERENNIAL broadleaf weeds,

such as Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, such as field bindweed and hedge bindweed, are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets, after the effective period for this product. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of this product see the RATES AND TIMINGS section under the SMALL GRAINS heading for details.

KAMBA 4SL RATES PER TREATED ACRE:

Weed Type	Amount of Product Per Treated
Annual	1/2 to 1 pts
Biennial	1 to 2 pts
Perennial	
Suppression	1 to 2 pts
Control - noted ⁽¹⁾ perennials	2 pts
Control - other perennials	2 pts

Retreatments may be made as needed; however, do not exceed a total of 4 pints of this product per treated acre during any given fallow period.

TANK MIX TREATMENTS

This product may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled and geographic or other restrictions.

BROADCAST RATE PER TREATED ACRE FOR ANNUAL WEED CONTROL:

Apply 1/4 to 1 pint of Kamba 4SL with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Aatrex 4L ¹	atrazine	4 lb/gal	1/2 to 6 pts.
Aatrex Nine-0 ¹	atrazine	90% DF	1/2 to 3.3 lbs.
Amber ²	trisulfuron	75% DF	0.28 to 0.35 oz.
Ally ²	metsulfuron-methyl	75% DF	0.1 oz.
Fallow Master	glyphosate + dicamba	2.0 lb/gal	22 to 44 fl. oz.
Finesse ²	chlorsulfuron + metsulfuron-methyl	75% DF	0.2 oz.
Gramoxone Extra	paraquat	2.5 lb/gal	1-1/2 pts.
Kerb'	pronamide	50-W	1/2 to 1.0 lb.
Landmaster BW	glyphosate + 2,4-D	2.4 lb/gal	27 to 54 fl oz.
Roundup	glyphosate	3 lb/gal	8 to 48 fl oz.

Sencor DF ¹	metribuzin	75% DF	1/2 to 1 lb.
Sencor 4 ¹	metribuzin	4 lb/gal	3/4 to 1-1/2 pts.
2,4-D	2,4-D	4 lb/gal	1 - 2 pts. (1/2 to 1 lb ai/A) ³

¹ Tank mixes of this product with these products may be subject to special restrictions. See the Product Label of the tank mix partner for intended use rates, restrictions and other precautions.

² When tank mixing with sulfonylurea herbicides refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1 to 2 quarts per 100 gallons of spray or not more than 0.25 to 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea resistant weeds may not be controlled by tank mixes of this product and a sulfonylurea. Refer to the this product tank mix section for alternative tank mixes.

³ When using formulations other than 4 lb/gal, use pounds active/acre listed.

**BROADCAST RATE PER TREATED ACRE
FOR BIENNIAL AND PERENNIAL WEED CONTROL:**

Apply 1 to 4 pints of Kamba 4SL with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Curtail	clopyralid + 2,4-D	2.38 lb/gal	2 to 4 pts.
2,4-D	2,4-D	4.0 lb/gal	2 to 4 pts. (1 - 2 lb ai/A) ¹
Landmaster BW	glyphosate + 2,4-D	2.4 lb/gal	54 fl oz.
Roundup	glyphosate	3.0 lb/gal	1 to 5 qts.
Tordon 22K	picloram	2.0 lb/gal	1/2 to 1pt.

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

SPECIAL TANK MIX TREATMENTS

For suppression of perennial weeds, apply 1/2 to 1 pint of this product with 8 to 16 fluid ounces of Roundup Herbicide per treated acre.

For wild garlic control, apply 1 pint of this product with 3 pints of 2,4-D LV Ester (4 lb/gal) per treated acre. Apply when wild garlic is 4 to 8 inches tall.

For Canada thistle control, use this product, this product plus Curtail, or this product plus Roundup Herbicide tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brume, jointed goatgrass, common rye and volunteer wheat when they are actively growing. Use 1 pint this product with 1/2 to 1 lb Kerb 50W. Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply this product plus Landmaster BW or Fallow Master Herbicide to give improved control of kochia, wild buckwheat, prickly lettuce, field bindweed and Canada thistle. Use 1/8 to 1/4 pint this product plus 22 to 54 fluid ounces of Landmaster BW or Fallow Master Herbicide for annual weed control or 1/4 to 1/2 pint this product plus 22 to 54 fluid ounces of Landmaster BW or Fallow Master Herbicide for perennial weed suppression.

CROPPING RESTRICTIONS

The following recommendations are based on this product use rates up to 4 pints per treated acre.

CORN, SORGHUM and SOYBEANS may be planted in the spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, treated areas should be cultivated to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of this product per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of this product per treated acre. Exclude days when ground is frozen.

WHEAT may be planted in the fall or spring following applications. Also, spot applications may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

East of the Mississippi River, the interval is 20 days per pint of this product per treated acre or 1.25 days per 1 ounce. Moisture is essential for this product degradation. Exclude days when ground is frozen.

West of the Mississippi River, the interval is 45 days per pint of this product per treated acre or 3 days per ounce. Moisture is essential for this product degradation. Exclude days when ground is frozen.

Following a normal harvest of barley, oats, or wheat, any rotational crop may be planted. If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow up with the planting of a sensitive crop.

**CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND
(SPOT APPLICATION ONLY)**

**FOR USE ONLY IN THE STATES OF IDAHO, MONTANA, NEVADA, OREGON, UTAH, AND
WASHINGTON**

IMPORTANT

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions.

Do not treat subirrigated cropland or areas where the soil remains saturated with water throughout the year.

Make only one application of this product per year.

WEEDS CONTROLLED

This product, when applied at recommended rates, will control many broadleaf weeds including:

Bindweed, Field	Knapweed
Dock, Broadleaf (Bitterdock)	Russian Ragwort
Dock, Curly	Tansy Spurge
Knapweed, Black	Leafy Thistle and Canada

RATES AND TIMINGS

This product may be applied at any time following a crop harvest to stubble fallow or other cropland. Application should be made when weeds are actively growing and prior to a killing frost.

Apply 1 quart (1 lb. a.i.) of this product per treated acre. Application may be made up to one month prior to the planting of wheat.

NOTE: Do not use unless injury to wheat or rotated barley will be acceptable.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application. Do not plant broadleaf crops such as alfalfa, beans, peas, potatoes, or sugarbeets into treated areas until two years after application.

In most cases, treatments will not kill perennial weed seedlings, which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

WIPER APPLICATION USES

IMPORTANT

Observe all PRECAUTIONS. Read and follow MIXING AND APPLICATION instructions .

This product may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution containing 1 part this product to 1 part water. Do not contact desirable vegetation with herbicide solution. Wiper application should only be made to crops (including pastures) and non-cropland areas described in this label with the exception of Grain Sorghum (Milo).

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL:

Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. If recycling or reconditioning not available, puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring

at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

(062909R)

Kamba is a registered trademark of Nufarm Americas Inc.
All other trademarks are the property of their respective owners.