



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

August 30, 2021

Ms. Rebecca M. Horton
Consultant/ Agent
Bluewater Chemgroup, Inc.
P.O. Box 11384
Fort Wayne, IN 46857

Subject: Notification per PRN 98-10 – Minor label revisions
Product Name: Liquid Harvest Mesotrione
EPA Registration Number: 83190-6
Application Date: June 14, 2021
Decision Number: 576827

Dear Ms. Horton:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.


The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 83190-6
Decision No. 576827

If you have any questions, you may contact BeWanda Alexander at (703)347-0313 or by email at alexander.bewanda@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Heather E. McFarley". The signature is written in a cursive style with a loop at the end of the last name.

Heather McFarley,
Acting Product Manager 24
Fungicide and Herbicide Branch,
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

83190-6

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

08/30/2021

MESOTRIONE	Group	27	Herbicide
------------	-------	-----------	-----------

Liquid Harvest Mesotrione

Optional Marketing Claims:

- [Provides Selective and Residual Control of Weeds in Ornamental Turfgrasses]
- [For Use On: Kentucky Bluegrass, Buffalograss, St. Augustinegrass & Other Species]
- [Controls Weeds in Lawns & Turfgrass]
- [Broadleaf Herbicide]

Active Ingredient:	By Weight
Mesotrione: 2-[4-(methylsulfonyl)-2-nitrobenzoyl]-1,3-cyclohexanedione	40.0%
Other Ingredients:	<u>60.0%</u>
TOTAL:	100.0%

Contains 4 lbs. Mesotrione per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back label [booklet] for additional Precautionary Statements, First Aid and complete Directions for Use.

Net Contents: 8 FL OZ (237 mL), 1 PT (473 mL), 1 QT (946 mL), 1 GAL (3.8 liters), 5 GAL (18.9 liters)

EPA Reg. No. 83190-6
EPA Est. No. 72838-IN-003

BLUEWATER CHEMGROUP, INC.
PO Box 11384
Fort Wayne, IN 46857

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF ON SKIN	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by the poison control center or doctor. • DO NOT give anything to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
HOTLINE NUMBER	
For Chemical Emergency Assistance (Spill, Leak, Fire, or Accident) call CHEMTREC: 1-800-424-9300 .	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid contact with skin, eyes, or clothing.

Personal Protection Equipment (PPE)

Applicators and Other Handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves (barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC), or Viton)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove and wash contaminated clothing before reuse.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Environmental Hazards

DO NOT apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water or rinsate.

Surface Water Advisory

This product may contaminate water through drift or spray in wind. This product has a high potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Physical and Chemical Hazards

DO NOT use or store near heat or open flame. **DO NOT** use with or store near any oxidizing or reducing agents. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- coveralls
- shoes plus socks
- chemical-resistant gloves (barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, and viton \geq 14 mils)

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard, 40 CFR Part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, and greenhouses.

DO NOT enter treated areas without protection clothing until sprays have dried.

Liquid Harvest Mesotrione is applied pre- and post-emergence to provide selective contact and residual control of turfgrass weeds. If applied pre-emergence, it is absorbed when emerging from the soil. Pre-emergence activity and control is reduced under dry conditions. Activate **Liquid Harvest Mesotrione** with 0.15 inches of irrigation if rain hasn't fallen within 10 days of application. Post-emergent control is obtained by absorption into the soil and contact with foliage. Growth ceases post-application, weeds turn white from chlorophyll loss, and will die within three weeks. Make a repeat application after 2-3 weeks to improve post-emergence weed control. Add a non-ionic surfactant when making post-emergence applications.

Turfgrass color can temporarily become white during treatment. Whitening typically occurs 5-7 days post-application and lasts for several weeks. A second application to the same site will cause less whitening of plant tissue.

Liquid Harvest Mesotrione controls weeds prior to and during seeding of certain turfgrasses during turf renovation (see **New Seedings**). If making pre-emergence application to established turf, tank mix **Liquid Harvest Mesotrione** with other pre-emergence herbicides for longer residual and broad spectrum control.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Applications

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For all applications, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Spray Drift Advisories

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform disposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potentially increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

WINDBLOWN SOIL PARTICLES

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product in prevailing conditions may be expected to result in off-site movement.

RESISTANCE MANAGEMENT

There is no known resistance to **Liquid Harvest Mesotrione** and no known instances of cross-resistance between the active ingredient in **Liquid Harvest Mesotrione** (mesotrione, a Group 27 HPPD inhibitor) and other classes of herbicides. **DO NOT** apply less than the specified label instructions and rotate to a herbicide product with a different mode of action to prevent the occurrence of weed resistance if the maximum number of applications and/or rate of product has been met and weeds are not controlled.

Suspected herbicide-resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of non-controlled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

To minimize the occurrence of resistant biotypes, observe the following weed management practices:

- Scout application site before and after herbicide applications.
- Start with a clean application site, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g., a selective and/or a residual herbicide) and cultural practices (e.g., tillage or crop rotation) where appropriate).
- Utilize the specified label rate for the most difficult to control weed in your field. Avoid tank mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture directions that encourage application rates of this product below the label directions.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Report any incidence of repeated non-performance of this product on a particular weed to local extension specialists, certified crop advisors or your Bluewater Chemgroup Inc. representative.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of actions for each target weed.

Approved Use Sites

Liquid Harvest Mesotrione can be applied to commercial and residential turfgrasses. Non-crop area use sites include golf courses, sod farms, athletic fields, parks, residential and commercial properties, cemeteries, airports, and lawns.

Apply **Liquid Harvest Mesotrione** at reduced rates of 4 fl. oz./A (0.125 lb. a.i./A) or less if tank mixing with atrazine, bentazon, or simazine. Before tank mixing **Liquid Harvest Mesotrione** with other herbicides, conduct a compatibility, safety, and efficacy test before treating larger areas. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Thoroughly clean application equipment after use to avoid injury to sensitive plants.

To avoid injury to sensitive species, keep traffic out of treated areas until sprays have dried; irrigate soil lightly to move **Liquid Harvest Mesotrione** from turf foliage before resuming normal irrigation.

Turfgrass Use Restrictions:

- **DO NOT** overspray or allow spray to drift to ornamentals or flower beds and gardens. Roses and daylilies are particularly sensitive to **Liquid Harvest Mesotrione**.
- **DO NOT** apply more than 8 fl. oz./A (0.25 lb. AI/A) in a single application
- **DO NOT** apply more than 16 fl. oz. (0.50 lb. AI per acre per year) per acre per year.
- **DO NOT** make more than 4 applications per year when using reduced application rates.
- If multiple applications are made, they must be made no closer than 14 days apart.
- **DO NOT** plant any crop other than turfgrass for 18 months post-application of **Liquid Harvest Mesotrione** to avoid turfgrass injury.

- **DO NOT** apply organophosphate or carbamate insecticides within 7 days of applying **Liquid Harvest Mesotrione**.
- **DO NOT** apply **Liquid Harvest Mesotrione** through any type of irrigation system.
- **DO NOT** make aerial applications.
- **DO NOT** use treated clippings to mulch trees or vegetable/flower gardens.
- **DO NOT** apply this product on Bentgrass, *Poa annua*, kikuyugrass, zoysiagrass, seashore paspalum, and bermudagrass, if plant injury is unacceptable. Maintain a 5-foot buffer between treated areas and bentgrass or *Poa annua* greens.
- **DO NOT** apply over the top of exposed roots of trees and ornamentals.
- **DO NOT** use on golf course putting greens; maintain a minimum of a 5-foot buffer between putting greens and treated areas.
- Residential Lawns: **DO NOT** make broadcast applications for pre- and post-emergent weed control unless the home lawn is being reseeded and/or renovated as whitening of some turfgrasses may occur.

Turfgrass Species & Application Rates

Species	Application Rate (Fl. Oz. per Acre)
Kentucky bluegrass (<i>Poa pratensis</i>) Centipedegrass (<i>Eremochloa ophiuroides</i>) Buffalograss (<i>Buchloe dactyloides</i>) Tall fescue (<i>Festuca arundinacea</i>)	5-8 (0.155-0.25 lb. AI/A)
Perennial ryegrass* (<i>Lolium perenne</i>) Fine fescue* (creeping red, chewings and hard) <i>Festuca</i> spp.	5 (0.155 lb. AI/A)
St. Augustinegrass* (grown for sod) (<i>Stenotaphrum secundatum</i>)	4 (0.125 lb. AI/A)

*See additional rate instructions below.

APPLICATION INSTRUCTIONS

Pre-Emergence Applications:

Apply 4-8 fl. oz. of **Liquid Harvest Mesotrione** per acre (0.125-0.25 lb. AI/A) in at least 30 gallons of water per acre before seeds germinate and as close to seed germination as possible. Combine this product with another pre-emergence herbicide for extended control of crabgrass and foxtail.

Pre-Emergence Application Precautions:

Liquid Harvest Mesotrione is most effective on established turf when applied post-emergence unless it is combined with another soil active herbicide.

Pre-Emergence Application Restrictions:

- **DO NOT** exceed 5 fl. oz. per acre (0.155 lb. AI/A) per application to perennial ryegrass, fine fescues, or mixed stands that consist of >50% perennial ryegrass and/or fine fescue.
- St. Augustinegrass sod: **DO NOT** exceed 4 fl. oz. per acre (0.125 lb. AI/A).

Application to New Seedlings/New Lawns

Apply 5-8 fl. oz. **Liquid Harvest Mesotrione** per acre (0.155-0.025 lb. AI/A) in at least 30 gallons of water per acre before seeding or after seeding of resistant turfgrass species listed below, except fine fescue, as application to fine fescue can reduce grass density. **Liquid Harvest Mesotrione** can be effectively used on grass seed blends that contain <20% by weight hard/fine fescue. For optimal control, apply at grass seeding or as close to seeding as possible.

New Seedlings/New Lawns Restrictions:

- **DO NOT** spray on newly germinated turfgrass. Delay treatment until grass has been mowed 2-4 times and/or 4 weeks after emergence (whichever is longer).

Post-Emergence Application Instructions:

Apply 4-8 fl. oz. of **Liquid Harvest Mesotrione** per acre (0.125-0.25 lb. AI/A) in at least 30 gallons of water per acre with a NIS surfactant. Make a repeat application 2-3 weeks later for optimal weed control. Apply to young, actively growing weeds.

Post-Emergence Application Precautions:

Moisture stress and application to mature weeds can reduce herbicide efficacy.

Bentgrass (*Agrostis* spp.)/Nimbleweed (*Muhlenbergia schreberi*) Control:

Apply 5 fl. oz. **Liquid Harvest Mesotrione** per acre (0.155 lb. AI/A) in at least 30 gallons of water per acre combined with a NIS surfactant at 2-3 week intervals for a maximum of 3 applications. For optimal Bentgrass control, apply in late summer/early fall just prior to new growth.

St. Augustinegrass (Sod uses only) and Centipedegrass Treatment:

Apply to established turf ONLY.

St. Augustinegrass (Sod uses only) and Centipedegrass Restrictions:

- **DO NOT** exceed 4 fl. oz. (0.125 lb. AI) **Liquid Harvest Mesotrione** if tank mixing with Atrazine or Simazine.
- **DO NOT** exceed 0.5 lb. atrazine or simazine active ingredient. See atrazine/simazine labels for precautions and restrictions.

Dormant Bermudagrass Application only:

Apply 5 fl. oz. per acre of **Liquid Harvest Mesotrione** (0.155 lb AI/A) to control winter weeds listed in the **Weeds Controlled** table below. Make a repeat application 2-3 weeks later. Application of **Liquid Harvest Mesotrione** to semi-dormant turf will cause bermudagrass whitening.

Spot Applications of this product

Spray Mix	Application Rate	Rate of this product	Rate of NIS adjuvant
2 gallons	1 gallon per 1,000 sq. ft.	1 teaspoon	3 teaspoons

Spot Application Restrictions:

- **DO NOT** apply more than 16 fl. oz. of **Liquid Harvest Mesotrione** per acre per year (equivalent to 0.5 lb. AI per acre per year).

WEEDS CONTROLLED USING PRE-EMERGENCE APPLICATION

Apply **Liquid Harvest Mesotrione** with a grass pre-emergence herbicide, except when used to control weeds in new seedings. **Liquid Harvest Mesotrione** will control the following weeds using pre-emergence application:

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crusgalli</i>
Bentgrass (Creeping)	<i>Agrostis stolonifera</i>
Bluegrass (Annual)*	<i>Poa annua</i> *
Buckhorn Plantain	<i>Plantago lanceolata</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed (Common)	<i>Stellaria media</i>
Chickweed (Mouseear)	<i>Cerastium vulgatum</i>
Clover (Large Hop)	<i>Trifolium aureum</i>
Clover (White)	<i>Trifolium repens</i>
Crabgrass (Large)	<i>Digitaria sanguinalis</i>
Crabgrass (Smooth)	<i>Digitaria ischaemum</i>
Crabgrass (Southern)	<i>Digitaria ciliaris</i>
Foxtail (Yellow)	<i>Setaria glauca</i>
Galinsoga	<i>Galinsoga ciliate</i>
Lambsquarters (Common)	<i>Chenopodium album</i>
Pigweed (Redroot)	<i>Amaranthus retroflexus</i>
Pigweed (Smooth)	<i>Amaranthus hybridus</i>
Purslane (Common)	<i>Portulaca oleracea</i>
Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Smartweed (Pale)	<i>Polygonum lapathifolium</i>
Smartweed (Pennsylvania)	<i>Polygonum pensylvanicum</i>
Speedwell (Persian)	<i>Veronica persica</i>
Speedwell (Purslane)	<i>Veronica peregrina</i>
Wild Carrot	<i>Daucus carota</i>

*Suppression only.

WEEDS CONTROLLED USING POST-EMERGENCE APPLICATION

Make a second application of **Liquid Harvest Mesotrione** 2-3 weeks after initial treatment. For optimal control add a NIS-type surfactant and apply to young, actively growing weeds. **Liquid Harvest Mesotrione** will control the following weeds using post-emergence application:

Common Name	Scientific Name
Barnyardgrass	<i>Echinochloa crusgalli</i>
Bentgrass (Creeping)	<i>Agrostis stolonifera</i>
Buckhorn Plantain	<i>Plantago lanceolata</i>
Carpetweed	<i>Mollugo verticillata</i>
Chickweed (Common)	<i>Stellaria media</i>
Chickweed (Mouseear)	<i>Cerastium vulgatum</i>
Clover (Large Hop)	<i>Trifolium aureum</i>
Clover (White)	<i>Trifolium repens</i>
Crabgrass (Large)*	<i>Digitaria sanguinalis*</i>
Crabgrass (Smooth)*	<i>Digitaria ischaemum*</i>
Crabgrass (Southern)*	<i>Digitaria ciliaris*</i>
Curly dock	<i>Rumex crispus</i>
Dandelion (Catsear)	<i>Hypochoeris radicata</i>
Dandelion (Common)	<i>Taraxacum officinale</i>
Florida Betony	<i>Stachys floridana</i>
Florida Pusley	<i>Richardia scabra</i>
Foxtail (Yellow)	<i>Setaria glauca</i>
Galinsoga	<i>Galinsoga ciliate</i>
Goosegrass*	<i>Eleusine indica*</i>
Ground Ivy	<i>Glechoma hederacea</i>
Heal-All	<i>Prunella vulgaris</i>
Henbit	<i>Lamium amplexicaule</i>
Lambsquarters (Common)	<i>Chenopodium album</i>
Lawn Burweed	<i>Soliva sessilis</i>
Lovegrass (Tufted)	<i>Eragrostis pectinacea</i>
Marestail	<i>Conyza Canadensis</i>
Nimblewill	<i>Muhlenbergia schreberi</i>
Nutsedge (Yellow)	<i>Cyperus esculentus</i>
Oxalis	<i>Oxalis stricta</i>
Pigweed (Redroot)	<i>Amaranthus retroflexus</i>
Pigweed (Smooth)	<i>Amaranthus hybridus</i>
Purslane (Common)	<i>Portulaca oleracea</i>
Shepherd's Purse	<i>Capsella bursa-pastoris</i>
Smartweed (Pale)	<i>Polygonum lapathifolium</i>
Smartweed (Pennsylvania)	<i>Polygonum pennsylvanicum</i>
Sowthistle	<i>Sonchus oleraceus</i>
Swinecress	<i>Coronopus didymus</i>
Thistle (Canada)	<i>Cirsium arvense</i>
Verbena	<i>Verbena hastate</i>
Wild Carrot	<i>Daucus carota</i>
Wild Violet	<i>Viola pratensis</i>
Windmill-grass	<i>Chloris verticillata</i>

*For optimal control, apply to less than 4 tiller crabgrass and goosegrass.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep container tightly closed when not in use. Keep away from heat and flame. **DO NOT** store near seed, fertilizers, or foodstuffs. Can be stored at temperatures as low as minus 20°F. Keep away from heat and flame.

Pesticide Disposal: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Open dumping is prohibited.

Container Handling: Non-refillable container. **DO NOT** reuse or refill this container. Offer for recycling if available.

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of BlueWater Chemgroup, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold BlueWater Chemgroup, Inc. and Seller harmless for any claims relating to such factors.

BlueWater Chemgroup, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or BlueWater Chemgroup, Inc., and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BLUEWATER CHEMGROUP, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither BlueWater Chemgroup, Inc. nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF BLUEWATER CHEMGROUP, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF BLUEWATER CHEMGROUP, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

BlueWater Chemgroup, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of BlueWater Chemgroup, Inc.

All trademarks are the property of their respective owners.