



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

December 14, 2016

Annelle Finstad  
Registration Specialist  
Winfield Solutions LLC  
P.O. Box 64589  
St. Paul, MN 55164-0589

Subject: Notification per PRN 98-10 and 2001-5 – Update mode of action, company name and address, emergency service phone number, referral statements, and warranty disclaimer and limitation of liability  
Product Name: TEBUCURE FUNGICIDE 3.6  
EPA Registration Number: 83222-19  
Application Date: 11/22/2016  
Decision Number: 523961

Dear Ms. Finstad:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notices (PRN) 98-10 and (PRN) 2007-4 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and PRN 2007-4 and finds that the action requested falls within the scope of PRN 98-10 and 2007-4. The label submitted with the application has been stamped “Notification” and will be placed in our records.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Page 2 of 2  
EPA Reg. No. 83222-19  
Decision No. 523961

If you have any questions, please contact Maryam K. Muhammad by phone at 703-347-0301, or via email at [Mohammad.maryam@epa.gov](mailto:Mohammad.maryam@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to be 'HJ', written in a cursive style.

Hope Johnson, Product Manager 21  
Fungicide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

**NOTIFICATION**

83222-19

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

12/14/2016

**MASTER LABEL**

**GROUP 3 FUNGICIDE**

**Tebucure Fungicide 3.6**

**Sublabel A: Agricultural Uses**

Asparagus, Beans (fresh & dry except succulent shelled), Cucurbit Vegetable Group, Dry Bulb Onion, Garlic, Great-Headed (Elephant) Garlic, Welsh Onion, Shallot, Green Onion, Leek, Spring Onion, Scallion, Japanese Bunching Onion, Green Shallots, Green Eschalots, Garden Beet, Leafy Brassica Greens Group, Okra and Turnip; Barley, Corn (sweet corn, field corn, field corn grown for seed and popcorn), Cotton, Grasses Grown for Seed, Peanuts, Soybeans, Sunflower, Wheat and Seed Treatment (sweet corn, field corn, field corn grown for seed and popcorn); Lychee, and Pecan; Hops and Leatherleaf Fern.

**Sublabel B: Turf and Ornamental Uses**

**A. Disease Control in Golf Course Turf**

**B. Disease Control in Field, Nursery and Container Ornamentals and Commercial and Residential Landscapes** including Roses, Flowers, Ornamental Crabapples, Dogwoods and Other Landscape Trees, Azaleas, Camellias, Rhododendrons and Other Landscape Ornamental Shrubs, Ground Covers, Vines and Leatherleaf Fern.

<b>ACTIVE INGREDIENT:</b>	<b>% BY WT.</b>
Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....	38.7%
<b>OTHER INGREDIENTS:</b> .....	<u>61.3%</u>
<b>TOTAL:</b>	100.0%

Contains 3.6 pounds tebuconazole per gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**Manufactured for By:**  
Direct AG Source, LLC  
30473 260<sup>th</sup> Street  
Eldora, IA 50627  
Winfield Solutions, LLC  
P.O. Box 64589  
St. Paul, MN 55164-0589

EPA Reg. No. 83222-19

EPA Est. No.

1/1025/6

## Sublabel A: Agricultural Uses

GROUP 3 FUNGICIDE

# Tebucure Fungicide 3.6

For control of specified diseases on [asparagus], [barley], [beans], [corn], [cotton], [cucurbit vegetables], [grasses grown for seed], [hops], [dry bulb onion and garlic], [green onions], [leafy Brassica greens], [garden beet roots and tops], [lychee], [okra], [pecan], [peanut], [soybean], [sunflower], [turnip], [and] [wheat].

### ACTIVE INGREDIENT:

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-  
1H-1,2,4-triazole-1-ethanol.....38.7%

**OTHER INGREDIENTS:**.....61.3%

**TOTAL** 100.0%

Contains 3.6 pounds tebuconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

FIRST AID	
<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>Note to Physician:</b> No specific antidote. Treat symptomatically.	
<b>Symptoms of Poisoning:</b> The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 7:30 AM-3:30 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, Contact 1-877-424-7452 for emergency medical treatment information.	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

**SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY.**

Manufactured for By:  
Direct AG Source, LLC

EPA Reg. No. 83222-19  
EPA Establishment No.

30473 260<sup>th</sup> Street  
Eldora, IA 50627  
Winfield Solutions, LLC  
P.O. Box 64589  
St. Paul, MN 55164-0589

Net Contents  
Lot No.

1/1025/6

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as bather laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in WPS.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Groundwater Advisory:** Tebuconazole is known to leach through soil into the ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water

features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

### PRODUCT INFORMATION

**Spray Volume:** Apply TEBUCURE FUNGICIDE 3.6 by ground in a minimum of 10 gallons of spray solution per acre or by air in a minimum of 5 gallons of spray solution per acre. Check equipment calibration frequently. For best disease control, complete coverage and uniform application are essential, especially when lower spray volumes are used. Use the appropriate spray volume to ensure complete coverage.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regular serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

**Chemigation:** Apply TEBUCURE FUNGICIDE 3.6 through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse

systems) used for pesticide application to a public water system unless the pesticide label-prescribed

safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

**Mixing:** Add specified amount of TEBUCURE FUNGICIDE 3.6 to the spray tank while filling with water to the desired level. Maintain agitation while mixing. If other materials are added to the spray tank, be sure that the TEBUCURE FUNGICIDE 3.6 is thoroughly dispersed before other materials are added. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

**Compatibility:** To determine the compatibility of TEBUCURE FUNGICIDE 3.6 with other mixing partners, use the following procedure: Pour the labeled proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information contact your local [Direct AG Source, LLC](#) [Winfield Solutions, LLC](#) representative.

### **Resistance Management Statement**

TEBUCURE FUNGICIDE 3.6 is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to TEBUCURE FUNGICIDE 3.6 and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. [Direct AG Source](#) [Winfield Solutions, LLC](#) encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

## AGRICULTURAL CROPS

### APPLICATION DIRECTIONS

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Asparagus</b>	Rust ( <i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre
	<p><b>Notes:</b> Apply TEBUCURE FUNGICIDE 3.6 as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre (0.11 lbs. ai — 0.17 lbs. ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl. oz/acre or 0.51 lbs. ai/acre).</p>	
<p><b>Comments:</b> Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 is a sterol demethylation inhibitor (DMI) fungicide (Group3). Alternating TEBUCURE FUNGICIDE 3.6 with other DMI fungicides may lead to resistance. Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Barley</b>	Rust ( <i>Puccinia</i> spp.) Head Blight ( <i>Fusarium</i> spp.) - Suppression	4 fl. oz. per acre
	<p><b>Notes:</b> Apply TEBUCURE FUNGICIDE 3.6 in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of TEBUCURE FUNGICIDE 3.6 may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of TEBUCURE FUNGICIDE 3.6. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p><b>Application timing directions:</b>  <b>Rusts:</b> Apply TEBUCURE FUNGICIDE 3.6 at the earliest sign of rust pustules on foliage.  <b>Fusarium head blight:</b> Optimal timing of TEBUCURE FUNGICIDE 3.6 for Fusarium Head Blight suppression is when stem heads have fully emerged (Feekes 10.5) on 50% of the plants.</p>	
<p><b>Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		



CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Beans</b> (fresh & dry except succulent shelled)	Rust ( <i>Uromyces appendiculatus</i> )	4 to 6 fl. oz. per acre
	<b>Notes:</b> Apply TEBUCURE FUNGICIDE 3.6 in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: TEBUCURE FUNGICIDE 3.6 may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per crop season. Beans, dry: TEBUCURE FUNGICIDE 3.6 may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per crop season.	
<b>Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Corn</b> (sweet corn, field corn, field corn grown for seed and popcorn)	Rust ( <i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre
	Northern leaf blight ( <i>Helminthosporium turcicum</i> )	
	Southern leaf blight ( <i>Helminthosporium maydis</i> )	
	Northern leaf spot ( <i>Helminthosporium carbonum</i> )	
	Gray leaf spot ( <i>Cercospora zea-maydis</i> )	
<b>Notes:</b> Apply TEBUCURE FUNGICIDE 3.6 in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14- day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of TEBUCURE FUNGICIDE 3.6 may be applied per acre per crop season. <b>Sweet corn:</b> TEBUCURE FUNGICIDE 3.6 may be applied up to 7 days before harvest of ears or forage and 49 days before the harvest of fodder. Field, seed or popcorn: TEBUCURE FUNGICIDE 3.6 may be applied up to 21 days before harvest of forage and 36 days before the harvest of grain or fodder.		
<b>Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 days. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

<b>Cotton</b>	Southwestern cotton rust ( <i>Puccinia cacabata</i> )	6 to 8 fl. oz. per acre
	<b>Notes:</b> Apply TEBUCURE FUNGICIDE 3.6 in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. TEBUCURE FUNGICIDE 3.6 may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per crop season.	
<b>Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (RED = 12 hours.		

<b>CROP</b>	<b>DISEASE</b>	<b>RATE OF TEBUCURE FUNGICIDE 3.6</b>
<b>Cucurbit Vegetables Group</b> Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )	4 to 6 fl. oz. per acre
	Gummy stem blight — suppression ( <i>Didymella bryonae</i> ) (watermelon, squash, pumpkin and melons only)	8 fl. oz. per acre
<b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. TEBUCURE FUNGICIDE 3.6 may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per crop season.		

**Comments:** For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3).  
Restricted-entry interval REI = 12 hours.

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Dry bulb onion</b> <b>Garlic</b> <b>Great-headed (elephant) garlic</b> <b>Shallot</b>	White rot <i>(Sclerotium cepivorum)</i>	<b>White rot:</b> 20.5 fl. oz. per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust <i>(Puccinia dill, Puccinia porii)</i> Purple blotch <i>(Alternaria porii)</i>	4 to 6 fl. oz. per acre
	<p><b>White rot:</b> For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz. TEBUCURE FUNGICIDE 3.6 per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl. oz./acre.</p> <p><b>Rust:</b> For the control of rust make foliar applications at the rate of 4 to 6 fl. oz. TEBUCURE FUNGICIDE 3.6 per acre per application. Repeat at 10- to 14-day intervals.</p> <p>Apply TEBUCURE FUNGICIDE 3.6 in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p><b>Notes:</b> Do not apply more than 32.5 fl. oz. TEBUCURE FUNGICIDE 3.6 per acre per season if an in-furrow treatment is made. If TEBUCURE FUNGICIDE 3.6 is not applied as an in-furrow treatment then do not apply more than 12 fl. oz. TEBUCURE FUNGICIDE 3.6 per acre per season as a foliar spray. Do not apply within 7 days of harvest.</p>	
<p><b>Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (RE1) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Grasses Grown for Seed	Rusts ( <i>Puccinia</i> spp)	4 to 8 fl. oz. per acre
	Apply the specified rate of TEBUCURE FUNGICIDE 3.6 as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications 14- to 16- day intervals. Under heavy disease pressure use 6 to 8 fl oz/acre and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
	Apply specified rate of TEBUCURE FUNGICIDE 3.6 when powdery mildew first appears on leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl. oz./acre and shorter spray intervals.	
<p><b>Comments:</b> Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.</p> <p>For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with TEBUCURE FUNGICIDE 3.6.</p> <p>A maximum of 16 fl. oz. (1 pint) may be applied per acre per crop season. TEBUCURE FUNGICIDE 3.6 may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots Welsh Onion	White rot ( <i>Sclerotium cepivorum</i> ) Suppression only	4 to 6 fl. oz. per acre
	Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> ) Purple blotch ( <i>Alternaria porii</i> )	
For control of diseases make foliar applications using an interval of 10- to 14-days. Apply TEBUCURE FUNGICIDE 3.6 in a protective spray schedule or when weather conditions are favorable for rust development.		
<b>Notes:</b> Do not apply more than 24 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per season. Do not apply within 7 days of harvest.		
<p><b>Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Hops</b>	Powdery mildew ( <i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i> )	4 to 8 fl. oz. per acre
	<b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. TEBUCURE FUNGICIDE 3.6 may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per crop season, Increase the spray volume and the application rate as vine growth increases during the season.	
<b>Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
<b>Leafy Brassica Greens</b> <b>Broccoli raab</b> Chinese cabbage (bok choy) Collards Kale Minima Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot ( <i>Cercospora brassicola</i> ) Powdery mildew ( <i>Erysiphe cruciferarum</i> ) Alternaria leaf spot ( <i>Alternaria brassicola</i> )	3 to 4 fl. oz. per acre
	<b>Notes:</b> Do not apply more than 16 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per season. Do not apply within 7 days of harvest. Do not apply more often than once every 10 days.	
<b>Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). <b>Restriction:</b> Application to turnip greens is limited to East of the Rockies. Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Garden beet, roots and tops (leaves)	Cercospora leaf spot ( <i>Cercospora beticola</i> )	3 to 7.2 fl. oz. per acre
	<b>Notes:</b> Make applications on 14-day intervals. Do not apply more than 28.8 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per season. Do not apply within 7 days of harvest.	
<p><b>Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval(REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Lychee	Anthrachnose ( <i>Colletotrichum gloeosporioides</i> )	4 to 6 fl. oz. per acre
	<b>Notes:</b> Begin first application of TEBUCURE FUNGICIDE 3.6 as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per season. TEBUCURE FUNGICIDE 3.6 can be applied up to and including the day of harvest PHI = 0 days).	
<p><b>Comments:</b> For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval(REI) = 2 days</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Okra	Cercospora leaf spot ( <i>Cercospora beticola</i> )	4 to 6 fl. oz. per acre
<p><b>Notes:</b> Apply specific dosage of TEBUCURE FUNGICIDE 3.6 in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per season.</p>		
<p><b>Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Peanut	<p><b>SOILBORNE:</b> Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)</p> <p><b>FOLIAR:</b> Early leaf spot Late leaf spot Leaf rust Web blotch (<i>Phoma</i>) Pepper spot (<i>Leptosphaerulina</i>)</p>	7.2 fl. oz. per acre
<p><b>FOUR-APPLICATION SPRAY PROGRAM:</b> Apply the specified rate in a preventative spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of TEBUCURE FUNGICIDE 3.6 to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6.</p> <p><b>LEAF SPOT ADVISORY SCHEDULE:</b> For control of soilborne diseases in an advisory schedule, apply TEBUCURE FUNGICIDE 3.6 in the first advisory spray in July and continue TEBUCURE FUNGICIDE 3.6 applications at 14-day intervals. Applications after August 15 should be tank-mixed with chlorothalonil for resistance management purposes.</p>		
<p><b>Use Instructions:</b> For optimum control of the specified soilborne diseases, four consecutive applications of TEBUCURE FUNGICIDE 3.6 must be made at 14-day intervals. A maximum of 28.8 fl. oz. of TEBUCURE FUNGICIDE 3.6 may be applied per crop season. TEBUCURE FUNGICIDE 3.6 may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas. TEBUCURE FUNGICIDE 3.6 is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Chlorothalonil may be tank-mixed at the rate of 12 oz of active ingredient with TEBUCURE FUNGICIDE 3.6 as a leaf spot resistance management strategy. A spray surfactant is not necessary when TEBUCURE FUNGICIDE 3.6 is tank-mixed with chlorothalonil. Mixing or alternating TEBUCURE FUNGICIDE 3.6 with other DMI fungicides may lead to resistance. TEBUCURE FUNGICIDE 3.6 must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i>. Drought conditions will decrease the effectiveness of TEBUCURE FUNGICIDE 3.6 against the root and pod rots.</p> <p>Use TEBUCURE FUNGICIDE 3.6 in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.</p> <p>Restricted-entry interval (REI) = 12 hours</p>		
<p><b>Timing of TEBUCURE FUNGICIDE 3.6</b> <b>Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot</b></p>		
<b>Spray Program</b>	<b>TEBUCURE FUNGICIDE 3.6 Application No.</b>	<b>Chlorothalonil Application No.</b>
7 applications	3, 4, 5 and 6	1, 2 and 7



CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Pecan	Brown leaf spot <i>(Sirosporium cliffusium)</i> Downy spot <i>(Mycosphaerella caryigena)</i> Liver spot <i>(Gnomonia caryae)</i> Scab <i>(Cladosporium caryigenum)</i> Vein spot <i>(Gnomonia ner<sup>v</sup> seda)</i> Zonate leaf spot <i>(Grovesinia pyramidalis)</i>	4 to 8 fl. oz. per acre
<p><b>Notes:</b> Apply TEBUCURE FUNGICIDE 3.6 in a preventative spray schedule beginning at early bud break (young leaves unfolding) and continue applications at 10- to 14-day intervals through the pollination period. TEBUCURE FUNGICIDE 3.6 should be applied at 4 fl. oz. per acre in a tank-mix with the recommended rate of Super-Tin<sup>®</sup> in cover sprays. Follow label directions for Super-Tin<sup>®</sup>. Do not add a surfactant to the spray solution when using Super-Tin<sup>®</sup>. Apply TEBUCURE FUNGICIDE 3.6 in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of TEBUCURE FUNGICIDE 3.6 to full-size mature trees and 4 to 6 fl. oz. per acre of TEBUCURE FUNGICIDE 3.6 to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of TEBUCURE FUNGICIDE 3.6 may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p>		
<p><b>Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy. Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Soybean	Rust ( <i>Phakopsora pachyrhizi</i> )  Powdery mildew ( <i>Microsphaera diffusa</i> )	3 to 4 fl. oz. per acre
<p><b>Use Instructions:</b> Apply TEBUCURE FUNGICIDE 3.6 as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label recommended rate of a spray surfactant must be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.</p> <p><b>Restrictions:</b> Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz. per acre per use season. Restricted-entry interval = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Sunflower	Rust ( <i>Puccinia helianthi</i> )	4 to 6 fl. oz. per acre
<p><b>Notes:</b> Apply specific dosage of TEBUCURE FUNGICIDE 3.6 at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per season or within 50 days of harvest.</p>		
<p><b>Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. Contact your state Extension Service or <b>Direct AG Source Winfield Solutions, LLC</b> representative for a list of approved surfactants. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot ( <i>Cercospora brassicicola</i> )	4 to 7.2 fl. oz. per acre
	<p><b>Notes:</b> Apply the specific dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. TEBUCURE FUNGICIDE 3.6 may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of TEBUCURE FUNGICIDE 3.6 per acre per crop season.</p>	
<p><b>Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6
Wheat	Rusts leaf, stem and stripe ( <i>Puccinia</i> spp.)	4 fl. oz. per acre
	Head blight or scab ( <i>Fusarium</i> spp.) - Suppression	
<p><b>Notes:</b> Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of TEBUCURE FUNGICIDE 3.6 may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with TEBUCURE FUNGICIDE 3.6. Apply TEBUCURE FUNGICIDE 3.6 in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><b>Application timing directions:</b>  <b>Rusts:</b> Apply TEBUCURE FUNGICIDE 3.6 at the earliest sign of rust pustules on foliage.  <b>Fusarium head blight:</b> Optimal timing of TEBUCURE FUNGICIDE 3.6 for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>		
<p><b>Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUCURE FUNGICIDE 3.6. TEBUCURE FUNGICIDE 3.6 must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUCURE FUNGICIDE 3.6 will be resistant to weathering. TEBUCURE FUNGICIDE 3.6 is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.</p>		

<b>SEED TREATMENT- Corn</b> (Sweet Corn, Field Corn, Field Corn Grown For Seed and Popcorn) For control of soilborne and seedborne Fusarium and soilborne and Seedborne head smut.		
<b>SEED LABELING:</b> To meet US Federal Seed Act requirements, all seed treated with TEBUCURE FUNGICIDE 3.6 must be labeled:  <p style="text-align: center;"><b>“TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES.”</b>  <b>“Treated with Tebuconazole.”</b>  <b>“Excess treated seed may be used for ethanol production</b>  <b>only if (1) by-products are not used for livestock feed and (2) no measurable residues</b>  <b>of pesticide remain in ethanol by-products that are used in agronomic practice.”</b></p>		
<b>USE PRECAUTION:</b> When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored with an EPA approved dye such as one of the dyes listed in 40 CFR Section 180.1001(c) and (d) to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.		
<b>DISEASE</b>	<b>RATE FI OZ/CWT</b>	<b>DIRECTIONS FOR USE</b>
<b>Soilborne and Seedborne Fusarium</b>	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with TEBUCURE FUNGICIDE 3.6. The length of control will vary depending on the rate used.
<b>Soilborne and Seedborne Head smut (<i>Sphacelotheca reiliana</i>)</b>	0.27 — 0.54	

<b>PLANT</b>	<b>DISEASE</b>	<b>RATE OF ORIUS 3.6 F</b>
<b>LEATHERLEAF FERN (FLORIDA ONLY)</b>	Anthracnose (suppression)	5 to 10 fl oz per acre
	<b>Notes:</b> Make the first application before anthracnose symptoms are present and continue at 12- to14-day intervals. <b>USE RESTRICTIONS:</b> A maximum of 5 pints of Tebucure Fungicide 3.6 may be applied per acre per year.	
<b>Comments:</b> Apply TEBUCURE FUNGICIDE 3.6 in a minimum of 5 gallons of spray solution per acre using ground equipment or chemigation. Restricted-entry interval (REI) = 12 hours. <b>USE LIMITATIONS:</b> TEBUCURE FUNGICIDE 3.6 can cause phytotoxicity to Leatherleaf Fern under certain environmental conditions. Applications in temperatures less than 70° F can cause phytotoxicity in the form of leaf burning and/or yellowing. Application followed by temperatures falling below 55°F can cause similar symptoms. Before using this product on Leatherleaf Fern, read the <b>LIMITATION OF WARRANTY AND LIABILITY WARRANTY DISCLAIMER and LIMITATION OF LIABILITY</b> section in its entirety.		

**OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

**Spray Drift Management:** For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.

Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground application when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperatures.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

#### **ROTATIONAL CROPS**

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through, spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below:

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling: Nonrefillable Container.** Do not reuse or refill container. Offer for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill, or incineration. Do not burn unless allowed by state and local authorities. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

**Containers 5 gallons or less:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once cleaned, offer for recycling if available.

**Containers larger than 5 gallons:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Once cleaned, offer for recycling or reconditioning if appropriate.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call  
CHEMTREC 1-800-424-9300**

### **IMPORTANT INFORMATION – READ BEFORE USING PRODUCT CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Direct AG Source, LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of Direct AG Source, LLC and Seller. To the fullest extent allowed by State law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Direct AG Source, LLC and Seller harmless for any claims relating to such factors.

Direct AG Source, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or Direct AG Source, LLC, and to the extent consistent with applicable law Buyer and User assume the risk of any such use. DIRECT AG SOURCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by law, Direct AG Source, LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF DIRECT AG SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE, AT THE ELECTION OF DIRECT AG SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT, OR COMPENSATION LIMITED TO DAMAGES NOT EXCEEDING THE FAIR MARKET PURCHASE PRICE, AND SHALL NOT INCLUDE INCIDENTAL OR CONSEQUENTIAL DAMAGES.**

Direct AG Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of Direct AG Source, LLC

#### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.**

#### LIMITATION OF LIABILITY

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.**

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

## Sublabel B: Turf and Ornamental Uses

**GROUP 3 FUNGICIDE**

# Tebucure Fungicide 3.6

**ACTIVE INGREDIENT:**

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol.....38.7%

**OTHER INGREDIENTS:**.....61.3%

**TOTAL** 100.0%

Contains 3.6 pounds tebuconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
<p><b>Note to Physician:</b> No specific antidote. Treat symptomatically.</p> <p><b>Symptoms of Poisoning:</b> The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 7:30 AM-3:30 PM PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, Contact 1-877-424-7452 for emergency medical treatment information.</p>	

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

**SEE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS, COMPLETE DIRECTIONS FOR USE, WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY.**

Manufactured for By:  
 Direct AG Source, LLC  
 30473 260<sup>th</sup> Street  
 Eldora, IA 50627  
 Winfield Solutions, LLC  
 P.O. Box 64589  
 St. Paul, MN 55164-0589

EPA Reg. No. 83222-19  
 EPA Establishment No.  
 Net Contents  
 Lot No.

1/1025/6



## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as bather laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Remove and wash contaminated clothing before reuse. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Groundwater Advisory:** Tebuconazole is known to leach through soil into the ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**Surface Water Advisory:** This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops.

PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Entry Restrictions for Non-WPS Uses:** Do not enter or allow others to enter the treated area until sprays have dried.

**Spray Volume:** For turf, apply Tebucure Fungicide 3.6 in 66-132 gallons of water per acre by ground sprayer. For ornamentals other than Leatherleaf Fern, use 50-300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at the time of application. For Leatherleaf Fern, apply Tebucure Fungicide 3.6 in a minimum of 5 gallons of finished spray per acre using ground equipment or chemigation. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

**Chemigation:** Apply Tebucure Fungicide 3.6 through irrigation equipment only to Leatherleaf Fern in Florida to suppress anthracnose. Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Contact State Extension Service specialist, equipment manufacturers or other experts if you have questions regarding calibration. Do not connect an irrigation systems (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjusts if the need arises.

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regular serves an average of at least

25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, back flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally dosed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

**Mixing:** Add labeled amount of Tebucure Fungicide 3.6 into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Tebucure Fungicide 3.6 should be thoroughly dispersed prior to the addition of other materials.

Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

**Compatibility:** To determine the compatibility of Tebucure Fungicide 3.6 with other products, use the following procedure: Pour the labeled proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be remixed readily, the mixture is considered physically compatible. For further information contact your local [Direct AG Source Winfield Solutions, LLC](#) representative.

#### **Resistance Management Statement**

Tebucure Fungicide 3.6 is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Tebucure Fungicide 3.6 and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistance isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of this product should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. [Direct AG Source Winfield Solutions, LLC](#) encourages responsible management to ensure effective long-term control of the fungal disease on this label.

## TURF AND ORNAMENTAL USES

### DISEASE CONTROL IN GOLF COURSE TURF

#### PRODUCT INFORMATION

For use on all Golf turf applications of cool season and warm season grasses (such as Bentgrasses, Bluegrasses, Fescues, Ryegrasses, St. Augustine grasses, and Zoysia) or their mixtures. Tebucure Fungicide 3.6 is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. **Note:** Bermudagrass can be sensitive to Tebucure Fungicide 3.6 under certain conditions. Do not apply consecutive applications during or just after dormancy break. Avoid applications when temperatures are expected to exceed 85 degrees F.

Tebucure Fungicide 3.6 can be used for the prevention and control of the diseases mentioned in table below. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Preventative treatments can be applied using 28 day intervals as indicated. When treating golf greens, always treat aprons and approaches. Spray uniformly over the area to be treated with properly calibrated equipment. Apply the specified amount of Tebucure Fungicide 3.6 Fungicide in sufficient water for thorough coverage. A volume of 66 – 132 gallons per acre (1.5 – 3.0 gallons per 1,000 sq ft) is recommended. Apply using properly calibrated low volume, hand held, mechanical or motorized ground broadcast equipment. Application to small areas may be made with low-pressure handwand or backpack equipment.

Depending on the disease, Tebucure Fungicide 3.6 should be watered into the crown and active root zone for best results. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. For best results use spray mixture the same day it is prepared.

#### USE RESTRICTIONS

- For use on golf course turf only.
- Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high school), campgrounds, churches, and theme parks.
- Not for residential use.
- Not for use on turf being grown for sale or commercial use as sod.
- Do not use clippings for animal feed.
- Do not exceed 3.6 oz of Tebucure Fungicide 3.6 Fungicide per 1,000 sq ft per year.
- Do not apply more than 6 applications per year in all states except New York, and do not apply more than 3 applications per year in New York State.

DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6 FL OZ/1000 SQ FT	REMARKS
<b>Dollar Spot</b> <i>(Sclerotinia homoeocarpa)</i> <b>Copper Spot</b> <i>(Gloeocercospora sorghi)</i> <b>Powdery Mildew</b> <i>(Erysiphe graminis)</i> <b>Corticium Red Thread</b> <i>(Laetisaria fuciformis)</i> <b>Rusts</b> <i>(Puccinia spp.)</i>	0.6	For prevention, begin applications when conditions are favorable for disease development. Do not make two consecutive applications of Tebucure Fungicide 3.6. Alternate with another fungicide with a different mode of action. A second application may be made after 28 days.
<b>Brown Patch/Rhizoctonia</b> <b>Blight, Large Patch</b> <i>(Rhizoctonia solani)</i> <b>Brown Ring Patch</b> <i>(R. circinata)</i>	0.6	
<b>Anthracnose -Basal and Foliar</b> <i>(Colletotrichum cereale)</i> <b>Red Thread</b> <i>(Laetisaria fuciformis)</i> <b>Pink Patch</b> <i>(Limonomyces rosipellis)</i>	0.6	
<b>Bermuda Grass decline</b> <i>(Gaeumannomyces graminis var. graminis)</i>	0.6	Immediately after fungicide is applied, the area should be irrigated with sufficient water to move the active ingredient down into the crown and root zone of the turf. The amount of water is dependent on the depth of the root zone. For prevention, begin applications two or four weeks prior to the historical appearance of disease symptoms. Initiate cultural control practices at the same time the fungicide is applied. Refer to your local County Extension Service for this information. Apply subsequent applications at 28 day intervals.
<b>Take All Patch</b> <i>(Gaeumannomyces graminis)</i>	0.6	For prevention, apply in the fall when soil temperature reaches 55-65° F and again in the spring under similar soil temperature conditions. Applications in both fall and spring may be necessary. Immediately after fungicide is applied, the area should be irrigated with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6 FL OZ/1000 SQ FT	REMARKS
<b>Gray Leaf Spot</b> <i>(Pyricularia grisea)</i>	0.6	Apply when conditions are favorable for disease development at 28 day intervals. If using 0.6 oz/1000 sq ft, or under conditions favoring moderate to heavy disease pressure, Tebucure Fungicide 3.6 can be tank mixed with a registered contact fungicide at label rate
<b>Stripe Smut</b> <i>(Ustilago striiformis)</i>	0.6	Make a single application to historical disease areas in spring as grass growth begins.
<b>Spring Dead Spot</b> <i>(Leptosphaeria korrea, L. narmari, Ophiosphaerella herpotricha, Gaeumannomyces graminis)</i> <b>Necrotic Ring Spot</b> <i>(Leptosphaeria korrea)</i>	0.6	For prevention, apply in fall when soil temperature reaches 65° F and again in spring under similar soil temp conditions or after dormancy break. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.
<b>Fusarium Patch</b> <i>(Fusarium roseum)</i>	0.6	Apply first application in mid-June or 28 days prior to time this blight normally becomes evident. Make applications at no less than 28 days intervals.
<b>Summer Patch</b> <i>(Magnaporthe poae)</i>	0.6	Apply beginning in the spring. Do not make two consecutive applications of Tebucure Fungicide 3.6. Alternate with another fungicide with a different mode of action. Second and third applications may be made at 28 day intervals. See local university recommendations for suggested timing. Immediately after fungicide is applied, irrigate the area with sufficient water to move the active ingredient down into the crown and active root zone of the turf. The amount of water is dependent on the depth of the root zone.

DISEASE	RATE OF TEBUCURE FUNGICIDE 3.6 FL OZ/1000 SQ FT	REMARKS
<b>Zoysia Patch, Large Patch of zoysia</b> <i>(Rhizoctonia solani)</i>	0.6	Make first application in early fall (mid-September to mid-October) prior to development of disease symptoms. A second application in early spring may be necessary in areas where disease pressure is known to be heavy.
<b>Gray Snow Mold/Typhula Blight</b> <i>(Typhula incarnate)</i>  <b>Pink Snow Mold/Microdochium Patch</b> <i>(Microdochium nivalis)</i>	0.6	Apply in the fall, before anticipated turf dormancy and before first snow cover. If turf breaks dormancy during winter months, a second application may be made. Do not apply over snow cover, or when turf is dormant. It is recommended that Tebucure Fungicide 3.6 be tank-mixed with other registered snow mold products for best season long results.
<b>COMMENTS:</b> Apply the specified amount of Tebucure Fungicide 3.6 in 1.5 to 3.0 gallons of water per 1000 sq ft. Make all applications after mowing and allow foliage to dry thoroughly before irrigation. Do not use clippings for animal feed. Do not exceed 3.6 fl oz of Tebucure Fungicide 3.6 per 1000 sq ft per year. Do not exceed 6 applications per year.		

**DISEASE CONTROL IN FIELD, NURSERY AND CONTAINER ORNAMENTALS AND COMMERCIAL and RESIDENTIAL LANDSCAPES**

**PRODUCT INFORMATION:** For use on ornamental plants only; not for woodlands or forest management.

Tebucure Fungicide 3.6 can be used in a preventative and curative disease control program for the listed plant types and disease in the table below. Optimum disease management is obtained when Tebucure Fungicide 3.6 is used in conjunction with sound disease management practices.

Apply material with properly calibrated hand held, mechanical or motorized spray equipment. Begin applications when disease first appears and repeat at 14-21 day intervals during the growing season. Use the shortest interval when conditions are unusually favorable for the development of disease. For hand held, mechanical, or motorized applications, mix as directed below and apply as a full coverage spray to drip for the prevention and control of the diseases listed below. Choose a finished spray volume appropriate for the size of the plants and amount of foliage, which will provide thorough coverage throughout the canopy. Allow sprays to dry before overhead irrigation is applied.

**USE RESTRICTIONS**

- Intended for use only by professional applicators.
- Apply Tebucure Fungicide 3.6 at rates of 4-10 fl oz per acre in 100 gallons of water. Spray volume may range from 50 up to 300 gallons of finished spray per acre depending upon equipment, plant species and plant growth stage at time of application.
- Do not apply more than 30 fl oz per acre in a single application.
- Do not apply more than 0.94 gallons (120 fl oz) of Tebucure Fungicide 3.6 (equal to 3.38 lbs of tebuconazole) per acre per year.
- Do not make more than 4 applications per year at highest rate.
- Do not apply to bearing fruit trees or vegetables.

**NOTE:** The **DIRECTIONS FOR USE** section of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on African Violets, Begonias, Boston Fern, and Geraniums.

**ORNAMENTALS DISEASE CONTROL**

PLANTS	DISEASE	APPLICATION	
		TO PREVENT DISEASE	TO TREAT DISEASE
Roses	Black Spot Powdery Mildew Rust	Apply every 14-21 days during the growing season, starting when leaves first appear.	Apply every 14 days for a total of 3 applications beginning at the first sign of disease.
Flowers	Leaf Spot Powdery Mildew Rust Southern Blight	Apply at least 3 times per year, 14-21 days apart, beginning with spring bud break.	
Crabapples (Ornamental), Dogwoods and other Landscape (ornamental) Trees	Anthrachnose Leaf Spot Powdery Mildew Rust Scab	Rotation or tank-mixing with barrier protectant fungicides is recommended for resistance management.	
Azaleas, Camellias, Rhododendrons and other Landscape (Ornamental) Shrubs	Anthrachnose Black Spot Leaf Spot Petal Blight	Petal Blight: Apply 2-3 times per week into the flowers as they open and develop color.	
Ground Covers and Vines	Powdery Mildew Rust Southern Blight		

For small plantings, add 1 teaspoon of Tebucure Fungicide 3.6 to 2.5 gallons of water.

**Pump Style Sprayers**

1. Add the appropriate amounts of concentrate and water to the sprayer tank.
2. Close the sprayer, shake well and pressurize
3. Adjust nozzle to a coarse spray pattern and apply.
4. Occasionally repressurize the sprayer, if needed, to maintain a good spray pattern.

PLANT	DISEASE	RATE OF ORIUS 3.6 F
LEATHERLEAF FERN (FLORIDA ONLY)	Anthrachnose (suppression)	5 to 10 fl oz per acre
	<b>Notes:</b> Make the first application before anthracnose symptoms are present and continue at 12- to 14-day intervals. <b>USE RESTRICTIONS:</b> A maximum of 5 pints of Tebucure Fungicide 3.6 may be applied per acre per year.	
<b>Comments:</b> Apply Tebucure Fungicide 3.6 in a minimum of 5 gallons of spray solution per acre using ground equipment or chemigation. <b>USE LIMITATIONS:</b> Tebucure Fungicide 3.6 can cause phytotoxicity to Leatherleaf Fern under certain environmental conditions. Applications in temperatures less than 70°F can cause phytotoxicity in the form of leaf burning and/or yellowing. Application followed by temperatures falling below 55°F can cause similar symptoms. Before using this product on Leatherleaf Fern, read the <b>LIMITATION OF WARRANTY AND LIABILITY WARRANTY DISCLAIMER and LIMITATION OF LIABILITY</b> section in its entirety		



**OBSERVE THE FOLLOWING RESTRICTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

- Apply only during alternate years in fields adjacent to aquatic areas listed above.
- Do not apply by ground or air within 100 feet of aquatic areas listed above.
- Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

**Spray Drift Management:** For aerial applications, mount the spray boom on the aircraft so as to minimize drift caused by wing tip vortices. Use the minimum practical boom length, and do not exceed 75% of the wing span or rotor diameter. Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Release the spray at the lowest possible height consistent with good pest control and flight safety. Avoid applications more than 10 feet above the crop canopy. Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area. Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature. Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If container is leaking, invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through, spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below:

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling: Nonrefillable Container.** Do not reuse or refill container. Offer for recycling or reconditioning if available, or puncture and dispose of in a sanitary landfill, or incineration. Do not burn unless allowed by state and local authorities. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

**Containers 5 gallons or less:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Once cleaned, offer for recycling if available.

**Containers larger than 5 gallons:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Once cleaned, offer for recycling or reconditioning if appropriate.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call  
CHEMTREC 1-800-424-9300**

**IMPORTANT INFORMATION - READ BEFORE USING PRODUCT  
CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Direct AG Source, LLC or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of Direct AG Source, LLC and Seller. To the fullest extent allowed by State law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Direct AG Source, LLC and Seller harmless for any claims relating to such factors.

Direct AG Source, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or Direct AG Source, LLC, and to the extent consistent with applicable law Buyer and User assume the risk of any such use. **DIRECT AG SOURCE, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, Direct AG Source, LLC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF DIRECT AG SOURCE, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE, AT THE ELECTION OF DIRECT AG SOURCE, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT, OR COMPENSATION LIMITED TO DAMAGES NOT EXCEEDING THE FAIR MARKET PURCHASE PRICE, AND SHALL NOT INCLUDE INCIDENTAL OR CONSEQUENTIAL DAMAGES.**

Direct AG Source, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of Direct AG Source, LLC

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.**

#### LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.