

84930-11

3/13/2014

1/32

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460



OFFICE OF
CHEMICAL SAFETY AND
POLLUTION PREVENTION

ARCANA, LLC
c/o Scott Baker
Lighthouse Product Services
1966 W 15th Street Suite 6
Loveland, CO 80538

MAR 13 2014

Subject: Amended label adding pollinator protection language
Product Name: ARC-IMIDA 2# Insecticide
EPA Reg. No. 84930-11
EPA Decision No. 488611
Submission dated March 7, 2014

Dear Mr. Baker:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act is acceptable. A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release the product for shipment with the new labeling. See 40 CFR 156.10(a)(6).

Under 40 CFR 152.130(d), EPA may establish dates by which all product distributed or sold by the registrant must bear revised labeling. The following paragraphs set forth the schedule for ensuring that that your product bears revised labeling within a reasonable time period.

- Any product released for shipment after 2/28/14 must bear the new label.

If these conditions are not complied with, EPA will take appropriate action against this registration. If you have any questions please contact Dr. Debra Rate at 703-306-0309 or rate.debra@epa.gov.

Regards,

A handwritten signature in cursive script that reads "Venus Eagle".

Venus Eagle, Product Manager (01)
Insecticide-Rodenticide Branch
Registration Division (7505P)

2/32

ACCEPTED
MAR 13 2014

Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:

ARC-IMIDA 2# Insecticide

EPA. Reg. No: 84930-11

For use in pest management and suppression of insect vectored diseases and maintenance of plant health for soil application

ACTIVE INGREDIENT

Imidacloprid -1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-Imidazolidinimine*	21.4%
Other ingredients	78.6%
Total	100.0%

*Contains 2 pounds of imidacloprid per gallon

KEEP OUT OF REACH OF CHILDREN CAUTION

EPA Reg. No. 84930-11

EPA Est. No. XXXXX-XX-XXX

Net Contents:

Manufactured For:
ARCANA, LLC
1966 W 15th Street Suite 6
Loveland, CO 80538

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>HOT LINE NUMBER: Have the product container or label with you when calling a poison control center, doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC Web site: www.npic.orst.edu). Outside of these times call</p>	

01/12/2014
3:00:21 PM
01/12/2014 3:00:21 PM
01/12/2014 3:00:21 PM

call your poison control center at 1-800-222-1222.
NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

Applicators and other handlers must wear:

- Long-sleeved shirts and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water: Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

- When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to inertial areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where

soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

TAKE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS; MARSHES OR NATURAL PONDS; ESTUARIES, AND COMMERCIAL FISH FARM PONDS.

Spray Drift Management

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150 - 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. Do not apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions

Do not make ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Mixing and Loading Requirements

To avoid potential contamination of groundwater, the use of a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment is recommended. If a containment pad is not used,

maintain a minimum distance of 25 feet between mixing and loading area and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes, or field drains.

No-Spray Zone Requirements for Soil Applications

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Run-off Management

Do not cultivate within 10 feet of the aquatic areas to allow growth of a vegetative filter strip. When using ARC-IMIDA 2# on erodible soils, Best Management Practices for minimizing run-off should be employed. Consult your local Natural Resources Conservation Service for recommendations in your use area.

Endangered Species Notices

Under the Endangered Species Act it is a federal offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

Resistance Management

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

ARC-IMIDA 2# Insecticide contains a Group 4A Insecticide. Insecticide biotypes with acquired or inherent resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

The active ingredient in ARC-IMIDA 2# belongs to the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to Imidacloprid 2 F. In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season: 1) only a single soil application of ARC-IMIDA 2# be made; 2) foliar applications of products from this same class not be made following a long residual soil application of ARC-IMIDA 2# or other neonicotinoid products.

Other Group 4A neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Gaucho, Intruder, Leverage, Provado, and Trimax.

Other Group 4A neonicotinoid products used as soil treatments include: Platinum.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations.

6/32

Also, for more information on Insect Resistance Management (IRM) visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

DO NOT apply this product, by any application method, to linden, basswood, or *Tilia* species trees in Oregon.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treatment areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any water material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

To Confine Spills: Handle and open container in a manner as to prevent spillage. If the container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile and dispose of as directed for pesticides below. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

DISPOSAL STATEMENTS:

Nonrefillable container: Do not reuse this container to hold materials other than pesticides or dilute pesticides (rinsate). After emptying and cleaning, it may be allowable to temporarily hold rinsate or other pesticide-related materials in the container. Contact your state regulatory agency to determine allowable practices in your state. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact The Agricultural Container Recycling Council (ACRC) at www.acrecycle.org.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For packages up to 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For packages greater than 5 gallons and less than 56 gallons: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to

drip.

For packages greater than 56 gallons: To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

For refillable containers: Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

APPLICATION INSTRUCTIONS

For best control, direct applications of ARC-IMIDA 2# Insecticide into the seed or root-zone of crop. ARC-IMIDA 2# may be applied with ground or chemigation application. Do not apply by air. Use only by broadcast or foliar applications on seedling flats or trays, or where product is meant to be washed from foliage to soil before drying on foliage within the planthouse/greenhouse.

For best result, apply Imidacloprid 2 F to the root-zone of plants. Since ARC-IMIDA 2# is continuously taken into the roots over a long period of time, the earlier ARC-IMIDA 2# is available to a developing plant, the earlier the protection begins. The systemic nature of ARC-IMIDA 2# allows movement from roots through the xylem tissue to all vegetative parts of the plant, which results in extended residual activity of ARC-IMIDA 2#, the control of insects and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate of ARC-IMIDA 2# applied affects the length of the plant protection period; use the higher rates within the specified rate range when infestations occur later in crop development, or where pest pressure is continuous. ARC-IMIDA 2# will generally not control insects infesting flowers, blooms or fruit, and additional crop protection may be required for insects feeding in, or on these plant parts and for insects not listed in the crop-specific, pests controlled sections of this label.

Suppression (less than complete control of certain diseases and insect pests including reduced feeding) may also be caused by an ARC-IMIDA 2# application. In such cases, pests/diseases may require supplemental control measures.

The use of ARC-IMIDA 2# on crops grown for production of true seeds is not intended for private or commercial planting unless allowed by state specific 24(c) labeling. Always take care to minimize exposure of ARC-IMIDA 2# to honey bees and other pollinators. Additional information on ARC-IMIDA 2# uses for these

crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants, or local representative.

Make applications only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool or other soil-less media or plants growing hydroponically. Mix ARC-IMIDA 2# with water or other appropriate diluent before application. Agitate the ARC-IMIDA 2# and water suspension to avoid settling.

Do not apply more than 0.50 lbs active ingredient per acre, per year, regardless of formulation or method of application, unless specified within a crop-specific applications section for a given crop.

Mixing Instructions

To prepare the application mixture, first add a portion of the required amount of water to the tank and then add ARC-IMIDA 2# while agitation is underway. Add the remainder of water needed. Agitate during both mixing and application. ARC-IMIDA 2# may also be used with other pesticides and/or fertilizer solutions. Please see compatibility note below. When tank mixtures of ARC-IMIDA 2# and other pesticides are involved, prepare the tank mixture as recommended above and follow suggested Mixing Order below.

Mixing Order

When making pesticide mixtures, add ARC-IMIDA 2# and other wettable powders or wettable granules first, followed by flowable (suspension concentrate) products, then emulsifiable concentrates last. Agitate as each ingredient is added. Do not add an ingredient until previous one is thoroughly mixed. If a fertilizer solution is added, a fertilizer/pesticide compatibility agent may be needed. To ensure a uniform spray mixture, continuous agitation is necessary during both mixing and application.

Compatibility Note

Test the compatibility of any intended mixture before adding ARC-IMIDA 2# to the spray or mix tank. To do this, add proportionate amounts of each ingredient in the appropriate order, to a suitable size jar, cap, shake the mixture for 5 minutes, and let set for 5 minutes. If the mixing is poor, or there is formation of precipitates that do not readily redisperse, then the blend is incompatible and should not be used. For further information, contact your local United Phosphorus, Inc. representative.

CHEMIGATION - DIRECTIONS FOR USE

Types of Irrigation Systems

Chemigation applications of ARC-IMIDA 2# may only be made to crops through chemigation systems as specified in crop-specific application sections and only through low-pressure systems unless specifically recommended for a given crop. Do not apply Imicloprid 2 F through any other type of irrigation system.

Uniform Water Distribution and System Calibration

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift

Do not apply when wind speed drift beyond the area intended for treatment.

Required System Safety Devices

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. System must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water From Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option of the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to the top or overflow rim of the reservoir tank of at least twice the inside

diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROTATIONAL CROPS*

Treated areas may be replanted with any crop specified on an imidacloprid label, any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops with no established tolerances for the active ingredient, observe a 12 month plant-back interval.

Immediate Plant-Back All crops on this label plus the following crops not on this label: barley, canola, corn (field, pop & sweet), rapeseed, sorghum, soybean, and wheat

30-Day Plant-Back Cereals (including buckwheat, millet, oats, rice, rye, and triticale), safflower

10-Month Plant-Back Onion and bulb vegetables

12-Month Plant-Back All other crops

* Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

FIELD CROPS

COTTON

PESTS	RATE	INSTRUCTIONS
Cotton Aphid Plant Bugs Thrips Whiteflies	1.3 fl oz/1000 row feet 17.0 - 21.1 fl oz/A (depending on row spacing)	Use one of the following methods to apply the specified dosage of Imidacloprid 2 F: 1. In-furrow spray during planting directed on or below seed; 2. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; 3. Chemigation into root-zone through low-pressure drip or trickle irrigation. RESTRICTIONS: Do not apply more than 21.1 fl oz (0.33 lb ai) ARC-IMIDA 2# per acre per year. No matter which formulation or method of

		<p>application is used, do not apply more than 0.5 lb active ingredient of ARC-IMIDA 2#, Provado, Trimax or Leverage per acre per season, including seed treatment such as Gaucho, soil <u>and</u> foliar uses. Do not graze treated fields after any application of ARC-IMIDA 2#.</p> <p>Please note the Resistance Management section of this label.</p>
--	--	--

POTATOES

PESTS	RATE	INSTRUCTIONS
<p>Controlled Aphids Colorado Potato Beetle Flea Beetles Leafhoppers Potato Psyllid</p> <p>Suppressed Symptoms of: Potato leaf roll Virus (PLRV) Potato yellows Net necrosis (PLRV) Wireworms (with in-furrow spray at planting)</p>	<p>0.9 - 1.3 fl oz/1000 row feet</p> <p>13.0 - 20.0 fl oz/A</p>	<p>Use one of the following methods to apply the specified dosage of Imidacloprid 2 :</p> <ol style="list-style-type: none"> 1. In-furrow spray during planting, directed onto seed pieces or seed potatoes; 2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil; 3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil; 4. Narrow band directly below the eventual seed row in a bedding operation no more than 7 days before planting. Place ARC-IMIDA 2# applications below soil surface and in contact with seed piece or within root-zone. For potatoes grown on highly permeable soils with a shallow water table, at-plant applications of ARC-IMIDA 2# may be made in 2 to 4 inch band (width of planter show opening) and completely covered. <p>RESTRICTIONS: Do not apply more than 20.0 fl oz (0.31 lb ai) ARC-IMIDA 2# per acre per year.</p>

POTATOES

(Seed Piece Treatment)

This use not allowed in California unless other directed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
<p>Controlled Aphids Colorado potato</p>	<p>0.4 - 0.8 fl oz/100 lbs seed</p>	<p>Apply specified rate as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water or less.</p>

beetle Flea beetles Leafhoppers Potato psyllid Wireworms (seed-piece protection)	8.0 – 16.0 fl oz/A (based on a seeding rate of 2000 lbs/A)	to 1 part ARC-IMIDA 2#. Agitate or stir spray solution as necessary. Apply fungicidal or inert absorbent dusts after ARC-IMIDA 2# is applied. Make applications only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Avoid extended exposure of seed-pieces to sunlight by planting as soon as possible after treating and in accordance with the recommendation of your local Extension specialist. Consult your local Arcana, LLC representative or crop protection product dealer for information relevant to your area
Suppressed Symptoms of: Potato leaf roll virus (PRLV) Potato yellows Net necrosis (PLRV)	0.8 fl oz/100 lbs seed 16.0 fl oz/A (based on a seeding rate of 2000 lbs/A)	RESTRICTIONS: Do not use treated seed-pieces for food, feed, or fodder. Do not apply any additional application of ARC-IMIDA 2# (in-furrow), Gaucho, Leverage, or Provado following an ARC-IMIDA 2# seed-piece treatment. Do not apply more than 20.0 fl oz (0.31 lb ai) ARC-IMIDA 2# per acre per year.

TOBACCO

PESTS	RATE	INSTRUCTIONS
Aphids Flea beetles	1.0 fl oz/1000 plants as seedling tray drench OR 1.4 fl oz/1000 plants 9in-furrow transplant water)	Use one of the following methods to apply specified dosage of Imidacloprid 2 : Uniform broadcast foliar spray to seedlings in trays (tray drench) no more than 7 days before transplanting, followed immediately by overhead irrigation to wash ARC-IMIDA 2# from foliage into potting media. Failure to wash the product from foliage may reduce pest control. Handle transplants carefully during setting to avoid dislodging treated potting media from roots. In-furrow spray or transplant-water drench during setting. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
Mole crickets Whiteflies Wireworms	1.4 – 2.8 oz/1000 plants as seedling tray	Important Note: Proper tray drench applications of ARC-IMIDA 2# have been shown to be the most effective method of

	drench OR 1.8 – 2.8 fl oz/1000 plants (in-furrow transplant water)	application. However, the specified rate of ARC-IMIDA 2# may be applied as combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of midacloprid 2 F into the plant and a delay in control.
Suppressed Cutworms Symptoms of: Tomato spotted wilt virus (TSWV)	1.4 – 2.8 oz/1000 plants as seedling tray drench OR 1.8 – 2.8 fl oz/1000 plants (in-furrow transplant water)	RESTRICTIONS: Do not apply more than 32.0 fl oz (0.50 lb ai) ARC-IMIDA 2# per acre per year. Pre-Harvest Interval (PHI): 14 days

VEGETABLE AND SMALL FRUIT CROPS

CUCURBIT VEGETABLES

Including: Chayote(fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melon* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, and Winter melon), Pumpkin, Squash, (includes summer squash types such as: butternut squash calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash, and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citullus lanatus*).

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

Not for use in California unless specifically allowed by state specific 24(c) labeling.

Field Applications

PESTS	RATE	INSTRUCTIONS
Control: Aphids Cucumber beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies	16.0 – 24.0 fl oz/A	Use one of the following methods to apply the specified dosage of ARC-IMIDA 2#: 1. Chemigation into root-zone by means of low pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application;
Suppression: Bacterial wilt (as vectored by various		

<p>cucumber beetles) Leaf silvering from whitefly feeding</p>		<p>4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. Incorporate product into the root zone.</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai) ARC-IMIDA 2# per acre, per crop season.</p>
---	--	---

Planthouse Applications

PESTS	RATE	INSTRUCTIONS
<p>Aphids Whiteflies</p>	<p>0.1 fl oz/1000 plants</p>	<p>Use one of the following methods to apply the specified dosage to seedling in trays in the planthouse. Target soil media (tray drench), not more than 7 days prior to transplanting.</p> <ol style="list-style-type: none"> 1. Uniform, broadcast high volume foliar spray, followed by sufficient overhead irrigation to wash ARC-IMIDA 2# from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash product from foliage may reduce pest control; 2. Injection into overhead irrigation system, using sufficient volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray. <p>Applications made in the planthouse will only provide short-term protection and should not substitute for a field application. Make an additional field/soil application within 2 weeks after transplanting to provide continuous protection. Applications of higher rates or increased number of application in planthouse may result in significant plant injury. Handle transplant carefully during setting to avoid dislodging treated potting media from roots.</p> <p>Important Note: Not all varieties of cucurbit vegetables have been tested for tolerance to ARC-IMIDA 2# applied to seedling flats. It is therefore recommended to treat a small number of plants and confirm tolerance for 7 days before treating entire planthouse.</p>

		<p>RESTRICTIONS: Do not apply more than 0.1 fl oz (0.0156 lb ai)/1000 plants ARC-IMIDA 2# for planthouse applications. Do not apply more than once as a planthouse application.</p>
--	--	--

GREENHOUSE VEGETABLES

(Mature plants in production greenhouses)

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

Cucumber, Tomato, only

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	1.4 fl oz/1000 plants	<p>Apply specified dosage in at least 16 gallons of water for tomatoes and 21 gallons of water for cucumbers. Use soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Make applications only to plants grown in field-type soils, potting media, or mixtures thereof. Do not apply to plants grown in non-soil medias such as perlite, vermiculite, rock wool or other soil-less media, or plants growing hydroponically. Do not apply to immature plants since phytotoxicity may occur.</p> <p>Apply when infestation pressure surpasses threshold and beneficial are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficial (<i>Orius</i> sp.) can occur when ARC-IMIDA 2# is applied.</p> <p>Many varieties of vegetables have been tested for tolerance to ARC-IMIDA 2# and show good safety. However, certain varieties may show more sensitivity to ARC-IMIDA 2#. Treatment of a few plants is recommended before treating a whole greenhouse.</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 0 days Do not apply ARC-IMIDA 2# more than once per crop season.</p>

FRUITING VEGETABLES

Including: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet), Tomato, Pepinos, Tomatillo

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

Field Application

PESTS	RATE	INSTRUCTIONS
<p>Controlled: Aphids Colorado potato beetle Flea beetles Leafhoppers Thrips (foliage-feeding thrips, only) Whiteflies</p> <p>Suppressed: Symptoms of: Tomato mottle virus Tomato spotted wilt virus Tomato yellow leaf curl virus</p>	<p>Okra and Pepper 16.0 – 32.0 fl oz/A</p> <p>Other Crops 16.0 – 24.0 fl oz/A</p>	<p>Use one of the following methods to apply specified dosage of ARC-IMIDA 2#:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. ARC-IMIDA 2# must be incorporated into root-zone. <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per crop season to pepper and okra; Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per crop season to other fruiting vegetables.</p>

Planthouse Application

PESTS	RATE	INSTRUCTIONS
<p>Aphids Whiteflies</p>	<p>0.1 fl oz/1000 plants</p>	<p>Use one of the following methods to apply the specified dosage to seedlings in trays in the planthouse. Target soil media (tray drench), not more than 7 days prior to transplanting.</p> <ol style="list-style-type: none"> 1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash ARC-IMIDA 2# from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash the product from

		<p>foliage may result in reduced pest control.</p> <p>2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray.</p> <p>Applications made in the planthouse will only provide short-term protection and should not substitute for a field application. Make an additional field/soil application within 2 weeks after transplanting to provide continuous protection. Applications of higher rates or increased number of application in planthouse may result in significant plant injury. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.</p> <p>Important Note: Not all varieties of fruiting vegetables have been tested for tolerance to ARC-IMIDA 2# applied to seedling flats. It is therefore recommended to treat a small number of plants and confirm tolerance for 7 day prior to treating entire planthouse.</p> <p>RESTRICTIONS: Do not apply more than 0.1 fl oz (0.0156 lb ai)/1000 plants ARC-IMIDA 2# for planthouse applications.</p> <p>Do not apply more than once as a planthouse application.</p>
--	--	---

HEAD AND STEM BRASSICA VEGETABLES

Including: Broccoli, Broccoli raab (rapini), Brussels sprouts, cabbage, Cauliflower, Cavalo, broccoli, Chinese (gai lon) broccoli, Chinese (box choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard Spinach, Rape greens, Turnip tops (leaves).

LEAFY GREENS VEGETABLES

Including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland,) yellow rocket, winter cress), Dandelion, dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach), Watercress (commercial production only, application must not be made to native cress growing in streams or other bodies of water), Watercress (upland).

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	10.0 – 24.0 fl oz/36 inch rows	<p>Use one of the following methods to apply the specified amount of ARC-IMIDA 2#:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow sprayer directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting. 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. ARC-IMIDA 2# must be incorporated into root-zone. <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per crop season.</p>

LEAFY PETIOLE VEGETABLES¹

Including: Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss Chard.

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling

PESTS	RATE	INSTRUCTIONS
Aphids Leafhoppers Whiteflies	10.0 – 24.0 fl oz/A	<p>Apply specified dosage of ARC-IMIDA 2# using one of the following methods:</p> <ol style="list-style-type: none"> 1. Chemigation into root zone by means of low-pressure drip, trickle, micro-sprinkler, or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench.

		<p>6. Subsurface side-dress on both sides of each row. ARC-IMIDA 2# must be incorporated into root-zone.</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 45 days</p> <p>Do not apply more than 24.0 fl oz (0.38 lb ai)/A Imidacloprid 2 F per crop season.</p>
--	--	--

LEGUME VEGETABLES except soybean, dry

Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean. Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin) Bean (*Phsaeolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean) Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea cathang, Chinese longbean cowpeas, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean) Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea) Other Beans and Peas (Broad Bean (fava), Chickpea (garbonzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean)

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
<p>Controlled: Aphids Leafhoppers Thrips (foliage-feeding thrips, only) Whiteflies</p> <p>Suppressed: Symptoms of: Bean common mosaic virus (BCMV) Bean golden mosaic virus (BGMV) Beet curly top hybrigeminivirus (BCTV)</p>	<p>16.0 – 24.0 fl oz/A</p>	<p>Apply specified dosage of ARC-IMIDA 2# using one of the following methods:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray at planting directed on or below seed; 3. In a narrow (2" or less) surface band over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation with 24 hours following application; 4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; 5. As a post-seeding drench, transplant drench, or hill drench. <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days</p> <p>Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per crop season.</p>

ROOT VEGETABLES

Including: (Beet (garden*), Burdock (edible*), Carrot*, Celeriac*, Chervil (turnip-rooted*), Chicory *, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip*, Radish*, Oriental radish (diakon*) Rutabaga*), Salsify (oyster plant), Salsify (black*), Salsify (Spanish*), Skirret and turnip*.

* Tops and Greens from these crops may be used for food or feed.

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling

PESTS	RATE	INSTRUCTIONS
Aphids Flea beetles Leafhoppers Whiteflies	0.7 – 1.7 fl oz/1000 row feet 10.0 – 24.0 fl oz/A	Apply specified dosage of ARC-IMIDA 2# using one of the following methods: 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray (rate specified per 1000 row-feet) or, shaken-in 1 to 2" below seed depth during planting; 3. In a narrow (2" or less) band directly (1 to 2") below the eventual seed row in a bedding operation 14 or fewer days before planting. Important Note: The rate used affects the length of control. Higher rates should be used where infestations occur later in crop development, or where pest pressure is continuous. Rates less than 0.7 fluid ounces/1000 row-feet will not provide adequate residual pest control. ARC-IMIDA 2# treated crops grown on very high organic matter soils (muck) may also require additional pest management control. RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per crop season. Do not apply more than once per crop season.

TUBEROUS AND CORM VEGETABLES

Including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet*) Chayote (root), Chufa, Dasheen (taro*), Ginger, Leren, Sweetpotato, Tanier (cocoyam*), Tumeric, Yam bean (jicama; manioc pea), Yam (true*)

(For applications on potato see Field Crops section)

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

* Tops and greens from these crops may be used for food or feed.

PESTS	RATE	INSTRUCTIONS
Aphids Flea beetles Leafhoppers Whiteflies	0.7 – 1.7 fl oz/1000 row feet	Use one of the following methods to apply specified dosage of ARC-IMIDA 2#: <ol style="list-style-type: none"> 1. In-furrow spray (rate specified per 1000 row-feet) over planting material (hulis) or shanked-in 1 to 2" below hulis depth at planting; 2. Side-dress not more than 0.6 fl oz/1000 row-feet no later than 45 days after planting. Observe same PHI as above. Important Note: The rate applied affects the length of control. Use higher rate within the specified rate range where infestations occur later in crop development, or where pest pressure is continuous. Rates less than 0.7 fl oz/1000 row-feet may not provide adequate residual pest control. ARC-IMIDA 2# treated crops grown on very high organic matter soils (muck) may also require additional pest management control. RESTRICTIONS: Pre-Harvest Interval (PHI) from planting application: 3 days (leaves); 125 days (corms). Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per crop season. Do not apply ARC-IMIDA 2# more than once per crop season.

STRAWBERRIES

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

Annual and Perennial Strawberries

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	24.0 – 32.0 fl oz/A	Use one of the following methods to apply specified dosage of ARC-IMIDA 2#: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler, or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening; 2. As a plant material or plant hole treatment just prior to, or during transplanting. The application rate affects the length of control. Use higher rate within the specified rate range where infestations may occur later in crop development or where pest pressure is continuous.

		<p>RESTRICTIONS: Pre-Harvest Interval (PHI): 14 days. Do not apply immediately prior to bud opening or during bloom or when bees are foraging. Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per crop season. Do not use both application methods on the same crop in the same season.</p>
--	--	---

Post-Harvest Use on Perennial Strawberries

PESTS	RATE	INSTRUCTIONS
White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	16.0 – 24.0 fl oz/A	<p>Make a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply specified rate of ARC-IMIDA 2# using one of the following methods:</p> <ol style="list-style-type: none"> 1. As a ground spray by means of boom or backpack sprayer in a minimum of 20 gallons of water per acre; 2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed; 3. As a chemigation application with 600 – 1000 gallons of water followed by 0.10 to 0.25" irrigation. <p>Important Note: All soil-surface applications must be followed by either 0.25" of rainfall over overhead irrigation per acre within 2 hours of application in order to adequately incorporate ARC-IMIDA 2# into egg-deposition zone.</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 14 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per year. Do not use both application methods on the same crop in the same season.</p>

SUGARBEETS

(for use only in California)

Not for use on crops grown for seed unless specifically allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
Controlled:	6.0 – 12.0	Applied specified rate of ARC-IMIDA 2# in

Aphids Leafhoppers Whiteflies Flea beetles	fl oz/A	sufficient carrier for uniform application, directly below each seed furrow. Apply either during the bedding operation immediately prior to planting or at the time of planting.
Suppressed: Symptoms of: Western yellows/Beet curly top hybridgeminivirus		The low rate may be used to aid establishment of stands in whitefly areas, or for early season control of the other pests listed. RESTRICTIONS: Do not apply more than 12.0 fl oz (0.18 lb ai)/A ARC-IMIDA 2# per crop year. Do not apply more than a total of 0.18 lb active ingredient (any formulation) on any row spacing per year.

ARC-IMIDA 2# INSECTICIDE CONVERSION CHART FOR LINEAR APPLICATION

RATE Fluid Ounces/A	RATE fluid ounces/1000 row-feet Based on average row spacing (in inches)							
	10	15	20	25	30	35	40	45
10	0.19	0.29	0.33	0.43	0.57	0.67	0.76	0.86
12	0.23	0.34	0.46	0.57	0.69	0.80	0.92	1.03
14	0.27	0.40	0.54	0.67	0.80	0.94	1.07	1.21
16	0.31	0.46	0.61	0.77	0.92	1.07	1.22	1.38
18	0.34	0.52	0.69	0.86	1.03	1.21	1.38	1.55
20	0.38	0.57	0.76	0.96	1.15	1.34	1.53	1.72
22	0.42	0.63	0.84	1.05	1.26	1.47	1.68	1.89
24	0.46	0.69	0.92	1.15	1.38	1.61	1.84	2.07
26	0.50	0.75	0.99	1.24	1.49	1.74	1.99	2.24
28	0.54	0.80	1.07	1.34	1.61	1.87	2.14	2.41
30	0.57	0.86	1.15	1.43	1.72	2.01	2.29	2.58
32	0.6	0.92	1.22	1.52	1.84	2.14	2.45	2.75

Important Note: The application rate affects the length of control and to a considerable extent, the degree of control of effect. Row-spacing X ARC-IMIDA 2# rate combinations in shaded blocks may not proved adequate residual pest control and are not recommended for long-term, residual control. Use higher labeled rates as specified where infestations may occur later in crop development or where pest pressure is continuous. United Phosphorus, Inc. offers no warranty for use of ARC-IMIDA 2# at rates below 0.07 fl oz/1000 row-feet.

TREE, BUSH, AND VINE CROPS

BUSHBERRIES

Including: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Lingonberry, Salal

PESTS	RATE	INSTRUCTIONS
Japanese beetle	16.0 – 32.0	Use one of the following methods to apply

<p>(adults, feeding on foliage) White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)</p>	<p>fl oz/A</p>	<p>specified dosage of ARC-IMIDA 2#: 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. 18-inch band on each side of the row follow with 0.25" of irrigation immediately after application. Apply ARC-IMIDA 2# to moist soil. If necessary, apply one hour of irrigation water immediately before application of ARC-IMIDA 2#. For best results, ½ to 1" of irrigation water or rainfall should be applied or received within 24 hours of application to facilitate movement into the soil and into the root-zone. <u>Grubs:</u> For best grub control, apply ARC-IMIDA 2# to controls 1st or 2nd instar larvae. Make application post-bloom up to 7 days before harvest, or post-harvest until October 1st. <u>Japanese beetle larvae:</u> apply from June 1 to July 15. Apply to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field to control resident grub populations. Applications directed to the root-zone will help protect berry plant roots from grub feeding. RESTRICTIONS: Pre-Harvest Interval (PHI): 7 days Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per year.</p>
---	----------------	--

CITRUS (Containerized)

Including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, and other cultivars and/or hybrids of these.

PESTS	RATE	INSTRUCTIONS
<p>Controlled: Aphids Asian citrus psyllid Black fly Citrus leafminer</p>	<p>0.75 mL/ft³ container media</p>	<p>Determine volume of container and Calculate required dosage based on 0.50 mls/0.1 ft³ potting media. Apply calculated dosage per container as described above. Do not exceed rate of 3.0 mls / plant per crop season</p>

Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies		regardless of container size. Apply calculated dosage of ARC-IMIDA 2# per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For best results, make treatment at planting before insect infestation occurs.
Citrus Root weevil (larval complex)	1.25 – 2.50 ml/ft ³ container media	Retreat if necessary.
Suppressed: Citrus thrips	2.50 ml/ft ³ container media	Citrus root weevil complex (larvae): apply before neonate larvae entering potting media. Use the higher rate within the specified rate range for heavy infestations. RESTRICTIONS: Maximum rate is 2.50 ml/ft ³ container media. Do not apply immediately prior to bud opening or during bloom or when bees are foraging.

CITRUS (Field)

Including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine, Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote (*Casimiroa* spp), and other cultivar and/or hybrids of these.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Termites (FL only) Whiteflies	16.0 – 32.0 fl oz/A	Apply specified rate of ARC-IMIDA 2# using one of the following methods: 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For best results, apply to newly planted trees or those previously trained to drip, trickle, or micro-sprinkler irrigation. Lightly pre-wet the soil to break soil surface tension before applying ARC-IMIDA 2#. Chemigation application can be made separate to normal irrigation but should be followed by 10 to 20 minutes of additional watering to move ARC-IMIDA 2# into the root-zone. Allow 24 hours before initiating subsequent irrigations;
Suppressed: Symptoms of: Citrus tristeza virus (CTV) through vector control Citrus yellows Thrips (foliage feeding thrips only)	32.0 fl oz/A	2. Soil surface band spray on both sides of the tree. Bands should overlap at the tree base to create a continuous band within the drip-line area of the tree, to be

		<p>followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root-zone. This method is suitable for very coarse soils with 0.75% organic matter or less;</p> <p>3. Drench to base of tree not exceeding one-quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only recommended for trees up to 8 feet tall;</p> <p>For control of existing termite infestations, apply specified dosage in 1 – 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surround soil in the immediate vicinity of the tree trunk</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 0 day Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per year.</p>
--	--	---

CRANBERRIES

PESTS	RATE	INSTRUCTIONS
<p>Rootgrubs (scarabaeiae) Rootworms (chrysomelidae)</p>	<p>16.0-32.0 fl oz/A</p>	<p>Apply ARC-IMIDA 2# to moist soil. Apply specified rate of ARC-IMIDA 2# using one of the following methods:</p> <ol style="list-style-type: none"> 1. As a soil spray (ground application) directed to the root and crown area using at least 20 gal of water per acre; 2. As a chemigation application with 600 to 1000 gal water. <p>Immediately upon application, incorporate ARC-IMIDA 2# into root-zone by 0.1 – 0.3" water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.</p> <p><u>Rootgrubs and Rootworms:</u> Best control may be achieved when application is made post-bloom immediately after bees are removed. Applications should target early instar larvae.</p> <p>ARC-IMIDA 2# has not been tested for crop</p>

		<p>response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of ARC-IMIDA 2# and the desired fungicides or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 30 days Do not apply pre-bloom or during bloom or when bees are foraging. Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per season.</p>
--	--	---

GRAPES, Including: American bunch grape, Muscadine grape and Vinifera grape.

PESTS	RATE	INSTRUCTIONS
<p>Controlled: Mealybugs Leafhoppers/S harpshooters <i>Phylloxera</i>* spp.</p>	<p>16.0 – 32.0 fl oz/A</p> <p>24.0 – 32.0 fl oz/A</p>	<p>Apply specified dosage of ARC-IMIDA 2# using one of the following methods:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation; 3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation. <p>For best results, apply between bud-break and the pea-berry stage.</p> <p>*Repeated and regular use of ARC-IMIDA 2# over several, consecutive growing seasons controls existing <i>Phylloxera</i> infestations over time or prevents <i>Phylloxera</i> from becoming established.</p> <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 30 days Do not apply more than 32.0 fl oz (0.50 lb)/A ARC-IMIDA 2# per year.</p>
<p>Suppressed: Pierce's disease</p>		

HOPS

Not for use in California unless directed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
Aphids	19.2 fl oz/A	Apply specified dosage of ARC-IMIDA 2# using one of the following methods:

		<ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation; 3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation. <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 60 day Do not apply more than 19.2 fl oz (0.3 lb ai)/A ARC-IMIDA 2# per year.</p>
--	--	--

PECANS

Not for this use in California except when allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
<p>Controlled: Aphids Two-lined spittlebug</p> <p>Supressed: Pecan scab (from reduction in honey deposition)</p>	<p>16.0 – 32.0 fl oz/A</p>	<p>Make applications from May 15 until July 15. Applications made later in the season may be less effective.</p> <p>Apply product to slightly moist soil and allow soil to dry before additional irrigation.</p> <p>Apply specified rate of ARC-IMIDA 2# using one of the following methods:</p> <ol style="list-style-type: none"> 1. Chemigation to root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site; 3. Subsurface side-dress shanked into the root-zone near emitter line. Treat distance, wetted by the emitter set of each tree. <p>RESTRICTIONS: Pre-Harvest Interval (PHI): 7 days Do not apply pre-bloom or during bloom or when bees are foraging. Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per year.</p>

POME FRUITS

Including Apple, crabapple, loquat, mayhaw, pear (including Oriental pear), Quince

Not for this use in California except when allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
Aphids (including wooly apple aphid) Leafhoppers	16.0 – 24.0 fl oz/A	Apply specified dosage of ARC-IMIDA 2# through chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days Do not apply pre-bloom or during bloom or when bees are foraging. Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# year.

STONE FRUITS

Including: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

Not for this use in California except when allowed by state specific 24(c) labeling.

In-field Soil Application

PESTS	RATE	INSTRUCTIONS
Aphids (including wooly apple aphid) Leafhoppers	16.0 – 24.0 fl oz/A	Applications Apply specified dosage of ARC-IMIDA 2# by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. RESTRICTIONS: Pre-Harvest Interval (PHI): 21 days Do not apply pre-bloom or during bloom or when bees are foraging. Do not apply more than 24.0 fl oz (0.38 lb ai)/A ARC-IMIDA 2# per year.

Pre-Plant Root Dip Application

PESTS	RATE	INSTRUCTIONS
Black peach aphid (infesting roots)	2.0 fl oz/10 gal root-dip solution	Mix ARC-IMIDA 2# at specified rate per 10 gallons of water. Completely wet bare-root transplant to slightly above the graft union by soaking them in the ARC-IMIDA 2# solution for up to 5 minutes. Allow solution to dry on roots then transplant trees as soon as possible following treatment.

TROPICAL FRUIT

Including: Acerola, Avocado, Black sapote, Canistel, Feijoa, Jaboticaba, Guava, Longan, Lychee, Marney sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan Rambutan, Sapodilla, Spanish lime, Star apple, Starfruit, Wax jambu
 Not for this use in California except when allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Leafhoppers Whiteflies	24.0 – 32.0 fl oz/A	Apply specified rate of ARC-IMIDA 2# by chemigation through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
Suppressed: Scales	32.0 fl oz/A	RESTRICTIONS: Pre-Harvest Interval (PHI): 6 days Do not apply pre-bloom or during bloom or when bees are foraging. Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per year.

POPLAR/COTTONWOOD

Includes members of the genus Populus grown for pulp or timber
 Not for this use in California except when allowed by state specific 24(c) labeling.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Cottonwood leaf beetle	16.0 – 32.0 fl oz/A	Apply specified rate of ARC-IMIDA 2# by chemigation through low-pressure drip irrigation.
Suppressed: <i>Phylloxera popularia</i>		<u>Cottonwood leaf beetle:</u> make application early, when the beetles first begin feeding, for best protection. Larger trees may need earlier treatment as a result of slower uptake. <u>Phylloxera:</u> apply early in the year, from break of dormancy through May. RESTRICTIONS: Do not apply pre-bloom or during bloom or when bees are foraging. Do not apply more than 32.0 fl oz (0.50 lb ai)/A ARC-IMIDA 2# per year.

IMPORTANT INFORMATION READ BEFORE USING PRODUCTS

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such

factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of ARCANA LLC or Seller, All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold ARCANA LLC and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, ARCANA LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or ARCANA LLC, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ARCANA LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by state law, neither ARCANA LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ARCANA LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ARCANA, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

ARCANA LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of Sale and Limitation of Warranty and Liability which may not be modified except by written agreement signed by a duly authorized representative of ARCANA LLC.