

# Annexes to the Inventory of U.S. GHG Emissions and Sinks

The following eight annexes provide additional information related to the material presented in the main body of this report as directed in the *UNFCCC Guidelines on Reporting and Review* (UNFCCC 2003). Annex I contains an analysis of the key categories of emissions discussed in this report and a review of the methodology used to identify those key categories. Annex 2 describes the methodologies used to estimate CO<sub>2</sub> emissions from fossil fuel combustion, the carbon content of fossil fuels, and the amount of carbon stored in products from non-energy uses of fossil fuels. Annex 3 discusses the methodologies used for a number of individual source categories in greater detail than was presented in the main body of the report and includes explicit activity data and emission factor tables. Annex 4 presents the IPCC reference approach for estimating CO<sub>2</sub> emissions from fossil fuel combustion. Annex 5 addresses the criteria for the inclusion of an emission source category and discusses some of the sources that are excluded from U.S. estimates. Annex 6 provides a range of additional information that is relevant to the contents of this report. Annex 7 provides data on the uncertainty of the emission estimates included in this report. Finally, Annex 8 provides information on the QA/QC methods and procedures used in the development of the Inventory.

|  |     |
|--|-----|
| Annexes to the Inventory of U.S. GHG Emissions and Sinks.....  | 1   |
| ANNEX 1 Key Category Analysis .....  | 2   |
| ANNEX 2 Methodology and Data for Estimating CO <sub>2</sub> Emissions from Fossil Fuel Combustion .....  | 30  |
| 2.1. Methodology for Estimating Emissions of CO <sub>2</sub> from Fossil Fuel Combustion .....   | 30  |
| 2.2. Methodology for Estimating the Carbon Content of Fossil Fuels.....  | 64  |
| 2.3. Methodology for Estimating Carbon Emitted from Non-Energy Uses of Fossil Fuels .....  | 102 |
| ANNEX 3 Methodological Descriptions for Additional Source or Sink Categories.....  | 129 |
| 3.1. Methodology for Estimating Emissions of CH <sub>4</sub> , N <sub>2</sub> O, and Indirect Greenhouse Gases from Stationary Combustion.....   | 129 |
| 3.2. Methodology for Estimating Emissions of CH <sub>4</sub> , N <sub>2</sub> O, and Indirect Greenhouse Gases from Mobile Combustion and Methodology for and Supplemental Information on Transportation-Related GHG Emissions ..... | 137 |
| 3.3. Methodology for Estimating Emissions from Commercial Aircraft Jet Fuel Consumption.....   | 176 |
| 3.4. Methodology for Estimating CH <sub>4</sub> Emissions from Coal Mining .....   | 180 |
| 3.5. Methodology for Estimating CH <sub>4</sub> and CO <sub>2</sub> Emissions from Petroleum Systems .....   | 189 |
| 3.6. Methodology for Estimating CH <sub>4</sub> and CO <sub>2</sub> Emissions from Natural Gas Systems.....  | 198 |
| 3.7. Methodology for Estimating CO <sub>2</sub> , N <sub>2</sub> O, and CH <sub>4</sub> Emissions from the Incineration of Waste .....   | 223 |
| 3.8. Methodology for Estimating Emissions from International Bunker Fuels used by the U.S. Military .....  | 230 |
| 3.9. Methodology for Estimating HFC and PFC Emissions from Substitution of Ozone Depleting Substances .....  | 237 |
| 3.10. Methodology for Estimating CH <sub>4</sub> Emissions from Enteric Fermentation.....  | 259 |
| 3.11. Methodology for Estimating CH <sub>4</sub> and N <sub>2</sub> O Emissions from Manure Management .....   | 283 |
| 3.12. Methodology for Estimating N <sub>2</sub> O Emissions, CH <sub>4</sub> Emissions and Soil Organic C Stock Changes from Agricultural Lands (Cropland and Grassland) .....   | 318 |
| 3.13. Methodology for Estimating Net Carbon Stock Changes in <i>Forest Land Remaining Forest Land</i> and <i>Land Converted to Forest Land</i> .....   | 385 |
| 3.14. Methodology for Estimating CH <sub>4</sub> Emissions from Landfills.....   | 420 |
| ANNEX 4 IPCC Reference Approach for Estimating CO <sub>2</sub> Emissions from Fossil Fuel Combustion .....   | 430 |
| ANNEX 5 Assessment of the Sources and Sinks of Greenhouse Gas Emissions Not Included .....   | 440 |
| ANNEX 6 Additional Information.....  | 443 |
| 6.1. Global Warming Potential Values .....   | 443 |
| 6.2. Ozone Depleting Substance Emissions.....  | 454 |
| 6.3. Sulfur Dioxide Emissions .....  | 456 |
| ANNEX 7 Uncertainty.....   | 472 |
| 7.1. Overview.....   | 472 |
| 7.2. Methodology and Results .....   | 472 |
| 7.3. Planned Improvements .....  | 479 |
| 7.4. Additional Information on Uncertainty Analyses by Source .....  | 480 |
| ANNEX 8 QA/QC Procedures.....  | 491 |
| 8.1. Background.....   | 491 |
| 8.2. Purpose.....  | 491 |
| 8.3. Assessment Factors .....  | 492 |

# ANNEX 1 Key Category Analysis

The United States has identified national key categories based on the estimates presented in this report. The *2006 Intergovernmental Panel on Climate Change (IPCC) Guidelines for National Greenhouse Gas Inventories* (IPCC 2006) describes a key category as a “[category] that is prioritized within the national inventory system because its estimate has a significant influence on a country’s total inventory of greenhouse gases in terms of the absolute level, the trend, or the uncertainty in emissions and removals.” By definition, key categories are sources or sinks that have the greatest contribution to the absolute overall level of national emissions in any of the years covered by the time series. In addition, when an entire time series of emission estimates is prepared, a determination of key categories must also account for the influence of the trends of individual categories. Therefore, a trend assessment is conducted to identify source and sink categories for which significant uncertainty in the estimate would have considerable effects on overall emission trends. Finally, a qualitative evaluation of key categories should be performed, in order to capture any key categories that were not identified in either of the quantitative analyses, but can be considered key because of the unique country-specific estimation methods.

The methodology for conducting a key category analysis, as defined by the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006), includes:

- Approach 1 (including both level and trend assessments);
- Approach 2 (including both level and trend assessments, and incorporating uncertainty analysis); and
- Qualitative approach.

This Annex presents an analysis of key categories, both for sources only and also for sources and sinks (i.e., including Land Use, Land-Use Change and Forestry [LULUCF]); discusses Approach 1, Approach 2, and qualitative approaches to identifying key categories; provides level and trend assessment equations; and provides a brief statistical evaluation of IPCC’s quantitative methodologies for defining key categories. Table A-1 presents the key categories for the United States (including and excluding LULUCF categories) using emissions and uncertainty data in this report, and ranked according to their sector and global warming potential (GWP)-weighted emissions in 2014. The table also indicates the criteria used in identifying these categories (i.e., level, trend, Approach 1, Approach 2, and/or qualitative assessments).

**Table A-1 : Key Source Categories for the United States (1990-2014)**

| IPCC Source Categories   | Gas             | Approach 1           |                      |                   |                   | Approach 2           |                      |                   |                   | Qual <sup>a</sup> | 2014 Emissions (MMT CO <sub>2</sub> Eq.) |
|--|-----------------|----------------------|----------------------|-------------------|-------------------|----------------------|----------------------|-------------------|-------------------|-------------------|--|
|  |                 | Level Without LULUCF | Trend Without LULUCF | Level With LULUCF | Trend With LULUCF | Level Without LULUCF | Trend Without LULUCF | Level With LULUCF | Trend With LULUCF |                   |  |
| <b>Energy</b>  |                 |                      |                      |                   |                   |                      |                      |                   |                   |                   |  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 1,570.4                                  |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                               | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 1,467.5                                  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial              | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 466.0                                    |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation  | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 443.2                                    |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential             | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 277.6                                    |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial              | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 271.9                                    |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial              | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 189.2                                    |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                           | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 150.1                                    |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                               | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 114.3                                    |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                              | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 92.0                                     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial             | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 75.3                                     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential             | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 67.5                                     |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                   | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 42.4                                     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial              | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 38.2                                     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories        | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 34.6                                     |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                             | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 28.0                                     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation  | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 25.3                                     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial             | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 4.5                                      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories        | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 3.0                                      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential            | CO <sub>2</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 0.0                                      |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                   | CH <sub>4</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 176.1                                    |
| CH <sub>4</sub> Emissions from Petroleum Systems                                     | CH <sub>4</sub> | .                    | .                    | .                 | .                 | .                    | .                    | .                 | .                 |                   | 68.1                                     |

|  |                  |   |   |   |   |   |   |   |   |                |
|--|------------------|---|---|---|---|---|---|---|---|----------------|
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub>  | • | • | • | • | • | • | • | • | 67.6           |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  |   |   |   |   |   |   | • | • | 5.0            |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | N <sub>2</sub> O |   | • |   | • |   |   | • | • | 19.6           |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                                  | N <sub>2</sub> O | • | • | • | • |   |   | • |   | 12.5           |
| International Bunker Fuels <sup>b</sup>  | Several          |   |   |   |   |   |   |   | • | 104.2          |
| <b>Industrial Processes and Product Use</b>  |                  |   |   |   |   |   |   |   |   |                |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | • | • | • | • |   |   | • | • | 55.4           |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | • |   | • |   |   |   |   |   | 38.8           |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  |   | • |   | • |   |   |   |   | 12.1           |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O |   | • |   | • |   |   |   |   | 5.4            |
| Emissions from Substitutes for Ozone Depleting Substances                                | HiGWP            | • | • | • | • |   |   | • | • | 161.2          |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | HiGWP            |   | • |   | • |   |   | • | • | 5.6            |
| HFC-23 Emissions from HCFC-22 Production   | HiGWP            | • | • | • | • |   |   | • |   | 5.0            |
| PFC Emissions from Aluminum Production   | HiGWP            |   | • |   | • |   |   |   |   | 2.5            |
| <b>Agriculture</b>   |                  |   |   |   |   |   |   |   |   |                |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>  | • | • | • | • |   |   | • | • | 164.3          |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | • | • | • | • |   |   | • | • | 61.2           |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                      | N <sub>2</sub> O | • |   | • |   |   |   | • |   | 261.0          |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O | • |   | • |   |   |   | • | • | 57.3           |
| <b>Waste</b>   |                  |   |   |   |   |   |   |   |   |                |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>  | • | • | • | • |   |   | • | • | 148.0          |
| <b>Land Use, Land Use Change, and Forestry</b>   |                  |   |   |   |   |   |   |   |   |                |
| CO <sub>2</sub> Emissions from Land Converted to Grassland                               | CO <sub>2</sub>  |   |   | • |   |   |   |   |   | 40.4           |
| CO <sub>2</sub> Emissions from Land Converted to Cropland                                | CO <sub>2</sub>  |   |   | • | • |   |   | • | • | 22.1           |
| CO <sub>2</sub> Emissions from Grassland Remaining Grassland                             | CO <sub>2</sub>  |   |   |   | • |   |   | • | • | 3.8            |
| CO <sub>2</sub> Emissions from Cropland Remaining Cropland                               | CO <sub>2</sub>  |   |   | • | • |   |   | • | • | (8.4)          |
| CO <sub>2</sub> Emissions from Landfilled Yard Trimmings and Food Scraps                 | CO <sub>2</sub>  |   |   |   | • |   |   | • | • | (11.6)         |
| CO <sub>2</sub> Emissions from Urban Trees   | CO <sub>2</sub>  |   |   | • | • |   |   | • | • | (90.6)         |
| CO <sub>2</sub> Emissions from Forest Land Remaining Forest Land                         | CO <sub>2</sub>  |   |   | • | • |   |   | • | • | (742.3)        |
| CH <sub>4</sub> Emissions from Forest Fires  | CH <sub>4</sub>  |   |   |   |   |   |   | • | • | 7.3            |
| <b>Subtotal Without LULUCF</b>   |                  |   |   |   |   |   |   |   |   | <b>6,687.8</b> |

|  |                |
|--|----------------|
| <b>Total Emissions Without LULUCF</b>  | <b>6,870.5</b> |
| <b>Percent of Total Without LULUCF</b> | <b>97%</b>     |
| <b>Subtotal With LULUCF</b>            | <b>5,905.5</b> |
| <b>Total Emissions With LULUCF</b>     | <b>6,108.0</b> |
| <b>Percent of Total With LULUCF</b>    | <b>97%</b>     |

<sup>a</sup> Qualitative criteria.

<sup>b</sup> Emissions from this source not included in totals.

Notes: Parentheses indicate negative values (or sequestration). Table A-2 provides a complete listing of source categories by IPCC sector, along with notations on the criteria used in identifying key categories, without LULUCF sources and sinks. Similarly, Table A-3 provides a complete listing of source and sink categories by IPCC sector, along with notations on the criteria used in identifying key categories, including LULUCF sources and sinks. The notations refer specifically to the year(s) in the inventory time series (i.e., 1990 to 2014) in which each source category reached the threshold for being a key category based on either an Approach 1 or Approach 2 level assessment.

In addition to conducting Approach 1 and 2 level and trend assessments, a qualitative assessment of the source categories, as described in the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006), was conducted to capture any key categories that were not identified by any quantitative method. One additional key category, international bunker fuels, was identified using this qualitative assessment. International bunker fuels are fuels consumed for aviation or marine international transport activities, and emissions from these fuels are reported separately from totals in accordance with IPCC guidelines. If these emissions were included in the totals, bunker fuels would qualify as a key category according to Approach 1. The amount of uncertainty associated with estimation of emissions from international bunker fuels also supports the qualification of this source category as key, which would qualify it as a key category according to Approach 2.

**Table A-2: U.S. Greenhouse Gas Inventory Source Categories without LULUCF**

| IPCC Source Categories   | Direct GHG      | 2014 Emissions<br>(MMT CO <sub>2</sub> Eq.) | Key<br>Category? | ID<br>Criteria <sup>a</sup>                                 | Level in which<br>year(s)? <sup>b</sup> |
|--|-----------------|---|------------------|---|---|
| <b>Energy</b>  |                 |   |                  |   |   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation | CO <sub>2</sub> | 1,570.4                                     | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                               | CO <sub>2</sub> | 1,467.5                                     | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial              | CO <sub>2</sub> | 466.0                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation  | CO <sub>2</sub> | 443.2                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990 <sub>1</sub> , 2014                |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential             | CO <sub>2</sub> | 277.6                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial              | CO <sub>2</sub> | 271.9                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial              | CO <sub>2</sub> | 189.2                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                           | CO <sub>2</sub> | 150.1                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014 <sub>1</sub>                 |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                               | CO <sub>2</sub> | 114.3                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                              | CO <sub>2</sub> | 92.0  | •                | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub>   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial             | CO <sub>2</sub> | 75.3  | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential             | CO <sub>2</sub> | 67.5  | •                | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub>   |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                   | CO <sub>2</sub> | 42.4  | •                | L <sub>1</sub> L <sub>2</sub>                               | 1990, 2014                              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial              | CO <sub>2</sub> | 38.2  | •                | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub>   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories        | CO <sub>2</sub> | 34.6  | •                | L <sub>1</sub>  | 2014 <sub>1</sub>                       |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                             | CO <sub>2</sub> | 28.0  | •                | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub>                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation  | CO <sub>2</sub> | 25.3  | •                | L <sub>1</sub> T <sub>1</sub> T <sub>2</sub>                | 1990 <sub>1</sub>                       |
| CO <sub>2</sub> Emissions from Incineration of Waste                                 | CO <sub>2</sub> | 9.4   |                  |   |   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial             | CO <sub>2</sub> | 4.5   | •                | T <sub>1</sub>  |   |
| CO <sub>2</sub> Emissions from Petroleum Systems                                     | CO <sub>2</sub> | 3.6   |                  |   |   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories       | CO <sub>2</sub> | 3.4   |                  |   |   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories        | CO <sub>2</sub> | 3.0   | •                | T <sub>2</sub>  |   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy             | CO <sub>2</sub> | 0.4   |                  |   |   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential            | CO <sub>2</sub> | 0.0   | •                | T <sub>2</sub>  |   |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                   | CH <sub>4</sub> | 176.1                                       | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| CH <sub>4</sub> Emissions from Petroleum Systems                                     | CH <sub>4</sub> | 68.1  | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                              |
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub> | 67.6  | •                | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014 <sub>1</sub>                 |
| Fugitive Emissions from Abandoned Underground Coal Mines                             | CH <sub>4</sub> | 6.3   |                  |   |   |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential               | CH <sub>4</sub> | 5.0   | •                | L <sub>2</sub>  | 1990 <sub>2</sub>                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                | CH <sub>4</sub> | 1.5   |                  |   |   |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                               | CH <sub>4</sub> | 1.4   |                  |   |   |

|   |                  |       |   |   |                                       |
|---|------------------|-------|---|---|---------------------------------------|
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial             | CH <sub>4</sub>  | 1.1   |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                           | CH <sub>4</sub>  | 0.5   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation | CH <sub>4</sub>  | 0.4   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories       | CH <sub>4</sub>  | 0.1   |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                        | CH <sub>4</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                          | CH <sub>4</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Incineration of Waste                              | CH <sub>4</sub>  | +     |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation | N <sub>2</sub> O | 19.6  | • | T <sub>1</sub> L <sub>2</sub> T <sub>2</sub>                | 1990 <sub>2</sub> , 2014 <sub>2</sub> |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                           | N <sub>2</sub> O | 12.5  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990                                  |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial             | N <sub>2</sub> O | 2.4   |   |   |                                       |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                          | N <sub>2</sub> O | 2.0   |   |   |                                       |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                       | N <sub>2</sub> O | 1.4   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential            | N <sub>2</sub> O | 1.0   |   |   |                                       |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                         | N <sub>2</sub> O | 0.5   |   |   |                                       |
| N <sub>2</sub> O Emissions from Incineration of Waste                             | N <sub>2</sub> O | 0.3   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial             | N <sub>2</sub> O | 0.3   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories       | N <sub>2</sub> O | 0.1   |   |   |                                       |
| International Bunker Fuels <sup>c</sup>   | Several          | 104.2 | • |   |                                       |

---

#### Industrial Processes and Product Use

---

|  |                  |      |   |   |                                       |
|--|------------------|------|---|---|---------------------------------------|
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | 55.4 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014 <sub>1</sub>               |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 38.8 | • | L <sub>1</sub>  | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 26.5 |   |   |                                       |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 14.1 |   |   |                                       |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  | 12.1 | • | T <sub>1</sub>  |                                       |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 9.4  |   |   |                                       |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 4.5  |   |   |                                       |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 4.0  |   |   |                                       |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 2.8  |   |   |                                       |
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8  |   |   |                                       |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 1.9  |   |   |                                       |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.8  |   |   |                                       |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.3  |   |   |                                       |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.1  |   |   |                                       |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 1.0  |   |   |                                       |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5  |   |   |                                       |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.2  |   |   |                                       |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +    |   |   |                                       |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.1  |   |   |                                       |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +    |   |   |                                       |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +    |   |   |                                       |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +    |   |   |                                       |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 10.9 |   |   |                                       |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 5.4  | • | T <sub>1</sub>  |                                       |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2  |   |   |                                       |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | 0.2  |   |   |                                       |

|  |                  |       |   |   |                          |
|--|------------------|-------|---|---|--------------------------|
| Emissions from Substitutes for Ozone Depleting Substances                                | HiGWP            | 161.2 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 2014                     |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | HiGWP            | 5.6   | • | L <sub>1</sub> T <sub>1</sub> T <sub>2</sub>                | 1990 <sub>1</sub>        |
| HFC-23 Emissions from HCFC-22 Production   | HiGWP            | 5.0   | • | T <sub>1</sub> T <sub>2</sub>                               |                          |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | HiGWP            | 4.5   |   |   |                          |
| PFC Emissions from Aluminum Production   | HiGWP            | 2.5   | • | T <sub>1</sub>  |                          |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | HiGWP            | 1.0   |   |   |                          |
| HFC-134a Emissions from Magnesium Production and Processing                              | HiGWP            | 0.1   |   |   |                          |
| <b>Agriculture</b>   |                  |       |   |   |                          |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>  | 164.3 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014               |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | 61.2  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990 <sub>1</sub> , 2014 |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 11.9  |   |   |                          |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.3   |   |   |                          |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                      | N <sub>2</sub> O | 261.0 | • | L <sub>1</sub> L <sub>2</sub>                               | 1990, 2014               |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O | 57.3  | • | L <sub>1</sub> L <sub>2</sub> T <sub>2</sub>                | 1990, 2014               |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 17.5  |   |   |                          |
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues                   | N <sub>2</sub> O | 0.1   |   |   |                          |
| <b>Waste</b>   |                  |       |   |   |                          |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>  | 148.0 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014               |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>  | 14.7  |   |   |                          |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 2.1   |   |   |                          |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 4.8   |   |   |                          |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 1.8   |   |   |                          |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> For the ID criteria, Q refers to "Qualitative", L refers to a key category identified through a level assessment; T refers to a key category identified through a trend assessment and the subscripted number refers to either an Approach 1 or Approach 2 assessment (e.g., L<sub>2</sub> designates a source is a key category for an Approach 2 level assessment).

<sup>b</sup> If the source is a key category for both L<sub>1</sub> and L<sub>2</sub> (as designated in the ID criteria column), it is a key category for both assessments in the years provided unless noted by a subscript, in which case it is a key category for that assessment in that year only (e.g., 1990<sub>2</sub> designates a source is a key category for the Approach 2 assessment only in 1990).

<sup>c</sup> Emissions from these sources not included in totals.

Note: LULUCF sources and sinks are not included in this analysis.

**Table A-3: U.S. Greenhouse Gas Inventory Source Categories with LULUCF**

| IPCC Source Categories   | Direct GHG      | 2014 Emissions (MMT CO <sub>2</sub> Eq.) | Key Category? | ID Criteria <sup>a</sup>                                    | Level in which year(s) <sup>b</sup>   |
|--|-----------------|--|---------------|---|---------------------------------------|
| <b>Energy</b>  |                 |  |               |   |                                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation | CO <sub>2</sub> | 1,570.4                                  | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                               | CO <sub>2</sub> | 1,467.5                                  | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial              | CO <sub>2</sub> | 466.0                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation  | CO <sub>2</sub> | 443.2                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990 <sub>1</sub> , 2014              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential             | CO <sub>2</sub> | 277.6                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial              | CO <sub>2</sub> | 271.9                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial              | CO <sub>2</sub> | 189.2                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990 <sub>1</sub> , 2014              |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                           | CO <sub>2</sub> | 150.1                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990, 2014 <sub>1</sub>               |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                               | CO <sub>2</sub> | 114.3                                    | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                              | CO <sub>2</sub> | 92.0                                     | •             | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial             | CO <sub>2</sub> | 75.3                                     | •             | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |

|   |                  |       |   |   |                                       |
|---|------------------|-------|---|---|---------------------------------------|
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential            | CO <sub>2</sub>  | 67.5  | • | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                  | CO <sub>2</sub>  | 42.4  | • | L <sub>1</sub> L <sub>2</sub>                               | 1990 <sub>1</sub> , 2014              |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial             | CO <sub>2</sub>  | 38.2  | • | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories       | CO <sub>2</sub>  | 34.6  | • | L <sub>1</sub> T <sub>1</sub>                               | 2014 <sub>1</sub>                     |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                            | CO <sub>2</sub>  | 28.0  | • | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation | CO <sub>2</sub>  | 25.3  | • | L <sub>1</sub> T <sub>1</sub> T <sub>2</sub>                | 1990 <sub>1</sub>                     |
| CO <sub>2</sub> Emissions from Incineration of Waste                                | CO <sub>2</sub>  | 9.4   |   |   |                                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial            | CO <sub>2</sub>  | 4.5   | • | T <sub>1</sub>  |                                       |
| CO <sub>2</sub> Emissions from Petroleum Systems                                    | CO <sub>2</sub>  | 3.6   |   |   |                                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories      | CO <sub>2</sub>  | 3.4   |   |   |                                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories       | CO <sub>2</sub>  | 3.0   |   |   |                                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy            | CO <sub>2</sub>  | 0.4   |   |   |                                       |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential           | CO <sub>2</sub>  | 0.0   |   |   |                                       |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                  | CH <sub>4</sub>  | 176.1 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CH <sub>4</sub> Emissions from Petroleum Systems                                    | CH <sub>4</sub>  | 68.1  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| Fugitive Emissions from Coal Mining   | CH <sub>4</sub>  | 67.6  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014 <sub>1</sub>               |
| Fugitive Emissions from Abandoned Underground Coal Mines                            | CH <sub>4</sub>  | 6.3   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential              | CH <sub>4</sub>  | 5.0   | • | L <sub>2</sub>  | 1990 <sub>2</sub> , 2014 <sub>2</sub> |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial               | CH <sub>4</sub>  | 1.5   |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                              | CH <sub>4</sub>  | 1.4   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial               | CH <sub>4</sub>  | 1.1   |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                             | CH <sub>4</sub>  | 0.5   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation   | CH <sub>4</sub>  | 0.4   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories         | CH <sub>4</sub>  | 0.1   |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                          | CH <sub>4</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                            | CH <sub>4</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Incineration of Waste                                | CH <sub>4</sub>  | +     |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation   | N <sub>2</sub> O | 19.6  | • | T <sub>1</sub> L <sub>2</sub> T <sub>2</sub>                | 1990 <sub>2</sub> , 2014 <sub>2</sub> |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                             | N <sub>2</sub> O | 12.5  | • | L <sub>1</sub> T <sub>1</sub> T <sub>2</sub>                | 1990 <sub>1</sub>                     |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial               | N <sub>2</sub> O | 2.4   |   |   |                                       |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                            | N <sub>2</sub> O | 2.0   |   |   |                                       |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                         | N <sub>2</sub> O | 1.4   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential              | N <sub>2</sub> O | 1.0   |   |   |                                       |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                           | N <sub>2</sub> O | 0.5   |   |   |                                       |
| N <sub>2</sub> O Emissions from Incineration of Waste                               | N <sub>2</sub> O | 0.3   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial               | N <sub>2</sub> O | 0.3   |   |   |                                       |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories         | N <sub>2</sub> O | 0.1   |   |   |                                       |
| International Bunker Fuels <sup>c</sup>   | Several          | 104.2 | • |   |                                       |

## Industrial Processes and Product Use

|  |                  |       |   |   |                                       |
|--|------------------|-------|---|---|---------------------------------------|
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | 55.4  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014 <sub>1</sub>               |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 38.8  | • | L <sub>1</sub>  | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 26.5  |   |   |                                       |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 14.1  |   |   |                                       |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  | 12.1  | • | T <sub>1</sub>  |                                       |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 9.4   |   |   |                                       |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 4.5   |   |   |                                       |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 4.0   |   |   |                                       |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 2.8   |   |   |                                       |
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8   |   |   |                                       |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 1.9   |   |   |                                       |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.8   |   |   |                                       |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.3   |   |   |                                       |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.1   |   |   |                                       |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 1.0   |   |   |                                       |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5   |   |   |                                       |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.2   |   |   |                                       |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.1   |   |   |                                       |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +     |   |   |                                       |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +     |   |   |                                       |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 10.9  |   |   |                                       |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 5.4   | • | T <sub>1</sub>  |                                       |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2   |   |   |                                       |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | 0.2   |   |   |                                       |
| Emissions from Substitutes for Ozone Depleting Substances                                | HiGWP            | 161.2 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 2014                                  |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | HiGWP            | 5.6   | • | L <sub>1</sub> T <sub>1</sub>                               | 1990 <sub>1</sub>                     |
| HFC-23 Emissions from HCFC-22 Production   | HiGWP            | 5.0   | • | T <sub>1</sub> T <sub>2</sub>                               |                                       |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | HiGWP            | 4.5   |   |   |                                       |
| PFC Emissions from Aluminum Production   | HiGWP            | 2.5   | • | T <sub>1</sub>  |                                       |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | HiGWP            | 1.0   |   |   |                                       |
| HFC-134a Emissions from Magnesium Production and Processing                              | HiGWP            | 0.1   |   |   |                                       |
| <b>Agriculture</b>   |                  |       |   |   |                                       |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>  | 164.3 | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990, 2014                            |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | 61.2  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub>                | 1990 <sub>1</sub> , 2014              |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 11.9  |   |   |                                       |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.3   |   |   |                                       |
| DirectN <sub>2</sub> O Emissions from Agricultural Soil Management                       | N <sub>2</sub> O | 261.0 | • | L <sub>1</sub> L <sub>2</sub>                               | 1990, 2014                            |
| IndirectN <sub>2</sub> O Emissions from Applied Nitrogen                                 | N <sub>2</sub> O | 57.3  | • | L <sub>1</sub> L <sub>2</sub> T <sub>2</sub>                | 1990, 2014                            |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 17.5  |   |   |                                       |

|  |                  |         |   |   |                                       |
|--|------------------|---------|---|---|---------------------------------------|
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues   | N <sub>2</sub> O | 0.1     |   |   |                                       |
| <b>Waste</b>   |                  |         |   |   |                                       |
| CH <sub>4</sub> Emissions from Landfills                                 | CH <sub>4</sub>  | 148.0   | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CH <sub>4</sub> Emissions from Wastewater Treatment                      | CH <sub>4</sub>  | 14.7    |   |   |                                       |
| CH <sub>4</sub> Emissions from Composting                                | CH <sub>4</sub>  | 2.1     |   |   |                                       |
| N <sub>2</sub> O Emissions from Wastewater Treatment                     | N <sub>2</sub> O | 4.8     |   |   |                                       |
| N <sub>2</sub> O Emissions from Composting                               | N <sub>2</sub> O | 1.8     |   |   |                                       |
| <b>Land Use, Land Use Change, and Forestry</b>                           |                  |         |   |   |                                       |
| CO <sub>2</sub> Emissions from Land Converted to Grassland               | CO <sub>2</sub>  | 40.4    | • | L <sub>1</sub>  | 1990 <sub>1</sub> , 2014 <sub>1</sub> |
| CO <sub>2</sub> Emissions from Land Converted to Cropland                | CO <sub>2</sub>  | 22.1    | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990                                  |
| CO <sub>2</sub> Emissions from Urea Fertilization                        | CO <sub>2</sub>  | 4.5     |   |   |                                       |
| CO <sub>2</sub> Emissions from Liming                                    | CO <sub>2</sub>  | 4.1     |   |   |                                       |
| CO <sub>2</sub> Emissions from Grassland Remaining Grassland             | CO <sub>2</sub>  | 3.8     | • | T <sub>1</sub> L <sub>2</sub> T <sub>2</sub>                | 1990 <sub>2</sub> , 2014 <sub>2</sub> |
| CO <sub>2</sub> Emissions from Peatlands Remaining Peatlands             | CO <sub>2</sub>  | 0.8     |   |   |                                       |
| CO <sub>2</sub> Emissions from Land Converted to Forest Land             | CO <sub>2</sub>  | (0.3)   |   |   |                                       |
| CO <sub>2</sub> Emissions from Cropland Remaining Cropland               | CO <sub>2</sub>  | (8.4)   | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014 <sub>2</sub>               |
| CO <sub>2</sub> Emissions from Landfilled Yard Trimmings and Food Scraps | CO <sub>2</sub>  | (11.6)  | • | T <sub>1</sub> L <sub>2</sub> T <sub>2</sub>                | 1990 <sub>2</sub>                     |
| CO <sub>2</sub> Emissions from Urban Trees                               | CO <sub>2</sub>  | (90.6)  | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CO <sub>2</sub> Emissions from Forest Land Remaining Forest Land         | CO <sub>2</sub>  | (742.3) | • | L <sub>1</sub> T <sub>1</sub> L <sub>2</sub> T <sub>2</sub> | 1990, 2014                            |
| CH <sub>4</sub> Emissions from Forest Fires                              | CH <sub>4</sub>  | 7.3     | • | L <sub>2</sub> T <sub>2</sub>                               | 2014 <sub>2</sub>                     |
| CH <sub>4</sub> Emissions from Peatlands Remaining Peatlands             | CH <sub>4</sub>  | +       |   |   |                                       |
| N <sub>2</sub> O Emissions from Forest Fires                             | N <sub>2</sub> O | 4.8     |   |   |                                       |
| N <sub>2</sub> O Emissions from Settlement Soils                         | N <sub>2</sub> O | 2.4     |   |   |                                       |
| N <sub>2</sub> O Emissions from Forest Soils                             | N <sub>2</sub> O | 0.5     |   |   |                                       |
| N <sub>2</sub> O Emissions from Peatlands Remaining Peatlands            | N <sub>2</sub> O | +       |   |   |                                       |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> For the ID criteria, Q refers to "Qualitative", L refers to a key category identified through a level assessment; T refers to a key category identified through a trend assessment and the subscripted number refers to either an Approach 1 or Approach 2 assessment (e.g., L<sub>2</sub> designates a source is a key category for an Approach 2 level assessment).

<sup>b</sup> If the source is a key category for both L<sub>1</sub> and L<sub>2</sub> (as designated in the ID criteria column), it is a key category for both assessments in the years provided unless noted by a subscript, in which case it is a key category only for that assessment in only that year (e.g., 1990<sub>2</sub> designates a source is a key category for the Approach 2 assessment only in 1990).

<sup>c</sup> Emissions from these sources not included in totals.

Note: Parentheses indicate negative values (or sequestration).

## Evaluation of Key Categories

### Level Assessment

When using an Approach 1 for the level assessment, a predetermined cumulative emissions threshold is used to identify key categories. When source and sink categories are sorted in order of decreasing absolute emissions, those that fall at the top of the list and cumulatively account for 95 percent of emissions are considered key categories. The 95 percent threshold in the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006) was designed to establish a general level where the key category analysis covers approximately 75 to 92 percent of inventory uncertainty.

Including the Approach 2 provides additional insight into why certain source categories are considered key, and how to prioritize inventory improvements. In the Approach 2, the level assessment for each category from the Approach 1 is multiplied by its percent relative uncertainty. If the uncertainty reported is asymmetrical, the absolute value of the larger uncertainty is used. Uncertainty is not estimated for the following sources: CO<sub>2</sub> emissions from stationary combustion – geothermal energy; CO<sub>2</sub> emissions from mobile combustion by mode of transportation; CH<sub>4</sub> and N<sub>2</sub>O emissions from mobile combustion by mode of off-road transportation; and CH<sub>4</sub> from the incineration of waste. While CO<sub>2</sub> emissions from geothermal energy are included in the overall emissions estimate, they are not an official IPCC source category. As a result, there are no guidelines to associate uncertainty with the emissions estimate; therefore, an uncertainty analysis was not conducted. The uncertainty associated with CO<sub>2</sub> from mobile combustion is applied to each mode's emissions estimate, and the uncertainty associated with off-road vehicle CH<sub>4</sub> and N<sub>2</sub>O emissions are applied to both CH<sub>4</sub> and N<sub>2</sub>O emissions from

aviation, marine, and other sources. No uncertainty was associated with CH<sub>4</sub> emissions from waste incineration because emissions are less than 0.05 kt CH<sub>4</sub> and an uncertainty analysis was not conducted. When source and sink categories are sorted in decreasing order of this calculation, those that fall at the top of the list and cumulatively account for 90 percent of emissions are considered key categories. The key categories identified by the Approach 2 level assessment may differ from those identified by the Approach 1 assessment. The final set of key categories includes all source and sink categories identified as key by either the Approach 1 or the Approach 2 assessment, keeping in mind that the two assessments are not mutually exclusive.

It is important to note that a key category analysis can be sensitive to the definitions of the source and sink categories. If a large source category is split into many subcategories, then the subcategories may have contributions to the total inventory that are too small for those source categories to be considered key. Similarly, a collection of small, non-key source categories adding up to less than 5 percent of total emissions could become key source categories if those source categories were aggregated into a single source category. The United States has attempted to define source and sink categories by the conventions which would allow comparison with other international key categories, while still maintaining the category definitions that constitute how the emissions estimates were calculated for this report. As such, some of the category names used in the key category analysis may differ from the names used in the main body of the report. Additionally, the United States accounts for some source categories, including fossil fuel feedstocks, international bunkers, and emissions from U.S. Territories, that are derived from unique data sources using country-specific methodologies.

Table A-4 through Table A-7 contain the 1990 and 2014 level assessments for both with and without LULUCF sources and sinks, and contain further detail on where each source falls within the analysis. Approach 1 key categories are shaded dark gray. Additional key categories identified by the Approach 2 assessment are shaded light gray.

### Trend Assessment

Approach 1 for trend assessment is defined as the product of the source or sink category level assessment and the absolute difference between the source or sink category trend and the total trend. In turn, the source or sink category trend is defined as the change in emissions from the base year to the current year, as a percentage of current year emissions from that source or sink category. The total trend is the percentage change in total inventory emissions from the base year to the current year.

Thus, the source or sink category trend assessment will be large if the source or sink category represents a large percentage of emissions and/or has a trend that is quite different from the overall inventory trend. To determine key categories, the trend assessments are sorted in decreasing order, so that the source or sink categories with the highest trend assessments appear first. The trend assessments are summed until the threshold of 95 percent is reached; all categories that fall within that cumulative 95 percent are considered key categories.

For Approach 2, the trend assessment for each category from Approach 1 is multiplied by its percent relative uncertainty. If the uncertainty reported is asymmetrical, the larger uncertainty is used. When source and sink categories are sorted in decreasing order of this calculation, those that fall at the top of the list and cumulatively account for 90 percent of emissions are considered key categories. The key categories identified by the Approach 2 trend assessment may differ from those identified by the Approach 1 assessment. The final set of key categories includes all source and sink categories identified as key by either the Approach 1 or the Approach 2 assessment, keeping in mind that the two assessments are not mutually exclusive.

Table A-8 and Table A-9 contain the 1990 through 2014 trend assessment for both with and without LULUCF sources and sinks, and contain further detail on where each source falls within the analysis. Approach 1 key categories are shaded dark gray. Additional key categories identified by the Approach 2 assessment are shaded light gray.

**Table A-4: 1990 Key Source Category Approach 1 and Approach 2 Analysis—Level Assessment, without LULUCF**

| IPCC Source Categories   | Direct GHG       | Approach 1                                 |                     |                     | Approach 2               |                     |
|--|------------------|--|---------------------|---------------------|--------------------------|---------------------|
|  |                  | 1990 Estimate<br>(MMT CO <sub>2</sub> Eq.) | Level<br>Assessment | Cumulative<br>Total | Uncertainty <sup>a</sup> | Level<br>Assessment |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation | CO <sub>2</sub>  | 1,547.6                                    | 0.24                | 0.24                | 10%                      | 0.023               |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                               | CO <sub>2</sub>  | 1,188.9                                    | 0.19                | 0.43                | 6%                       | 0.012               |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial              | CO <sub>2</sub>  | 408.9                                      | 0.06                | 0.49                | 7%                       | 0.005               |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial              | CO <sub>2</sub>  | 278.3                                      | 0.04                | 0.54                | 19%                      | 0.008               |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                  | N <sub>2</sub> O | 245.0                                      | 0.04                | 0.57                | 24%                      | 0.009               |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential             | CO <sub>2</sub>  | 238.0                                      | 0.04                | 0.61                | 7%                       | 0.003               |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                   | CH <sub>4</sub>  | 206.8                                      | 0.03                | 0.64                | 30%                      | 0.010               |

|  |                  |       |       |      |      |        |
|--|------------------|-------|-------|------|------|--------|
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                               | CO <sub>2</sub>  | 187.4 | 0.03  | 0.67 | 6%   | 0.002  |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>  | 179.6 | 0.03  | 0.70 | 61%  | 0.017  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation      | CO <sub>2</sub>  | 175.3 | 0.03  | 0.73 | 5%   | 0.001  |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>  | 164.2 | 0.03  | 0.75 | 18%  | 0.005  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial                 | CO <sub>2</sub>  | 155.3 | 0.02  | 0.78 | 16%  | 0.004  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial                  | CO <sub>2</sub>  | 142.1 | 0.02  | 0.80 | 7%   | 0.002  |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                                   | CO <sub>2</sub>  | 118.1 | 0.02  | 0.82 | 42%  | 0.008  |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | 99.7  | 0.02  | 0.83 | 15%  | 0.002  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation      | CO <sub>2</sub>  | 97.5  | 0.02  | 0.85 | 8%   | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential                 | CO <sub>2</sub>  | 97.4  | 0.02  | 0.86 | 5%   | 0.001  |
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub>  | 96.5  | 0.02  | 0.88 | 15%  | 0.002  |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                                  | CO <sub>2</sub>  | 73.3  | 0.01  | 0.89 | 6%   | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial                  | CO <sub>2</sub>  | 63.3  | 0.01  | 0.90 | 5%   | 0.001  |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O | 58.2  | 0.01  | 0.91 | 139% | 0.013  |
| HFC-23 Emissions from HCFC-22 Production   | HFCs             | 46.1  | 0.01  | 0.92 | 10%  | 0.001  |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                                 | CO <sub>2</sub>  | 44.3  | 0.01  | 0.92 | 6%   | <0.001 |
| CH <sub>4</sub> Emissions from Petroleum Systems   | CH <sub>4</sub>  | 38.7  | 0.01  | 0.93 | 149% | 0.009  |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                       | CO <sub>2</sub>  | 37.7  | 0.01  | 0.94 | 30%  | 0.002  |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                                  | N <sub>2</sub> O | 37.7  | 0.01  | 0.94 | 27%  | 0.002  |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | 37.2  | 0.01  | 0.95 | 20%  | 0.001  |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 33.3  | 0.01  | 0.95 | 6%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories            | CO <sub>2</sub>  | 27.2  | <0.01 | 0.96 | 11%  | <0.001 |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | SF <sub>6</sub>  | 25.4  | <0.01 | 0.96 | 23%  | 0.001  |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 21.6  | <0.01 | 0.96 | 5%   | <0.001 |
| PFC Emissions from Aluminum Production   | PFCs             | 21.5  | <0.01 | 0.97 | 2%   | <0.001 |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>  | 15.7  | <0.01 | 0.97 | 39%  | 0.001  |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 15.2  | <0.01 | 0.97 | 4%   | <0.001 |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 14.0  | <0.01 | 0.97 | 24%  | 0.001  |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 13.1  | <0.01 | 0.98 | 17%  | <0.001 |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 13.0  | <0.01 | 0.98 | 8%   | <0.001 |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 12.1  | <0.01 | 0.98 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial                 | CO <sub>2</sub>  | 12.0  | <0.01 | 0.98 | 15%  | <0.001 |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 11.7  | <0.01 | 0.98 | 3%   | <0.001 |
| CO <sub>2</sub> Emissions from Incineration of Waste                                     | CO <sub>2</sub>  | 8.0   | <0.01 | 0.99 | 14%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | N <sub>2</sub> O | 7.4   | <0.01 | 0.99 | 173% | 0.002  |
| Fugitive Emissions from Abandoned Underground Coal Mines                                 | CH <sub>4</sub>  | 7.2   | <0.01 | 0.99 | 24%  | <0.001 |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 6.8   | <0.01 | 0.99 | 2%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  | 5.2   | <0.01 | 0.99 | 234% | 0.002  |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                                   | CH <sub>4</sub>  | 5.2   | <0.01 | 0.99 | 18%  | <0.001 |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | SF <sub>6</sub>  | 5.2   | <0.01 | 0.99 | 11%  | <0.001 |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  | 4.9   | <0.01 | 0.99 | 15%  | <0.001 |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2   | <0.01 | 0.99 | 24%  | <0.001 |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 3.8   | <0.01 | 0.99 | 12%  | <0.001 |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | Several          | 3.6   | <0.01 | 0.99 | 6%   | <0.001 |
| CO <sub>2</sub> Emissions from Petroleum Systems   | CO <sub>2</sub>  | 3.6   | <0.01 | 0.99 | 149% | 0.001  |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 3.4   | <0.01 | 1.00 | 108% | 0.001  |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | N <sub>2</sub> O | 3.1   | <0.01 | 1.00 | 222% | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential                | CO <sub>2</sub>  | 3.0   | <0.01 | 1.00 | NE   | <0.001 |

|  |                  |     |       |      |      |        |
|--|------------------|-----|-------|------|------|--------|
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8 | <0.01 | 1.00 | 7%   | <0.001 |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 2.2 | <0.01 | 1.00 | 12%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | CH <sub>4</sub>  | 1.8 | <0.01 | 1.00 | 50%  | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                              | N <sub>2</sub> O | 1.7 | <0.01 | 1.00 | 3%   | <0.001 |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.5 | <0.01 | 1.00 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.5 | <0.01 | 1.00 | 20%  | <0.001 |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 1.5 | <0.01 | 1.00 | 13%  | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                                 | N <sub>2</sub> O | 1.2 | <0.01 | 1.00 | 2%   | <0.001 |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.2 | <0.01 | 1.00 | 13%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | N <sub>2</sub> O | 1.0 | <0.01 | 1.00 | 206% | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | CH <sub>4</sub>  | 1.0 | <0.01 | 1.00 | 140% | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories           | CO <sub>2</sub>  | 0.6 | <0.01 | 1.00 | 19%  | <0.001 |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 0.6 | <0.01 | 1.00 | 21%  | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                                | N <sub>2</sub> O | 0.6 | <0.01 | 1.00 | 31%  | <0.001 |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5 | <0.01 | 1.00 | 16%  | <0.001 |
| N <sub>2</sub> O Emissions from Incineration of Waste                                    | N <sub>2</sub> O | 0.5 | <0.01 | 1.00 | 163% | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy                 | CO <sub>2</sub>  | 0.4 | <0.01 | 1.00 | NA   | <0.001 |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 0.4 | <0.01 | 1.00 | 50%  | <0.001 |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.4 | <0.01 | 1.00 | 9%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | N <sub>2</sub> O | 0.4 | <0.01 | 1.00 | 42%  | <0.001 |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 0.3 | <0.01 | 1.00 | 50%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                                  | CH <sub>4</sub>  | 0.3 | <0.01 | 1.00 | 2%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | CH <sub>4</sub>  | 0.3 | <0.01 | 1.00 | 22%  | <0.001 |
| Emissions from Substitutes for Ozone Depleting Substances                                | Several          | 0.3 | <0.01 | 1.00 | 10%  | <0.001 |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.2 | <0.01 | 1.00 | 40%  | <0.001 |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.2 | <0.01 | 1.00 | 55%  | <0.001 |
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues                   | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 29%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 200% | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                               | CH <sub>4</sub>  | 0.1 | <0.01 | 1.00 | 3%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 56%  | <0.001 |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | +   | <0.01 | 1.00 | 12%  | <0.001 |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 10%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                                 | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 8%   | <0.001 |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 19%  | <0.001 |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 12%  | <0.001 |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +   | <0.01 | 1.00 | 5%   | <0.001 |
| CH <sub>4</sub> Emissions from Incineration of Waste                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | NE   | <0.001 |
| HFC-134a Emissions from Magnesium Production and Processing                              | HFCs             | 0.0 | <0.01 | 1.00 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories            | CO <sub>2</sub>  | 0.0 | <0.01 | 1.00 | 17%  | <0.001 |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Note: LULUCF sources and sinks are not included in this analysis.

<sup>a</sup> Percent relative uncertainty. If the corresponding uncertainty is asymmetrical, the uncertainty given here is the larger and always positive.

NE (Not Estimated)

NA (Not Available)

**Table A-5: 1990 Key Source Category Approach 1 and Approach 2 Analysis—Level Assessment, with LULUCF**

| IPCC Source Categories   | Direct GHG       | 1990 Estimate             | Approach 1 | Cumulative | Uncertainty <sup>a</sup> | Approach 2 |
|--|------------------|---------------------------|------------|------------|--------------------------|------------|
|  |                  | (MMT CO <sub>2</sub> Eq.) | Level      |            |                          | Level      |
|  |                  |                           | Assessment | Total      |                          | Assessment |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation     | CO <sub>2</sub>  | 1,547.6                   | 0.21       | 0.21       | 10%                      | 0.020      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                                   | CO <sub>2</sub>  | 1,188.9                   | 0.16       | 0.37       | 6%                       | 0.010      |
| CO <sub>2</sub> Emissions from Forest Land Remaining Forest Land                         | CO <sub>2</sub>  | 723.5                     | 0.10       | 0.47       | 37%                      | 0.037      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial                  | CO <sub>2</sub>  | 408.9                     | 0.06       | 0.52       | 7%                       | 0.004      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial                  | CO <sub>2</sub>  | 278.3                     | 0.04       | 0.56       | 19%                      | 0.007      |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                      | N <sub>2</sub> O | 245.0                     | 0.03       | 0.60       | 24%                      | 0.008      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential                 | CO <sub>2</sub>  | 238.0                     | 0.03       | 0.63       | 7%                       | 0.002      |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                       | CH <sub>4</sub>  | 206.8                     | 0.03       | 0.66       | 30%                      | 0.008      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                               | CO <sub>2</sub>  | 187.4                     | 0.03       | 0.68       | 6%                       | 0.002      |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>  | 179.6                     | 0.02       | 0.71       | 61%                      | 0.015      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation      | CO <sub>2</sub>  | 175.3                     | 0.02       | 0.73       | 5%                       | 0.001      |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>  | 164.2                     | 0.02       | 0.75       | 18%                      | 0.004      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial                 | CO <sub>2</sub>  | 155.3                     | 0.02       | 0.77       | 16%                      | 0.003      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial                  | CO <sub>2</sub>  | 142.1                     | 0.02       | 0.79       | 7%                       | 0.001      |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                                   | CO <sub>2</sub>  | 118.1                     | 0.02       | 0.81       | 42%                      | 0.007      |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | 99.7                      | 0.01       | 0.82       | 15%                      | 0.002      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation      | CO <sub>2</sub>  | 97.5                      | 0.01       | 0.83       | 8%                       | 0.001      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential                 | CO <sub>2</sub>  | 97.4                      | 0.01       | 0.85       | 5%                       | 0.001      |
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub>  | 96.5                      | 0.01       | 0.86       | 15%                      | 0.002      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                                  | CO <sub>2</sub>  | 73.3                      | 0.01       | 0.87       | 6%                       | 0.001      |
| CO <sub>2</sub> Emissions from Land Converted to Cropland                                | CO <sub>2</sub>  | 65.7                      | 0.01       | 0.88       | 52%                      | 0.005      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial                  | CO <sub>2</sub>  | 63.3                      | 0.01       | 0.89       | 5%                       | <0.001     |
| CO <sub>2</sub> Emissions from Urban Trees   | CO <sub>2</sub>  | 60.4                      | 0.01       | 0.90       | 51%                      | 0.004      |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O | 58.2                      | 0.01       | 0.90       | 139%                     | 0.011      |
| HFC-23 Emissions from HCFC-22 Production   | HFCs             | 46.1                      | 0.01       | 0.91       | 10%                      | 0.001      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                                 | CO <sub>2</sub>  | 44.3                      | 0.01       | 0.92       | 6%                       | <0.001     |
| CO <sub>2</sub> Emissions from Land Converted to Grassland                               | CO <sub>2</sub>  | 39.1                      | 0.01       | 0.92       | 27%                      | 0.001      |
| CH <sub>4</sub> Emissions from Petroleum Systems   | CH <sub>4</sub>  | 38.7                      | 0.01       | 0.93       | 149%                     | 0.008      |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                       | CO <sub>2</sub>  | 37.7                      | 0.01       | 0.93       | 30%                      | 0.002      |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                                  | N <sub>2</sub> O | 37.7                      | 0.01       | 0.94       | 27%                      | 0.001      |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | 37.2                      | 0.01       | 0.94       | 20%                      | 0.001      |
| CO <sub>2</sub> Emissions from Cropland Remaining Cropland                               | CO <sub>2</sub>  | 34.3                      | <0.01      | 0.95       | 414%                     | 0.019      |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 33.3                      | <0.01      | 0.95       | 6%                       | <0.001     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories            | CO <sub>2</sub>  | 27.2                      | <0.01      | 0.96       | 11%                      | <0.001     |
| CO <sub>2</sub> Emissions from Landfilled Yard Trimmings and Food Scraps                 | CO <sub>2</sub>  | 26.0                      | <0.01      | 0.96       | 64%                      | 0.002      |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | SF <sub>6</sub>  | 25.4                      | <0.01      | 0.96       | 23%                      | 0.001      |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 21.6                      | <0.01      | 0.97       | 5%                       | <0.001     |
| PFC Emissions from Aluminum Production   | PFCs             | 21.5                      | <0.01      | 0.97       | 2%                       | <0.001     |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>  | 15.7                      | <0.01      | 0.97       | 39%                      | 0.001      |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 15.2                      | <0.01      | 0.97       | 4%                       | <0.001     |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 14.0                      | <0.01      | 0.97       | 24%                      | <0.001     |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 13.1                      | <0.01      | 0.98       | 17%                      | <0.001     |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 13.0                      | <0.01      | 0.98       | 8%                       | <0.001     |

|  |                  |      |       |      |       |        |
|--|------------------|------|-------|------|-------|--------|
| CO <sub>2</sub> Emissions from Grassland Remaining Grassland                             | CO <sub>2</sub>  | 12.9 | <0.01 | 0.98 | 1013% | 0.018  |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 12.1 | <0.01 | 0.98 | 5%    | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial                 | CO <sub>2</sub>  | 12.0 | <0.01 | 0.98 | 15%   | <0.001 |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 11.7 | <0.01 | 0.98 | 3%    | <0.001 |
| CO <sub>2</sub> Emissions from Incineration of Waste                                     | CO <sub>2</sub>  | 8.0  | <0.01 | 0.99 | 14%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | N <sub>2</sub> O | 7.4  | <0.01 | 0.99 | 173%  | 0.002  |
| Fugitive Emissions from Abandoned Underground Coal Mines                                 | CH <sub>4</sub>  | 7.2  | <0.01 | 0.99 | 24%   | <0.001 |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 6.8  | <0.01 | 0.99 | 2%    | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  | 5.2  | <0.01 | 0.99 | 234%  | 0.002  |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                                   | CH <sub>4</sub>  | 5.2  | <0.01 | 0.99 | 18%   | <0.001 |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | SF <sub>6</sub>  | 5.2  | <0.01 | 0.99 | 11%   | <0.001 |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  | 4.9  | <0.01 | 0.99 | 15%   | <0.001 |
| CO <sub>2</sub> Emissions from Liming  | CO <sub>2</sub>  | 4.7  | <0.01 | 0.99 | 111%  | 0.001  |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2  | <0.01 | 0.99 | 24%   | <0.001 |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 3.8  | <0.01 | 0.99 | 12%   | <0.001 |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | Several          | 3.6  | <0.01 | 0.99 | 6%    | <0.001 |
| CO <sub>2</sub> Emissions from Petroleum Systems   | CO <sub>2</sub>  | 3.6  | <0.01 | 0.99 | 149%  | 0.001  |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 3.4  | <0.01 | 0.99 | 108%  | <0.001 |
| CH <sub>4</sub> Emissions from Forest Fires  | CH <sub>4</sub>  | 3.3  | <0.01 | 0.99 | 174%  | 0.001  |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | N <sub>2</sub> O | 3.1  | <0.01 | 1.00 | 222%  | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential                | CO <sub>2</sub>  | 3.0  | <0.01 | 1.00 | NE    | <0.001 |
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8  | <0.01 | 1.00 | 7%    | <0.001 |
| CO <sub>2</sub> Emissions from Urea Fertilization  | CO <sub>2</sub>  | 2.4  | <0.01 | 1.00 | 42%   | <0.001 |
| N <sub>2</sub> O Emissions from Forest Fires   | N <sub>2</sub> O | 2.2  | <0.01 | 1.00 | 157%  | <0.001 |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 2.2  | <0.01 | 1.00 | 12%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | CH <sub>4</sub>  | 1.8  | <0.01 | 1.00 | 50%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                              | N <sub>2</sub> O | 1.7  | <0.01 | 1.00 | 3%    | <0.001 |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.5  | <0.01 | 1.00 | 5%    | <0.001 |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.5  | <0.01 | 1.00 | 20%   | <0.001 |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 1.5  | <0.01 | 1.00 | 13%   | <0.001 |
| N <sub>2</sub> O Emissions from Settlement Soils   | N <sub>2</sub> O | 1.4  | <0.01 | 1.00 | 268%  | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                                 | N <sub>2</sub> O | 1.2  | <0.01 | 1.00 | 2%    | <0.001 |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.2  | <0.01 | 1.00 | 13%   | <0.001 |
| CO <sub>2</sub> Emissions from Peatlands Remaining Peatlands                             | CO <sub>2</sub>  | 1.1  | <0.01 | 1.00 | 19%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | N <sub>2</sub> O | 1.0  | <0.01 | 1.00 | 206%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | CH <sub>4</sub>  | 1.0  | <0.01 | 1.00 | 140%  | <0.001 |
| CO <sub>2</sub> Emissions from Land Converted to Forest Land                             | CO <sub>2</sub>  | 0.7  | <0.01 | 1.00 | 70%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories           | CO <sub>2</sub>  | 0.6  | <0.01 | 1.00 | 19%   | <0.001 |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 0.6  | <0.01 | 1.00 | 21%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                                | N <sub>2</sub> O | 0.6  | <0.01 | 1.00 | 31%   | <0.001 |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5  | <0.01 | 1.00 | 16%   | <0.001 |
| N <sub>2</sub> O Emissions from Incineration of Waste                                    | N <sub>2</sub> O | 0.5  | <0.01 | 1.00 | 163%  | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy                 | CO <sub>2</sub>  | 0.4  | <0.01 | 1.00 | NA    | <0.001 |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 0.4  | <0.01 | 1.00 | 50%   | <0.001 |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.4  | <0.01 | 1.00 | 9%    | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | N <sub>2</sub> O | 0.4  | <0.01 | 1.00 | 42%   | <0.001 |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 0.3  | <0.01 | 1.00 | 50%   | <0.001 |

|  |                  |     |       |      |      |        |
|--|------------------|-----|-------|------|------|--------|
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                                  | CH <sub>4</sub>  | 0.3 | <0.01 | 1.00 | 2%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | CH <sub>4</sub>  | 0.3 | <0.01 | 1.00 | 22%  | <0.001 |
| Emissions from Substitutes for Ozone Depleting Substances                                | Several          | 0.3 | <0.01 | 1.00 | 10%  | <0.001 |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.2 | <0.01 | 1.00 | 40%  | <0.001 |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.2 | <0.01 | 1.00 | 55%  | <0.001 |
| N <sub>2</sub> O Emissions from Forest Soils   | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 318% | <0.001 |
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues                   | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 29%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 200% | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                               | CH <sub>4</sub>  | 0.1 | <0.01 | 1.00 | 3%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 56%  | <0.001 |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | +   | <0.01 | 1.00 | 12%  | <0.001 |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 10%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                                 | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 8%   | <0.001 |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 19%  | <0.001 |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 12%  | <0.001 |
| CH <sub>4</sub> Emissions from Peatlands Remaining Peatlands                             | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 62%  | <0.001 |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +   | <0.01 | 1.00 | 5%   | <0.001 |
| N <sub>2</sub> O Emissions from Peatlands Remaining Peatlands                            | N <sub>2</sub> O | +   | <0.01 | 1.00 | 61%  | <0.001 |
| CH <sub>4</sub> Emissions from Incineration of Waste                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | NE   | <0.001 |
| HFC-134a Emissions from Magnesium Production and Processing                              | HFCs             | 0.0 | <0.01 | 1.00 | 5%   | <0.001 |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> Percent relative uncertainty. If the corresponding uncertainty is asymmetrical, the uncertainty given here is the larger and always positive.

NE (Not Estimated)

NA (Not Available)

**Table A-6: 2014 Key Source Category Approach 1 and Approach 2 Analysis—Level Assessment, without LULUCF**

| IPCC Source Categories   | Direct GHG       | Approach 1                              |                  |                  | Approach 2 Level         |            |
|--|------------------|---|------------------|------------------|--------------------------|------------|
|  |                  | 2014 Estimate (MMT CO <sub>2</sub> Eq.) | Level Assessment | Cumulative Total | Uncertainty <sup>a</sup> | Assessment |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation | CO <sub>2</sub>  | 1,570.4                                 | 0.23             | 0.23             | 10%                      | 0.022      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                               | CO <sub>2</sub>  | 1,467.5                                 | 0.21             | 0.44             | 6%                       | 0.014      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial              | CO <sub>2</sub>  | 466.0                                   | 0.07             | 0.51             | 7%                       | 0.005      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation  | CO <sub>2</sub>  | 443.2                                   | 0.06             | 0.57             | 5%                       | 0.003      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential             | CO <sub>2</sub>  | 277.6                                   | 0.04             | 0.61             | 7%                       | 0.003      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial              | CO <sub>2</sub>  | 271.9                                   | 0.04             | 0.65             | 19%                      | 0.008      |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                  | N <sub>2</sub> O | 261.0                                   | 0.04             | 0.69             | 24%                      | 0.009      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial              | CO <sub>2</sub>  | 189.2                                   | 0.03             | 0.72             | 7%                       | 0.002      |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                   | CH <sub>4</sub>  | 176.1                                   | 0.03             | 0.75             | 30%                      | 0.008      |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                  | CH <sub>4</sub>  | 164.3                                   | 0.02             | 0.77             | 18%                      | 0.004      |
| Emissions from Substitutes for Ozone Depleting Substances                            | Several          | 161.2                                   | 0.02             | 0.79             | 10%                      | 0.002      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                           | CO <sub>2</sub>  | 150.1                                   | 0.02             | 0.81             | 6%                       | 0.001      |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>  | 148.0                                   | 0.02             | 0.84             | 61%                      | 0.013      |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                               | CO <sub>2</sub>  | 114.3                                   | 0.02             | 0.85             | 42%                      | 0.007      |

|  |                  |      |       |      |      |        |
|--|------------------|------|-------|------|------|--------|
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                                  | CO <sub>2</sub>  | 92.0 | 0.01  | 0.87 | 6%   | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial                 | CO <sub>2</sub>  | 75.3 | 0.01  | 0.88 | 16%  | 0.002  |
| CH <sub>4</sub> Emissions from Petroleum Systems   | CH <sub>4</sub>  | 68.1 | 0.01  | 0.89 | 149% | 0.015  |
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub>  | 67.6 | 0.01  | 0.90 | 15%  | 0.002  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential                 | CO <sub>2</sub>  | 67.5 | 0.01  | 0.91 | 5%   | 0.001  |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | 61.2 | 0.01  | 0.92 | 20%  | 0.002  |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O | 57.3 | 0.01  | 0.92 | 139% | 0.012  |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | 55.4 | 0.01  | 0.93 | 15%  | 0.001  |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                       | CO <sub>2</sub>  | 42.4 | 0.01  | 0.94 | 30%  | 0.002  |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 38.8 | 0.01  | 0.94 | 6%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial                  | CO <sub>2</sub>  | 38.2 | 0.01  | 0.95 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories            | CO <sub>2</sub>  | 34.6 | 0.01  | 0.95 | 11%  | 0.001  |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                                 | CO <sub>2</sub>  | 28.0 | <0.01 | 0.96 | 6%   | <0.001 |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 26.5 | <0.01 | 0.96 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation      | CO <sub>2</sub>  | 25.3 | <0.01 | 0.97 | 8%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | N <sub>2</sub> O | 19.6 | <0.01 | 0.97 | 173% | 0.005  |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 17.5 | <0.01 | 0.97 | 24%  | 0.001  |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>  | 14.7 | <0.01 | 0.97 | 39%  | 0.001  |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 14.1 | <0.01 | 0.98 | 3%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                                  | N <sub>2</sub> O | 12.5 | <0.01 | 0.98 | 27%  | <0.001 |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  | 12.1 | <0.01 | 0.98 | 15%  | <0.001 |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 11.9 | <0.01 | 0.98 | 17%  | <0.001 |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 10.9 | <0.01 | 0.98 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 9.4  | <0.01 | 0.98 | 8%   | <0.001 |
| CO <sub>2</sub> Emissions from Incineration of Waste                                     | CO <sub>2</sub>  | 9.4  | <0.01 | 0.99 | 14%  | <0.001 |
| Fugitive Emissions from Abandoned Underground Coal Mines                                 | CH <sub>4</sub>  | 6.3  | <0.01 | 0.99 | 24%  | <0.001 |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | SF <sub>6</sub>  | 5.6  | <0.01 | 0.99 | 23%  | <0.001 |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 5.4  | <0.01 | 0.99 | 4%   | <0.001 |
| HFC-23 Emissions from HCFC-22 Production   | HFCs             | 5.0  | <0.01 | 0.99 | 10%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  | 5.0  | <0.01 | 0.99 | 234% | 0.002  |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 4.8  | <0.01 | 0.99 | 108% | 0.001  |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | Several          | 4.5  | <0.01 | 0.99 | 6%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial                 | CO <sub>2</sub>  | 4.5  | <0.01 | 0.99 | 15%  | <0.001 |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 4.5  | <0.01 | 0.99 | 13%  | <0.001 |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2  | <0.01 | 0.99 | 24%  | <0.001 |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 4.0  | <0.01 | 0.99 | 12%  | <0.001 |
| CO <sub>2</sub> Emissions from Petroleum Systems   | CO <sub>2</sub>  | 3.6  | <0.01 | 0.99 | 149% | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories           | CO <sub>2</sub>  | 3.4  | <0.01 | 0.99 | 19%  | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories            | CO <sub>2</sub>  | 3.0  | <0.01 | 0.99 | 17%  | <0.001 |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 2.8  | <0.01 | 1.00 | 2%   | <0.001 |
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8  | <0.01 | 1.00 | 7%   | <0.001 |
| PFC Emissions from Aluminum Production   | PFCs             | 2.5  | <0.01 | 1.00 | 2%   | <0.001 |

|  |                  |     |       |      |      |        |
|--|------------------|-----|-------|------|------|--------|
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | N <sub>2</sub> O | 2.4 | <0.01 | 1.00 | 222% | 0.001  |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 2.1 | <0.01 | 1.00 | 50%  | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                                 | N <sub>2</sub> O | 2.0 | <0.01 | 1.00 | 2%   | <0.001 |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 1.9 | <0.01 | 1.00 | 12%  | <0.001 |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 1.8 | <0.01 | 1.00 | 50%  | <0.001 |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.8 | <0.01 | 1.00 | 13%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | CH <sub>4</sub>  | 1.5 | <0.01 | 1.00 | 50%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                                   | CH <sub>4</sub>  | 1.4 | <0.01 | 1.00 | 18%  | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                              | N <sub>2</sub> O | 1.4 | <0.01 | 1.00 | 3%   | <0.001 |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.3 | <0.01 | 1.00 | 5%   | <0.001 |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.1 | <0.01 | 1.00 | 20%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | CH <sub>4</sub>  | 1.1 | <0.01 | 1.00 | 140% | <0.001 |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | SF <sub>6</sub>  | 1.0 | <0.01 | 1.00 | 11%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | N <sub>2</sub> O | 1.0 | <0.01 | 1.00 | 206% | <0.001 |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 1.0 | <0.01 | 1.00 | 21%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                                  | CH <sub>4</sub>  | 0.5 | <0.01 | 1.00 | 2%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                                | N <sub>2</sub> O | 0.5 | <0.01 | 1.00 | 31%  | <0.001 |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5 | <0.01 | 1.00 | 16%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | CH <sub>4</sub>  | 0.4 | <0.01 | 1.00 | 22%  | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy                 | CO <sub>2</sub>  | 0.4 | <0.01 | 1.00 | NA   | <0.001 |
| N <sub>2</sub> O Emissions from Incineration of Waste                                    | N <sub>2</sub> O | 0.3 | <0.01 | 1.00 | 163% | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | N <sub>2</sub> O | 0.3 | <0.01 | 1.00 | 42%  | <0.001 |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.3 | <0.01 | 1.00 | 40%  | <0.001 |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | 0.2 | <0.01 | 1.00 | 12%  | <0.001 |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.2 | <0.01 | 1.00 | 9%   | <0.001 |
| HFC-134a Emissions from Magnesium Production and Processing                              | HFCs             | 0.1 | <0.01 | 1.00 | 5%   | <0.001 |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.1 | <0.01 | 1.00 | 55%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 200% | <0.001 |
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues                   | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 29%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | CH <sub>4</sub>  | 0.1 | <0.01 | 1.00 | 56%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                               | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 3%   | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                                 | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 8%   | <0.001 |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 12%  | <0.001 |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 10%  | <0.001 |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 19%  | <0.001 |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +   | <0.01 | 1.00 | 5%   | <0.001 |
| CH <sub>4</sub> Emissions from Incineration of Waste                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | NE   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential                | CO <sub>2</sub>  | 0.0 | <0.01 | 1.00 | NE   | <0.001 |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Note: LULUCF sources and sinks are not included in this analysis.

<sup>a</sup> Percent relative uncertainty. If the corresponding uncertainty is asymmetrical, the uncertainty given here is the larger and always positive.

NE - Not Estimated

NA - Not Available

**Table A-7: 2014 Key Source Category Approach 1 and Approach 2 Analysis—Level Assessment with LULUCF**

| IPCC Source Categories   | Direct GHG (MMT CO <sub>2</sub> Eq.) | Approach 1    |            |            | Approach 2               |            |
|--|--------------------------------------|---------------|------------|------------|--------------------------|------------|
|  |                                      | 2014 Estimate | Level      | Cumulative | Level                    | Assessment |
|  |                                      |               | Assessment | Total      | Uncertainty <sup>a</sup> | Assessment |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation     | CO <sub>2</sub>                      | 1,570.4       | 0.20       | 0.20       | 10%                      | 0.019      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                                   | CO <sub>2</sub>                      | 1,467.5       | 0.19       | 0.39       | 6%                       | 0.012      |
| CO <sub>2</sub> Emissions from Forest Land Remaining Forest Land                         | CO <sub>2</sub>                      | 742.3         | 0.09       | 0.48       | 37%                      | 0.035      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial                  | CO <sub>2</sub>                      | 466.0         | 0.06       | 0.54       | 7%                       | 0.004      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation      | CO <sub>2</sub>                      | 443.2         | 0.06       | 0.60       | 5%                       | 0.003      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential                 | CO <sub>2</sub>                      | 277.6         | 0.04       | 0.64       | 7%                       | 0.002      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial                  | CO <sub>2</sub>                      | 271.9         | 0.03       | 0.67       | 19%                      | 0.007      |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                      | N <sub>2</sub> O                     | 261.0         | 0.03       | 0.70       | 24%                      | 0.008      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial                  | CO <sub>2</sub>                      | 189.2         | 0.02       | 0.73       | 7%                       | 0.002      |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                       | CH <sub>4</sub>                      | 176.1         | 0.02       | 0.75       | 30%                      | 0.007      |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>                      | 164.3         | 0.02       | 0.77       | 18%                      | 0.004      |
| Emissions from Substitutes for Ozone Depleting Substances                                | Several                              | 161.2         | 0.02       | 0.79       | 10%                      | 0.002      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                               | CO <sub>2</sub>                      | 150.1         | 0.02       | 0.81       | 6%                       | 0.001      |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>                      | 148.0         | 0.02       | 0.83       | 61%                      | 0.012      |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                                   | CO <sub>2</sub>                      | 114.3         | 0.01       | 0.84       | 42%                      | 0.006      |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                                  | CO <sub>2</sub>                      | 92.0          | 0.01       | 0.86       | 6%                       | 0.001      |
| CO <sub>2</sub> Emissions from Urban Trees   | CO <sub>2</sub>                      | 90.6          | 0.01       | 0.87       | 51%                      | 0.006      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial                 | CO <sub>2</sub>                      | 75.3          | 0.01       | 0.88       | 16%                      | 0.002      |
| CH <sub>4</sub> Emissions from Petroleum Systems   | CH <sub>4</sub>                      | 68.1          | 0.01       | 0.89       | 149%                     | 0.013      |
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub>                      | 67.6          | 0.01       | 0.90       | 15%                      | 0.001      |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential                 | CO <sub>2</sub>                      | 67.5          | 0.01       | 0.90       | 5%                       | <0.001     |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>                      | 61.2          | 0.01       | 0.91       | 20%                      | 0.002      |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O                     | 57.3          | 0.01       | 0.92       | 139%                     | 0.010      |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>                      | 55.4          | 0.01       | 0.93       | 15%                      | 0.001      |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                       | CO <sub>2</sub>                      | 42.4          | 0.01       | 0.93       | 30%                      | 0.002      |
| CO <sub>2</sub> Emissions from Land Converted to Grassland                               | CO <sub>2</sub>                      | 40.4          | 0.01       | 0.94       | 27%                      | 0.001      |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>                      | 38.8          | <0.01      | 0.94       | 6%                       | <0.001     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial                  | CO <sub>2</sub>                      | 38.2          | <0.01      | 0.95       | 5%                       | <0.001     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories            | CO <sub>2</sub>                      | 34.6          | <0.01      | 0.95       | 11%                      | <0.001     |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                                 | CO <sub>2</sub>                      | 28.0          | <0.01      | 0.95       | 6%                       | <0.001     |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>                      | 26.5          | <0.01      | 0.96       | 5%                       | <0.001     |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation      | CO <sub>2</sub>                      | 25.3          | <0.01      | 0.96       | 8%                       | <0.001     |
| CO <sub>2</sub> Emissions from Land Converted to Cropland                                | CO <sub>2</sub>                      | 22.1          | <0.01      | 0.96       | 52%                      | 0.001      |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | N <sub>2</sub> O                     | 19.6          | <0.01      | 0.97       | 173%                     | 0.004      |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O                     | 17.5          | <0.01      | 0.97       | 24%                      | 0.001      |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>                      | 14.7          | <0.01      | 0.97       | 39%                      | 0.001      |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>                      | 14.1          | <0.01      | 0.97       | 3%                       | <0.001     |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                                  | N <sub>2</sub> O                     | 12.5          | <0.01      | 0.97       | 27%                      | <0.001     |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>                      | 12.1          | <0.01      | 0.98       | 15%                      | <0.001     |

|  |                  |      |       |      |       |        |
|--|------------------|------|-------|------|-------|--------|
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 11.9 | <0.01 | 0.98 | 17%   | <0.001 |
| CO <sub>2</sub> Emissions from Landfilled Yard Trimmings and Food Scraps                 | CO <sub>2</sub>  | 11.6 | <0.01 | 0.98 | 64%   | 0.001  |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 10.9 | <0.01 | 0.98 | 5%    | <0.001 |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 9.4  | <0.01 | 0.98 | 8%    | <0.001 |
| CO <sub>2</sub> Emissions from Incineration of Waste                                     | CO <sub>2</sub>  | 9.4  | <0.01 | 0.98 | 14%   | <0.001 |
| CO <sub>2</sub> Emissions from Cropland Remaining Cropland                               | CO <sub>2</sub>  | 8.4  | <0.01 | 0.98 | 414%  | 0.004  |
| CH <sub>4</sub> Emissions from Forest Fires  | CH <sub>4</sub>  | 7.3  | <0.01 | 0.98 | 174%  | 0.002  |
| Fugitive Emissions from Abandoned Underground Coal Mines                                 | CH <sub>4</sub>  | 6.3  | <0.01 | 0.99 | 24%   | <0.001 |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | SF <sub>6</sub>  | 5.6  | <0.01 | 0.99 | 23%   | <0.001 |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 5.4  | <0.01 | 0.99 | 4%    | <0.001 |
| HFC-23 Emissions from HCFC-22 Production   | HFCs             | 5.0  | <0.01 | 0.99 | 10%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  | 5.0  | <0.01 | 0.99 | 234%  | 0.001  |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 4.8  | <0.01 | 0.99 | 108%  | 0.001  |
| N <sub>2</sub> O Emissions from Forest Fires   | N <sub>2</sub> O | 4.8  | <0.01 | 0.99 | 157%  | 0.001  |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | Several          | 4.5  | <0.01 | 0.99 | 6%    | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial                 | CO <sub>2</sub>  | 4.5  | <0.01 | 0.99 | 15%   | <0.001 |
| CO <sub>2</sub> Emissions from Urea Fertilization  | CO <sub>2</sub>  | 4.5  | <0.01 | 0.99 | 42%   | <0.001 |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 4.5  | <0.01 | 0.99 | 13%   | <0.001 |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2  | <0.01 | 0.99 | 24%   | <0.001 |
| CO <sub>2</sub> Emissions from Liming  | CO <sub>2</sub>  | 4.1  | <0.01 | 0.99 | 111%  | 0.001  |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 4.0  | <0.01 | 0.99 | 12%   | <0.001 |
| CO <sub>2</sub> Emissions from Grassland Remaining Grassland                             | CO <sub>2</sub>  | 3.8  | <0.01 | 0.99 | 1013% | 0.005  |
| CO <sub>2</sub> Emissions from Petroleum Systems   | CO <sub>2</sub>  | 3.6  | <0.01 | 0.99 | 149%  | 0.001  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories           | CO <sub>2</sub>  | 3.4  | <0.01 | 0.99 | 19%   | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories            | CO <sub>2</sub>  | 3.0  | <0.01 | 0.99 | 17%   | <0.001 |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 2.8  | <0.01 | 1.00 | 2%    | <0.001 |
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8  | <0.01 | 1.00 | 7%    | <0.001 |
| PFC Emissions from Aluminum Production   | PFCs             | 2.5  | <0.01 | 1.00 | 2%    | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | N <sub>2</sub> O | 2.4  | <0.01 | 1.00 | 222%  | 0.001  |
| N <sub>2</sub> O Emissions from Settlement Soils   | N <sub>2</sub> O | 2.4  | <0.01 | 1.00 | 268%  | 0.001  |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 2.1  | <0.01 | 1.00 | 50%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                                 | N <sub>2</sub> O | 2.0  | <0.01 | 1.00 | 2%    | <0.001 |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 1.9  | <0.01 | 1.00 | 12%   | <0.001 |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 1.8  | <0.01 | 1.00 | 50%   | <0.001 |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.8  | <0.01 | 1.00 | 13%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | CH <sub>4</sub>  | 1.5  | <0.01 | 1.00 | 50%   | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                                   | CH <sub>4</sub>  | 1.4  | <0.01 | 1.00 | 18%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                              | N <sub>2</sub> O | 1.4  | <0.01 | 1.00 | 3%    | <0.001 |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.3  | <0.01 | 1.00 | 5%    | <0.001 |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.1  | <0.01 | 1.00 | 20%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | CH <sub>4</sub>  | 1.1  | <0.01 | 1.00 | 140%  | <0.001 |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | SF <sub>6</sub>  | 1.0  | <0.01 | 1.00 | 11%   | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | N <sub>2</sub> O | 1.0  | <0.01 | 1.00 | 206%  | <0.001 |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 1.0  | <0.01 | 1.00 | 21%   | <0.001 |

|  |                  |     |       |      |      |        |
|--|------------------|-----|-------|------|------|--------|
| CO <sub>2</sub> Emissions from Peatlands Remaining Peatlands                             | CO <sub>2</sub>  | 0.8 | <0.01 | 1.00 | 19%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                                  | CH <sub>4</sub>  | 0.5 | <0.01 | 1.00 | 2%   | <0.001 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                                | N <sub>2</sub> O | 0.5 | <0.01 | 1.00 | 31%  | <0.001 |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5 | <0.01 | 1.00 | 16%  | <0.001 |
| N <sub>2</sub> O Emissions from Forest Soils   | N <sub>2</sub> O | 0.5 | <0.01 | 1.00 | 318% | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | CH <sub>4</sub>  | 0.4 | <0.01 | 1.00 | 22%  | <0.001 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy                 | CO <sub>2</sub>  | 0.4 | <0.01 | 1.00 | NA   | <0.001 |
| CO <sub>2</sub> Emissions from Land Converted to Forest Land                             | CO <sub>2</sub>  | 0.3 | <0.01 | 1.00 | 70%  | <0.001 |
| N <sub>2</sub> O Emissions from Incineration of Waste                                    | N <sub>2</sub> O | 0.3 | <0.01 | 1.00 | 163% | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | N <sub>2</sub> O | 0.3 | <0.01 | 1.00 | 42%  | <0.001 |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.3 | <0.01 | 1.00 | 40%  | <0.001 |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | 0.2 | <0.01 | 1.00 | 12%  | <0.001 |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.2 | <0.01 | 1.00 | 9%   | <0.001 |
| HFC-134a Emissions from Magnesium Production and Processing                              | HFCs             | 0.1 | <0.01 | 1.00 | 5%   | <0.001 |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.1 | <0.01 | 1.00 | 55%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 200% | <0.001 |
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues                   | N <sub>2</sub> O | 0.1 | <0.01 | 1.00 | 29%  | <0.001 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | CH <sub>4</sub>  | 0.1 | <0.01 | 1.00 | 56%  | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                               | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 3%   | <0.001 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                                 | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 8%   | <0.001 |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 12%  | <0.001 |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 10%  | <0.001 |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 19%  | <0.001 |
| CH <sub>4</sub> Emissions from Peatlands Remaining Peatlands                             | CH <sub>4</sub>  | +   | <0.01 | 1.00 | 62%  | <0.001 |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +   | <0.01 | 1.00 | 5%   | <0.001 |
| N <sub>2</sub> O Emissions from Peatlands Remaining Peatlands                            | N <sub>2</sub> O | +   | <0.01 | 1.00 | 61%  | <0.001 |
| CH <sub>4</sub> Emissions from Incineration of Waste                                     | CH <sub>4</sub>  | +   | <0.01 | 1.00 | NE   | <0.001 |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> Percent relative uncertainty. If the corresponding uncertainty is asymmetrical, the uncertainty given here is the larger and always positive.

NE (Not Estimated)

NA (Not Available)

**Table A-8: 1990-2014 Key Source Category Approach 1 and 2 Analysis—Trend Assessment, without LULUCF**

| IPCC Source Categories   | Direct GHG      | 1990 Estimate             |                           | 2014 Estimate    |                  | Approach 1            | Approach 2 | % | % Cumulative Total |
|--|-----------------|---------------------------|---------------------------|------------------|------------------|-----------------------|------------|---|--------------------|
|  |                 | (MMT CO <sub>2</sub> Eq.) | (MMT CO <sub>2</sub> Eq.) | Trend Assessment | Trend Assessment | Contribution to Trend |            |   |                    |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation                              | CO <sub>2</sub> | 175.3                     | 443.2                     | 0.04             | 0.002            | 15.8                  | 16         |   |                    |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road Emissions from Substitutes for Ozone Depleting Substances | CO <sub>2</sub> | 1,188.9                   | 1,467.5                   | 0.03             | 0.002            | 11.8                  | 28         |   |                    |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation                             | Several         | 0.3                       | 161.2                     | 0.03             | 0.002            | 10.0                  | 38         |   |                    |
|  | CO <sub>2</sub> | 1,547.6                   | 1,570.4                   | 0.01             | 0.001            | 5.7                   | 43         |   |                    |

|  |                  |       |       |       |        |     |    |
|--|------------------|-------|-------|-------|--------|-----|----|
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial                 | CO <sub>2</sub>  | 155.3 | 75.3  | 0.01  | 0.002  | 5.7 | 49 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation      | CO <sub>2</sub>  | 97.5  | 25.3  | 0.01  | 0.001  | 4.9 | 54 |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub>  | 99.7  | 55.4  | 0.01  | 0.001  | 3.2 | 57 |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                               | CO <sub>2</sub>  | 187.4 | 150.1 | 0.01  | <0.001 | 3.2 | 60 |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                       | CH <sub>4</sub>  | 206.8 | 176.1 | 0.01  | 0.002  | 2.9 | 63 |
| CH <sub>4</sub> Emissions from Landfills   | CH <sub>4</sub>  | 179.6 | 148.0 | 0.01  | 0.004  | 2.8 | 66 |
| HFC-23 Emissions from HCFC-22 Production   | HFCs             | 46.1  | 5.0   | 0.01  | 0.001  | 2.8 | 69 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential                 | CO <sub>2</sub>  | 97.4  | 67.5  | 0.01  | <0.001 | 2.3 | 71 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial                  | CO <sub>2</sub>  | 142.1 | 189.2 | 0.01  | <0.001 | 2.3 | 73 |
| Fugitive Emissions from Coal Mining  | CH <sub>4</sub>  | 96.5  | 67.6  | 0.01  | 0.001  | 2.2 | 76 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial                  | CO <sub>2</sub>  | 63.3  | 38.2  | <0.01 | <0.001 | 1.8 | 77 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                                  | N <sub>2</sub> O | 37.7  | 12.5  | <0.01 | 0.001  | 1.7 | 79 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial                  | CO <sub>2</sub>  | 278.3 | 271.9 | <0.01 | 0.001  | 1.7 | 81 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial                  | CO <sub>2</sub>  | 408.9 | 466.0 | <0.01 | <0.001 | 1.7 | 82 |
| CH <sub>4</sub> Emissions from Petroleum Systems   | CH <sub>4</sub>  | 38.7  | 68.1  | <0.01 | 0.006  | 1.6 | 84 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential                 | CO <sub>2</sub>  | 238.0 | 277.6 | <0.01 | <0.001 | 1.4 | 85 |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution                  | SF <sub>6</sub>  | 25.4  | 5.6   | <0.01 | 0.001  | 1.3 | 87 |
| CH <sub>4</sub> Emissions from Manure Management   | CH <sub>4</sub>  | 37.2  | 61.2  | <0.01 | 0.001  | 1.3 | 88 |
| PFC Emissions from Aluminum Production   | PFCs             | 21.5  | 2.5   | <0.01 | <0.001 | 1.3 | 89 |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                                 | CO <sub>2</sub>  | 44.3  | 28.0  | <0.01 | <0.001 | 1.2 | 91 |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                                  | CO <sub>2</sub>  | 73.3  | 92.0  | <0.01 | <0.001 | 0.8 | 91 |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                                   | CO <sub>2</sub>  | 118.1 | 114.3 | <0.01 | 0.001  | 0.8 | 92 |
| CH <sub>4</sub> Emissions from Enteric Fermentation                                      | CH <sub>4</sub>  | 164.2 | 164.3 | <0.01 | <0.001 | 0.7 | 93 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | N <sub>2</sub> O | 7.4   | 19.6  | <0.01 | 0.003  | 0.7 | 94 |
| N <sub>2</sub> O Emissions from Adipic Acid Production                                   | N <sub>2</sub> O | 15.2  | 5.4   | <0.01 | <0.001 | 0.7 | 94 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial                 | CO <sub>2</sub>  | 12.0  | 4.5   | <0.01 | <0.001 | 0.5 | 95 |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                          | CO <sub>2</sub>  | 4.9   | 12.1  | <0.01 | <0.001 | 0.4 | 95 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories            | CO <sub>2</sub>  | 27.2  | 34.6  | <0.01 | <0.001 | 0.3 | 96 |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                                | N <sub>2</sub> O | 58.2  | 57.3  | <0.01 | 0.001  | 0.3 | 96 |
| CO <sub>2</sub> Emissions from Ammonia Production  | CO <sub>2</sub>  | 13.0  | 9.4   | <0.01 | <0.001 | 0.3 | 96 |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                       | SF <sub>6</sub>  | 5.2   | 1.0   | <0.01 | <0.001 | 0.3 | 97 |
| CO <sub>2</sub> Emissions from Aluminum Production                                       | CO <sub>2</sub>  | 6.8   | 2.8   | <0.01 | <0.001 | 0.3 | 97 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                                   | CH <sub>4</sub>  | 5.2   | 1.4   | <0.01 | <0.001 | 0.3 | 97 |
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 21.6  | 26.5  | <0.01 | <0.001 | 0.2 | 97 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential                | CO <sub>2</sub>  | 3.0   | 0.0   | <0.01 | 0.001  | 0.2 | 97 |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 33.3  | 38.8  | <0.01 | <0.001 | 0.2 | 98 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories            | CO <sub>2</sub>  | 0.0   | 3.0   | <0.01 | <0.001 | 0.2 | 98 |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 1.5   | 4.5   | <0.01 | <0.001 | 0.2 | 98 |

|  |                  |       |       |       |        |      |     |
|--|------------------|-------|-------|-------|--------|------|-----|
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories           | CO <sub>2</sub>  | 0.6   | 3.4   | <0.01 | <0.001 | 0.2  | 98  |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 14.0  | 17.5  | <0.01 | <0.001 | 0.2  | 98  |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 13.1  | 11.9  | <0.01 | <0.001 | 0.1  | 98  |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                      | N <sub>2</sub> O | 245.0 | 261.0 | <0.01 | <0.001 | 0.1  | 99  |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>  | 15.7  | 14.7  | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 12.1  | 10.9  | <0.01 | <0.001 | 0.1  | 99  |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                       | CO <sub>2</sub>  | 37.7  | 42.4  | <0.01 | <0.001 | 0.1  | 99  |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 0.4   | 2.1   | <0.01 | <0.001 | 0.1  | 99  |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 11.7  | 14.1  | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 0.3   | 1.8   | <0.01 | <0.001 | 0.1  | 99  |
| Fugitive Emissions from Abandoned Underground Coal Mines                                 | CH <sub>4</sub>  | 7.2   | 6.3   | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 3.4   | 4.8   | <0.01 | <0.001 | 0.1  | 99  |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | N <sub>2</sub> O | 3.1   | 2.4   | <0.01 | <0.001 | 0.1  | 99  |
| CO <sub>2</sub> Emissions from Incineration of Waste                                     | CO <sub>2</sub>  | 8.0   | 9.4   | <0.01 | <0.001 | 0.1  | 100 |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | Several          | 3.6   | 4.5   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                                 | N <sub>2</sub> O | 1.2   | 2.0   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  | 5.2   | 5.0   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.5   | 1.1   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                              | N <sub>2</sub> O | 1.7   | 1.4   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.2   | 1.8   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | CH <sub>4</sub>  | 1.8   | 1.5   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 2.2   | 1.9   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2   | 4.2   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.5   | 1.3   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 0.6   | 1.0   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Petroleum Systems   | CO <sub>2</sub>  | 3.6   | 3.6   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Silicon Carbide Production and Consumption                | CO <sub>2</sub>  | 0.4   | 0.2   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Soda Ash Production and Consumption                       | CO <sub>2</sub>  | 2.8   | 2.8   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Incineration of Waste                                    | N <sub>2</sub> O | 0.5   | 0.3   | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Other                                  | CH <sub>4</sub>  | 0.3   | 0.5   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Semiconductor Manufacture                                | N <sub>2</sub> O | +     | 0.2   | <0.01 | <0.001 | <0.1 | 100 |
| HFC-134a Emissions from Magnesium Production and Processing                              | HFCs             | 0.0   | 0.1   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | N <sub>2</sub> O | 1.0   | 1.0   | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Petrochemical Production                                  | CH <sub>4</sub>  | 0.2   | 0.1   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Marine                                | N <sub>2</sub> O | 0.6   | 0.5   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | N <sub>2</sub> O | 0.4   | 0.3   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation        | CH <sub>4</sub>  | 0.3   | 0.4   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Urea Consumption for Non-Ag Purposes                      | CO <sub>2</sub>  | 3.8   | 4.0   | <0.01 | <0.001 | <0.1 | 100 |

|  |                  |     |     |       |        |      |     |
|--|------------------|-----|-----|-------|--------|------|-----|
| CH <sub>4</sub> Emissions from Mobile Combustion: Aviation                               | CH <sub>4</sub>  | 0.1 | +   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5 | 0.5 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | N <sub>2</sub> O | 0.1 | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Field Burning of Agricultural Residues                    | CH <sub>4</sub>  | 0.2 | 0.3 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Silicon Carbide Production and Consumption                | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Geothermal Energy                 | CO <sub>2</sub>  | 0.4 | 0.4 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Commercial                    | CH <sub>4</sub>  | 1.0 | 1.1 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Field Burning of Agricultural Residues                   | N <sub>2</sub> O | 0.1 | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - U.S. Territories              | CH <sub>4</sub>  | +   | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Ferroalloy Production                                     | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Marine                                 | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Magnesium Production and Processing                       | CO <sub>2</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Incineration of Waste                                     | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Note: LULUCF sources and sinks are not included in this analysis.

**Table A-9: 1990-2014 Key Source Category Approach 1 and 2 Analysis—Trend Assessment, with LULUCF**

| IPCC Source Categories   | Direct GHG      | 1990 Estimate<br>(MM TCO <sub>2</sub> Eq.) | 2014 Estimate<br>(MMT CO <sub>2</sub> Eq.) | Approach 1          | Approach 2          | Percent<br>Contribution to<br>Trend (%) | Cumulative<br>Contribution to<br>Trend (%) |
|--|-----------------|--|--|---------------------|---------------------|---|--|
|  |                 |  |  | Trend<br>Assessment | Trend<br>Assessment |   |  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation      | CO <sub>2</sub> | 175.3                                      | 443.2                                      | 0.03                | 0.002               | 14.6                                    | 15   |
| CO <sub>2</sub> Emissions from Mobile Combustion: Road                                   | CO <sub>2</sub> | 1,188.9                                    | 1,467.5                                    | 0.03                | 0.002               | 11.8                                    | 26   |
| Emissions from Substitutes for Ozone Depleting Substances                                | Several         | 0.3  | 161.2                                      | 0.02                | 0.002               | 9.2                                     | 36   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Industrial                 | CO <sub>2</sub> | 155.3                                      | 75.3                                       | 0.01                | 0.002               | 5.1                                     | 41   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation      | CO <sub>2</sub> | 97.5                                       | 25.3                                       | 0.01                | 0.001               | 4.4                                     | 45   |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation     | CO <sub>2</sub> | 1,547.6                                    | 1,570.4                                    | 0.01                | 0.001               | 3.9                                     | 49   |
| CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub> | 99.7                                       | 55.4                                       | 0.01                | 0.001               | 2.9                                     | 52   |
| CO <sub>2</sub> Emissions from Mobile Combustion: Aviation                               | CO <sub>2</sub> | 187.4                                      | 150.1                                      | 0.01                | <0.001              | 2.8                                     | 55   |
| CO <sub>2</sub> Emissions from Land Converted to Cropland                                | CO <sub>2</sub> | 65.7                                       | 22.1                                       | 0.01                | 0.003               | 2.7                                     | 57   |
| HFC-23 Emissions from HCFC-22 Production   | HFCs            | 46.1                                       | 5.0  | 0.01                | 0.001               | 2.5                                     | 60   |
| CH <sub>4</sub> Emissions from Natural Gas Systems                                       | CH <sub>4</sub> | 206.8                                      | 176.1                                      | 0.01                | 0.002               | 2.4                                     | 62   |

|   |                  |       |       |       |        |     |    |
|---|------------------|-------|-------|-------|--------|-----|----|
| CH <sub>4</sub> Emissions from Landfills  | CH <sub>4</sub>  | 179.6 | 148.0 | 0.01  | 0.004  | 2.4 | 65 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial           | CO <sub>2</sub>  | 142.1 | 189.2 | 0.01  | <0.001 | 2.2 | 67 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential          | CO <sub>2</sub>  | 97.4  | 67.5  | <0.01 | <0.001 | 2.0 | 69 |
| Fugitive Emissions from Coal Mining   | CH <sub>4</sub>  | 96.5  | 67.6  | <0.01 | 0.001  | 2.0 | 71 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial           | CO <sub>2</sub>  | 408.9 | 466.0 | <0.01 | <0.001 | 1.9 | 73 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial           | CO <sub>2</sub>  | 63.3  | 38.2  | <0.01 | <0.001 | 1.6 | 74 |
| CO <sub>2</sub> Emissions from Cropland Remaining Cropland                        | CO <sub>2</sub>  | 34.3  | 8.4   | <0.01 | 0.016  | 1.6 | 76 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Road                           | N <sub>2</sub> O | 37.7  | 12.5  | <0.01 | 0.001  | 1.6 | 78 |
| CH <sub>4</sub> Emissions from Petroleum Systems                                  | CH <sub>4</sub>  | 38.7  | 68.1  | <0.01 | 0.005  | 1.5 | 79 |
| CO <sub>2</sub> Emissions from Urban Trees  | CO <sub>2</sub>  | 60.4  | 90.6  | <0.01 | 0.002  | 1.5 | 81 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Residential          | CO <sub>2</sub>  | 238.0 | 277.6 | <0.01 | <0.001 | 1.4 | 82 |
| CO <sub>2</sub> Emissions from Forest Land Remaining Forest Land                  | CO <sub>2</sub>  | 723.5 | 742.3 | <0.01 | 0.001  | 1.4 | 83 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Industrial           | CO <sub>2</sub>  | 278.3 | 271.9 | <0.01 | 0.001  | 1.3 | 85 |
| CH <sub>4</sub> Emissions from Manure Management                                  | CH <sub>4</sub>  | 37.2  | 61.2  | <0.01 | 0.001  | 1.2 | 86 |
| SF <sub>6</sub> Emissions from Electrical Transmission and Distribution           | SF <sub>6</sub>  | 25.4  | 5.6   | <0.01 | 0.001  | 1.2 | 87 |
| PFC Emissions from Aluminum Production  | PFCs             | 21.5  | 2.5   | <0.01 | <0.001 | 1.1 | 88 |
| CO <sub>2</sub> Emissions from Mobile Combustion: Marine                          | CO <sub>2</sub>  | 44.3  | 28.0  | <0.01 | <0.001 | 1.1 | 89 |
| CO <sub>2</sub> Emissions from Landfilled Yard Trimmings and Food Scraps          | CO <sub>2</sub>  | 26.0  | 11.6  | <0.01 | 0.001  | 0.9 | 90 |
| CO <sub>2</sub> Emissions from Mobile Combustion: Other                           | CO <sub>2</sub>  | 73.3  | 92.0  | <0.01 | <0.001 | 0.8 | 91 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation | N <sub>2</sub> O | 7.4   | 19.6  | <0.01 | 0.003  | 0.7 | 92 |
| CO <sub>2</sub> Emissions from Non-Energy Use of Fuels                            | CO <sub>2</sub>  | 118.1 | 114.3 | <0.01 | 0.001  | 0.6 | 92 |
| N <sub>2</sub> O Emissions from Adipic Acid Production                            | N <sub>2</sub> O | 15.2  | 5.4   | <0.01 | <0.001 | 0.6 | 93 |
| CO <sub>2</sub> Emissions from Grassland Remaining Grassland                      | CO <sub>2</sub>  | 12.9  | 3.8   | <0.01 | 0.014  | 0.6 | 94 |
| CH <sub>4</sub> Emissions from Enteric Fermentation                               | CH <sub>4</sub>  | 164.2 | 164.3 | <0.01 | <0.001 | 0.5 | 94 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial          | CO <sub>2</sub>  | 12.0  | 4.5   | <0.01 | <0.001 | 0.5 | 95 |
| CO <sub>2</sub> Emissions from Other Process Uses of Carbonates                   | CO <sub>2</sub>  | 4.9   | 12.1  | <0.01 | <0.001 | 0.4 | 95 |
| CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories     | CO <sub>2</sub>  | 27.2  | 34.6  | <0.01 | <0.001 | 0.3 | 95 |
| SF <sub>6</sub> Emissions from Magnesium Production and Processing                | SF <sub>6</sub>  | 5.2   | 1.0   | <0.01 | <0.001 | 0.3 | 96 |
| CO <sub>2</sub> Emissions from Aluminum Production                                | CO <sub>2</sub>  | 6.8   | 2.8   | <0.01 | <0.001 | 0.3 | 96 |
| CO <sub>2</sub> Emissions from Ammonia Production                                 | CO <sub>2</sub>  | 13.0  | 9.4   | <0.01 | <0.001 | 0.2 | 96 |
| Indirect N <sub>2</sub> O Emissions from Applied Nitrogen                         | N <sub>2</sub> O | 58.2  | 57.3  | <0.01 | 0.001  | 0.2 | 96 |
| CH <sub>4</sub> Emissions from Mobile Combustion: Road                            | CH <sub>4</sub>  | 5.2   | 1.4   | <0.01 | <0.001 | 0.2 | 97 |
| CH <sub>4</sub> Emissions from Forest Fires                                       | CH <sub>4</sub>  | 3.3   | 7.3   | <0.01 | 0.001  | 0.2 | 97 |

|  |                  |       |       |       |        |      |     |
|--|------------------|-------|-------|-------|--------|------|-----|
| CO <sub>2</sub> Emissions from Petrochemical Production                                  | CO <sub>2</sub>  | 21.6  | 26.5  | <0.01 | <0.001 | 0.2  | 97  |
| CO <sub>2</sub> Emissions from Cement Production   | CO <sub>2</sub>  | 33.3  | 38.8  | <0.01 | <0.001 | 0.2  | 97  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Residential                | CO <sub>2</sub>  | 3.0   | 0.0   | <0.01 | <0.001 | 0.2  | 97  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Gas - U.S. Territories            | CO <sub>2</sub>  | 0.0   | 3.0   | <0.01 | <0.001 | 0.2  | 98  |
| CO <sub>2</sub> Emissions from Carbon Dioxide Consumption                                | CO <sub>2</sub>  | 1.5   | 4.5   | <0.01 | <0.001 | 0.2  | 98  |
| CO <sub>2</sub> Emissions from Stationary Combustion - Coal - U.S. Territories           | CO <sub>2</sub>  | 0.6   | 3.4   | <0.01 | <0.001 | 0.2  | 98  |
| N <sub>2</sub> O Emissions from Manure Management  | N <sub>2</sub> O | 14.0  | 17.5  | <0.01 | <0.001 | 0.2  | 98  |
| N <sub>2</sub> O Emissions from Forest Fires   | N <sub>2</sub> O | 2.2   | 4.8   | <0.01 | 0.001  | 0.1  | 98  |
| CO <sub>2</sub> Emissions from Natural Gas Systems                                       | CO <sub>2</sub>  | 37.7  | 42.4  | <0.01 | <0.001 | 0.1  | 98  |
| CH <sub>4</sub> Emissions from Rice Cultivation  | CH <sub>4</sub>  | 13.1  | 11.9  | <0.01 | <0.001 | 0.1  | 98  |
| CO <sub>2</sub> Emissions from Urea Fertilization  | CO <sub>2</sub>  | 2.4   | 4.5   | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Nitric Acid Production                                   | N <sub>2</sub> O | 12.1  | 10.9  | <0.01 | <0.001 | 0.1  | 99  |
| CH <sub>4</sub> Emissions from Wastewater Treatment                                      | CH <sub>4</sub>  | 15.7  | 14.7  | <0.01 | <0.001 | 0.1  | 99  |
| CO <sub>2</sub> Emissions from Lime Production   | CO <sub>2</sub>  | 11.7  | 14.1  | <0.01 | <0.001 | 0.1  | 99  |
| CH <sub>4</sub> Emissions from Composting  | CH <sub>4</sub>  | 0.4   | 2.1   | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Composting   | N <sub>2</sub> O | 0.3   | 1.8   | <0.01 | <0.001 | 0.1  | 99  |
| Direct N <sub>2</sub> O Emissions from Agricultural Soil Management                      | N <sub>2</sub> O | 245.0 | 261.0 | <0.01 | <0.001 | 0.1  | 99  |
| Fugitive Emissions from Abandoned Underground Coal Mines                                 | CH <sub>4</sub>  | 7.2   | 6.3   | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Wastewater Treatment                                     | N <sub>2</sub> O | 3.4   | 4.8   | <0.01 | <0.001 | 0.1  | 99  |
| CO <sub>2</sub> Emissions from Land Converted to Grassland                               | CO <sub>2</sub>  | 39.1  | 40.4  | <0.01 | <0.001 | 0.1  | 99  |
| N <sub>2</sub> O Emissions from Settlement Soils   | N <sub>2</sub> O | 1.4   | 2.4   | <0.01 | <0.001 | 0.1  | 99  |
| CO <sub>2</sub> Emissions from Incineration of Waste                                     | CO <sub>2</sub>  | 8.0   | 9.4   | <0.01 | <0.001 | 0.1  | 99  |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | N <sub>2</sub> O | 3.1   | 2.4   | <0.01 | <0.001 | <0.1 | 99  |
| CO <sub>2</sub> Emissions from Liming  | CO <sub>2</sub>  | 4.7   | 4.1   | <0.01 | <0.001 | <0.1 | 100 |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> Emissions from Semiconductor Manufacture | Several          | 3.6   | 4.5   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Other                                 | N <sub>2</sub> O | 1.2   | 2.0   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential                   | CH <sub>4</sub>  | 5.2   | 5.0   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Phosphoric Acid Production                                | CO <sub>2</sub>  | 1.5   | 1.1   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Titanium Dioxide Production                               | CO <sub>2</sub>  | 1.2   | 1.8   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Mobile Combustion: Aviation                              | N <sub>2</sub> O | 1.7   | 1.4   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial                    | CH <sub>4</sub>  | 1.8   | 1.5   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Forest Soils   | N <sub>2</sub> O | 0.1   | 0.5   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Land Converted to Forest Land                             | CO <sub>2</sub>  | 0.7   | 0.3   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Ferroalloy Production                                     | CO <sub>2</sub>  | 2.2   | 1.9   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Glass Production  | CO <sub>2</sub>  | 1.5   | 1.3   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Zinc Production   | CO <sub>2</sub>  | 0.6   | 1.0   | <0.01 | <0.001 | <0.1 | 100 |

|  |                  |     |     |       |        |      |     |
|--|------------------|-----|-----|-------|--------|------|-----|
| CO <sub>2</sub> Emissions from Peatlands<br>Remaining Peatlands                                | CO <sub>2</sub>  | 1.1 | 0.8 | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Product Uses   | N <sub>2</sub> O | 4.2 | 4.2 | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Silicon Carbide<br>Production and Consumption                   | CO <sub>2</sub>  | 0.4 | 0.2 | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Petroleum Systems   | CO <sub>2</sub>  | 3.6 | 3.6 | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Incineration of Waste  | N <sub>2</sub> O | 0.5 | 0.3 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Mobile Combustion:<br>Other                                     | CH <sub>4</sub>  | 0.3 | 0.5 | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Semiconductor<br>Manufacture                                   | N <sub>2</sub> O | +   | 0.2 | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Soda Ash Production<br>and Consumption                          | CO <sub>2</sub>  | 2.8 | 2.8 | <0.01 | <0.001 | <0.1 | 100 |
| HFC-134a Emissions from Magnesium<br>Production and Processing                                 | HFCs             | 0.0 | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary<br>Combustion - Residential                      | N <sub>2</sub> O | 1.0 | 1.0 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Petrochemical<br>Production                                     | CH <sub>4</sub>  | 0.2 | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary<br>Combustion - Electricity Generation           | CH <sub>4</sub>  | 0.3 | 0.4 | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Mobile Combustion:<br>Marine                                   | N <sub>2</sub> O | 0.6 | 0.5 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary<br>Combustion - Commercial                       | N <sub>2</sub> O | 0.4 | 0.3 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Mobile Combustion:<br>Aviation                                  | CH <sub>4</sub>  | 0.1 | +   | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Field Burning of<br>Agricultural Residues                       | CH <sub>4</sub>  | 0.2 | 0.3 | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Lead Production   | CO <sub>2</sub>  | 0.5 | 0.5 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary<br>Combustion - U.S. Territories                 | N <sub>2</sub> O | 0.1 | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Silicon Carbide<br>Production and Consumption                   | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Field Burning of<br>Agricultural Residues                      | N <sub>2</sub> O | 0.1 | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Iron and Steel<br>Production & Metallurgical Coke<br>Production | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Stationary<br>Combustion - Geothermal Energy                    | CO <sub>2</sub>  | 0.4 | 0.4 | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary<br>Combustion - U.S. Territories                 | CH <sub>4</sub>  | +   | 0.1 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Ferroalloy<br>Production  | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| Non-CO <sub>2</sub> Emissions from Stationary<br>Combustion - Commercial                       | CH <sub>4</sub>  | 1.0 | 1.1 | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Urea Consumption<br>for Non-Ag Purposes                         | CO <sub>2</sub>  | 3.8 | 4.0 | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Peatlands<br>Remaining Peatlands                                | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CO <sub>2</sub> Emissions from Magnesium<br>Production and Processing                          | CO <sub>2</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| CH <sub>4</sub> Emissions from Mobile Combustion:<br>Marine                                    | CH <sub>4</sub>  | +   | +   | <0.01 | <0.001 | <0.1 | 100 |
| N <sub>2</sub> O Emissions from Peatlands<br>Remaining Peatlands                               | N <sub>2</sub> O | +   | +   | <0.01 | <0.001 | <0.1 | 100 |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

## References

IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Negara, and K. Tanabe (eds.). Hayman, Kanagawa, Japan.

# ANNEX 2 Methodology and Data for Estimating CO<sub>2</sub> Emissions from Fossil Fuel Combustion

## 2.1. Methodology for Estimating Emissions of CO<sub>2</sub> from Fossil Fuel Combustion

Carbon dioxide (CO<sub>2</sub>) emissions from fossil fuel combustion were estimated using a “bottom-up” methodology characterized by eight steps. These steps are described below.

### Step 1: Determine Total Fuel Consumption by Fuel Type and Sector

The bottom-up methodology used by the United States for estimating CO<sub>2</sub> emissions from fossil fuel combustion is conceptually similar to the approach recommended by the Intergovernmental Panel on Climate Change (IPCC) for countries that intend to develop detailed, sector-based emission estimates in line with a Tier 2 method in the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006). Total consumption data and adjustments to consumption are presented in Columns 2 through 13 of Table A-10.

Adjusted consumption data are presented in Columns 2 through 8 of Table A-11 through Table A-35 with totals by fuel type in Column 8 and totals by end-use sector in the last rows. Fuel consumption data for the bottom-up approach were obtained directly from the Energy Information Administration (EIA) of the U.S. Department of Energy. These data were first gathered in physical units, and then converted to their energy equivalents (see the Constants, Units, and Conversions Annex). The EIA data were collected through a variety of consumption surveys at the point of delivery or use and qualified with survey data on fuel production, imports, exports, and stock changes. Individual data elements were supplied by a variety of sources within EIA. Most information was taken from published reports, although some data were drawn from unpublished energy studies and databases maintained by EIA.

Energy consumption data were aggregated by sector (i.e., residential, commercial, industrial, transportation, electricity generation, and U.S. Territories), primary fuel type (e.g., coal, natural gas, and petroleum), and secondary fuel type (e.g., motor gasoline, distillate fuel). The 2014 total adjusted energy consumption across all sectors, including U.S. Territories, and energy types was 73,677.7 trillion British thermal units (Tbtu), as indicated in the last entry of column 13 in Table A-10. This total excludes fuel used for non-energy purposes and fuel consumed as international bunkers, both of which were deducted in earlier steps.

Electricity consumption information was allocated to each sector based on EIA’s distribution of electricity retail sales to ultimate customers (i.e., residential, commercial, industrial, and other). Because the “other” fuel use includes sales to both the commercial and transportation sectors, EIA’s limited transportation electricity use data were subtracted from “other” electricity use and also reported separately. This total was consequently combined with the commercial electricity data. Further information on these electricity end uses is described in EIA’s *Monthly Energy Review* (EIA 2016).

There are also three basic differences between the consumption data presented in Table A-10 and Table A-11 through Table A-35 and those recommended in the IPCC (2006) emission inventory methodology.

First, consumption data in the U.S. Inventory are presented using higher heating values (HHV)<sup>1</sup> rather than the lower heating values (LHV)<sup>2</sup> reflected in the IPCC (2006) emission inventory methodology. This convention is followed because data obtained from EIA are based on HHV. Of note, however, is that EIA renewable energy statistics are often published using LHV. The difference between the two conventions relates to the treatment of the heat energy that is consumed in the process of evaporating the water contained in the fuel. The simplified convention used by the International Energy Agency for converting from HHV to LHV is to multiply the energy content by 0.95 for petroleum and coal and by 0.9 for natural gas.

Second, while EIA's energy use data for the United States includes only the 50 U.S. states and the District of Columbia, the data reported to the United Nations Framework Convention on Climate Change (UNFCCC) are to include

<sup>1</sup> Also referred to as Gross Calorific Values (GCV).

<sup>2</sup> Also referred to as Net Calorific Values (NCV).

energy consumption within U.S. Territories. Therefore, consumption estimates for U.S. Territories<sup>3</sup> were added to domestic consumption of fossil fuels. Energy consumption data from U.S. Territories are presented in Column 7 of Table A-11 through Table A-35. It is reported separately from domestic sectoral consumption, because it is collected separately by EIA with no sectoral disaggregation.

Third, there were a number of modifications made in this report that may cause consumption information herein to differ from figures given in the cited literature. These are (1) the reallocation of select amounts of coking coal, petroleum coke, natural gas, residual fuel oil, and other oil (>401 degrees Fahrenheit) for processes accounted for in the Industrial Processes and Product Use chapter, (2) corrections for synthetic natural gas production, (3) subtraction of other fuels used for non-energy purposes, and (4) subtraction of international bunker fuels. These adjustments are described in the following steps.

### **Step 2: Subtract uses accounted for in the Industrial Processes and Product Use chapter.**

Portions of the fuel consumption data for seven fuel categories—coking coal, distillate fuel, industrial other coal, petroleum coke, natural gas, residual fuel oil, and other oil (>401 degrees Fahrenheit)—were reallocated to the Industrial Processes and Product Use chapter, as these portions were consumed as raw materials during non-energy related industrial processes. Emissions from these fuels used as raw materials are presented in the Industrial Processes and Product Use chapter, and are removed from the energy and non-energy consumption estimates within the Energy chapter.

- Coking coal is used as a raw material (specifically as a reducing agent) in the blast furnace process to produce iron and steel, lead, and zinc and therefore is not used as a fuel for this process.
- Similarly, petroleum coke is used in multiple processes as a raw material, and is thus not used as a fuel in those applications. The processes in which petroleum coke is used include (1) ferroalloy production, (2) aluminum production (for the production of C anodes and cathodes), (3) titanium dioxide production (in the chloride process), (4) ammonia production, and (5) silicon carbide.
- Natural gas consumption is used for the production of ammonia, and blast furnace and coke oven gas used in iron and steel production.
- Residual fuel oil and other oil (>401 degrees Fahrenheit) are both used in the production of C black.
- Natural gas, distillate fuel, coal, and metallurgical coke are used to produce pig iron through the reduction of iron ore in the production of iron and steel.

### **Step 3: Adjust for Conversion of Fossil Fuels and Exports**

First, a portion of industrial “other” coal that is accounted for in EIA coal combustion statistics is actually used to make “synthetic natural gas” via coal gasification at the Dakota Gasification Plant, a synthetic natural gas plant. The plant produces synthetic natural gas and byproduct CO<sub>2</sub>. The synthetic natural gas enters the natural gas distribution system. Since October 2000, a portion of the CO<sub>2</sub> produced by the coal gasification plant has been exported to Canada by pipeline. The remainder of the CO<sub>2</sub> byproduct from the plant is released to the atmosphere. The energy in this synthetic natural gas enters the natural gas distribution stream, and is accounted for in EIA natural gas combustion statistics. Because this energy of the synthetic natural gas is already accounted for as natural gas combustion, this amount of energy is deducted from the industrial coal consumption statistics to avoid double counting. The exported CO<sub>2</sub> is not emitted to the atmosphere in the United States, and therefore the energy used to produce this amount of CO<sub>2</sub> is subtracted from industrial other coal.

### **Step 4: Adjust Sectoral Allocation of Distillate Fuel Oil and Motor Gasoline**

EPA conducted a separate bottom-up analysis of transportation fuel consumption based on data from the Federal Highway Administration (FHWA). The FHWA data indicated that the amount of distillate and motor gasoline consumption allocated to the transportation sector in the EIA statistics should be adjusted. Therefore, for the estimates presented in the U.S. Inventory, the transportation sector’s distillate fuel and motor gasoline consumption was adjusted to match the value obtained from the bottom-up analysis. As the total distillate and motor gasoline consumption estimate from EIA are considered to be accurate at the national level, the distillate and motor gasoline consumption totals for the residential, commercial, and industrial sectors were adjusted proportionately.

---

<sup>3</sup>Fuel consumption by U.S. Territories (i.e., American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands) is included in this report.

### **Step 5: Subtract Consumption for Non-Energy Use**

U.S. aggregate energy statistics include consumption of fossil fuels for non-energy purposes. Depending on the end-use, non-energy uses of fossil fuels can result in long term storage of some or all of the C contained in the fuel. For example, asphalt made from petroleum can sequester up to 100 percent of the C contained in the petroleum feedstock for extended periods of time. Other non-energy fossil fuel products, such as lubricants or plastics also store C, but can lose or emit some of this C when they are used and/or burned as waste. As the emission pathways of C used for non-energy purposes are vastly different than fuel combustion, these emissions are estimated separately in the Carbon Emitted in Products from Non-Energy Uses of Fossil Fuels section in this chapter. Therefore, the amount of fuels used for non-energy purposes, shown in Table A-36, was subtracted from total fuel consumption.

### **Step 6: Subtract Consumption of International Bunker Fuels**

Emissions from international transport activities, or international bunker fuel consumption, are not included in national totals and instead reported separately, as required by the IPCC (2006) and UNFCCC (2014) inventory reporting guidelines. EIA energy statistics, however, include these bunker fuels—jet fuel for aircraft, and distillate fuel oil and residual fuel oil for marine shipping—as part of fuel consumption by the transportation end-use sector. Therefore, the amount of consumption for international bunker fuels was estimated and subtracted from total fuel consumption (see Table A-37). Emissions from international bunker fuels have been estimated separately and not included in national totals.<sup>4</sup>

### **Step 7: Determine the C Content of All Fuels**

The C content of combusted fossil fuels was estimated by multiplying adjusted energy consumption (Columns 2 through 8 of (Table A-11 through Table A-35) by fuel-specific C content coefficients (see Table A-38 and Table A-39) that reflect the amount of C per unit of energy in each fuel. The C content coefficients used in the U.S. Inventory were derived by EIA from detailed fuel information and are similar to the C content coefficients contained in the IPCC's default methodology (IPCC 2006), with modifications reflecting fuel qualities specific to the United States.

### **Step 8: Estimate CO<sub>2</sub> Emissions**

Actual CO<sub>2</sub> emissions in the United States were summarized by major fuel (i.e., coal, petroleum, natural gas, geothermal) and consuming sector (i.e., residential, commercial, industrial, transportation, electricity generation, and U.S. Territories). Emission estimates are expressed in million metric tons of carbon dioxide equivalents (MMT CO<sub>2</sub> Eq.). To convert from C content to CO<sub>2</sub> emissions, the fraction of C that is oxidized was applied. This fraction was 100 percent based on guidance in IPCC (2006).

To determine total emissions by final end-use sector, emissions from electricity generation were distributed to each end-use sector according to its share of aggregate electricity consumption (see Table A-40). This pro-rated approach to allocating emissions from electricity generation may overestimate or underestimate emissions for particular sectors due to differences in the average C content of fuel mixes burned to generate electricity.

To provide a more detailed accounting of emissions from transportation, fuel consumption data by vehicle type and transportation mode were used to allocate emissions by fuel type calculated for the transportation end-use sector. Additional information on the allocation is available in Annex 3.2.

[BEGIN BOX]

#### **Box 1. Uses of Greenhouse Gas Reporting Program Data in Reporting Emissions from Industrial Sector Fossil Fuel Combustion**

As described in the calculation methodology, total fossil fuel consumption for each year is based on aggregated end-use sector consumption published by the EIA. The availability of facility-level combustion emissions through EPA's Greenhouse Gas Reporting Program (GHGRP) has provided an opportunity to better characterize the industrial sector's energy consumption and emissions in the United States, through a disaggregation of EIA's industrial sector fuel consumption data from select industries.

---

<sup>4</sup> Refer to the International Bunker Fuels section of the Energy chapter and Annex 3.3 for a description of the methodology for distinguishing between international and domestic fuel consumption.

For EPA's GHGRP 2010, 2011, 2012, 2013, and 2014 reporting years, facility-level fossil fuel combustion emissions reported through the GHGRP were categorized and distributed to specific industry types by utilizing facility-reported NAICS codes (as published by the U.S. Census Bureau). As noted previously in this report, the definitions and provisions for reporting fuel types in EPA's GHGRP include some differences from the Inventory's use of EIA national fuel statistics to meet the UNFCCC reporting guidelines. The IPCC has provided guidance on aligning facility-level reported fuels and fuel types published in national energy statistics, which guided this exercise.<sup>5</sup>

This year's effort represents an attempt to align, reconcile, and coordinate the facility-level reporting of fossil fuel combustion emissions under EPA's GHGRP with the national-level approach presented in this report. Consistent with recommendations for reporting the Inventory to the UNFCCC, progress was made on certain fuel types for specific industries and has been included in the Common Reporting Format (CRF) tables that are submitted to the UNFCCC along with this report.<sup>6</sup> However, a full mapping was not completed this year due to fuel category differences between national statistics published by EIA and facility-level reported GHGRP data. Efforts in reconciling fuels focused on standard, common fuel types (e.g., natural gas, distillate fuel oil) where the fuels in EIA's national statistics aligned well with facility-level GHGRP data. For these reasons, the current information presented in the CRF tables should be viewed as an initial attempt at this exercise. Additional efforts will be made for future Inventory reports to improve the mapping of fuel types, and examine ways to reconcile and coordinate any differences between facility-level data and national statistics.

Additionally, this year's analysis expanded this effort through the full time series presented in this report, by linking EPA's GHGRP facility-level reporting with the information published by EIA in its Manufacturing Energy Consumption Survey (MECS) data. The GHGRP and MECS data provided disaggregated industrial fuel consumption by NAICS codes for certain years of the 1990-2014 time series. The GHGRP data are available for 2010, 2011, 2012, 2013, and 2014. The MECS data are available for 1991, 1994, 1998, 2002, 2006, and 2010. In order to disaggregate NAICS codes for the full entire time series, the GHGRP and MECS data were used in conjunction with the annual aggregate EIA MER data to implement splicing techniques as advised in the 2006 IPCC Guidelines "Volume 1 General Guidance and Reporting, Chapter 5 Time Series Consistency" to provide the full time series<sup>7</sup>. The results of the industrial sector disaggregation effort are provided in the CRF tables submitted with this Inventory. It is believed that the current analysis has led to improvements in the presentation of data in the Inventory, but further work will be conducted, and future improvements will be realized in subsequent Inventory reports.

Additionally, to assist in the disaggregation of industrial fuel consumption, EIA will now synthesize energy consumption data using the same procedure as is used for the last historical (benchmark) year of the Annual Energy Outlook (AEO). This procedure reorganizes the most recent data from the MECS (conducted every four years) into the nominal data submission year using the same energy-economy integrated model used to produce the AEO projections, the National Energy Modeling System (NEMS). EIA believes this "nowcasting" technique provides an appropriate estimate of energy consumption for the CRF.

To address gaps in the time series, EIA performs a NEMS model projection, using the MECS baseline sub-sector energy consumption. The NEMS model accounts for changes in factors that influence industrial sector energy consumption, and has access to data which may be more recent than MECS, such as industrial sub-sector macro industrial output (i.e., shipments) and fuel prices. By evaluating the impact of these factors on industrial subsector energy consumption, NEMS can anticipate changes to the energy shares occurring post-MECS and can provide a way to appropriately disaggregate the energy-related emissions data into the CRF.

While the fuel consumption values for the various manufacturing sub-sectors are not directly surveyed for all years, they represent EIA's best estimate of historical consumption values for non-MECS years. Moreover, as an integral part of each AEO publication, this synthetic data series is likely to be maintained consistent with all available EIA and non-EIA data sources even as the underlying data sources evolve for both manufacturing and non-manufacturing industries alike.

Other sectors' fuel consumption (i.e., commercial, residential, transportation) will be benchmarked with the latest aggregate values from the *Monthly Energy Review*.<sup>8</sup> EIA will work with the EPA to back cast these values to 1990.

[END BOX]

---

<sup>5</sup> See Section 4 "Use of Facility-Level Data in Good Practice National Greenhouse Gas Inventories" of the IPCC meeting report, and specifically the section on using facility-level data in conjunction with energy data, at <[http://www.ipcc-nggip.iges.or.jp/meeting/pdffiles/1008\\_Model\\_and\\_Facility\\_Level\\_Data\\_Report.pdf](http://www.ipcc-nggip.iges.or.jp/meeting/pdffiles/1008_Model_and_Facility_Level_Data_Report.pdf)>.

<sup>6</sup> See <<http://www.epa.gov/climatechange/ghgemissions/usinventoryreport.html>>.

<sup>7</sup> Available online at: <[http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/1\\_Volume1/V1\\_5\\_Ch5\\_Timeseries.pdf](http://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/1_Volume1/V1_5_Ch5_Timeseries.pdf)>

<sup>8</sup> See <<http://www.eia.gov/totalenergy/data/monthly/>>.

**Table A-10: 2014 Energy Consumption Data by Fuel Type (TBtu) and Adjusted Energy Consumption Data**

|                           | 1                                     | 2              | 3               | 4               | 5               | 6            | 7               | 8                               | 9                          | 10           | 11          | 12   | 13                         |
|---------------------------|---------------------------------------|----------------|-----------------|-----------------|-----------------|--------------|-----------------|---------------------------------|----------------------------|--------------|-------------|------|----------------------------|
| Fuel Type                 | Total Consumption (TBtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Adjustments (TBtu) <sup>b</sup> |                            |              |             |      | Total Adjusted Consumption |
|                           | Res.                                  | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Bunker Fuel                     | Unadjusted NEU Consumption |              |             |      |                            |
|                           |                                       |                |                 |                 |                 |              |                 |                                 | Ind.                       | Trans.       | Terr.       |      |                            |
| <b>Total Coal</b>         | NE                                    | 47.9           | 828.9           | NE              | 16,441.0        | 36.9         | 17,354.7        |                                 | 33.4                       |              |             |      | 17,321.3                   |
| Residential Coal          | NE                                    |                |                 |                 |                 |              | NE              |                                 |                            |              |             |      | NE                         |
| Commercial Coal           |                                       | 47.9           |                 |                 |                 |              | 47.9            |                                 |                            |              |             |      | 47.9                       |
| Industrial Coking Coal    |                                       |                | 23.0            |                 |                 |              | 23.0            |                                 | 23.0                       |              |             |      | 23.0                       |
| Industrial Other Coal     |                                       |                | 805.8           |                 |                 |              | 805.8           |                                 | 10.3                       |              |             |      | 795.5                      |
| Transportation Coal       |                                       |                |                 | NE              |                 |              | NE              |                                 |                            |              |             |      | NE                         |
| Electric Power Coal       |                                       |                |                 |                 | 16,441.0        |              | 16,441.0        |                                 |                            |              |             |      | 16,441.0                   |
| U.S. Territory Coal (bit) |                                       |                |                 |                 |                 | 36.9         | 36.9            |                                 |                            |              |             |      | 36.9                       |
| <b>Natural Gas</b>        | 5,237.3                               | 3,568.8        | 9,104.3         | 898.9           | 8,361.7         | 56.6         | 27,227.5        |                                 | 311.8                      |              |             |      | 26,915.7                   |
| <b>Total Petroleum</b>    | 989.6                                 | 543.2          | 7,993.7         | 25,016.3        | 295.5           | 528.9        | 35,367.1        | 1,436.3                         | 4,354.8                    | 136.0        | 53.6        |      | 29,386.5                   |
| Asphalt & Road Oil        |                                       |                | 792.6           |                 |                 |              | 792.6           |                                 | 792.6                      |              |             |      |                            |
| Aviation Gasoline         |                                       |                |                 | 21.7            |                 |              | 21.7            |                                 |                            |              |             |      | 21.7                       |
| Distillate Fuel Oil       | 513.9                                 | 343.7          | 1,314.7         | 6,134.3         | 82.2            | 74.9         | 8,463.6         | 82.0                            | 5.8                        |              |             |      | 8,375.7                    |
| Jet Fuel                  |                                       |                |                 | 3,042.1         | NA              | 55.2         | 3,097.3         | 985.1                           |                            |              |             |      | 2,112.2                    |
| Kerosene                  | 13.7                                  | 2.0            | 2.8             |                 |                 | 7.2          | 25.8            |                                 |                            |              |             |      | 25.8                       |
| LPG                       | 462.0                                 | 151.2          | 2,433.3         | 43.7            |                 | 13.4         | 3,103.5         |                                 | 2,148.7                    |              |             |      | 954.8                      |
| Lubricants                |                                       |                | 144.0           | 136.0           |                 | 1.0          | 281.1           |                                 | 144.0                      | 136.0        | 1.0         |      |                            |
| Motor Gasoline            |                                       | 37.9           | 134.7           | 15,192.0        |                 | 175.9        | 15,540.5        |                                 |                            |              |             |      | 15,540.5                   |
| Residual Fuel             |                                       | 7.9            |                 | 446.5           | 95.1            | 148.7        | 698.3           | 369.2                           |                            |              |             |      | 329.2                      |
| Other Petroleum           |                                       |                |                 |                 |                 |              |                 |                                 |                            |              |             |      |                            |
| AvGas Blend Components    |                                       |                | (0.1)           |                 |                 |              | (0.1)           |                                 |                            |              |             |      | (0.1)                      |
| Crude Oil                 |                                       |                |                 |                 |                 |              |                 |                                 |                            |              |             |      |                            |
| MoGas Blend Components    |                                       |                |                 |                 |                 |              |                 |                                 |                            |              |             |      |                            |
| Misc. Products            |                                       |                | 182.7           |                 |                 | 52.6         | 235.3           |                                 | 182.7                      |              |             | 52.6 |                            |
| Naphtha (<401 deg. F)     |                                       |                | 442.6           |                 |                 |              | 442.6           |                                 | 442.6                      |              |             |      |                            |
| Other Oil (>401 deg. F)   |                                       |                | 247.2           |                 |                 |              | 247.2           |                                 | 247.2                      |              |             |      |                            |
| Pentanes Plus             |                                       |                | 88.4            |                 |                 |              | 88.4            |                                 | 44.2                       |              |             |      | 44.2                       |
| Petroleum Coke            |                                       | 0.5            | 653.5           |                 | 118.2           |              | 772.1           |                                 | 61.4                       |              |             |      | 710.8                      |
| Still Gas                 |                                       |                | 1,516.9         |                 |                 |              | 1,516.9         |                                 | 164.6                      |              |             |      | 1,352.4                    |
| Special Naphtha           |                                       |                | 106.1           |                 |                 |              | 106.1           |                                 | 106.1                      |              |             |      |                            |
| Unfinished Oils           |                                       |                | (80.6)          |                 |                 |              | (80.6)          |                                 |                            |              |             |      | (80.6)                     |
| Waxes                     |                                       |                | 14.8            |                 |                 |              | 14.8            |                                 | 14.8                       |              |             |      |                            |
| <b>Geothermal</b>         |                                       |                |                 |                 | 54.2            |              | 54.2            |                                 |                            |              |             |      | 54.2                       |
| <b>Total (All Fuels)</b>  | <b>6,226.8</b>                        | <b>4,159.9</b> | <b>17,926.8</b> | <b>25,915.2</b> | <b>25,152.4</b> | <b>622.3</b> | <b>80,003.5</b> | <b>1,436.3</b>                  | <b>4,699.9</b>             | <b>136.0</b> | <b>53.6</b> |      | <b>73,677.7</b>            |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37). Note: Parentheses indicate negative values. NE (Not Estimated); NA (Not Available).

**Table A-11: 2014 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 47.9           | 795.5           | NE              | 16,441.0        | 36.9         | 17,321.3        | NE   | 4.5          | 75.3         | NE             | 1,570.4        | 3.4         | 1,653.7        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 47.9           |                 |                 |                 |              | 47.9            |  | 4.5          |              |                |                |             | 4.5            |    |
| Industrial Other Coal     |  |                | 795.5           |                 |                 |              | 795.5           |  |              | 75.3         |                |                |             | 75.3           |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 16,441.0        |              | 16,441.0        |  |              |              |                | 1,570.4        |             | 1,570.4        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 5,237.3                                  | 3,568.8        | 8,792.5         | 898.9           | 8,361.7         | 56.6         | 26,915.7        | 277.6  | 189.2        | 466.0        | 47.6           | 443.2          | 3.0         | 1,426.6        |    |
| <b>Total Petroleum</b>    | 989.6                                    | 543.2          | 3,638.9         | 23,444.0        | 295.5           | 475.3        | 29,386.5        | 67.5   | 38.2         | 271.9        | 1,690.0        | 25.3           | 34.6        | 2,127.5        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 21.7            |                 |              | 21.7            |  |              |              | 1.5            |                |             | 1.5            |    |
| Distillate Fuel Oil       | 513.9                                    | 343.7          | 1,308.9         | 6,052.2         | 82.2            | 74.9         | 8,375.7         | 38.0   | 25.4         | 96.8         | 447.6          | 6.1            | 5.5         | 619.4          |    |
| Jet Fuel                  |  |                |                 | 2,057.0         | NA              | 55.2         | 2,112.2         |  |              |              | 148.6          |                | 4.0         | 152.5          |    |
| Kerosene                  | 13.7                                     | 2.0            | 2.8             |                 |                 | 7.2          | 25.8            | 1.0  | 0.1          | 0.2          |                |                | 0.5         | 1.9            |    |
| LPG                       | 462.0                                    | 151.2          | 284.6           | 43.7            |                 | 13.4         | 954.8           | 28.5   | 9.3          | 17.6         | 2.7            |                | 0.8         | 58.9           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 37.9           | 134.7           | 15,192.0        |                 | 175.9        | 15,540.5        |  | 2.7          | 9.6          | 1,083.8        |                | 12.5        | 1,108.6        |    |
| Residual Fuel             |  | 7.9            |                 | 77.4            | 95.1            | 148.7        | 329.2           |  | 0.6          |              | 5.8            | 7.1            | 11.2        | 24.7           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | (0.1)           |                 |                 |              | (0.1)           |  |              | (0.0)        |                |                |             | (0.0)          |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 44.2            |                 |                 |              | 44.2            |  |              | 3.1          |                |                |             | 3.1            |    |
| Petroleum Coke            |  | 0.5            | 592.1           |                 | 118.2           |              | 710.8           |  | 0.1          | 60.5         |                | 12.1           |             | 72.6           |    |
| Still Gas                 |  |                | 1,352.4         |                 |                 |              | 1,352.4         |  |              | 90.2         |                |                |             | 90.2           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (80.6)          |                 |                 |              | (80.6)          |  |              | (6.0)        |                |                |             | (6.0)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 54.2            |              | 54.2            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>6,226.8</b>                           | <b>4,159.9</b> | <b>13,226.9</b> | <b>24,342.9</b> | <b>25,152.4</b> | <b>568.7</b> | <b>73,677.7</b> | <b>345.1</b>   | <b>231.9</b> | <b>813.3</b> | <b>1,737.6</b> | <b>2,039.3</b> | <b>41.0</b> | <b>5,208.2</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

NE (Not Estimated)

NA (Not Available)

Note: Parentheses indicate negative values.

**Table A-12: 2013 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 41.4           | 800.0           | NE              | 16,450.6        | 36.9         | 17,328.8        | NE   | 3.9          | 75.7         | NE             | 1,571.3        | 3.4         | 1,654.4        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 41.4           |                 |                 |                 |              | 41.4            |  | 3.9          |              |                |                |             | 3.9            |    |
| Industrial Other Coal     |  |                | 800.0           |                 |                 |              | 800.0           |  |              | 75.7         |                |                |             | 75.7           |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 16,450.6        |              | 16,450.6        |  |              |              |                | 1,571.3        |             | 1,571.3        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 5,022.9                                  | 3,379.8        | 8,525.3         | 887.3           | 8,376.3         | 56.6         | 26,248.2        | 266.2  | 179.1        | 451.9        | 47.0           | 444.0          | 3.0         | 1,391.2        |    |
| <b>Total Petroleum</b>    | 935.9                                    | 540.1          | 3,823.7         | 23,108.6        | 255.2           | 510.1        | 29,173.6        | 63.4   | 38.0         | 284.6        | 1,666.0        | 22.4           | 37.1        | 2,111.6        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 22.4            |                 |              | 22.4            |  |              |              | 1.5            |                |             | 1.5            |    |
| Distillate Fuel Oil       | 457.2                                    | 319.6          | 1,169.8         | 5,866.3         | 55.4            | 80.4         | 7,948.6         | 33.8   | 23.6         | 86.5         | 433.9          | 4.1            | 5.9         | 587.9          |    |
| Jet Fuel                  |  |                |                 | 2,036.9         | NA              | 59.2         | 2,096.1         |  |              |              | 147.1          |                | 4.3         | 151.4          |    |
| Kerosene                  | 8.3                                      | 1.0            | 1.5             |                 |                 | 7.7          | 18.4            | 0.6  | 0.1          | 0.1          |                |                | 0.6         | 1.3            |    |
| LPG                       | 470.5                                    | 153.9          | 348.9           | 44.5            |                 | 14.4         | 1,032.2         | 29.0   | 9.5          | 21.5         | 2.7            |                | 0.9         | 63.7           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 40.8           | 268.7           | 14,937.2        |                 | 188.8        | 15,435.6        |  | 2.9          | 19.2         | 1,065.6        |                | 13.5        | 1,101.2        |    |
| Residual Fuel             |  | 24.4           |                 | 201.4           | 77.2            | 159.6        | 462.6           |  | 1.8          |              | 15.1           | 5.8            | 12.0        | 34.7           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | (0.4)           |                 |                 |              | (0.4)           |  |              | (0.0)        |                |                |             | (0.0)          |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 47.1            |                 |                 |              | 47.1            |  |              | 3.3          |                |                |             | 3.3            |    |
| Petroleum Coke            |  | 0.4            | 600.9           |                 | 122.5           |              | 723.7           |  | 0.0          | 61.4         |                | 12.5           |             | 73.9           |    |
| Still Gas                 |  |                | 1,370.6         |                 |                 |              | 1,370.6         |  |              | 91.4         |                |                |             | 91.4           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 16.7            |                 |                 |              | 16.7            |  |              | 1.2          |                |                |             | 1.2            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 53.8            |              | 53.8            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>5,958.8</b>                           | <b>3,961.3</b> | <b>13,149.0</b> | <b>23,995.9</b> | <b>25,135.8</b> | <b>603.6</b> | <b>72,804.4</b> | <b>329.7</b>   | <b>221.0</b> | <b>812.2</b> | <b>1,713.0</b> | <b>2,038.1</b> | <b>43.5</b> | <b>5,157.6</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

Note: Parentheses indicate negative values.

NE (Not Estimated)

NA (Not Available)

**Table A-13: 2012 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 43.6           | 782.3           | NE              | 15,821.2        | 36.9         | 16,684.0        | NE   | 4.1          | 74.1         | NE             | 1,511.2        | 3.4         | 1,592.8        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 43.6           |                 |                 |                 |              | 43.6            |  | 4.1          |              |                |                |             | 4.1            |    |
| Industrial Other Coal     |  |                | 782.3           |                 |                 |              | 782.3           |  |              | 74.1         |                |                |             | 74.1           |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 15,821.2        |              | 15,821.2        |  |              |              |                | 1,511.2        |             | 1,511.2        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 4,242.1                                  | 2,959.5        | 8,203.0         | 779.8           | 9,286.8         | 49.2         | 25,520.3        | 224.8  | 156.9        | 434.8        | 41.3           | 492.2          | 2.6         | 1,352.6        |    |
| <b>Total Petroleum</b>    | 846.7                                    | 506.0          | 3,650.4         | 22,963.0        | 214.2           | 516.1        | 28,696.4        | 57.7   | 35.7         | 274.1        | 1,655.4        | 18.3           | 37.6        | 2,078.9        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 25.1            |                 |              | 25.1            |  |              |              | 1.7            |                |             | 1.7            |    |
| Distillate Fuel Oil       | 437.5                                    | 322.1          | 1,144.7         | 5,780.1         | 52.4            | 81.3         | 7,818.1         | 32.4   | 23.8         | 84.7         | 427.5          | 3.9            | 6.0         | 578.2          |    |
| Jet Fuel                  |  |                |                 | 1,985.2         | NA              | 59.9         | 2,045.1         |  |              |              | 143.4          |                | 4.3         | 147.7          |    |
| Kerosene                  | 7.7                                      | 1.2            | 2.0             |                 |                 | 7.8          | 18.7            | 0.6  | 0.1          | 0.1          |                |                | 0.6         | 1.4            |    |
| LPG                       | 401.6                                    | 137.6          | 345.0           | 37.2            |                 | 14.6         | 935.9           | 24.8   | 8.5          | 21.3         | 2.3            |                | 0.9         | 57.8           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 13.3           | 87.1            | 14,924.3        |                 | 191.1        | 15,215.9        |  | 1.0          | 6.2          | 1,064.7        |                | 13.6        | 1,085.5        |    |
| Residual Fuel             |  | 31.4           |                 | 211.1           | 76.7            | 161.5        | 480.6           |  | 2.4          |              | 15.8           | 5.8            | 12.1        | 36.1           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | (0.0)           |                 |                 |              | (0.0)           |  |              | (0.0)        |                |                |             | (0.0)          |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 42.2            |                 |                 |              | 42.2            |  |              | 3.0          |                |                |             | 3.0            |    |
| Petroleum Coke            |  | 0.4            | 649.1           |                 | 85.1            |              | 734.6           |  | 0.0          | 66.3         |                | 8.7            |             | 75.0           |    |
| Still Gas                 |  |                | 1,320.2         |                 |                 |              | 1,320.2         |  |              | 88.1         |                |                |             | 88.1           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 60.1            |                 |                 |              | 60.1            |  |              | 4.5          |                |                |             | 4.5            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 53.1            |              | 53.1            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>5,088.8</b>                           | <b>3,509.1</b> | <b>12,635.7</b> | <b>23,742.8</b> | <b>25,375.3</b> | <b>602.1</b> | <b>70,953.8</b> | <b>282.5</b>   | <b>196.7</b> | <b>782.9</b> | <b>1,696.8</b> | <b>2,022.2</b> | <b>43.6</b> | <b>5,024.7</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

Note: Parentheses indicate negative values. NE (Not Estimated); NA (Not Available).

**Table A-14: 2011 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 61.7           | 866.1           | NE              | 18,035.2        | 36.9         | 18,999.9        | NE   | 5.8          | 82.0         | NE             | 1,722.7        | 3.4         | 1,813.9        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 61.7           |                 |                 |                 |              | 61.7            |  | 5.8          |              |                |                |             | 5.8            |    |
| Industrial Other Coal     |  |                | 866.1           |                 |                 |              | 866.1           |  |              | 82.0         |                |                |             | 82.0           |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 18,035.2        |              | 18,035.2        |  |              |              |                | 1,722.7        |             | 1,722.7        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 4,804.6                                  | 3,216.1        | 7,873.4         | 733.5           | 7,712.2         | 27.1         | 24,366.9        | 254.7  | 170.5        | 417.3        | 38.9           | 408.8          | 1.4         | 1,291.5        |    |
| <b>Total Petroleum</b>    | 1,059.7                                  | 625.4          | 3,645.5         | 23,145.2        | 295.0           | 503.8        | 29,274.6        | 72.2   | 44.5         | 274.0        | 1,668.8        | 25.8           | 36.7        | 2,121.9        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 27.1            |                 |              | 27.1            |  |              |              | 1.9            |                |             | 1.9            |    |
| Distillate Fuel Oil       | 534.8                                    | 400.5          | 1,255.8         | 5,814.5         | 63.7            | 79.3         | 8,148.5         | 39.6   | 29.6         | 92.9         | 430.0          | 4.7            | 5.9         | 602.6          |    |
| Jet Fuel                  |  |                |                 | 2,029.9         | NA              | 58.5         | 2,088.4         |  |              |              | 146.6          |                | 4.2         | 150.8          |    |
| Kerosene                  | 18.5                                     | 3.2            | 3.6             |                 |                 | 7.6          | 32.9            | 1.4  | 0.2          | 0.3          |                |                | 0.6         | 2.4            |    |
| LPG                       | 506.4                                    | 146.5          | 208.8           | 34.0            |                 | 14.2         | 909.8           | 31.2   | 9.0          | 12.9         | 2.1            |                | 0.9         | 56.1           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 21.3           | 123.2           | 14,981.8        |                 | 186.6        | 15,313.0        |  | 1.5          | 8.8          | 1,068.8        |                | 13.3        | 1,092.4        |    |
| Residual Fuel             |  | 53.7           | 46.9            | 258.0           | 93.1            | 157.6        | 609.2           |  | 4.0          | 3.5          | 19.4           | 7.0            | 11.8        | 45.7           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 0.0             |                 |                 |              | 0.0             |  |              | 0.0          |                |                |             | 0.0            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 27.3            |                 |                 |              | 27.3            |  |              | 1.9          |                |                |             | 1.9            |    |
| Petroleum Coke            |  | 0.2            | 600.3           |                 | 138.3           |              | 738.8           |  | 0.0          | 61.3         |                | 14.1           |             | 75.4           |    |
| Still Gas                 |  |                | 1,323.4         |                 |                 |              | 1,323.4         |  |              | 88.3         |                |                |             | 88.3           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 56.1            |                 |                 |              | 56.1            |  |              | 4.2          |                |                |             | 4.2            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 52.3            |              | 52.3            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>5,864.2</b>                           | <b>3,903.3</b> | <b>12,385.0</b> | <b>23,878.7</b> | <b>26,094.7</b> | <b>567.8</b> | <b>72,693.6</b> | <b>326.8</b>   | <b>220.7</b> | <b>773.3</b> | <b>1,707.6</b> | <b>2,157.7</b> | <b>41.5</b> | <b>5,227.7</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

Note: Parentheses indicate negative values. NE (Not Estimated); NA (Not Available).

**Table A-15: 2010 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (TBtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 69.7           | 951.6           | NE              | 19,133.5        | 36.9         | 20,191.6        | NE   | 6.6          | 90.1         | NE             | 1,827.6        | 3.4         | 1,927.7        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 69.7           |                 |                 |                 |              | 69.7            |  | 6.6          |              |                |                |             | 6.6            |    |
| Industrial Other Coal     |  |                | 951.6           |                 |                 |              | 951.6           |  |              | 90.1         |                |                |             | 90.1           |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 19,133.5        |              | 19,133.5        |  |              |              |                | 1,827.6        |             | 1,827.6        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 4,878.1                                  | 3,164.7        | 7,683.2         | 719.0           | 7,527.6         | 27.8         | 24,000.4        | 258.6  | 167.7        | 407.2        | 38.1           | 399.0          | 1.5         | 1,272.1        |    |
| <b>Total Petroleum</b>    | 1,116.2                                  | 643.5          | 3,699.2         | 23,447.1        | 370.3           | 501.6        | 29,777.9        | 76.0   | 45.8         | 278.2        | 1,690.2        | 31.4           | 36.5        | 2,158.2        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 27.0            |                 |              | 27.0            |  |              |              | 1.9            |                |             | 1.9            |    |
| Distillate Fuel Oil       | 557.0                                    | 388.0          | 1,134.4         | 5,706.1         | 79.7            | 78.9         | 7,944.0         | 41.2   | 28.7         | 83.9         | 422.0          | 5.9            | 5.8         | 587.5          |    |
| Jet Fuel                  |  |                |                 | 2,097.5         | NA              | 58.2         | 2,155.6         |  |              |              | 151.5          |                | 4.2         | 155.7          |    |
| Kerosene                  | 29.1                                     | 4.8            | 7.3             |                 |                 | 7.5          | 48.8            | 2.1  | 0.4          | 0.5          |                |                | 0.6         | 3.6            |    |
| LPG                       | 530.1                                    | 140.1          | 219.7           | 29.5            |                 | 14.2         | 933.6           | 32.7   | 8.6          | 13.6         | 1.8            |                | 0.9         | 57.6           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 48.5           | 243.1           | 15,314.8        |                 | 186.1        | 15,792.5        |  | 3.5          | 17.3         | 1,092.5        |                | 13.3        | 1,126.6        |    |
| Residual Fuel             |  | 61.7           | 32.2            | 272.2           | 154.1           | 156.8        | 676.9           |  | 4.6          | 2.4          | 20.4           | 11.6           | 11.8        | 50.8           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | (0.2)           |                 |                 |              | (0.2)           |  |              | (0.0)        |                |                |             | (0.0)          |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 77.8            |                 |                 |              | 77.8            |  |              | 5.4          |                |                |             | 5.4            |    |
| Petroleum Coke            |  | 0.3            | 633.0           |                 | 136.6           |              | 770.0           |  | 0.0          | 64.6         |                | 13.9           |             | 78.6           |    |
| Still Gas                 |  |                | 1,324.0         |                 |                 |              | 1,324.0         |  |              | 88.3         |                |                |             | 88.3           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 28.0            |                 |                 |              | 28.0            |  |              | 2.1          |                |                |             | 2.1            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 51.9            |              | 51.9            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>5,994.3</b>                           | <b>3,877.8</b> | <b>12,334.0</b> | <b>24,166.0</b> | <b>27,083.3</b> | <b>566.2</b> | <b>74,021.7</b> | <b>334.6</b>   | <b>220.1</b> | <b>775.5</b> | <b>1,728.3</b> | <b>2,258.4</b> | <b>41.4</b> | <b>5,358.3</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

Note: Parentheses indicate negative values. NE (Not Estimated); NA (Not Available).

**Table A-16: 2009 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 73.4           | 877.3           | NE              | 18,225.3        | 36.9         | 19,212.8        | NE   | 6.9          | 83.0         | NE             | 1,740.9        | 3.4         | 1,834.2        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 73.4           |                 |                 |                 |              | 73.4            |  | 6.9          |              |                |                |             | 6.9            |    |
| Industrial Other Coal     |  |                | 877.3           |                 |                 |              | 877.3           |  |              | 83.0         |                |                |             | 83.0           |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 18,225.3        |              | 18,225.3        |  |              |              |                | 1,740.9        |             | 1,740.9        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 4,883.1                                  | 3,186.6        | 7,125.1         | 714.9           | 7,022.4         | 27.4         | 22,959.4        | 258.8  | 168.9        | 377.6        | 37.9           | 372.2          | 1.5         | 1,216.9        |    |
| <b>Total Petroleum</b>    | 1,138.2                                  | 668.6          | 3,550.0         | 23,303.6        | 382.4           | 531.0        | 29,573.7        | 77.5   | 47.6         | 267.0        | 1,679.1        | 32.2           | 38.6        | 2,142.0        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 26.6            |                 |              | 26.6            |  |              |              | 1.8            |                |             | 1.8            |    |
| Distillate Fuel Oil       | 563.4                                    | 382.4          | 1,018.1         | 5,488.2         | 69.6            | 83.4         | 7,605.1         | 41.7   | 28.3         | 75.3         | 405.9          | 5.1            | 6.2         | 562.4          |    |
| Jet Fuel                  |  |                |                 | 2,134.2         | NA              | 61.5         | 2,195.7         |  |              |              | 154.1          |                | 4.4         | 158.6          |    |
| Kerosene                  | 27.7                                     | 4.2            | 4.4             |                 |                 | 8.0          | 44.3            | 2.0  | 0.3          | 0.3          |                |                | 0.6         | 3.2            |    |
| LPG                       | 547.1                                    | 138.9          | 201.7           | 28.0            |                 | 15.0         | 930.7           | 33.8   | 8.6          | 12.4         | 1.7            |                | 0.9         | 57.4           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 71.6           | 328.2           | 15,440.8        |                 | 197.5        | 16,038.2        |  | 5.1          | 23.4         | 1,101.5        |                | 14.1        | 1,144.1        |    |
| Residual Fuel             |  | 71.3           | 67.3            | 185.7           | 181.0           | 165.7        | 670.9           |  | 5.4          | 5.1          | 13.9           | 13.6           | 12.4        | 50.4           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | (0.8)           |                 |                 |              | (0.8)           |  |              | (0.1)        |                |                |             | (0.1)          |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 63.8            |                 |                 |              | 63.8            |  |              | 4.5          |                |                |             | 4.5            |    |
| Petroleum Coke            |  | 0.2            | 624.0           |                 | 131.8           |              | 756.1           |  | 0.0          | 63.7         |                | 13.5           |             | 77.2           |    |
| Still Gas                 |  |                | 1,321.1         |                 |                 |              | 1,321.1         |  |              | 88.1         |                |                |             | 88.1           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (77.8)          |                 |                 |              | (77.8)          |  |              | (5.8)        |                |                |             | (5.8)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 51.2            |              | 51.2            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>6,021.3</b>                           | <b>3,928.5</b> | <b>11,552.3</b> | <b>24,018.5</b> | <b>25,681.3</b> | <b>595.2</b> | <b>71,797.2</b> | <b>336.3</b>   | <b>223.5</b> | <b>727.7</b> | <b>1,717.0</b> | <b>2,145.7</b> | <b>43.5</b> | <b>5,193.5</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37). Note: Parentheses indicate negative values. NE (Not Estimated); NA (Not Available).

**Table A-17: 2008 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | NE                                       | 80.8           | 1,081.5         | NE              | 20,513.0        | 36.9         | 21,712.0        | NE   | 7.6          | 102.4        | NE             | 1,959.4        | 3.4         | 2,072.8        |    |
| Residential Coal          | NE                                       |                |                 |                 |                 |              | NE              | NE   |              |              |                |                |             | NE             |    |
| Commercial Coal           |  | 80.8           |                 |                 |                 |              | 80.8            |  | 7.6          |              |                |                |             | 7.6            |    |
| Industrial Other Coal     |  |                | 1,081.5         |                 |                 |              | 1,081.5         |  |              | 102.4        |                |                |             | 102.4          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 20,513.0        |              | 20,513.0        |  |              |              |                | 1,959.4        |             | 1,959.4        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 5,010.1                                  | 3,228.4        | 7,571.4         | 692.1           | 6,828.9         | 29.3         | 23,360.2        | 265.5  | 171.1        | 401.3        | 36.7           | 361.9          | 1.6         | 1,238.1        |    |
| <b>Total Petroleum</b>    | 1,201.5                                  | 635.0          | 3,994.0         | 24,342.5        | 459.3           | 489.9        | 31,122.2        | 82.1   | 44.9         | 299.2        | 1,755.1        | 38.4           | 35.8        | 2,255.5        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 28.3            |                 |              | 28.3            |  |              |              | 2.0            |                |             | 2.0            |    |
| Distillate Fuel Oil       | 627.5                                    | 321.3          | 1,103.1         | 6,058.8         | 72.5            | 110.3        | 8,293.5         | 46.4   | 23.8         | 81.6         | 448.1          | 5.4            | 8.2         | 613.4          |    |
| Jet Fuel                  |  |                |                 | 2,396.1         | NA              | 35.0         | 2,431.1         |  |              |              | 173.0          |                | 2.5         | 175.6          |    |
| Kerosene                  | 21.3                                     | 4.4            | 3.8             |                 |                 | 5.9          | 35.4            | 1.6  | 0.3          | 0.3          |                |                | 0.4         | 2.6            |    |
| LPG                       | 552.7                                    | 158.0          | 226.7           | 40.1            |                 | 15.7         | 993.3           | 34.1   | 9.7          | 14.0         | 2.5            |                | 1.0         | 61.3           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 80.0           | 437.3           | 15,547.9        |                 | 133.7        | 16,198.9        |  | 5.7          | 31.2         | 1,109.2        |                | 9.5         | 1,155.6        |    |
| Residual Fuel             |  | 71.0           | 131.5           | 271.3           | 240.4           | 189.3        | 903.5           |  | 5.3          | 9.9          | 20.4           | 18.1           | 14.2        | 67.8           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 0.1             |                 |                 |              | 0.1             |  |              | 0.0          |                |                |             | 0.0            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 76.5            |                 |                 |              | 76.5            |  |              | 5.4          |                |                |             | 5.4            |    |
| Petroleum Coke            |  | 0.3            | 645.7           |                 | 146.4           |              | 792.3           |  | 0.0          | 65.9         |                | 14.9           |             | 80.9           |    |
| Still Gas                 |  |                | 1,423.0         |                 |                 |              | 1,423.0         |  |              | 94.9         |                |                |             | 94.9           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (53.7)          |                 |                 |              | (53.7)          |  |              | (4.0)        |                |                |             | (4.0)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 50.6            |              | 50.6            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>6,211.6</b>                           | <b>3,944.2</b> | <b>12,646.8</b> | <b>25,034.6</b> | <b>27,851.8</b> | <b>556.0</b> | <b>76,245.1</b> | <b>347.6</b>   | <b>223.6</b> | <b>802.8</b> | <b>1,791.8</b> | <b>2,360.1</b> | <b>40.8</b> | <b>5,566.7</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values).

<sup>b</sup> Adjustments are subtracted from total consumption estimates and include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

Note: Parentheses indicate negative values. NE (Not Estimated); NA (Not Available).

**Table A-18: 2007 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>7.8</b>                               | <b>70.0</b>    | <b>1,130.8</b>  | <b>NE</b>       | <b>20,807.7</b> | <b>36.9</b>  | <b>22,053.2</b> | <b>0.7</b>   | <b>6.7</b>   | <b>107.0</b> | <b>NE</b>      | <b>1,987.3</b> | <b>3.4</b>  | <b>2,105.1</b> |    |
| Residential Coal          | 7.8                                      |                |                 |                 |                 |              | 7.8             | 0.7  |              |              |                |                |             | 0.7            |    |
| Commercial Coal           |  | 70.0           |                 |                 |                 |              | 70.0            |  | 6.7          |              |                |                |             | 6.7            |    |
| Industrial Other Coal     |  |                | 1,130.8         |                 |                 |              | 1,130.8         |  |              | 107.0        |                |                |             | 107.0          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 20,807.7        |              | 20,807.7        |  |              |              |                | 1,987.3        |             | 1,987.3        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | <b>4,835.4</b>                           | <b>3,085.1</b> | <b>7,521.3</b>  | <b>663.5</b>    | <b>7,005.2</b>  | <b>26.7</b>  | <b>23,137.2</b> | <b>256.3</b>   | <b>163.5</b> | <b>398.6</b> | <b>35.2</b>    | <b>371.3</b>   | <b>1.4</b>  | <b>1,226.3</b> |    |
| <b>Total Petroleum</b>    | <b>1,220.1</b>                           | <b>679.8</b>   | <b>4,584.7</b>  | <b>25,578.2</b> | <b>647.8</b>    | <b>549.8</b> | <b>33,260.3</b> | <b>84.3</b>  | <b>48.6</b>  | <b>342.2</b> | <b>1,851.0</b> | <b>52.9</b>    | <b>40.3</b> | <b>2,419.3</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 31.6            |                 |              | 31.6            |  |              |              | 2.2            |                |             | 2.2            |    |
| Distillate Fuel Oil       | 692.5                                    | 366.2          | 1,177.2         | 6,394.7         | 88.7            | 136.5        | 8,855.7         | 51.2   | 27.1         | 87.1         | 472.9          | 6.6            | 10.1        | 654.9          |    |
| Jet Fuel                  |  |                |                 | 2,485.0         | NA              | 55.5         | 2,540.4         |  |              |              | 179.5          |                | 4.0         | 183.5          |    |
| Kerosene                  | 43.9                                     | 9.2            | 13.4            |                 |                 | 5.2          | 71.8            | 3.2  | 0.7          | 1.0          |                |                | 0.4         | 5.3            |    |
| LPG                       | 483.7                                    | 121.4          | 379.1           | 21.9            |                 | 11.6         | 1,017.7         | 29.8   | 7.5          | 23.4         | 1.4            |                | 0.7         | 62.8           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 107.3          | 536.9           | 16,258.9        |                 | 155.1        | 17,058.1        |  | 7.7          | 38.5         | 1,166.0        |                | 11.1        | 1,223.4        |    |
| Residual Fuel             |  | 75.4           | 130.4           | 386.1           | 396.6           | 185.9        | 1,174.4         |  | 5.7          | 9.8          | 29.0           | 29.8           | 14.0        | 88.2           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 1.8             |                 |                 |              | 1.8             |  |              | 0.1          |                |                |             | 0.1            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 89.7            |                 |                 |              | 89.7            |  |              | 6.3          |                |                |             | 6.3            |    |
| Petroleum Coke            |  | 0.4            | 708.4           |                 | 162.6           |              | 871.3           |  | 0.0          | 72.3         |                | 16.6           |             | 89.0           |    |
| Still Gas                 |  |                | 1,482.6         |                 |                 |              | 1,482.6         |  |              | 98.9         |                |                |             | 98.9           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 65.2            |                 |                 |              | 65.2            |  |              | 4.8          |                |                |             | 4.8            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>49.9</b>     |              | <b>49.9</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,063.3</b>                           | <b>3,834.9</b> | <b>13,236.8</b> | <b>26,241.6</b> | <b>28,510.7</b> | <b>613.3</b> | <b>78,500.7</b> | <b>341.3</b>   | <b>218.8</b> | <b>847.9</b> | <b>1,886.1</b> | <b>2,411.9</b> | <b>45.1</b> | <b>5,751.2</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-19: 2006 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | 6.4                                      | 64.8           | 1,188.8         | NE              | 20,461.9        | 36.9         | 21,758.7        | 0.6  | 6.2          | 112.6        | NE             | 1,953.7        | 3.4         | 2,076.6        |    |
| Residential Coal          | 6.4                                      |                |                 |                 |                 |              | 6.4             | 0.6  |              |              |                |                |             | 0.6            |    |
| Commercial Coal           |  | 64.8           |                 |                 |                 |              | 64.8            |  | 6.2          |              |                |                |             | 6.2            |    |
| Industrial Other Coal     |  |                | 1,188.8         |                 |                 |              | 1,188.8         |  |              | 112.6        |                |                |             | 112.6          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 20,461.9        |              | 20,461.9        |  |              |              |                | 1,953.7        |             | 1,953.7        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 36.9         | 36.9            |  |              |              |                |                | 3.4         | 3.4            |    |
| <b>Natural Gas</b>        | 4,475.9                                  | 2,901.7        | 7,323.2         | 625.0           | 6,375.1         | 26.14        | 21,727.0        | 237.3  | 153.8        | 388.2        | 33.1           | 338.0          | 1.4         | 1,151.8        |    |
| <b>Total Petroleum</b>    | 1,202.3                                  | 677.6          | 4,709.0         | 25,651.7        | 637.0           | 619.9        | 33,497.4        | 83.4   | 48.5         | 351.7        | 1,849.5        | 53.2           | 45.4        | 2,431.8        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 33.4            |                 |              | 33.4            |  |              |              | 2.3            |                |             | 2.3            |    |
| Distillate Fuel Oil       | 690.4                                    | 389.0          | 1,194.7         | 6,334.2         | 73.4            | 90.2         | 8,771.8         | 51.1   | 28.8         | 88.4         | 468.5          | 5.4            | 6.7         | 648.7          |    |
| Jet Fuel                  |  |                |                 | 2,523.8         | NA              | 76.1         | 2,599.9         |  |              |              | 182.3          |                | 5.5         | 187.8          |    |
| Kerosene                  | 66.4                                     | 15.2           | 29.6            |                 |                 | 4.4          | 115.5           | 4.9  | 1.1          | 2.2          |                |                | 0.3         | 8.5            |    |
| LPG                       | 445.5                                    | 123.2          | 369.7           | 27.5            |                 | 6.6          | 972.5           | 27.5   | 7.6          | 22.8         | 1.7            |                | 0.4         | 60.0           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 74.7           | 577.2           | 16,426.4        |                 | 187.9        | 17,266.1        |  | 5.3          | 41.2         | 1,171.8        |                | 13.4        | 1,231.7        |    |
| Residual Fuel             |  | 75.3           | 176.4           | 306.3           | 360.5           | 254.8        | 1,173.3         |  | 5.7          | 13.2         | 23.0           | 27.1           | 19.1        | 88.1           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 0.6             |                 |                 |              | 0.6             |  |              | 0.0          |                |                |             | 0.0            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 70.0            |                 |                 |              | 70.0            |  |              | 4.9          |                |                |             | 4.9            |    |
| Petroleum Coke            |  | 0.3            | 724.3           |                 | 203.0           |              | 927.6           |  | 0.0          | 74.0         |                | 20.7           |             | 94.7           |    |
| Still Gas                 |  |                | 1,496.2         |                 |                 |              | 1,496.2         |  |              | 99.8         |                |                |             | 99.8           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 70.3            |                 |                 |              | 70.3            |  |              | 5.2          |                |                |             | 5.2            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 49.7            |              | 49.7            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>5,684.6</b>                           | <b>3,644.0</b> | <b>13,221.0</b> | <b>26,276.6</b> | <b>27,523.7</b> | <b>682.9</b> | <b>77,032.8</b> | <b>321.3</b>   | <b>208.6</b> | <b>852.6</b> | <b>1,882.6</b> | <b>2,345.3</b> | <b>50.2</b> | <b>5,660.6</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-20: 2005 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>8.4</b>                               | <b>97.0</b>    | <b>1,219.1</b>  | <b>NE</b>       | <b>20,737.2</b> | <b>32.7</b>  | <b>22,094.5</b> | <b>0.8</b>   | <b>9.3</b>   | <b>115.3</b> | <b>NE</b>      | <b>1,983.8</b> | <b>3.0</b>  | <b>2,112.3</b> |    |
| Residential Coal          | 8.4                                      |                |                 |                 |                 |              | 8.4             | 0.8  |              |              |                |                |             | 0.8            |    |
| Commercial Coal           |  | 97.0           |                 |                 |                 |              | 97.0            |  | 9.3          |              |                |                |             | 9.3            |    |
| Industrial Other Coal     |  |                | 1,219.1         |                 |                 |              | 1,219.1         |  |              | 115.3        |                |                |             | 115.3          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 20,737.2        |              | 20,737.2        |  |              |              |                | 1,983.8        |             | 1,983.8        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 32.7         | 32.7            |  |              |              |                |                | 3.0         | 3.0            |    |
| <b>Natural Gas</b>        | <b>4,946.4</b>                           | <b>3,073.2</b> | <b>7,329.7</b>  | <b>623.9</b>    | <b>6,014.5</b>  | <b>24.3</b>  | <b>22,012.0</b> | <b>262.2</b>   | <b>162.9</b> | <b>388.5</b> | <b>33.1</b>    | <b>318.8</b>   | <b>1.3</b>  | <b>1,166.7</b> |    |
| <b>Total Petroleum</b>    | <b>1,368.3</b>                           | <b>715.6</b>   | <b>4,327.7</b>  | <b>25,799.2</b> | <b>1,222.1</b>  | <b>622.7</b> | <b>34,055.5</b> | <b>94.9</b>  | <b>51.3</b>  | <b>324.2</b> | <b>1,854.0</b> | <b>97.9</b>    | <b>45.6</b> | <b>2,467.8</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 35.4            |                 |              | 35.4            |  |              |              | 2.4            |                |             | 2.4            |    |
| Distillate Fuel Oil       | 771.6                                    | 404.2          | 1,127.5         | 6,186.2         | 114.5           | 121.3        | 8,725.4         | 57.1   | 29.9         | 83.4         | 457.5          | 8.5            | 9.0         | 645.3          |    |
| Jet Fuel                  |  |                |                 | 2,621.7         | NA              | 66.0         | 2,687.7         |  |              |              | 189.3          |                | 4.8         | 194.1          |    |
| Kerosene                  | 83.8                                     | 21.6           | 39.1            |                 |                 | 5.8          | 150.2           | 6.1  | 1.6          | 2.9          |                |                | 0.4         | 11.0           |    |
| LPG                       | 512.9                                    | 131.4          | 349.6           | 28.2            |                 | 0.8          | 1,022.8         | 31.7   | 8.1          | 21.6         | 1.7            |                | 0.0         | 63.1           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 42.3           | 328.9           | 16,671.3        |                 | 193.6        | 17,236.1        |  | 3.0          | 23.4         | 1,183.7        |                | 13.7        | 1,223.8        |    |
| Residual Fuel             |  | 115.8          | 237.4           | 256.4           | 876.5           | 235.2        | 1,721.3         |  | 8.7          | 17.8         | 19.3           | 65.8           | 17.7        | 129.3          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 8.3             |                 |                 |              | 8.3             |  |              | 0.6          |                |                |             | 0.6            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 98.1            |                 |                 |              | 98.1            |  |              | 6.9          |                |                |             | 6.9            |    |
| Petroleum Coke            |  | 0.3            | 706.6           |                 | 231.1           |              | 938.0           |  | 0.0          | 72.1         |                | 23.6           |             | 95.8           |    |
| Still Gas                 |  |                | 1,429.4         |                 |                 |              | 1,429.4         |  |              | 95.4         |                |                |             | 95.4           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | 2.8             |                 |                 |              | 2.8             |  |              | 0.2          |                |                |             | 0.2            |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>50.1</b>     |              | <b>50.1</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,323.1</b>                           | <b>3,885.8</b> | <b>12,876.6</b> | <b>26,423.1</b> | <b>28,024.0</b> | <b>679.7</b> | <b>78,212.2</b> | <b>357.8</b>   | <b>223.5</b> | <b>828.0</b> | <b>1,887.0</b> | <b>2,400.9</b> | <b>49.9</b> | <b>5,747.1</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-21: 2004 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10              | 11           | 12             | 13             | 14          | 15             |            |                |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|-----------------|--------------|----------------|----------------|-------------|----------------|------------|----------------|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |                 |              |                |                |             |                |            |                |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.            | Trans.       | Elec.          | Terr.          | Total       |                |            |                |
| <b>Total Coal</b>         | <b>11.4</b>                              | <b>102.9</b>   | <b>1,262.0</b>  |                 |                 | <b>NE</b>    | <b>20,305.0</b> |  | <b>32.0</b>  | <b>21,713.3</b> | <b>1.1</b>   | <b>9.8</b>     | <b>118.3</b>   | <b>NE</b>   | <b>1,943.1</b> | <b>2.9</b> | <b>2,075.1</b> |
| Residential Coal          | 11.4                                     |                |                 |                 |                 |              |                 |  |              | 11.4            | 1.1          |                |                |             |                |            | 1.1            |
| Commercial Coal           |  | 102.9          |                 |                 |                 |              |                 |  |              | 102.9           | 9.8          |                |                |             |                |            | 9.8            |
| Industrial Other Coal     |  |                | 1,262.0         |                 |                 |              |                 |  |              | 1,262.0         |              | 118.3          |                |             |                |            | 118.3          |
| Transportation Coal       |  |                |                 |                 |                 | NE           |                 |  |              | NE              |              |                | NE             |             |                |            | NE             |
| Electric Power Coal       |  |                |                 |                 |                 |              | 20,305.0        |  |              | 20,305.0        |              |                |                |             | 1,943.1        |            | 1,943.1        |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 |              |                 |  | 32.0         | 32.0            |              |                |                |             |                | 2.9        | 2.9            |
| <b>Natural Gas</b>        | <b>4,980.8</b>                           | <b>3,201.0</b> | <b>7,913.5</b>  | <b>602.0</b>    | <b>5,594.9</b>  | <b>24.6</b>  | <b>22,316.9</b> |  | <b>264.1</b> | <b>169.7</b>    | <b>419.6</b> | <b>31.9</b>    | <b>296.7</b>   | <b>1.3</b>  | <b>1,183.4</b> |            |                |
| <b>Total Petroleum</b>    | <b>1,467.8</b>                           | <b>764.3</b>   | <b>4,197.3</b>  | <b>25,533.5</b> | <b>1,201.0</b>  | <b>653.1</b> | <b>33,817.0</b> |  | <b>102.2</b> | <b>54.7</b>     | <b>314.4</b> | <b>1,835.2</b> | <b>95.8</b>    | <b>47.9</b> | <b>2,450.3</b> |            |                |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Aviation Gasoline         |  |                |                 |                 | 31.2            |              |                 |  |              | 31.2            |              |                | 2.2            |             |                |            | 2.2            |
| Distillate Fuel Oil       | 871.3                                    | 443.6          | 1,130.6         | 5,910.4         | 111.2           | 134.4        | 8,601.5         |  | 64.4         | 32.8            | 83.6         | 437.1          | 8.2            | 9.9         | 636.1          |            |                |
| Jet Fuel                  |  |                |                 |                 |                 | 2,584.8      | NA              |  |              |                 |              | 186.7          |                | 5.0         | 191.6          |            |                |
| Kerosene                  | 84.8                                     | 20.5           | 28.2            |                 |                 |              |                 |  | 6.2          | 1.5             | 2.1          |                |                | 0.4         | 10.2           |            |                |
| LPG                       | 511.7                                    | 152.0          | 372.7           | 19.1            |                 |              |                 |  | 31.6         | 9.4             | 23.0         | 1.2            |                | 0.0         | 65.2           |            |                |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Motor Gasoline            |  | 25.5           | 212.4           | 16,801.6        |                 |              |                 |  |              |                 | 1.8          | 15.1           | 1,194.1        |             | 14.1           | 1,225.1    |                |
| Residual Fuel             |  | 122.5          | 204.7           | 186.4           | 879.0           | 245.0        | 1,637.6         |  |              | 9.2             | 15.4         | 14.0           | 66.0           | 18.4        | 123.0          |            |                |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| AvGas Blend Components    |  |                | 10.6            |                 |                 |              |                 |  |              |                 |              | 0.7            |                |             |                | 0.7        |                |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Pentanes Plus             |  |                | 111.2           |                 |                 |              |                 |  |              |                 |              | 7.8            |                |             |                |            | 7.8            |
| Petroleum Coke            |  | 0.3            | 719.1           |                 | 210.8           |              |                 |  |              |                 | 0.0          | 73.4           | 21.5           |             |                |            | 95.0           |
| Still Gas                 |  |                | 1,483.3         |                 |                 |              |                 |  |              |                 |              | 99.0           |                |             |                |            | 99.0           |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| Unfinished Oils           |  |                | (75.6)          |                 |                 |              |                 |  |              |                 |              | (5.6)          |                |             |                |            | (5.6)          |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |                 |              |                |                |             |                |            |                |
| <b>Geothermal</b>         |  |                |                 |                 |                 |              | <b>50.5</b>     |  |              | <b>50.5</b>     |              |                | <b>0.4</b>     |             |                |            | <b>0.4</b>     |
| <b>Total (All Fuels)</b>  | <b>6,460.1</b>                           | <b>4,068.2</b> | <b>13,372.8</b> | <b>26,135.5</b> | <b>27,151.5</b> | <b>709.7</b> | <b>77,897.7</b> |  | <b>367.4</b> | <b>234.2</b>    | <b>852.3</b> | <b>1,867.1</b> | <b>2,335.9</b> | <b>52.1</b> | <b>5,709.2</b> |            |                |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-22: 2003 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | 12.2                                     | 82.0           | 1,248.8         | NE              | 20,184.7        | 33.9         | 21,561.7        | 1.2  | 7.8          | 117.0        | NE             | 1,931.0        | 3.1         | 2,060.1        |    |
| Residential Coal          | 12.2                                     |                |                 |                 |                 |              | 12.2            | 1.2  |              |              |                |                |             | 1.2            |    |
| Commercial Coal           |  | 82.0           |                 |                 |                 |              | 82.0            |  | 7.8          |              |                |                |             | 7.8            |    |
| Industrial Other Coal     |  |                | 1,248.8         |                 |                 |              | 1,248.8         |  |              | 117.0        |                |                |             | 117.0          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 20,184.7        |              | 20,184.7        |  |              |              |                | 1,931.0        |             | 1,931.0        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 33.9         | 33.9            |  |              |              |                |                | 3.1         | 3.1            |    |
| <b>Natural Gas</b>        | 5,209.4                                  | 3,260.9        | 7,845.1         | 627.4           | 5,246.2         | 26.9         | 22,216.0        | 275.9  | 172.7        | 415.4        | 33.2           | 277.8          | 1.4         | 1,176.4        |    |
| <b>Total Petroleum</b>    | 1,468.3                                  | 764.6          | 3,959.1         | 24,919.1        | 1,204.8         | 621.7        | 32,937.5        | 101.9  | 54.7         | 297.2        | 1,789.0        | 95.0           | 45.3        | 2,383.2        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 30.2            |                 |              | 30.2            |  |              |              | 2.1            |                |             | 2.1            |    |
| Distillate Fuel Oil       | 853.5                                    | 454.2          | 1,058.2         | 5,704.9         | 160.8           | 120.5        | 8,352.1         | 63.1   | 33.6         | 78.3         | 421.9          | 11.9           | 8.9         | 617.7          |    |
| Jet Fuel                  |  |                |                 | 2,482.5         | NA              | 76.1         | 2,558.5         |  |              |              | 179.3          |                | 5.5         | 184.8          |    |
| Kerosene                  | 70.3                                     | 18.6           | 24.1            |                 |                 | 10.7         | 123.7           | 5.1  | 1.4          | 1.8          |                |                | 0.8         | 9.1            |    |
| LPG                       | 544.5                                    | 156.9          | 326.8           | 17.9            |                 | 10.5         | 1,056.6         | 33.7   | 9.7          | 20.2         | 1.1            |                | 0.7         | 65.3           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 23.5           | 126.9           | 16,584.5        |                 | 209.9        | 16,944.8        |  | 1.7          | 9.0          | 1,177.2        |                | 14.9        | 1,202.8        |    |
| Residual Fuel             |  | 111.1          | 176.4           | 99.1            | 869.4           | 193.9        | 1,450.0         |  | 8.3          | 13.2         | 7.4            | 65.3           | 14.6        | 108.9          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 7.5             |                 |                 |              | 7.5             |  |              | 0.5          |                |                |             | 0.5            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 110.4           |                 |                 |              | 110.4           |  |              | 7.7          |                |                |             | 7.7            |    |
| Petroleum Coke            |  | 0.3            | 701.9           |                 | 174.7           |              | 876.8           |  | 0.0          | 71.7         |                | 17.8           |             | 89.5           |    |
| Still Gas                 |  |                | 1,477.3         |                 |                 |              | 1,477.3         |  |              | 98.6         |                |                |             | 98.6           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (50.4)          |                 |                 |              | (50.4)          |  |              | (3.7)        |                |                |             | (3.7)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 49.2            |              | 49.2            |  |              |              |                | 0.4            |             | 0.4            |    |
| <b>Total (All Fuels)</b>  | <b>6,689.9</b>                           | <b>4,107.5</b> | <b>13,052.9</b> | <b>25,546.6</b> | <b>26,685.0</b> | <b>682.4</b> | <b>76,764.4</b> | <b>378.9</b>   | <b>235.2</b> | <b>829.6</b> | <b>1,822.3</b> | <b>2,304.2</b> | <b>49.9</b> | <b>5,620.0</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-23: 2002 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>12.2</b>                              | <b>89.8</b>    | <b>1,243.7</b>  | <b>NE</b>       | <b>19,782.8</b> | <b>10.8</b>  | <b>21,139.3</b> | <b>1.2</b>   | <b>8.6</b>   | <b>116.6</b> | <b>NE</b>      | <b>1,889.9</b> | <b>1.0</b>  | <b>2,017.2</b> |    |
| Residential Coal          | 12.2                                     |                |                 |                 |                 |              | 12.2            | 1.2  |              |              |                |                |             | 1.2            |    |
| Commercial Coal           |  | 89.8           |                 |                 |                 |              | 89.8            |  | 8.6          |              |                |                |             | 8.6            |    |
| Industrial Other Coal     |  |                | 1,243.7         |                 |                 |              | 1,243.7         |  |              | 116.6        |                |                |             | 116.6          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 19,782.8        |              | 19,782.8        |  |              |              |                | 1,889.9        |             | 1,889.9        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.8         | 10.8            |  |              |              |                |                | 1.0         | 1.0            |    |
| <b>Natural Gas</b>        | <b>4,995.0</b>                           | <b>3,212.5</b> | <b>8,086.3</b>  | <b>698.9</b>    | <b>5,766.8</b>  | <b>22.8</b>  | <b>22,782.3</b> | <b>264.7</b>   | <b>170.3</b> | <b>428.6</b> | <b>37.0</b>    | <b>305.7</b>   | <b>1.2</b>  | <b>1,207.5</b> |    |
| <b>Total Petroleum</b>    | <b>1,360.9</b>                           | <b>647.4</b>   | <b>3,794.7</b>  | <b>24,961.9</b> | <b>961.2</b>    | <b>556.9</b> | <b>32,283.0</b> | <b>94.1</b>  | <b>46.2</b>  | <b>284.6</b> | <b>1,793.6</b> | <b>76.8</b>    | <b>40.6</b> | <b>2,335.9</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 33.7            |                 |              | 33.7            |  |              |              | 2.3            |                |             | 2.3            |    |
| Distillate Fuel Oil       | 763.8                                    | 394.5          | 1,051.2         | 5,590.0         | 127.3           | 92.8         | 8,019.6         | 56.5   | 29.2         | 77.7         | 413.4          | 9.4            | 6.9         | 593.1          |    |
| Jet Fuel                  |  |                |                 | 2,565.5         | NA              | 61.8         | 2,627.3         |  |              |              | 185.3          |                | 4.5         | 189.7          |    |
| Kerosene                  | 59.9                                     | 15.9           | 13.8            |                 |                 | 8.2          | 97.9            | 4.4  | 1.2          | 1.0          |                |                | 0.6         | 7.2            |    |
| LPG                       | 537.1                                    | 140.8          | 393.3           | 14.3            |                 | 11.2         | 1,096.7         | 33.2   | 8.7          | 24.3         | 0.9            |                | 0.7         | 67.8           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 16.2           | 110.8           | 16,530.5        |                 | 189.5        | 16,847.0        |  | 1.1          | 7.9          | 1,174.6        |                | 13.5        | 1,197.1        |    |
| Residual Fuel             |  | 79.8           | 146.1           | 227.9           | 658.7           | 193.6        | 1,306.1         |  | 6.0          | 11.0         | 17.1           | 49.5           | 14.5        | 98.1           |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 7.5             |                 |                 |              | 7.5             |  |              | 0.5          |                |                |             | 0.5            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 111.9           |                 |                 |              | 111.9           |  |              | 7.8          |                |                |             | 7.8            |    |
| Petroleum Coke            |  | 0.2            | 696.3           |                 | 175.2           |              | 871.7           |  | 0.0          | 71.1         |                | 17.9           |             | 89.0           |    |
| Still Gas                 |  |                | 1,399.4         |                 |                 |              | 1,399.4         |  |              | 93.4         |                |                |             | 93.4           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (135.7)         |                 |                 |              | (135.7)         |  |              | (10.1)       |                |                |             | (10.1)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>49.4</b>     |              | <b>49.4</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,368.1</b>                           | <b>3,949.7</b> | <b>13,124.6</b> | <b>25,660.9</b> | <b>26,560.2</b> | <b>590.5</b> | <b>76,254.1</b> | <b>360.0</b>   | <b>225.0</b> | <b>829.8</b> | <b>1,830.6</b> | <b>2,272.7</b> | <b>42.8</b> | <b>5,560.9</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-24: 2001 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>12.0</b>                              | <b>96.9</b>    | <b>1,358.4</b>  | <b>NE</b>       | <b>19,613.7</b> | <b>3.8</b>   | <b>21,084.8</b> | <b>1.1</b>   | <b>9.2</b>   | <b>127.8</b> | <b>NE</b>      | <b>1,869.8</b> | <b>0.4</b>  | <b>2,008.4</b> |    |
| Residential Coal          | 12.0                                     |                |                 |                 |                 |              | 12.0            | 1.1  |              |              |                |                |             | 1.1            |    |
| Commercial Coal           |  | 96.9           |                 |                 |                 |              | 96.9            |  | 9.2          |              |                |                |             | 9.2            |    |
| Industrial Other Coal     |  |                | 1,358.4         |                 |                 |              | 1,358.4         |  |              | 127.8        |                |                |             | 127.8          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 19,613.7        |              | 19,613.7        |  |              |              |                | 1,869.8        |             | 1,869.8        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 3.8          | 3.8             |  |              |              |                |                | 0.4         | 0.4            |    |
| <b>Natural Gas</b>        | <b>4,889.0</b>                           | <b>3,097.3</b> | <b>7,949.0</b>  | <b>658.0</b>    | <b>5,458.1</b>  | <b>22.9</b>  | <b>22,074.3</b> | <b>259.1</b>   | <b>164.2</b> | <b>421.3</b> | <b>34.9</b>    | <b>289.3</b>   | <b>1.2</b>  | <b>1,170.0</b> |    |
| <b>Total Petroleum</b>    | <b>1,464.9</b>                           | <b>720.4</b>   | <b>3,919.6</b>  | <b>24,453.7</b> | <b>1,276.4</b>  | <b>632.4</b> | <b>32,467.4</b> | <b>101.9</b>   | <b>51.6</b>  | <b>293.8</b> | <b>1,754.6</b> | <b>98.4</b>    | <b>46.2</b> | <b>2,346.5</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 34.9            |                 |              | 34.9            |  |              |              | 2.4            |                |             | 2.4            |    |
| Distillate Fuel Oil       | 844.1                                    | 472.4          | 1,184.5         | 5,411.3         | 170.3           | 109.4        | 8,192.1         | 62.4   | 34.9         | 87.6         | 400.2          | 12.6           | 8.1         | 605.9          |    |
| Jet Fuel                  |  |                |                 | 2,626.3         | NA              | 98.9         | 2,725.2         |  |              |              | 189.7          |                | 7.1         | 196.8          |    |
| Kerosene                  | 95.1                                     | 31.4           | 23.2            |                 |                 | 0.9          | 150.6           | 7.0  | 2.3          | 1.7          |                |                | 0.1         | 11.0           |    |
| LPG                       | 525.7                                    | 142.7          | 372.1           | 13.7            |                 | 7.0          | 1,061.2         | 32.5   | 8.8          | 23.0         | 0.8            |                | 0.4         | 65.6           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 3.8            | 30.3            | 16,208.0        |                 | 187.8        | 16,429.9        |  | 0.3          | 2.1          | 1,149.4        |                | 13.3        | 1,165.2        |    |
| Residual Fuel             |  | 69.9           | 146.7           | 159.5           | 1,002.8         | 228.4        | 1,607.2         |  | 5.2          | 11.0         | 12.0           | 75.3           | 17.2        | 120.7          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 6.1             |                 |                 |              | 6.1             |  |              | 0.4          |                |                |             | 0.4            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 131.6           |                 |                 |              | 131.6           |  |              | 9.2          |                |                |             | 9.2            |    |
| Petroleum Coke            |  | 0.2            | 683.3           |                 | 103.2           |              | 786.7           |  | 0.0          | 69.8         |                | 10.5           |             | 80.3           |    |
| Still Gas                 |  |                | 1,417.3         |                 |                 |              | 1,417.3         |  |              | 94.6         |                |                |             | 94.6           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (75.4)          |                 |                 |              | (75.4)          |  |              | (5.6)        |                |                |             | (5.6)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>46.9</b>     |              | <b>46.9</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,365.9</b>                           | <b>3,914.5</b> | <b>13,227.1</b> | <b>25,111.8</b> | <b>26,395.0</b> | <b>659.0</b> | <b>75,673.3</b> | <b>362.2</b>   | <b>225.0</b> | <b>843.0</b> | <b>1,789.4</b> | <b>2,257.9</b> | <b>47.8</b> | <b>5,525.2</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-25: 2000 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>11.4</b>                              | <b>91.9</b>    | <b>1,348.8</b>  | <b>NE</b>       | <b>20,220.2</b> | <b>10.3</b>  | <b>21,682.4</b> | <b>1.1</b>   | <b>8.8</b>   | <b>127.3</b> | <b>NE</b>      | <b>1,927.4</b> | <b>0.9</b>  | <b>2,065.5</b> |    |
| Residential Coal          | 11.4                                     |                |                 |                 |                 |              | 11.4            | 1.1  |              |              |                |                |             | 1.1            |    |
| Commercial Coal           |  | 91.9           |                 |                 |                 |              | 91.9            |  | 8.8          |              |                |                |             | 8.8            |    |
| Industrial Other Coal     |  |                | 1,348.8         |                 |                 |              | 1,348.8         |  |              | 127.3        |                |                |             | 127.3          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 20,220.2        |              | 20,220.2        |  |              |              |                | 1,927.4        |             | 1,927.4        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.3         | 10.3            |  |              |              |                |                | 0.9         | 0.9            |    |
| <b>Natural Gas</b>        | <b>5,104.6</b>                           | <b>3,251.5</b> | <b>8,656.0</b>  | <b>672.0</b>    | <b>5,293.4</b>  | <b>12.7</b>  | <b>22,990.2</b> | <b>270.7</b>   | <b>172.5</b> | <b>459.1</b> | <b>35.6</b>    | <b>280.8</b>   | <b>0.7</b>  | <b>1,219.4</b> |    |
| <b>Total Petroleum</b>    | <b>1,429.5</b>                           | <b>695.0</b>   | <b>3,577.9</b>  | <b>24,659.5</b> | <b>1,144.1</b>  | <b>471.8</b> | <b>31,977.9</b> | <b>99.0</b>  | <b>49.7</b>  | <b>267.7</b> | <b>1,769.8</b> | <b>88.4</b>    | <b>34.2</b> | <b>2,308.8</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 36.3            |                 |              | 36.3            |  |              |              | 2.5            |                |             | 2.5            |    |
| Distillate Fuel Oil       | 780.0                                    | 423.3          | 1,006.4         | 5,436.7         | 174.7           | 71.3         | 7,892.3         | 57.7   | 31.3         | 74.4         | 402.1          | 12.9           | 5.3         | 583.7          |    |
| Jet Fuel                  |  |                |                 | 2,700.3         | NA              | 74.1         | 2,774.3         |  |              |              | 195.0          |                | 5.3         | 200.4          |    |
| Kerosene                  | 94.6                                     | 29.7           | 15.6            |                 |                 | 2.4          | 142.2           | 6.9  | 2.2          | 1.1          |                |                | 0.2         | 10.4           |    |
| LPG                       | 554.9                                    | 150.4          | 468.7           | 11.9            |                 | 8.0          | 1,193.9         | 34.4   | 9.3          | 29.0         | 0.7            |                | 0.5         | 74.0           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 16,030.9        |                 | 185.2        | 16,216.1        |  |              |              | 1,136.2        |                | 13.1        | 1,149.3        |    |
| Residual Fuel             |  | 91.6           | 184.1           | 443.5           | 870.8           | 130.9        | 1,720.8         |  | 6.9          | 13.8         | 33.3           | 65.4           | 9.8         | 129.2          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 3.8             |                 |                 |              | 3.8             |  |              | 0.3          |                |                |             | 0.3            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 171.6           |                 |                 |              | 171.6           |  |              | 12.0         |                |                |             | 12.0           |    |
| Petroleum Coke            |  | 0.2            | 697.6           |                 | 98.6            |              | 796.4           |  | 0.0          | 71.2         |                | 10.1           |             | 81.3           |    |
| Still Gas                 |  |                | 1,431.2         |                 |                 |              | 1,431.2         |  |              | 95.5         |                |                |             | 95.5           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (401.2)         |                 |                 |              | (401.2)         |  |              | (29.7)       |                |                |             | (29.7)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>48.1</b>     |              | <b>48.1</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,545.4</b>                           | <b>4,038.5</b> | <b>13,582.7</b> | <b>25,331.5</b> | <b>26,705.8</b> | <b>494.7</b> | <b>76,698.6</b> | <b>370.8</b>   | <b>230.9</b> | <b>854.1</b> | <b>1,805.5</b> | <b>2,296.9</b> | <b>35.9</b> | <b>5,594.0</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-26: 1999 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10              | 11             | 12             | 13           | 14             | 15             |             |                |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|-----------------|----------------|----------------|--------------|----------------|----------------|-------------|----------------|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |                 |                |                |              |                |                |             |                |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.            | Trans.         | Elec.          | Terr.        | Total          |                |             |                |
| <b>Total Coal</b>         | <b>14.0</b>                              | <b>102.5</b>   | <b>1,372.8</b>  |                 |                 | <b>NE</b>    | <b>19,279.5</b> |  | <b>10.2</b>  | <b>20,779.0</b> | <b>1.3</b>     | <b>9.8</b>     | <b>129.9</b> | <b>NE</b>      | <b>1,836.4</b> | <b>0.9</b>  | <b>1,978.3</b> |
| Residential Coal          | 14.0                                     |                |                 |                 |                 |              |                 |  |              | 14.0            | 1.3            |                |              |                |                |             | 1.3            |
| Commercial Coal           |  | 102.5          |                 |                 |                 |              |                 |  |              | 102.5           | 9.8            |                |              |                |                |             | 9.8            |
| Industrial Other Coal     |  |                | 1,372.8         |                 |                 |              |                 |  |              | 1,372.8         |                | 129.9          |              |                |                |             | 129.9          |
| Transportation Coal       |  |                |                 |                 |                 | NE           |                 |  |              | NE              |                |                | NE           |                |                |             | NE             |
| Electric Power Coal       |  |                |                 |                 |                 |              | 19,279.5        |  |              | 19,279.5        |                |                |              | 1,836.4        |                |             | 1,836.4        |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 |              |                 |  | 10.2         | 10.2            |                |                |              |                |                | 0.9         | 0.9            |
| <b>Natural Gas</b>        | <b>4,834.9</b>                           | <b>3,115.0</b> | <b>8,424.6</b>  |                 | <b>675.3</b>    |              | <b>4,902.1</b>  |  |              | <b>21,952.0</b> | <b>256.3</b>   | <b>165.1</b>   | <b>446.6</b> | <b>35.8</b>    | <b>259.9</b>   |             | <b>1,163.8</b> |
| <b>Total Petroleum</b>    | <b>1,344.0</b>                           | <b>614.9</b>   | <b>3,580.6</b>  | <b>24,062.3</b> | <b>1,211.2</b>  |              | <b>461.1</b>    |  |              | <b>31,274.1</b> | <b>92.9</b>    | <b>43.8</b>    | <b>269.4</b> | <b>1,725.3</b> | <b>93.8</b>    | <b>33.5</b> | <b>2,258.9</b> |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Aviation Gasoline         |  |                |                 |                 | 39.2            |              |                 |  |              | 39.2            |                |                |              | 2.7            |                |             | 2.7            |
| Distillate Fuel Oil       | 706.9                                    | 374.4          | 986.1           | 5,245.8         | 140.0           | 79.4         | 7,532.5         | 52.3   | 27.7         | 72.9            | 388.0          | 10.4           | 5.9          | 557.1          |                |             |                |
| Jet Fuel                  |  |                |                 | 2,664.8         | NA              | 59.5         | 2,724.4         |  |              |                 | 192.5          |                | 4.3          | 196.8          |                |             |                |
| Kerosene                  | 111.2                                    | 26.9           | 12.8            |                 |                 | 3.7          | 154.7           | 8.1  | 2.0          | 0.9             |                |                | 0.3          | 11.3           |                |             |                |
| LPG                       | 526.0                                    | 140.2          | 395.9           | 14.3            |                 | 8.3          | 1,084.6         | 32.5   | 8.7          | 24.5            | 0.9            |                | 0.5          | 67.1           |                |             |                |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Motor Gasoline            |  |                |                 | 15,922.5        |                 | 164.1        | 16,086.6        |  |              |                 | 1,128.0        |                | 11.6         | 1,139.7        |                |             |                |
| Residual Fuel             |  | 73.3           | 150.9           | 175.7           | 958.7           | 146.0        | 1,504.6         |  | 5.5          | 11.3            | 13.2           | 72.0           | 11.0         | 113.0          |                |             |                |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| AvGas Blend Components    |  |                | 6.4             |                 |                 |              | 6.4             |  |              |                 | 0.4            |                |              | 0.4            |                |             |                |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Pentanes Plus             |  |                | 182.5           |                 |                 |              | 182.5           |  |              |                 | 12.8           |                |              | 12.8           |                |             |                |
| Petroleum Coke            |  | 0.1            | 719.8           |                 | 112.5           |              | 832.4           |  | 0.0          | 73.5            |                | 11.5           |              | 85.0           |                |             |                |
| Still Gas                 |  |                | 1,414.1         |                 |                 |              | 1,414.1         |  |              | 94.3            |                |                |              | 94.3           |                |             |                |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| Unfinished Oils           |  |                | (287.9)         |                 |                 |              | (287.9)         |  |              |                 | (21.3)         |                |              | (21.3)         |                |             |                |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |                 |                |                |              |                |                |             |                |
| <b>Geothermal</b>         |  |                |                 |                 | <b>50.6</b>     |              | <b>50.6</b>     |  |              |                 |                | <b>0.4</b>     |              | <b>0.4</b>     |                |             |                |
| <b>Total (All Fuels)</b>  | <b>6,192.9</b>                           | <b>3,832.5</b> | <b>13,378.0</b> | <b>24,737.6</b> | <b>25,443.4</b> | <b>471.3</b> | <b>74,055.7</b> | <b>350.6</b>   | <b>218.8</b> | <b>845.9</b>    | <b>1,761.1</b> | <b>2,190.5</b> | <b>34.5</b>  | <b>5,401.3</b> |                |             |                |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-27: 1998 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>11.5</b>                              | <b>93.4</b>    | <b>1,470.8</b>  | <b>NE</b>       | <b>19,215.7</b> | <b>10.5</b>  | <b>20,802.0</b> | <b>1.1</b>   | <b>8.9</b>   | <b>139.1</b> | <b>NE</b>      | <b>1,828.2</b> | <b>1.0</b>  | <b>1,978.3</b> |    |
| Residential Coal          | 11.5                                     |                |                 |                 |                 |              | 11.5            | 1.1  |              |              |                |                |             | 1.1            |    |
| Commercial Coal           |  | 93.4           |                 |                 |                 |              | 93.4            |  | 8.9          |              |                |                |             | 8.9            |    |
| Industrial Other Coal     |  |                | 1,470.8         |                 |                 |              | 1,470.8         |  |              | 139.1        |                |                |             | 139.1          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 19,215.7        |              | 19,215.7        |  |              |              |                | 1,828.2        |             | 1,828.2        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.5         | 10.5            |  |              |              |                |                | 1.0         | 1.0            |    |
| <b>Natural Gas</b>        | <b>4,646.1</b>                           | <b>3,083.0</b> | <b>8,826.0</b>  | <b>666.1</b>    | <b>4,674.9</b>  |              | <b>21,896.1</b> | <b>246.0</b>   | <b>163.3</b> | <b>467.4</b> | <b>35.3</b>    | <b>247.6</b>   |             | <b>1,159.5</b> |    |
| <b>Total Petroleum</b>    | <b>1,209.0</b>                           | <b>610.1</b>   | <b>3,473.3</b>  | <b>23,285.5</b> | <b>1,306.1</b>  | <b>445.5</b> | <b>30,329.5</b> | <b>84.1</b>  | <b>43.7</b>  | <b>262.6</b> | <b>1,671.4</b> | <b>101.3</b>   | <b>32.5</b> | <b>2,195.6</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 35.5            |                 |              | 35.5            |  |              |              | 2.5            |                |             | 2.5            |    |
| Distillate Fuel Oil       | 676.8                                    | 376.0          | 1,030.5         | 4,949.9         | 135.6           | 71.9         | 7,240.7         | 50.1   | 27.8         | 76.2         | 366.1          | 10.0           | 5.3         | 535.5          |    |
| Jet Fuel                  |  |                |                 | 2,608.0         | NA              | 59.9         | 2,667.8         |  |              |              | 188.4          |                | 4.3         | 192.7          |    |
| Kerosene                  | 108.3                                    | 31.2           | 22.1            |                 |                 | 6.3          | 167.8           | 7.9  | 2.3          | 1.6          |                |                | 0.5         | 12.3           |    |
| LPG                       | 423.9                                    | 117.6          | 271.6           | 17.6            |                 | 5.9          | 836.7           | 26.1   | 7.2          | 16.7         | 1.1            |                | 0.4         | 51.6           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 15,595.6        |                 | 160.4        | 15,756.0        |  |              |              | 1,107.5        |                | 11.4        | 1,118.9        |    |
| Residual Fuel             |  | 85.2           | 173.3           | 78.9            | 1,047.0         | 141.1        | 1,525.5         |  | 6.4          | 13.0         | 5.9            | 78.6           | 10.6        | 114.6          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 4.0             |                 |                 |              | 4.0             |  |              | 0.3          |                |                |             | 0.3            |    |
| Crude Oil                 |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 147.0           |                 |                 |              | 147.0           |  |              | 10.3         |                |                |             | 10.3           |    |
| Petroleum Coke            |  | 0.1            | 707.7           |                 | 123.6           |              | 831.4           |  | 0.0          | 72.3         |                | 12.6           |             | 84.9           |    |
| Still Gas                 |  |                | 1,431.0         |                 |                 |              | 1,431.0         |  |              | 95.5         |                |                |             | 95.5           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (313.9)         |                 |                 |              | (313.9)         |  |              | (23.3)       |                |                |             | (23.3)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>50.4</b>     |              | <b>50.4</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>5,866.6</b>                           | <b>3,786.5</b> | <b>13,770.1</b> | <b>23,951.6</b> | <b>25,247.1</b> | <b>456.1</b> | <b>73,078.0</b> | <b>331.2</b>   | <b>215.9</b> | <b>869.1</b> | <b>1,706.6</b> | <b>2,177.4</b> | <b>33.4</b> | <b>5,333.7</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-28: 1997 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>16.0</b>                              | <b>129.4</b>   | <b>1,457.6</b>  | <b>NE</b>       | <b>18,904.5</b> | <b>10.4</b>  | <b>20,518.0</b> | <b>1.5</b>   | <b>12.3</b>  | <b>137.6</b> | <b>NE</b>      | <b>1,797.0</b> | <b>1.0</b>  | <b>1,949.5</b> |    |
| Residential Coal          | 16.0                                     |                |                 |                 |                 |              | 16.0            | 1.5  |              |              |                |                |             | 1.5            |    |
| Commercial Coal           |  | 129.4          |                 |                 |                 |              | 129.4           |  | 12.3         |              |                |                |             | 12.3           |    |
| Industrial Other Coal     |  |                | 1,457.6         |                 |                 |              | 1,457.6         |  |              | 137.6        |                |                |             | 137.6          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 18,904.5        |              | 18,904.5        |  |              |              |                | 1,797.0        |             | 1,797.0        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.4         | 10.4            |  |              |              |                |                | 1.0         | 1.0            |    |
| <b>Natural Gas</b>        | <b>5,092.9</b>                           | <b>3,285.3</b> | <b>9,032.5</b>  | <b>780.3</b>    | <b>4,125.5</b>  |              | <b>22,316.6</b> | <b>270.1</b>   | <b>174.2</b> | <b>479.0</b> | <b>41.4</b>    | <b>218.8</b>   |             | <b>1,183.4</b> |    |
| <b>Total Petroleum</b>    | <b>1,335.0</b>                           | <b>655.9</b>   | <b>3,898.2</b>  | <b>22,698.3</b> | <b>926.7</b>    | <b>445.4</b> | <b>29,959.4</b> | <b>93.1</b>  | <b>47.1</b>  | <b>290.2</b> | <b>1,628.8</b> | <b>72.2</b>    | <b>32.4</b> | <b>2,163.9</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 39.7            |                 |              | 39.7            |  |              |              | 2.7            |                |             | 2.7            |    |
| Distillate Fuel Oil       | 787.3                                    | 399.6          | 1,059.3         | 4,797.9         | 110.5           | 81.6         | 7,236.3         | 58.2   | 29.6         | 78.3         | 354.8          | 8.2            | 6.0         | 535.2          |    |
| Jet Fuel                  |  |                |                 | 2,553.8         | NA              | 62.1         | 2,615.9         |  |              |              | 184.4          |                | 4.5         | 188.9          |    |
| Kerosene                  | 92.9                                     | 24.6           | 18.8            |                 |                 | 4.0          | 140.3           | 6.8  | 1.8          | 1.4          |                |                | 0.3         | 10.3           |    |
| LPG                       | 454.8                                    | 120.2          | 429.9           | 14.2            |                 | 6.5          | 1,025.7         | 28.1   | 7.4          | 26.5         | 0.9            |                | 0.4         | 63.3           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 15,156.2        |                 | 160.1        | 15,316.2        |  |              |              | 1,075.6        |                | 11.4        | 1,087.0        |    |
| Residual Fuel             |  | 111.2          | 240.1           | 136.5           | 714.6           | 131.1        | 1,333.5         |  | 8.4          | 18.0         | 10.3           | 53.7           | 9.8         | 100.1          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 9.1             |                 |                 |              | 9.1             |  |              | 0.6          |                |                |             | 0.6            |    |
| Crude Oil                 |  |                | 4.6             |                 |                 |              | 4.6             |  |              | 0.3          |                |                |             | 0.3            |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 164.5           |                 |                 |              | 164.5           |  |              | 11.5         |                |                |             | 11.5           |    |
| Petroleum Coke            |  | 0.1            | 639.9           |                 | 101.6           |              | 741.6           |  | 0.0          | 65.3         |                | 10.4           |             | 75.7           |    |
| Still Gas                 |  |                | 1,435.0         |                 |                 |              | 1,435.0         |  |              | 95.7         |                |                |             | 95.7           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (102.9)         |                 |                 |              | (102.9)         |  |              | (7.6)        |                |                |             | (7.6)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>50.2</b>     |              | <b>50.2</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,443.9</b>                           | <b>4,070.6</b> | <b>14,388.4</b> | <b>23,478.6</b> | <b>24,007.0</b> | <b>455.8</b> | <b>72,844.3</b> | <b>364.7</b>   | <b>233.6</b> | <b>906.8</b> | <b>1,670.1</b> | <b>2,088.4</b> | <b>33.4</b> | <b>5,297.1</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-29: 1996 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>16.6</b>                              | <b>121.6</b>   | <b>1,454.9</b>  | <b>NE</b>       | <b>18,429.0</b> | <b>10.3</b>  | <b>20,032.4</b> | <b>1.6</b>   | <b>11.6</b>  | <b>137.4</b> | <b>NE</b>      | <b>1,752.4</b> | <b>1.0</b>  | <b>1,903.9</b> |    |
| Residential Coal          | 16.6                                     |                |                 |                 |                 |              | 16.6            | 1.6  |              |              |                |                |             | 1.6            |    |
| Commercial Coal           |  | 121.6          |                 |                 |                 |              | 121.6           |  | 11.6         |              |                |                |             | 11.6           |    |
| Industrial Other Coal     |  |                | 1,454.9         |                 |                 |              | 1,454.9         |  |              | 137.4        |                |                |             | 137.4          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 18,429.0        |              | 18,429.0        |  |              |              |                | 1,752.4        |             | 1,752.4        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.3         | 10.3            |  |              |              |                |                | 1.0         | 1.0            |    |
| <b>Natural Gas</b>        | <b>5,354.4</b>                           | <b>3,226.3</b> | <b>9,020.3</b>  | <b>736.9</b>    | <b>3,862.4</b>  |              | <b>22,200.4</b> | <b>283.9</b>   | <b>171.1</b> | <b>478.3</b> | <b>39.1</b>    | <b>204.8</b>   |             | <b>1,177.2</b> |    |
| <b>Total Petroleum</b>    | <b>1,398.0</b>                           | <b>719.1</b>   | <b>3,914.4</b>  | <b>22,506.2</b> | <b>817.3</b>    | <b>434.7</b> | <b>29,789.7</b> | <b>97.6</b>  | <b>51.8</b>  | <b>291.7</b> | <b>1,615.2</b> | <b>63.4</b>    | <b>31.6</b> | <b>2,151.3</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 37.4            |                 |              | 37.4            |  |              |              | 2.6            |                |             | 2.6            |    |
| Distillate Fuel Oil       | 840.5                                    | 438.3          | 1,050.9         | 4,594.9         | 109.3           | 76.5         | 7,110.4         | 62.2   | 32.4         | 77.7         | 339.8          | 8.1            | 5.7         | 525.9          |    |
| Jet Fuel                  |  |                |                 | 2,556.0         | NA              | 78.5         | 2,634.5         |  |              |              | 184.6          |                | 5.7         | 190.3          |    |
| Kerosene                  | 88.8                                     | 21.0           | 18.3            |                 |                 | 3.0          | 131.1           | 6.5  | 1.5          | 1.3          |                |                | 0.2         | 9.6            |    |
| LPG                       | 468.7                                    | 122.4          | 401.7           | 15.6            |                 | 7.3          | 1,015.8         | 28.9   | 7.5          | 24.8         | 1.0            |                | 0.5         | 62.6           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 14,987.5        |                 | 151.4        | 15,138.9        |  |              |              | 1,063.6        |                | 10.7        | 1,074.4        |    |
| Residual Fuel             |  | 137.2          | 284.7           | 314.9           | 628.4           | 118.0        | 1,483.1         |  | 10.3         | 21.4         | 23.6           | 47.2           | 8.9         | 111.4          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 7.0             |                 |                 |              | 7.0             |  |              | 0.5          |                |                |             | 0.5            |    |
| Crude Oil                 |  |                | 13.7            |                 |                 |              | 13.7            |  |              | 1.0          |                |                |             | 1.0            |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 177.5           |                 |                 |              | 177.5           |  |              | 12.4         |                |                |             | 12.4           |    |
| Petroleum Coke            |  | 0.1            | 638.6           |                 | 79.6            |              | 718.3           |  | 0.0          | 65.2         |                | 8.1            |             | 73.3           |    |
| Still Gas                 |  |                | 1,434.9         |                 |                 |              | 1,434.9         |  |              | 95.7         |                |                |             | 95.7           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (112.8)         |                 |                 |              | (112.8)         |  |              | (8.4)        |                |                |             | (8.4)          |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>48.9</b>     |              | <b>48.9</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total (All Fuels)</b>  | <b>6,769.0</b>                           | <b>4,067.0</b> | <b>14,389.7</b> | <b>23,243.1</b> | <b>23,157.6</b> | <b>445.0</b> | <b>72,071.4</b> | <b>383.1</b>   | <b>234.5</b> | <b>907.4</b> | <b>1,654.3</b> | <b>2,021.0</b> | <b>32.6</b> | <b>5,232.8</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-30: 1995 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | 17.5                                     | 116.8          | 1,526.9         | NE              | 17,466.3        | 10.2         | 19,137.7        | 1.7  | 11.2         | 144.4        | NE             | 1,660.7        | 0.9         | 1,819.0        |    |
| Residential Coal          | 17.5                                     |                |                 |                 |                 |              | 17.5            | 1.7  |              |              |                |                |             | 1.7            |    |
| Commercial Coal           |  | 116.8          |                 |                 |                 |              | 116.8           |  | 11.2         |              |                |                |             | 11.2           |    |
| Industrial Other Coal     |  |                | 1,526.9         |                 |                 |              | 1,526.9         |  |              | 144.4        |                |                |             | 144.4          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 17,466.3        |              | 17,466.3        |  |              |              |                | 1,660.7        |             | 1,660.7        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.2         | 10.2            |  |              |              |                |                | 0.9         | 0.9            |    |
| <b>Natural Gas</b>        | 4,954.2                                  | 3,096.0        | 8,722.5         | 724.0           | 4,302.0         |              | 21,798.6        | 262.7  | 164.2        | 462.5        | 38.4           | 228.1          |             | 1,155.9        |    |
| <b>Total Petroleum</b>    | 1,262.3                                  | 695.0          | 3,542.8         | 21,939.9        | 754.5           | 461.9        | 28,656.3        | 88.5   | 50.2         | 263.6        | 1,571.5        | 58.7           | 33.6        | 2,066.0        |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 39.6            |                 |              | 39.6            |  |              |              | 2.7            |                |             | 2.7            |    |
| Distillate Fuel Oil       | 793.2                                    | 419.8          | 969.7           | 4,379.4         | 108.0           | 89.5         | 6,759.5         | 58.7   | 31.0         | 71.7         | 323.9          | 8.0            | 6.6         | 499.9          |    |
| Jet Fuel                  |  |                |                 | 2,428.8         | NA              | 75.7         | 2,504.5         |  |              |              | 172.2          |                | 5.4         | 177.6          |    |
| Kerosene                  | 74.3                                     | 22.1           | 15.4            |                 |                 | 3.6          | 115.4           | 5.4  | 1.6          | 1.1          |                |                | 0.3         | 8.4            |    |
| LPG                       | 394.8                                    | 108.7          | 403.4           | 17.7            |                 | 5.6          | 930.2           | 24.4   | 6.7          | 24.9         | 1.1            |                | 0.3         | 57.4           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 2.7            | 30.1            | 14,687.1        |                 | 146.8        | 14,866.6        |  | 0.2          | 2.1          | 1,042.4        |                | 10.4        | 1,055.2        |    |
| Residual Fuel             |  | 141.5          | 286.2           | 387.3           | 566.0           | 140.7        | 1,521.6         |  | 10.6         | 21.5         | 29.1           | 42.5           | 10.6        | 114.3          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 5.3             |                 |                 |              | 5.3             |  |              | 0.4          |                |                |             | 0.4            |    |
| Crude Oil                 |  |                | 14.5            |                 |                 |              | 14.5            |  |              | 1.1          |                |                |             | 1.1            |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 169.0           |                 |                 |              | 169.0           |  |              | 11.8         |                |                |             | 11.8           |    |
| Petroleum Coke            |  | 0.1            | 600.7           |                 | 80.6            |              | 681.4           |  | 0.0          | 61.3         |                | 8.2            |             | 69.6           |    |
| Still Gas                 |  |                | 1,369.5         |                 |                 |              | 1,369.5         |  |              | 91.4         |                |                |             | 91.4           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (320.9)         |                 |                 |              | (320.9)         |  |              | (23.8)       |                |                |             | (23.8)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | 45.6            |              | 45.6            |  |              |              |                | 0.3            |             | 0.3            |    |
| <b>Total Coal</b>         | <b>6,233.9</b>                           | <b>3,907.8</b> | <b>13,792.3</b> | <b>22,663.8</b> | <b>22,568.4</b> | <b>472.1</b> | <b>69,638.3</b> | <b>352.8</b>   | <b>225.6</b> | <b>870.5</b> | <b>1,609.9</b> | <b>1,947.9</b> | <b>34.5</b> | <b>5,041.2</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-31: 1994 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>20.8</b>                              | <b>118.1</b>   | <b>1,594.9</b>  | <b>NE</b>       | <b>17,260.9</b> | <b>10.0</b>  | <b>19,004.7</b> | <b>2.0</b>   | <b>11.3</b>  | <b>150.7</b> | <b>NE</b>      | <b>1,638.8</b> | <b>0.9</b>  | <b>1,803.7</b> |    |
| Residential Coal          | 20.8                                     |                |                 |                 |                 |              | 20.8            | 2.0  |              |              |                |                |             | 2.0            |    |
| Commercial Coal           |  | 118.1          |                 |                 |                 |              | 118.1           |  | 11.3         |              |                |                |             | 11.3           |    |
| Industrial Other Coal     |  |                | 1,594.9         |                 |                 |              | 1,594.9         |  |              | 150.7        |                |                |             | 150.7          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 17,260.9        |              | 17,260.9        |  |              |              |                | 1,638.8        |             | 1,638.8        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 10.0         | 10.0            |  |              |              |                |                | 0.9         | 0.9            |    |
| <b>Natural Gas</b>        | <b>4,959.8</b>                           | <b>2,962.0</b> | <b>8,290.3</b>  | <b>708.5</b>    | <b>3,977.3</b>  |              | <b>20,897.9</b> | <b>262.9</b>   | <b>157.0</b> | <b>439.4</b> | <b>37.6</b>    | <b>210.8</b>   |             | <b>1,107.6</b> |    |
| <b>Total Petroleum</b>    | <b>1,306.7</b>                           | <b>746.7</b>   | <b>3,699.5</b>  | <b>21,494.2</b> | <b>1,058.7</b>  | <b>506.3</b> | <b>28,812.1</b> | <b>91.9</b>  | <b>54.1</b>  | <b>274.7</b> | <b>1,539.4</b> | <b>81.2</b>    | <b>36.9</b> | <b>2,078.3</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 38.1            |                 |              | 38.1            |  |              |              | 2.6            |                |             | 2.6            |    |
| Distillate Fuel Oil       | 858.1                                    | 447.9          | 977.4           | 4,183.3         | 120.0           | 118.8        | 6,705.4         | 63.5   | 33.1         | 72.3         | 309.4          | 8.9            | 8.8         | 495.9          |    |
| Jet Fuel                  |  |                |                 | 2,473.8         | NA              | 65.8         | 2,539.5         |  |              |              | 175.5          |                | 4.7         | 180.2          |    |
| Kerosene                  | 64.9                                     | 19.5           | 16.9            |                 |                 | 3.0          | 104.3           | 4.8  | 1.4          | 1.2          |                |                | 0.2         | 7.6            |    |
| LPG                       | 383.7                                    | 107.3          | 423.1           | 34.0            |                 | 7.3          | 955.3           | 23.7   | 6.6          | 26.1         | 2.1            |                | 0.4         | 59.0           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 14,407.0        |                 | 147.4        | 14,554.4        |  |              |              | 1,022.9        |                | 10.5        | 1,033.4        |    |
| Residual Fuel             |  | 171.9          | 368.4           | 358.1           | 869.0           | 164.1        | 1,931.5         |  | 12.9         | 27.7         | 26.9           | 65.3           | 12.3        | 145.0          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 6.1             |                 |                 |              | 6.1             |  |              | 0.4          |                |                |             | 0.4            |    |
| Crude Oil                 |  |                | 18.7            |                 |                 |              | 18.7            |  |              | 1.4          |                |                |             | 1.4            |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 169.4           |                 |                 |              | 169.4           |  |              | 11.9         |                |                |             | 11.9           |    |
| Petroleum Coke            |  | 0.1            | 594.9           |                 | 69.7            |              | 664.7           |  | 0.0          | 60.7         |                | 7.1            |             | 67.9           |    |
| Still Gas                 |  |                | 1,404.0         |                 |                 |              | 1,404.0         |  |              | 93.7         |                |                |             | 93.7           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (279.2)         |                 |                 |              | (279.2)         |  |              | (20.7)       |                |                |             | (20.7)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>53.0</b>     |              | <b>53.0</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total Coal</b>         | <b>6,287.4</b>                           | <b>3,826.8</b> | <b>13,584.7</b> | <b>22,202.8</b> | <b>22,349.9</b> | <b>516.3</b> | <b>68,767.8</b> | <b>356.8</b>   | <b>222.4</b> | <b>864.8</b> | <b>1,577.0</b> | <b>1,931.2</b> | <b>37.8</b> | <b>4,990.0</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-32: 1993 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>25.7</b>                              | <b>117.3</b>   | <b>1,585.0</b>  | <b>NE</b>       | <b>17,195.9</b> | <b>9.6</b>   | <b>18,933.5</b> | <b>2.5</b>   | <b>11.3</b>  | <b>149.8</b> | <b>NE</b>      | <b>1,632.5</b> | <b>0.9</b>  | <b>1,796.9</b> |    |
| Residential Coal          | 25.7                                     |                |                 |                 |                 |              | 25.7            | 2.5  |              |              |                |                |             | 2.5            |    |
| Commercial Coal           |  | 117.3          |                 |                 |                 |              | 117.3           |  | 11.3         |              |                |                |             | 11.3           |    |
| Industrial Other Coal     |  |                | 1,585.0         |                 |                 |              | 1,585.0         |  |              | 149.8        |                |                |             | 149.8          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 17,195.9        |              | 17,195.9        |  |              |              |                | 1,632.5        |             | 1,632.5        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 9.6          | 9.6             |  |              |              |                |                | 0.9         | 0.9            |    |
| <b>Natural Gas</b>        | <b>5,063.3</b>                           | <b>2,923.3</b> | <b>8,272.5</b>  | <b>644.7</b>    | <b>3,537.5</b>  |              | <b>20,441.3</b> | <b>268.4</b>   | <b>155.0</b> | <b>438.6</b> | <b>34.2</b>    | <b>187.5</b>   |             | <b>1,083.7</b> |    |
| <b>Total Petroleum</b>    | <b>1,348.5</b>                           | <b>743.4</b>   | <b>3,588.7</b>  | <b>20,866.3</b> | <b>1,123.8</b>  | <b>459.4</b> | <b>28,129.9</b> | <b>94.9</b>  | <b>53.8</b>  | <b>267.4</b> | <b>1,498.2</b> | <b>86.4</b>    | <b>33.6</b> | <b>2,034.3</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 38.4            |                 |              | 38.4            |  |              |              | 2.7            |                |             | 2.7            |    |
| Distillate Fuel Oil       | 883.3                                    | 447.2          | 989.9           | 3,889.4         | 86.5            | 104.9        | 6,401.1         | 65.3   | 33.1         | 73.2         | 287.6          | 6.4            | 7.8         | 473.4          |    |
| Jet Fuel                  |  |                |                 | 2,368.4         | NA              | 62.1         | 2,430.5         |  |              |              | 168.2          |                | 4.4         | 172.6          |    |
| Kerosene                  | 75.6                                     | 14.0           | 13.1            |                 |                 | 3.8          | 106.5           | 5.5  | 1.0          | 1.0          |                |                | 0.3         | 7.8            |    |
| LPG                       | 389.6                                    | 109.2          | 412.2           | 20.2            |                 | 4.9          | 936.2           | 24.0   | 6.7          | 25.4         | 1.2            |                | 0.3         | 57.8           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 14,182.3        |                 | 127.8        | 14,310.1        |  |              |              | 1,010.9        |                | 9.1         | 1,020.0        |    |
| Residual Fuel             |  | 172.7          | 382.9           | 367.5           | 958.6           | 155.9        | 2,037.7         |  | 13.0         | 28.8         | 27.6           | 72.0           | 11.7        | 153.0          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 0.2             |                 |                 |              | 0.2             |  |              | 0.0          |                |                |             | 0.0            |    |
| Crude Oil                 |  |                | 21.2            |                 |                 |              | 21.2            |  |              | 1.6          |                |                |             | 1.6            |    |
| MoGas Blend Components    |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 166.1           |                 |                 |              | 166.1           |  |              | 11.6         |                |                |             | 11.6           |    |
| Petroleum Coke            |  | 0.2            | 614.6           |                 | 78.6            |              | 693.4           |  | 0.0          | 62.8         |                | 8.0            |             | 70.8           |    |
| Still Gas                 |  |                | 1,384.6         |                 |                 |              | 1,384.6         |  |              | 92.4         |                |                |             | 92.4           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (396.0)         |                 |                 |              | (396.0)         |  |              | (29.3)       |                |                |             | (29.3)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>57.3</b>     |              | <b>57.3</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total Coal</b>         | <b>6,437.5</b>                           | <b>3,783.9</b> | <b>13,446.1</b> | <b>21,511.0</b> | <b>21,914.5</b> | <b>469.0</b> | <b>67,562.0</b> | <b>365.8</b>   | <b>220.1</b> | <b>855.7</b> | <b>1,532.4</b> | <b>1,906.9</b> | <b>34.4</b> | <b>4,915.3</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-33: 1992 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>25.6</b>                              | <b>116.6</b>   | <b>1,554.6</b>  | <b>NE</b>       | <b>16,465.6</b> | <b>8.8</b>   | <b>18,171.1</b> | <b>2.5</b>   | <b>11.3</b>  | <b>147.4</b> | <b>NE</b>      | <b>1,569.6</b> | <b>0.8</b>  | <b>1,731.6</b> |    |
| Residential Coal          | 25.6                                     |                |                 |                 |                 |              | 25.6            | 2.5  |              |              |                |                |             | 2.5            |    |
| Commercial Coal           |  | 116.6          |                 |                 |                 |              | 116.6           |  | 11.3         |              |                |                |             | 11.3           |    |
| Industrial Other Coal     |  |                | 1,554.6         |                 |                 |              | 1,554.6         |  |              | 147.4        |                |                |             | 147.4          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 16,465.6        |              | 16,465.6        |  |              |              |                | 1,569.6        |             | 1,569.6        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 8.8          | 8.8             |  |              |              |                |                | 0.8         | 0.8            |    |
| <b>Natural Gas</b>        | <b>4,804.6</b>                           | <b>2,871.2</b> | <b>8,125.3</b>  | <b>608.1</b>    | <b>3,511.5</b>  |              | <b>19,920.7</b> | <b>254.5</b>   | <b>152.1</b> | <b>430.5</b> | <b>32.2</b>    | <b>186.0</b>   |             | <b>1,055.4</b> |    |
| <b>Total Petroleum</b>    | <b>1,365.8</b>                           | <b>788.9</b>   | <b>3,759.2</b>  | <b>20,361.3</b> | <b>990.7</b>    | <b>444.9</b> | <b>27,710.8</b> | <b>96.5</b>  | <b>57.3</b>  | <b>279.5</b> | <b>1,464.6</b> | <b>75.5</b>    | <b>32.5</b> | <b>2,005.9</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 41.1            |                 |              | 41.1            |  |              |              | 2.8            |                |             | 2.8            |    |
| Distillate Fuel Oil       | 931.4                                    | 481.7          | 1,028.5         | 3,665.7         | 73.5            | 91.8         | 6,272.6         | 68.9   | 35.6         | 76.1         | 271.1          | 5.4            | 6.8         | 463.9          |    |
| Jet Fuel                  |  |                |                 | 2,343.8         | NA              | 61.3         | 2,405.1         |  |              |              | 166.6          |                | 4.4         | 171.0          |    |
| Kerosene                  | 65.0                                     | 11.1           | 9.8             |                 |                 | 3.3          | 89.2            | 4.8  | 0.8          | 0.7          |                |                | 0.2         | 6.5            |    |
| LPG                       | 369.4                                    | 106.9          | 441.8           | 19.4            |                 | 11.9         | 949.4           | 22.8   | 6.6          | 27.3         | 1.2            |                | 0.7         | 58.7           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 13,891.2        |                 | 122.1        | 14,013.3        |  |              |              | 992.8          |                | 8.7         | 1,001.5        |    |
| Residual Fuel             |  | 189.1          | 323.9           | 400.1           | 872.2           | 154.6        | 1,939.8         |  | 14.2         | 24.3         | 30.0           | 65.5           | 11.6        | 145.7          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 0.2             |                 |                 |              | 0.2             |  |              | 0.0          |                |                |             | 0.0            |    |
| Crude Oil                 |  |                | 27.4            |                 |                 |              | 27.4            |  |              | 2.0          |                |                |             | 2.0            |    |
| MoGas Blend Components    |  |                | 75.7            |                 |                 |              | 75.7            |  |              | 5.4          |                |                |             | 5.4            |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 161.3           |                 |                 |              | 161.3           |  |              | 11.3         |                |                |             | 11.3           |    |
| Petroleum Coke            |  | 0.1            | 627.2           |                 | 45.0            |              | 672.2           |  | 0.0          | 64.0         |                | 4.6            |             | 68.6           |    |
| Still Gas                 |  |                | 1,418.4         |                 |                 |              | 1,418.4         |  |              | 94.6         |                |                |             | 94.6           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (354.8)         |                 |                 |              | (354.8)         |  |              | (26.3)       |                |                |             | (26.3)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>55.1</b>     |              | <b>55.1</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total Coal</b>         | <b>6,196.0</b>                           | <b>3,776.7</b> | <b>13,439.1</b> | <b>20,969.4</b> | <b>21,022.9</b> | <b>453.7</b> | <b>65,857.7</b> | <b>353.5</b>   | <b>220.6</b> | <b>857.4</b> | <b>1,496.9</b> | <b>1,831.5</b> | <b>33.3</b> | <b>4,793.2</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-34: 1991 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>25.4</b>                              | <b>115.5</b>   | <b>1,602.7</b>  | <b>NE</b>       | <b>16,249.7</b> | <b>7.7</b>   | <b>18,001.0</b> | <b>2.4</b>   | <b>11.1</b>  | <b>152.1</b> | <b>NE</b>      | <b>1,548.2</b> | <b>0.7</b>  | <b>1,714.6</b> |    |
| Residential Coal          | 25.4                                     |                |                 |                 |                 |              | 25.4            | 2.4  |              |              |                |                |             | 2.4            |    |
| Commercial Coal           |  | 115.5          |                 |                 |                 |              | 115.5           |  | 11.1         |              |                |                |             | 11.1           |    |
| Industrial Other Coal     |  |                | 1,602.7         |                 |                 |              | 1,602.7         |  |              | 152.1        |                |                |             | 152.1          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 16,249.7        |              | 16,249.7        |  |              |              |                | 1,548.2        |             | 1,548.2        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 7.7          | 7.7             |  |              |              |                |                | 0.7         | 0.7            |    |
| <b>Natural Gas</b>        | <b>4,667.2</b>                           | <b>2,795.4</b> | <b>7,827.8</b>  | <b>620.3</b>    | <b>3,377.4</b>  |              | <b>19,288.1</b> | <b>247.3</b>   | <b>148.1</b> | <b>414.7</b> | <b>32.9</b>    | <b>178.9</b>   |             | <b>1,021.8</b> |    |
| <b>Total Petroleum</b>    | <b>1,381.5</b>                           | <b>881.8</b>   | <b>3,430.9</b>  | <b>19,715.8</b> | <b>1,198.3</b>  | <b>425.4</b> | <b>27,033.6</b> | <b>97.5</b>  | <b>64.0</b>  | <b>255.7</b> | <b>1,414.7</b> | <b>90.7</b>    | <b>30.9</b> | <b>1,953.5</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 41.7            |                 |              | 41.7            |  |              |              | 2.9            |                |             | 2.9            |    |
| Distillate Fuel Oil       | 931.0                                    | 517.7          | 1,050.8         | 3,449.7         | 83.6            | 71.4         | 6,104.1         | 68.9   | 38.3         | 77.7         | 255.1          | 6.2            | 5.3         | 451.4          |    |
| Jet Fuel                  |  |                |                 | 2,373.6         | NA              | 78.2         | 2,451.8         |  |              |              | 168.8          |                | 5.6         | 174.4          |    |
| Kerosene                  | 72.3                                     | 12.1           | 11.4            |                 |                 | 2.8          | 98.6            | 5.3  | 0.9          | 0.8          |                |                | 0.2         | 7.2            |    |
| LPG                       | 378.1                                    | 108.2          | 342.2           | 21.1            |                 | 13.8         | 863.5           | 23.3   | 6.7          | 21.1         | 1.3            |                | 0.9         | 53.3           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  | 31.8           | 72.3            | 13,605.3        |                 | 124.7        | 13,834.1        |  | 2.3          | 5.2          | 969.7          |                | 8.9         | 986.0          |    |
| Residual Fuel             |  | 211.9          | 270.9           | 224.4           | 1,085.3         | 134.6        | 1,927.2         |  | 15.9         | 20.3         | 16.9           | 81.5           | 10.1        | 144.7          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | (0.1)           |                 |                 |              | (0.1)           |  |              | (0.0)        |                |                |             | (0.0)          |    |
| Crude Oil                 |  |                | 39.0            |                 |                 |              | 39.0            |  |              | 2.9          |                |                |             | 2.9            |    |
| MoGas Blend Components    |  |                | (25.9)          |                 |                 |              | (25.9)          |  |              | (1.8)        |                |                |             | (1.8)          |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 147.0           |                 |                 |              | 147.0           |  |              | 10.3         |                |                |             | 10.3           |    |
| Petroleum Coke            |  |                | 587.6           |                 | 29.3            |              | 616.9           |  |              | 60.0         |                | 3.0            |             | 63.0           |    |
| Still Gas                 |  |                | 1,385.9         |                 |                 |              | 1,385.9         |  |              | 92.5         |                |                |             | 92.5           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (450.2)         |                 |                 |              | (450.2)         |  |              | (33.3)       |                |                |             | (33.3)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>54.5</b>     |              | <b>54.5</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total Coal</b>         | <b>6,074.0</b>                           | <b>3,792.7</b> | <b>12,861.4</b> | <b>20,336.1</b> | <b>20,879.8</b> | <b>433.2</b> | <b>64,377.2</b> | <b>347.2</b>   | <b>223.3</b> | <b>822.5</b> | <b>1,447.6</b> | <b>1,818.2</b> | <b>31.6</b> | <b>4,690.3</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-35: 1990 Energy Consumption Data and CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Fuel Type**

|                           | 1  | 2              | 3               | 4               | 5               | 6            | 7               | 8  | 9            | 10           | 11             | 12             | 13          | 14             | 15 |
|---------------------------|--|----------------|-----------------|-----------------|-----------------|--------------|-----------------|--|--------------|--------------|----------------|----------------|-------------|----------------|----|
| Fuel Type                 | Adjusted Consumption (Tbtu) <sup>a</sup> |                |                 |                 |                 |              |                 | Emissions <sup>b</sup> (MMT CO <sub>2</sub> Eq.) from Energy Use |              |              |                |                |             |                |    |
|                           | Res.                                     | Comm.          | Ind.            | Trans.          | Elec.           | Terr.        | Total           | Res.   | Comm.        | Ind.         | Trans.         | Elec.          | Terr.       | Total          |    |
| <b>Total Coal</b>         | <b>31.1</b>                              | <b>124.5</b>   | <b>1,640.5</b>  | <b>NE</b>       | <b>16,261.0</b> | <b>7.0</b>   | <b>18,064.0</b> | <b>3.0</b>   | <b>12.0</b>  | <b>155.3</b> | <b>NE</b>      | <b>1,547.6</b> | <b>0.6</b>  | <b>1,718.4</b> |    |
| Residential Coal          | 31.1                                     |                |                 |                 |                 |              | 31.1            | 3.0  |              |              |                |                |             | 3.0            |    |
| Commercial Coal           |  | 124.5          |                 |                 |                 |              | 124.5           |  | 12.0         |              |                |                |             | 12.0           |    |
| Industrial Other Coal     |  |                | 1,640.5         |                 |                 |              | 1,640.5         |  |              | 155.3        |                |                |             | 155.3          |    |
| Transportation Coal       |  |                |                 | NE              |                 |              | NE              |  |              |              | NE             |                |             | NE             |    |
| Electric Power Coal       |  |                |                 |                 | 16,261.0        |              | 16,261.0        |  |              |              |                | 1,547.6        |             | 1,547.6        |    |
| U.S. Territory Coal (bit) |  |                |                 |                 |                 | 7.0          | 7.0             |  |              |              |                |                | 0.6         | 0.6            |    |
| <b>Natural Gas</b>        | <b>4,490.9</b>                           | <b>2,682.2</b> | <b>7,716.4</b>  | <b>679.9</b>    | <b>3,308.5</b>  |              | <b>18,877.9</b> | <b>238.0</b>   | <b>142.1</b> | <b>408.9</b> | <b>36.0</b>    | <b>175.3</b>   |             | <b>1,000.3</b> |    |
| <b>Total Petroleum</b>    | <b>1,375.2</b>                           | <b>869.4</b>   | <b>3,743.8</b>  | <b>20,323.1</b> | <b>1,289.4</b>  | <b>374.8</b> | <b>27,975.7</b> | <b>97.4</b>  | <b>63.3</b>  | <b>278.3</b> | <b>1,457.7</b> | <b>97.5</b>    | <b>27.2</b> | <b>2,021.5</b> |    |
| Asphalt & Road Oil        |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Aviation Gasoline         |  |                |                 | 45.0            |                 |              | 45.0            |  |              |              | 3.1            |                |             | 3.1            |    |
| Distillate Fuel Oil       | 959.2                                    | 525.4          | 1,098.5         | 3,554.8         | 96.5            | 74.0         | 6,308.4         | 70.9   | 38.9         | 81.2         | 262.9          | 7.1            | 5.5         | 466.5          |    |
| Jet Fuel                  |  |                |                 | 2,590.1         | NA              | 61.0         | 2,651.1         |  |              |              | 184.2          |                | 4.3         | 188.6          |    |
| Kerosene                  | 63.9                                     | 11.8           | 12.3            |                 |                 | 2.6          | 90.6            | 4.7  | 0.9          | 0.9          |                |                | 0.2         | 6.6            |    |
| LPG                       | 352.1                                    | 102.3          | 380.2           | 22.9            |                 | 14.4         | 871.9           | 21.8   | 6.3          | 23.5         | 1.4            |                | 0.9         | 53.9           |    |
| Lubricants                |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Motor Gasoline            |  |                |                 | 13,810.1        |                 | 101.0        | 13,911.1        |  |              |              | 983.5          |                | 7.2         | 990.7          |    |
| Residual Fuel             |  | 229.8          | 364.1           | 300.3           | 1,162.6         | 121.8        | 2,178.7         |  | 17.3         | 27.3         | 22.6           | 87.3           | 9.2         | 163.6          |    |
| Other Petroleum           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| AvGas Blend Components    |  |                | 0.2             |                 |                 |              | 0.2             |  |              | 0.0          |                |                |             | 0.0            |    |
| Crude Oil                 |  |                | 50.9            |                 |                 |              | 50.9            |  |              | 3.8          |                |                |             | 3.8            |    |
| MoGas Blend Components    |  |                | 53.7            |                 |                 |              | 53.7            |  |              | 3.8          |                |                |             | 3.8            |    |
| Misc. Products            |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Naphtha (<401 deg. F)     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Other Oil (>401 deg. F)   |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Pentanes Plus             |  |                | 125.2           |                 |                 |              | 125.2           |  |              | 8.8          |                |                |             | 8.8            |    |
| Petroleum Coke            |  |                | 591.2           |                 | 30.4            |              | 621.5           |  |              | 60.4         |                | 3.1            |             | 63.5           |    |
| Still Gas                 |  |                | 1,436.5         |                 |                 |              | 1,436.5         |  |              | 95.8         |                |                |             | 95.8           |    |
| Special Naphtha           |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| Unfinished Oils           |  |                | (369.0)         |                 |                 |              | (369.0)         |  |              | (27.3)       |                |                |             | (27.3)         |    |
| Waxes                     |  |                |                 |                 |                 |              |                 |  |              |              |                |                |             |                |    |
| <b>Geothermal</b>         |  |                |                 |                 | <b>52.7</b>     |              | <b>52.7</b>     |  |              |              |                | <b>0.4</b>     |             | <b>0.4</b>     |    |
| <b>Total Coal</b>         | <b>5,897.2</b>                           | <b>3,676.0</b> | <b>13,100.6</b> | <b>21,003.0</b> | <b>20,911.6</b> | <b>381.9</b> | <b>64,970.3</b> | <b>338.3</b>   | <b>217.4</b> | <b>842.5</b> | <b>1,493.8</b> | <b>1,820.8</b> | <b>27.9</b> | <b>4,740.7</b> |    |

<sup>a</sup> Expressed as gross calorific values (i.e., higher heating values). Adjustments include biofuels, conversion of fossil fuels, non-energy use (see Table A-36), and international bunker fuel consumption (see Table A-37).

<sup>b</sup> Consumption and/or emissions of select fuels are shown as negative due to differences in EIA energy balancing accounting. These are designated with parentheses.

NE (Not Estimated); NA (Not Available)

**Table A-36: Unadjusted Non-Energy Fuel Consumption (Tbtu)**

| Sector/Fuel Type                              | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Industry</b>                               | <b>4,544.0</b> | <b>5,089.7</b> | <b>5,576.5</b> | <b>5,263.7</b> | <b>5,425.7</b> | <b>5,342.9</b> | <b>5,847.2</b> | <b>5,483.3</b> | <b>5,470.0</b> | <b>5,225.2</b> | <b>4,769.7</b> | <b>4,510.2</b> | <b>4,753.0</b> | <b>4,665.0</b> | <b>4,611.3</b> | <b>4,823.4</b> | <b>4,699.9</b> |
| Industrial Coking Coal                        | 0.0            | 37.8           | 53.5           | 24.8           | 40.3           | 51.9           | 167.8          | 80.4           | 62.9           | 2.3            | 29.2           | 6.4            | 64.8           | 60.8           | 132.5          | 119.3          | 23.0           |
| Industrial Other Coal                         | 8.2            | 11.3           | 12.4           | 11.3           | 12.0           | 11.9           | 11.9           | 11.9           | 11.9           | 11.9           | 11.9           | 11.9           | 10.3           | 10.3           | 10.3           | 10.3           | 10.3           |
| Natural Gas to Chemical<br>Plants, Other Uses | 305.9          | 371.0          | 401.7          | 391.8          | 380.7          | 345.3          | 306.6          | 270.4          | 233.4          | 233.6          | 233.6          | 233.6          | 311.8          | 311.8          | 311.8          | 311.8          | 311.8          |
| Asphalt & Road Oil                            | 1,170.2        | 1,178.2        | 1,275.7        | 1,256.9        | 1,240.0        | 1,219.5        | 1,303.8        | 1,323.2        | 1,261.2        | 1,197.0        | 1,012.0        | 873.1          | 877.8          | 859.5          | 826.7          | 783.3          | 792.6          |
| LPG   | 1,201.4        | 1,586.9        | 1,759.3        | 1,642.3        | 1,766.3        | 1,701.6        | 1,768.5        | 1,659.5        | 1,734.6        | 1,726.7        | 1,596.6        | 1,748.0        | 1,901.6        | 1,943.4        | 1,990.5        | 2,149.0        | 2,148.7        |
| Lubricants                                    | 186.3          | 177.8          | 189.9          | 174.0          | 171.9          | 159.0          | 161.0          | 160.2          | 156.1          | 161.2          | 149.6          | 134.5          | 149.5          | 141.8          | 130.5          | 138.1          | 144.0          |
| Pentanes Plus                                 | 125.2          | 169.0          | 171.6          | 131.6          | 111.9          | 110.4          | 111.2          | 98.1           | 70.1           | 89.7           | 76.5           | 63.8           | 77.7           | 27.3           | 42.2           | 47.1           | 44.2           |
| Naphtha (<401 deg. F)                         | 347.8          | 373.0          | 613.5          | 493.7          | 582.6          | 613.0          | 749.4          | 698.7          | 628.9          | 562.5          | 477.2          | 471.9          | 490.6          | 487.3          | 453.9          | 517.8          | 442.6          |
| Other Oil (>401 deg. F)                       | 753.9          | 801.0          | 722.2          | 662.5          | 632.1          | 699.4          | 779.5          | 708.0          | 790.6          | 744.1          | 647.8          | 424.8          | 452.5          | 388.5          | 287.2          | 223.9          | 247.2          |
| Still Gas                                     | 36.7           | 47.9           | 17.0           | 49.3           | 61.7           | 59.0           | 62.9           | 67.7           | 57.2           | 44.2           | 47.3           | 133.9          | 147.8          | 163.6          | 160.6          | 166.7          | 164.6          |
| Petroleum Coke                                | 123.1          | 120.6          | 98.5           | 174.3          | 145.8          | 122.8          | 217.7          | 186.9          | 213.6          | 201.2          | 224.5          | 180.7          | 61.0           | 62.4           | 67.6           | 62.4           | 61.4           |
| Special Naphtha                               | 107.1          | 70.8           | 97.4           | 78.5           | 102.4          | 80.5           | 51.0           | 62.5           | 70.1           | 78.0           | 84.9           | 46.2           | 26.1           | 22.6           | 14.7           | 100.0          | 106.1          |
| Other (Wax/Misc.)                             |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |
| Distillate Fuel Oil                           | 7.0            | 6.8            | 11.7           | 11.7           | 11.7           | 11.7           | 11.7           | 11.7           | 17.5           | 17.5           | 17.5           | 17.5           | 5.8            | 5.8            | 5.8            | 5.8            | 5.8            |
| Waxes   | 33.3           | 40.6           | 33.1           | 36.3           | 32.2           | 31.1           | 30.8           | 31.4           | 26.2           | 21.9           | 19.1           | 12.2           | 17.1           | 15.1           | 15.3           | 16.5           | 14.8           |
| Miscellaneous Products                        | 137.8          | 97.1           | 119.2          | 124.9          | 134.2          | 126.0          | 113.4          | 112.8          | 136.0          | 133.5          | 142.0          | 151.8          | 158.7          | 164.7          | 161.6          | 171.2          | 182.7          |
| <b>Transportation</b>                         | <b>176.0</b>   | <b>167.9</b>   | <b>179.4</b>   | <b>164.3</b>   | <b>162.4</b>   | <b>150.1</b>   | <b>152.1</b>   | <b>151.3</b>   | <b>147.4</b>   | <b>152.2</b>   | <b>141.3</b>   | <b>127.1</b>   | <b>141.2</b>   | <b>133.9</b>   | <b>123.2</b>   | <b>130.4</b>   | <b>136.0</b>   |
| Lubricants                                    | 176.0          | 167.9          | 179.4          | 164.3          | 162.4          | 150.1          | 152.1          | 151.3          | 147.4          | 152.2          | 141.3          | 127.1          | 141.2          | 133.9          | 123.2          | 130.4          | 136.0          |
| <b>U.S. Territories</b>                       | <b>86.7</b>    | <b>90.8</b>    | <b>152.2</b>   | <b>80.3</b>    | <b>140.2</b>   | <b>123.5</b>   | <b>110.8</b>   | <b>121.9</b>   | <b>133.4</b>   | <b>108.4</b>   | <b>132.1</b>   | <b>59.6</b>    | <b>56.4</b>    | <b>56.7</b>    | <b>58.1</b>    | <b>57.4</b>    | <b>53.6</b>    |
| Lubricants                                    | 0.7            | 2.0            | 3.1            | 0.0            | 3.0            | 4.9            | 5.1            | 4.6            | 6.2            | 5.9            | 2.7            | 1.0            | 1.0            | 1.0            | 1.0            | 1.0            | 1.0            |
| Other Petroleum (Misc.<br>Prod.)              | 86.0           | 88.8           | 149.1          | 80.3           | 137.2          | 118.6          | 105.7          | 117.3          | 127.2          | 102.5          | 129.4          | 58.5           | 55.4           | 55.7           | 57.1           | 56.4           | 52.6           |
| <b>Total</b>                                  | <b>4,806.7</b> | <b>5,348.5</b> | <b>5,908.0</b> | <b>5,508.4</b> | <b>5,728.3</b> | <b>5,616.5</b> | <b>6,110.1</b> | <b>5,756.6</b> | <b>5,750.8</b> | <b>5,485.9</b> | <b>5,043.0</b> | <b>4,696.8</b> | <b>4,950.6</b> | <b>4,855.7</b> | <b>4,792.7</b> | <b>5,011.2</b> | <b>4,889.5</b> |

Note: These values are unadjusted non-energy fuel use provided by EIA. They have not yet been adjusted to remove petroleum feedstock exports and processes accounted for in the Industrial Processes and Product Use Chapter.

**Table A-37: International Bunker Fuel Consumption (Tbtu)**

| Fuel Type                             | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|---------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Marine Residual Fuel Oil              | 715.7          | 523.2          | 444.1          | 426.0          | 448.9          | 471.8          | 553.1          | 581.0          | 599.4          | 607.5          | 654.6          | 604.8          | 619.8          | 518.4          | 459.5          | 379.8          | 369.2          |
| Marine Distillate Fuel Oil &<br>Other | 158.0          | 125.7          | 85.9           | 72.4           | 82.6           | 103.9          | 143.6          | 126.9          | 119.3          | 111.3          | 122.2          | 111.0          | 128.2          | 107.4          | 91.7           | 75.4           | 82.0           |
| Aviation Jet Fuel                     | 539.4          | 703.4          | 880.1          | 799.7          | 774.8          | 783.0          | 797.7          | 853.1          | 855.6          | 872.7          | 796.8          | 749.1          | 865.4          | 919.9          | 916.3          | 931.6          | 985.1          |
| <b>Total</b>                          | <b>1,413.1</b> | <b>1,352.3</b> | <b>1,410.0</b> | <b>1,298.1</b> | <b>1,306.3</b> | <b>1,358.7</b> | <b>1,494.4</b> | <b>1,561.0</b> | <b>1,574.2</b> | <b>1,591.5</b> | <b>1,573.6</b> | <b>1,464.9</b> | <b>1,613.4</b> | <b>1,545.7</b> | <b>1,467.4</b> | <b>1,386.9</b> | <b>1,436.3</b> |

Note: Further information on the calculation of international bunker fuel consumption of aviation jet fuel is provided in Annex 0: Methodology for Estimating Emissions from Commercial Aircraft Jet Fuel Consumption.

**Table A-38: Key Assumptions for Estimating CO<sub>2</sub> Emissions**

| <b>Fuel Type</b>                       | <b>C Content Coefficient<br/>(MMT C/QBtu)</b> |
|--|---|
| <b>Coal</b>                            |   |
| Residential Coal                       | [c]   |
| Commercial Coal                        | [c]   |
| Industrial Coking Coal                 | [c]   |
| Industrial Other Coal                  | [c]   |
| Electric Power Coal                    | [c]   |
| U.S. Territory Coal (bit)              | 25.14   |
| <b>Pipeline Natural Gas</b>            |   |
| Flare Gas <sup>a</sup>                 | 14.92   |
| <b>Petroleum</b>                       |   |
| Asphalt & Road Oil                     | 20.55   |
| Aviation Gasoline                      | 18.86   |
| Distillate Fuel Oil No. 1              | 19.98   |
| Distillate Fuel Oil No. 2 <sup>b</sup> | 20.17   |
| Distillate Fuel Oil No. 4              | 20.47   |
| Jet Fuel                               | [c]   |
| Kerosene                               | 19.96   |
| LPG (energy use)                       | [c]   |
| LPG (non-energy use)                   | [c]   |
| Lubricants                             | 20.20   |
| Motor Gasoline                         | [c]   |
| Residual Fuel Oil No. 5                | 19.89   |
| Residual Fuel Oil No. 6 <sup>b</sup>   | 20.48   |
| <b>Other Petroleum</b>                 |   |
| AvGas Blend Components                 | 18.87   |
| Crude Oil                              | [c]   |
| MoGas Blend Components                 | [c]   |
| Misc. Products                         | [c]   |
| Misc. Products (Territories)           | 20.00   |
| Naphtha (<401 deg. F)                  | 18.55   |
| Other Oil (>401 deg. F)                | 20.17   |
| Pentanes Plus                          | 19.10   |
| Petroleum Coke                         | 27.85   |
| Still Gas                              | 18.20   |
| Special Naphtha                        | 19.74   |
| Unfinished Oils                        | [c]   |
| Waxes                                  | 19.80   |
| <b>Geothermal</b>                      | <b>2.05</b>                                   |

<sup>a</sup> Flare gas is not used in the CO<sub>2</sub> from fossil fuel combustion calculations and is presented for informational purposes only.

<sup>b</sup> Distillate fuel oil No.2 and residual fuel oil No. 6 are used in the CO<sub>2</sub> from fossil fuel combustion calculations, and other oil types are presented for informational purposes only. An additional discussion on the derivation of these carbon content coefficients is presented in Annex 2.2.

Sources: C coefficients from EIA (2009b) and EPA (2010a).

[c] These coefficients vary annually due to fluctuations in fuel quality (see Table A-39).

**Table A-39: Annually Variable C Content Coefficients by Year (MMT C/QBtu)**

| Fuel Type              | 1990  | 1995  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  | 2008               | 2009               | 2010               | 2011               | 2012               | 2013               | 2014               |
|------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| Residential Coal       | 26.20 | 26.13 | 26.01 | 26.00 | 25.98 | 26.04 | 25.91 | 26.09 | 26.29 | 25.94 | 25.71 <sup>a</sup> |
| Commercial Coal        | 26.00 | 26.13 | 26.01 | 26.00 | 25.98 | 26.04 | 25.91 | 26.09 | 26.29 | 25.94 | 25.71              | 25.71              | 25.71              | 25.71              | 25.71              | 25.71              | 25.71              |
| Industrial Coking Coal | 31.00 | 31.00 | 31.00 | 31.00 | 31.00 | 31.00 | 31.00 | 31.00 | 31.00 | 31.00 | 31.00              | 31.00              | 31.00              | 31.00              | 31.00              | 31.00              | 31.00              |
| Industrial Other Coal  | 25.82 | 25.80 | 25.74 | 25.66 | 25.57 | 25.55 | 25.56 | 25.80 | 25.84 | 25.82 | 25.82              | 25.82              | 25.82              | 25.82              | 25.82              | 25.82              | 25.82              |
| Electric Power Coal    | 25.96 | 25.93 | 26.00 | 26.00 | 26.05 | 26.09 | 26.10 | 26.09 | 26.04 | 26.05 | 26.05              | 26.05              | 26.05              | 26.05              | 26.05              | 26.05              | 26.05              |
| Pipeline Natural Gas   | 14.45 | 14.46 | 14.47 | 14.46 | 14.46 | 14.44 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46              | 14.46              | 14.46              | 14.46              | 14.46              | 14.46              | 14.46              |
| LPG (energy use)       | 16.86 | 16.82 | 16.89 | 16.87 | 16.85 | 16.86 | 16.84 | 16.84 | 16.83 | 16.82 | 16.83              | 16.83              | 16.83              | 16.83              | 16.83              | 16.83              | 16.83              |
| LPG (non-energy use)   | 17.06 | 17.09 | 17.09 | 17.10 | 17.09 | 17.09 | 17.07 | 17.06 | 17.06 | 17.05 | 17.06              | 17.06              | 17.06              | 17.06              | 17.06              | 17.06              | 17.06              |
| Motor Gasoline         | 19.42 | 19.36 | 19.33 | 19.34 | 19.38 | 19.36 | 19.38 | 19.36 | 19.45 | 19.56 | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              |
| Jet Fuel               | 19.40 | 19.34 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70              | 19.70              | 19.70              | 19.70              | 19.70              | 19.70              | 19.70              |
| MoGas Blend            |       |       |       |       |       |       |       |       |       |       |                    |                    |                    |                    |                    |                    |                    |
| Components             | 19.42 | 19.36 | 19.33 | 19.34 | 19.38 | 19.36 | 19.38 | 19.36 | 19.45 | 19.56 | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              |
| Misc. Products         | 20.15 | 20.21 | 20.22 | 20.27 | 20.28 | 20.25 | 20.31 | 20.31 | 20.28 | 20.28 | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              |
| Unfinished Oils        | 20.15 | 20.21 | 20.22 | 20.27 | 20.28 | 20.25 | 20.31 | 20.31 | 20.28 | 20.28 | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              |
| Crude Oil              | 20.15 | 20.21 | 20.22 | 20.27 | 20.28 | 20.25 | 20.31 | 20.31 | 20.28 | 20.28 | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              |

<sup>a</sup> EIA discontinued collection of residential sector coal consumption data in 2008, because consumption of coal in the residential sector is extremely limited. Therefore, the number cited here is developed from commercial/institutional consumption.

Source: EPA (2010a).

**Table A-40: Electricity Consumption by End-Use Sector (Billion Kilowatt-Hours)**

| End-Use Sector | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Residential    | 924          | 1,043        | 1,192        | 1,202        | 1,265        | 1,276        | 1,292        | 1,359        | 1,352        | 1,392        | 1,381        | 1,365        | 1,446        | 1,423        | 1,375        | 1,395        | 1,407        |
| Commercial     | 838          | 953          | 1,159        | 1,191        | 1,205        | 1,199        | 1,230        | 1,275        | 1,300        | 1,336        | 1,336        | 1,307        | 1,330        | 1,328        | 1,327        | 1,337        | 1,352        |
| Industrial     | 1,070        | 1,163        | 1,235        | 1,159        | 1,156        | 1,181        | 1,186        | 1,169        | 1,158        | 1,154        | 1,142        | 1,044        | 1,103        | 1,124        | 1,123        | 1,129        | 1,136        |
| Transportation | 5            | 5            | 5            | 6            | 6            | 7            | 7            | 8            | 7            | 8            | 8            | 8            | 8            | 8            | 7            | 8            | 8            |
| <b>Total</b>   | <b>2,837</b> | <b>3,164</b> | <b>3,592</b> | <b>3,557</b> | <b>3,632</b> | <b>3,662</b> | <b>3,716</b> | <b>3,811</b> | <b>3,817</b> | <b>3,890</b> | <b>3,866</b> | <b>3,724</b> | <b>3,887</b> | <b>3,883</b> | <b>3,832</b> | <b>3,868</b> | <b>3,903</b> |

Note: Does not include the U.S. Territories.

Source: EIA (2016).

## References

- EIA (2016) *Monthly Energy Review*. February 2016, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2016/02)
- EIA (2013) *EIA Manufacturing Consumption of Energy (MECS) 2010*. U.S. Department of Energy, Energy Information Administration, Washington, D.C.
- EIA (2009b) *Manufacturing Consumption of Energy 2006*. Energy Information Administration, U.S. Department of Energy. Washington, DC. Released July, 2009.
- EPA (2010a) Carbon Content Coefficients Developed for EPA's Mandatory Reporting Rule. Office of Air and Radiation, Office of Atmospheric Programs, U.S. Environmental Protection Agency, Washington, D.C.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- UNFCCC (2014) *Decision 24/CP.19, Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention*. United Nations Framework Convention on Climate Change (UNFCCC) Conference of the Parties Nineteenth session, Warsaw, Poland. November 23, 2013. Available online at: <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>.

## 2.2. Methodology for Estimating the Carbon Content of Fossil Fuels

This sub-annex presents the background and methodology for estimating the carbon (C) content of fossil fuels combusted in the United States. The C content of a particular fossil fuel represents the maximum potential emissions to the atmosphere if all C in the fuel is oxidized during combustion. The C content coefficients used in past editions of this report were developed using methods first outlined in the U.S. Energy Information Administration's (EIA) *Emissions of Greenhouse Gases in the United States: 1987-1992* (1994) and were developed primarily by EIA. For this report, EPA has updated many of the C content coefficients based on carbon dioxide (CO<sub>2</sub>) emission factors developed for the Mandatory Reporting of Greenhouse Gases Rule, signed in September 2009 (EPA 2009b, 2010). This sub-annex describes an updated methodology for estimating the C content of natural gas, and presents a time-series analysis of changes in U.S. C content coefficients for coal, petroleum products, and natural gas. A summary of C content coefficients used in this report appears in Table A-41.

Though the methods for estimating C contents for coal, natural gas, and petroleum products differ in their details, they each follow the same basic approach. First, because C coefficients are presented in terms of mass per unit energy (i.e., million metric tons C per quadrillion Btu or MMT C/QBtu), those fuels that are typically described in volumetric units (i.e., petroleum products and natural gas) are converted to units of mass using an estimated density. Second, C contents are derived from fuel sample data, using descriptive statistics to estimate the C share of the fuel by weight. The heat content of the fuel is then estimated based on the sample data, or where sample data are unavailable or unrepresentative, by default values that reflect the characteristics of the fuel as defined by market requirements. A discussion of each fuel appears below.

The C content of coal is described first because approximately one-third of all U.S. C emissions from fossil fuel combustion are associated with coal consumption. The methods and sources for estimating the C content of natural gas are provided next. Approximately one-quarter of U.S. greenhouse gas emissions from fossil fuel combustion are attributable to natural gas consumption. Finally, this sub-annex examines C contents of petroleum products. U.S. energy consumption statistics account for more than 20 different petroleum products.

**Table A-41: Carbon Content Coefficients Used in this Report (MMT Carbon/QBtu)**

| Fuel Type                               | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008               | 2009               | 2010               | 2011               | 2012               | 2013               | 2014               |
|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
| <b>Coal</b>                             |              |              |              |              |              |              |              |              |              |              |                    |                    |                    |                    |                    |                    |                    |
| Residential Coal <sup>a</sup>           | 26.20        | 26.13        | 26.01        | 26.00        | 25.98        | 26.04        | 25.91        | 26.09        | 26.29        | 25.94        | 25.71 <sup>b</sup> |
| Commercial Coal <sup>a</sup>            | 26.20        | 26.13        | 26.01        | 26.00        | 25.98        | 26.04        | 25.91        | 26.09        | 26.29        | 25.94        | 25.71              | 25.71              | 25.71              | 25.71              | 25.71              | 25.71              | 25.71              |
| Industrial Coking Coal <sup>a</sup>     | 25.53        | 25.57        | 25.63        | 25.63        | 25.65        | 25.63        | 25.63        | 25.60        | 25.60        | 25.61        | 25.61              | 25.61              | 25.61              | 25.61              | 25.61              | 25.61              | 25.61              |
| Industrial Other Coal <sup>a</sup>      | 25.82        | 25.80        | 25.74        | 25.66        | 25.57        | 25.55        | 25.56        | 25.80        | 25.84        | 25.82        | 25.82              | 25.82              | 25.82              | 25.82              | 25.82              | 25.82              | 25.82              |
| Utility Coal <sup>a,c</sup>             | 25.96        | 25.93        | 26.00        | 26.00        | 26.05        | 26.09        | 26.10        | 26.09        | 26.04        | 26.05        | 26.05              | 26.05              | 26.05              | 26.05              | 26.05              | 26.05              | 26.05              |
| <b>Pipeline Natural Gas<sup>d</sup></b> | <b>14.45</b> | <b>14.46</b> | <b>14.47</b> | <b>14.46</b> | <b>14.46</b> | <b>14.44</b> | <b>14.46</b> | <b>14.46</b> | <b>14.46</b> | <b>14.46</b> | <b>14.46</b>       | <b>14.46</b>       | <b>14.46</b>       | <b>14.46</b>       | <b>14.46</b>       | <b>14.46</b>       | <b>14.46</b>       |
| <b>Flare Gas</b>                        | <b>15.31</b>       | <b>15.31</b>       | <b>15.31</b>       | <b>15.31</b>       | <b>15.31</b>       | <b>15.31</b>       | <b>15.31</b>       |
| <b>Petroleum</b>                        |              |              |              |              |              |              |              |              |              |              |                    |                    |                    |                    |                    |                    |                    |
| Asphalt and Road Oil                    | 20.55        | 20.55        | 20.55        | 20.55        | 20.55        | 20.55        | 20.55        | 20.55        | 20.55        | 20.55        | 20.55              | 20.55              | 20.55              | 20.55              | 20.55              | 20.55              | 20.55              |
| Aviation Gasoline                       | 18.86        | 18.86        | 18.86        | 18.86        | 18.86        | 18.86        | 18.86        | 18.86        | 18.86        | 18.86        | 18.86              | 18.86              | 18.86              | 18.86              | 18.86              | 18.86              | 18.86              |
| Distillate Fuel Oil No. 1               | 19.98        | 19.98        | 19.98        | 19.98        | 19.98        | 19.98        | 19.98        | 19.98        | 19.98        | 19.98        | 19.98              | 19.98              | 19.98              | 19.98              | 19.98              | 19.98              | 19.98              |
| Distillate Fuel Oil No. 2               | 20.17        | 20.17        | 20.17        | 20.17        | 20.17        | 20.17        | 20.17        | 20.17        | 20.17        | 20.17        | 20.17              | 20.17              | 20.17              | 20.17              | 20.17              | 20.17              | 20.17              |
| Distillate Fuel Oil No. 4               | 20.47        | 20.47        | 20.47        | 20.47        | 20.47        | 20.47        | 20.47        | 20.47        | 20.47        | 20.47        | 20.47              | 20.47              | 20.47              | 20.47              | 20.47              | 20.47              | 20.47              |
| Jet Fuel <sup>a</sup>                   | 19.40        | 19.34        | 19.70        | 19.70        | 19.70        | 19.70        | 19.70        | 19.70        | 19.70        | 19.70        | 19.70              | 19.70              | 19.70              | 19.70              | 19.70              | 19.70              | 19.70              |
| Kerosene                                | 19.96        | 19.96        | 19.96        | 19.96        | 19.96        | 19.96        | 19.96        | 19.96        | 19.96        | 19.96        | 19.96              | 19.96              | 19.96              | 19.96              | 19.96              | 19.96              | 19.96              |
| LPG (energy use) <sup>d</sup>           | 16.86        | 16.82        | 16.89        | 16.87        | 16.85        | 16.86        | 16.84        | 16.84        | 16.83        | 16.82        | 16.83              | 16.83              | 16.83              | 16.83              | 16.83              | 16.83              | 16.83              |
| LPG (non-energy use) <sup>d</sup>       | 17.06        | 17.09        | 17.09        | 17.10        | 17.09        | 17.09        | 17.07        | 17.06        | 17.06        | 17.05        | 17.06              | 17.06              | 17.06              | 17.06              | 17.06              | 17.06              | 17.06              |
| Lubricants                              | 20.20        | 20.20        | 20.20        | 20.20        | 20.20        | 20.20        | 20.20        | 20.20        | 20.20        | 20.20        | 20.20              | 20.20              | 20.20              | 20.20              | 20.20              | 20.20              | 20.20              |
| Motor Gasoline <sup>d</sup>             | 19.42        | 19.36        | 19.33        | 19.34        | 19.38        | 19.36        | 19.38        | 19.36        | 19.45        | 19.56        | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              |
| Residual Fuel No. 5                     | 19.89        | 19.89        | 19.89        | 19.89        | 19.89        | 19.89        | 19.89        | 19.89        | 19.89        | 19.89        | 19.89              | 19.89              | 19.89              | 19.89              | 19.89              | 19.89              | 19.89              |
| Residual Fuel No. 6                     | 20.48        | 20.48        | 20.48        | 20.48        | 20.48        | 20.48        | 20.48        | 20.48        | 20.48        | 20.48        | 20.48              | 20.48              | 20.48              | 20.48              | 20.48              | 20.48              | 20.48              |
| <b>Other Petroleum</b>                  |              |              |              |              |              |              |              |              |              |              |                    |                    |                    |                    |                    |                    |                    |
| Av. Gas Blend Comp.                     | 18.87        | 18.87        | 18.87        | 18.87        | 18.87        | 18.87        | 18.87        | 18.87        | 18.87        | 18.87        | 18.87              | 18.87              | 18.87              | 18.87              | 18.87              | 18.87              | 18.87              |
| Mo. Gas Blend Comp <sup>c</sup>         | 19.42        | 19.36        | 19.33        | 19.34        | 19.38        | 19.36        | 19.38        | 19.36        | 19.45        | 19.56        | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              | 19.46              |
| Crude Oil <sup>d</sup>                  | 20.15        | 20.21        | 20.22        | 20.27        | 20.28        | 20.25        | 20.31        | 20.31        | 20.28        | 20.28        | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              |
| Misc. Products <sup>d</sup>             | 20.15        | 20.21        | 20.22        | 20.27        | 20.28        | 20.25        | 20.31        | 20.31        | 20.28        | 20.28        | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              | 20.31              |

|                              |             |             |             |             |             |             |             |             |             |             |             |             |             |             |             |             |             |
|------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Misc. Products (Terr.)       | 20.15       | 20.21       | 20.22       | 20.27       | 20.28       | 20.25       | 20.31       | 20.31       | 20.28       | 20.28       | 20.31       | 20.31       | 20.31       | 20.31       | 20.31       | 20.31       | 20.31       |
| Naphtha (<401 deg. F)        | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       | 18.55       |
| Other oil (>401 deg. F)      | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       | 20.17       |
| Pentanes Plus                | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       | 19.10       |
| Petroleum Coke               | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       | 27.85       |
| Still Gas                    | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       | 18.20       |
| Special Naphtha              | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       | 19.74       |
| Unfinished Oils <sup>d</sup> | 20.15       | 20.21       | 20.22       | 20.27       | 20.28       | 20.25       | 20.31       | 20.31       | 20.28       | 20.28       | 20.31       | 20.31       | 20.31       | 20.31       | 20.31       | 20.31       | 20.31       |
| Waxes                        | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       |
| Other Wax and Misc.          | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       | 19.80       |
| <b>Geothermal</b>            | <b>2.05</b> |

<sup>a</sup>C contents vary annually based on changes in annual mix of production and end-use consumption of coal from each producing state.

<sup>b</sup>EIA discontinued collection of residential sector coal consumption data in 2008, because consumption of coal in the residential sector is extremely limited. Therefore, the number cited here is developed from commercial/institutional consumption.

<sup>c</sup>C content for utility coal used in the electric power calculations. All coefficients based on higher heating value. Higher heating value (gross heating value) is the total amount of heat released when a fuel is burned. Coal, crude oil, and natural gas all include chemical compounds of carbon and hydrogen. When those fuels are burned, the carbon and hydrogen combine with oxygen in the air to produce CO<sub>2</sub> and water. Some of the energy released in burning goes into transforming the water into steam and is usually lost. The amount of heat spent in transforming the water into steam is counted as part of gross heat content. Lower heating value (net heating value), in contrast, does not include the heat spent in transforming the water into steam. Using a simplified methodology based on International Energy Agency defaults, higher heating value can be converted to lower heating value for coal and petroleum products by multiplying by 0.95 and for natural gas by multiplying by 0.90. Carbon content coefficients are presented in higher heating value because U.S. energy statistics are reported by higher heating value.

<sup>d</sup>C contents vary annually based on changes in fuel composition.

## Coal

Approximately one-third of all U.S. CO<sub>2</sub> emissions from fossil fuel combustion are associated with coal consumption. Although the IPCC (2006) guidelines provide C contents for coal according to rank, it was necessary to develop C content coefficients by consuming sector to match the format in which coal consumption is reported by EIA. Because the C content of coal varies by the state in which it was mined and by coal rank, and because the sources of coal for each consuming sector vary by year, the weighted average C content for coal combusted in each consuming sector also varies over time. A time series of C contents by coal rank and consuming sector appears in Table A-42.<sup>9</sup>

### Methodology

The methodology for developing C contents for coal by consuming sector consists of four steps. An additional step has been taken to calculate C contents by coal rank to facilitate comparison with IPCC default values.

#### Step 1. Determine Carbon Contents by Rank and by State of Origin

Carbon contents by rank and state of origin are estimated on the basis of 7,092 coal samples, 6,588 of which were collected by the U.S. Geological Survey (USGS 1998) and 504 samples that come from the Pennsylvania State University database (PSU 2010). These coal samples are classified according to rank and state of origin. For each rank in each state, the average heat content and C content of the coal samples are calculated based on the proximate (heat) and ultimate (percent carbon) analyses of the samples. Dividing the C content (reported in pounds CO<sub>2</sub>) by the heat content (reported in million Btu or MMBtu) yields an average C content coefficient. This coefficient is then converted into units of MMT C/QBtu.

#### Step 2: Determine Weighted Average Carbon Content by State

Carbon contents by rank and origin calculated in Step 1 are then weighted by the annual share of state production that was each rank. State production by rank is obtained from the EIA. This step yields a single carbon content per state that varies annually based on production. However, most coal-producing states produce only one rank of coal. For these states the weighted factor equals the carbon content calculated in Step 1 and is constant across the time series.

#### Step 3: Allocate Sectoral Consumption by State of Origin

U.S. energy statistics<sup>10</sup> through 2007 provide data on the origin of coal used in four areas: 1) the electric power industry, 2) industrial coking, 3) all other industrial uses, and 4) the residential and commercial end-use sectors.<sup>11</sup> Because U.S. energy statistics do not provide the distribution of coal rank consumed by each consuming sector, it is assumed that each sector consumes a representative mixture of coal ranks from a particular state that matches the mixture of all coal produced in that state during the year. Thus, the weighted state-level factor developed in Step 2 is applied.

#### Step 4: Weight Sectoral Carbon Contents to Reflect the Rank and State of Origin of Coal Consumed

Sectoral C contents are calculated by multiplying the share of coal purchased from each state by the state's weighted C content estimated in Step 2. The resulting partial C contents are then totaled across all states to generate a national sectoral C content.

$$C_{\text{sector}} = S_{\text{state1}} \times C_{\text{state1}} + S_{\text{state2}} \times C_{\text{state2}} + \dots + S_{\text{state50}} \times C_{\text{state50}}$$

where,

<sup>9</sup> For a comparison to earlier estimated carbon contents see *Chronology and Explanation of Changes in Individual Carbon Content Coefficients of Fossil Fuels* near the end of this annex.

<sup>10</sup> U.S. Energy Information Administration (EIA). *Coal Distribution – Annual* (2001-2008); and *Coal Industry Annual* (1990-2000).

<sup>11</sup> Beginning in 2008, the EIA collects and reports data on commercial and institutional coal consumption, rather than residential and commercial consumption. Thus, the residential / commercial coal coefficient reported in Table A-41 for 2009 represents the mix of coal consumed by commercial and institutional users. Currently, only an extremely small amount of coal is consumed in the U.S. residential sector.

$C_{\text{sector}}$  = The C content by consuming sector;  
 $S_{\text{state}}$  = The portion of consuming sector coal consumption attributed to production from a given state;  
 $C_{\text{state}}$  = The estimated weighted C content of all ranks produced in a given state.

**Table A-42: Carbon Content Coefficients for Coal by Consuming Sector and Coal Rank (MMT C/QBtu) (1990-2014)**

| Consuming Sector           | 1990  | 1995  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  | 2008  | 2009  | 2010  | 2011  | 2012  | 2013  | 2014  |
|----------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Electric Power             | 25.96 | 25.93 | 26.00 | 26.00 | 26.05 | 26.09 | 26.10 | 26.09 | 26.04 | 26.05 | 26.05 | 26.05 | 26.05 | 26.05 | 26.05 | 26.05 | 26.05 |
| Industrial Coking          | 25.53 | 25.57 | 25.63 | 25.63 | 25.65 | 25.63 | 25.63 | 25.60 | 25.60 | 25.61 | 25.61 | 25.61 | 25.61 | 25.61 | 25.61 | 25.61 | 25.61 |
| Other Industrial           | 25.82 | 25.80 | 25.74 | 25.66 | 25.57 | 25.55 | 25.56 | 25.80 | 25.84 | 25.82 | 25.82 | 25.82 | 25.82 | 25.82 | 25.82 | 25.82 | 25.82 |
| Residential/<br>Commercial | 26.20 | 26.13 | 26.01 | 26.00 | 25.98 | 26.04 | 25.91 | 26.09 | 26.29 | 25.94 | 25.71 | 25.71 | 25.71 | 25.71 | 25.71 | 25.71 | 25.71 |
| <b>Coal Rank</b>           |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Anthracite                 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 | 28.28 |
| Bituminous                 | 25.38 | 25.42 | 25.45 | 25.46 | 25.46 | 25.45 | 25.45 | 25.45 | 25.45 | 25.45 | 25.44 | 25.44 | 25.44 | 25.44 | 25.44 | 25.44 | 25.44 |
| Sub-bituminous             | 26.50 | 26.50 | 26.49 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 | 26.50 |
| Lignite                    | 26.58 | 26.59 | 26.61 | 26.62 | 26.63 | 26.62 | 26.62 | 26.62 | 26.62 | 26.64 | 26.65 | 26.65 | 26.65 | 26.65 | 26.65 | 26.65 | 26.65 |

<sup>a</sup> In 2008, the EIA began collecting consumption data for commercial and institutional consumption rather than commercial and residential consumption.  
Sources: C content coefficients calculated from USGS (1998) and PSU (2010); data presented in EPA (2010b).

### Step 5: Develop National-Level Carbon Contents by Rank for Comparison to IPCC Defaults

Although not used to calculate emissions, national-level C contents by rank are more easily compared to C contents of other countries than are sectoral C contents. This step requires weighting the state-level C contents by rank developed under Step 1 by overall coal production by state and rank. Each state-level C content by rank is multiplied by the share of national production of that rank that each state represents. The resulting partial C contents are then summed across all states to generate an overall C content for each rank.

$$N_{\text{rank}} = P_{\text{rank1}} \times C_{\text{rank1}} + P_{\text{rank2}} \times C_{\text{rank2}} + \dots + P_{\text{rankn}} \times C_{\text{rankn}}$$

where,

|                   |   |   |
|-------------------|---|---|
| $N_{\text{rank}}$ | = | The national C content by rank;   |
| $P_{\text{rank}}$ | = | The portion of U.S. coal production of a given rank attributed to each state; and |
| $C_{\text{rank}}$ | = | The estimated C content of a given rank in each state.                            |

### Data Sources

The ultimate analysis of coal samples was based on the 7,092 coal samples, 6,588 of which are from USGS (1998) and 504 that come from the Pennsylvania State University Coal Database (PSU 2010). Data contained in the USGS's CoalQual Database are derived primarily from samples taken between 1973 and 1989, and were largely reported in State Geological Surveys. Data in the PSU Coal Database are mainly from samples collected by PSU since 1967 and are housed at the PSU Sample Bank. Only the subset of PSU samples that are whole-seam channel samples are included in the development of carbon factors in order to increase data accuracy.

Data on coal consumption by sector and state of origin, as well as coal production by state and rank, were obtained from EIA. The EIA's *Annual Coal Report* (EPA 2001-2009a) is the source for state coal production by rank from 2001 through 2008. In prior years, the EIA reported this data in its *Coal Industry Annual* (EPA 1990 through 2001). Data for coal consumption by state of origin and consuming sector for 2001 through 2008 was obtained from the EIA's *Coal Distribution – Annual* (EPA 2001 through 2009b). For 1990-2000, end-use data was obtained from the *Coal Industry Annual* (EPA 1990 through 2001).

### Uncertainty

Carbon contents vary considerably by state. Bituminous coal production and sub-bituminous coal production represented 47.3 percent and 46.1 percent of total U.S. supply in 2008, respectively. State average C content coefficients for bituminous coal vary from a low of 85.59 kg CO<sub>2</sub> per MMBtu in Texas to a high of 105.21 kg CO<sub>2</sub> per MMBtu in Montana. However, Texas bituminous coal is considered anomalous,<sup>12</sup> has not been produced since 2004 and production since 1990 peaked at just 446,000 short tons in 1996. The next lowest average emission factor for bituminous coal is found in Western Kentucky (91.36 kg CO<sub>2</sub> per MMBtu). In 2000, Montana produced no bituminous coal and Western Kentucky production accounted for just 4.5 percent of overall bituminous production. In 2008, more than 60 percent of bituminous coal was produced in three states: West Virginia, Kentucky (predominantly from the Eastern production region), and Pennsylvania, and this share has remained fairly constant since 1990. These three states show a variation in C content for bituminous coals of ±0.7 percent, based on more than 2,000 samples (see Table A-43).

Similarly, the C content coefficients for sub-bituminous coal range from 91.29 kg CO<sub>2</sub> per MMBtu in Utah to 98.10 kg CO<sub>2</sub> per MMBtu in Alaska. However, Utah has no recorded production of sub-bituminous coal since 1990. Production of sub-bituminous coal in Alaska has made up less than 0.7 percent of total sub-bituminous production since 1990, with even this small share declining over time. Wyoming has represented between 75 percent and 87 percent of total sub-bituminous coal production in the United States in each year since 1990. Thus, the C content coefficient for Wyoming (97.22 kg CO<sub>2</sub> per MMBtu), based on 455 samples, dominates the national average.

The interquartile range of C content coefficients among samples of sub-bituminous coal in Wyoming was ±1.5 percent from the mean. Similarly, this range among samples of bituminous coal from West Virginia, Kentucky, and Pennsylvania was ±1.2 percent or less for each state. The large number of samples and the low variability within the sample set of the states that represent the predominant source of supply of U.S. coal suggest that the uncertainty in this factor is very low, on the order of ±1.0 percent.

---

<sup>12</sup> See, for example: San Filippo, 1999. USGS. (U.S. Geological Survey Open-File Report 99-301), Ch. 4.

For comparison, J. Quick (2010) completed an analysis similar in methodology to that used here, in order to generate national average C emission factors as well as county-level factors. This study's rank-based national average factors have a maximum deviation from the factors developed in this Inventory report of -0.55 percent, which is for lignite (range: -0.55 to +0.1 percent). This corroboration further supports the assertion of minimal uncertainty in the application of the rank-based factors derived for the purposes of this Inventory.

**Table A-43: Variability in Carbon Content Coefficients by Rank Across States (Kilograms CO<sub>2</sub> Per MMBtu)**

| State               | Number of Samples | Bituminous   | Sub-bituminous | Anthracite    | Lignite      |
|---------------------|-------------------|--------------|----------------|---------------|--------------|
| Alabama             | 951               | 92.84        | -              | -             | 99.10        |
| Alaska              | 91                | 98.33        | 98.10          | -             | 98.65        |
| Arizona             | 15                | 93.94        | 97.34          | -             | -            |
| Arkansas            | 80                | 96.36        | -              | -             | 94.97        |
| Colorado            | 318               | 94.37        | 96.52          | -             | 101.10       |
| Georgia             | 35                | 95.01        | -              | -             | -            |
| Idaho               | 1                 | -            | 94.90          | -             | -            |
| Illinois            | 57                | 92.33        | -              | -             | -            |
| Indiana             | 146               | 92.65        | -              | -             | -            |
| Iowa                | 100               | 91.87        | -              | -             | -            |
| Kansas              | 29                | 90.91        | -              | -             | -            |
| Kentucky            | 897               | 92.61        | -              | -             | -            |
| Louisiana           | 1                 | -            | -              | -             | 96.01        |
| Maryland            | 47                | 94.29        | -              | -             | -            |
| Massachusetts       | 3                 | -            | -              | 114.82        | -            |
| Michigan            | 3                 | 92.88        | -              | -             | -            |
| Mississippi         | 8                 | -            | -              | -             | 98.19        |
| Missouri            | 111               | 91.71        | -              | -             | -            |
| Montana             | 309               | 105.21       | 97.73          | 103.60        | 99.40        |
| Nevada              | 2                 | 94.41        | -              | -             | 99.86        |
| New Mexico          | 185               | 94.29        | 94.89          | 103.92        | -            |
| North Dakota        | 202               | -            | 93.97          | -             | 99.48        |
| Ohio                | 674               | 91.84        | -              | -             | -            |
| Oklahoma            | 63                | 92.33        | -              | -             | -            |
| Pennsylvania        | 861               | 93.33        | -              | 103.68        | -            |
| Tennessee           | 61                | 92.82        | -              | -             | -            |
| Texas               | 64                | 85.59        | 94.19          | -             | 94.47        |
| Utah                | 169               | 95.75        | 91.29          | -             | -            |
| Virginia            | 470               | 93.51        | -              | 98.54         | -            |
| Washington          | 18                | 94.53        | 97.36          | 102.53        | 106.55       |
| West Virginia       | 612               | 93.84        | -              | -             | -            |
| Wyoming             | 503               | 94.80        | 97.22          | -             | -            |
| <b>U.S. Average</b> | <b>7,092</b>      | <b>93.13</b> | <b>96.94</b>   | <b>104.29</b> | <b>98.63</b> |

Note: "-" Indicates no sample data available.

Sources: Calculated from USGS (1998), and PSU (2010); data presented in EPA (2010).

## Natural Gas

Natural gas is predominantly composed of methane (CH<sub>4</sub>), which is 75 percent C by weight and contains 14.2 MMT C/QBtu (higher heating value), but it may also contain many other compounds that can lower or raise its overall C content. These other compounds may be divided into two classes: (1) natural gas liquids (NGLs) and (2) non-hydrocarbon gases. The most common NGLs are ethane (C<sub>2</sub>H<sub>6</sub>), propane (C<sub>3</sub>H<sub>8</sub>), butane (C<sub>4</sub>H<sub>10</sub>), and, to a lesser extent, pentane (C<sub>5</sub>H<sub>12</sub>) and hexane (C<sub>6</sub>H<sub>14</sub>). Because the NGLs have more C atoms than CH<sub>4</sub> (which has only one), their presence increases the overall C content of natural gas. NGLs have a commercial value greater than that of CH<sub>4</sub>, and therefore are usually separated from raw natural gas at gas processing plants and sold as separate products. Ethane is typically used as a petrochemical

feedstock, propane and butane have diverse uses, and natural gasoline<sup>13</sup> contributes to the gasoline/naphtha "octane pool," used primarily to make motor gasoline.

Raw natural gas can also contain varying amounts of non-hydrocarbon gases, such as CO<sub>2</sub>, nitrogen, helium and other noble gases, and hydrogen sulfide. The share of non-hydrocarbon gases is usually less than 5 percent of the total, but there are individual natural gas reservoirs where the share can be much larger. The treatment of non-hydrocarbon gases in raw gas varies. Hydrogen sulfide is always removed. Inert gases are removed if their presence is substantial enough to reduce the energy content of the gas below pipeline specifications (see Step 1, below). Otherwise, inert gases will usually be left in the natural gas. Because the raw gas that is usually flared (see Step 2, below) contains NGLs and CO<sub>2</sub>, it will typically have a higher overall C content than gas that has been processed and moved to end-use customers via transmission and distribution pipelines.

## **Methodology**

The methodology for estimating the C contents of pipeline and flared natural gas can be described in five steps.

### **Step 1: Define pipeline-quality natural gas**

In the United States, pipeline-quality natural gas is required to have an energy content greater than 970 Btu per cubic foot, but less than 1,100 Btu per cubic foot. Hydrogen sulfide content must be negligible. Typical pipeline-quality natural gas is about 95 percent CH<sub>4</sub>, 3 percent NGLs, and 2 percent non-hydrocarbon gases, of which approximately half is CO<sub>2</sub>.

However, there remains a range of gas compositions that are consistent with pipeline specifications. The minimum C content coefficient for natural gas would match that for pure CH<sub>4</sub>, which equates to an energy content of 1,005 Btu per standard cubic foot. Gas compositions with higher or lower Btu content tend to have higher C emission factors, because the "low" Btu gas has a higher content of inert gases (including CO<sub>2</sub> offset with more NGLs), while "high" Btu gas tends to have more NGLs.

### **Step 2: Define flared gas**

Every year, a certain amount of natural gas is flared in the United States. There are several reasons that gas is flared:

- There may be no market for some batches of natural gas, the amount may be too small or too variable, or the quality might be too poor to justify treating the gas and transporting it to market (such is the case when gas contains large shares of CO<sub>2</sub>). Most natural gas that is flared for these reasons is "rich" associated gas, with relatively high energy content, high NGL content, and a high C content.
- Gas treatment plants may flare substantial volumes of natural gas because of "process upsets," because the gas is "off spec," or possibly as part of an emissions control system. Gas flared at processing plants may be of variable quality.

Data on the energy content of flare gas, as reported by states to EIA, indicate an average energy content of 1,130 Btu per standard cubic foot (EIA 1994). Flare gas may have an even higher energy content than reported by EIA since rich associated gas can have energy contents as high as 1,300 to 1,400 Btu per cubic foot.

### **Step 3: Determine a relationship between carbon content and heat content**

A relationship between C content and heat content may be used to develop a C content coefficient for natural gas consumed in the United States. In 1994, EIA examined the composition (including C contents) of 6,743 samples of pipeline-quality natural gas from utilities and/or pipeline companies in 26 cities located in 19 states. To demonstrate that these samples were representative of actual natural gas "as consumed" in the United States, their heat content was compared to that of the national average. For the most recent year, the average heat content of natural gas consumed in the United States was 1,029 Btu per cubic foot, and has varied by less than 1 percent (1,027 to 1,029 Btu per cubic foot) over the past 5 years. Meanwhile, the average heat content of the 6,743 samples was 1,027 Btu per cubic foot, and the median heat content was 1,031 Btu per cubic foot. Thus, the average heat content of the sample set falls well within the typical range of natural gas

---

<sup>13</sup> A term used in the gas processing industry to refer to a mixture of liquid hydrocarbons (mostly pentanes and heavier hydrocarbons) extracted from natural gas.

consumed in the United States, suggesting that these samples continue to be representative of natural gas “as consumed” in the United States. The average and median composition of these samples appear in Table A-44.

**Table A-44: Composition of Natural Gas (Percent)**

| Compound                                  | Average | Median |
|---|---------|--------|
| Methane                                   | 93.07   | 95.00  |
| Ethane                                    | 3.21    | 2.79   |
| Propane                                   | 0.59    | 0.48   |
| Higher Hydrocarbons                       | 0.32    | 0.30   |
| Non-hydrocarbons                          | 2.81    | 1.43   |
| Higher Heating Value (Btu per cubic foot) | 1,027   | 1,031  |

Source: Gas Technology Institute (1992).

Carbon contents were calculated for a series of sub-samples based on their CO<sub>2</sub> content and heat content. Carbon contents were calculated for the groups of samples with less than 1.0 percent (n=5,181) and less than 1.5 percent CO<sub>2</sub> only (n=6,522) and those with less than 1.0 or 1.5 percent CO<sub>2</sub> and less than 1,050 Btu/cf (n=4,888 and 6,166, respectively). These stratifications were chosen to exclude samples with CO<sub>2</sub> content and heat contents outside the range of pipeline-quality natural gas. In addition, hexane was removed from the samples since it is usually stripped out of raw natural gas before delivery because it is a valuable natural gas liquid used as a feedstock for gasoline. The average carbon contents for the four separate sub-samples are shown below in Table A-45.

**Table A-45: Carbon Content of Pipeline-Quality Natural Gas by CO<sub>2</sub> and Heat Content (MMT C/QBtu)**

| Sample                                    | Average Carbon Content |
|---|------------------------|
| Full Sample                               | 14.48                  |
| < 1.0% CO <sub>2</sub>                    | 14.43                  |
| < 1.5% CO <sub>2</sub>                    | 14.47                  |
| < 1.0 % CO <sub>2</sub> and <1,050 Btu/cf | 14.42                  |
| < 1.5 % CO <sub>2</sub> and <1,050 Btu/cf | 14.47                  |

Source: EPA (2010).

#### Step 4. Apply carbon content coefficients developed in Step 3 to pipeline natural gas

A regression analysis was performed on the sub-samples in to further examine the relationship between carbon (C) content and heat content. The regression used carbon content as the dependent variable and heat content as the independent variable. The resulting R-squared values<sup>14</sup> for each of the sub-samples ranged from 0.79 for samples with less than 1.5 percent CO<sub>2</sub> and under 1,050 Btu/cf to 0.91 for samples containing less than 1.0 percent CO<sub>2</sub> only. However, the sub-sample with less than 1.5 percent CO<sub>2</sub> and 1,050 Btu/cf was chosen as the representative sample for two reasons. First, it most accurately reflects the range of CO<sub>2</sub> content and heat content of pipeline quality natural gas. Secondly, the R-squared value, although it is the lowest of the sub-groups tested, remains relatively high. This high R-squared indicates a low percentage of variation in C content as related to heat content. The regression for this sub-sample resulted in the following equation:

$$C \text{ Content} = (0.011 \times \text{Heat Content}) + 3.5341$$

This equation was used to estimate the annual predicted carbon content of natural gas from 1990 to 2010 based on the EIA’s national average pipeline-quality gas heat content for each year. The table of average C contents for each year is shown below in Table A-46.

**Table A-46: Carbon Content Coefficients for Natural Gas (MMT Carbon/QBtu)**

| Fuel Type   | 1990  | 1995  | 2000  | 2005  | 2006  | 2007  | 2008  | 2009  | 2010  | 2011  | 2012  | 2013  | 2014  |
|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Natural Gas | 14.45 | 14.46 | 14.47 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46 | 14.46 |

Source: EPA (2010).

<sup>14</sup> R-squared represents the percentage of variation in the dependent variable (in this case carbon content) explained by variation in the independent variables.

### Step 5. Apply carbon content coefficients developed in Step 3 to flare gas

Selecting a C content coefficient for flare gas was much more difficult than for pipeline natural gas, because of the uncertainty of its composition and of the combustion efficiency of the flare. Because EIA estimates the heat content of flare gas at 1,130 Btu per cubic foot, the average C content for samples with more than 1,100 Btu per cubic foot (n=18) was chosen as the relevant sub-sample from which to calculate a flare gas carbon content. The sample dataset did not include any samples with more than 1,130 Btu per cubic foot.

Hexane was not removed from flare gas samples since it is assumed that natural gas liquids are present in samples with higher heat contents. Carbon contents were calculated for each sample with a heat content of more than 1,100 Btu per cubic foot. The simple average C content for the sample sub-set representing flare gas is shown below in Table A-47.

**Table A-47: Carbon Content of Flare Gas (MMT C/QBtu)**

| Relevant Sub-Sample | Average Carbon Content |
|---------------------|------------------------|
| >1,100 Btu/cf       | 15.31                  |

Source: EPA (2010).

### Data Sources


Natural gas samples were obtained from the Gas Technology Institute (1992). Average heat content data for natural gas consumed in the United States was taken from EIA (2009a).

### Uncertainty

The assignment of C content coefficients for natural gas, and particularly for flare gas, requires more subjective judgment than the methodology used for coal. This subjective judgment may introduce additional uncertainty.

Figure A-1 shows the relationship between the calculated C content for each natural gas sample and its energy content. This figure illustrates the relatively restricted range of variation in both the energy content (which varies by about 6 percent from average) and the C emission coefficient of natural gas (which varies by about 5 percent). Thus, the knowledge that gas has been sold via pipeline to an end-use consumer allows its C emission coefficient to be predicted with an accuracy of  $\pm 5.0$  percent.

**Figure A-1: Carbon Content for Samples of Pipeline-Quality Natural Gas Included in the Gas Technology Institute Database**



Source: EIA (1994) Energy Information Administration, Emissions of Greenhouse Gases in the United States 1987-1992, U.S. Department of Energy, Washington, DC, November, 1994, DOE/EIA 0573, Appendix A.

Natural gas suppliers may achieve the same overall energy content from a wide variety of methane, higher hydrocarbon, and non-hydrocarbon gas combinations. Thus, the plot reveals large variations in C content for a single Btu value. In fact, the variation in C content for a single Btu value may be nearly as great as the variation for the whole sample. As a result, while energy content has some predictive value, the specific energy content does not substantially improve the accuracy of an estimated C content coefficient beyond the  $\pm 5.0$  percent offered with the knowledge that it is of pipeline-quality.

The plot of C content also reveals other interesting anomalies. Samples with the lowest emissions coefficients tend to have energy contents of about 1,000 Btu per cubic foot. They are composed of almost pure CH<sub>4</sub>. Samples with a greater proportion of NGLs (e.g., ethane, propane, and butane) tend to have energy contents greater than 1,000 Btu per cubic foot, along with higher emissions coefficients. Samples with a greater proportion of inert gases tend to have lower energy content, but they usually contain CO<sub>2</sub> as one of the inert gases and, consequently, also tend to have higher emission coefficients (see left side of Figure A-1).

For the full sample (n=6,743), the average C content of a cubic foot of gas was 14.48 MMT C/QBtu (see Table A-46). Additionally, a regression analysis using the full sample produced a predicted C content of 14.49 MMT C/QBtu based on a heat content of 1,029 Btu/cf (the average heat content in the United States for the most recent year). However, these two values include an upward influence on the resulting carbon content that is caused by inclusion in the sample set of the samples that contain large amounts of inert carbon dioxide and those samples with more than 1,050 Btu per cubic foot that contain an unusually large amount of NGLs. Because typical gas consumed in the United States does not contain such a large amount of carbon dioxide or natural gas liquids, a C content of 14.47 MMT C/QBtu, based on samples with less than 1.5 percent carbon dioxide and less than 1,050 Btu per cubic foot, better represents the pipeline-quality fuels typically consumed.

## Petroleum

There are four critical determinants of the C content coefficient for a petroleum-based fuel:

- The density of the fuel (e.g., the weight in kilograms of one barrel of fuel);
- The fraction by mass of the product that consists of hydrocarbons, and the fraction of non-hydrocarbon impurities;
- The specific types of “families” of hydrocarbons that make up the hydrocarbon portion of the fuel; and
- The heat content of the fuel.

$$\text{where,} \quad C_{\text{fuel}} = (D_{\text{fuel}} \times S_{\text{fuel}}) / E_{\text{fuel}}$$

|                   |   |                                       |
|-------------------|---|---------------------------------------|
| $C_{\text{fuel}}$ | = | The C content coefficient of the fuel |
| $D_{\text{fuel}}$ | = | The density of the fuel               |
| $S_{\text{fuel}}$ | = | The share of the fuel that is C       |
| $E_{\text{fuel}}$ | = | The heat content of the fuel          |

Most of the density, carbon share, or heat contents applied to calculate the carbon coefficients for petroleum products that are described in this sub-Annex and applied to this emissions Inventory have been updated for this edition of the report. These changes have been made where necessary to increase the accuracy of the underlying data or to align the petroleum properties data used in this report with that developed for use in the *Mandatory Reporting of Greenhouse Gases Rule* (EPA 2009b).

Petroleum products vary between 5.6 degrees API gravity<sup>15</sup> (dense products such as asphalt and road oil) and 247 degrees (ethane). This is a range in density of 60 to 150 kilograms per barrel, or  $\pm 50$  percent. The variation in C content, however, is much smaller ( $\pm 5$  to 7 percent) for products produced by standard distillation refining: ethane is 80 percent C by weight, while petroleum coke is 90 to 92 percent C. This tightly bound range of C contents can be explained by basic petroleum chemistry (see below). Additional refining can increase carbon contents. Calcined coke, for example, is formed by heat treating petroleum coke to about 1600 degrees Kelvin (calcining), to expel volatile materials and increase the percentage of elemental C. This product can contain as much as 97 to 99 percent carbon. Calcined coke is mainly used in the aluminum and steel industry to produce C anodes.

### Petroleum Chemistry


Crude oil and petroleum products are typically mixtures of several hundred distinct compounds, predominantly hydrocarbons. All hydrocarbons contain hydrogen and C in various proportions. When crude oil is distilled into petroleum products, it is sorted into fractions by the boiling temperature of these hundreds of organic compounds. Boiling temperature is strongly correlated with the number of C atoms in each molecule. Petroleum products consisting of relatively simple molecules and few C atoms have low boiling temperatures, while larger molecules with more C atoms have higher boiling temperatures.

Products that boil off at higher temperatures are usually denser, which implies greater C content as well. Petroleum products with higher C contents, in general, have lower energy content per unit mass and higher energy content per unit volume than products with lower C contents. Empirical research led to the establishment of a set of quantitative relationships between density, energy content per unit weight and volume, and C and hydrogen content. Figure A-2 compares C content coefficients calculated on the basis of the derived formula with actual C content coefficients for a range of crude oils, fuel oils, petroleum products, and pure hydrocarbons. The actual fuel samples were drawn from the sources described below in the discussions of individual petroleum products.

---

<sup>15</sup> API gravity is an arbitrary scale expressing the gravity or density of liquid petroleum products, as established by the American Petroleum Institute (API). The measuring scale is calibrated in terms of degrees API. The higher the API gravity, the lighter the compound. Light crude oils generally exceed 38 degrees API and heavy crude oils are all crude oils with an API gravity of 22 degrees or below. Intermediate crude oils fall in the range of 22 degrees to 38 degrees API gravity. API gravity can be calculated with the following formula:  $\text{API Gravity} = (141.5 / \text{Specific Gravity}) - 131.5$ . Specific gravity is the density of a material relative to that of water. At standard temperature and pressure, there are 62.36 pounds of water per cubic foot, or 8.337 pounds water per gallon.

**Figure A-2: Estimated and Actual Relationships Between Petroleum Carbon Content Coefficients and Hydrocarbon Density**



Source: Carbon content factors for paraffins are calculated based on the properties of hydrocarbons in V. Guthrie (ed.), *Petroleum Products Handbook* (New York: McGraw Hill, 1960) p. 33. Carbon content factors from other petroleum products are drawn from sources described below. Relationship between density and emission factors based on the relationship between density and energy content in U.S. Department of Commerce, National Bureau of Standards, *Thermal Properties of Petroleum Products*, Miscellaneous Publication, No. 97 (Washington, D.C., 1929), pp.16-21, and relationship between energy content and fuel composition in S. Ringen, J. Lanum, and F.P. Miknis, "Calculating Heating Values from the Elemental Composition of Fossil Fuels," *Fuel*, Vol. 58 (January 1979), p.69.

The derived empirical relationship between C content per unit heat and density is based on the types of hydrocarbons most frequently encountered. Petroleum fuels can vary from this relationship due to non-hydrocarbon impurities and variations in molecular structure among classes of hydrocarbons. In the absence of more exact information, this empirical relationship offers a good indication of C content.

#### **Non-hydrocarbon Impurities**

Most fuels contain a certain share of non-hydrocarbon material. This is also primarily true of crude oils and fuel oils. The most common impurity is sulfur, which typically accounts for between 0.5 and 4 percent of the mass of most crude oils, and can form an even higher percentage of heavy fuel oils. Some crude oils and fuel oils also contain appreciable quantities of oxygen and nitrogen, typically in the form of asphaltenes or various acids. The nitrogen and oxygen content of crude oils can range from near zero to a few percent by weight. Lighter petroleum products have much lower levels of impurities, because the refining process tends to concentrate all of the non-hydrocarbons in the residual oil fraction. Light products usually contain less than 0.5 percent non-hydrocarbons by mass. Thus, the C content of heavy fuel oils can often be several percent lower than that of lighter fuels, due entirely to the presence of non-hydrocarbons.

#### **Variations in Hydrocarbon Classes**

Hydrocarbons can be divided into five general categories, each with a distinctive relationship between density and C content and physical properties. Refiners tend to control the mix of hydrocarbon types in particular products in order to give petroleum products distinct properties. The main classes of hydrocarbons are described below.

*Paraffins.* Paraffins are the most common constituent of crude oil, usually comprising 60 percent by mass. Paraffins are straight-chain hydrocarbons with the general formula  $C_nH_{2n+2}$ . Paraffins include ethane ( $C_2H_6$ ), propane ( $C_3H_8$ ), butane ( $C_4H_{10}$ ), and octane ( $C_8H_{18}$ ). As the chemical formula suggests, the C content of the paraffins increases with their C number: ethane is 79.89 percent C by weight, octane 84.12 percent. As the size of paraffin molecules increases, the C content approaches the limiting value of 85.7 percent asymptotical (see Figure A-3).

*Cycloparaffins.* Cycloparaffins are similar to paraffins, except that the C molecules form ring structures rather than straight chains, and consequently require two fewer hydrogen molecules than paraffins. Cycloparaffins always have the general formula  $C_nH_{2n}$  and are 85.63 percent C by mass, regardless of molecular size.


*Olefins.* Olefins are a very reactive and unstable form of paraffin: a straight chain with two carbon atoms double bonded together (thus are unsaturated) compared to the carbon atoms in a paraffin (which are saturated with hydrogen). They are never found in crude oil but are created in moderate quantities by the refining process. Gasoline, for example, may contain between 2 and 20 percent olefins. They also have the general formula  $C_nH_{2n}$ , and hence are also always 85.63 percent C by weight. Propylene ( $C_3H_6$ ), a common intermediate petrochemical product, is an olefin.

*Aromatics.* Aromatics are very reactive hydrocarbons that are relatively uncommon in crude oil (10 percent or less). Light aromatics increase the octane level in gasoline, and consequently are deliberately created by catalytic reforming of heavy naphtha. Aromatics also take the form of ring structures with some double bonds between C atoms. The most common aromatics are benzene ( $C_6H_6$ ), toluene ( $C_7H_8$ ), and xylene ( $C_8H_{10}$ ). The general formula for aromatics is  $C_nH_{2n-6}$ . Benzene is 92.26 percent C by mass, while xylene is 90.51 percent C by mass and toluene is 91.25 percent C by mass. Unlike the other hydrocarbon families, the C content of aromatics declines asymptotically toward 85.7 percent with increasing C number and density (see Figure A-3).

*Polynuclear Aromatics.* Polynuclear aromatics are large molecules with a multiple ring structure and few hydrogen atoms, such as naphthalene ( $C_{10}H_8$  and 93.71 percent C by mass) and anthracene ( $C_{14}H_{10}$  and 97.7 percent C). They are relatively rare but do appear in heavier petroleum products.

Figure A-3 illustrates the share of C by weight for each class of hydrocarbon. Hydrocarbon molecules containing 2 to 4 C atoms are all natural gas liquids; hydrocarbons with 5 to 10 C atoms are predominantly found in naphtha and gasoline; and hydrocarbon compounds with 12 to 20 C atoms comprise "middle distillates," which are used to make diesel fuel, kerosene and jet fuel. Larger molecules which can be vacuum distilled may be used as lubricants, waxes, and residual fuel oil or cracked and blended into the gasoline or distillate pools.

**Figure A-3: Carbon Content of Pure Hydrocarbons as a Function of Carbon Number**



Source: J.M. Hunt, *Petroleum Geochemistry and Geology* (San Francisco, CA, W.H. Freeman and Company, 1979), pp. 31-37.

If nothing is known about the composition of a particular petroleum product, assuming that it is 85.7 percent C by mass is not an unreasonable first approximation. Since denser products have higher C numbers, this guess would be most likely to be correct for crude oils and fuel oils. The C content of lighter products is more affected by the shares of paraffins and aromatics in the blend.

**Energy Content of Petroleum Products**

The exact energy content (gross heat of combustion) of petroleum products is not generally known. EIA estimates energy consumption in Btu on the basis of a set of industry-standard conversion factors. These conversion factors are generally accurate to within 3 to 5 percent.

**Individual Petroleum Products**

The United States maintains data on the consumption of more than twenty separate petroleum products and product categories. The C contents, heat contents, and density for each product are provided below in Table A-48. A description of the methods and data sources for estimating the key parameters for each individual petroleum product appears below.

**Table A-48: Carbon Content Coefficients and Underlying Data for Petroleum Products**

| Fuel                                  | 2008 Carbon Content<br>(MMT C/QBtu) | Gross Heat of Combustion<br>(MMBtu/Barrel) | Density<br>(API Gravity) | Percent<br>Carbon |
|---------------------------------------|-------------------------------------|--|--------------------------|-------------------|
| Motor Gasoline                        | 19.46                               | [a]  | [a]                      | [a]               |
| LPG(total)                            | 16.97                               | [b]  | [b]                      | [b]               |
| LPG (energy use)                      | 16.83                               | [b]  | [b]                      | [b]               |
| LPG (non-energy use)                  | 17.06                               | [b]  | [b]                      | [b]               |
| Jet Fuel                              | 19.70                               | 5.670                                      | 42.0                     | 86.30             |
| Distillate Fuel No. 1                 | 19.98                               | 5.822                                      | 35.3                     | 86.40             |
| Distillate Fuel No. 2                 | 20.17                               | 5.809                                      | 35.8                     | 87.30             |
| Distillate Fuel No. 4                 | 20.47                               | 6.135                                      | 23.2                     | 86.47             |
| Residual Fuel No. 5                   | 19.89                               | 5.879                                      | 33.0                     | 85.67             |
| Residual Fuel No. 6                   | 20.48                               | 6.317                                      | 15.5                     | 84.67             |
| Asphalt and Road Oil                  | 20.55                               | 6.636                                      | 5.6                      | 83.47             |
| Lubricants                            | 20.20                               | 6.065                                      | 25.7                     | 85.80             |
| Naphtha (< 400 deg. F) <sup>a</sup>   | 18.55                               | 5.248                                      | 62.4                     | 84.11             |
| Other Oils (>400 deg. F) <sup>a</sup> | 20.17                               | 5.825                                      | 35.8                     | 87.30             |
| Aviation Gas                          | 18.86                               | 5.048                                      | 69.0                     | 85.00             |
| Kerosene                              | 19.96                               | 5.825                                      | 35.3                     | 86.40             |
| Petroleum Coke                        | 27.85                               | 6.024                                      | -                        | 92.28             |
| Special Naphtha                       | 19.74                               | 5.248                                      | 52.0                     | 84.75             |
| Petroleum Waxes                       | 19.80                               | 5.537                                      | 43.3                     | 85.30             |
| Still Gas                             | 18.20                               | 6.000                                      | -                        | 77.70             |
| Crude Oil                             | 20.31                               | 5.800                                      | 31.2                     | 85.49             |
| Unfinished Oils                       | 20.31                               | 5.825                                      | 31.2                     | 85.49             |
| Miscellaneous Products                | 20.31                               | 5.796                                      | 31.2                     | 85.49             |
| Pentanes Plus                         | 19.10                               | 4.620                                      | 81.3                     | 83.63             |

[a] Calculation of the carbon content coefficient for motor gasoline in 2008 uses separate higher heating values for conventional and reformulated gasoline of 5.253 and 5.150, respectively (EIA 2008a). Densities and carbon shares (percent carbon) are annually variable and separated by both fuel formulation and grade, see Motor Gasoline and Blending Components, below, for details.

[b] LPG is a blend of multiple paraffinic hydrocarbons: ethane, propane, isobutane, and normal butane, each with their own heat content, density and C content, see Table A-51.

<sup>a</sup> Petrochemical feedstocks have been split into naphthas and other oils for this Inventory report. Parameters presented are for naphthas with a boiling temperature less than 400 degrees Fahrenheit. Other oils are petrochemical feedstocks with higher boiling points. They are assumed to have the same characteristics as distillate fuel oil no. 2.

Note: "-" Indicates no sample data available.

Sources: EIA (1994); EIA (2009a); EPA (2009b); and EPA (2010).

#### *Motor Gasoline and Motor Gasoline Blending Components*

Motor gasoline is a complex mixture of relatively volatile hydrocarbons with or without small quantities of additives, blended to form a fuel suitable for use in spark-ignition engines.<sup>16</sup> "Motor Gasoline" includes conventional gasoline; all types of oxygenated gasoline, including gasohol; and reformulated gasoline; but excludes aviation gasoline.

Gasoline is the most widely used petroleum product in the United States, and its combustion accounts for nearly 20 percent of all U.S. CO<sub>2</sub> emissions. EIA collects consumption data (i.e., "petroleum products supplied" to end-users) for several types of finished gasoline over the 1990 through 2014 time period: regular, mid-grade and premium conventional gasoline (all years) and regular, mid-grade and premium reformulated gasoline (November 1994 to 2014). Leaded and oxygenated gasoline are not separately included in the data used for this report.<sup>17</sup>

<sup>16</sup> Motor gasoline, as defined in ASTM Specification D 4814 or Federal Specification VV-G-1690C, is characterized as having a boiling range of 122 degrees to 158 degrees Fahrenheit at the 10-percent recovery point to 365 degrees to 374 degrees Fahrenheit at the 90-percent recovery point.

<sup>17</sup> Oxygenated gasoline volumes are included in the conventional gasoline data provided by EIA from 2007 onwards. Leaded gasoline was included in total gasoline by EIA until October 1993.

The American Society for Testing and Materials (ASTM) standards permit a broad range of densities for gasoline, ranging from 50 to 70 degrees API gravity, or 111.52 to 112.65 kilograms per barrel (EIA 1994), which implies a range of possible C and energy contents per barrel. Table A-49 reflects changes in the density of gasoline over time and across grades and formulations of gasoline through 2014.

**Table A-49: Motor Gasoline Density, 1990 – 2014 (Degrees API)**

| Fuel Grade                         | 1990 | 1995 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 |
|------------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>Conventional - Winter Grade</b> |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| Low Octane                         | 62.0 | 59.8 | 61.6 | 61.7 | 61.6 | 61.8 | 62.4 | 62.6 | 62.7 | 63.1 | 63.0 | 63.0 | 63.0 | 63.0 | 63.0 | 63.0 | 63.0 |
| High Octane                        | 59.0 | 58.0 | 59.7 | 59.1 | 59.0 | 59.9 | 60.7 | 60.9 | 60.0 | 60.3 | 60.9 | 60.9 | 60.9 | 60.9 | 60.9 | 60.9 | 60.9 |
| <b>Conventional - Summer Grade</b> |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| Low Octane                         | 58.2 | 56.1 | 56.8 | 57.2 | 56.5 | 56.8 | 57.4 | 57.9 | 57.8 | 57.5 | 58.6 | 58.6 | 58.6 | 58.6 | 58.6 | 58.6 | 58.6 |
| High Octane                        | 55.5 | 55.1 | 55.8 | 55.5 | 55.7 | 56.0 | 57.0 | 57.0 | 57.4 | 56.9 | 58.0 | 58.0 | 58.0 | 58.0 | 58.0 | 58.0 | 58.0 |
| <b>Reformulated - Winter Grade</b> |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| Low Octane                         | NA   | 61.9 | 62.7 | 62.6 | 61.9 | 62.1 | 62.7 | 62.8 | 62.3 | 62.1 | 62.4 | 62.4 | 62.4 | 62.4 | 62.4 | 62.4 | 62.4 |
| High Octane                        | NA   | 59.9 | 61.1 | 61.0 | 61.8 | 61.9 | 61.8 | 61.8 | 61.7 | 62.1 | 62.5 | 62.5 | 62.5 | 62.5 | 62.5 | 62.5 | 62.5 |
| <b>Reformulated - Summer Grade</b> |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| Low Octane                         | NA   | 58.5 | 58.4 | 58.8 | 58.2 | 59.1 | 58.1 | 58.4 | 58.7 | 58.5 | 59.1 | 59.1 | 59.1 | 59.1 | 59.1 | 59.1 | 59.1 |
| High Octane                        | NA   | 56.7 | 58.3 | 58.2 | 58.0 | 58.7 | 58.9 | 58.1 | 59.0 | 59.3 | 59.8 | 59.8 | 59.8 | 59.8 | 59.8 | 59.8 | 59.8 |

Notes: NA – Not Applicable, fuel type was not analyzed.

Source: National Institute of Petroleum and Energy Research (1990 through 2014).

The density of motor gasoline increased across all grades through 1994, partly as a result of the leaded gasoline phase-out. In order to maintain the “anti-knock” quality and octane ratings of gasoline in the absence of lead, the portion of aromatic hydrocarbons blended into gasoline through the refining process was increased. As discussed above, aromatic hydrocarbons have a lower ratio of hydrogen to C than other hydrocarbons typically found in gasoline, and therefore increase fuel density.

The trend in gasoline density was reversed beginning in 1996 with the development of fuel additives that raised oxygen content. In 1995, a requirement for reformulated gasoline in non-attainment areas implemented under the Clean Air Act Amendments further changed the composition of gasoline consumed in the United States. Through 2005, methyl tertiary butyl ether (MTBE), ethanol, ethyl tertiary butyl ether (ETBE), and tertiary amyl methyl ether (TAME) were added to reformulated and sometimes to conventional gasoline to boost its oxygen content, reduce its toxics impacts and increase its octane. The increased oxygen reduced the emissions of carbon monoxide and unburned hydrocarbons. These oxygen-rich blending components are also much lower in C than standard gasoline. The average gallon of reformulated gasoline consumed in 2005 contained over 10 percent MTBE and 0.6 percent TAME (by volume). The characteristics of reformulated fuel additives appear in Table A-50.

**Table A-50: Characteristics of Major Reformulated Fuel Additives**

| Additive       | Density (Degrees API) | Carbon Share (Percent) |
|----------------|-----------------------|------------------------|
| MTBE           | 58.6                  | 68.13                  |
| ETBE           | 58.5                  | 70.53                  |
| TAME           | 51.2                  | 70.53                  |
| DIPE           | 62.7                  | 70.53                  |
| Ethanol (100%) | 45.8                  | 52.14                  |

Source: EPA (2009b).

Since 2005, due to concerns about the potential environmental consequences of the use of MTBE in fuels, there has been a shift away from the addition of MTBE, TAME, ETBE, and DIPE and towards the use of ethanol as a fuel oxygenate.<sup>18</sup> Ethanol, also called ethyl alcohol, is an anhydrous alcohol with molecular formula C<sub>2</sub>H<sub>5</sub>OH. Ethanol has a lower C share than other oxygenates, approximately 52 percent compared to about 70 percent for MTBE and TAME. The density of ethanol was calculated by fitting density data at 10 degree intervals to a polynomial of order two and then using

<sup>18</sup> The annual motor gasoline carbon contents that are applied for this Inventory do not include the carbon contributed by the ethanol contained in reformulated fuels. Ethanol is a biofuel, and net carbon fluxes from changes in biogenic carbon reservoirs in croplands are accounted for in the estimates for Land Use, Land-Use Change and Forestry.

the fit to interpolate the value of the density at 15 degrees Celsius. A common fuel mixture of 10 percent denatured ethanol (denatured by 2 percent hydrocarbons) and 90 percent gasoline, known as E10, is widely used in the United States and does not require any modification to vehicle engines or fuel systems. The average gallon of reformulated alcohol blend gasoline in 2008 contained 8.6 percent ethanol (by volume). As of 2010, ten states require the use of ethanol-blended fuel.<sup>19</sup> Ethanol blends up to E85 (85 percent ethanol, 15 percent gasoline) are in use in the United States but can only be used in specially designed vehicles called flexible fuel vehicles (FFVs). Most ethanol fuel in the United States is produced using corn as feedstock,<sup>20</sup> although production pathways utilizing agricultural waste, woody biomass and other resources are in development.

#### Methodology

##### *Step 1. Disaggregate U.S. gasoline consumption by grade and type*

Separate monthly data for U.S. sales to end users of finished gasoline by product grade and season for both standard gasoline and reformulated gasoline were obtained from the EIA.

##### *Step 2. Develop carbon content coefficients for each grade and type*

Annual C content coefficients for each gasoline grade, type, and season are derived from four parameters for each constituent of the finished gasoline blend: the volumetric share of each constituent,<sup>21</sup> the density of the constituent, share of the constituent<sup>22</sup> that is C; and the energy content of a gallon of the relevant formulation of gasoline. The percent by mass of each constituent of each gasoline type was calculated using percent by volume data from the National Institute for Petroleum and Energy Research (NIPER) and the density of each constituent. The ether additives listed in Table A-50 are accounted for in both reformulated fuels and conventional fuels, to the extent that they were present in the fuel. From 2006 onward, reformulated fuel mass percentages are calculated from their constituents, net of the share provided by ethanol. C content coefficients were then derived from the calculated percent by mass values by weighting the C share of each constituent by its contribution to the total mass of the finished motor gasoline product.

##### *Step 3. Weight overall gasoline carbon content coefficient for consumption of each grade and type*

The C content for each grade, type, and season of fuel is multiplied by the share of annual consumption represented by the grade and fuel type during the relevant time period. Individual coefficients are then summed and totaled to yield an overall C content coefficient for each year.

#### Data Sources

Data for the density of motor gasoline were derived from NIPER (1990 through 2009). Data on the characteristics of reformulated gasoline, including C share, were also taken from NIPER (1990 through 2009).

Standard heat contents for motor gasoline of 5.253 MMBtu per barrel conventional gasoline and 5.150 MMBtu per barrel reformulated gasoline<sup>23</sup> were adopted from EIA (2009a).

#### Uncertainty

The uncertainty underlying the C content coefficients for motor gasoline has three underlying sources: (1) the uncertainty in the averages published by NIPER, (2) uncertainty in the C shares assumed in the EPA's analysis to be representative of the constituent hydrocarbon classes within gasoline (aromatics, olefins and saturates), and (3) uncertainty in the heat contents applied.

A variable number of samples are used each year to determine the average percent by volume share of each hydrocarbon within each grade, season and formulation of gasoline that are obtained from NIPER. The total number of samples analyzed for each seasonal NIPER report varies from approximately 730 to over 1,800 samples over the period from 1990 through 2014. The number of samples analyzed that underlie the calculation of the average make-up of each seasonal formulation and grade varies from approximately 50 to over 400, with the greatest number of samples each season

<sup>19</sup> Ethanol.org. Available at <<http://www.ethanol.org/index.php?id=79&parentid=26>>. Retrieved 2-19-2010.

<sup>20</sup> "Ethanol Market Penetration." Alternative Fuels and Advanced Vehicles Data Center, U.S. DOE. Available at <<http://www.afdc.energy.gov/afdc/ethanol/market.html>>. Retrieved 2-19-2010.

<sup>21</sup> Calculations account for the properties of the individual constituents of gasoline, including, as applicable to the fuel grade and type: aromatics (excluding benzene), olefins, benzene, saturates, MTBE, TAME, ETBE, DIPE and ethanol.

<sup>22</sup> Saturates are assumed to be octane and aromatics are assumed to be toluene.

<sup>23</sup> The reformulated gasoline heat content is applied to both reformulated blends containing ethers and those containing ethanol.

being of conventional, regular or premium gasoline. Further, not all sample data submitted to NIPER contains data for each of the properties, such that the number of samples underlying each constituent average value for each season, grade and formulation may be variable within the single gasoline type (e.g., of the 1,073 samples for which some data was obtained for gasoline sold in Winter 1995 through 1996, benzene content was provided for all samples, while olefin, aromatic and saturate content was provided for just 736 of those samples).

The distribution of sample origin collected for the NIPER report and the calculation of national averages are not reflective of sales volumes. The publication of simple, rather than sales-weighted averages to represent national average values increases the uncertainty in their application to the calculation of carbon content factors for the purposes of this Inventory. Further, data for each sample is submitted voluntarily, which may also affect their representativeness.

Additionally, because the simple average constituent shares are calculated based upon data that have been renormalized to account for the share of ethers and alcohols, total average volume shares may not equal 100 percent.

The simple average for each hydrocarbon constituent is contained within a range of values that are as wide as -63.0/+74.5 percent of the mean across the Winter 2007 through 2008 and -51.3/+49.6 percent across the Summer 2008 samples of conventional, regular grade gasoline. However, these wide ranges exist for benzene, which generally accounts for only 1 percent, by volume, of each gallon. In contrast, saturates, the class of hydrocarbon that contribute the largest share, by volume, ranges only -6.5/+6.4 percent for the same set of winter samples and -8.8/+15.7 percent for the summer samples.

Secondly, EPA's calculation of C content factors for each gasoline type includes the following assumptions: for the purposes of assigning a carbon share to each compound in the blend, aromatic content (other than benzene) is assumed to be toluene and saturated hydrocarbons are assumed to be octane. All olefins have the same carbon share because they all have a molecular formula in the form  $C_nH_{2n}$ , so the C share applied to the olefin portion of the total gasoline blend does not increase the level of uncertainty in the calculation. These assumptions are based upon the use of octane and octane isomers as the primary saturates and toluene as the primary non-benzene aromatic in U.S. motor gasoline blends. The octane rating of a particular blend is based upon the equivalent iso-octane to heptane ratio, which is achieved through significant octane content relative to the other saturates. Aside from benzene, U.S. gasolines will include toluene as a major aromatic component, so toluene may be assumed a reasonable representative of total non-benzene aromatic content (EPA 2009a).

For each hydrocarbon category, the assumed C content lies within a range of possible values for all such hydrocarbons. Among saturated hydrocarbons, the C share of octane (84.12 percent) is at the high end of the range while ethane is represents the low end of the range (79.89 percent C). Total saturates constitute from 40 to 95 percent by volume of a given gasoline blend. For aromatics, toluene (91.25 percent C) lies in the middle of the possible range. This range is bounded by cumene (89.94 percent C) and naphthalene (93.71 percent C). Total aromatics may make up between 3 and 50 percent by volume of any given gasoline blend. The range of these potential values contributes to the uncertainty surrounding the final calculated C factors.

However, as demonstrated above in Figure A-3, the amount of variation in C content of gasoline is restricted by the compounds in the fuel to  $\pm 4$  percent. Further, despite variation in sampling survey response, sample size and annually variable fuel formulation requirements, the observed variation in the annual weighted motor gasoline coefficients estimated for this Inventory is  $\pm 0.8$  percent over 1990 through 2014.

The third primary contributor to uncertainty is the assumed heat content. The heat contents are industry standards established many years ago. The heat contents are standard conversion factors used by EIA to convert volumetric energy data to energy units. Because the heat contents of fuels change over time, without necessarily and directly altering their volume, the conversion of known volumetric data to energy units may introduce bias. Thus, a more precise approach to estimating emissions factors would be to calculate C content per unit of volume, rather than per unit of energy. Adopting this approach, however, makes it difficult to compare U.S. C content coefficients with those of other nations.

The changes in density of motor gasoline over the last decade suggest that the heat content of the fuels is also changing. However, that change within any season grade has been less than 1 percent over the decade. Of greater concern is the use of a standardized heat content across grades that show a variation in density of  $\pm 1.5$  percent from the mean for conventional gasoline and  $\pm 1.0$  percent for reformulated fuels.

#### *Jet Fuel*

Jet fuel is a refined petroleum product used in jet aircraft engines. There are two classes of jet fuel used in the United States: "naphtha-based" jet fuels and "kerosene-based" jet fuels. In 1989, 13 percent of U.S. consumption was naphtha-based fuel, with the remainder kerosene-based jet fuel. In 1993, the U.S. Department of Defense began a conversion

from naphtha-based JP-4 jet fuel to kerosene-based jet fuel, because of the possibility of increased demand for reformulated motor gasoline limiting refinery production of naphtha-based jet fuel. By 1996, naphtha-based jet fuel represented less than one-half of one percent of all jet fuel consumption. The C content coefficient for jet fuel used in this report prior to 1996 represents a consumption-weighted combination of the naphtha-based and kerosene-based coefficients. From 1996 to 2014, only the kerosene-based portion of total consumption is considered significant.

#### Methodology

##### *Step 1. Estimate the carbon content for naphtha-based jet fuels*

Because naphtha-based jet fuels are used on a limited basis in the United States, sample data on its characteristics are limited. The density of naphtha-based jet fuel (49 degrees) was estimated as the central point of the acceptable API gravity range published by ASTM. The heat content of the fuel was assumed to be 5.355 MMBtu per barrel based on EIA industry standards. The C fraction was derived from an estimated hydrogen content of 14.1 percent (Martel and Angello 1977), and an estimated content of sulfur and other non-hydrocarbons of 0.1 percent.

##### *Step 2. Estimate the carbon content for kerosene-based jet fuels*

The density of kerosene-based jet fuels was estimated at 42 degrees API and the carbon share at 86.3 percent. The density estimate was based on 38 fuel samples examined by NIPER. Carbon share was estimated on the basis of a hydrogen content of 13.6 percent found in fuel samples taken in 1959 and reported by Martel and Angello, and on an assumed sulfur content of 0.1 percent. The EIA's standard heat content of 5.670 MMBtu per barrel was adopted for kerosene-based jet fuel.

##### *Step 3. Weight the overall jet fuel carbon content coefficient for consumption of each type of fuel (1990-1995 only)*

For years 1990 through 1995, the C content for each jet fuel type (naphtha-based, kerosene-based) is multiplied by the share of overall consumption of that fuel type, as reported by EIA (2009a). Individual coefficients are then summed and totaled to yield an overall C content coefficient. Only the kerosene-based C coefficient is reflected in the overall jet fuel coefficient for 1996 through 2014.

#### Data Sources

Data on the C content of naphtha-based jet fuel was taken from C.R. Martel and L.C. Angello (1977). Data on the density of naphtha-based jet fuel was taken from ASTM (1985). Standard heat contents for kerosene and naphtha-based jet fuels were adopted from EIA (2009a). Data on the C content of kerosene-based jet fuel is based on C.R. Martel and L.C. Angello (1977) and the density is derived from NIPER (1993).

#### Uncertainty

Variability in jet fuel is relatively small with the average C share of kerosene-based jet fuel varying by less than  $\pm 1$  percent and the density varying by  $\pm 1$  percent. This is because the ratio of fuel mass to useful energy must be tightly bounded to maximize safety and range. There is more uncertainty associated with the density and C share of naphtha-based jet fuel because sample data were unavailable and default values were used. This uncertainty has only a small impact on the overall uncertainty of the C content coefficient for jet fuels, however, because naphtha-based jet fuel represents a small and declining share of total jet fuel consumption in the United States and is treated as negligible when calculating C content factors for 1996 onward.

#### *Distillate Fuel*

Distillate fuel is a general classification for diesel fuels and fuel oils. Products known as No. 1, No. 2, and No. 4 diesel fuel are used in on-highway diesel engines, such as those in trucks and automobiles, as well as off-highway engines, such as those in railroad locomotives and agricultural machinery. No. 1, No. 2, and No. 4 fuel oils are also used for space heating and electric power generation.

#### Methodology

For this Inventory, separate C coefficients have been estimated for each of the three distillates, although the level of aggregation of U.S. energy statistics requires that a single coefficient is used to represent all three grades in inventory calculations. In past Inventories, the emission coefficient was only determined for distillate No. 2. Distillate No. 2 remains the representative grade applied to the distillate class for calculation purposes. Coefficients developed for No. 1 and No. 4 distillate are provided for informational purposes. The C share each distillate is drawn from *Perry's Chemical Engineers' Handbook, 8<sup>th</sup> Ed.* (Green & Perry 2008). Each C share was combined with individual heat contents of 5.822, 5.809 and

6.135 MMBtu per barrel, respectively for distillates No. 1, No. 2, and No. 4, and densities of 35.3, 35.8, and 23.2 degrees API to calculate C coefficients for each distillate type.

#### Data Sources

Densities for distillate No. 1 and No. 2 were derived from *Alliance of Automobile Manufacturers, Diesel Survey – Winter 2008* (AAM 2009). Densities are based on four, and 144 samples, respectively. The density of distillate fuel oil No. 4 is taken from *Perry's Chemical Engineer's Handbook, 8<sup>th</sup> Ed.* (Green & Perry 2008), Table 24-6.

Heat contents are adopted from EPA (2009b). And carbon shares for each distillate are from *Perry's Chemical Engineers' Handbook* (Green & Perry 2008), Table 24-6.

#### Uncertainty

The primary source of uncertainty for the estimated C content of distillate fuel is the selection of No. 2 distillate as the typical distillate fuel oil or diesel fuel. No. 2 fuel oil is generally consumed for home heating. No. 1 distillate is generally less dense and if it is consumed in large portions for mobile sources, the application of the C content estimated for No. 2 for this report is likely to be too high when applied to both No. 1 and No. 2 distillates. The opposite is true of the application of a coefficient based upon the properties of No. 2 to the consumption of No. 4 distillate, which is of a significantly higher density and thus, has a higher C coefficient despite its lower C share. The overall effect on uncertainty from applying a single factor will depend on the relative annual consumption of each distillate.

The densities applied to the calculation of each carbon factor are an underlying source of uncertainty. While the density of No. 1 distillate is based upon just four samples, the factor applied to all distillates in the Inventory estimates (that for No. 2 oil) is based on a much larger sample size (144). Given the range of densities for these three distillate fuel classes (0.1342 to 0.1452 MT/bbl at 60°F), the uncertainty associated with the assumed density of distillate fuels is predominately a result of the use of No. 2 to represent all distillate consumption. There is also a small amount of uncertainty in the No. 2 distillate density itself. This is due to the possible variation across seasonal diesel formulations and fuel grades and between stationary and transport applications within the No. 2 distillate classification. The range of the density of the samples of No. 2 diesel (regular grade, 15 ppm sulfur) is  $\pm 2.5$  percent from the mean, while the range in density across the small sample set of No. 1 diesel is -2.1 to +1.6 percent of the mean. Samples from AAM (2009) of Premium No. 2 diesel (n=5) and higher sulfur (500 ppm S) regular diesel (n=2), which are also consumed in the United States, each have nominally higher average densities (+1.3 percent and +0.6 percent, respectively) than do the low-sulfur regular diesel samples that underlie the density applied in this Inventory.

The use of the 144 AAM samples to define the density of No. 2 distillate (and those four samples used to define that of No. 1 distillate) may introduce additional uncertainty because the samples were collected from just one season of on-road fuel production (Winter 2008). Despite the limited sample frame, the average No. 2 density calculated from the samples is applied to the calculation of a uniform C coefficient applicable for all years of the Inventory and for all types of distillate consumption. The ASTM standards for each grade of diesel fuel oil do not include a required range in which the density must lie, and the density (as well as heat content and carbon share) may vary according to the additives in each seasonal blend and the sulfur content of each sub-grade.

However, previous studies also show relatively low variation in density across samples of No. 2 and across all distillates, supporting the application of a single No. 2 density to all U.S. distillate consumption. The average density calculated from samples analyzed by the EIA in 1994 (n=7) differs only very slightly from the value applied for the purposes of this Inventory (-0.12 percent for No. 2 distillate). Further, the difference between the mean density applied to this Inventory (No. 2 only) and that calculated from EIA samples of all distillates, regardless of grade, is also near zero (-0.06 percent, based on n=14, of distillates No. 1, No. 2 and No. 4 combined).

A C share of 87.30 percent is applied to No. 2 distillate, while No. 1 and No. 4 have C shares estimated at 86.40 and 86.47 percent, respectively. Again, the application of parameters specific to No. 2 to the consumption of all three distillates contributes to an increased level of uncertainty in the overall coefficient and emissions estimate and its broad application. For comparison, four No. 1 fuel oil samples obtained by EIA (1994) contained an average of 86.19 percent C, while seven samples No. 2 fuel oil from the same EIA analysis showed an average of 86.60 percent C. Additionally, three samples of No. 4 distillate indicate an average C share of 85.81 percent. The range of C share observed across the seven No. 2 samples is 86.1 to 87.5 percent, and across all samples (all three grades, n=14) the range is 85.3 to 87.5 percent C. There also exists an uncertainty of  $\pm 1$  percent in the share of C in No. 2 based on the limited sample size.

### *Residual Fuel*

Residual fuel is a general classification for the heavier oils, known as No. 5 and No. 6 fuel oils, that remain after the distillate fuel oils and lighter hydrocarbons are distilled away in refinery operations. Residual fuel conforms to ASTM Specifications D 396 and D 975 and Federal Specification VV-F-815C. No. 5, a residual fuel oil of medium viscosity, is also known as Navy Special and is defined in Military Specification MIL-F-859E, including Amendment 2 (NATO Symbol F-770). It is used in steam-powered vessels in government service and inshore power plants. No. 6 fuel oil includes Bunker C fuel oil and is used for the production of electric power, space heating, vessel bunkering, and various industrial purposes.

In the United States, electric utilities purchase about one-third of the residual oil consumed. A somewhat larger share is used for vessel bunkering, and the balance is used in the commercial and industrial sectors. The residual oil (defined as No. 6 fuel oil) consumed by electric utilities has an energy content of 6.287 MMBtu per barrel (EIA 2008a) and an average sulfur content of 1 percent (EIA 2001). This implies a density of about 17 degrees API.

#### Methodology

Because U.S. energy consumption statistics are available only as an aggregate of No. 5 and No. 6 residual oil, a single coefficient must be used to represent the full residual fuel category. As in earlier editions of this report, residual fuel oil has been defined as No. 6 fuel oil, due to the majority of residual consumed in the United States being No. 6. However, for this report, a separate coefficient for fuel oil No. 5 has also been developed for informational purposes. Densities of 33.0 and 15.5 degrees API were adopted when developing the C content coefficients for Nos. 5 and 6, respectively (Wauquier, J.-P., ed. 1995; Green & Perry, ed. 2008).

The estimated C share of fuel oil No. 5 is 85.67 percent, based on an average of 12 ultimate analyses of samples of fuel oil (EIA 1994). An average share of C in No. 6 residual oil of 84.67 percent by mass was used, based on Perry's, 8<sup>th</sup> Ed. (Green & Perry 2008).

#### Data Sources

Data on the C share and density of residual fuel oil No. 6 were obtained from Green & Perry, ed. (2008). Data on the C share of fuel oil No. 5 was adopted from EIA (1994), and the density of No. 5 was obtained from Wauquier, J.-P., ed. (1995). Heat contents for both No. 5 and No. 6 fuel oil are adopted from EPA (2009b).

#### Uncertainty

Beyond the application of a C factor based upon No. 6 oil to all residual oil consumption, the largest source of uncertainty in estimating the C content of residual fuel centers on the estimates of density. Fuel oils are likely to differ depending on the application of the fuel (i.e., power generation or as a marine vessel fuel). Slight differences between the density of residual fuel used by utilities and that used in mobile applications are likely attributable to non-sulfur impurities, which reduce the energy content of the fuel, but do not greatly affect the density of the product. Impurities of several percent are commonly observed in residual oil. The extent of the presence of impurities has a greater effect on the uncertainty of C share estimation than it does on density. This is because these impurities do provide some Btu content to the fuel, but they are absent of carbon. Fuel oils with significant sulfur, nitrogen and heavy metals contents would have a different total carbon share than a fuel oil that is closer to pure hydrocarbon. This contributes to the uncertainty of the estimation of an average C share and C coefficient for these varied fuels.

The 12 samples of residual oil (EIA 1994) cover a density range from 4.3 percent below to 8.2 percent above the mean density. The observed range of C share in these samples is -2.5 to +1.8 percent of the mean. Overall, the uncertainty associated with the C content of residual fuel is probably  $\pm 1$  percent.

#### *Liquefied Petroleum Gases (LPG)*

EIA identifies four categories of paraffinic hydrocarbons as LPG: ethane, propane, isobutane, and n-butane. Because each of these compounds is a pure paraffinic hydrocarbon, their C shares are easily derived by taking into account the atomic weight of C (12.01) and the atomic weight of hydrogen (1.01). Thus, for example, the C share of propane, C<sub>3</sub>H<sub>8</sub>, is 81.71 percent. The densities and heat contents of the compounds are also well known, allowing C content coefficients to be calculated directly. Table A-51 summarizes the physical characteristic of LPG.

**Table A-51: Physical Characteristics of Liquefied Petroleum Gases**

| Compound  | Chemical Formula               | Density (Barrels Per Metric Ton) | Carbon Content (Percent) | Energy Content (MMBtu/Barrel) | Carbon Content Coefficient (MMT C/QBtu) |
|-----------|--------------------------------|----------------------------------|--------------------------|-------------------------------|---|
| Ethane    | C <sub>2</sub> H <sub>6</sub>  | 11.55                            | 79.89                    | 3.082                         | 17.16                                   |
| Propane   | C <sub>3</sub> H <sub>8</sub>  | 12.76                            | 81.71                    | 3.836                         | 16.76                                   |
| Isobutane | C <sub>4</sub> H <sub>10</sub> | 11.42                            | 82.66                    | 3.974                         | 17.77                                   |
| n-butane  | C <sub>4</sub> H <sub>10</sub> | 10.98                            | 82.66                    | 4.326                         | 17.75                                   |

Source: Densities – CRC Handbook of Chemistry and Physics (2008/09); Carbon Contents – derived from the atomic weights of the elements; Energy Contents – EPA (2009b). All values are for the compound in liquid form. The density and energy content of ethane are for refrigerated ethane (-89 degrees C). Values for n-butane are for pressurized butane (-25 degrees C).

#### Methodology

##### *Step 1. Assign carbon content coefficients to each pure paraffinic compound*

Based on their known physical characteristics, a C content coefficient is assigned to each compound contained in the U.S. energy statistics category, Liquefied Petroleum Gases.

##### *Step 2. Weight individual LPG coefficients for share of fuel use consumption*

A C content coefficient for LPG used as fuel is developed based on the consumption mix of the individual compounds reported in U.S. energy statistics.

##### *Step 3. Weight individual LPG coefficients for share of non-fuel use consumption*

The mix of LPG consumed for non-fuel use differs significantly from the mix of LPG that is combusted. While the majority of LPG consumed for fuel use is propane, ethane is the largest component of LPG used for non-fuel applications. A C content coefficient for LPG used for non-fuel applications is developed based on the consumption mix of the individual compounds reported in U.S. energy statistics.

##### *Step 4. Weight the carbon content coefficients for fuel use and non-fuel use by their respective shares of consumption*

The changing shares of LPG fuel use and non-fuel use consumption appear below in Table A-52.

#### Data Sources

Data on C share was derived via calculations based on atomic weights of each element of the four individual compounds densities are from the CRC Handbook of Chemistry and Physics, 89<sup>th</sup> Edition. The energy content of each LPG is from the EPA (2009b). LPG consumption was based on data obtained from API (1990 through 2014) and EIA (2009b). Non-fuel use of LPG was obtained from API (1990 through 2008).

#### Uncertainty

Because LPG consists of pure paraffinic compounds whose density, heat content and C share are physical constants, there is limited uncertainty associated with the C content coefficient for this petroleum product. Any uncertainty is associated with the collection of data tabulating fuel- and non-fuel consumption in U.S. energy statistics. This uncertainty is likely less than ±3 percent.

**Table A-52: Consumption and Carbon Content Coefficients of Liquefied Petroleum Gases, 1990-2014**

|                                  | 1990        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|----------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>Energy Consumption (QBtu)</b> |             |             |             |             |             |             |             |             |             |             |             |             |             |             |             |             |
| <b>Fuel Use</b>                  | <b>0.88</b> | <b>1.31</b> | <b>1.16</b> | <b>1.25</b> | <b>1.22</b> | <b>1.26</b> | <b>1.21</b> | <b>1.19</b> | <b>1.20</b> | <b>1.13</b> | <b>1.13</b> | <b>1.16</b> | <b>1.16</b> | <b>1.16</b> | <b>1.16</b> | <b>1.16</b> |
| Ethane                           | 0.04        | 0.10        | 0.06        | 0.06        | 0.06        | 0.06        | 0.06        | 0.06        | 0.07        | 0.06        | 0.07        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        |
| Propane                          | 0.77        | 1.07        | 1.00        | 1.10        | 1.07        | 1.12        | 1.08        | 1.07        | 1.09        | 1.02        | 1.02        | 1.02        | 1.02        | 1.02        | 1.02        | 1.02        |
| Butane                           | 0.06        | 0.07        | 0.06        | 0.05        | 0.06        | 0.06        | 0.05        | 0.05        | 0.05        | 0.05        | 0.03        | 0.05        | 0.05        | 0.05        | 0.05        | 0.05        |
| Isobutane                        | 0.01        | 0.06        | 0.04        | 0.04        | 0.03        | 0.01        | 0.01        | 0.01        | 0.00        | 0.00        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        |
| <b>Non-Fuel Use</b>              | <b>1.35</b> | <b>1.90</b> | <b>1.77</b> | <b>1.85</b> | <b>1.75</b> | <b>1.80</b> | <b>1.70</b> | <b>1.74</b> | <b>1.78</b> | <b>1.67</b> | <b>1.80</b> | <b>1.96</b> | <b>1.96</b> | <b>1.96</b> | <b>1.96</b> | <b>1.96</b> |
| Ethane                           | 0.71        | 1.04        | 0.96        | 1.00        | 0.92        | 0.97        | 0.91        | 0.98        | 1.03        | 0.95        | 1.12        | 1.22        | 1.22        | 1.22        | 1.22        | 1.22        |
| Propane                          | 0.51        | 0.65        | 0.59        | 0.64        | 0.63        | 0.66        | 0.63        | 0.63        | 0.64        | 0.60        | 0.60        | 0.60        | 0.60        | 0.60        | 0.60        | 0.60        |
| Butane                           | 0.11        | 0.11        | 0.13        | 0.12        | 0.13        | 0.13        | 0.12        | 0.12        | 0.11        | 0.12        | 0.08        | 0.12        | 0.12        | 0.12        | 0.12        | 0.12        |

|                                    |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
|------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Isobutane                          | 0.02  | 0.09  | 0.09  | 0.08  | 0.07  | 0.03  | 0.03  | 0.02  | 0.01  | 0.00  | 0.01  | 0.03  | 0.03  | 0.03  | 0.03  | 0.03  |
| <b>Carbon Content (MMT C/QBtu)</b> |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Fuel Use                           | 16.86 | 16.89 | 16.87 | 16.85 | 16.86 | 16.84 | 16.84 | 16.83 | 16.82 | 16.83 | 16.83 | 16.83 | 16.83 | 16.83 | 16.83 | 16.83 |
| Non-Fuel Use                       | 17.06 | 17.09 | 17.10 | 17.09 | 17.09 | 17.07 | 17.06 | 17.06 | 17.05 | 17.06 | 17.06 | 17.06 | 17.06 | 17.06 | 17.06 | 17.06 |

Sources: Fuel use of LPG based on data from EIA (2009b) and API (1990 through 2007). Non-fuel use of LPG from API (1990 through 2008). Volumes converted using the energy contents provided in Table A-51. C contents from EPA (2010).

### Aviation Gasoline

Aviation gasoline is used in piston-powered airplane engines. It is a complex mixture of relatively volatile hydrocarbons with or without small quantities of additives, blended to form a fuel suitable for use in aviation reciprocating engines. Fuel specifications are provided in ASTM Specification D910 and Military Specification MIL-G-5572. Aviation gas is a relatively minor contributor to greenhouse gas emissions compared to other petroleum products, representing approximately 0.1 percent of all consumption.

The ASTM standards for boiling and freezing points in aviation gasoline effectively limit the aromatics content to a maximum of 25 percent (ASTM D910). Because weight is critical in the operation of an airplane, aviation gas must have as many Btu per pound (implying a lower density) as possible, given other requirements of piston engines such as high anti-knock quality.

### Methodology

A C content coefficient for aviation gasoline was calculated on the basis of the EIA standard heat content of 5.048 MMBtu per barrel. This implies a density of approximately 69 degrees API gravity or 5.884 pounds per gallon, based on the relationship between heat content and density of petroleum liquids, as described in *Thermal Properties of Petroleum Products* (DOC 1929). To estimate the share of C in the fuel, it was assumed that aviation gasoline is 87.5 percent iso-octane, 9.0 percent toluene, and 3.5 percent xylene. The maximum allowable sulfur content in aviation gasoline is 0.05 percent, and the maximum allowable lead content is 0.1 percent. These amounts were judged negligible and excluded for the purposes of this analysis. This yielded a C share of 85.00 percent and a C content coefficient of 18.86 MMT C/QBtu.

### Data Sources

Data sources include ASTM (1985). A standard heat content for aviation gas was adopted from EIA (2009a).

### Uncertainty

The relationship used to calculate density from heat content has an accuracy of five percent at 1 atm. The uncertainty associated with the C content coefficient for aviation gasoline is larger than that for other liquid petroleum products examined because no ultimate analyses of samples are available. Given the requirements for safe operation of piston-powered aircraft the composition of aviation gas is well bounded and the uncertainty of the C content coefficient is likely to be  $\pm 5$  percent.

### Still Gas

Still gas, or refinery gas, is composed of light hydrocarbon gases that are released as petroleum is processed in a refinery. The composition of still gas is highly variable, depending primarily on the nature of the refining process and secondarily on the composition of the product being processed. Petroleum refineries produce still gas from many different processes. Still gas can be used as a fuel or feedstock within the refinery, sold as a petrochemical feedstock, or purified and sold as pipeline-quality natural gas. For the purposes of this Inventory, the coefficient derived here is only applied to still gas that is consumed as a fuel. In general, still gas tends to include large amounts of free hydrogen and methane, as well as smaller amounts of heavier hydrocarbons. Because different refinery operations result in different gaseous by-products, it is difficult to determine what represents typical still gas.

### Methodology

The properties of still gas used to calculate the carbon content are taken from the literature. The carbon share of still gas was calculated from its net calorific value and carbon content from IPCC (2006). This calculation yields a carbon share of 77.7 percent. The density of still gas was estimated to be 0.1405 metric tons per barrel based on its heat content (EIA 2008a) and the relationship between heat content and density that is described by the U.S. Department of Commerce, Bureau of Standards (DOC 1929).

### Data Sources

The carbon share of still gas is calculated from data provided by IPCC (2006). Density is estimated at 0.1405 metric tons per barrel, approximately 28.3 degrees API, based on the heat content of 6.00 MMBtu/barrel of still gas from EIA (2009a).

### Uncertainty

The EIA obtained data on four samples of still gas. Table A-53 below shows the composition of those samples.

**Table A-53: Composition, Energy Content, and Carbon Content Coefficient for Four Samples of Still Gas**

| Sample | Hydrogen (%) | Methane (%) | Ethane (%) | Propane (%) | Btu Per Cubic Foot | Carbon Content (MMT C/QBtu) |
|--------|--------------|-------------|------------|-------------|--------------------|-----------------------------|
| One    | 12.7         | 28.1        | 17.1       | 11.9        | 1,388              | 17.51                       |
| Two    | 34.7         | 20.5        | 20.5       | 6.7         | 1,143              | 14.33                       |
| Three  | 72.0         | 12.8        | 10.3       | 3.8         | 672                | 10.23                       |
| Four   | 17.0         | 31.0        | 16.2       | 2.4         | 1,100              | 15.99                       |

Sources: EIA (2008b).

Because the composition of still gas is highly heterogeneous, the C content coefficient for this product is highly uncertain. Gas streams with a large, free-hydrogen content are likely to be used as refinery or chemical feedstocks. Therefore, the sample cited above with the very high H content of 72 percent (and the lowest calculated C content) is less likely to be representative of the still gas streams to which the calculated coefficient is applied. The C content coefficient used for this report is probably at the high end of the plausible range given that it is higher than the greatest sample-based C content in Table A-53.

### *Asphalt*

Asphalt is used to pave roads. Because most of its C is retained in those roads, it is a small source of carbon dioxide emissions. It is derived from a class of hydrocarbons called "asphaltenes," which are abundant in some crude oils but not in others. Asphaltenes have oxygen and nitrogen atoms bound into their molecular structure, so that they tend to have lower C contents than do other hydrocarbons.

### Methodology

Ultimate analyses of twelve samples of asphalts showed an average C content of 83.47 percent. The EIA standard Btu content for asphalt of 6.636 MMBtu per barrel was assumed. The ASTM petroleum measurement tables show a density of 5.6 degrees API or 8.605 pounds per gallon for asphalt. Together, these variables generate C content coefficient of 20.55 MMT C/QBtu.

### Data Sources

A standard heat content for asphalt was adopted from EIA (2009a). The density of asphalt was determined by the ASTM (1985). C share is adopted from analyses in EIA (2008b).

### Uncertainty

The share of C in asphalt ranges from 79 to 88 percent by weight. Also present in the mixture are hydrogen and sulfur, with shares by weight ranging from seven to 13 percent for hydrogen, and from trace levels to eight percent for sulfur. Because C share and total heat content in asphalts do vary systematically, the overall C content coefficient is likely to be accurate to  $\pm 5$  percent.

### *Lubricants*

Lubricants are substances used to reduce friction between bearing surfaces, or incorporated into processing materials used in the manufacture of other products, or used as carriers of other materials. Petroleum lubricants may be produced either from distillates or residues. Lubricants include all grades of lubricating oils, from spindle oil to cylinder oil to those used in greases. Lubricant consumption is dominated by motor oil for automobiles, but there is a large range of product compositions and end uses within this category.

### Methodology

The ASTM Petroleum Measurement tables give the density of lubricants at 25.6 degrees API, or 0.1428 metric tons per barrel. Ultimate analysis of a single sample of motor oil yielded a C content of 85.80 percent. A standard heat

content of 6.065 MMBtu per barrel was adopted from EIA. These factors produce a C content coefficient of 20.20 MMT C/QBtu.

#### Data Sources

A standard heat content was adopted from the EIA (2009a). The carbon content of lubricants is adopted from ultimate analysis of one sample of motor oil (EPA 2009a). The density of lubricating oils was determined by ASTM (1985).

#### Uncertainty

Uncertainty in the estimated C content coefficient for lubricants is driven by the large range of product compositions and end uses in this category combined with an inability to establish the shares of the various products captured under this category in U.S. energy statistics. Because lubricants may be produced from either the distillate or residual fractions during refineries, the possible C content coefficients range from 19.89 MMT C/QBtu to 21.48 MMT C/QBtu or an uncertainty band from -1.5 percent to +1.4 percent of the estimated value.

#### *Petrochemical Feedstocks*

U.S. energy statistics distinguish between two different kinds of petrochemical feedstocks: those with a boiling temperature below 400 degrees Fahrenheit, generally called “naphtha,” and those with a boiling temperature 401 degrees Fahrenheit and above, referred to as “other oils” for the purposes of this Inventory.

#### Methodology

The C content of these petrochemical feedstocks are estimated independently according to the following steps.

##### *Step 1. Estimate the carbon content coefficient for naphtha*

Because reformed naphtha is used to make motor gasoline (hydrogen is released to raise aromatics content and octane rating), “straight-run” naphtha is assumed to be used as a petrochemical feedstock. Ultimate analyses of five samples of naphtha were examined and showed an average C share of 84.11 percent. A density of 62.4 degrees API gravity was taken from the Handbook of Petroleum Refining Processes, 3<sup>rd</sup> ed. The standard EIA heat content of 5.248 MMBtu per barrel is used to estimate a C content coefficient of 18.55 MMT C/QBtu.

##### *Step 2. Estimate the carbon content coefficient for petrochemical feedstocks with a boiling temperature 400 degrees Fahrenheit and above (“other oils”)*

The boiling temperature of this product places it into the “middle distillate” fraction in the refining process, and EIA estimates that these petrochemical feedstocks have the same heat content as distillate fuel No. 2. Thus, the C content coefficient of 20.17 MMT C/QBtu used for distillate fuel No. 2 is also adopted for this portion of the petrochemical feedstocks category.

#### Data Sources

Naphthas: Data on the C content was taken from Unzelman (1992). Density is from Meyers (2004). A standard heat content for naphthas was adopted from EIA (2009a). Other oils: See Distillate Fuel, Distillate No.2.

#### Uncertainty

Petrochemical feedstocks are not so much distinguished on the basis of chemical composition as on the identity of the purchaser, who are presumed to be a chemical company or a petrochemical unit co-located on the refinery grounds. Naphthas are defined, for the purposes of U.S. energy statistics, as those naphtha products destined for use as a petrochemical feedstock. Because naphthas are also commonly used to produce motor gasoline, there exists a considerable degree of uncertainty about the exact composition of petrochemical feedstocks.

Different naphthas are distinguished by their density and by the share of paraffins, isoparaffins, olefins, naphthenes and aromatics contained in the oil. Naphtha from the same crude oil fraction may have vastly different properties depending on the source of the crude. Two different samples of Egyptian crude, for example, produced two straight run naphthas having naphthene and paraffin contents (percent volume) that differ by 18.1 and 17.5 percent, respectively (Matar and Hatch 2000).

Naphthas are typically used either as a petrochemical feedstock or a gasoline feedstock, with lighter paraffinic naphthas going to petrochemical production. Naphthas that are rich in aromatics and naphthenes tend to be reformed or blended into gasoline. Thus, the product category encompasses a range of possible fuel compositions, creating a range of possible C shares and densities. The uncertainty associated with the calculated C content of naphthas is primarily a function

of the uncertainty that underlies the average carbon share calculation, which is based on a limited number of samples. Two additional samples cited by the EIA (1994) have a range of 83.80 to 84.42 percent C.

The uncertainty of the C content for other oils is based upon the assumption of distillate oil No. 2 as a product representative of the ill-defined classification of “other oils,” and from the calculation of the C content of No. 2 itself (see “Distillate Fuels,” above). While No. 2 distillate is used as a proxy for “other oils” for the purposes of this Inventory’s carbon coefficient, important differences exist between these two petroleum products, contributing some uncertainty to the cross-application. Other oils are defined herein as those “oils with a boiling range equal to or greater than 401°F that are generally intended for use as a petrochemical feedstock and are not defined elsewhere.” For comparison, various material safety data sheets (MSDSs) published by producers of distillate No. 2 indicate a boiling range for this product of 320-700 degrees Fahrenheit. The relatively open definition of the classification “other oils” leaves room for potentially significant variation in the heating value, density and carbon share properties of each feedstock oil having a boiling point above 400 degrees Fahrenheit, creating a large band of uncertainty beyond that associated with the C factor for distillate No. 2.

#### *Kerosene*

A light petroleum distillate that is used in space heaters, cook stoves, and water heaters and is suitable for use as a light source when burned in wick-fed lamps, kerosene is drawn from the same petroleum fraction as jet fuel. Kerosene is generally comparable to No. 1 distillate oil.

#### Methodology

The average density and C share of kerosene are assumed to be the same as those for distillate No. 1 since the physical characteristics of the products are very similar. Thus, a density of 35.3 degrees API and average C share of 86.40 percent were applied to a standard heat content for distillate No. 1 of 5.825 MMBtu per barrel to yield a C content coefficient of 19.96 MMT C/QBtu.

#### Data Sources

A standard heat content for distillate No. 1 was adopted from EIA (2009a).

#### Uncertainty

Uncertainty in the estimated C content for kerosene is driven by the selection of distillate No. 1 as a proxy for kerosene. If kerosene is more like kerosene-based jet fuel, the true C content coefficient is likely to be some 1.3 percent lower. If kerosene is more aptly compared to No. 2 distillate oil, then the true C content coefficient is likely to be about 1.1 percent higher. While kerosene is a light petroleum distillate, like distillate No. 1, the two oil classes do have some variation in their properties. For example, the boiling range of kerosene is 250 to 550 degrees Fahrenheit, whereas No. 1 oils typically boil over a range from 350 to 615 degrees Fahrenheit. The properties of individual kerosenes will vary with their use and particular crude origin, as well. Both kerosene and fuel oil No. 1 are primarily composed of hydrocarbons having 9 to 16 carbon atoms per molecule. However, kerosene is a straight-run No. 1 fuel oil, additional cracking processes and additives contribute to the range of possible fuels that make up the broader distillate No. 1 oil category.

#### *Petroleum Coke*

Petroleum coke is the solid residue by-product of the extensive processing of crude oil. It is a coal-like solid, usually has a C content greater than 90 percent, and is used as a boiler fuel and industrial raw material.

#### Methodology

Ultimate analyses of two samples of petroleum coke showed an average C share of 92.28 percent. The ASTM standard density of 9.543 pounds per gallon was adopted and the EIA standard energy content of 6.024 MMBtu per barrel assumed. Together, these factors produced an estimated C content coefficient of 27.85 MMT C/QBtu.

#### Data Sources

C content was derived from two samples from Martin, S.W. (1960). The density of petroleum coke was taken from the ASTM (1985). A standard heat content for petroleum coke was adopted from EIA (2009a).

#### Uncertainty

The uncertainty associated with the estimated C content coefficient of petroleum coke can be traced to two factors: the use of only two samples to establish C contents and a standard heat content which may be too low. Together, these uncertainties are likely to bias the C content coefficient upwards by as much as 6 percent.

### *Special Naphtha*

Special naphtha is defined as a light petroleum product to be used for solvent applications, including commercial hexane and four classes of solvent: (1) Stoddard solvent, used in dry cleaning; (2) high flash point solvent, used as an industrial paint because of its slow evaporative characteristics; (3) odorless solvent, most often used for residential paints; and (4) high solvency mineral spirits, used for architectural finishes. These products differ in both density and C percentage, requiring the development of multiple coefficients.

#### Methodology

The method for estimating the C content coefficient of special naphtha includes three steps.

##### *Step 1. Estimate the carbon content coefficient for hexane*

Hexane is a pure paraffin containing 6 C atoms and 14 hydrogen atoms; thus, it is 83.63 percent C. Its density is 83.7 degrees API or 5.477 pounds per gallon and its derived C content coefficient is 21.40 MMT C/QBtu.

##### *Step 2. Estimate the carbon contents of non-hexane special naphthas*

The hydrocarbon compounds in special naphthas are assumed to be either paraffinic or aromatic (see discussion above). The portion of aromatics in odorless solvents is estimated at less than 1 percent, Stoddard and high flash point solvents contain 15 percent aromatics and high solvency mineral spirits contain 30 percent aromatics (Boldt and Hall 1977). These assumptions, when combined with the relevant densities, yield the C content factors contained in Table A-54, below.

**Table A-54: Characteristics of Non-hexane Special Naphthas**

| Special Naphtha  | Aromatic Content<br>(Percent) | Density<br>(Degrees API) | Carbon Share<br>(Percent Mass) | Carbon Content<br>(MMT C/QBtu) |
|------------------|-------------------------------|--------------------------|--------------------------------|--------------------------------|
| Odorless Solvent | 1                             | 55.0                     | 84.51                          | 19.41                          |
| Stoddard Solvent | 15                            | 47.9                     | 84.44                          | 20.11                          |
| High Flash Point | 15                            | 47.6                     | 84.70                          | 20.17                          |
| Mineral Spirits  | 30                            | 43.6                     | 85.83                          | 20.99                          |

Sources: EIA (2008b) and Boldt and Hall (1977).

##### *Step 3. Develop weighted carbon content coefficient based on consumption of each special naphtha*

EIA reports only a single consumption figure for special naphtha. The C contents of the five special naphthas are weighted according to the following formula: approximately 10 percent of all special naphtha consumed is hexane; the remaining 90 percent is assumed to be distributed evenly among the four other solvents. The resulting emissions coefficient for special naphthas is 19.74 MMT C/QBtu.

#### Data Sources

A standard heat content for special naphtha was adopted from EIA (2009a). Density and aromatic contents were adopted from Boldt and Hall (1977).

#### Uncertainty

The principal uncertainty associated with the estimated C content coefficient for special naphtha is the allocation of overall consumption across individual solvents. The overall uncertainty is bounded on the low end by the C content of odorless solvent and on the upper end by the C content of hexane. This implies an uncertainty band of -1.7 percent to +8.4 percent.

### *Petroleum Waxes*

The ASTM standards define petroleum wax as a product separated from petroleum that is solid or semi-solid at 77 degrees Fahrenheit (25 degrees Celsius). The two classes of petroleum wax are paraffin waxes and microcrystalline waxes. They differ in the number of C atoms and the type of hydrocarbon compounds. Microcrystalline waxes have longer C chains and more variation in their chemical bonds than paraffin waxes.

#### Methodology

The method for estimating the C content coefficient for petroleum waxes includes three steps.

### *Step 1. Estimate the carbon content of paraffin waxes*

For the purposes of this analysis, paraffin waxes are assumed to be composed of 100 percent paraffinic compounds with a chain of 25 C atoms. The resulting C share for paraffinic wax is 85.23 percent and the density is estimated at 45 degrees API or 6.684 pounds per gallon.

### *Step 2. Estimate the carbon content of microcrystalline waxes*

Microcrystalline waxes are assumed to consist of 50 percent paraffinic and 50 percent cycloparaffinic compounds with a chain of 40 C atoms, yielding a C share of 85.56 percent. The density of microcrystalline waxes is estimated at 36.7 degrees API, based on a sample of 10 microcrystalline waxes found in the *Petroleum Products Handbook*.

### *Step 3. Develop a carbon content coefficient for petroleum waxes by weighting the density and carbon content of paraffinic and microcrystalline waxes*

A weighted average density and C content was calculated for petroleum waxes, assuming that wax consumption is 80 percent paraffin wax and 20 percent microcrystalline wax. The weighted average C content is 85.30 percent, and the weighted average density is 6.75 pounds per gallon. EIA's standard heat content for waxes is 5.537 MMBtu per barrel. These inputs yield a C content coefficient for petroleum waxes of 19.80 MMT C/QBtu.

### Data Sources

Density of paraffin wax was taken from ASTM (1985). Density of microcrystalline waxes was derived from 10 samples found in Guthrie (1960). A standard heat content for petroleum waxes was adopted from EIA (2009a).

### Uncertainty

Although there is considerable qualitative uncertainty associated with the allocation of petroleum waxes and microcrystalline waxes, the quantitative variation in the C contents for all waxes is limited to  $\pm 1$  percent because of the nearly uniform relationship between C and other elements in petroleum waxes broadly defined.

### *Crude Oil, Unfinished Oils, and Miscellaneous Products*

U.S. energy statistics include several categories of petroleum products designed to ensure that reported refinery accounts "balance" and cover any "loopholes" in the taxonomy of petroleum products. These categories include crude oil, unfinished oils, and miscellaneous products. Crude oil is rarely consumed directly, miscellaneous products account for less than one percent of oil consumption, and unfinished oils are a balancing item that may show negative consumption. For C accounting purposes, it was assumed that all these products have the same C content as crude oil.

### Methodology

EIA reports on the average density and sulfur content of U.S. crude oil purchased by refineries. To develop a method of estimating C content based on this information, results of ultimate analyses of 182 crude oil samples were collected. Within the sample set, C content ranged from 82 to 88 percent C, but almost all samples fell between 84 percent and 86 percent C. The density and sulfur content of the crude oil data were regressed on the C content, producing the following equation:

$$\text{Percent C} = 76.99 + (10.19 \times \text{Specific Gravity}) + (-0.76 \times \text{Sulfur Content})$$

Absent the term representing sulfur content, the equation had an R-squared of only 0.35.<sup>24</sup> When C content was adjusted to exclude sulfur, the R-squared value rose to 0.65. While sulfur is the most important non-hydrocarbon impurity, nitrogen and oxygen can also be significant, but they do not seem to be correlated with either density or sulfur content. Restating these results, density accounts for about 35 percent of the variation in C content, impurities account for about 30 percent of the variation, and the remaining 35 percent is accounted for by other factors, including (presumably) the degree to which aromatics and polynuclear aromatics are present in the crude oil. Applying this equation to the 2014 crude oil quality data (30.21 degrees API and 1.47 percent sulfur) produces an estimated C content of 84.79 percent. Applying the density and C content to the EIA standard energy content for crude oil of 5.800 MMBtu per barrel produced an emissions coefficient of 20.31 MMT C/QBtu.

---

<sup>24</sup> R-squared represents the percentage of variation in the dependent variable (in this case carbon content) explained by variation in the independent variables.

### Data Sources

Carbon content was derived from 182 crude oil samples, including 150 samples from U.S. National Research Council (1927). A standard heat content for crude oil was adopted from EIA (2009a).

### Uncertainty

The uncertainty of the estimated C content for crude oil centers on the 35 percent of variation that cannot be explained by density and sulfur content. This variation is likely to alter the C content coefficient by  $\pm 3$  percent. Since unfinished oils and miscellaneous products are impossible to define, the uncertainty of applying a crude oil C content is likely to be bounded by the range of petroleum products described in this chapter at  $\pm 10$  percent.

## **Chronology and Explanation of Changes in Individual Carbon Content Coefficients of Fossil Fuels**

### **Coal**

#### Original 1994 Analysis

A set of 5,426 coal samples from the EIA coal analysis file were used to develop C content estimates. The results from that sample set appear below in Table A-55. The EIA Coal Analysis File was originally developed by the U.S. Bureau of Mines and contained over 60,000 coal samples obtained through numerous coal seams throughout the United States. Many of the samples were collected starting in the 1940s and 1950s through the 1980s and analyzed in U.S. government laboratories.

**Table A-55: Carbon Content Coefficients for Coal by Consuming Sector and Coal Rank, 1990 – 2000 (MMT C/QBtu)**

|                             | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  |
|-----------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>Consuming Sector</b>     |       |       |       |       |       |       |       |       |       |       |       |
| Electric Power              | 25.68 | 25.69 | 25.69 | 26.71 | 25.72 | 25.74 | 25.74 | 25.76 | 25.76 | 25.76 | 25.76 |
| Industrial Coking           | 25.51 | 25.51 | 25.51 | 25.51 | 25.52 | 25.53 | 25.55 | 25.56 | 25.56 | 25.56 | 25.56 |
| Other Industrial            | 25.58 | 25.59 | 25.62 | 25.61 | 25.63 | 25.63 | 25.61 | 25.63 | 25.63 | 25.63 | 25.63 |
| Residential /<br>Commercial | 25.92 | 26.00 | 26.13 | 25.97 | 25.95 | 26.00 | 25.92 | 26.00 | 26.00 | 26.00 | 26.00 |
| <b>Coal Rank</b>            |       |       |       |       |       |       |       |       |       |       |       |
| Anthracite                  | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 | 28.13 |
| Bituminous                  | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 | 25.37 |
| Sub-bituminous              | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 | 26.24 |
| Lignite                     | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 | 26.62 |

Sources: Emission factors by consuming sector from B.D. Hong and E.R. Slatnick, "Carbon Dioxide Emission Factors for Coal," U.S. Energy Information Administration, *Quarterly Coal Report*, January-March 1994 (Washington, DC, 1994); and emission factors by rank from Science Applications International Corporation, "Analysis of the Relationship Between Heat and Carbon Content of U.S. Fuels: Final Task Report," Prepared for the U.S. Energy Information Administration, Office of Coal, Nuclear, Electric and Alternative Fuels (Washington, DC 1992).

#### 2002 Update

The methodology employed for these estimates was unchanged from previous years; however, the underlying coal data sample set was updated. A new database, CoalQual 2.0 (1998), compiled by the U.S. Geological Survey (USGS) was adopted for the updated analysis. The updated sample set included 6,588 coal samples collected by the USGS and its state affiliates between 1973 and 1989. The decision to switch to the sample data contained in the USGS CoalQual database from the EIA database was made because the samples contained in the USGS database were collected and analyzed more recently than those obtained by EIA from the Bureau of Mines. The new coefficients developed in the 2002 revision were in use through the 1990 through 2007 Inventory and are provided in Table A-57.

**Table A-56: Carbon Content Coefficients for Coal by Consuming Sector and Coal Rank, 1990 – 2000 (MMT C/QBtu)**

|                         | 1990  | 1991  | 1992  | 1993  | 1994  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>Consuming Sector</b> |       |       |       |       |       |       |       |       |       |       |       |
| Electric Power          | 25.68 | 25.69 | 25.69 | 25.71 | 25.72 | 25.74 | 25.74 | 25.76 | 25.76 | 25.76 | 25.76 |
| Industrial Coking       | 25.51 | 25.51 | 25.51 | 25.51 | 25.52 | 25.53 | 25.55 | 25.56 | 25.56 | 25.56 | 25.56 |
| Other Industrial        | 25.58 | 25.60 | 25.62 | 25.61 | 25.63 | 25.63 | 25.61 | 25.63 | 25.63 | 25.63 | 25.63 |
| Residential/ Commercial | 25.92 | 26.00 | 26.13 | 25.97 | 25.95 | 26.00 | 25.92 | 26.00 | 26.00 | 26.00 | 26.00 |
| <b>Coal Rank</b>        |       |       |       |       |       |       |       |       |       |       |       |
| Anthracite              | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 |
| Bituminous              | 25.43 | 25.45 | 25.44 | 25.45 | 25.46 | 25.47 | 25.47 | 25.48 | 25.47 | 25.48 | 25.49 |
| Sub-bituminous          | 26.50 | 26.49 | 26.49 | 26.48 | 26.49 | 26.49 | 26.49 | 26.49 | 26.49 | 26.49 | 26.48 |
| Lignite                 | 26.19 | 26.21 | 26.22 | 26.21 | 26.24 | 26.22 | 26.17 | 26.20 | 26.23 | 26.26 | 26.30 |

Sources: Data from USGS, U.S. Coal Quality Database Version 2.0 (1998) and analysis prepared by SAIC (2007).

#### 2007 Update

The analysis of the USGS Coal Qual data was updated in 2007 to make a technical correction that affected the value for lignite and those sectors which consume lignite Table A-57 contains the annual coefficients that resulted from the 2007 analysis.

**Table A-57: Carbon Content Coefficients for Coal by Consuming Sector and Coal Rank, 1990-2007 (MMT C/QBtu)**

|                         | 1990  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  |
|-------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>Consuming Sector</b> |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Electric Power          | 25.68 | 25.74 | 25.74 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 | 25.76 |
| Industrial Coking       | 25.51 | 25.53 | 25.55 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 | 25.56 |
| Other Industrial        | 25.58 | 25.63 | 25.61 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 | 25.63 |
| Residential/Commercial  | 25.92 | 26.00 | 25.92 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 | 26.00 |
| <b>Coal Rank</b>        |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Anthracite              | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 | 28.26 |
| Bituminous              | 25.43 | 25.47 | 25.47 | 25.48 | 25.47 | 25.48 | 25.49 | 25.49 | 25.49 | 25.49 | 25.49 | 25.49 | 25.49 | 25.49 |
| Sub-bituminous          | 26.50 | 26.49 | 26.49 | 26.49 | 26.49 | 26.49 | 26.48 | 26.48 | 26.48 | 26.48 | 26.48 | 26.48 | 26.48 | 26.48 |
| Lignite                 | 26.19 | 26.22 | 26.17 | 26.20 | 26.23 | 26.26 | 26.30 | 26.30 | 26.30 | 26.30 | 26.30 | 26.30 | 26.30 | 26.57 |

Sources: Data from USGS, U.S. Coal Quality Database Version 2.0 (1998) and analysis prepared by (SAIC 2007).

## 2010 Update

The estimated annual C content coefficients for coal by rank and sector of consumption were updated again in 2010. Sample data from the Energy Institute at Pennsylvania State University (504 samples) were added to the 6,588 USGS samples to create a new database of 7,092 samples. The same analytical method used in the 2002 update was applied using these additional samples to calculate revised state-level carbon contents for each coal rank and then for national average consumption by end-use sector and by rank.

### **Natural Gas**

A revised analytical methodology underlies the natural gas coefficients used in this report. Prior to the current Inventory, descriptive statistics were used to stratify 6,743 samples of pipeline quality natural gas by heat content and then to determine the average C content of natural gas at the national average heat content (EIA 1994). The same coefficient was applied to all pipeline natural gas consumption for all years, because U.S. energy statistics showed a range of national average heat contents of pipeline gas of only 1,025 to 1,031 Btu per cubic foot (1 percent) from 1990 through 1994. A separate factor was developed in the same manner for all flared gas. In the previous Inventory, a weighted national average C content was calculated using the average C contents for each sub-sample of gas that conformed with an individual state's typical cubic foot of natural gas since there is regional variation in energy content. The result was a weighted national average of 14.47 MMT C/QBtu.

The current Inventory is revised to make use of the same set of samples, but utilizes a regression equation, as described above, of sample-based heat content and carbon content data in order to calculate annually-variable national average C content coefficients based on annual national average heat contents for pipeline natural gas and for flare gas. In addition, the revised analysis calculates an average C content from all samples with less than 1.5 percent CO<sub>2</sub> and less than 1,050 Btu/cf (samples most closely approximating the makeup of pipeline quality natural gas). The result was identical to the previous weighted national average of 14.47 MMT C/QBtu. The average C contents from the 1994 calculations are presented in Table A-58 below for comparison.

**Table A-58: Carbon Content of Pipeline-Quality Natural Gas by Energy Content (MMT C/QBtu)**

| <b>Sample</b>             | <b>Average Carbon Content</b> |
|---------------------------|-------------------------------|
| GRI Full Sample           | 14.51                         |
| Greater than 1,000 Btu    | 14.47                         |
| 1,025 to 1,035 Btu        | 14.45                         |
| 975 to 1,000 Btu          | 14.73                         |
| 1,000 to 1,025 Btu        | 14.43                         |
| 1,025 to 1,050 Btu        | 14.47                         |
| 1,050 to 1,075 Btu        | 14.58                         |
| 1,075 to 1,100 Btu        | 14.65                         |
| Greater than 1,100 Btu    | 14.92                         |
| Weighted National Average | 14.47                         |

Source: EIA (1994).

### **Petroleum Products**

All of the petroleum product C coefficients except that for Aviation Gasoline Blending Components have been updated for the current Inventory. EPA is updating these factors to better align the fuel properties data that underlie the Inventory factors with those published in the *Mandatory Reporting of Greenhouse Gases Rule* (EPA 2009b), Suppliers of Petroleum Products (MM) and Stationary Combustion (C) subparts. The coefficients that were applied in the previous report are provided in Table A-59 below. Specifically, each of the coefficients used in this report have been calculated from updated density and C share data, largely adopted from analyses undertaken for the Rule (EPA 2009b). In some cases, the heat content applied to the conversion to a carbon-per-unit-energy basis has also been updated. Additionally, the category Misc. Products (U.S. Territories), which is based upon the coefficients calculated for crude oil, has been allowed to vary annually with the crude oil coefficient. The petrochemical feedstock category has been eliminated for this report because the constituent products—naphthas and other oils—are estimated independently. Further, although the level of aggregation of U.S. energy statistics currently limits the application of coefficients for residual and distillate fuels to these two generic classifications, individual coefficients for the five major types of fuel oil (Nos. 1, 2, 4, 5 and 6) have been estimated for the current report and are presented in Table A-48 above. Each of the C coefficients applied in the previous Inventory is provided below for comparison (Table A-59).

**Table A-59: Carbon Content Coefficients and Underlying Data for Petroleum Products**

| Fuel                     | 2007 Carbon Content (MMT C/QBtu) | Gross Heat of Combustion (MMBtu/Barrel) | Density (API Gravity) | Percent Carbon     |
|--------------------------|----------------------------------|---|-----------------------|--------------------|
| Motor Gasoline           | 19.33                            | 5.219                                   | 59.1                  | 86.60              |
| LPG (total) <sup>a</sup> | 16.99                            | (See b)                                 | (See b)               | (See b)            |
| LPG (energy use)         | 17.18                            | (See b)                                 | (See b)               | (See b)            |
| LPG (non-energy use)     | 16.76                            | (See b)                                 | (See b)               | (See b)            |
| Jet Fuel                 | 19.33                            | 5.670                                   | 42.0                  | 86.30              |
| Distillate Fuel          | 19.95                            | 5.825                                   | 35.5                  | 86.34              |
| Residual Fuel            | 21.49                            | 6.287                                   | 11.0                  | 85.68              |
| Asphalt and Road Oil     | 20.62                            | 6.636                                   | 5.6                   | 83.47              |
| Lubricants               | 20.24                            | 6.065                                   | 25.6                  | 85.80              |
| Petrochemical Feedstocks | 19.37                            | 5.248 <sup>c</sup>                      | 67.1 <sup>c</sup>     | 84.11 <sup>c</sup> |
| Aviation Gas             | 18.87                            | 5.048                                   | 69.0                  | 85.00              |
| Kerosene                 | 19.72                            | 5.670                                   | 41.4                  | 86.01              |
| Petroleum Coke           | 27.85                            | 6.024                                   | -                     | 92.28              |
| Special Naphtha          | 19.86                            | 5.248                                   | 51.2                  | 84.76              |
| Petroleum Waxes          | 19.81                            | 5.537                                   | 43.3                  | 85.29              |
| Still Gas                | 17.51                            | 6.000                                   | -                     | -                  |
| Crude Oil                | 20.33                            | 5.800                                   | 30.5                  | 85.49              |
| Unfinished Oils          | 20.33                            | 5.825                                   | 30.5                  | 85.49              |
| Miscellaneous Products   | 20.33                            | 5.796                                   | 30.5                  | 85.49              |
| Pentanes Plus            | 18.24                            | 4.620                                   | 81.7                  | 83.70              |
| Natural Gasoline         | 18.24                            | 4.620                                   | 81.7                  | 83.70              |

<sup>a</sup> LPG is a blend of multiple paraffinic hydrocarbons: ethane, propane, isobutane, and normal butane, each with their own heat content, density and C content, see Table A-51.

<sup>b</sup> Heat, density, and percent carbon values are provided separately for ethane, propane and isobutene.

<sup>c</sup> Parameters presented are for naphthas with a boiling temperature less than 400 degrees Fahrenheit. Petrochemical feedstocks with higher boiling points are assumed to have the same characteristics as distillate fuel.

Note: "-" Indicates no sample data available.

Sources: EIA (1994); EIA (2008a); SAIC (2007).

Additional revisions to the Inventory's C coefficients since 1990 are detailed below.

### **Jet Fuel**

#### 1995 Update

Between 1994 and 1995, the C content coefficient for kerosene-based jet fuel was revised downward from 19.71 MMT C/QBtu to 19.33 MMT C/QBtu. This downward revision was the result of a shift in the sample set used from one collected between 1959 and 1972 and reported on by Martel and Angello in 1977 to one collected by Boeing in 1989 and published by Hadaller and Momenthy in 1990. The downward revision was a result of a decrease in density, as well as slightly lower C shares than in the earlier samples. However, the assumed heat content is unchanged because it is based on an EIA standard and probably yields a downward bias in the revised C content coefficient.

#### 1990 through 2008 Inventory Update

The coefficient was revised again for the 1990 through 2008 Inventory, returning to Martel and Angello and NIPER as the source of the carbon share and density data, respectively, for kerosene-based fuels. This change was made in order to align the coefficients used for this report with the values used in the *Mandatory Reporting of Greenhouse Gases Rule* (EPA 2009b). The return to the use of the Martel and Angello and NIPER coefficients was deemed more appropriate for the Rule as it was considered a more conservative coefficient given the uncertainty and variability in coefficients across the types of jet fuel in use in the United States. The factor will be revisited in future Inventories in light of data received from reporting entities in response to the Rule.

### Liquefied Petroleum Gases (LPG)

The C content coefficient of LPG is updated annually to reflect changes in the consumption mix of the underlying compounds: ethane; propane; isobutane; and normal butane. In 1994, EIA included pentanes plus—assumed to have the characteristics of hexane—in the mix of compounds broadly described as LPG. In 1995, EIA removed pentanes plus from this fuel category. Because pentanes plus is relatively rich in C per unit of energy, its removal from the consumption mix lowered the C content coefficient for LPG from 17.26 MMT C/QBtu to 16.99 MMT C/QBtu. In 1998, EIA began separating LPG consumption into two categories: energy use and non-fuel use and providing individual coefficients for each. Because LPG for fuel use typically contains higher proportions of propane than LPG for non-fuel use, the C content coefficient for fuel use was 1.8 to 2.5 percent higher than the coefficient for non-fuel use in previous inventories (see Table A-59).

However, for the current update of the LPG coefficients, the assumptions that underlie the selection of density and heat content data for each pure LPG compound have been updated, leading to a significant revision of the assumed properties of ethane. For this report, the physical characteristics of ethane, which constitutes over 90 percent of LPG consumption for non-fuel uses, have been updated to reflect ethane that is in (refrigerated) liquid form. Previously, the share of ethane was included using the density and energy content of gaseous ethane. Table A-60, below, compares the values applied for each of the compounds under the two sets of coefficient calculations. The C share of each pure compound was also updated by using more precise values for each compound's molecular weight.

Due in large part to the revised assumptions for ethane, the weighted C content for non-fuel use is now higher than that of the weighted coefficient for fuel use, which is dominated by the consumption of more dense propane. Under the revised assumptions, each annual weighted coefficient for non-fuel LPG consumption is 1.2 to 1.7 percent higher each year than is that for LPGs consumed for fuel (energy) uses.

**Table A-60: Physical Characteristics of Liquefied Petroleum Gases**

| Compound  | Chemical Formula               | 1990-2007          | Updated            | 1990-2007                  | Updated                    | 1990-2007                          | Updated                            |
|-----------|--------------------------------|--------------------|--------------------|----------------------------|----------------------------|------------------------------------|------------------------------------|
|           |                                | Density (bbl / MT) | Density (bbl / MT) | Energy Content (MMBtu/bbl) | Energy Content (MMBtu/bbl) | C Content Coefficient (MMT C/QBtu) | C Content Coefficient (MMT C/QBtu) |
| Ethane    | C <sub>2</sub> H <sub>6</sub>  | 16.88              | 11.55              | 2.916                      | 3.082                      | 16.25                              | 17.16                              |
| Propane   | C <sub>3</sub> H <sub>8</sub>  | 12.44              | 12.76              | 3.824                      | 3.836                      | 17.20                              | 16.76                              |
| Isobutane | C <sub>4</sub> H <sub>10</sub> | 11.20              | 11.42              | 4.162                      | 3.974                      | 17.75                              | 17.77                              |
| n-butane  | C <sub>4</sub> H <sub>10</sub> | 10.79              | 10.98              | 4.328                      | 4.326                      | 17.72                              | 17.75                              |

Sources: Updated: Densities – CRC Handbook of Chemistry and Physics, 89<sup>th</sup> Ed. (2008/09); Energy Contents – EPA (2009b). All values are for the compound in liquid form. The density and energy content of ethane are for refrigerated ethane (-89 degrees C). Values for n-butane are for pressurized butane (-25 degrees C). Values in previous editions of this Inventory: Gurthrie (1960).

### Motor Gasoline

The C content coefficient for motor gasoline varies annually based on the density of and proportion of additives in a representative sample of motor gasoline examined each year. However, in 1997 EIA began incorporating the effects of the introduction of reformulated gasoline into its estimate of C content coefficients for motor gasoline. This change resulted in a downward step function in C content coefficients for gasoline of approximately 0.3 percent beginning in 1995. In 2005 through 2006 reformulated fuels containing ethers began to be phased out nationally. Ethanol was added to gasoline blends as a replacement oxygenate, leading to another shift in gasoline density (see Table A-49), in the list and proportion of constituents that form the blend and in the blended C share based on those constituents.

**Table A-61: Carbon Content Coefficients for Petroleum Products, 1990-2007 (IMMT C/QBtu)**

| Fuel Type                         | 1990  | 1995  | 1996  | 1997  | 1998  | 1999  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  |
|-----------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>Petroleum</b>                  |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Asphalt and Road Oil              | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 | 20.62 |
| Aviation Gasoline                 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 |
| Distillate Fuel Oil               | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 |
| Jet Fuel <sup>a</sup>             | 19.40 | 19.34 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 |
| Kerosene                          | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 | 19.72 |
| LPG (energy use) <sup>a</sup>     | 17.21 | 17.20 | 17.20 | 17.18 | 17.23 | 17.25 | 17.20 | 17.21 | 17.20 | 17.21 | 17.20 | 17.19 | 17.19 | 17.18 |
| LPG (non-energy use) <sup>a</sup> | 16.83 | 16.87 | 16.86 | 16.88 | 16.88 | 16.84 | 16.81 | 16.83 | 16.82 | 16.84 | 16.81 | 16.81 | 16.78 | 16.76 |
| Lubricants                        | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 | 20.24 |
| Motor Gasoline <sup>a</sup>       | 19.41 | 19.38 | 19.36 | 19.35 | 19.33 | 19.33 | 19.34 | 19.34 | 19.35 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 |
| Residual Fuel                     | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 | 21.49 |
| <b>Other Petroleum</b>            |       |       |       |       |       |       |       |       |       |       |       |       |       |       |
| Av Gas Blend Comp.                | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 | 18.87 |
| Mo Gas Blend Comp <sup>a</sup>    | 19.41 | 19.38 | 19.36 | 19.35 | 19.33 | 19.33 | 19.34 | 19.34 | 19.35 | 19.33 | 19.33 | 19.33 | 19.33 | 19.33 |
| Crude Oil <sup>a</sup>            | 20.16 | 20.23 | 20.25 | 20.24 | 20.24 | 20.19 | 20.23 | 20.29 | 20.30 | 20.28 | 20.33 | 20.33 | 20.33 | 20.33 |
| Misc. Products <sup>a</sup>       | 20.16 | 20.23 | 20.25 | 20.24 | 20.24 | 20.19 | 20.23 | 20.29 | 20.30 | 20.28 | 20.33 | 20.33 | 20.33 | 20.33 |
| Misc. Products (Terr.)            | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 | 20.00 |
| Naphtha (<401 deg. F)             | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 | 18.14 |
| Other oil (>401 deg. F)           | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 | 19.95 |
| Pentanes Plus                     | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 | 18.24 |
| Petrochemical Feed.               | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 | 19.37 |
| Petroleum Coke                    | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 | 27.85 |
| Still Gas                         | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 | 17.51 |
| Special Naphtha                   | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 | 19.86 |
| Unfinished Oils <sup>a</sup>      | 20.16 | 20.23 | 20.25 | 20.24 | 20.24 | 20.19 | 20.23 | 20.29 | 20.30 | 20.28 | 20.33 | 20.33 | 20.33 | 20.33 |
| Waxes                             | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 |
| Other Wax and Misc.               | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 | 19.81 |

<sup>a</sup>C contents vary annually based on changes in fuel composition.

## References

- AAM (2009) *Diesel Survey*. Alliance of Automobile Manufacturers, Winter 2008.
- American Gas Association (1974) *Gas Engineer's Handbook*, New York, NY, Industrial Press, pp. 3/71, 3.87.
- API (1990 through 2008) *Sales of Natural Gas Liquids and Liquefied Refinery Gases*, American Petroleum Institute.
- API (1988) *Alcohols and Ethers: A Technical Assessment of Their Applications as Fuels and Fuel Components*, American Petroleum Institute, API 4261.
- ASTM (1985) *ASTM and Other Specifications for Petroleum Products and Lubricants*, American Society for Testing and Materials. Philadelphia, PA.
- Black, F. and L. High (1979) "Methodology for Determining Particulate and Gaseous Diesel Emissions," in, *The Measurement and Control of Diesel Particulate Emissions*, Society of Automotive Engineers, p. 128.
- Boldt, K. and B.R. Hall (1977) *Significance of Tests for Petroleum Products*, Philadelphia, PA, American Society for Testing and Materials, p. 30.
- Chemical Rubber Company (CRC), (2008/2009), *Handbook of Chemistry and Physics*, 89th Ed., editor D. Lide, Cleveland, OH: CRC Press.
- DOC (1929) *Thermal Properties of Petroleum Products*, U.S. Department of Commerce, National Bureau of Standards. Washington, DC. pp.16-21.
- EIA (2001-2009b) *Coal Distribution - Annual*, U.S. Department of Energy, Energy Information Administration. Washington, DC. DOE/EIA.
- EIA (2008a) *Monthly Energy Review, September 2006* and Published Supplemental Tables on Petroleum Product detail. Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2007/9).
- EIA (2008b) *Documentation for Emissions of Greenhouse Gases in the United States 2006*. DOE/EIA-0638(2006). October 2008.
- EIA (2009a) *Annual Energy Review*, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0384(2008).
- EIA (2009b) *Petroleum Supply Annual*, Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at [http://www.eia.doe.gov/oil\\_gas/petroleum/data\\_publications/petroleum\\_supply\\_annual/psa\\_volume1/psa\\_volume1.html](http://www.eia.doe.gov/oil_gas/petroleum/data_publications/petroleum_supply_annual/psa_volume1/psa_volume1.html).
- EIA (2001-2009a) *Annual Coal Report*, U.S. Department of Energy, Energy Information Administration. Washington, DC. DOE/EIA 0584.
- EIA (2001) *Cost and Quality of Fuels for Electric Utility Plants 2000*, Energy Information Administration. Washington, DC. August 2001. Available online at [http://www.eia.doe.gov/cneaf/electricity/cq/cq\\_sum.html](http://www.eia.doe.gov/cneaf/electricity/cq/cq_sum.html).
- EIA (1990-2001) *Coal Industry Annual*, U.S. Department of Energy, Energy Information Administration. Washington, DC. DOE/EIA 0584.
- EIA (1994) *Emissions of Greenhouse Gases in the United States 1987-1992*, Energy Information Administration, U.S. Department of Energy. Washington, DC. November, 1994. DOE/EIA 0573.
- EIA (1993) *Btu Tax on Finished Petroleum Products*, Energy Information Administration, Petroleum Supply Division (unpublished manuscript, April 1993).
- EPA (2010) *Carbon Content Coefficients Developed for EPA's Inventory of Greenhouse Gases and Sinks*. Office of Air and Radiation, Office of Atmospheric Programs, U.S. Environmental Protection Agency, Washington, D.C.
- EPA (2009a), "Industry Overview and Current Reporting Requirements for Petroleum Refining and Petroleum Imports," Petroleum Product Suppliers Technical Support Document for the Proposed Mandatory Reporting Rule. Office of Air and Radiation. 30 January, 2009.
- EPA (2009b) *Mandatory Reporting of Greenhouse Gases Rule*. Federal Register Docket ID EPA-HQ-OAR-2008-0508-2278, 30 September, 2009.

- Gas Technology Institute (1992) Database as documented in W.E. Liss, W.H. Thrasher, G.F. Steinmetz, P. Chowdiah, and A. Atari, *Variability of Natural Gas Composition in Select Major Metropolitan Areas of the United States*. GRI-92/0123. March 1992.
- Guthrie, V.B. (ed.) (1960) *Characteristics of Compounds*, Petroleum Products Handbook, p.3-3. New York, NY, McGraw-Hill.
- Hadaller, O.J. and A.M. Momeny (1990) *The Characteristics of Future Fuels*, Part 1, "Conventional Heat Fuels". Seattle, WA, Boeing Corp. September 1990. pp. 46-50.
- Intergovernmental Panel on Climate Change (IPCC) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. Prepared by the National Greenhouse Gas Inventories Programme (Japan, 2006).
- Matar, S. and L. Hatch (2000) *Chemistry of Petrochemical Processes*, 2<sup>nd</sup> Ed. Gulf Publishing Company: Houston.
- Martel, C.R., and L.C. Angello (1977) "Hydrogen Content as a Measure of the Combustion Performance of Hydrocarbon Fuels," in *Current Research in Petroleum Fuels*, Volume I. New York, NY, MSS Information Company, p. 116.
- Martin, S.W. (1960) "Petroleum Coke," in Virgil Guthrie (ed.), *Petroleum Processing Handbook*, New York, NY, McGraw-Hill, pp. 14-15.
- Meyers, (2004) *Handbook of Petroleum Refining Processes*, 3<sup>rd</sup> ed., NY, NY: McGraw Hill.
- National Institute for Petroleum and Energy Research (1990 through 2009) *Motor Gasolines, Summer and Motor Gasolines, Winter*.
- NIPER (1993) C. Dickson, *Aviation Turbine Fuels, 1992*, NIPER-179 PPS93/2 (Bartlesville, OK: National Institute for Petroleum and Energy Research, March 1993).
- Pennsylvania State University (PSU) (2010) Coal Sample Bank and Database. Data received by SAIC 18 February 2010 from Gareth Mitchell, The Energy Institute, Pennsylvania State University.
- Quick, Jeffrey (2010) "Carbon Dioxide Emission Factors for U.S. Coal by Origin and Destination," *Environmental Science & Technology*, Forthcoming.
- SAIC (2007) Analysis prepared by Science Applications International Corporation for EPA, Office of Air and Radiation, Market Policies Branch.
- U.S. National Research Council (1927) *International Critical Tables of Numerical Data, Physics, Chemistry, and Technology*, New York, NY, McGraw-Hill.
- Unzelman, G.H. (1992) "A Sticky Point for Refiners: FCC Gasoline and the Complex Model," *Fuel Reformulation*, July/August 1992, p. 29.
- USGS (1998) *CoalQual Database Version 2.0*, U.S. Geological Survey.
- Wauquier, J., ed. (1995) *Petroleum Refining, Crude Oil, Petroleum Products and Process Flowsheets* (Editions Technip - Paris, 1995) pg.225, Table 5.16.

## 2.3. Methodology for Estimating Carbon Emitted from Non-Energy Uses of Fossil Fuels

Carbon (C) storage associated with the non-energy use of fossil fuels was calculated by multiplying each fuel's potential emissions (i.e., each fuel's total C content) by a fuel-specific storage factor, as listed in Table A-62. The remaining C—i.e., that which is not stored—is emitted. This sub-annex explains the methods and data sources employed in developing the storage factors for petrochemical feedstocks (industrial other coal, natural gas for non-fertilizer uses, liquefied petroleum gases (LPG), pentanes plus, naphthas, other oils, still gas, special naphtha), asphalt and road oil, lubricants, waxes, and miscellaneous products. The storage factors<sup>25</sup> for the remaining non-energy fuel uses are either based on values recommended for use by IPCC (2006), or when these were not available, assumptions based on the potential fate of C in the respective non-energy use (NEU) products.

**Table A-62: Fuel Types and Percent of C Stored for Non-Energy Uses**

| Sector/Fuel Type                            | Storage Factor (%) |
|---|--------------------|
| <b>Industry</b>                             |                    |
| Industrial Coking Coal <sup>a</sup>         | 10%                |
| Industrial Other Coal <sup>b</sup>          | 65%                |
| Natural Gas to Chemical Plants <sup>b</sup> | 65%                |
| Asphalt & Road Oil                          | 100%               |
| LPG <sup>b</sup>                            | 65%                |
| Lubricants                                  | 9%                 |
| Pentanes Plus <sup>b</sup>                  | 65%                |
| Naphtha (<401 deg. F) <sup>b</sup>          | 65%                |
| Other Oil (>401 deg. F) <sup>b</sup>        | 65%                |
| Still Gas <sup>b</sup>                      | 65%                |
| Petroleum Coke <sup>c</sup>                 | 30%                |
| Special Naphtha <sup>b</sup>                | 65%                |
| Distillate Fuel Oil                         | 50%                |
| Waxes                                       | 58%                |
| Miscellaneous Products                      | 0%                 |
| <b>Transportation</b>                       |                    |
| Lubricants                                  | 9%                 |
| <b>U.S. Territories</b>                     |                    |
| Lubricants                                  | 9%                 |
| Other Petroleum (Misc. Prod.)               | 10%                |

<sup>a</sup> Includes processes for which specific coking coal consumption and emission factor data are not available. Consumption of coking coal for production of iron and steel is covered in the Industrial Processes and Product Use chapter.

<sup>b</sup> The storage factor listed is the value for 2014. As described in this annex, the factor varies over time.

<sup>c</sup> Assumes petroleum coke consumption is for pigments. Consumption of petroleum coke for production of primary aluminum anodes, electric arc furnace anodes, titanium dioxide, ammonia, urea, and ferroalloys is covered in the Industrial Processes and Product Use chapter.

The following sections describe the non-energy uses in greater detail, outlining the methods employed and data used in estimating each storage factor. Several of the fuel types tracked by EIA are used in organic chemical synthesis and in other manufacturing processes, and are referred to collectively as “petrochemical feedstocks.” Because the methods and data used to analyze them overlap, they are handled as a group and are discussed first. Discussions of the storage factors for asphalt and road oil, lubricants, waxes, and miscellaneous products follow.

<sup>25</sup> Throughout this section, references to “storage factors” represent the proportion of carbon stored.

## Petrochemical Feedstocks

Petrochemical feedstocks—industrial other coal, natural gas for non-fertilizer uses, LPG, pentanes plus, naphthas, other oils, still gas, special naphtha—are used in the manufacture of a wide variety of man-made chemicals and products. Plastics, rubber, synthetic fibers, solvents, paints, fertilizers, pharmaceuticals, and food additives are just a few of the derivatives of these fuel types. Chemically speaking, these fuels are diverse, ranging from simple natural gas (i.e., predominantly CH<sub>4</sub>) to heavier, more complex naphthas and other oils.<sup>26</sup>

After adjustments for (1) use in industrial processes and (2) net exports, these eight fuel categories constituted approximately 217.3 MMT CO<sub>2</sub> Eq., or 68 percent, of the 319.5 MMT CO<sub>2</sub> Eq. of non-energy fuel consumption in 2014. For 2014, the storage factor for the eight fuel categories was 65 percent. In other words, of the net consumption, 65 percent was destined for long-term storage in products—including products subsequently combusted for waste disposal—while the remaining 35 percent was emitted to the atmosphere directly as CO<sub>2</sub> (e.g., through combustion of industrial by-products) or indirectly as CO<sub>2</sub> precursors (e.g., through evaporative product use). The indirect emissions include a variety of organic gases such as volatile organic compounds (VOCs) and carbon monoxide (CO), which eventually oxidize into CO<sub>2</sub> in the atmosphere. The derivation of the storage factor is described in the following sections.

### Methodology and Data Sources

The petrochemical feedstocks storage factor is equal to the ratio of C stored in the final products to total C content for the non-energy fossil fuel feedstocks used in industrial processes, after adjusting for net exports of feedstocks. One aggregate storage factor was calculated to represent all eight fuel feedstock types. The feedstocks were grouped because of the overlap of their derivative products. Due to the many reaction pathways involved in producing petrochemical products (or wastes), it becomes extraordinarily complex to link individual products (or wastes) to their parent fuel feedstocks.

Import and export data for feedstocks were obtained from the Energy Information Administration (EIA) for the major categories of petrochemical feedstocks. EIA's *Petroleum Supply Annual* publication tracks imports and exports of petrochemical feedstocks, including butanes, butylenes, ethane, ethylene, propane, propylene, LPG, and naphthas (i.e., most of the large volume primary chemicals produced by petroleum refineries). These imports and exports are already factored into the U.S. fuel consumption statistics. However, EIA does not track imports and exports of chemical intermediates and products produced by the chemical industry (e.g., xylenes, vinyl chloride), which are derived from the primary chemicals produced by the refineries. These products represent very large flows of C derived from fossil fuels (i.e., fossil C), so estimates of net flows not already considered in EIA's dataset were developed for the entire time series from 1990 to 2014.

The approach to estimate imports and exports involves three steps, listed here and then described in more detail below:

- Step 1.* Identify commodities derived from petrochemical feedstocks, and calculate net import/export for each.
- Step 2.* Estimate the C content for each commodity.
- Step 3.* Sum the net C imports/exports across all commodities.

Step 1 relies heavily on information provided by the National Petrochemical and Refiners Association (NPRAs) and U.S. Bureau of the Census (BoC) trade statistics published by the U.S. International Trade Commission (USITC). NPRAs provided a spreadsheet of the ten-digit BoC Harmonized Tariff Schedule (HTS) Commodity Codes used to compile import-export data for periodic reports issued to NPRAs's membership on trade issues. Additional feedstock commodities were identified by HTS code in the BoC data system and included in the net import/export analysis.

One of the difficulties in analyzing trade data is that a large portion of the outputs from the refining industry are fuels and fuel components, and it was difficult to segregate these from the outputs used for non-energy uses. The NPRAs-supplied codes identify fuels and fuel components, thus providing a sound basis for isolating net imports/exports of petrochemical feedstocks. Although MTBE and related ether imports are included in the published NPRAs data, these commodities are not included in the total net imports/exports calculated here, because it is assumed that they are fuel additives and do not contribute to domestic petrochemical feedstocks. Net exports of MTBE and related ethers are also not included in the totals, as these commodities are considered to be refinery products that are already accounted for in the EIA

---

<sup>26</sup> Naphthas are compounds distilled from petroleum containing 4 to 12 carbon atoms per molecule and having a boiling point less than 401 degrees Fahrenheit. "Other oils" are distillates containing 12 to 25 carbon atoms per molecule and having a boiling point greater than 401 degrees Fahrenheit.

data. Imports and exports of commodities for which production and consumption data are provided by EIA (e.g., butane, ethylene, and liquefied petroleum gases) are also not included in the totals, to avoid double-counting.

Another difficulty is that one must be careful to assure that there is not double-counting of imports and exports in the data set. Other parts of the mass balance (described later) provide information on C flows, in some cases based on production data and in other cases based on consumption data. Production data relates only to production within the country; consumption data incorporates information on imports and exports as well as production. Because many commodities are emissive in their use, but not necessarily their production, consumption data is appropriately used in calculations for emissive fates. For purposes of developing an overall mass balance on U.S. non-energy uses of C, for those materials that are non-emissive (e.g., plastics), production data is most applicable. And for purposes of adjusting the mass balance to incorporate C flows associated with imports and exports, it was necessary to carefully review whether or not the mass balance already incorporated cross-boundary flows (through the use of consumption data), and to adjust the import/export balance accordingly.

The BoC trade statistics are publicly available<sup>27</sup> and cover a complete time series from 1990 to 2014. These statistics include information on imports and exports of thousands of commodities. After collecting information on annual flows of the more than 100 commodities identified by NPRA, Step 2 involves calculating the C content for each commodity from its chemical formula. In cases where the imports and exports were expressed in units of volume, rather than mass, they were converted to mass based on the commodities' densities.

Step 3 involves summing the net C imports/exports across all commodities. The results of this step are shown in Table A-63. As shown in the table, the United States has been a net exporter of chemical intermediates and products throughout the 1990 to 2014 period.

**Table A-63: Net Exports of Petrochemical Feedstocks, 1990 – 2014 (MMT CO<sub>2</sub> Eq.)**

|             | 1990 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------|------|------|------|------|------|------|------|
| Net Exports | 12.0 | 6.5  | 7.2  | 7.9  | 10.1 | 8.5  | 3.8  |

After adjusting for imports and exports, the C budget is adjusted for the quantity of C that is used in the Industrial Processes and Product Use sector of the Inventory. Fossil fuels used for non-energy purposes in industrial processes—and for which C emissions and storage have been characterized through mass balance calculations and/or emission factors that directly link the non-energy use fossil fuel raw material and the industrial process product—are not included in the non-energy use sector. These industrial processes (and their non-energy use fossil fuel raw materials) include iron and steel (coal coke), primary aluminum (petroleum coke), titanium oxide (petroleum coke), ferroalloys (petroleum coke), and ammonia and urea (petroleum coke and natural gas).

For each year of the Inventory, the total C content of non-energy uses was calculated by starting with the EIA estimate of non-energy use, and reducing it by the adjustment factor for net exports (see Table A-63) to yield net domestic fuel consumption for non-energy. The balance was apportioned to either stored C or emissive C, based on a storage factor.

The overall storage factor for the feedstocks was determined by developing a mass balance on the C in feedstocks, and characterizing products, uses, and environmental releases as resulting in either storage or emissions. The total C in the system was estimated by multiplying net domestic consumption for non-energy by the C content of each of the feedstocks (i.e., industrial other coal, natural gas for non-fertilizer uses, LPG, pentanes plus, naphthas, other oils, still gas, special naphtha). Carbon content values for the fuel feedstocks are discussed in the Estimating Emissions from Fossil Fuel Combustion and Estimating the Carbon Content from Fossil Fuel Combustion Annexes.

Next, C pools and releases in a variety of industrial releases, energy recovery processes, and products were characterized. The C fate categories are plastics, energy recovery, synthetic rubber, synthetic fibers, organic solvents, C black, detergents and personal cleansers, industrial non-methane volatile organic compound (NMVOC) emissions, hazardous waste incineration, industrial toxic chemical (i.e., TRI) releases, pesticides, food additives, antifreeze and deicers (glycols), and silicones.<sup>28</sup>

<sup>27</sup> See the U.S International Trade Commission (USITC) Trade Dataweb at <<http://dataweb.usitc.gov/>>.

<sup>28</sup> For the most part, the releases covered by the U.S. Toxic Release Inventory (TRI) represent air emissions or water discharges associated with production facilities. Similarly, VOC emissions are generally associated with production facilities. These emissions could have been accounted for as part of the Waste chapter, but because they are not necessarily associated with waste management, they were included here. Toxic releases are not a “product” category, but they are referred to as such for ease of discussion.

The C in each product or waste produced was categorized as either stored or emitted. The aggregate storage factor is the C-weighted average of storage across fuel types. As discussed later in the section on uncertainty, the sum of stored C and emitted C (i.e., the outputs of the system) exceeded total C consumption (i.e., the inputs to the system) for some years in the time series.<sup>29</sup> To address this mass imbalance, the storage factor was calculated as C storage divided by total C outputs (rather than C storage divided by C inputs).

Note that the system boundaries for the storage factor do not encompass the entire life-cycle of fossil-based C consumed in the United States insofar as emissions of CO<sub>2</sub> from waste combustion are accounted for separately in the Inventory and are discussed in the Incineration of Waste section of the Energy chapter.

The following sections provide details on the calculation steps, assumptions, and data sources employed in estimating and classifying the C in each product and waste shown in Table A-64. Summing the C stored and dividing it by total C outputs yields the overall storage factor, as shown in the following equation for 2014:

$$\text{Overall Storage Factor} = \text{C Stored} / (\text{C Stored} + \text{C Emitted}) = \\ 142.3 \text{ MMT CO}_2 \text{ Eq.} / (142.3 + 75.1) \text{ MMT CO}_2 \text{ Eq.} = 65\%$$

**Table A-64: C Stored and Emitted by Products from Feedstocks in 2014 (MMT CO<sub>2</sub> Eq.)**

| Product/Waste Type          | C Stored<br>(MMT CO <sub>2</sub> Eq.) | C Emitted<br>(MMT CO <sub>2</sub> Eq.) |
|-----------------------------|---------------------------------------|--|
| <b>Industrial Releases</b>  | <b>0.1</b>                            | <b>5.5</b>                             |
| TRI Releases                | 0.1                                   | 1.0                                    |
| Industrial VOCs             | NA                                    | 3.6                                    |
| Non-combustion CO           | NA                                    | 0.5                                    |
| Hazardous Waste Incin.      | NA                                    | 0.4                                    |
| <b>Energy Recovery</b>      | <b>NA</b>                             | <b>41.0</b>                            |
| <b>Products</b>             | <b>142.2</b>                          | <b>12.5</b>                            |
| Plastics                    | 119.3                                 | NA                                     |
| Synthetic Rubber            | 14.8                                  | NA                                     |
| Antifreeze and deicers      | NA                                    | 0.8                                    |
| Abraded tire rubber         | NA                                    | 0.3                                    |
| Food additives              | NA                                    | 1.1                                    |
| Silicones                   | 0.5                                   | NA                                     |
| Synthetic Fiber             | 7.3                                   | NA                                     |
| Pesticides                  | 0.2                                   | 0.2                                    |
| Soaps, shampoos, detergents | NA                                    | 4.8                                    |
| Solvent VOCs                | NA                                    | 5.3                                    |
| <b>Total</b>                | <b>142.3</b>                          | <b>59.0</b>                            |

NA (Not Applicable)

Note: Totals may not sum due to independent rounding.

The three categories of C accounted for in the table are industrial releases, energy recovery, and products. Each is discussed below.

### **Industrial Releases**

Industrial releases include toxic chemicals reported through the Toxics Release Inventory (TRI), industrial emissions of volatile organic compounds (VOCs), CO emissions (other than those related to fuel combustion), and emissions from hazardous waste incineration.

#### *TRI Releases*

Fossil-derived C is found in many toxic substances released by industrial facilities. The TRI, maintained by EPA, tracks these releases by chemical and environmental release medium (i.e., land, air, or water) on a biennial basis (EPA

<sup>29</sup> Overall, there was fairly close agreement between inputs and outputs: for the entire 1990 through 2012 time series, inputs exceeded outputs by a time-weighted average of 0.2 percent. During the period 1990 through 2000, carbon inputs exceeded carbon outputs (i.e., the sum of carbon stored and carbon emitted) by a time-weighted average of 10 percent. For those years, the assumption was made that the “missing” carbon was lost through fates leading to emissions. This discrepancy shifted during the period from 2001 through 2012, in which carbon outputs exceeded carbon inputs by a time-weighted average of 12 percent.

2000b). By examining the C contents and receiving media for the top 35 toxic chemicals released, which account for 90 percent of the total mass of chemicals, the quantity of C stored and emitted in the form of toxic releases can be estimated.

The TRI specifies releases by chemical, so C contents were assigned to each chemical based on molecular formula. The TRI also classifies releases by disposal location as either off-site or on-site. The on-site releases are further subdivided into air emissions, surface water discharges, underground injection, and releases to land; the latter is further broken down to disposal in a RCRA Subtitle C (i.e., hazardous waste) landfill or to “Other On-Site Land Disposal.”<sup>30</sup> The C released in each disposal location is provided in Table A-65.

Each on-site classification was assigned a storage factor. A 100 percent storage factor was applied to disposition of C to underground injection and to disposal to RCRA-permitted landfills, while the other disposition categories were assumed to result in an ultimate fate of emission as CO<sub>2</sub> (i.e., a storage factor of zero was applied to these categories). The release allocation is not reported for off-site releases; therefore, the approach was to develop a C-weighted average storage factor for the on-site C and apply it to the off-site releases.

For the remaining 10 percent of the TRI releases, the weights of all chemicals were added and an average C content value, based upon the top 35 chemicals’ C contents, was applied. The storage and emission allocation for the remaining 10 percent of the TRI releases was carried out in the same fashion as for the 35 major chemicals.

Data on TRI releases for the full 1990 through 2014 time series were not readily available. Since this category is small (less than 1 MMT C emitted and stored), the 1998 value was applied for the entire time series.

**Table A-65: 1998 TRI Releases by Disposal Location (kt CO<sub>2</sub> Eq.)**

| Disposal Location                 | Carbon Stored<br>(kt CO <sub>2</sub> Eq.) | Carbon Emitted<br>(kt CO <sub>2</sub> Eq.) |
|-----------------------------------|---|--|
| Air Emissions                     | NA  | 924.0                                      |
| Surface Water Discharges          | NA  | 6.7  |
| Underground Injection             | 89.4                                      | NA   |
| RCRA Subtitle C Landfill Disposal | 1.4                                       | NA   |
| Other On-Site Land Releases       | NA  | 15.9                                       |
| Off-site Releases                 | 6.4                                       | 36.0                                       |
| <b>Total</b>                      | <b>97.2</b>                               | <b>982.6</b>                               |

NA (Not Applicable)

Note: Totals may not sum due to independent rounding.

#### *Volatile Organic Compound Emissions from Industrial Processes and Solvent Evaporation Emissions*

Data on annual non-methane volatile organic compound (NMVOC) emissions were obtained from preliminary data (EPA 2015b), and disaggregated based on EPA (2003), which, in its final iteration, will be published on the National Emission Inventory (NEI) Air Pollutant Emission Trends web site. The 1990 through 2014 Trends data include information on NMVOC emissions by end-use category; some of these fall into the heading of “industrial releases” in Table A-64 above, and others are related to “product use;” for ease of discussion, both are covered here. The end-use categories that represent “Industrial NMVOC Emissions” include some chemical and allied products, certain petroleum related industries, and other industrial processes. NMVOC emissions from solvent utilization (product use) were considered to be a result of non-energy use of petrochemical feedstocks. These categories were used to distinguish non-energy uses from energy uses; other categories where VOCs could be emitted due to combustion of fossil fuels were excluded to avoid double counting. Data for 2012, 2013, or 2014 are not yet available for most of the sources, so they have been set equal to 2011 values.

Because solvent evaporation and industrial NMVOC emission data are provided in tons of total NMVOCs, assumptions were made concerning the average C content of the NMVOCs for each category of emissions. The assumptions for calculating the C fraction of industrial and solvent utilization emissions were made separately and differ significantly. For industrial NMVOC emissions, a C content of 85 percent was assumed. This value was chosen to reflect the C content of an average volatile organic compound based on the list of the most abundant NMVOCs provided in the Trends Report. The list contains only pure hydrocarbons, including saturated alkanes (C contents ranging from 80 to 85 percent based upon

<sup>30</sup> Only the top nine chemicals had their land releases separated into RCRA Landfills and Other Land Disposal. For the remaining chemicals, it was assumed that the ratio of disposal in these two categories was equal to the carbon-weighted average of the land disposal fate of the top nine chemicals (i.e., 8 percent attributed to RCRA Landfills and 92 percent in the “Other” category).

C number), alkenes (C contents approximately 85 percent), and some aromatics (C contents approximately 90 percent, depending upon substitution).

An EPA solvent evaporation emissions dataset (Tooly 2001) was used to estimate the C content of solvent emissions. The dataset identifies solvent emissions by compound or compound category for six different solvent end-use categories: degreasing, graphic arts, dry cleaning, surface coating, other industrial processes, and non-industrial processes. The percent C of each compound identified in the dataset was calculated based on the molecular formula of the individual compound (e.g., the C content of methylene chloride is 14 percent; the C content of toluene is 91 percent). For solvent emissions that are identified in the EPA dataset only by chemical category (e.g., butanediol derivatives) a single individual compound was selected to represent each category, and the C content of the category was estimated based on the C content of the representative compound. The overall C content of the solvent evaporation emissions for 1998, estimated to be 56 percent, is assumed to be constant across the entire time series.

The results of the industrial and solvent NMVOC emissions analysis are provided in Table A-66 for 1990 through 2014. Industrial NMVOC emissions in 2014 were 3.6 MMT CO<sub>2</sub> Eq. and solvent evaporation emissions in 2014 were 5.3 MMT CO<sub>2</sub> Eq.

**Table A-66: Industrial and Solvent NMVOC Emissions**

|  | 1990  | 1995  | 2000  | 2005  | 2010  | 2011  | 2012  | 2013  | 2014  |
|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| <b>Industrial NMVOCs<sup>a</sup></b>     |       |       |       |       |       |       |       |       |       |
| NMVOCs ('000 Short Tons)                 | 1,279 | 1,358 | 802   | 825   | 1,189 | 1,289 | 1,289 | 1,289 | 1,289 |
| Carbon Content (%)                       | 85%   | 85%   | 85%   | 85%   | 85%   | 85%   | 85%   | 85%   | 85%   |
| Carbon Emitted (MMT CO <sub>2</sub> Eq.) | 3.6   | 3.8   | 2.3   | 2.3   | 3.4   | 3.6   | 3.6   | 3.6   | 3.6   |
| <b>Solvent Evaporation<sup>b</sup></b>   |       |       |       |       |       |       |       |       |       |
| Solvents ('000 Short Tons)               | 5,750 | 6,183 | 4,832 | 4,245 | 2,974 | 2,811 | 2,811 | 2,811 | 2,811 |
| Carbon Content (%)                       | 56%   | 56%   | 56%   | 56%   | 56%   | 56%   | 56%   | 56%   | 56%   |
| Carbon Emitted (MMT CO <sub>2</sub> Eq.) | 10.8  | 11.6  | 9.0   | 7.9   | 5.6   | 5.3   | 5.3   | 5.3   | 5.3   |

<sup>a</sup> Includes emissions from chemical and allied products, petroleum and related industries, and other industrial processes categories.

<sup>b</sup> Includes solvent usage and solvent evaporation emissions from degreasing, graphic arts, dry cleaning, surface coating, other industrial processes, and non-industrial processes.

#### *Non-Combustion Carbon Monoxide Emissions*

Carbon monoxide (CO) emissions data were also obtained from the NEI preliminary data (EPA 2015b), and disaggregated based on EPA (2003). There are three categories of CO emissions in the report that are classified as process-related emissions not related to fuel combustion. These include chemical and allied products manufacturing, metals processing, and other industrial processes. Some of these CO emissions are accounted for in the Industrial Processes and Product Use section of this report, and are therefore not accounted for in this section. These include total C emissions from the primary aluminum, titanium dioxide, iron and steel, and ferroalloys production processes. The total C (CO and CO<sub>2</sub>) emissions from oil and gas production, petroleum refining, and asphalt manufacturing are also accounted for elsewhere in this Inventory. Biogenic emissions (e.g., pulp and paper process emissions) are accounted for in the Land Use, Land-Use Change and Forestry chapter and excluded from calculation of CO emissions in this section. Those CO emissions that are not accounted for elsewhere are considered to be by-products of non-fuel use of feedstocks, and are thus included in the calculation of the petrochemical feedstocks storage factor. Table A-67 lists the CO emissions that remain after taking into account the exclusions listed above. As data for 2012, 2013, and 2014 are not yet available for most sources, they have been set equal to 2011 values.

**Table A-67: Non-Combustion Carbon Monoxide Emissions**

|  | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|--|------|------|------|------|------|------|------|------|------|
| CO Emissions ('000 Short Tons)           | 489  | 481  | 623  | 461  | 405  | 349  | 349  | 349  | 349  |
| Carbon Emitted (MMT CO <sub>2</sub> Eq.) | 0.7  | 0.7  | 0.9  | 0.7  | 0.6  | 0.5  | 0.5  | 0.5  | 0.5  |

Note: Includes emissions from chemical and allied products, petroleum and related industries, metals processing, and other industrial processes categories.

#### *Hazardous Waste Incineration*

Hazardous wastes are defined by the EPA under the Resource Conservation and Recovery Act (RCRA).<sup>31</sup> Industrial wastes, such as rejected products, spent reagents, reaction by-products, and sludges from wastewater or air

<sup>31</sup> [42 U.S.C. §6924, SDWA §3004]

pollution control, are federally regulated as hazardous wastes if they are found to be ignitable, corrosive, reactive, or toxic according to standardized tests or studies conducted by the EPA.

Hazardous wastes must be treated prior to disposal according to the federal regulations established under the authority of RCRA. Combustion is one of the most common techniques for hazardous waste treatment, particularly for those wastes that are primarily organic in composition or contain primarily organic contaminants. Generally speaking, combustion devices fall into two categories: incinerators that burn waste solely for the purpose of waste management, and boilers and industrial furnaces (BIFs) that burn waste in part to recover energy from the waste. More than half of the hazardous waste combusted in the United States is burned in BIFs; because these processes are included in the energy recovery calculations described below, they are not included as part of hazardous waste incineration.

EPA’s Office of Solid Waste requires biennial reporting of hazardous waste management activities, and these reports provide estimates of the amount of hazardous waste burned for incineration or energy recovery. EPA stores this information in its Resource Conservation and Recovery Act (RCRA) Information system (EPA 2013a), formerly reported in its Biennial Reporting System (BRS) database (EPA 2000a, 2009, 2015a). Combusted hazardous wastes are identified based on EPA-defined management system types M041 through M049 (incineration). Combusted quantities are grouped into four representative waste form categories based on the form codes reported in the BRS: aqueous liquids, organic liquids and sludges, organic solids, and inorganic solids. To relate hazardous waste quantities to C emissions, “fuel equivalent” factors were derived for hazardous waste by assuming that the hazardous wastes are simple mixtures of a common fuel, water, and noncombustible ash. For liquids and sludges, crude oil is used as the fuel equivalent and coal is used to represent solids.

Fuel equivalent factors were multiplied by the tons of waste incinerated to obtain the tons of fuel equivalent. Multiplying the tons of fuel equivalent by the C content factors (discussed in the Estimating the Carbon Content from Fossil Fuel Combustion Annex) yields tons of C emitted. Implied C content is calculated by dividing the tons of C emitted by the associated tons of waste incinerated.

Waste quantity data for hazardous wastes were obtained from EPA’s RCRA Information/BRS database for reporting years 1989, 1991, 1993, 1995, 1997, 1999, 2001, 2003, 2005, 2007, 2009, 2011, 2013, and 2014 (EPA 2000a, 2009, 2013a, 2015a). Combusted waste quantities were obtained from Form GM (Generation and Management) for wastes burned on site and Form WR (Wastes Received) for waste received from off-site for combustion. For each of the waste types, assumptions were developed on average waste composition (see Table A-68). Regulations require incinerators to achieve at least 99.99 percent destruction of organics; this formed the basis for assuming the fraction of C oxidized. Emissions from hazardous waste incineration in 2014 were 0.4 MMT CO<sub>2</sub> Eq. Table A-69 lists the CO<sub>2</sub> emissions from hazardous waste incineration.

**Table A-68: Assumed Composition of Combusted Hazardous Waste by Weight (Percent)**

| Waste Type                  | Water (%) | Noncombustibles (%) | Fuel Equivalent (%) |
|-----------------------------|-----------|---------------------|---------------------|
| Aqueous Waste               | 90        | 5                   | 5                   |
| Organic Liquids and Sludges | 40        | 20                  | 40                  |
| Organic Solids              | 20        | 40                  | 40                  |
| Inorganic Solids            | 20        | 70                  | 10                  |

**Table A-69: CO<sub>2</sub> Emitted from Hazardous Waste Incineration (MMT CO<sub>2</sub> Eq.)**

|             | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------|------|------|------|------|------|------|------|------|------|
| C Emissions | 1.1  | 1.7  | 1.0  | 0.6  | 0.3  | 0.4  | 0.4  | 0.4  | 0.4  |

### **Energy Recovery**

The amount of feedstocks combusted for energy recovery was estimated from data included in EIA’s Manufacturers Energy Consumption Survey (MECS) for 1991, 1994, 1998, 2002, 2006, and 2010 (EIA 1994, 1997, 2001, 2005, 2010, 2013b). Some fraction of the fossil C exiting refineries and designated for use for feedstock purposes actually ends up being combusted for energy recovery (despite the designation of feedstocks as a “non-energy” use) because the chemical reactions in which fuel feedstocks are used are not 100 percent efficient. These chemical reactions may generate unreacted raw material feedstocks or generate by-products that have a high energy content. The chemical industry and many downstream industries are energy-intensive and often have boilers or other energy recovery units on-site, and thus these unreacted feedstocks or by-products are often combusted for energy recovery. Also, as noted above in the section on hazardous waste incineration, regulations provide a strong incentive—and in some cases require—burning of organic wastes generated from chemical production processes.

Information available from the MECS include data on the consumption for energy recovery of “other” fuels in the petroleum and coal products, chemicals, primary metals, nonmetallic minerals, and other manufacturing sectors. These “other” fuels include refinery still gas; waste gas; waste oils, tars, and related materials; petroleum coke, coke oven and blast furnace gases; scrap tires; liquor or black liquor; woodchips and bark; and other uncharacterized fuels. Fuel use of petroleum coke is included separately in the fuel use data provided annually by EIA, and energy recovery of coke oven gas and blast furnace gas (i.e., by-products of the iron and steel production process) is addressed in the Iron and Steel production section in the Industrial Processes and Product Use chapter. Consumption of refinery still gas in the refinery sector is also included separately in the fuel use data from EIA. The combustion of scrap tires in cement kilns, lime kilns, and electric arc furnaces is accounted for in the Waste Incineration chapter; data from the Rubber Manufacturers Association (RMA 2009a) were used to subtract out energy recovery from scrap tires in these industries. Consumption of net steam, assumed to be generated from fossil fuel combustion, is also included separately in the fuel use data from EIA. Therefore, these categories of “other” fuels are addressed elsewhere in the Inventory and not considered as part of the petrochemical feedstocks energy recovery analysis. Liquor or black liquor and woodchips and bark are assumed to be biogenic fuels, in accordance with IPCC (2006), and therefore are not included in the Inventory. The remaining categories of fuels, including waste gas; waste oils, tars, and related materials; and other uncharacterized fuels are assumed to be petrochemical feedstocks burned for energy recovery (see Table A-70). The conversion factors listed in the Estimating Emissions from Fossil Fuel Combustion Annex were used to convert the Btu values for each fuel feedstock to MMT CO<sub>2</sub>. Petrochemical feedstocks combusted for energy recovery corresponded to 42.7 MMT CO<sub>2</sub> Eq. in 1991, 35.7 MMT CO<sub>2</sub> Eq. in 1994, 58.7 MMT CO<sub>2</sub> Eq. in 1998, 71.6 MMT CO<sub>2</sub> Eq. in 2002, 73.8 MMT CO<sub>2</sub> Eq. in 2006, and 41.0 MMT CO<sub>2</sub> Eq. in 2010. Values for petrochemical feedstocks burned for energy recovery for years between 1991 and 1994, between 1994 and 1998, between 1998 and 2002, between 2002 and 2006, and between 2007 and 2010 have been estimated by linear interpolation. The value for 1990 is assumed to be the same as the value for 1991, and the value for 2011, 2012, and 2013, 2014 are assumed to be the same as the value for 2010 (Table A-71).

**Table A-70: Summary of 2010 MECS Data for Other Fuels Used in Manufacturing/Energy Recovery (Trillion Btu)**

| Subsector and Industry                             | NAICS CODE | Waste Gas <sup>a</sup> | Waste Oils/Tars <sup>b</sup> | Refinery Still   |                        |                          |
|--|------------|------------------------|------------------------------|------------------|------------------------|--------------------------|
|  |            |                        |                              | Gas <sup>c</sup> | Net Steam <sup>d</sup> | Other Fuels <sup>e</sup> |
| Printing and Related Support                       | 323        | 0                      | 0                            | 0                | 0                      | 0                        |
| Petroleum and Coal Products                        | 324        | 0                      | 6                            | 1,349            | 153                    | 54                       |
| Chemicals  | 325        | 376                    | 7                            | 0                | 266                    | 110                      |
| Plastics and Rubber Products                       | 326        | 0                      | 0                            | 0                | 0                      | 0                        |
| Nonmetallic Mineral Products                       | 327        | 1                      | 7                            | 0                | 0                      | 0                        |
| Primary Metals                                     | 331        | 0                      | 0                            | 0                | 12                     | 34                       |
| Fabricated Metal Products                          | 332        | 0                      | 0                            | 0                | 0                      | 0                        |
| Machinery  | 333        | 0                      | 0                            | 0                | 0                      | 1                        |
| Computer and Electronic Products                   | 334        | 0                      | 0                            | 0                | 0                      | 0                        |
| Electrical Equip., Appliances, Components          | 335        | 0                      | 0                            | 0                | 0                      | 0                        |
| Transportation Equipment                           | 336        | 2                      | 0                            | 0                | 0                      | 3                        |
| Furniture and Related Products                     | 337        | 0                      | 0                            | 0                | 0                      | 0                        |
| Miscellaneous                                      | 339        | 0                      | 0                            | 0                | 0                      | 0                        |
| <b>Total (Trillion Btu)</b>                        |            | <b>379</b>             | <b>20</b>                    | <b>1,349</b>     | <b>432</b>             | <b>201</b>               |
| Average C Content (MMT/QBtu)                       |            | 18.14                  | 20.62                        | 17.51            | 0                      | 19.37                    |
| Fraction Oxidized                                  |            | 1                      | 1                            | 1                | 0                      | 1                        |
| <b>Total C (MMT)</b>                               |            | <b>6.88</b>            | <b>0.41</b>                  | <b>23.62</b>     | <b>0.00</b>            | <b>3.90</b>              |
| <b>Total C (MMT) (ex. still gas from refining)</b> |            | <b>6.88</b>            | <b>0.41</b>                  | <b>NA</b>        | <b>NA</b>              | <b>3.90</b>              |

NA (Not Applicable)

<sup>a</sup> C content: Waste Gas is assumed to be same as naphtha <401 deg. F.

<sup>b</sup> C content: Waste Oils/Tars is assumed to be same as asphalt/road oil.

<sup>c</sup> Refinery "still gas" fuel consumption is reported elsewhere in the Inventory and is excluded from the total C content estimate.

<sup>d</sup> Net steam fuel consumption is reported elsewhere in the Inventory and is excluded from the total C content estimate.

<sup>e</sup> C content: "Other" is assumed to be the same as petrochemical feedstocks.

**Table A-71: Carbon Emitted from Fuels Burned for Energy Recovery (MMT CO<sub>2</sub> Eq.)**

|             | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------|------|------|------|------|------|------|------|------|------|
| C Emissions | 41.9 | 41.5 | 65.2 | 73.3 | 41.0 | 41.0 | 41.0 | 41.0 | 41.0 |

## Products

More C is found in products than in industrial releases or energy recovery. The principal types of products are plastics; synthetic rubber; synthetic fiber; C black; pesticides; soaps, detergents, and cleansers; food additives; antifreeze and deicers (glycols); silicones; and solvents. Solvent evaporation was discussed previously along with industrial releases of NMVOCs; the other product types are discussed below.

### Plastics

Data on annual production of plastics through 2005 were taken from the American Plastics Council (APC), as published in *Chemical & Engineering News* and on the APC and Society of Plastics Industry (SPI) websites, and through direct communication with the APC (APC 2000, 2001, 2003 through 2006; SPI 2000; Eldredge-Roebuck 2000). Data for 2006 through 2014 were taken directly or derived from the American Chemistry Council (ACC 2007 through 2015 supplemented by Vallianos 2011, 2012, 2013, 2014, 2015). In 2009, the American Chemistry Council consolidated the resin categories for which it reports plastics production. Production numbers in the original categories were provided via personal correspondence for 2009, 2011, 2012, 2013, and 2014 (Vallianos 2011, 2012, 2013, 2014, 2015). Production figures for the consolidated resin categories in 2010 were linearly interpolated from 2009 and 2011 data. Production was organized by resin type (see Table A-72) and by year.

Several of the resin categories included production from Canada and/or Mexico, in addition to the U.S. values for part of the time series. The production data for the affected resins and years were corrected using an economic adjustment factor, based on the percent of North American production value in this industry sector accounted for by the United States. A C content was then assigned for each resin. These C contents were based on molecular formulae and are listed in Table A-73 and Table A-74. In cases where the resin type is generic, referring to a group of chemicals and not a single polymer (e.g., phenolic resins, other styrenic resins), a representative compound was chosen. For other resins, a weighted C content of 68 percent was assumed (i.e., it was assumed that these resins had the same content as those for which a representative compound could be assigned).

There were no emissive uses of plastics identified, so 100 percent of the C was considered stored in products. As noted in the chapter, an estimate of emissions related to the combustion of these plastics in the municipal solid waste stream can be found in the Incineration of Waste section of the Energy chapter; those emissions are not incorporated in the mass balance for feedstocks (described in this annex) to avoid double-counting.

**Table A-72: 2014 Plastic Resin Production (MMT dry weight) and C Stored (MMT CO<sub>2</sub> Eq.)**

| Resin Type                                    | 2014 Production <sup>a</sup><br>(MMT dry weight) | Carbon Stored<br>(MMT CO <sub>2</sub> Eq.) |
|---|--|--|
| Epoxy   | 0.3  | 0.7  |
| Urea  | 0.5  | 0.6  |
| Melamine                                      | 0.5  | 0.5  |
| Phenolic                                      | 1.6  | 4.4  |
| Low-Density Polyethylene (LDPE)               | 2.9  | 9.2  |
| Linear Low-Density Polyethylene (LLDPE)       | 5.7  | 17.9                                       |
| High Density Polyethylene (HDPE)              | 7.2  | 22.6                                       |
| Polypropylene (PP)                            | 6.3  | 19.9                                       |
| Acrylonitrile-butadiene-styrene (ABS)         | 0.4  | 1.3  |
| Other Styrenics <sup>b</sup>                  | 0.5  | 1.7  |
| Polystyrene (PS)                              | 2.2  | 7.5  |
| Nylon   | 0.5  | 1.2  |
| Polyvinyl chloride (PVC) <sup>c</sup>         | 6.2  | 8.7  |
| Thermoplastic Polyester                       | 3.4  | 7.8  |
| All Other (including Polyester (unsaturated)) | 6.1  | 15.4                                       |
| <b>Total</b>                                  | <b>44.2</b>                                      | <b>119.3</b>                               |

<sup>a</sup> Production estimates provided by the American Chemistry Council include Canadian production for Urea, Melamine, Phenolic, LDPE, LLDPE, HDPE, PP, ABS, SAN, Other Styrenics, PS, Nylon, PVC, and Thermoplastic Polyester, and Mexican production for PP, ABS, SAN, Other Styrenics, Nylon, and Thermoplastic Polyester. Values have been adjusted to account just for U.S. production.

<sup>b</sup> Includes Styrene-acrylonitrile (SAN).

<sup>c</sup> Includes copolymers.

Note: Totals may not sum due to independent rounding.

**Table A-73: Assigned C Contents of Plastic Resins (% by weight)**

| Resin Type                              | C Content | Source of C Content Assumption                               |
|---|-----------|--|
| Epoxy                                   | 76%       | Typical epoxy resin made from epichlorhydrin and bisphenol A |
| Polyester (Unsaturated)                 | 63%       | Poly (ethylene terephthalate) (PET)                          |
| Urea                                    | 34%       | 50% carbamal, 50% N-(hydroxymethyl) urea *                   |
| Melamine                                | 29%       | Trimethylol melamine *                                       |
| Phenolic                                | 77%       | Phenol   |
| Low-Density Polyethylene (LDPE)         | 86%       | Polyethylene   |
| Linear Low-Density Polyethylene (LLDPE) | 86%       | Polyethylene   |
| High Density Polyethylene (HDPE)        | 86%       | Polyethylene   |
| Polypropylene (PP)                      | 86%       | Polypropylene  |
| Acrylonitrile-Butadiene-Styrene (ABS)   | 85%       | 50% styrene, 25% acrylonitrile, 25% butadiene                |
| Styrene-Acrylonitrile (SAN)             | 80%       | 50% styrene, 50% acrylonitrile                               |
| Other Styrenics                         | 92%       | Polystyrene  |
| Polystyrene (PS)                        | 92%       | Polystyrene  |
| Nylon                                   | 65%       | Average of nylon resins (see Table A-74 )                    |
| Polyvinyl Chloride (PVC)                | 38%       | Polyvinyl chloride   |
| Thermoplastic Polyester                 | 63%       | Polyethylene terephthalate                                   |
| All Other                               | 69%       | Weighted average of other resin production                   |

\*Does not include alcoholic hydrogens.

**Table A-74: Major Nylon Resins and their C Contents (% by weight)**

| Resin      | C Content |
|------------|-----------|
| Nylon 6    | 64%       |
| Nylon 6,6  | 64%       |
| Nylon 4    | 52%       |
| Nylon 6,10 | 68%       |
| Nylon 6,11 | 69%       |
| Nylon 6,12 | 70%       |
| Nylon 11   | 72%       |

#### Synthetic Rubber

Data on synthetic rubber in tires were derived from data on the scrap tire market and the composition of scrap tires from the Rubber Manufacturers' Association (RMA). The market information is presented in the report *2013 U.S. Scrap Tire Management Summary* (RMA 2014), while the tire composition information is from the "Scrap Tires, Facts and Figures" section of the organization's website (RMA 2009). The most recent data available were for 2013, so it was assumed that the 2014 consumption was equal to that of 2013. Data on synthetic rubber in other products (durable goods, nondurable goods, and containers and packaging) were obtained from EPA's *Municipal Solid Waste in the United States* reports (1996 through 2003a, 2005, 2007b, 2008, 2009a, 2011a, 2013b; 2014) and detailed unpublished backup data for some years not shown in the *Characterization of Municipal Solid Waste in the United States* reports (Schneider 2007). The abraded rubber from scrap passenger tires was assumed to be 2.5 pounds per scrap tire, while the abraded rubber from scrap commercial tires was assumed to be 10 pounds per scrap tire. Data on abraded rubber weight were obtained by calculating the average weight difference between new and scrap tires (RMA 2014). Import and export data were obtained from the published by the U.S. International Trade Commission (U.S. International Trade Commission 2015).

A C content for synthetic rubber (90 percent for tire synthetic rubber and 85 percent for non-tire synthetic rubber) was assigned based on the weighted average of C contents (based on molecular formula) by elastomer type consumed in 1998, 2001, and 2002 (see Table A-75). The 1998 consumption data were obtained from the International Institute of Synthetic Rubber Producers (IISRP) press release "Synthetic Rubber Use Growth to Continue Through 2004, Says IISRP and RMA" (IISRP 2000). The 2001 and 2002 consumption data were obtained from the IISRP press release, "IISRP Forecasts Moderate Growth in North America to 2007" (IISRP 2003).

The rubber in tires that is abraded during use (the difference between new tire and scrap tire rubber weight) was considered to be 100 percent emitted. Other than abraded rubber, there were no emissive uses of scrap tire and non-tire rubber identified, so 100 percent of the non-abraded amount was assumed stored. Emissions related to the combustion of rubber in scrap tires and consumer goods can be found in the Incineration of Waste section of the Energy chapter.

**Table A-75: 2002 Rubber Consumption (kt) and C Content (%)**

| Elastomer Type          | 2002 Consumption (kt) <sup>a</sup> | C Content  |
|-------------------------|------------------------------------|------------|
| SBR Solid               | 768                                | 91%        |
| Polybutadiene           | 583                                | 89%        |
| Ethylene Propylene      | 301                                | 86%        |
| Polychloroprene         | 54                                 | 59%        |
| NBR Solid               | 84                                 | 77%        |
| Polyisoprene            | 58                                 | 88%        |
| Others                  | 367                                | 88%        |
| <b>Weighted Average</b> | <b>NA</b>                          | <b>90%</b> |
| <b>Total</b>            | <b>2,215</b>                       | <b>NA</b>  |

<sup>a</sup> Includes consumption in Canada.

NA (Not Applicable)

Note: Totals may not sum due to independent rounding.

### Synthetic Fibers

Annual synthetic fiber production data were obtained from the Fiber Economics Bureau, as published in *Chemical & Engineering News* (FEB 2001, 2003, 2005, 2007, 2009, 2010, 2011, 2012, 2013). The most recent data available were for 2012, so it was assumed that the 2013 and 2014 consumption was equal to that of 2012. These data are organized by year and fiber type. For each fiber, a C content was assigned based on molecular formula (see Table A-76). For polyester, the C content for poly (ethylene terephthalate) (PET) was used as a representative compound. For nylon, the average C content of nylon 6 and nylon 6.6 was used, since these are the most widely produced nylon fibers. Cellulosic fibers, such as acetate and rayon, have been omitted from the synthetic fibers' C accounting displayed here because much of their C is of biogenic origin and carbon fluxes from biogenic compounds are accounted for in the Land Use, Land-Use Change and Forestry chapter. These fibers account for only 4 percent of overall fiber production by weight.

There were no emissive uses of fibers identified, so 100 percent of the C was considered stored. Note that emissions related to the combustion of textiles in municipal solid waste are accounted for under the Incineration of Waste section of the Energy chapter.

**Table A-76: 2014 Fiber Production (MMT), C Content (%), and C Stored (MMT CO<sub>2</sub> Eq.)**

| Fiber Type   | Production (MMT) | C Content | C Stored (MMT CO <sub>2</sub> Eq.) |
|--------------|------------------|-----------|------------------------------------|
| Polyester    | 1.2              | 63%       | 2.8                                |
| Nylon        | 0.6              | 64%       | 1.3                                |
| Olefin       | 1.0              | 86%       | 3.2                                |
| Acrylic      | +                | 68%       | 0.1                                |
| <b>Total</b> | <b>2.8</b>       | <b>NA</b> | <b>7.3</b>                         |

+ Less than 0.05 MMT

NA (Not Applicable)

Note: Totals may not sum due to independent rounding.

### Pesticides

Pesticide consumption data were obtained from the *1994/1995, 1996/1997, 1998/1999, 2000/2001, 2006/2007 Pesticides Industry Sales and Usage Market Estimates* (EPA 1998, 1999, 2002, 2004, 2011b) reports. The most recent data available were for 2007, so it was assumed that the 2008 through 2014 consumption was equal to that of 2007. Active ingredient compound names and consumption weights were available for the top 25 agriculturally-used pesticides and top 10 pesticides used in the home and garden and the industry/commercial/government categories. The report provides a range of consumption for each active ingredient; the midpoint was used to represent actual consumption. Each of these compounds was assigned a C content value based on molecular formula. If the compound contained aromatic rings substituted with chlorine or other halogens, then the compound was considered persistent and the C in the compound was assumed to be stored. All other pesticides were assumed to release their C to the atmosphere. Over one-third of 2007 total pesticide active ingredient consumption was not specified by chemical type in the *Sales and Usage* report (EPA 2011b). This unspecified portion of the active ingredient consumption was treated as a single chemical and assigned a C content and a storage factor based on the weighted average of the known chemicals' values.

**Table A-77: Active Ingredient Consumption in Pesticides (Million lbs.) and C Emitted and Stored (MMT CO<sub>2</sub> Eq.) in 2007**

| Pesticide Use <sup>a</sup> | Active Ingredient<br>(Million lbs.) | C Emitted<br>(MMT CO <sub>2</sub> Eq.) | C Stored<br>(MMT CO <sub>2</sub> Eq.) |
|----------------------------|-------------------------------------|--|---------------------------------------|
| Agricultural Uses          | 473.5                               | 0.1                                    | 0.1                                   |
| Non-Agricultural Uses      | 76.8                                | +                                      | +                                     |
| Home & Garden              | 30.3                                | +                                      | +                                     |
| Industry/Gov't/Commercial  | 46.5                                | +                                      | +                                     |
| Other                      | 337.7                               | 0.1                                    | 0.1                                   |
| <b>Total</b>               | <b>888.0</b>                        | <b>0.2</b>                             | <b>0.2</b>                            |

+ Less than 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> 2007 estimates (EPA 2011b).

Note: Totals may not sum due to independent rounding.

### *Soaps, Shampoos, and Detergents*

Cleansers—soaps, shampoos, and detergents—are among the major consumer products that may contain fossil C. All of the C in cleansers was assumed to be fossil-derived, and, as cleansers eventually biodegrade, all of the C was assumed to be emitted. The first step in estimating C flows was to characterize the “ingredients” in a sample of cleansers. For this analysis, cleansers were limited to the following personal household cleaning products: bar soap, shampoo, laundry detergent (liquid and granular), dishwasher detergent, and dishwashing liquid. Data on the annual consumption of household personal cleansers were obtained from the U.S. Census Bureau 1992, 1997, 2002, 2007, 2012 Economic Census (U.S. Bureau of the Census 1994, 1999, 2004, 2009, 2014). Consumption values for 1990 and 1991 were assumed to be the same as the 1992 value; consumption was interpolated between 1992 and 1997, 1997 and 2002, and 2002 and 2007; consumption for 2008 through 2014 was assumed to equal the 2007 value. Cleanser consumption values were adjusted by import and export data to develop U.S. consumption estimates.

Chemical formulae were used to determine C contents (as percentages) of the ingredients in the cleansers. Each product’s overall C content was then derived from the composition and contents of its ingredients. From these values the mean C content for cleansers was calculated to be 21.9 percent.

The Census Bureau presents consumption data in terms of quantity (in units of million gallons or million pounds) and/or terms of value (thousands of dollars) for eight specific categories, such as “household liquid laundry detergents, heavy duty” and “household dry alkaline automatic dishwashing detergents.” Additionally, the report provides dollar values for the total consumption of “soaps, detergents, etc.—dry” and “soaps, detergents, etc.—liquid.” The categories for which both quantity and value data are available is a subset of total production. Those categories that presented both quantity and value data were used to derive pounds per dollar and gallons per dollar conversion rates, and they were extrapolated (based on the Census Bureau estimate of total value) to estimate the total quantity of dry and liquid<sup>32</sup> cleanser categories, respectively.

Next, the total tonnage of cleansers was calculated (wet and dry combined) for 1997. Multiplying the mean C content (21.9 percent) by this value yielded an estimate of 4.6 MMT CO<sub>2</sub> Eq. in cleansers for 1997. For all subsequent years, it was assumed that the ratio of value of shipments to total carbon content remained constant. For 1998 through 2014, value of shipments was adjusted to 1997 dollars using the producer price index for soap and other detergent manufacturing (Bureau of Labor Statistics 2014). The ratio of value of shipments to carbon content was then applied to arrive at total carbon content of cleansers. Estimates are shown in Table A-78.

**Table A-78: C Emitted from Utilization of Soaps, Shampoos, and Detergents (MMT CO<sub>2</sub> Eq.)**

|             | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------|------|------|------|------|------|------|------|------|------|
| C Emissions | 3.6  | 4.2  | 4.5  | 6.7  | 5.8  | 5.4  | 4.7  | 4.7  | 4.8  |

### *Antifreeze and Deicers*

Glycol compounds, including ethylene glycol, propylene glycol, diethylene glycol, and triethylene glycol, are used as antifreeze in motor vehicles, deicing fluids for commercial aircraft, and other similar uses. These glycol compounds are assumed to ultimately enter wastewater treatment plants where they are degraded by the wastewater treatment process to CO<sub>2</sub> or to otherwise biodegrade to CO<sub>2</sub>. Glycols are water soluble and degrade rapidly in the environment (Howard 1993).

<sup>32</sup> A density of 1.05 g/mL—slightly denser than water—was assumed for liquid cleansers.

Annual production data for each glycol compound used as antifreeze and deicers were obtained from the *Guide to the Business of Chemistry* (ACC 2015b) and the EPA Chemical Data Access Tool (CDAT) (EPA 2014). Import and export data were used to adjust annual production data to annual consumption data. The percentage of the annual consumption of each glycol compound used for antifreeze and deicing applications was estimated from Chemical Profiles data published on The Innovation Group website<sup>33</sup> and from similar data published in the Chemical Market Reporter, which became ICIS Chemical Business in 2005.<sup>34</sup> Production data for propylene glycol, diethylene glycol, and triethylene glycol are no longer reported in the Guide to the Business of Chemistry, so data from ICIS Chemical Business on total demand was used with import and export data to estimate production of these chemicals. ICIS last reported total demand for propylene glycol and diethylene glycol in 2006, and triethylene glycol demand in 2007. EPA reported total U.S. production of propylene glycol, diethylene glycol, and triethylene glycol in 2012 in the CDAT (EPA 2014). Total demand for these compounds for 2012 was calculated from the 2012 production data using import and export data. Demand for propylene glycol and diethylene glycol was interpolated for years between 2006 and 2012, and demand for triethylene glycol was interpolated for years between 2007 and 2012, using the calculated 2012 total demand values for each compound and the most recently reported total demand data from ICIS. Values for 2013 and 2014 for these compounds were proxied from the 2012 values. Production data for ethylene glycol was provided by the 2015 *Guide to the Business of Chemistry* (ACC 2015b).

The glycol compounds consumed in antifreeze and deicing applications is assumed to be 100 percent emitted as CO<sub>2</sub>. Emissions of CO<sub>2</sub> from utilization of antifreeze and deicers are summarized in Table A-79.

**Table A-79: C Emitted from Utilization of Antifreeze and Deicers (MMT CO<sub>2</sub> Eq.)**

|             | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------|------|------|------|------|------|------|------|------|------|
| C Emissions | 1.2  | 1.4  | 1.5  | 1.2  | 0.9  | 0.7  | 0.9  | 0.8  | 0.8  |

#### *Food Additives*

Petrochemical feedstocks are used to manufacture synthetic food additives, including preservatives, flavoring agents, and processing agents. These compounds include glycerin, propylene glycol, benzoic acid, and other compounds. These compounds are incorporated into food products, and are assumed to ultimately enter wastewater treatment plants where they are degraded by the wastewater treatment processes to CO<sub>2</sub> or to otherwise biodegrade to CO<sub>2</sub>. Certain food additives, e.g., glycerin, are manufactured both from petrochemical feedstocks and from biogenic feedstocks. Food additives that are derived from biogenic feedstocks are accounted for in the Land Use, Land-Use Change and Forestry chapter.

Annual production data for food additive compounds were obtained from the *Guide to the Business of Chemistry* (ACC 2015b). Historical values for adipic acid were adjusted according to the most recent data in the 2015 *Guide to the Business of Chemistry*. Import and export data were used to adjust annual production data to annual consumption data. The percentage of the annual consumption of food additive compounds was estimated from Chemical Profiles data published on The Innovation Group website<sup>35</sup> and from similar data published in the Chemical Market Reporter, which became ICIS Chemical Business in 2005.<sup>36</sup> Production data for several food additive compounds are no longer reported in the *Guide to the Business of Chemistry*, so data from ICIS Chemical Business on total demand was used with import and export data to estimate production of these chemicals.

ICIS last reported total demand for glycerin and benzoic acid in 2007, and demand for propionic acid in 2008. Total demand for acetic acid and maleic anhydride were last reported by ICIS in 2005, and dipropylene glycol demand in 2004. ICIS last reported cresylic acid demand in 1999. EPA reported total U.S. production of these compounds in 2012 in the CDAT (EPA 2014). Total demand for these compounds for 2012 was calculated from the 2012 production data using import and export data. Demand for each of these compounds was interpolated for years between the most recently reported total demand data from ICIS and 2012, using the calculated 2012 total demand values for each compound. Values for 2013 and 2014 for these compounds were proxied from the 2012 values.

The consumption of synthetic food additives is assumed to be 100 percent emitted as CO<sub>2</sub>. Emissions of CO<sub>2</sub> from utilization of synthetic food additives are summarized in Table A-80.

<sup>33</sup> See <<http://www.the-innovation-group.com/ChemProfiles>>.

<sup>34</sup> See <<http://www.icis.com/home/default.aspx>>.

<sup>35</sup> See <<http://www.the-innovation-group.com/ChemProfiles>>.

<sup>36</sup> See <<http://www.icis.com/home/default.aspx>>.

**Table A-80: C Emitted from Utilization of Food Additives (MMT CO<sub>2</sub> Eq.)**

|             | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------|------|------|------|------|------|------|------|------|------|
| C Emissions | 0.6  | 0.7  | 0.7  | 0.8  | 0.9  | 1.0  | 1.1  | 1.1  | 1.1  |

### Silicones

Silicone compounds (e.g., polymethyl siloxane) are used as sealants and in manufactured products. Silicone compounds are manufactured from petrochemical feedstocks including methyl chloride. It is assumed that petrochemical feedstocks used to manufacture silicones are incorporated into the silicone products and not emitted as CO<sub>2</sub> in the manufacturing process. It is also assumed that the C contained in the silicone products is stored, and not emitted as CO<sub>2</sub>.

Annual production data for each silicone manufacturing compound were obtained from the Guide to the Business of Chemistry (ACC 2015b). Import and export data were used to adjust annual production data to annual consumption data. The percentage of the annual consumption of each silicone manufacturing compound was estimated from Chemical Profiles data published on The Innovation Group website and from similar data published in the Chemical Market Reporter, which became ICIS Chemical Business in 2005.<sup>37</sup> ICIS last reported production of methyl chloride in 2007. EPA reported total U.S. production of methyl chloride in 2012 in the CDAT (EPA 2014). Total consumption of methyl chloride for 2012 was calculated from the 2012 production data using import and export data. Production of methyl chloride was interpolated for years between 2007 and 2012, using the calculated 2012 total production value for methyl chloride and the most recently reported total production data from ICIS. The production value for 2013 and 2014 was proxied from the 2012 value.

The consumption of silicone manufacturing compounds is assumed to be 100 percent stored, and not emitted as CO<sub>2</sub>. Storage of silicone manufacturing compounds is summarized in Table A-81.

**Table A-81: C Stored in Silicone Products (MMT CO<sub>2</sub> Eq.)**

|           | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-----------|------|------|------|------|------|------|------|------|------|
| C Storage | 0.3  | 0.4  | 0.4  | 0.4  | 0.5  | 0.5  | 0.5  | 0.5  | 0.5  |

### Uncertainty

A Tier 2 Monte Carlo analysis was performed using @RISK software to determine the level of uncertainty surrounding the estimates of the feedstocks C storage factor and the quantity of C emitted from feedstocks in 2014. The Tier 2 analysis was performed to allow the specification of probability density functions for key variables, within a computational structure that mirrors the calculation of the Inventory estimate. Statistical analyses or expert judgments of uncertainty were not available directly from the information sources for the activity variables; thus, uncertainty estimates were determined using assumptions based on source category knowledge. Uncertainty estimates for production data (the majority of the variables) were assumed to exhibit a normal distribution with a relative error of ±20 percent in the underlying EIA estimates, plus an additional ±15 percent to account for uncertainty in the assignment of imports and exports. An additional 10 percent (for a total of ±45 percent) was applied to the production of other oils (>401 degrees Fahrenheit) to reflect the additional uncertainty in the assignment of part of the production quantity to industrial processes. A relatively narrow uniform distribution ±1 percent to ±15 percent, depending on the fuel type) was applied to each C coefficient.

The Monte Carlo analysis produced a storage factor distribution with a mean of 65 percent, a standard deviation of 5.8 percent, and the 95 percent confidence interval of 52 percent and 72 percent. This compares to the calculated Inventory estimate of 65 percent. The analysis produced a C emission distribution with a mean of 75.1 MMT CO<sub>2</sub> Eq., standard deviation of 21.3 MMT CO<sub>2</sub> Eq., and 95 percent confidence limits of 49.6 and 125.3 MMT CO<sub>2</sub> Eq. This compares with a calculated Inventory estimate of 75.1 MMT CO<sub>2</sub> Eq.

The apparently tight confidence limits for the storage factor and C storage probably understate uncertainty, as a result of the way this initial analysis was structured. As discussed above, the storage factor for feedstocks is based on an analysis of six fates that result in long-term storage (e.g., plastics production), and eleven that result in emissions (e.g., volatile organic compound emissions). Rather than modeling the total uncertainty around all 17 of these fate processes, the current analysis addresses only the storage fates, and assumes that all C that is not stored is emitted. As the production statistics that drive the storage factors are relatively well-characterized, this approach yields a result that is probably biased toward understating uncertainty.

As far as specific sources of uncertainty, there are several cross-cutting factors that pervade the characterization of C flows for feedstocks. The aggregate storage factor for petrochemical feedstocks (industrial other coal, natural gas for non-

<sup>37</sup> Ibid.

fertilizer uses, LPG, pentanes plus, naphthas, other oils, still gas, special naphtha) is based on assuming that the ultimate fates of all of these fuel types—in terms of storage and emissions—are similar. In addition, there are uncertainties associated with the simplifying assumptions made for each end use category C estimate. Generally, the estimate for a product is subject to one or more of the following uncertainties:

- The value used for estimating the C content has been assumed or assigned based upon a representative compound.
- The split between C storage and emission has been assumed based on an examination of the environmental fate of the products in each end use category.
- Environmental fates leading to emissions are assumed to operate rapidly, i.e., emissions are assumed to occur within one year of when the fossil C enters the non-energy mass balance. Some of the pathways that lead to emissions as CO<sub>2</sub> may actually take place on a time-scale of several years or decades. By attributing the emissions to the year in which the C enters the mass balance (i.e., the year in which it leaves refineries as a non-energy fuel use and thus starts being tracked by EIA), this approach has the effect of “front-end loading” the emission profile.

Another cross-cutting source of uncertainty is that for several sources the amount of C stored or emitted was calculated based on data for only a single year. This specific year may not be representative of storage for the entire Inventory period. Sources of uncertainty associated with specific elements of the analysis are discussed below.

Import and export data for petrochemical feedstocks were obtained from EIA, the National Petroleum Refiners Association, and the BoC for the major categories of petrochemical feedstocks (EIA 2001; NPRA 2001; and U.S. Bureau of the Census 2015). The complexity of the organic chemical industry, with multiple feedstocks, intermediates, and subtle differences in nomenclature, makes it difficult to ensure that the adjustments to the EIA data for imports and exports is accurate and the approach used here may underestimate or overestimate net exports of C.

Oxidation factors have been applied to non-energy uses of petrochemical feedstocks in the same manner as for energy uses. However, for those fuels where IPCC storage factors are used, this “oxidation factor” may be inherent in the storage factor applied when calculating emissions from non-energy consumption, which would result in a double-counting of the unoxidized C. Oxidation factors are small corrections, on the order of 1 percent, and therefore application of oxidation factors to non-energy uses may result in a slight underestimation of C emissions from non-energy uses.

The major uncertainty in using the TRI data is the possibility of double counting emissions that are already accounted for in the NMVOC data (see above) and in the storage and emission assumptions used. The approach for predicting environmental fate simplifies some complex processes, and the balance between storage and emissions is very sensitive to the assumptions on fate. Extrapolating from known to unknown characteristics also introduces uncertainty. The two extrapolations with the greatest uncertainty are: (1) that the release media and fate of the off-site releases were assumed to be the same as for on-site releases, and (2) that the C content of the least frequent 10 percent of TRI releases was assumed to be the same as for the chemicals comprising 90 percent of the releases. However, the contribution of these chemicals to the overall estimate is small. The off-site releases only account for 3 percent of the total releases, by weight, and, by definition, the less frequent compounds only account for 10 percent of the total releases.

The principal sources of uncertainty in estimating CO<sub>2</sub> emissions from solvent evaporation and industrial NMVOC emissions are in the estimates of (a) total emissions and (b) their C content. Solvent evaporation and industrial NMVOC emissions reported by EPA are based on a number of data sources and emission factors, and may underestimate or overestimate emissions. The C content for solvent evaporation emissions is calculated directly from the specific solvent compounds identified by EPA as being emitted, and is thought to have relatively low uncertainty. The C content for industrial emissions has more uncertainty, however, as it is calculated from the average C content of an average volatile organic compound based on the list of the most abundant measured NMVOCs provided in EPA (2002a).

Uncertainty in the hazardous waste combustion analysis is introduced by the assumptions about the composition of combusted hazardous wastes, including the characterization that hazardous wastes are similar to mixtures of water, noncombustibles, and fuel equivalent materials. Another limitation is the assumption that all of the C that enters hazardous waste combustion is emitted—some small fraction is likely to be sequestered in combustion ash—but given that the destruction and removal efficiency for hazardous organics is required to meet or exceed 99.99 percent, this is a very minor source of uncertainty. C emission estimates from hazardous waste should be considered central value estimates that are likely to be accurate to within ±50 percent.

The amount of feedstocks combusted for energy recovery was estimated from data included in the *Manufacturers Energy Consumption Surveys* (MECS) for 1991, 1994, 1998, 2002, 2006, and 2010 (EIA 1994, 1997, 2001, 2005, 2010, 2013b). MECS is a comprehensive survey that is conducted every four years and intended to represent U.S. industry as a whole, but because EIA does not receive data from all manufacturers (i.e., it is a sample rather than a census), EIA must

extrapolate from the sample. Also, the “other” fuels are identified in the MECS data in broad categories, including refinery still gas; waste gas; waste oils, tars, and related materials; petroleum coke, coke oven and blast furnace gases; and other uncharacterized fuels. Moreover, the industries using these “other” fuels are also identified only in broad categories, including the petroleum and coal products, chemicals, primary metals, nonmetallic minerals, and other manufacturing sectors. The “other” fuel consumption data are reported in BTUs (energy units) and there is uncertainty concerning the selection of a specific conversion factor for each broad “other” fuel category to convert energy units to mass units. Taken as a whole, the estimate of energy recovery emissions probably introduces more uncertainty than any other element of the non-energy analysis.

Uncertainty in the C storage estimate for plastics arises primarily from four factors. First, production of some plastic resins is not tracked directly and must be estimated based on other market data. Second, the raw data on production for several resins include Canadian and/or Mexican production and may overestimate the amount of plastic produced from U.S. fuel feedstocks; this analysis includes adjustments to “back out” the Canadian and Mexican values, but these adjustments are approximate. Third, the assumed C content values are estimates for representative compounds, and thus do not account for the many formulations of resins available. This uncertainty is greater for resin categories that are generic (e.g., phenolics, other styrenics, nylon) than for resins with more specific formulations (e.g., polypropylene, polyethylene). Fourth, the assumption that all of the C contained in plastics is stored ignores certain end uses (e.g., adhesives and coatings) where the resin may be released to the atmosphere; however, these end-uses are likely to be small relative to use in plastics.

The quantity of C stored in synthetic rubber only accounts for the C stored in scrap tire synthetic rubber. The value does not take into account the rubber stored in other durable goods, clothing, footwear, and other non-durable goods, or containers and packaging. This adds uncertainty to the total mass balance of C stored. There are also uncertainties as to the assignment of C content values; however, they are much smaller than in the case of plastics. There are probably fewer variations in rubber formulations than in plastics, and the range of potential C content values is much narrower. Lastly, assuming that all of the C contained in rubber is stored ignores the possibility of volatilization or degradation during product lifetimes. However, the proportion of the total C that is released to the atmosphere during use is probably negligible.

A small degree of uncertainty arises from the assignment of C content values in textiles; however, the magnitude of this uncertainty is less than that for plastics or rubber. Although there is considerable variation in final textile products, the stock fiber formulations are standardized and proscribed explicitly by the Federal Trade Commission.

For pesticides, the largest source of uncertainty involves the assumption that an active ingredient’s C is either zero percent stored or 100 percent stored. This split is a generalization of chemical behavior, based upon active-ingredient molecular structure, and not on compound-specific environmental data. The mechanism by which a compound is bound or released from soils is very complicated and can be affected by many variables, including the type of crop, temperature, application method, and harvesting practice. Another smaller source of uncertainty arises from the C content values applied to the unaccounted for portion of active ingredient. C contents vary widely among pesticides, from 7 to 72 percent, and the remaining pesticides may have a chemical make-up that is very different from the 32 pesticides that have been examined. Additionally, pesticide consumption data were only available for 1987, 1993, 1995, 1997, 1999, 2001, and 2007; the majority of the time series data were interpolated or held constant at the latest (2007) value. Another source of uncertainty is that only the “active” ingredients of pesticides are considered in the calculations; the “inactive” ingredients may also be derived from petrochemical feedstocks.

It is important to note that development of this uncertainty analysis is a multi-year process. The current feedstocks analysis examines NEU fuels that end in storage fates. Thus, only C stored in pesticides, plastics, synthetic fibers, synthetic rubbers, silicones, and TRI releases to underground injection and Subtitle C landfills is accounted for in the uncertainty estimate above. In the future this analysis will be expanded to include the uncertainty surrounding emitted fates in addition to the storage fates. Estimates of variable uncertainty will also be refined where possible to include fewer assumptions. With these major changes in future Inventories, the uncertainty estimate is expected to change, and likely increase. An increase in the uncertainty estimate in the coming years will not indicate that the Inventory calculations have become less certain, but rather that the methods for estimating uncertainty have become more comprehensive; thus, potential future changes in the results of this analysis will reflect a change in the uncertainty analysis, not a change in the Inventory quality.

## **Asphalt and Road Oil**

Asphalt is one of the principal non-energy uses of fossil fuels. The term “asphalt” generally refers to a mixture of asphalt cement and a rock material aggregate, a volatile petroleum distillate, or water. For the purposes of this analysis, “asphalt” is used interchangeably with asphalt cement, a residue of crude oil. Though minor amounts of C are emitted during production, asphalt has an overall C storage factor of almost 100 percent, as discussed below.

Paving is the primary application of asphalt cement, comprising 86 percent of production. The three types of asphalt paving produced in the United States are hot mix asphalt (HMA), cut-backs, and emulsified asphalt. HMA, which makes up 90 percent of total asphalt paving (EPA 2001), contains asphalt cement mixed with an aggregate of rock materials. Cut-back asphalt is composed of asphalt cement thinned with a volatile petroleum distillate (e.g., naphtha). Emulsified asphalt contains only asphalt cement and water. Roofing products are the other significant end use of asphalt cement, accounting for approximately 14 percent of U.S. production (Kelly 2000). No data were available on the fate of C in asphalt roofing; it was assumed that it has the same fate as C in asphalt paving applications.

### Methodology and Data Sources

A C storage factor was calculated for each type of asphalt paving. The fraction of C emitted by each asphalt type was multiplied by consumption data for asphalt paving (EPA 2001) to estimate a weighted average C storage factor for asphalt as a whole.

The fraction of C emitted by HMA was determined by first calculating the organic emissions (volatile organic compounds [VOCs], carbon monoxide [CO], polycyclic aromatic hydrocarbons [PAHs], hazardous air pollutants [HAPs], and phenol) from HMA paving, using emission factors reported in EPA (2001) and total HMA production.<sup>38</sup> The next step was to estimate the C content of the organic emissions. This calculation was based on the C content of CO and phenol, and an assumption of 85 percent C content for PAHs and HAPs. The C content of asphalt paving is a function of (1) the proportion of asphalt cement in asphalt paving, assumed to be 8 percent asphalt cement content based on EPA (2001), and (2) the proportion of C in asphalt cement. For the latter factor, all paving types were characterized as having a mass fraction of 85 percent C in asphalt cement, based on the assumption that asphalt is primarily composed of saturated paraffinic hydrocarbons. By combining these estimates, the result is that over 99.56 percent of the C in asphalt cement was retained (i.e., stored), and less than 0.44 percent was emitted.

Cut-back asphalt is produced in three forms: rapid, medium, and slow cure. The production processes for all three forms emit C primarily from the volatile petroleum distillate used in the process as a diluent to thin the asphalt cement so that it can be applied more readily (EPA 2001).

A mass balance on C losses from asphalt was constructed by first estimating the amount of carbon emitted as VOCs. Values for medium cure asphalt are used to represent all cut-back asphalt. The average weight of distillates used in medium cure cut-back asphalt (35 percent) is multiplied by the loss rate (as emissions of VOCs) of 70 percent from the *Emissions Inventory Guidebook* to arrive at an estimate that 25 percent of the diluent is emitted (Environment Canada 2006). Next, the fraction of C in the asphalt/ diluent mix that is emitted was estimated, assuming 85 percent C content; this yields an overall storage factor of 93.5 percent for cut-back asphalt.

One caveat associated with this calculation is that it is possible that the carbon flows for asphalt and diluent (volatile petroleum distillate) are accounted for separately in the EIA statistics on fossil fuel flows, and thus the mass balance calculation may need to re-map the system boundaries to correctly account for carbon flows. EPA plans to re-evaluate this calculation in the future.

It was assumed that there was no loss of C from emulsified asphalt (i.e., the storage factor is 100 percent) based on personal communication with an expert from Akzo Nobel Coatings, Inc. (James 2000).

Data on asphalt and road oil consumption and C content factors were supplied by EIA. Hot mix asphalt production and emissions factors, and the asphalt cement content of HMA were obtained from "Hot Mix Asphalt Plants Emissions Assessment Report" from EPA's *AP-42* (EPA 2001) publication. The consumption data for cut-back and emulsified asphalts were taken from a Moulthrop, et al. study used as guidance for estimating air pollutant emissions from paving processes (EIIP 2001). "Asphalt Paving Operation" *AP-42* (EPA 2001) provided the emissions source information used in the calculation of the C storage factor for cut-back asphalt. The storage factor for emulsified asphalt was provided by Alan James of Akzo Nobel Coatings, Inc. (James 2000).

### Uncertainty

A Tier 2 Monte Carlo analysis was performed using @RISK software to determine the level of uncertainty surrounding the estimates of the asphalt C storage factor and the quantity of C stored in asphalt in 2014. The Tier 2 analysis was performed to allow the specification of probability density functions for key variables, within a computational structure that mirrors the calculation of the Inventory estimate. Statistical analyses or expert judgments of uncertainty were not

---

<sup>38</sup> The emission factors are expressed as a function of asphalt paving tonnage (i.e., including the rock aggregate as well as the asphalt cement).

available directly from the information sources for the activity variables; thus, uncertainty estimates were determined using assumptions based on source category knowledge. Uncertainty estimates for asphalt production were assumed to be  $\pm 20$  percent, while the asphalt property variables were assumed to have narrower distributions. A narrow uniform distribution, with maximum 5 percent uncertainty ( $\pm 5$  percent) around the mean, was applied to the C content coefficient.

The Monte Carlo analysis produced a tight distribution of storage factor values, with the 95 percent confidence interval of 99 percent and 100 percent, with the mean value of 99 percent. This compares to the storage factor value used in the Inventory of 99.6 percent. The analysis produced a C emission distribution with a mean of 0.26 MMT CO<sub>2</sub> Eq., standard deviation of 0.12 and 95 percent confidence limits of 0.11 MMT CO<sub>2</sub> Eq. and 0.57 MMT CO<sub>2</sub> Eq. This compares to an Inventory calculated estimate of 0.26 MMT CO<sub>2</sub> Eq.

The principal source of uncertainty is that the available data are from short-term studies of emissions associated with the production and application of asphalt. As a practical matter, the cement in asphalt deteriorates over time, contributing to the need for periodic re-paving. Whether this deterioration is due to physical erosion of the cement and continued storage of C in a refractory form or physicochemical degradation and eventual release of CO<sub>2</sub> is uncertain. Long-term studies may reveal higher lifetime emissions rates associated with degradation.

Many of the values used in the analysis are also uncertain and are based on estimates and professional judgment. For example, the asphalt cement input for hot mix asphalt was based on expert advice indicating that the range is variable—from about 3 to 5 percent—with actual content based on climate and geographical factors (Connolly 2000). Over this range, the effect on the calculated C storage factor is minimal (on the order of 0.1 percent). Similarly, changes in the assumed C content of asphalt cement would have only a minor effect.

The consumption figures for cut-back and emulsified asphalts are based on information reported for 1994. More recent trends indicate a decrease in cut-back use due to high VOC emission levels and a related increase in emulsified asphalt use as a substitute. This change in trend would indicate an overestimate of emissions from asphalt.

Future improvements to this uncertainty analysis, and to the overall estimation of a storage factor for asphalt, include characterizing the long-term fate of asphalt.

## Lubricants

Lubricants are used in industrial and transportation applications. They can be subdivided into oils and greases, which differ in terms of physical characteristics (e.g., viscosity), commercial applications, and environmental fate. According to EIA (2015b), the C content from U.S. production of lubricants in 2014 was approximately 5.7 MMT C. Based on apportioning oils and greases to various environmental fates, and characterizing those fates as resulting in either long-term storage or emissions, the overall C storage factor was estimated to be 9.2 percent; thus, emissions in 2014 were about 5.2 MMT C, or 18.9 MMT CO<sub>2</sub> Eq.

### Methodology and Data Sources

For each lubricant category, a storage factor was derived by identifying disposal fates and applying assumptions as to the disposition of the C for each practice. An overall lubricant C storage factor was calculated by taking a production-weighted average of the oil and grease storage factors.

### Oils

Regulation of used oil in the United States has changed dramatically over the past 20 years.<sup>39</sup> The effect of these regulations and policies has been to restrict landfilling and dumping, and to encourage collection of used oil. The economics of the petroleum industry have generally not favored re-refining—instead, most of the used oil that has been collected has been combusted.

Table A-82 provides an estimated allocation of the fates of lubricant oils (Rinehart 2000), along with an estimate of the proportion of C stored in each fate. The ultimate fate of the majority of oils (about 84 percent) is combustion, either during initial use or after collection as used oil. Combustion results in 99 percent oxidation to CO<sub>2</sub> (EIIIP 1999), with correspondingly little long-term storage of C in the form of ash. Dumping onto the ground or into storm sewers, primarily by “do-it-yourselfers” who change their own oil, is another fate that results in conversion to CO<sub>2</sub> given that the releases are generally small and most of the oil is biodegraded (based on the observation that land farming—application to soil—is one

---

<sup>39</sup> For example, the U.S. EPA “RCRA (Resource Conservation and Recovery Act) On-line” web site (<<http://www.epa.gov/rcraonline/>>) has over 50 entries on used oil regulation and policy for 1994 through 2000.

of the most frequently used methods for degrading refinery wastes). In the landfill environment, which tends to be anaerobic within municipal landfills, it is assumed that 90 percent of the oil persists in an undegraded form, based on analogy with the persistence of petroleum in native petroleum-bearing strata, which is also anaerobic. Re-refining adds a recycling loop to the fate of oil. Re-refined oil was assumed to have a storage factor equal to the weighted average for the other fates (i.e., after re-refining, the oil would have the same probability of combustion, landfilling, or dumping as virgin oil), that is, it was assumed that about 97 percent of the C in re-refined oil is ultimately oxidized. Because of the dominance of fates that result in eventual release as CO<sub>2</sub>, only about 3 percent of the C in oil lubricants goes into long-term storage.

**Table A-82: Commercial and Environmental Fate of Oil Lubricants (Percent)**

| Fate of Oil  | Portion of Total Oil | C Stored    |
|--|----------------------|-------------|
| <b>Combusted During Use</b>                            | <b>20%</b>           | <b>0.2%</b> |
| <b>Not Combusted During Use</b>                        | <b>80%</b>           | <b>2.7%</b> |
| Combusted as Used Oil <sup>a</sup>                     | 64%                  | 0.6%        |
| Dumped on the ground or in storm sewers                | 6%                   | NA          |
| Landfilled   | 2%                   | 1.8%        |
| Re-refined into lube oil base stock and other products | 8%                   | 0.2%        |
| <b>Weighted Average</b>                                | <b>NA</b>            | <b>2.9%</b> |

<sup>a</sup> For example, in boilers or space heaters.

NA (Not Applicable)

### Greases

Table A-83 provides analogous estimates for lubricant greases. Unlike oils, grease is generally not combusted during use, and combustion for energy recovery and re-refining is thought to be negligible. Although little is known about the fate of waste grease, it was assumed that 90 percent of the non-combusted portion is landfilled, and the remainder is dumped onto the ground or storm sewers. Because much of the waste grease will be in containers that render it relatively inaccessible to biodegradation, and because greases contain longer chain paraffins, which are more persistent than oils, it was assumed that 90 percent and 50 percent of the C in landfilled and dumped grease, respectively, would be stored. The overall storage factor is 82 percent for grease.

**Table A-83: Commercial and Environmental Fate of Grease Lubricants (Percent)**

| Fate of Grease                          | Portion of Total |              |
|---|------------------|--------------|
|   | Grease           | C Stored     |
| <b>Combusted During Use</b>             | <b>5%</b>        | <b>0.1%</b>  |
| <b>Not Combusted During Use</b>         | <b>95%</b>       | <b>81.7%</b> |
| Landfilled                              | 86%              | 77.0%        |
| Dumped on the ground or in storm sewers | 10%              | 4.8%         |
| <b>Weighted Average</b>                 | <b>NA</b>        | <b>81.8%</b> |

NA (Not Applicable)

Having derived separate storage factors for oil and grease, the last step was to estimate the weighted average for lubricants as a whole. No data were found apportioning the mass of lubricants into these two categories, but the U.S. Census Bureau does maintain records of the value of production of lubricating oils and lubricating greases. These were retrieved from the relevant industry series summaries from the *1997 Economic Census* (U.S. Bureau of the Census 1999). Assuming that the mass of lubricants can be allocated according to the proportion of value of production (92 percent oil, 8 percent grease), applying these weights to the storage factors for oils and greases (3 percent and 82 percent) yields an overall storage factor of 9.2 percent.

### Uncertainty

A Tier 2 Monte Carlo analysis was performed using @RISK software to determine the level of uncertainty surrounding the estimates of the lubricants weighted average C storage factor and the quantity of C emitted from lubricants in 2014. The Tier 2 analysis was performed to allow the specification of probability density functions for key variables, within a computational structure that mirrors the calculation of the Inventory estimate. Statistical analyses or expert judgments of uncertainty were not available directly from the information sources for the activity variables; thus, uncertainty estimates were determined using assumptions based on source category knowledge. Uncertainty estimates for oil and grease variables were assumed to have a moderate variance, in triangular or uniform distribution. Uncertainty estimates for

lubricants production were assumed to be rather high ( $\pm 20$  percent). A narrow uniform distribution, with 6 percent uncertainty ( $\pm 6$  percent) around the mean, was applied to the lubricant C content coefficient.

The Monte Carlo analysis produced a storage factor distribution with the 95 percent confidence interval of 4 percent and 17 percent around a mean value of 10 percent. This compares to the calculated Inventory estimate of 9 percent. The analysis produced a C emission distribution approximating a normal curve with a mean of 18.9 MMT CO<sub>2</sub> Eq., standard deviation of 1.5 and 95 percent confidence limits of 15.5 MMT CO<sub>2</sub> Eq. and 21.9 MMT CO<sub>2</sub> Eq. This compares to an inventory calculated estimate of 18.9 MMT CO<sub>2</sub> Eq.

The principal sources of uncertainty for the disposition of lubricants are the estimates of the commercial use, post-use, and environmental fate of lubricants, which, as noted above, are largely based on assumptions and judgment. There is no comprehensive system to track used oil and greases, which makes it difficult to develop a verifiable estimate of the commercial fates of oil and grease. The environmental fate estimates for percent of C stored are less uncertain, but also introduce uncertainty in the estimate.

The assumption that the mass of oil and grease can be divided according to their value also introduces uncertainty. Given the large difference between the storage factors for oil and grease, changes in their share of total lubricant production have a large effect on the weighted storage factor.

Future improvements to the analysis of uncertainty surrounding the lubricants C storage factor and C stored include further refinement of the uncertainty estimates for the individual activity variables.

## Waxes

Waxes are organic substances that are solid at ambient temperature, but whose viscosity decreases as temperature increases. Most commercial waxes are produced from petroleum refining, though “mineral” waxes derived from animals, plants, and lignite (coal) are also used. An analysis of wax end uses in the United States, and the fate of C in these uses, suggests that about 42 percent of C in waxes is emitted, and 58 percent is stored.

### Methodology and Data Sources

The National Petroleum Refiners Association (NPRA) considers the exact amount of wax consumed each year by end use to be proprietary (Maguire 2004). In general, about thirty percent of the wax consumed each year is used in packaging materials, though this percentage has declined in recent years. The next highest wax end use, and fastest growing end use, is candles, followed by construction materials and firelogs. Table A-84 categorizes some of the wax end uses, which the NPRA generally classifies into cosmetics, plastics, tires and rubber, hot melt (adhesives), chemically modified wax substances, and other miscellaneous wax uses (NPRA 2002).

**Table A-84: Emissive and Non-emissive (Storage) Fates of Waxes: Uses by Fate and Percent of Total Mass**

| Use                    | Emissive   | Non-emissive |
|------------------------|------------|--------------|
| Packaging              | 6%         | 24%          |
| Non-packaging          | 36%        | 34%          |
| Candles                | 18%        | 2%           |
| Construction Materials | 4%         | 14%          |
| Firelogs               | 7%         | +            |
| Cosmetics              | 1%         | 2%           |
| Plastics               | 1%         | 2%           |
| Tires/Rubber           | 1%         | 1%           |
| Hot Melts              | 1%         | 1%           |
| Chemically Modified    | 0%         | 1%           |
| Other                  | 2%         | 9%           |
| <b>Total</b>           | <b>42%</b> | <b>58%</b>   |

+ Does not exceed 0.5 percent

A C storage factor for each wax end use was estimated and then summed across all end uses to provide an overall C storage factor for wax. Because no specific data on C contents of wax used in each end use were available, all wax products are assumed to have the same C content. Table A-85 categorizes wax end uses identified by the NPRA, and lists the estimated C storage factor of each end use.

**Table A-85: Wax End-Uses by Fate, Percent of Total Mass, Percent C Stored, and Percent of Total C Mass Stored**

| Use                    | Percent of Total Wax Mass | Percent of C Stored | Percent of Total C Mass Stored |
|------------------------|---------------------------|---------------------|--------------------------------|
| <b>Packaging</b>       | <b>30%</b>                | <b>79%</b>          | <b>24%</b>                     |
| <b>Non-Packaging</b>   |                           |                     |                                |
| Candles                | 20%                       | 10%                 | 2%                             |
| Construction Materials | 18%                       | 79%                 | 14%                            |
| Firelogs               | 7%                        | 1%                  | +                              |
| Cosmetics              | 3%                        | 79%                 | 2%                             |
| Plastics               | 3%                        | 79%                 | 2%                             |
| Tires/Rubber           | 3%                        | 47%                 | 1%                             |
| Hot Melts              | 3%                        | 50%                 | 1%                             |
| Chemically Modified    | 1%                        | 79%                 | 1%                             |
| Other                  | 12%                       | 79%                 | 9%                             |
| <b>Total</b>           | <b>100%</b>               | <b>NA</b>           | <b>58%</b>                     |

+ Does not exceed 0.5 percent

Notes: Totals may not sum due to independent rounding. Estimates of percent stored are based on professional judgment, ICF International.

Source mass percentages: NPRA (2002).

Emissive wax end-uses include candles, firelogs (synthetic fireplace logs), hotmelts (adhesives), matches, and explosives. At about 20 percent, candles consume the greatest portion of wax among emissive end uses. As candles combust during use, they release emissions to the atmosphere. For the purposes of the Inventory, it is assumed that 90 percent of C contained in candles is emitted as CO<sub>2</sub>. In firelogs, petroleum wax is used as a binder and as a fuel, and is combusted during product use, likely resulting in the emission of nearly all C contained in the product. Similarly, C contained in hotmelts is assumed to be emitted as CO<sub>2</sub> as heat is applied to these products during use. It is estimated that 50 percent of the C contained in hot melts is stored. Together, candles, firelogs, and hotmelts constitute approximately 30 percent of annual wax production (NPRA 2002).

All of the wax utilized in the production of packaging, cosmetics, plastics, tires and rubber, and other products is assumed to remain in the product (i.e., it is assumed that there are no emissions of CO<sub>2</sub> from wax during the production of the product). Wax is used in many different packaging materials including wrappers, cartons, papers, paperboard, and corrugated products (NPRA 2002). Davie (1993) and Davie et al. (1995) suggest that wax coatings in packaging products degrade rapidly in an aerobic environment, producing CO<sub>2</sub>; however, because packaging products ultimately enter landfills typically having an anaerobic environment, most of the C from this end use is assumed to be stored in the landfill.

In construction materials, petroleum wax is used as a water repellent on wood-based composite boards, such as particle board (IGI 2002). Wax used for this end-use should follow the life-cycle of the harvested wood used in product, which is classified into one of 21 categories, evaluated by life-cycle, and ultimately assumed to either be disposed of in landfills or be combusted (EPA 2003).

The fate of wax used for packaging, in construction materials, and for most remaining end uses is ultimately to enter the municipal solid waste (MSW) stream, where it is either combusted or sent to landfill for disposal. Most of the C contained in these wax products will be stored. It is assumed that approximately 21 percent of the C contained in these products will be emitted through combustion or at landfill. With the exception of tires and rubber, these end-uses are assigned a C storage factor of 79 percent.

Waxes used in tires and rubber follow the life cycle of the tire and rubber products. Used tires are ultimately recycled, landfilled, or combusted. The life-cycle of tires is addressed elsewhere in this annex as part of the discussion of rubber products derived from petrochemical feedstocks. For the purposes of the estimation of the C storage factor for waxes, wax contained in tires and rubber products is assigned a C storage factor of 47 percent.

### Uncertainty

A Tier 2 Monte Carlo analysis was performed using @RISK software to determine the level of uncertainty surrounding the estimates of the wax C storage factor and the quantity of C emitted from wax in 2014. A Tier 2 analysis was performed to allow the specification of probability density functions for key variables, within a computational structure that mirrors the calculation of the Inventory estimate. Statistical analyses or expert judgments of uncertainty were not available directly from the information sources for the activity variables; thus, uncertainty estimates were determined using assumptions based on source category knowledge. Uncertainty estimates for wax variables were assumed to have a moderate

variance, in normal, uniform, or triangular distribution; uniform distributions were applied to total consumption of waxes and the C content coefficients.

The Monte Carlo analysis produced a storage factor distribution, whose 95 percent confidence interval values fell within the range of 49 percent and 70 percent, around the mean value of 58 percent. This compares to the calculated Inventory estimate of 58 percent. The analysis produced an emission distribution, with the 95 percent confidence interval values of 0.3 MMT CO<sub>2</sub> Eq. and 0.7 MMT CO<sub>2</sub> Eq., with a mean value of 0.4 MMT CO<sub>2</sub> Eq. This compares with a calculated Inventory estimate of 0.5 MMT CO<sub>2</sub> Eq., which falls within the range of 95 percent confidence limits established by this quantitative uncertainty analysis. Uncertainty associated with the wax storage factor is considerable due to several assumptions pertaining to wax imports/exports, consumption, and fates.

## Miscellaneous Products

Miscellaneous products are defined by the U.S. Energy Information Administration as: “all finished [petroleum] products not classified elsewhere, e.g., petrolatum; lube refining by-products (e.g., aromatic extracts and tars); absorption oils; ram-jet fuel; petroleum rocket fuel; synthetic natural gas feedstocks; and specialty oils.”

### Methodology and Data Sources

Data are not available concerning the distribution of each of the above-listed subcategories within the “miscellaneous products” category. However, based on the anticipated disposition of the products in each subcategory, it is assumed that all of the C content of miscellaneous products is emitted rather than stored. Petrolatum and specialty oils (which include greases) are likely to end up in solid waste or wastewater streams rather than in durable products, and would be emitted through waste treatment. Absorption oil is used in natural gas processing and is not a feedstock for manufacture of durable products. Jet fuel and rocket fuel are assumed to be combusted in use, and synthetic natural gas feedstocks are assumed to be converted to synthetic natural gas that is also combusted in use. Lube refining by-products could potentially be used as feedstocks for manufacture of durable goods, but such by-products are more likely to be used in emissive uses. Lube refining by-products and absorption oils are liquids and are precluded from disposal in landfills. Because no sequestering end uses of any of the miscellaneous products subcategories have been identified, a zero percent storage factor is assigned to miscellaneous products. According to EIA (2015), the C content of miscellaneous petroleum products in 2013 was approximately 20.3 MMT C/QBtu. One hundred percent of the C content is assumed to be emitted to the atmosphere, where it is oxidized to CO<sub>2</sub>.

### Uncertainty

A separate uncertainty analysis was not conducted for miscellaneous products, though this category was included in the uncertainty analysis of other non-energy uses discussed in the following section.

## Other Non-Energy Uses

The remaining fuel types use storage factors that are not based on U.S.-specific analysis. For industrial coking coal and distillate fuel oil, storage factors were taken from IPCC (2006), which in turn draws from Marland and Rotty (1984). These factors are 0.1 and 0.5, respectively.

IPCC does not provide guidance on storage factors for the remaining fuel types (petroleum coke, miscellaneous products, and other petroleum), and assumptions were made based on the potential fate of C in the respective NEUs. Specifically, the storage factor for petroleum coke is 0.3, based on information from Huurman (2006) indicating that petroleum coke is used in the Netherlands for production of pigments, with 30 percent being stored long-term. EIA (2014) defines “miscellaneous products” as “all finished products not classified elsewhere (e.g., petrolatum, lube refining by-products (aromatic extracts and tars), absorption oils, ram-jet fuel, petroleum rocket fuels, synthetic natural gas feedstocks, and specialty oils).” All of these uses are emissive, and therefore the storage factor for miscellaneous products is set at zero. The “other petroleum” category is reported by U.S. Territories and accounts mostly for the same products as miscellaneous products, but probably also includes some asphalt, known to be non-emissive. The exact amount of asphalt or any of the other miscellaneous products is confidential business information, but based on judgment the storage factor for this category was estimated at 0.1.

For all these fuel types, the overall methodology simply involves multiplying C content by a storage factor, yielding an estimate of the mass of C stored. To provide a complete analysis of uncertainty for the entire NEU subcategory, the uncertainty around the estimate of “other” NEUs was characterized, as discussed below.

## Uncertainty

A Tier 2 Monte Carlo analysis was performed using @RISK software to determine the level of uncertainty surrounding the weighted average of the remaining fuels' C storage factors and the total quantity of C emitted from these other fuels in 2014. A Tier 2 analysis was performed to allow the specification of probability density functions for key variables, within a computational structure that mirrors the calculation of the Inventory estimate. Statistical analyses or expert judgments of uncertainty were not available directly from the information sources for some of the activity variables; thus, uncertainty estimates were determined using assumptions based on source category knowledge. A uniform distribution was applied to coking coal consumption, while the remaining consumption inputs were assumed to be normally distributed. The C content coefficients were assumed to have a uniform distribution; the greatest uncertainty range of 10 percent ( $\pm 10$  percent) around the Inventory value, was applied to coking coal and miscellaneous products. C coefficients for distillate fuel oil ranged from 18.5 to 21.1 MMT C/QBtu. The fuel-specific storage factors were assigned wide triangular distributions indicating greater uncertainty.

The Monte Carlo analysis produced a storage factor distribution with 95 percent confidence limits of 4 percent and 24 percent, with a mean of 4 percent. This compares to the Inventory calculation of weighted average (across the various fuels) storage factor of about 4 percent. The analysis produced an emission distribution, with the 95 percent confidence limit of 14.2 MMT CO<sub>2</sub> Eq. and 21.7 MMT CO<sub>2</sub> Eq., and a mean of 19.6 MMT CO<sub>2</sub> Eq. This compares with the Inventory estimate of 19.6 MMT CO<sub>2</sub> Eq., which falls closer to the upper boundary of the 95 percent confidence limit. The uncertainty analysis results are driven primarily by the very broad uncertainty inputs for the storage factors.

## References

- ACC (2015a) "U.S. Resin Production & Sales: 2014 vs. 2013," American Chemistry Council. Available online at: <<http://www.americanchemistry.com/Jobs/EconomicStatistics/Plastics-Statistics/Production-and-Sales-Data-by-Resin.pdf>>.
- ACC (2015b) "*Guide to the Business of Chemistry, 2015*," American Chemistry Council.
- ACC (2014) "U.S. Resin Production & Sales: 2013 vs. 2012," American Chemistry Council. Available online at: <<http://www.americanchemistry.com/Jobs/EconomicStatistics/Plastics-Statistics/Production-and-Sales-Data-by-Resin.pdf>>.
- ACC (2012) "*Guide to the Business of Chemistry, 2012*," American Chemistry Council.
- ACC (2011) "*Guide to the Business of Chemistry, 2011*," American Chemistry Council.
- ACC (2010) "*Guide to the Business of Chemistry, 2010*," American Chemistry Council.
- ACC (2007-2011) "PIPS Year-End Resin Statistics: Production, Sales and Captive Use." Available online at: <<http://www.americanchemistry.com/Jobs/EconomicStatistics/Plastics-Statistics/Production-and-Sales-Data-by-Resin.pdf>>.
- APC (2003-2006) "APC Year-End Statistics." Available online at: <[http://www.americanplasticscouncil.org/s\\_apc/docs/1700/1678.pdf](http://www.americanplasticscouncil.org/s_apc/docs/1700/1678.pdf)>.
- APC (2001) *as cited in ACS (2001) "Production: slow gains in output of chemicals and products lagged behind U.S. economy as a whole"* Chemical & Engineering News.
- APC (2000) Facts and Figures, Chemical & Engineering News, June 26, 2000.
- Bank of Canada (2013) Financial Markets Department Year Average of Exchange Rates. Available online at: <<http://www.bankofcanada.ca/stats/assets/pdf/nraa-2012.pdf>>.
- Bank of Canada (2012) Financial Markets Department Year Average of Exchange Rates. Available online at: <<http://www.bankofcanada.ca/stats/assets/pdf/nraa-2011.pdf>>.
- Bank of Canada (2009) Financial Markets Department Year Average of Exchange Rates. Available online at: <<http://www.bankofcanada.ca/pdf/nraa09.pdf>>.
- Bureau of Labor Statistics (2015) Producer Price Index Industry Data: Soap and Other Detergent Manufacturing. Available online at: <<http://data.bls.gov:8080/PDQ/outside.jsp?survey=pc>>.
- Davie, I.N., J.P. Winter, and R.P. Varoney (1995) "Decomposition of Coated Papers from a Quick Service Restaurant." *Technical Association for Pulp and Paper Industry Journal*. Vol 78 (5): 127-130.

- Davie, I.N. (1993) "Compostability of Petroleum Wax-based Coatings." *Technical Association for Pulp and Paper Industry Journal*. Vol 76 (2): 167-170.
- EIA (2015b) Supplemental Tables on Petroleum Product detail. *Monthly Energy Review, December 2015*. Energy Information Administration, U.S. Department of Energy, Washington, D.C. DOE/EIA-0035(2015/12).
- EIA (2015a) Supplemental Tables on Petroleum Product detail. *Monthly Energy Review, February 2015*, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2014/02)
- EIA (2013b) *EIA Manufacturing Consumption of Energy (MECS) 2010*, U.S. Department of Energy, Energy Information Administration, Washington, DC.
- EIA (2010) *EIA Manufacturing Consumption of Energy (MECS) 2006*, U.S. Department of Energy, Energy Information Administration, Washington, DC. EIA (2005) *EIA Manufacturing Consumption of Energy (MECS) 2002*, U.S. Department of Energy, Energy Information Administration, Washington, DC.
- EIA (2001) *EIA Manufacturing Consumption of Energy (MECS) 1998*, U.S. Department of Energy, Energy Information Administration, Washington, DC.
- EIA (1997) *EIA Manufacturing Consumption of Energy (MECS) 1994*, U.S. Department of Energy, Energy Information Administration, Washington, DC.
- EIA (1994) *EIA Manufacturing Consumption of Energy (MECS) 1991*, U.S. Department of Energy, Energy Information Administration, Washington, DC.
- Eldredge-Roebuck (2000) Personal communication between Joe Casola, ICF Consulting and Brandt Eldredge-Roebuck, American Plastics Council, 11 July 2000.
- EIIP (2001) "Area Sources" *Asphalt Paving*, Emissions Inventory Improvement Program: State and Territorial Air Pollution Program Administrators/Association of Local Air Pollution Control Officials and U.S. EPA, EIIP Document Series Vol. III, Ch. 17. (STAPPA/ALAPCO/EPA), Washington DC, January 2001. Available online at <<http://www.epa.gov/ttn/chief/eiip/techreport/volume03/index.html>>.
- EIIP (1999) *Methods for Estimating Greenhouse Gas Emissions from Combustion of Fossil Fuels*. Emissions Inventory Improvement Program: State and Territorial Air Pollution Program Administrators/Association of Local Air Pollution Control Officials and U.S. Environmental Protection Agency, EIIP Document Series Volume VIII, Chapter 1, STAPPA/ALAPCO/EPA, Washington, DC. August 2000. Available online at <<http://www.epa.gov/ttn/chief/eiip/viii01.pdf>>. 28 August 2000.
- Environment Canada (2006) *Emissions Inventory Guidebook v1.3*. Criteria Air Contaminants Division: Quebec, Canada. Available online at: <<http://www.eea.europa.eu/publications/EMEPCORINAIR5/B4611vs1.3.pdf>>.
- EPA (2013a, 2015a) RCRAInfo, Biennial Report, Generation and Management (GM) Form (Section 2 - Onsite Management) and Waste Received from Offsite (WR) Form.
- EPA (2015b) "1970 - 2014 Average annual emissions, all criteria pollutants in MS Excel." *National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data*. Office of Air Quality Planning and Standards, December 2013. Available online at: <<http://www.epa.gov/ttn/chief/trends/index.html>>.
- EPA (2014) Chemical Data Access Tool (CDAT). U.S. Environmental Protection Agency, June 2014. Available online at <[http://java.epa.gov/oppt\\_chemical\\_search/](http://java.epa.gov/oppt_chemical_search/)>. Accessed January 2015.
- EPA (1996-2003a, 2005, 2007b, 2008, 2009a, 2011a, 2013b; 2014) *Municipal Solid Waste in the United States: Facts and Figures*. Office of Solid Waste and Emergency Response, U.S. Environmental Protection Agency, Washington, DC. Available online at: <<http://www.epa.gov/epaoswer/non-hw/muncpl/msw99.htm>>.
- EPA (2011b) EPA's Pesticides Industry Sales and Usage, 2006 and 2007 Market Estimates. Available online at <<http://www.epa.gov/oppead1/pestsales/>>. Accessed January 2012>.
- EPA (2009) Biennial Reporting System (BRS) Database. U.S. Environmental Protection Agency, Envirofacts Warehouse. Washington, DC. Available online at <<http://www.epa.gov/enviro/html/brs/>>.
- EPA (2006) *Air Emissions Trends - Continued Progress Through 2005*. U.S. Environmental Protection Agency, Washington DC. December 19, 2006. Available online at: <<http://www.epa.gov/air/airtrends/index.html>>.

- EPA (2004) EPA's Pesticides Industry Sales and Usage, 2000 and 2001 Market Estimates. Available online at <<http://www.epa.gov/oppbead1/pestsales/>>. Accessed September 2006.
- EPA (2003) E-mail correspondence containing preliminary ambient air pollutant data. Office of Air Pollution and the Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency. December 22, 2003.
- EPA (2002) EPA's Pesticides Industry Sales and Usage, 1998 and 1999 Market Estimates, table 3.6. Available online at <[http://www.epa.gov/oppbead1/pestsales/99pestsales/market\\_estimates1999.pdf](http://www.epa.gov/oppbead1/pestsales/99pestsales/market_estimates1999.pdf)>. Accessed July 2003.
- EPA (2001) AP 42, Volume I, Fifth Edition. Chapter 11: Mineral Products Industry. Available online at <<http://www.epa.gov/ttn/chief/ap42/ch11/index.html>>.
- EPA (2000a) *Biennial Reporting System (BRS)*. U.S. Environmental Protection Agency, Envirofacts Warehouse. Washington, DC. Available online at <<http://www.epa.gov/enviro/html/brs/>>.
- EPA (2000b) *Toxics Release Inventory, 1998*. U.S. Environmental Protection Agency, Office of Environmental Information, Office of Information Analysis and Access, Washington, DC. Available online at <<http://www.epa.gov/triexplorer/chemical.htm>>.
- EPA (1999) EPA's Pesticides Industry Sales and Usage, 1996-1997 Market Estimates and Available online at: <[http://www.epa.gov/oppbead1/pestsales/97pestsales/market\\_estimates1997.pdf](http://www.epa.gov/oppbead1/pestsales/97pestsales/market_estimates1997.pdf)>.
- EPA (1998) EPA's Pesticides Industry Sales and Usage, 1994-1995 Market Estimates. Available online at <[http://www.epa.gov/oppbead1/pestsales/95pestsales/market\\_estimates1995.pdf](http://www.epa.gov/oppbead1/pestsales/95pestsales/market_estimates1995.pdf)>.
- FEB (2013) *Fiber Economics Bureau, as cited in C&EN (2013) Lackluster Year for Chemical Output: Production stayed flat or dipped in most world regions in 2012*. Chemical & Engineering News, American Chemical Society, 1 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2012) *Fiber Economics Bureau, as cited in C&EN (2012) Too Quiet After the Storm: After a rebound in 2010, chemical production hardly grew in 2011*. Chemical & Engineering News, American Chemical Society, 2 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2011) *Fiber Economics Bureau, as cited in C&EN (2011) Output Ramps up in all Regions*. Chemical & Engineering News, American Chemical Society, 4 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2010) *Fiber Economics Bureau, as cited in C&EN (2010) Output Declines in U.S., Europe*. Chemical & Engineering News, American Chemical Society, 6 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2009) *Fiber Economics Bureau, as cited in C&EN (2009) Chemical Output Slipped In Most Regions*. Chemical & Engineering News, American Chemical Society, 6 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2007) *Fiber Economics Bureau, as cited in C&EN (2007) Gains in Chemical Output Continue*. Chemical & Engineering News, American Chemical Society. July 2, 2007. Available online at: <<http://www.cen-online.org>>.
- FEB (2005) *Fiber Economics Bureau, as cited in C&EN (2005) Production: Growth in Most Regions*. Chemical & Engineering News, American Chemical Society, 11 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2003) *Fiber Economics Bureau, as cited in C&EN (2003) Production Inches Up in Most Countries*. Chemical & Engineering News, American Chemical Society, 7 July. Available online at: <<http://www.cen-online.org>>.
- FEB (2001) *Fiber Economics Bureau, as cited in ACS (2001) Production: slow gains in output of chemicals and products lagged behind U.S. economy as a whole*. Chemical & Engineering News, American Chemical Society, 25 June. Available online at: <<http://pubs.acs.org/cen>>.
- Financial Planning Association (2006) Canada/US Cross-Border Tools: US/Canada Exchange Rates. Available online at: <[http://www.fpanet.org/global/planners/US\\_Canada\\_ex\\_rates.cfm](http://www.fpanet.org/global/planners/US_Canada_ex_rates.cfm)>. Accessed August 16, 2006.
- Gosselin, Smith, and Hodge (1984) *Clinical Toxicology of Commercial Products*. Fifth Edition, Williams & Wilkins, Baltimore.
- Huurman, J.W.F. (2006) *Recalculation of Dutch Stationary Greenhouse Gas Emissions Based on sectoral Energy Statistics 1990-2002*. Statistics Netherlands, Voorburg, The Netherlands.
- IGI (2002) *100 Industry Applications*. The International Group Inc. Available online at <[http://www.igiwax.com/100\\_apps.html](http://www.igiwax.com/100_apps.html) Toronto, Ontario>.

- IISRP (2003) "*IISRP Forecasts Moderate Growth in North America to 2007*" International Institute of Synthetic Rubber Producers, Inc. New Release; available online at: <<http://www.iisrp.com/press-releases/2003-Press-Releases/IISRP-NA-Forecast-03-07.html>>.
- IISRP (2000) *Synthetic Rubber Use Growth to Continue Through 2004, Says IISRP and RMA*. International Institute of Synthetic Rubber Producers press release.
- INEGI (2006) Producción bruta total de las unidades económicas manufactureras por Subsector, Rama, Subrama y Clase de actividad. Available online at: <[http://www.inegi.gob.mx/est/contenidos/espanol/proyectos/censos/ce2004/tb\\_manufacturas.asp](http://www.inegi.gob.mx/est/contenidos/espanol/proyectos/censos/ce2004/tb_manufacturas.asp)>. Accessed August 15.
- IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories. The National Greenhouse Gas Inventories Programme, H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe, eds.; Institute for Global Environmental Strategies (IGES). Hayama, Kanagawa, Japan.
- James, A. (2000) Personal communication between Suzanne Bratis of ICF International and Alan James of Akzo Nobel Coatings, Inc. July 2000. (Tel: 614-294-3361).
- Kelly (2000) Personal communication between Tom Smith, ICF Consulting and Peter Kelly, Asphalt Roofing Manufacturers Association, August 2000.
- Maguire (2004) Personal communication with J. Maguire, National Petrochemicals and Refiners Association. August – September 2004.
- Marland, G., and R.M. Rotty (1984) *Carbon dioxide emissions from fossil fuels: A procedure for estimation and results for 1950-1982*, Tellus 36b:232-261.
- NPRA (2002) North American Wax - A Report Card. Available online at: <<http://www.npra.org/members/publications/papers/lubes/LW-02-126.pdf>>.
- Rinehart, T. (2000) Personal communication between Thomas Rinehart of U.S. Environmental Protection Agency, Office of Solid Waste, and Randall Freed of ICF International. July 2000. (Tel: 703-308-4309).
- RMA (2014) 2013 *U.S. Scrap Tire Management Summary*. Rubber Manufacturers Association, Washington, DC. November 2014.
- RMA (2011) *U.S. Scrap Tire Management Summary: 2005-2009*. Rubber Manufacturers Association, Washington, DC. October 2011, updated September 2013.
- RMA (2009) "Scrap Tire Markets: Facts and Figures – Scrap Tire Characteristics." Available online at: <[http://www.rma.org/scrap\\_tires/scrap\\_tire\\_markets/scrap\\_tire\\_characteristics/](http://www.rma.org/scrap_tires/scrap_tire_markets/scrap_tire_characteristics/)>. Accessed 17 September 2009.
- Schneider, S. (2007) E-mail between Shelly Schneider of Franklin Associates (a division of ERG) and Sarah Shapiro of ICF International, January 10, 2007.
- SPI (2000) The Society of the Plastics Industry Website, <http://www.plasticsindustry.org/industry/stat3.htm>, Accessed 28 June 2000.
- U.S. Bureau of the Census (2015) U.S International Trade Commission (USITC) Trade Dataweb. Available online at: <<http://dataweb.usitc.gov>>.
- U.S. Bureau of the Census (1994, 1999, 2004, 2009, 2014) *1992, 1997, 2002, 2007, 2012 Economic Census*. Available online at <[http://factfinder.census.gov/servlet/DatasetMainPageServlet?\\_program=ECN&\\_submenuId=&\\_lang=en&\\_ts=>](http://factfinder.census.gov/servlet/DatasetMainPageServlet?_program=ECN&_submenuId=&_lang=en&_ts=>)>.
- U.S. International Trade Commission (1990-2015) "Interactive Tariff and Trade DataWeb: Quick Query." Available online at <<http://dataweb.usitc.gov/>>. Accessed November 2014.
- Vallianos, Jean (2015) Personal communication between Tyler Fitch of ICF International and Jean Vallianos of the American Chemistry Council, December 20, 2015.
- Vallianos, Jean (2014) Personal communication between Sarah Biggar of ICF International and Jean Vallianos of the American Chemistry Council, November 13, 2014.
- Vallianos, Jean (2013) Personal communication between Sarah Biggar of ICF International and Jean Vallianos of the American Chemistry Council, November 8, 2013.

Vallianos, Jean (2012) Personal communication between Ben Eskin of ICF International and Jean Vallianos of the American Chemistry Council, September 14, 2012.

Vallianos, Jean (2011) Personal communication between Joe Indvik of ICF International and Jean Vallianos of the American Chemistry Council, January 4, 2011.

# ANNEX 3 Methodological Descriptions for Additional Source or Sink Categories

## 3.1. Methodology for Estimating Emissions of CH<sub>4</sub>, N<sub>2</sub>O, and Indirect Greenhouse Gases from Stationary Combustion

### Estimates of CH<sub>4</sub> and N<sub>2</sub>O Emissions

Methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) emissions from stationary combustion were estimated using emission factors and methods from the Intergovernmental Panel on Climate Change (IPCC). Estimates were obtained by multiplying emission factors—by sector and fuel type—by fossil fuel and wood consumption data. This “top-down” methodology is characterized by two basic steps, described below. Data are presented in Table A-86 through Table A-91.

#### Step 1: Determine Energy Consumption by Sector and Fuel Type

Energy consumption from stationary combustion activities was grouped by sector: industrial, commercial, residential, electric power, and U.S. Territories. For CH<sub>4</sub> and N<sub>2</sub>O from industrial, commercial, residential, and U.S. Territories, estimates were based upon consumption of coal, gas, oil, and wood. Energy consumption and wood consumption data for the United States were obtained from the Energy Information Administration’s (EIA) *Monthly Energy Review, February 2016* and Published Supplemental Tables on Petroleum Product detail (EIA 2016). Because the United States does not include U.S. Territories in its national energy statistics, fuel consumption data for U.S. Territories were collected separately from the EIA’s *International Energy Statistics* (Jacobs 2010).<sup>40</sup> Fuel consumption for the industrial sector was adjusted to subtract out construction and agricultural use, which is reported under mobile sources.<sup>41</sup> Construction and agricultural fuel use was obtained from EPA (2013). The energy consumption data by sector were then adjusted from higher to lower heating values by multiplying by 0.90 for natural gas and wood and by 0.95 for coal and petroleum fuel. This is a simplified convention used by the International Energy Agency (IEA). Table A-86 provides annual energy consumption data for the years 1990 through 2014.

In this Inventory, the emission estimation methodology for the electric power sector was revised from Tier 1 to Tier 2 as fuel consumption by technology-type for the electricity generation sector was obtained from the Acid Rain Program Dataset (EPA 2015a). This combustion technology-and fuel-use data was available by facility from 1996 to 2014. Since there was a difference between the EPA (2015a) and EIA (2016) total energy consumption estimates, the remainder between total energy consumption using EPA (2015a) and EIA (2016) was apportioned to each combustion technology type and fuel combination using a ratio of energy consumption by technology type from 1996 to 2014.

Energy consumption estimates were not available from 1990 to 1995 in the EPA (2015a) dataset, and as a result, consumption was calculated using total electric power consumption from EIA (2016) and the ratio of combustion technology and fuel types from EPA 2015a. The consumption estimates from 1990 to 1995 were estimated by applying the 1996 consumption ratio by combustion technology type to the total EIA consumption for each year from 1990 to 1995.

Lastly, there were significant differences between wood biomass consumption in the electric power sector between the EPA (2015a) and EIA (2016) datasets. The difference in wood biomass consumption in the electric power sector was distributed to the residential, commercial, and industrial sectors according to their percent share of wood biomass energy consumption calculated from EIA (2016).

#### Step 2: Determine the Amount of CH<sub>4</sub> and N<sub>2</sub>O Emitted

Activity data for industrial, commercial, residential, and U.S. Territories and fuel type for each of these sectors were then multiplied by default Tier 1 emission factors to obtain emission estimates. Emission factors for the residential, commercial, and industrial sectors were taken from the *2006 IPCC Guidelines for National Greenhouse Gas Inventories*

<sup>40</sup> U.S. Territories data also include combustion from mobile activities because data to allocate U.S. Territories’ energy use were unavailable. For this reason, CH<sub>4</sub> and N<sub>2</sub>O emissions from combustion by U.S. Territories are only included in the stationary combustion totals.

<sup>41</sup> Though emissions from construction and farm use occur due to both stationary and mobile sources, detailed data was not available to determine the magnitude from each. Currently, these emissions are assumed to be predominantly from mobile sources.

(IPCC 2006). These N<sub>2</sub>O emission factors by fuel type (consistent across sectors) were also assumed for U.S. Territories. The CH<sub>4</sub> emission factors by fuel type for U.S. Territories were estimated based on the emission factor for the primary sector in which each fuel was combusted. Table A-87 provides emission factors used for each sector and fuel type. For the electric power sector, emissions were estimated by multiplying fossil fuel and wood consumption by technology- and fuel-specific Tier 2 IPCC emission factors shown in Table A-88. Emission factors were used from the *2006 IPCC Guidelines* as the factors presented in these IPCC guidelines were taken directly from U.S. Environmental Protection Agency (EPA) publications on emissions rates for combustion sources.

### Estimates of NO<sub>x</sub>, CO, and NMVOC Emissions

Emissions estimates for NO<sub>x</sub>, CO, and NMVOCs were obtained from data published on the National Emission Inventory (NEI) Air Pollutant Emission Trends web site (EPA 2015b), and disaggregated based on EPA (2003).

For indirect greenhouse gases, the major source categories included coal, fuel oil, natural gas, wood, other fuels (i.e., bagasse, liquefied petroleum gases, coke, coke oven gas, and others), and stationary internal combustion, which includes emissions from internal combustion engines not used in transportation. EPA periodically estimates emissions of NO<sub>x</sub>, CO, and NMVOCs by sector and fuel type using a "bottom-up" estimating procedure. In other words, the emissions were calculated either for individual sources (e.g., industrial boilers) or for many sources combined, using basic activity data (e.g., fuel consumption or deliveries, etc.) as indicators of emissions. The national activity data used to calculate the individual categories were obtained from various sources. Depending upon the category, these activity data may include fuel consumption or deliveries of fuel, tons of refuse burned, raw material processed, etc. Activity data were used in conjunction with emission factors that relate the quantity of emissions to the activity.

The basic calculation procedure for most source categories presented in EPA (2003) and EPA (2009) is represented by the following equation:

$$E_{p,s} = A_s \times EF_{p,s} \times (1 - C_{p,s}/100)$$

where,

|    |   |                            |
|----|---|----------------------------|
| E  | = | Emissions                  |
| p  | = | Pollutant                  |
| s  | = | Source category            |
| A  | = | Activity level             |
| EF | = | Emission factor            |
| C  | = | Percent control efficiency |

The EPA currently derives the overall emission control efficiency of a category from a variety of sources, including published reports, the 1985 National Acid Precipitation and Assessment Program (NAPAP) emissions inventory, and other EPA databases. The U. S. approach for estimating emissions of NO<sub>x</sub>, CO, and NMVOCs from stationary combustion as described above is similar to the methodology recommended by the IPCC (IPCC 2006).

**Table A-86: Fuel Consumption by Stationary Combustion for Calculating CH<sub>4</sub> and N<sub>2</sub>O Emissions (Tbtu)**

| Fuel/End-Use Sector | 1990          | 1995          | 2000          | 2001          | 2002          | 2003          | 2004          | 2005          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|---------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Coal</b>         | <b>19,610</b> | <b>20,888</b> | <b>23,080</b> | <b>22,391</b> | <b>22,343</b> | <b>22,576</b> | <b>22,636</b> | <b>22,949</b> | <b>22,458</b> | <b>22,710</b> | <b>22,225</b> | <b>19,670</b> | <b>20,697</b> | <b>18,989</b> | <b>16,715</b> | <b>17,399</b> | <b>17,363</b> |
| Residential         | 31            | 17            | 11            | 12            | 12            | 12            | 11            | 8             | 6             | 8             | 0             | 0             | 0             | 0             | 0             | 0             | 0             |
| Commercial          | 124           | 117           | 92            | 97            | 90            | 82            | 103           | 97            | 65            | 70            | 81            | 73            | 70            | 62            | 44            | 41            | 48            |
| Industrial          | 1,640         | 1,527         | 1,349         | 1,358         | 1,244         | 1,249         | 1,262         | 1,219         | 1,189         | 1,131         | 1,081         | 877           | 952           | 866           | 782           | 800           | 796           |
| Electric Power      | 17,807        | 19,217        | 21,618        | 20,920        | 20,987        | 21,199        | 21,228        | 21,591        | 21,161        | 21,465        | 21,026        | 18,682        | 19,639        | 18,024        | 15,852        | 16,521        | 16,483        |
| U.S. Territories    | 7             | 10            | 10            | 4             | 11            | 34            | 32            | 33            | 37            | 37            | 37            | 37            | 37            | 37            | 37            | 37            | 37            |
| <b>Petroleum</b>    | <b>6,166</b>  | <b>5,659</b>  | <b>6,148</b>  | <b>6,632</b>  | <b>6,010</b>  | <b>6,394</b>  | <b>6,561</b>  | <b>6,486</b>  | <b>6,201</b>  | <b>6,064</b>  | <b>5,241</b>  | <b>4,670</b>  | <b>4,728</b>  | <b>4,417</b>  | <b>4,065</b>  | <b>4,204</b>  | <b>4,030</b>  |
| Residential         | 1,375         | 1,262         | 1,429         | 1,465         | 1,361         | 1,468         | 1,468         | 1,368         | 1,202         | 1,220         | 1,202         | 1,138         | 1,116         | 1,060         | 847           | 936           | 990           |
| Commercial          | 869           | 695           | 695           | 720           | 647           | 765           | 764           | 716           | 678           | 680           | 635           | 669           | 643           | 625           | 506           | 540           | 543           |
| Industrial          | 2,750         | 2,380         | 2,283         | 2,535         | 2,371         | 2,496         | 2,669         | 2,776         | 3,111         | 2,996         | 2,427         | 1,949         | 2,054         | 1,962         | 1,924         | 2,037         | 1,869         |
| Electric Power      | 797           | 860           | 1,269         | 1,279         | 1,074         | 1,043         | 1,007         | 1,004         | 590           | 618           | 488           | 383           | 412           | 266           | 273           | 180           | 153           |
| U.S. Territories    | 375           | 462           | 472           | 632           | 557           | 622           | 653           | 623           | 620           | 550           | 490           | 531           | 502           | 504           | 516           | 510           | 475           |
| <b>Natural Gas</b>  | <b>17,266</b> | <b>19,337</b> | <b>20,919</b> | <b>20,224</b> | <b>20,908</b> | <b>20,894</b> | <b>21,152</b> | <b>20,938</b> | <b>20,626</b> | <b>22,019</b> | <b>22,286</b> | <b>21,952</b> | <b>22,912</b> | <b>23,115</b> | <b>24,137</b> | <b>24,949</b> | <b>25,692</b> |
| Residential         | 4,491         | 4,954         | 5,105         | 4,889         | 4,995         | 5,209         | 4,981         | 4,946         | 4,476         | 4,835         | 5,010         | 4,883         | 4,878         | 4,805         | 4,242         | 5,023         | 5,237         |
| Commercial          | 2,682         | 3,096         | 3,252         | 3,097         | 3,212         | 3,261         | 3,201         | 3,073         | 2,902         | 3,085         | 3,228         | 3,187         | 3,165         | 3,216         | 2,960         | 3,380         | 3,569         |
| Industrial          | 7,716         | 8,723         | 8,656         | 7,949         | 8,086         | 7,845         | 7,914         | 7,330         | 7,323         | 7,521         | 7,571         | 7,125         | 7,683         | 7,873         | 8,203         | 8,525         | 8,792         |
| Electric Power      | 2,376         | 2,564         | 3,894         | 4,266         | 4,591         | 4,551         | 5,032         | 5,565         | 5,899         | 6,550         | 6,447         | 6,730         | 7,159         | 7,194         | 8,683         | 7,964         | 8,033         |
| U.S. Territories    | 0             | 0             | 13            | 23            | 23            | 27            | 25            | 24            | 26            | 27            | 29            | 27            | 28            | 27            | 49            | 57            | 61            |
| <b>Wood</b>         | <b>2,216</b>  | <b>2,370</b>  | <b>2,262</b>  | <b>2,006</b>  | <b>1,995</b>  | <b>2,002</b>  | <b>2,121</b>  | <b>2,137</b>  | <b>2,099</b>  | <b>2,089</b>  | <b>2,059</b>  | <b>1,931</b>  | <b>1,981</b>  | <b>2,010</b>  | <b>2,010</b>  | <b>2,170</b>  | <b>2,230</b>  |
| Residential         | 580           | 520           | 420           | 370           | 380           | 400           | 410           | 430           | 380           | 420           | 470           | 500           | 440           | 450           | 420           | 580           | 580           |
| Commercial          | 66            | 72            | 71            | 67            | 69            | 71            | 70            | 70            | 65            | 70            | 73            | 73            | 72            | 69            | 61            | 70            | 73            |
| Industrial          | 1,442         | 1,652         | 1,636         | 1,443         | 1,396         | 1,363         | 1,476         | 1,452         | 1,472         | 1,413         | 1,339         | 1,178         | 1,273         | 1,309         | 1,339         | 1,312         | 1,325         |
| Electric Power      | 129           | 125           | 134           | 126           | 150           | 167           | 165           | 185           | 182           | 186           | 177           | 180           | 196           | 182           | 190           | 207           | 251           |
| U.S. Territories    | NE            |

NE – Not Estimated

Note: Totals may not sum due to independent rounding.

**Table A-87: CH<sub>4</sub> and N<sub>2</sub>O Emission Factors by Fuel Type and Sector (g/GJ)<sup>a</sup>**

| Fuel/End-Use Sector | CH <sub>4</sub> | N <sub>2</sub> O |
|---------------------|-----------------|------------------|
| <b>Coal</b>         |                 |                  |
| Residential         | 300             | 1.5              |
| Commercial          | 10              | 1.5              |
| Industrial          | 10              | 1.5              |
| Electric Power      | 1               | 1.5              |
| U.S. Territories    | 1               | 1.5              |
| <b>Petroleum</b>    |                 |                  |
| Residential         | 10              | 0.6              |
| Commercial          | 10              | 0.6              |
| Industrial          | 3               | 0.6              |
| Electric Power      | 3               | 0.6              |
| U.S. Territories    | 5               | 0.6              |
| <b>Natural Gas</b>  |                 |                  |
| Residential         | 5               | 0.1              |
| Commercial          | 5               | 0.1              |
| Industrial          | 1               | 0.1              |
| Electric Power      | 4               | 0.1              |
| U.S. Territories    | 1               | 0.1              |
| <b>Wood</b>         |                 |                  |
| Residential         | 300             | 4.0              |
| Commercial          | 300             | 4.0              |
| Industrial          | 30              | 4.0              |
| Electric Power      | 30              | 4.0              |
| U.S. Territories    | NA              | NA               |

NA - Not Applicable

<sup>a</sup>GJ (Gigajoule) = 10<sup>9</sup> joules. One joule = 9.486×10<sup>-4</sup> Btu.**Table A-88: CH<sub>4</sub> and N<sub>2</sub>O Emission Factors by Technology Type and Fuel Type for the Electric Power Sector (g/GJ)<sup>a</sup>**

| Technology                                | Configuration                  | CH <sub>4</sub> | N <sub>2</sub> O |
|---|--------------------------------|-----------------|------------------|
| <b>Liquid Fuels</b>                       |                                |                 |                  |
| Residual Fuel Oil/Shale Oil Boilers       | Normal Firing                  | 0.8             | 0.3              |
|   | Tangential Firing              | 0.8             | 0.3              |
| Gas/Diesel Oil Boilers                    | Normal Firing                  | 0.9             | 0.4              |
|   | Tangential Firing              | 0.9             | 0.4              |
| Large Diesel Oil Engines >600 hp (447kW)  |                                | 4               | NA               |
| <b>Solid Fuels</b>                        |                                |                 |                  |
| Pulverized Bituminous Combination Boilers | Dry Bottom, wall fired         | 0.7             | 0.5              |
|   | Dry Bottom, tangentially fired | 0.7             | 1.4              |
|   | Wet bottom                     | 0.9             | 1.4              |
| Bituminous Spreader Stoker Boilers        | With and without re-injection  | 1               | 0.7              |
| Bituminous Fluidized Bed Combustor        | Circulating Bed                | 1               | 61               |
|   | Bubbling Bed                   | 1               | 61               |
| Bituminous Cyclone Furnace                |                                | 0.2             | 0.6              |
| Lignite Atmospheric Fluidized Bed         |                                | NA              | 71               |
| <b>Natural Gas</b>                        |                                |                 |                  |
| Boilers                                   |                                | 1               | 1                |
| Gas-Fired Gas Turbines >3MW               |                                | 1               | 1                |
| Large Dual-Fuel Engines                   |                                | 258             | NA               |
| Combined Cycle                            |                                | 4               | 3                |
| <b>Peat</b>                               |                                |                 |                  |
| Peat Fluidized Bed Combustion             | Circulating Bed                | 3               | 7                |
|   | Bubbling Bed                   | 3               | 3                |
| <b>Biomass</b>                            |                                |                 |                  |
| Wood/Wood Waste Boilers                   |                                | 11              | 7                |
| Wood Recovery Boilers                     |                                | 1               | 1                |

NA - Not Applicable

<sup>a</sup>Ibid.

**Table A-89: NO<sub>x</sub> Emissions from Stationary Combustion (kt)**

| Sector/Fuel Type         | 1990          | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|--------------------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Electric Power</b>    | <b>6,045</b>  | <b>5,792</b> | <b>4,829</b> | <b>4,454</b> | <b>4,265</b> | <b>3,930</b> | <b>3,595</b> | <b>3,434</b> | <b>3,249</b> | <b>3,064</b> | <b>2,847</b> | <b>2,552</b> | <b>2,226</b> | <b>1,893</b> | <b>1,654</b> | <b>1,665</b> | <b>1,609</b> |
| Coal                     | 5,119         | 5,061        | 4,130        | 3,802        | 3,634        | 3,349        | 3,063        | 2,926        | 2,768        | 2,611        | 2,426        | 2,175        | 1,896        | 1,613        | 1,409        | 1,419        | 1,371        |
| Fuel Oil                 | 200           | 87           | 147          | 149          | 142          | 131          | 120          | 114          | 108          | 102          | 95           | 85           | 74           | 63           | 55           | 55           | 54           |
| Natural gas              | 513           | 510          | 376          | 325          | 310          | 286          | 262          | 250          | 236          | 223          | 207          | 186          | 162          | 138          | 120          | 121          | 117          |
| Wood                     | NA            | NA           | 36           | 37           | 36           | 33           | 30           | 29           | 27           | 26           | 24           | 21           | 19           | 16           | 14           | 14           | 13           |
| Other Fuels <sup>a</sup> | NA            | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           |
| Internal Combustion      | 213           | 134          | 140          | 140          | 143          | 132          | 121          | 115          | 109          | 103          | 95           | 86           | 75           | 63           | 55           | 56           | 54           |
| <b>Industrial</b>        | <b>2,559</b>  | <b>2,650</b> | <b>2,278</b> | <b>2,296</b> | <b>1,699</b> | <b>1,641</b> | <b>1,580</b> | <b>1,515</b> | <b>1,400</b> | <b>1,285</b> | <b>1,165</b> | <b>1,126</b> | <b>1,087</b> | <b>1,048</b> | <b>1,048</b> | <b>1,048</b> | <b>1,048</b> |
| Coal                     | 530           | 541          | 484          | 518          | 384          | 371          | 357          | 342          | 316          | 290          | 263          | 254          | 245          | 237          | 237          | 237          | 237          |
| Fuel Oil                 | 240           | 224          | 166          | 153          | 114          | 110          | 106          | 101          | 94           | 86           | 78           | 75           | 73           | 70           | 70           | 70           | 70           |
| Natural gas              | 877           | 999          | 710          | 711          | 526          | 508          | 489          | 469          | 433          | 398          | 361          | 348          | 336          | 324          | 324          | 324          | 324          |
| Wood                     | NA            | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           |
| Other Fuels <sup>a</sup> | 119           | 111          | 109          | 116          | 86           | 83           | 80           | 76           | 70           | 65           | 59           | 57           | 55           | 53           | 53           | 53           | 53           |
| Internal Combustion      | 792           | 774          | 809          | 798          | 591          | 570          | 549          | 527          | 486          | 446          | 405          | 391          | 378          | 364          | 364          | 364          | 364          |
| <b>Commercial</b>        | <b>671</b>    | <b>607</b>   | <b>507</b>   | <b>428</b>   | <b>438</b>   | <b>408</b>   | <b>378</b>   | <b>490</b>   | <b>471</b>   | <b>452</b>   | <b>433</b>   | <b>445</b>   | <b>456</b>   | <b>548</b>   | <b>548</b>   | <b>548</b>   | <b>548</b>   |
| Coal                     | 36            | 35           | 21           | 21           | 19           | 19           | 19           | 19           | 18           | 17           | 15           | 15           | 15           | 15           | 15           | 15           | 15           |
| Fuel Oil                 | 88            | 94           | 52           | 52           | 50           | 49           | 49           | 49           | 46           | 43           | 39           | 39           | 38           | 37           | 37           | 37           | 37           |
| Natural gas              | 181           | 210          | 161          | 165          | 157          | 156          | 155          | 145          | 135          | 124          | 122          | 120          | 118          | 118          | 118          | 118          | 118          |
| Wood                     | NA            | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           |
| Other Fuels <sup>a</sup> | 366           | 269          | 273          | 189          | 212          | 183          | 154          | 267          | 263          | 258          | 254          | 269          | 284          | 378          | 378          | 378          | 378          |
| <b>Residential</b>       | <b>749</b>    | <b>813</b>   | <b>439</b>   | <b>446</b>   | <b>422</b>   | <b>422</b>   | <b>420</b>   | <b>418</b>   | <b>390</b>   | <b>363</b>   | <b>335</b>   | <b>329</b>   | <b>324</b>   | <b>318</b>   | <b>318</b>   | <b>318</b>   | <b>318</b>   |
| Coal <sup>b</sup>        | NA            | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           |
| Fuel Oil <sup>b</sup>    | NA            | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           |
| Natural Gas <sup>b</sup> | NA            | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           | NA           |
| Wood                     | 42            | 44           | 21           | 22           | 21           | 21           | 21           | 20           | 19           | 18           | 16           | 16           | 16           | 16           | 16           | 16           | 16           |
| Other Fuels <sup>a</sup> | 707           | 769          | 417          | 424          | 402          | 401          | 400          | 398          | 371          | 345          | 318          | 313          | 308          | 302          | 302          | 302          | 302          |
| <b>Total</b>             | <b>10,023</b> | <b>9,862</b> | <b>8,053</b> | <b>7,623</b> | <b>6,825</b> | <b>6,401</b> | <b>5,973</b> | <b>5,858</b> | <b>5,511</b> | <b>5,163</b> | <b>4,780</b> | <b>4,452</b> | <b>4,092</b> | <b>3,807</b> | <b>3,567</b> | <b>3,579</b> | <b>3,522</b> |

NA - Not Applicable

<sup>a</sup> Other Fuels include LPG, waste oil, coke oven gas, coke, and non-residential wood (EPA 20145).<sup>b</sup> Residential coal, fuel oil, and natural gas emissions are included in the Other Fuels category (EPA 2015b).

Note: Totals may not sum due to independent rounding.

**Table A-90: CO Emissions from Stationary Combustion (kt)**

| Sector/Fuel Type         | 1990       | 1995       | 2000       | 2001       | 2002       | 2003       | 2004       | 2005       | 2006       | 2007       | 2008       | 2009       | 2010       | 2011       | 2012       | 2013       | 2014       |
|--------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| <b>Electric Power</b>    | <b>329</b> | <b>337</b> | <b>439</b> | <b>439</b> | <b>594</b> | <b>591</b> | <b>586</b> | <b>582</b> | <b>609</b> | <b>637</b> | <b>660</b> | <b>676</b> | <b>693</b> | <b>710</b> | <b>710</b> | <b>710</b> | <b>710</b> |
| Coal                     | 213        | 227        | 221        | 220        | 298        | 296        | 294        | 292        | 305        | 319        | 330        | 339        | 347        | 356        | 356        | 356        | 356        |
| Fuel Oil                 | 18         | 9          | 27         | 28         | 38         | 37         | 37         | 37         | 38         | 40         | 42         | 43         | 44         | 45         | 45         | 45         | 45         |
| Natural gas              | 46         | 49         | 96         | 92         | 125        | 124        | 123        | 122        | 128        | 134        | 138        | 142        | 145        | 149        | 149        | 149        | 149        |
| Wood                     | NA         |
| Other Fuels <sup>a</sup> | NA         | NA         | 31         | 32         | 44         | 43         | 43         | 43         | 45         | 47         | 48         | 50         | 51         | 52         | 52         | 52         | 52         |
| Internal Combustion      | 52         | 52         | 63         | 67         | 91         | 90         | 90         | 89         | 93         | 97         | 101        | 103        | 106        | 108        | 108        | 108        | 108        |

|                          |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |
|--------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Industrial</b>        | <b>797</b>   | <b>958</b>   | <b>1,106</b> | <b>1,137</b> | <b>1,150</b> | <b>1,116</b> | <b>1,081</b> | <b>1,045</b> | <b>968</b>   | <b>892</b>   | <b>815</b>   | <b>834</b>   | <b>853</b>   | <b>872</b>   | <b>872</b>   | <b>872</b>   | <b>872</b>   |
| Coal                     | 95           | 88           | 118          | 125          | 127          | 123          | 119          | 115          | 107          | 98           | 90           | 92           | 94           | 96           | 96           | 96           | 96           |
| Fuel Oil                 | 67           | 64           | 48           | 45           | 46           | 44           | 43           | 42           | 39           | 35           | 32           | 33           | 34           | 35           | 35           | 35           | 35           |
| Natural gas              | 205          | 313          | 355          | 366          | 370          | 359          | 348          | 336          | 312          | 287          | 262          | 268          | 274          | 281          | 281          | 281          | 281          |
| Wood                     | NA           |
| Other Fuels <sup>a</sup> | 253          | 270          | 300          | 321          | 325          | 316          | 306          | 295          | 274          | 252          | 230          | 236          | 241          | 247          | 247          | 247          | 247          |
| Internal Combustion      | 177          | 222          | 285          | 279          | 282          | 274          | 266          | 257          | 238          | 219          | 200          | 205          | 209          | 214          | 214          | 214          | 214          |
| <b>Commercial</b>        | <b>205</b>   | <b>211</b>   | <b>151</b>   | <b>154</b>   | <b>177</b>   | <b>173</b>   | <b>169</b>   | <b>166</b>   | <b>156</b>   | <b>146</b>   | <b>137</b>   | <b>138</b>   | <b>140</b>   | <b>142</b>   | <b>142</b>   | <b>142</b>   | <b>142</b>   |
| Coal                     | 13           | 14           | 14           | 13           | 15           | 15           | 15           | 14           | 14           | 13           | 12           | 12           | 12           | 12           | 12           | 12           | 12           |
| Fuel Oil                 | 16           | 17           | 17           | 17           | 20           | 19           | 19           | 19           | 18           | 16           | 15           | 16           | 16           | 16           | 16           | 16           | 16           |
| Natural gas              | 40           | 49           | 83           | 84           | 97           | 95           | 93           | 91           | 86           | 80           | 75           | 76           | 77           | 78           | 78           | 78           | 78           |
| Wood                     | NA           |
| Other Fuels <sup>a</sup> | 136          | 132          | 36           | 38           | 44           | 43           | 42           | 41           | 39           | 37           | 34           | 35           | 35           | 35           | 35           | 35           | 35           |
| <b>Residential</b>       | <b>3,668</b> | <b>3,877</b> | <b>2,644</b> | <b>2,648</b> | <b>3,044</b> | <b>2,982</b> | <b>2,919</b> | <b>2,856</b> | <b>2,690</b> | <b>2,524</b> | <b>2,357</b> | <b>2,387</b> | <b>2,416</b> | <b>2,446</b> | <b>2,446</b> | <b>2,446</b> | <b>2,446</b> |
| Coal <sup>b</sup>        | NA           |
| Fuel Oil <sup>b</sup>    | NA           |
| Natural Gas <sup>b</sup> | NA           |
| Wood                     | 3,430        | 3,629        | 2,416        | 2,424        | 2,787        | 2,730        | 2,673        | 2,615        | 2,463        | 2,310        | 2,158        | 2,185        | 2,212        | 2,239        | 2,239        | 2,239        | 2,239        |
| Other Fuels <sup>a</sup> | 238          | 248          | 228          | 224          | 257          | 252          | 247          | 241          | 227          | 213          | 199          | 202          | 204          | 207          | 207          | 207          | 207          |
| <b>Total</b>             | <b>5,000</b> | <b>5,383</b> | <b>4,340</b> | <b>4,377</b> | <b>4,965</b> | <b>4,862</b> | <b>4,756</b> | <b>4,648</b> | <b>4,423</b> | <b>4,198</b> | <b>3,969</b> | <b>4,036</b> | <b>4,103</b> | <b>4,170</b> | <b>4,170</b> | <b>4,170</b> | <b>4,169</b> |

NA - Not Applicable

<sup>a</sup> Other Fuels include LPG, waste oil, coke oven gas, coke, and non-residential wood (EPA 2015b).

<sup>b</sup> Residential coal, fuel oil, and natural gas emissions are included in the Other Fuels category (EPA 2015b).

Note: Totals may not sum due to independent rounding.

**Table A-91: NMVOC Emissions from Stationary Combustion (kt)**

| Sector/Fuel Type         | 1990       | 1995       | 2000       | 2001       | 2002       | 2003       | 2004       | 2005       | 2006       | 2007       | 2008      | 2009      | 2010       | 2011       | 2012       | 2013       | 2014       |
|--------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|-----------|-----------|------------|------------|------------|------------|------------|
| <b>Electric Power</b>    | <b>43</b>  | <b>40</b>  | <b>56</b>  | <b>55</b>  | <b>45</b>  | <b>45</b>  | <b>44</b>  | <b>44</b>  | <b>42</b>  | <b>41</b>  | <b>40</b> | <b>39</b> | <b>38</b>  | <b>37</b>  | <b>37</b>  | <b>37</b>  | <b>37</b>  |
| Coal                     | 24         | 26         | 27         | 26         | 21         | 21         | 21         | 21         | 20         | 20         | 19        | 18        | 18         | 18         | 18         | 18         | 18         |
| Fuel Oil                 | 5          | 2          | 4          | 4          | 4          | 4          | 4          | 3          | 3          | 3          | 3         | 3         | 3          | 3          | 3          | 3          | 3          |
| Natural Gas              | 2          | 2          | 12         | 12         | 10         | 10         | 10         | 10         | 9          | 9          | 9         | 9         | 8          | 8          | 8          | 8          | 8          |
| Wood                     | NA         | NA        | NA        | NA         | NA         | NA         | NA         | NA         |
| Other Fuels <sup>a</sup> | NA         | NA         | 2          | 2          | 1          | 1          | 1          | 1          | 1          | 1          | 1         | 1         | 1          | 1          | 1          | 1          | 1          |
| Internal Combustion      | 11         | 9          | 11         | 10         | 9          | 9          | 8          | 8          | 8          | 8          | 8         | 7         | 7          | 7          | 7          | 7          | 7          |
| <b>Industrial</b>        | <b>165</b> | <b>187</b> | <b>157</b> | <b>159</b> | <b>138</b> | <b>132</b> | <b>126</b> | <b>120</b> | <b>113</b> | <b>105</b> | <b>97</b> | <b>99</b> | <b>100</b> | <b>101</b> | <b>101</b> | <b>101</b> | <b>102</b> |
| Coal                     | 7          | 5          | 9          | 10         | 9          | 9          | 8          | 8          | 7          | 7          | 6         | 6         | 7          | 7          | 7          | 7          | 7          |
| Fuel Oil                 | 11         | 11         | 9          | 9          | 7          | 7          | 7          | 6          | 6          | 6          | 5         | 5         | 5          | 5          | 5          | 5          | 5          |
| Natural Gas              | 52         | 66         | 53         | 54         | 47         | 45         | 43         | 41         | 38         | 36         | 33        | 33        | 34         | 34         | 34         | 34         | 34         |
| Wood                     | NA         | NA        | NA        | NA         | NA         | NA         | NA         | NA         |
| Other Fuels <sup>a</sup> | 46         | 45         | 27         | 29         | 25         | 24         | 23         | 22         | 21         | 19         | 18        | 18        | 18         | 19         | 19         | 19         | 19         |
| Internal Combustion      | 49         | 60         | 58         | 57         | 49         | 47         | 45         | 43         | 40         | 37         | 35        | 35        | 36         | 36         | 36         | 36         | 36         |
| <b>Commercial</b>        | <b>18</b>  | <b>21</b>  | <b>28</b>  | <b>29</b>  | <b>61</b>  | <b>54</b>  | <b>48</b>  | <b>33</b>  | <b>34</b>  | <b>35</b>  | <b>36</b> | <b>38</b> | <b>40</b>  | <b>42</b>  | <b>42</b>  | <b>42</b>  | <b>42</b>  |
| Coal                     | 1          | 1          | 1          | 1          | 1          | 1          | 1          | 1          | 1          | +          | +         | +         | +          | +          | +          | +          | +          |

|                          |            |            |              |              |              |              |              |            |            |            |            |            |            |            |            |            |            |
|--------------------------|------------|------------|--------------|--------------|--------------|--------------|--------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Fuel Oil                 | 3          | 3          | 4            | 4            | 6            | 5            | 3            | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          |
| Natural Gas              | 7          | 10         | 14           | 14           | 23           | 18           | 14           | 9          | 8          | 7          | 6          | 7          | 7          | 7          | 7          | 7          | 7          |
| Wood                     | NA         | NA         | NA           | NA           | NA           | NA           | NA           | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         |
| Other Fuels <sup>a</sup> | 8          | 8          | 9            | 10           | 31           | 30           | 30           | 22         | 24         | 26         | 28         | 29         | 31         | 32         | 32         | 32         | 32         |
| <b>Residential</b>       | <b>686</b> | <b>725</b> | <b>837</b>   | <b>836</b>   | <b>1,341</b> | <b>1,067</b> | <b>793</b>   | <b>518</b> | <b>465</b> | <b>411</b> | <b>358</b> | <b>378</b> | <b>399</b> | <b>419</b> | <b>419</b> | <b>419</b> | <b>419</b> |
| Coal <sup>b</sup>        | NA         | NA         | NA           | NA           | NA           | NA           | NA           | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         |
| Fuel Oil <sup>b</sup>    | NA         | NA         | NA           | NA           | NA           | NA           | NA           | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         |
| Natural Gas <sup>b</sup> | NA         | NA         | NA           | NA           | NA           | NA           | NA           | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         | NA         |
| Wood                     | 651        | 688        | 809          | 809          | 1,297        | 1,032        | 767          | 502        | 450        | 398        | 346        | 366        | 386        | 406        | 406        | 406        | 406        |
| Other Fuels <sup>a</sup> | 35         | 37         | 27           | 27           | 43           | 35           | 26           | 17         | 15         | 13         | 12         | 12         | 13         | 14         | 14         | 14         | 14         |
| <b>Total</b>             | <b>912</b> | <b>973</b> | <b>1,077</b> | <b>1,080</b> | <b>1,585</b> | <b>1,298</b> | <b>1,011</b> | <b>716</b> | <b>654</b> | <b>593</b> | <b>531</b> | <b>553</b> | <b>576</b> | <b>599</b> | <b>599</b> | <b>599</b> | <b>599</b> |

NA - Not Applicable

+ Does not exceed 0.5 kt.

<sup>a</sup> "Other Fuels" include LPG, waste oil, coke oven gas, coke, and non-residential wood (EPA 2015b).

<sup>b</sup> Residential coal, fuel oil, and natural gas emissions are included in the "Other Fuels" category (EPA 2015b).

Note: Totals may not sum due to independent rounding.

## References

- EIA (2016) Monthly Energy Review, February 2016, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2016/02).
- EPA (2015a) Acid Rain Program Dataset 1996 - 2014. Office of Air and Radiation, Office of Atmospheric Programs, U.S. Environmental Protection Agency, Washington, D.C.
- EPA (2015b) "1970 - 2013 Average annual emissions, all criteria pollutants in MS Excel." National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data. Office of Air Quality Planning and Standards, October 2014. Available online at <<http://www.epa.gov/ttn/chief/trends/index.html>>.
- EPA (2013) NONROAD 2008a Model. Office of Transportation and Air Quality, U.S. Environmental Protection Agency. Available online at <<http://www.epa.gov/oms/nonrdmdl.htm>>.
- IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- Jacobs, G. (2010) Personal communication. Gwendolyn Jacobs, Energy Information Administration and Rubaab Bhangu, ICF International. U.S. Territories Fossil Fuel Consumption. Unpublished. U.S. Energy Information Administration. Washington, DC.

## 3.2. Methodology for Estimating Emissions of CH<sub>4</sub>, N<sub>2</sub>O, and Indirect Greenhouse Gases from Mobile Combustion and Methodology for and Supplemental Information on Transportation-Related GHG Emissions

### Estimating CO<sub>2</sub> Emissions by Transportation Mode

Transportation-related CO<sub>2</sub> emissions, as presented in the CO<sub>2</sub> Emissions from Fossil Fuel Combustion section of the Energy chapter, were calculated using the methodology described in Annex 2.1. This section provides additional information on the data sources and approach used for each transportation fuel type. As noted in Annex 2.1, CO<sub>2</sub> emissions estimates for the transportation sector were calculated directly for on-road diesel fuel and motor gasoline based on data sources for individual modes of transportation (considered a bottom up approach). For most other fuel and energy types (aviation gasoline, residual fuel oil, natural gas, LPG, and electricity), CO<sub>2</sub> emissions were calculated based on transportation sector-wide fuel consumption estimates from the Energy Information Administration (EIA 2016 and EIA 2015) and apportioned to individual modes (considered a “top down” approach). Carbon dioxide emissions from commercial jet fuel use are obtained directly from the Federal Aviation Administration (FAA 2016), while CO<sub>2</sub> emissions from other aircraft jet fuel consumption is determined using a top down approach.

Based on interagency discussions between EPA, EIA, and FHWA beginning in 2005, it was agreed that use of “bottom up” data would be more accurate for diesel fuel and motor gasoline consumption in the transportation sector, based on the availability of reliable data sources. A “bottom up” diesel calculation was first implemented in the 1990 through 2005 Inventory, and a bottom-up gasoline calculation was introduced in the 1990 through 2006 Inventory for the calculation of emissions from on-road vehicles. Estimated motor gasoline and diesel consumption data for on-road vehicles by vehicle type come from FHWA’s *Highway Statistics*, Table VM-1 (FHWA 1996 through 2015),<sup>42</sup> and are based on federal and state fuel tax records. These fuel consumption estimates were then combined with estimates of fuel shares by vehicle type from DOE’s Transportation Energy Data Book Annex Tables A.1 through A.6 (DOE 1993 through 2015) to develop an estimate of fuel consumption for each vehicle type (i.e., passenger cars, light-duty trucks, buses, medium- and heavy-duty trucks, motorcycles). The on-road gas and diesel fuel consumption estimates by vehicle type were then adjusted for each year so that the sum of gasoline and diesel fuel consumption across all on-road vehicle categories matched the fuel consumption estimates in *Highway Statistics*’ Table MF-27 (FHWA 1996 through 2015). This resulted in a final “bottom up” estimate of motor gasoline and diesel fuel use by vehicle type, consistent with the FHWA total for on-road motor gasoline and diesel fuel use.

A primary challenge to switching from a top-down approach to a bottom-up approach for the transportation sector relates to potential incompatibilities with national energy statistics. From a multi-sector national standpoint, EIA develops the most accurate estimate of total motor gasoline and diesel fuel supplied and consumed in the United States. EIA then allocates this total fuel consumption to each major end-use sector (residential, commercial, industrial and transportation) using data from the *Fuel Oil and Kerosene Sales* (FOKS) report for distillate fuel oil and FHWA for motor gasoline. However, the “bottom-up” approach used for the on-road and non-road fuel consumption estimate, as described above, is considered to be the most representative of the transportation sector’s share of the EIA total consumption. Therefore, for years in which there was a disparity between EIA’s fuel allocation estimate for the transportation sector and the “bottom-up” estimate, adjustments were made to other end-use sector fuel allocations (residential, commercial and industrial) in order for the consumption of all sectors combined to equal the “top-down” EIA value.

In the case of motor gasoline, estimates of fuel use by recreational boats come from the NONROAD component of EPA’s MOVES2014a model (EPA 2015d), and these estimates, along with those from other sectors (e.g., commercial sector, industrial sector), were adjusted for years in which the bottom-up on-road motor gasoline consumption estimate exceeded the EIA estimate for total gasoline consumption of all sectors. Similarly, to ensure consistency with EIA’s total diesel estimate for all sectors, the diesel consumption totals for the residential, commercial, and industrial sectors were adjusted proportionately.

---

<sup>42</sup> In 2011 FHWA changed its methods for estimating vehicle miles traveled (VMT) and related data. These methodological changes included how vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. These changes were first incorporated for the 2010 Inventory and apply to the 2007-14 time period. This resulted in large changes in VMT and fuel consumption data by vehicle class, thus leading to a shift in emissions among on-road vehicle classes. For example, the category “Passenger Cars” has been replaced by “Light-duty Vehicles-Short Wheelbase” and “Other 2 axle-4 Tire Vehicles” has been replaced by “Light-duty Vehicles, Long Wheelbase.” This change in vehicle classification has moved some smaller trucks and sport utility vehicles from the light truck category to the passenger vehicle category in this emission inventory. These changes are reflected in a large drop in light-truck emissions between 2006 and 2007.

Estimates of diesel fuel consumption from rail were taken from the Association of American Railroads (AAR 2008 through 2015) for Class I railroads, the American Public Transportation Association (APTA 2007 through 2015 and APTA 2006) and Gaffney (2007) for commuter rail, the Upper Great Plains Transportation Institute (Benson 2002 through 2004) and Whorton (2006 through 2013) for Class II and III railroads, and DOE's *Transportation Energy Data Book* (DOE 1993 through 2015) for passenger rail. Estimates of diesel from ships and boats were taken from EIA's *Fuel Oil and Kerosene Sales* (1991 through 2015).

As noted above, for fuels other than motor gasoline and diesel, EIA's transportation sector total was apportioned to specific transportation sources. For jet fuel, estimates come from: FAA (2016) for domestic and international commercial aircraft, and DESC (2015) for domestic and international military aircraft. General aviation jet fuel consumption is calculated as the difference between total jet fuel consumption as reported by EIA and the total consumption from commercial and military jet fuel consumption. Commercial jet fuel CO<sub>2</sub> estimates are obtained directly from the Federal Aviation Administration (FAA 2016), while CO<sub>2</sub> emissions from domestic military and general aviation jet fuel consumption is determined using a top down approach. Domestic commercial jet fuels CO<sub>2</sub> from FAA is subtracted from total domestic jet fuel CO<sub>2</sub> emissions, and this remaining value is apportioned among domestic military and domestic general aviation based on their relative proportion of energy consumption. Estimates for biofuels, including ethanol and biodiesel were discussed separately in Chapter 3.2 Carbon Emitted from Non-Energy Uses of Fossil Fuels under the methodology for Estimating CO<sub>2</sub> from Fossil Combustion, and in Chapter 3.10 Wood Biomass and Ethanol Consumption and were not apportioned to specific transportation sources. Consumption estimates for biofuels were calculated based on data from the Energy Information Administration (EIA 2016).

Table A-92 displays estimated fuel consumption by fuel and vehicle type. Table A-93 displays estimated energy consumption by fuel and vehicle type. The values in both of these tables correspond to the figures used to calculate CO<sub>2</sub> emissions from transportation. Except as noted above, they are estimated based on EIA transportation sector energy estimates by fuel type, with activity data used to apportion consumption to the various modes of transport. The motor gasoline and diesel fuel consumption volumes published by EIA and FHWA include ethanol blended with gasoline and biodiesel blended with diesel. Biofuels blended with conventional fuels were subtracted from these consumption totals in order to be consistent with IPCC methodological guidance and UNFCCC reporting obligations, for which net carbon fluxes in biogenic carbon reservoirs in croplands are accounted for in the estimates for Land Use, Land-Use Change and Forestry chapter, not in Energy chapter totals. Ethanol fuel volumes were removed from motor gasoline consumption estimates for years 1990 through 2014 and biodiesel fuel volumes were removed from diesel fuel consumption volumes for years 2001 through 2014, as there was negligible use of biodiesel as a diesel blending component prior to 2001. The subtraction or removal of biofuels blended into motor gasoline and diesel were conducted following the methodology outlined in Step 2 ("Remove Biofuels from Petroleum") of the EIA's *Monthly Energy Review* (MER) Section 12 notes.

In order to remove the volume of biodiesel blended into diesel fuel, the refinery and blender net volume inputs of renewable diesel fuel sourced from EIA Petroleum Supply Annual (EIA 2015) *Table 18 - Refinery Net Input of Crude Oil and Petroleum Products* and *Table 20 - Blender Net Inputs of Petroleum Products* were subtracted from the transportation sector's total diesel fuel consumption volume (for both the "top-down" EIA and "bottom-up" FHWA estimates). To remove the fuel ethanol blended into motor gasoline, ethanol energy consumption data sourced from MER *Table 10.2b - Renewable Energy Consumption: Industrial and Transportation Sectors* (EIA 2016) were subtracted from the total EIA and FHWA transportation motor gasoline energy consumption estimates.

Total ethanol and biodiesel consumption estimates are shown separately in Table A-94.<sup>43</sup>

---

<sup>43</sup> Note that the refinery and blender net volume inputs of renewable diesel fuel sourced from EIA's Petroleum Supply Annual (PSA) differs from the biodiesel volume presented in Table A-94. The PSA data is representative of the amount of biodiesel that refineries and blenders added to diesel fuel to make low level biodiesel blends. This is the appropriate value to subtract from total diesel fuel volume, as it represents the amount of biofuel blended into diesel to create low-level biodiesel blends. The biodiesel consumption value presented in Table A-93 is representative of the total biodiesel consumed and includes biodiesel components in all types of fuel formulations, from low level (<5%) to high level (6-20%, 100%) blends of biodiesel. This value is sourced from MER Table 10.4 and is calculated as biodiesel production plus biodiesel net imports minus biodiesel stock exchange.

**Table A-92. Fuel Consumption by Fuel and Vehicle Type (million gallons unless otherwise specified)**

| Fuel/Vehicle Type                                      | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007 <sup>a</sup> | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Motor Gasoline<sup>b,c</sup></b>                    | <b>110,417</b> | <b>117,429</b> | <b>128,174</b> | <b>129,590</b> | <b>132,169</b> | <b>132,600</b> | <b>134,336</b> | <b>133,294</b> | <b>131,337</b> | <b>130,768</b>    | <b>125,050</b> | <b>124,189</b> | <b>123,175</b> | <b>120,497</b> | <b>120,035</b> | <b>120,138</b> | <b>122,188</b> |
| Passenger Cars   | 69,763         | 67,496         | 72,320         | 72,920         | 74,313         | 71,931         | 71,504         | 73,856         | 70,791         | 88,607            | 84,714         | 83,918         | 83,230         | 82,621         | 82,464         | 82,463         | 82,694         |
| Light-Duty Trucks                                      | 34,698         | 44,074         | 50,398         | 50,871         | 52,023         | 55,417         | 57,540         | 53,733         | 54,798         | 34,933            | 33,074         | 33,473         | 33,262         | 31,612         | 31,270         | 31,305         | 33,086         |
| Motorcycles  | 194            | 199            | 208            | 192            | 189            | 184            | 193            | 182            | 210            | 472               | 487            | 468            | 411            | 401            | 459            | 437            | 427            |
| Buses  | 39             | 41             | 43             | 40             | 38             | 36             | 49             | 41             | 41             | 79                | 81             | 84             | 82             | 80             | 92             | 95             | 100            |
| Medium- and Heavy-Duty Trucks                          | 4,350          | 4,044          | 4,065          | 3,961          | 4,006          | 3,446          | 3,475          | 3,922          | 3,961          | 5,164             | 5,220          | 4,798          | 4,773          | 4,383          | 4,358          | 4,455          | 4,506          |
| Recreational Boats <sup>d</sup>                        | 1,374          | 1,575          | 1,140          | 1,606          | 1,601          | 1,587          | 1,574          | 1,560          | 1,536          | 1,514             | 1,474          | 1,448          | 1,417          | 1,401          | 1,391          | 1,382          | 1,376          |
| <b>Distillate Fuel Oil (Diesel Fuel)<sup>b,c</sup></b> | <b>25,631</b>  | <b>31,604</b>  | <b>39,241</b>  | <b>39,057</b>  | <b>40,347</b>  | <b>41,176</b>  | <b>42,667</b>  | <b>44,658</b>  | <b>45,844</b>  | <b>46,427</b>     | <b>44,026</b>  | <b>39,873</b>  | <b>41,477</b>  | <b>42,280</b>  | <b>42,045</b>  | <b>42,672</b>  | <b>44,031</b>  |
| Passenger Cars   | 771            | 765            | 356            | 357            | 364            | 412            | 419            | 414            | 403            | 403               | 363            | 354            | 367            | 399            | 401            | 399            | 408            |
| Light-Duty Trucks                                      | 1,119          | 1,452          | 1,961          | 2,029          | 2,133          | 2,652          | 2,822          | 2,518          | 2,611          | 1,327             | 1,184          | 1,180          | 1,227          | 1,277          | 1,271          | 1,265          | 1,365          |
| Buses  | 781            | 851            | 997            | 906            | 860            | 930            | 1,316          | 1,030          | 1,034          | 1,520             | 1,436          | 1,335          | 1,326          | 1,419          | 1,515          | 1,525          | 1,634          |
| Medium- and Heavy-Duty Trucks                          | 18,574         | 23,240         | 30,179         | 30,124         | 31,417         | 31,539         | 32,598         | 35,159         | 36,089         | 37,518            | 35,726         | 32,364         | 33,683         | 33,859         | 33,877         | 34,426         | 35,541         |
| Recreational Boats                                     | 190            | 228            | 270            | 278            | 286            | 294            | 302            | 311            | 319            | 327               | 335            | 343            | 351            | 357            | 364            | 368            | 376            |
| Ships and Other Boats                                  | 735            | 1,204          | 1,372          | 1,244          | 1,197          | 1,173          | 802            | 780            | 724            | 794               | 767            | 768            | 726            | 993            | 733            | 741            | 606            |
| Rail <sup>e</sup>                                      | 3,461          | 3,863          | 4,106          | 4,119          | 4,089          | 4,176          | 4,407          | 4,446          | 4,664          | 4,538             | 4,215          | 3,529          | 3,798          | 3,975          | 3,884          | 3,948          | 4,101          |
| <b>Jet Fuel<sup>f</sup></b>                            | <b>19,186</b>  | <b>17,991</b>  | <b>20,002</b>  | <b>19,454</b>  | <b>19,004</b>  | <b>18,389</b>  | <b>19,147</b>  | <b>19,420</b>  | <b>18,695</b>  | <b>18,407</b>     | <b>17,749</b>  | <b>15,809</b>  | <b>15,537</b>  | <b>15,036</b>  | <b>14,705</b>  | <b>15,088</b>  | <b>15,237</b>  |
| Commercial Aircraft                                    | 11,569         | 12,136         | 14,672         | 13,121         | 12,774         | 12,943         | 13,147         | 13,976         | 14,426         | 14,708            | 13,400         | 12,588         | 11,931         | 12,067         | 11,932         | 12,031         | 12,131         |
| General Aviation Aircraft                              | 4,034          | 3,361          | 3,163          | 3,975          | 4,119          | 3,323          | 3,815          | 3,583          | 2,590          | 2,043             | 2,682          | 1,787          | 2,322          | 1,895          | 1,659          | 2,033          | 1,676          |
| Military Aircraft                                      | 3,583          | 2,495          | 2,167          | 2,359          | 2,110          | 2,123          | 2,185          | 1,860          | 1,679          | 1,656             | 1,667          | 1,434          | 1,283          | 1,074          | 1,114          | 1,024          | 1,430          |
| <b>Aviation Gasoline<sup>f</sup></b>                   | <b>374</b>     | <b>329</b>     | <b>302</b>     | <b>291</b>     | <b>281</b>     | <b>251</b>     | <b>260</b>     | <b>294</b>     | <b>278</b>     | <b>263</b>        | <b>235</b>     | <b>221</b>     | <b>225</b>     | <b>225</b>     | <b>209</b>     | <b>186</b>     | <b>181</b>     |
| General Aviation Aircraft                              | 374            | 329            | 302            | 291            | 281            | 251            | 260            | 294            | 278            | 263               | 235            | 221            | 225            | 225            | 209            | 186            | 181            |
| <b>Residual Fuel Oil<sup>f,g</sup></b>                 | <b>2,006</b>   | <b>2,587</b>   | <b>2,963</b>   | <b>1,066</b>   | <b>1,522</b>   | <b>662</b>     | <b>1,245</b>   | <b>1,713</b>   | <b>2,046</b>   | <b>2,579</b>      | <b>1,812</b>   | <b>1,241</b>   | <b>1,818</b>   | <b>1,723</b>   | <b>1,410</b>   | <b>1,345</b>   | <b>517</b>     |
| Ships and Other Boats                                  | 2,006          | 2,587          | 2,963          | 1,066          | 1,522          | 662            | 1,245          | 1,713          | 2,046          | 2,579             | 1,812          | 1,241          | 1,818          | 1,723          | 1,410          | 1,345          | 517            |
| <b>Natural Gas<sup>f</sup> (trillion cubic feet)</b>   | <b>0.7</b>     | <b>0.7</b>     | <b>0.7</b>     | <b>0.6</b>     | <b>0.7</b>     | <b>0.6</b>     | <b>0.6</b>     | <b>0.6</b>     | <b>0.6</b>     | <b>0.6</b>        | <b>0.7</b>     | <b>0.7</b>     | <b>0.7</b>     | <b>0.7</b>     | <b>0.8</b>     | <b>0.9</b>     | <b>0.9</b>     |
| Passenger Cars   | -              | -              | -              | -              | -              | -              | -              | -              | -              | -                 | -              | -              | -              | -              | -              | -              | -              |
| Light-Duty Trucks                                      | -              | -              | -              | -              | -              | -              | -              | -              | -              | -                 | -              | -              | -              | -              | -              | -              | -              |
| Buses  | -              | -              | -              | -              | -              | -              | -              | -              | -              | -                 | -              | -              | -              | -              | -              | -              | -              |
| Pipelines  | 0.7            | 0.7            | 0.6            | 0.6            | 0.7            | 0.6            | 0.6            | 0.6            | 0.6            | 0.6               | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            | 0.8            | 0.9            |
| <b>LPG<sup>f</sup></b>                                 | <b>265</b>     | <b>206</b>     | <b>138</b>     | <b>159</b>     | <b>166</b>     | <b>207</b>     | <b>222</b>     | <b>327</b>     | <b>320</b>     | <b>257</b>        | <b>468</b>     | <b>331</b>     | <b>348</b>     | <b>404</b>     | <b>442</b>     | <b>525</b>     | <b>519</b>     |
| Buses  | -              | 1.6            | 1.5            | 0.3            | 0.6            | 0.7            | 0.7            | 1.0            | 1.0            | -                 | -              | -              | -              | -              | -              | 4.6            | 4.6            |
| Light-Duty Trucks                                      | 106            | 98             | 88             | 108            | 117            | 144            | 167            | 247            | 229            | 185               | 340            | 228            | 243            | 283            | 316            | 368            | 364            |
| Medium- and Heavy-Duty Trucks                          | 159            | 106            | 49             | 51             | 49             | 62             | 55             | 79             | 89             | 72                | 128            | 103            | 106            | 121            | 126            | 152            | 150            |
| <b>Electricity<sup>f,h</sup></b>                       | <b>4,751</b>   | <b>4,975</b>   | <b>5,382</b>   | <b>5,724</b>   | <b>5,517</b>   | <b>6,810</b>   | <b>7,224</b>   | <b>7,506</b>   | <b>7,358</b>   | <b>8,173</b>      | <b>7,653</b>   | <b>7,768</b>   | <b>7,712</b>   | <b>7,672</b>   | <b>7,320</b>   | <b>7,625</b>   | <b>7,758</b>   |
| Rail   | 4,751          | 4,975          | 5,382          | 5,724          | 5,517          | 6,810          | 7,224          | 7,506          | 7,358          | 8,173             | 7,653          | 7,768          | 7,712          | 7,672          | 7,320          | 7,625          | 7,758          |

- Unreported or zero

<sup>a</sup> In 2011, FHWA changed its methodology for Table VM-1, which impacts estimates for the 2007-2014 time period. These methodological changes include how on-road vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. This resulted in large changes in fuel consumption data by vehicle class between 2006 and 2007.

<sup>b</sup> Figures do not include ethanol blended in motor gasoline or biodiesel blended into distillate fuel oil. Net carbon fluxes associated with ethanol are accounted for in the Land Use, Land-Use Change and Forestry chapter. This table is calculated with the heat content for gasoline without ethanol (from Table A.2 in the EIA Annual Energy Review) rather than the annually variable quantity-weighted heat content for gasoline with ethanol, which varies by year. In addition, updates to the distillate fuel oil heat content data from EIA for years 1993 through present resulted in changes to the time series for energy consumption and emissions compared to the previous Inventory.

<sup>c</sup> Gasoline and diesel highway vehicle fuel consumption estimates are based on data from FHWA Highway Statistics Table VM-1 and MF-27 (FHWA 1996 through 2015). These fuel consumption estimates are combined with estimates of fuel shares by vehicle type from DOE's TEDB Annex Tables A.1 through A.6 (DOE 1993 through 2015). TEDB data for 2014 has not been published yet, therefore 2013 data is used as a proxy.

<sup>d</sup> Fluctuations in recreational boat gasoline estimates reflect the use of this category to reconcile bottom-up values with EIA total gasoline estimates.

<sup>e</sup> Class II and Class II diesel consumption data for 2014 is not available yet, therefore 2013 data is used as a proxy.

<sup>f</sup> Estimated based on EIA transportation sector energy estimates by fuel type, with bottom-up activity data used for apportionment to modes.

<sup>g</sup> Fluctuations in reported fuel consumption may reflect data collection problems.

<sup>h</sup> Million Kilowatt-hours

**Table A-93: Energy Consumption by Fuel and Vehicle Type (Tbtu)**

| Fuel/Vehicle Type <sup>a</sup>                          | 1990          | 1995          | 2000          | 2001          | 2002          | 2003          | 2004          | 2005          | 2006          | 2007 <sup>b</sup> | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|-------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Motor Gasoline<sup>c, d</sup></b>                    | <b>13,810</b> | <b>14,687</b> | <b>16,031</b> | <b>16,208</b> | <b>16,531</b> | <b>16,585</b> | <b>16,802</b> | <b>16,671</b> | <b>16,426</b> | <b>16,259</b>     | <b>15,548</b> | <b>15,441</b> | <b>15,315</b> | <b>14,982</b> | <b>14,924</b> | <b>14,937</b> | <b>15,192</b> |
| Passenger Cars  | 8,725         | 8,442         | 9,045         | 9,120         | 9,294         | 8,996         | 8,943         | 9,237         | 8,854         | 11,017            | 10,533        | 10,434        | 10,348        | 10,272        | 10,253        | 10,253        | 10,282        |
| Light-Duty Trucks                                       | 4,340         | 5,512         | 6,303         | 6,363         | 6,507         | 6,931         | 7,197         | 6,720         | 6,854         | 4,343             | 4,112         | 4,162         | 4,136         | 3,930         | 3,888         | 3,892         | 4,114         |
| Motorcycles   | 24            | 25            | 26            | 24            | 24            | 23            | 24            | 23            | 26            | 59                | 61            | 58            | 51            | 50            | 57            | 54            | 53            |
| Buses   | 5             | 5             | 5             | 5             | 5             | 4             | 6             | 5             | 5             | 10                | 10            | 10            | 10            | 10            | 11            | 12            | 12            |
| Medium- and Heavy-Duty Trucks                           | 544           | 506           | 508           | 495           | 501           | 431           | 435           | 491           | 495           | 642               | 649           | 597           | 593           | 545           | 542           | 554           | 560           |
| Recreational Boats <sup>e</sup>                         | 172           | 197           | 143           | 201           | 200           | 199           | 197           | 195           | 192           | 188               | 183           | 180           | 176           | 174           | 173           | 172           | 171           |
| <b>Distillate Fuel Oil (Diesel Fuel)<sup>a, d</sup></b> | <b>3,555</b>  | <b>4,379</b>  | <b>5,437</b>  | <b>5,411</b>  | <b>5,590</b>  | <b>5,705</b>  | <b>5,910</b>  | <b>6,186</b>  | <b>6,334</b>  | <b>6,395</b>      | <b>6,059</b>  | <b>5,488</b>  | <b>5,706</b>  | <b>5,814</b>  | <b>5,780</b>  | <b>5,866</b>  | <b>6,052</b>  |
| Passenger Cars  | 107           | 106           | 49            | 50            | 50            | 57            | 58            | 57            | 56            | 55                | 50            | 49            | 51            | 55            | 55            | 55            | 56            |
| Light-Duty Trucks                                       | 155           | 201           | 272           | 281           | 296           | 367           | 391           | 349           | 361           | 183               | 163           | 162           | 169           | 176           | 175           | 174           | 188           |
| Buses   | 108           | 118           | 138           | 126           | 119           | 129           | 182           | 143           | 143           | 209               | 198           | 184           | 182           | 195           | 208           | 210           | 225           |
| Medium- and Heavy-Duty Trucks                           | 2,576         | 3,220         | 4,181         | 4,174         | 4,353         | 4,370         | 4,516         | 4,870         | 4,986         | 5,168             | 4,917         | 4,455         | 4,634         | 4,656         | 4,657         | 4,733         | 4,885         |
| Recreational Boats                                      | 26            | 32            | 37            | 39            | 40            | 41            | 42            | 43            | 44            | 45                | 46            | 47            | 48            | 49            | 50            | 51            | 52            |
| Ships and Other Boats                                   | 102           | 167           | 190           | 172           | 166           | 163           | 111           | 108           | 100           | 109               | 106           | 106           | 100           | 137           | 101           | 102           | 83            |
| Rail <sup>f</sup>                                       | 480           | 535           | 569           | 571           | 566           | 579           | 610           | 616           | 644           | 625               | 580           | 486           | 523           | 547           | 534           | 543           | 564           |
| <b>Jet Fuel<sup>g</sup></b>                             | <b>2,590</b>  | <b>2,429</b>  | <b>2,700</b>  | <b>2,626</b>  | <b>2,565</b>  | <b>2,482</b>  | <b>2,585</b>  | <b>2,622</b>  | <b>2,524</b>  | <b>2,485</b>      | <b>2,396</b>  | <b>2,134</b>  | <b>2,097</b>  | <b>2,030</b>  | <b>1,985</b>  | <b>2,037</b>  | <b>2,057</b>  |
| Commercial Aircraft                                     | 1,562         | 1,638         | 1,981         | 1,771         | 1,725         | 1,747         | 1,775         | 1,887         | 1,948         | 1,986             | 1,809         | 1,699         | 1,611         | 1,629         | 1,611         | 1,624         | 1,638         |
| General Aviation Aircraft                               | 545           | 454           | 427           | 537           | 556           | 449           | 515           | 484           | 350           | 276               | 362           | 241           | 314           | 256           | 224           | 274           | 226           |
| Military Aircraft <sup>a</sup>                          | 484           | 337           | 293           | 318           | 285           | 287           | 295           | 251           | 227           | 224               | 225           | 194           | 173           | 145           | 150           | 138           | 193           |
| <b>Aviation Gasoline<sup>g</sup></b>                    | <b>45</b>     | <b>40</b>     | <b>36</b>     | <b>35</b>     | <b>34</b>     | <b>30</b>     | <b>31</b>     | <b>35</b>     | <b>33</b>     | <b>32</b>         | <b>28</b>     | <b>27</b>     | <b>27</b>     | <b>27</b>     | <b>25</b>     | <b>22</b>     | <b>22</b>     |
| General Aviation Aircraft                               | 45            | 40            | 36            | 35            | 34            | 30            | 31            | 35            | 33            | 32                | 28            | 27            | 27            | 27            | 25            | 22            | 22            |
| <b>Residual Fuel Oil<sup>g, h</sup></b>                 | <b>300</b>    | <b>387</b>    | <b>443</b>    | <b>159</b>    | <b>228</b>    | <b>99</b>     | <b>186</b>    | <b>256</b>    | <b>306</b>    | <b>386</b>        | <b>271</b>    | <b>186</b>    | <b>272</b>    | <b>258</b>    | <b>211</b>    | <b>201</b>    | <b>73</b>     |
| Ships and Other Boats                                   | 300           | 387           | 443           | 159           | 228           | 99            | 186           | 256           | 306           | 386               | 271           | 186           | 272           | 258           | 211           | 201           | 73            |
| <b>Natural Gas<sup>g</sup></b>                          | <b>680</b>    | <b>724</b>    | <b>672</b>    | <b>658</b>    | <b>699</b>    | <b>627</b>    | <b>602</b>    | <b>624</b>    | <b>625</b>    | <b>663</b>        | <b>692</b>    | <b>715</b>    | <b>719</b>    | <b>734</b>    | <b>780</b>    | <b>887</b>    | <b>899</b>    |
| Passenger Cars  | -             | 2             | -             | -             | -             | -             | -             | -             | -             | -                 | -             | -             | -             | -             | -             | -             | -             |
| Light-Duty Trucks                                       | -             | -             | -             | -             | -             | -             | -             | -             | -             | -                 | -             | -             | -             | -             | -             | -             | -             |
| Buses   | -             | 1             | 8             | 9             | 12            | 14            | 16            | 16            | 16            | 19                | 21            | 22            | 20            | 20            | 20            | 21            | 21            |
| Pipelines   | 680           | 721           | 664           | 649           | 687           | 614           | 586           | 608           | 609           | 645               | 672           | 693           | 699           | 713           | 760           | 867           | 878           |
| <b>LPG<sup>g</sup></b>                                  | <b>23</b>     | <b>18</b>     | <b>12</b>     | <b>14</b>     | <b>14</b>     | <b>18</b>     | <b>19</b>     | <b>28</b>     | <b>27</b>     | <b>22</b>         | <b>40</b>     | <b>28</b>     | <b>29</b>     | <b>34</b>     | <b>37</b>     | <b>44</b>     | <b>44</b>     |
| Buses   | -             | -             | -             | -             | -             | -             | -             | -             | -             | -                 | -             | -             | -             | -             | -             | -             | -             |
| Light-Duty Trucks                                       | 9             | 8             | 8             | 9             | 10            | 12            | 14            | 21            | 20            | 16                | 29            | 19            | 21            | 24            | 27            | 31            | 31            |
| Medium- and Heavy-Duty Trucks                           | 14            | 9             | 4             | 4             | 4             | 5             | 5             | 7             | 8             | 6                 | 11            | 9             | 9             | 10            | 11            | 13            | 13            |
| <b>Electricity<sup>g</sup></b>                          | <b>16</b>     | <b>17</b>     | <b>18</b>     | <b>20</b>     | <b>19</b>     | <b>23</b>     | <b>25</b>     | <b>26</b>     | <b>25</b>     | <b>28</b>         | <b>26</b>     | <b>27</b>     | <b>26</b>     | <b>26</b>     | <b>25</b>     | <b>26</b>     | <b>26</b>     |
| Rail  | 16            | 17            | 18            | 20            | 19            | 23            | 25            | 26            | 25            | 28                | 26            | 27            | 26            | 26            | 25            | 26            | 26            |
| <b>Total</b>  | <b>21,019</b> | <b>22,681</b> | <b>25,350</b> | <b>25,131</b> | <b>25,680</b> | <b>25,570</b> | <b>26,160</b> | <b>26,449</b> | <b>26,302</b> | <b>26,270</b>     | <b>25,061</b> | <b>24,045</b> | <b>24,192</b> | <b>23,905</b> | <b>23,768</b> | <b>24,022</b> | <b>24,369</b> |

-Unreported or zero

<sup>a</sup> Note that updates to the distillate fuel oil heat content data from EIA for years 1993 through present resulted in changes to the time series for energy consumption and emissions compared to the previous Inventory.

<sup>b</sup> In 2011, FHWA changed its methodology for Table VM-1, which impacts estimates for the 2007-2014 time period. These methodological changes include how on-road vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. This resulted in large changes in fuel consumption data by vehicle class between 2006 and 2007.

<sup>c</sup> Figures do not include ethanol blended in motor gasoline or biodiesel blended into distillate fuel oil. Net carbon fluxes associated with ethanol are accounted for in the Land Use, Land-Use Change and Forestry chapter.

<sup>d</sup> Gasoline and diesel highway vehicle fuel consumption estimates are based on data from FHWA Highway Statistics Table VM-1 and MF-27 (FHWA 1996 through 2015). These fuel consumption estimates are combined with estimates of fuel shares by vehicle type from DOE's TEDB Annex Tables A.1 through A.6 (DOE 1993 through 2015). TEDB data for 2014 has not been published yet, therefore 2013 data is used as a proxy.

<sup>e</sup> Fluctuations in recreational boat gasoline estimates reflect the use of this category to reconcile bottom-up values with EIA total gasoline estimates.

<sup>f</sup> Class II and Class III diesel consumption data for 2014 is not available yet, therefore 2013 data is used as a proxy.

<sup>g</sup> Estimated based on EIA transportation sector energy estimates, with bottom-up data used for apportionment to modes.

<sup>h</sup> Fluctuations in reported fuel consumption may reflect data collection problems. Residual fuel oil for ships and other boats data is based on EIA's December 2015 Monthly Energy Review data.

**Table A-94: Transportation Sector Biofuel Consumption by Fuel Type (million gallons)**

| Fuel Type | 1990 | 1995  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  | 2008  | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   |
|-----------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------|--------|--------|--------|--------|--------|
| Ethanol   | 712  | 1,326 | 1,590 | 1,660 | 1,975 | 2,689 | 3,375 | 3,860 | 5,207 | 6,563 | 9,263 | 10,537 | 12,282 | 12,329 | 12,324 | 12,646 | 12,900 |
| Biodiesel | NA   | NA    | NA    | 10    | 16    | 14    | 27    | 91    | 261   | 354   | 304   | 322    | 260    | 886    | 899    | 1,429  | 1,417  |

NA – Not Available

Note: According to the MER, there was no biodiesel consumption prior to 2001.

## Estimates of CH<sub>4</sub> and N<sub>2</sub>O Emissions

Mobile source emissions of greenhouse gases other than CO<sub>2</sub> are reported by transport mode (e.g., road, rail, aviation, and waterborne), vehicle type, and fuel type. Emissions estimates of CH<sub>4</sub> and N<sub>2</sub>O were derived using a methodology similar to that outlined in the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006).

Activity data were obtained from a number of U.S. government agencies and other publications. Depending on the category, these basic activity data included fuel consumption and vehicle miles traveled (VMT). These estimates were then multiplied by emission factors, expressed as grams per unit of fuel consumed or per vehicle mile.

### Methodology for On-Road Gasoline and Diesel Vehicles

#### Step 1: Determine Vehicle Miles Traveled by Vehicle Type, Fuel Type, and Model Year

VMT by vehicle type (e.g., passenger cars, light-duty trucks, medium- and heavy-duty trucks,<sup>44</sup> buses, and motorcycles) were obtained from the Federal Highway Administration's (FHWA) *Highway Statistics* (FHWA 1996 through 2015).<sup>45</sup> As these vehicle categories are not fuel-specific, VMT for each vehicle type was disaggregated by fuel type (gasoline, diesel) so that the appropriate emission factors could be applied. VMT from *Highway Statistics* Table VM-1 (FHWA 1996 through 2015) was allocated to fuel types (gasoline, diesel, other) using historical estimates of fuel shares reported in the Appendix to the *Transportation Energy Data Book, Tables A.5 and A.6* (DOE 1993 through 2015). These fuel shares are drawn from various sources, including the Vehicle Inventory and Use Survey, the National Vehicle Population Profile, and the American Public Transportation Association. Fuel shares were first adjusted proportionately such that gasoline and diesel shares for each vehicle/fuel type category equaled 100 percent of national VMT. VMT for alternative fuel vehicles (AFVs) was calculated separately, and the methodology is explained in the following section on AFVs. Estimates of VMT from AFVs were then subtracted from the appropriate total VMT estimates to develop the final VMT estimates by vehicle/fuel type category.<sup>46</sup> The resulting national VMT estimates for gasoline and diesel on-road vehicles are presented in Table A-95 and Table A-96, respectively.

Total VMT for each on-road category (i.e., gasoline passenger cars, light-duty gasoline trucks, heavy-duty gasoline vehicles, diesel passenger cars, light-duty diesel trucks, medium- and heavy-duty diesel vehicles, and motorcycles) were distributed across 30 model years shown for 2014 in Table A-97. This distribution was derived by weighting the appropriate age distribution of the U.S. vehicle fleet according to vehicle registrations by the average annual age-specific vehicle mileage accumulation of U.S. vehicles. Age distribution values were obtained from EPA's MOBILE6 model for all years before 1999 (EPA 2000) and EPA's MOVES2014a model for years 2009 forward (EPA 2015d).<sup>47</sup> Age-specific vehicle mileage accumulations were also obtained from EPA's MOVES2014a model (EPA 2015d).<sup>48</sup>

#### Step 2: Allocate VMT Data to Control Technology Type

VMT by vehicle type for each model year was distributed across various control technologies as shown in Table A-103 through Table A-106. The categories "EPA Tier 0" and "EPA Tier 1" were used instead of the early three-way catalyst and advanced three-way catalyst categories, respectively, as defined in the *Revised 1996 IPCC Guidelines*. EPA Tier 0, EPA Tier 1, EPA Tier 2, and LEV refer to U.S. emission regulations, rather than control technologies; however, each does correspond to particular combinations of control technologies and engine design. EPA Tier 2 and its predecessors EPA

---

<sup>44</sup> Medium- and heavy-duty trucks correspond to FHWA's reporting categories of single-unit trucks and combination trucks. Single-unit trucks are defined as single frame trucks that have 2-axes and at least 6 tires or a gross vehicle weight rating (GVWR) exceeding 10,000 lbs.

<sup>45</sup> In 2011 FHWA changed its methods for estimated vehicle miles traveled (VMT) and related data. These methodological changes included how vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. These changes were first incorporated for the 2010 Inventory and apply to the 2007-14 time period. This resulted in large changes in VMT data by vehicle class, thus leading to a shift in emissions among on-road vehicle classes. For example, the category "Passenger Cars" has been replaced by "Light-duty Vehicles-Short Wheelbase" and "Other 2 axle-4 Tire Vehicles" has been replaced by "Light-duty Vehicles, Long Wheelbase." This change in vehicle classification has moved some smaller trucks and sport utility vehicles from the light truck category to the passenger vehicle category in this emission inventory. These changes are reflected in a large drop in light-truck emissions between 2006 and 2007.

<sup>46</sup> In Inventories through 2002, gasoline-electric hybrid vehicles were considered part of an "alternative fuel and advanced technology" category. However, vehicles are now only separated into gasoline, diesel, or alternative fuel categories, and gas-electric hybrids are now considered within the gasoline vehicle category.

<sup>47</sup> Age distributions were held constant for the period 1990-1998, and reflect a 25-year vehicle age span. EPA (2015b) provides a variable age distribution and 31-year vehicle age span beginning in year 1999.

<sup>48</sup> The updated vehicle distribution and mileage accumulation rates by vintage obtained from the MOVES 2014a model resulted in a decrease in emissions due to more miles driven by newer light-duty gasoline vehicles.

Tier 1 and Tier 0 apply to vehicles equipped with three-way catalysts. The introduction of “early three-way catalysts,” and “advanced three-way catalysts,” as described in the *Revised 1996 IPCC Guidelines*, roughly correspond to the introduction of EPA Tier 0 and EPA Tier 1 regulations (EPA 1998b).<sup>49</sup> EPA Tier 2 regulations affect vehicles produced starting in 2004 and are responsible for a noticeable decrease in N<sub>2</sub>O emissions compared EPA Tier 1 emissions technology (EPA 1999b).

Control technology assignments for light and heavy-duty conventional fuel vehicles for model years 1972 (when regulations began to take effect) through 1995 were estimated in EPA (1998). Assignments for 1998 through 2014 were determined using confidential engine family sales data submitted to EPA (EPA 2015f). Vehicle classes and emission standard tiers to which each engine family was certified were taken from annual certification test results and data (EPA 2015e). This information was used to determine the fraction of sales of each class of vehicle that met EPA Tier 0, EPA Tier 1, Tier 2, and LEV standards. Assignments for 1996 and 1997 were estimated based on the fact that EPA Tier 1 standards for light-duty vehicles were fully phased in by 1996. Tier 2 began initial phase-in by 2004.

### **Step 3: Determine CH<sub>4</sub> and N<sub>2</sub>O Emission Factors by Vehicle, Fuel, and Control Technology Type**

Emission factors for gasoline and diesel on-road vehicles utilizing Tier 2 and Low Emission Vehicle (LEV) technologies were developed by ICF (2006b); all other gasoline and diesel on-road vehicle emissions factors were developed by ICF (2004). These factors were based on EPA, CARB and Environment Canada laboratory test results of different vehicle and control technology types. The EPA, CARB and Environment Canada tests were designed following the Federal Test Procedure (FTP), which covers three separate driving segments, since vehicles emit varying amounts of GHGs depending on the driving segment. These driving segments are: (1) a transient driving cycle that includes cold start and running emissions, (2) a cycle that represents running emissions only, and (3) a transient driving cycle that includes hot start and running emissions. For each test run, a bag was affixed to the tailpipe of the vehicle and the exhaust was collected; the content of this bag was later analyzed to determine quantities of gases present. The emission characteristics of Segment 2 was used to define running emissions, and subtracted from the total FTP emissions to determine start emissions. These were then recombined based upon MOBILE6.2’s ratio of start to running emissions for each vehicle class to approximate average driving characteristics.

### **Step 4: Determine the Amount of CH<sub>4</sub> and N<sub>2</sub>O Emitted by Vehicle, Fuel, and Control Technology Type**

Emissions of CH<sub>4</sub> and N<sub>2</sub>O were then calculated by multiplying total VMT by vehicle, fuel, and control technology type by the emission factors developed in Step 3.

### **Methodology for Alternative Fuel Vehicles (AFVs)**

#### **Step 1: Determine Vehicle Miles Traveled by Vehicle and Fuel Type**

VMT for alternative fuel and advanced technology vehicles were calculated from “Methodology for Highway Vehicle Alternative Fuel GHG Projections Estimates” (Browning, 2015). Alternative Fuels include Compressed Natural Gas (CNG), Liquid Natural Gas (LNG), Liquefied Petroleum Gas (LPG), Ethanol, Methanol, Biodiesel, Hydrogen and Electricity. Most of the vehicles that use these fuels run on an Internal Combustion Engine (ICE) powered by the alternative fuel, although many of the vehicles can run on either the alternative fuel or gasoline (or diesel), or some combination.<sup>50</sup> Except for electric vehicles and plug-in hybrid vehicles, the alternative fuel vehicle VMT were calculated using the Energy Information Administration (EIA) Alternative Fuel Vehicle Data. The EIA data provides vehicle counts and fuel use for fleet vehicles used by electricity providers, federal agencies, natural gas providers, propane providers, state agencies and transit agencies for calendar years 2003 through 2013. For 1992 to 2002, EIA Data Tables were used to estimate fuel consumption and vehicle counts by vehicle type. These tables give total vehicle fuel use and vehicle counts by fuel and calendar year for the United States over the period 1992 through 2010. Breakdowns by vehicle type for 1992 through 2002 (both fuel consumed and vehicle counts) were assumed to be at the same ratio as for 2003 where data existed. For 1990, 1991 and 2014, fuel consumed by alternative fuel and vehicle type were extrapolated based on a regression analysis using the best curve fit based upon R<sup>2</sup> using the nearest 5 years of data.

For the current Inventory, counts of electric vehicles (EVs) and plug-in hybrid-electric vehicles (PHEVs) were taken from data compiled by the Electric Drive Transportation Association from 2011 to 2014 (EDTA 2015). EVs were

<sup>49</sup> For further description, see “Definitions of Emission Control Technologies and Standards” section of this annex below.

<sup>50</sup> Fuel types used in combination depend on the vehicle class. For light-duty vehicles, gasoline is generally blended with ethanol and diesel is blended with biodiesel; dual-fuel vehicles can run on gasoline or an alternative fuel – either natural gas or LPG – but not at the same time, while flex-fuel vehicles are designed to run on E85 (85 percent ethanol) or gasoline, or any mixture of the two in between. Heavy-duty vehicles are more likely to run on diesel fuel, natural gas, or LPG.

divided into cars and trucks using confidential engine family sales data submitted to EPA (EPA 2015f). Fuel use per vehicle for personal EVs and PHEVs were assumed to be the same as those for the public fleet vehicles surveyed by EIA and provided in their data tables.

Because AFVs run on different fuel types, their fuel use characteristics are not directly comparable. Accordingly, fuel economy for each vehicle type is expressed in gasoline equivalent terms, i.e., how much gasoline contains the equivalent amount of energy as the alternative fuel. Energy economy ratios (the ratio of the gasoline equivalent fuel economy of a given technology to that of conventional gasoline or diesel vehicles) were taken from the Argonne National Laboratory's GREET2015 model (ANL 2015). These ratios were used to estimate fuel economy in miles per gasoline gallon equivalent for each alternative fuel and vehicle type. Energy use per fuel type was then divided among the various weight categories and vehicle technologies that use that fuel. Total VMT per vehicle type for each calendar year was then determined by dividing the energy usage by the fuel economy. Note that for AFVs capable of running on both/either traditional and alternative fuels, the VMT given reflects only those miles driven that were powered by the alternative fuel, as explained in Browning (2015). VMT estimates for AFVs by vehicle category (passenger car, light-duty truck, medium-duty and heavy-duty vehicles) are shown in Table A-97, while more detailed estimates of VMT by control technology are shown in Table A-98.

### **Step 2: Determine CH<sub>4</sub> and N<sub>2</sub>O Emission Factors by Vehicle and Alternative Fuel Type**

Methane and N<sub>2</sub>O emission factors for alternative fuel vehicles (AFVs) are calculated according to studies by Argonne National Laboratory (2006) and Lipman & Delucchi (2002), and are reported in ICF (2006a). In these studies, N<sub>2</sub>O and CH<sub>4</sub> emissions for AFVs were expressed as a multiplier corresponding to conventional vehicle counterpart emissions. Emission estimates in these studies represent the current AFV fleet and were compared against Tier 1 emissions from light-duty gasoline vehicles to develop new multipliers. Alternative fuel heavy-duty vehicles were compared against gasoline heavy-duty vehicles as most alternative fuel heavy-duty vehicles use catalytic after treatment and perform more like gasoline vehicles than diesel vehicles. These emission factors are shown in Table A-108.

### **Step 3: Determine the Amount of CH<sub>4</sub> and N<sub>2</sub>O Emitted by Vehicle and Fuel Type**

Emissions of CH<sub>4</sub> and N<sub>2</sub>O were calculated by multiplying total VMT for each vehicle and fuel type (Step 1) by the appropriate emission factors (Step 2).

### **Methodology for Non-Road Mobile Sources**

Methane and N<sub>2</sub>O emissions from non-road mobile sources were estimated by applying emission factors to the amount of fuel consumed by mode and vehicle type.

Activity data for non-road vehicles include annual fuel consumption statistics by transportation mode and fuel type, as shown in Table A-102. Consumption data for ships and other boats (i.e., vessel bunkering) were obtained from DHS (2008) and EIA (1991 through 2015) for distillate fuel, and DHS (2008) and EIA (2015) for residual fuel; marine transport fuel consumption data for U.S. Territories (EIA 2015) were added to domestic consumption, and this total was reduced by the amount of fuel used for international bunkers.<sup>51</sup> Gasoline consumption by recreational boats was obtained from the NONROAD component of EPA's MOVES2014a model (EPA 2015d). Annual diesel consumption for Class I rail was obtained from the Association of American Railroads (AAR 2008 through 2015), diesel consumption from commuter rail was obtained from APTA (2007 through 2015) and Gaffney (2007), and consumption by Class II and III rail was provided by Benson (2002 through 2004) and Whorton (2006 through 2013).<sup>52</sup> Diesel consumption by commuter and intercity rail was obtained from DOE (1993 through 2015). Data on the consumption of jet fuel and aviation gasoline in aircraft were obtained from EIA (2016) and FAA (2016), as described in Annex 2.1: Methodology for Estimating Emissions of CO<sub>2</sub> from Fossil Fuel Combustion, and were reduced by the amount allocated to international bunker fuels (DESC 2015 and FAA 2016). Pipeline fuel consumption was obtained from EIA (2007 through 2015) (note: pipelines are a transportation source but are stationary, not mobile, sources). Data on fuel consumption by all non-transportation mobile sources were obtained from the NONROAD component of EPA's MOVES2014a model (EPA 2015d) and from FHWA (1996 through 2015) for gasoline consumption for trucks used off-road.<sup>53</sup>

---

<sup>51</sup> See International Bunker Fuels section of the Energy Chapter.

<sup>52</sup> Diesel consumption from Class II and Class III railroad were unavailable for 2014. Values are proxied from 2013, which is the last year the data was available.

<sup>53</sup> "Non-transportation mobile sources" are defined as any vehicle or equipment not used on the traditional road system, but excluding aircraft, rail and watercraft. This category includes snowmobiles, golf carts, riding lawn mowers, agricultural equipment, and trucks used for off-road purposes, among others.

Emissions of CH<sub>4</sub> and N<sub>2</sub>O from non-road mobile sources were calculated by multiplying U.S. default emission factors in the 2006 IPCC Guidelines by activity data for each source type (see Table A-109).

### Estimates of NO<sub>x</sub>, CO, and NMVOC Emissions

The emission estimates of NO<sub>x</sub>, CO, and NMVOCs from mobile combustion (transportation) were obtained from preliminary data (EPA 2015g), which, in final iteration, will be published on the EPA's National Emission Inventory (NEI) Air Pollutant Emission Trends web site. This EPA report provides emission estimates for these gases by fuel type using a procedure whereby emissions were calculated using basic activity data, such as amount of fuel delivered or miles traveled, as indicators of emissions. Table A-110 through Table A-112 provides complete emission estimates for 1990 through 2014.

**Table A-95: Vehicle Miles Traveled for Gasoline On-Road Vehicles (million miles)**

| Year              | Passenger Cars | Light-Duty Trucks | Heavy-Duty Vehicles <sup>b</sup> | Motorcycles |
|-------------------|----------------|-------------------|----------------------------------|-------------|
| 1990              | 1,391.4        | 554.8             | 25.8                             | 9.6         |
| 1991              | 1,341.9        | 627.8             | 25.4                             | 9.2         |
| 1992              | 1,355.1        | 683.4             | 25.1                             | 9.6         |
| 1993              | 1,356.8        | 721.0             | 24.9                             | 9.9         |
| 1994              | 1,387.7        | 739.2             | 25.3                             | 10.2        |
| 1995              | 1,421.0        | 763.0             | 25.1                             | 9.8         |
| 1996              | 1,455.1        | 788.6             | 24.4                             | 9.9         |
| 1997              | 1,489.0        | 821.6             | 24.0                             | 10.1        |
| 1998              | 1,537.1        | 837.7             | 24.1                             | 10.3        |
| 1999              | 1,559.6        | 868.3             | 24.3                             | 10.6        |
| 2000              | 1,592.2        | 887.6             | 24.2                             | 10.5        |
| 2001              | 1,620.0        | 905.9             | 23.9                             | 9.6         |
| 2002              | 1,650.0        | 926.8             | 23.9                             | 9.6         |
| 2003              | 1,663.5        | 944.1             | 24.2                             | 9.6         |
| 2004              | 1,691.1        | 985.5             | 24.6                             | 10.1        |
| 2005              | 1,699.6        | 998.8             | 24.8                             | 10.5        |
| 2006              | 1,681.8        | 1,038.6           | 24.8                             | 12.0        |
| 2007              | 2,093.7        | 562.8             | 34.2                             | 21.4        |
| 2008              | 2,014.5        | 580.9             | 35.0                             | 20.8        |
| 2009              | 2,005.5        | 592.4             | 32.5                             | 20.8        |
| 2010              | 2,015.4        | 597.4             | 32.3                             | 18.5        |
| 2011 <sup>a</sup> | 2,035.7        | 579.6             | 30.2                             | 18.5        |
| 2012              | 2,051.6        | 576.8             | 30.5                             | 21.4        |
| 2013              | 2,062.1        | 578.7             | 31.2                             | 20.4        |
| 2014              | 2,058.2        | 612.4             | 31.7                             | 20.0        |

Source: Derived from FHWA (1996 through 2014), Browning (2015).

<sup>a</sup> In 2011, FHWA changed its methodology for Table VM-1, which impacts estimates for the 2007 to 2014 time period. These methodological changes include how on-road vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. This resulted in large changes in VMT data by vehicle class between 2006 and 2007.

<sup>b</sup> Heavy-Duty Vehicles includes Medium-Duty Trucks, Heavy-Duty Trucks, and Buses.

Note: In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data. EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated in this year's inventory and apply to the 1990-2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes.

Note: Gasoline and diesel highway vehicle mileage are based on data from FHWA Highway Statistics Table VM-1 (FHWA 1996 through 2015). These mileage consumption estimates are combined with estimates of fuel shares by vehicle type from DOE's TEDB Annex Tables A.1 through A.6 (DOE 1993 through 2015). TEDB data for 2014 has not been published yet, therefore 2013 data is used as a proxy.

**Table A-96: Vehicle Miles Traveled for Diesel On-Road Vehicles (million miles)**

| Year              | Passenger Cars | Light-Duty Trucks | Heavy-Duty Vehicles <sup>a</sup> |
|-------------------|----------------|-------------------|----------------------------------|
| 1990              | 16.9           | 19.7              | 125.7                            |
| 1991              | 16.3           | 21.6              | 129.4                            |
| 1992              | 16.5           | 23.4              | 133.6                            |
| 1993              | 17.9           | 24.7              | 140.6                            |
| 1994              | 18.3           | 25.3              | 150.8                            |
| 1995              | 17.3           | 26.9              | 159.0                            |
| 1996              | 14.7           | 27.8              | 164.6                            |
| 1997              | 13.5           | 29.0              | 173.8                            |
| 1998              | 12.4           | 30.5              | 178.8                            |
| 1999              | 9.4            | 32.6              | 185.6                            |
| 2000              | 8.0            | 35.2              | 188.4                            |
| 2001              | 8.1            | 37.0              | 191.5                            |
| 2002              | 8.3            | 38.9              | 196.7                            |
| 2003              | 8.4            | 39.7              | 199.6                            |
| 2004              | 8.5            | 41.4              | 202.1                            |
| 2005              | 8.5            | 41.9              | 203.4                            |
| 2006              | 8.4            | 43.4              | 202.3                            |
| 2007              | 10.5           | 23.3              | 281.8                            |
| 2008              | 10.1           | 24.1              | 288.1                            |
| 2009              | 10.0           | 24.6              | 267.6                            |
| 2010              | 10.1           | 24.8              | 265.8                            |
| 2011 <sup>b</sup> | 10.1           | 23.3              | 245.6                            |
| 2012              | 10.1           | 23.1              | 247.9                            |
| 2013              | 10.1           | 22.6              | 250.5                            |
| 2014              | 10.0           | 24.0              | 255.0                            |

Source: Derived from FHWA (1996 through 2015), Browning (2015).

<sup>a</sup> Heavy-Duty Vehicles includes Medium-Duty Trucks, Heavy-Duty Trucks, and Buses.

<sup>b</sup> In 2011, FHWA changed its methodology for Table VM-1, which impacts estimates for the 2007 to 2014 time period. These methodological changes include how on-road vehicles are classified, moving from a system based on body-type to one that is based on wheelbase. This resulted in large changes in VMT data by vehicle class between 2006 and 2007.

Note: In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data. EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated in this year's inventory and apply to the 1990 to 2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes.

Note: Gasoline and diesel highway vehicle mileage are based on data from FHWA Highway Statistics Table VM-1 (FHWA 1996 through 2015). These mileage consumption estimates are combined with estimates of fuel shares by vehicle type from DOE's TEDB Annex Tables A.1 through A.6 (DOE 1993 through 2015).

**Table A-97: Vehicle Miles Traveled for Alternative Fuel On-Road Vehicles (million miles)**

| Year | Passenger Cars | Light-Duty Trucks | Heavy-Duty Vehicles <sup>a</sup> |
|------|----------------|-------------------|----------------------------------|
| 1990 | 0.0            | 0.1               | 0.5                              |
| 1991 | 0.0            | 0.1               | 0.5                              |
| 1992 | 0.0            | 0.1               | 0.4                              |
| 1993 | 0.0            | 0.1               | 0.6                              |
| 1994 | 0.1            | 0.1               | 0.5                              |
| 1995 | 0.1            | 0.1               | 0.5                              |
| 1996 | 0.1            | 0.1               | 0.5                              |
| 1997 | 0.1            | 0.1               | 0.5                              |
| 1998 | 0.1            | 0.1               | 0.5                              |
| 1999 | 0.1            | 0.1               | 0.5                              |
| 2000 | 0.1            | 0.2               | 0.6                              |
| 2001 | 0.2            | 0.2               | 0.7                              |
| 2002 | 0.2            | 0.3               | 0.8                              |
| 2003 | 0.3            | 0.3               | 0.9                              |
| 2004 | 0.2            | 0.3               | 1.0                              |
| 2005 | 0.3            | 0.3               | 1.3                              |
| 2006 | 0.3            | 0.5               | 2.2                              |
| 2007 | 0.2            | 0.6               | 2.7                              |

|      |     |     |     |
|------|-----|-----|-----|
| 2008 | 0.2 | 0.5 | 2.4 |
| 2009 | 0.2 | 0.5 | 2.5 |
| 2010 | 0.2 | 0.5 | 2.2 |
| 2011 | 0.5 | 1.2 | 5.6 |
| 2012 | 1.1 | 1.3 | 5.6 |
| 2013 | 2.3 | 2.0 | 8.5 |
| 2014 | 3.8 | 2.1 | 8.4 |

Source: Derived from Browning (2015).

<sup>a</sup> Heavy Duty-Vehicles includes medium-duty trucks, heavy-duty trucks, and buses.

Note: In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data. EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated in this year's inventory and apply to the 1990-2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes.

**Table A-98: Detailed Vehicle Miles Traveled for Alternative Fuel On-Road Vehicles (10<sup>6</sup> Miles)**

| Vehicle Type/Year         | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|---------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Light-Duty Cars</b>    | <b>4.0</b>   | <b>61.5</b>  | <b>115.9</b> | <b>150.0</b> | <b>222.9</b> | <b>267.6</b> | <b>249.4</b> | <b>254.7</b> | <b>296.3</b>   | <b>225.6</b>   | <b>219.5</b>   | <b>217.2</b>   | <b>235.5</b>   | <b>537.9</b>   | <b>1,065.8</b> | <b>2,314.6</b> | <b>3,815.1</b> |
| Methanol-Flex Fuel ICE    | +            | 48.9         | 15.2         | 11.5         | 8.8          | +            | +            | +            | +              | +              | +              | +              | +              | +              | +              | +              | +              |
| Ethanol-Flex Fuel ICE     | +            | 0.3          | 20.9         | 25.4         | 31.0         | 35.2         | 42.2         | 51.0         | 59.2           | 72.8           | 84.2           | 96.2           | 122.2          | 118.5          | 148.9          | 173.5          | 193.7          |
| CNG ICE                   | +            | 0.1          | 5.5          | 9.1          | 12.3         | 14.5         | 17.3         | 15.9         | 14.7           | 14.6           | 13.1           | 12.3           | 11.7           | 12.3           | 12.5           | 13.3           | 12.5           |
| CNG Bi-fuel               | +            | 0.2          | 18.0         | 29.5         | 40.3         | 47.6         | 53.5         | 40.6         | 25.4           | 19.3           | 13.0           | 10.1           | 8.1            | 7.1            | 4.5            | 3.5            | 2.3            |
| LPG ICE                   | 1.1          | 1.2          | 1.2          | 1.2          | 1.3          | 1.3          | 0.3          | 0.1          | 0.2            | 1.7            | 1.7            | 1.7            | +              | 0.2            | 0.2            | 0.4            | 1.0            |
| LPG Bi-fuel               | 2.8          | 3.0          | 3.0          | 3.0          | 3.2          | 3.3          | 7.9          | 3.3          | 3.7            | 1.7            | 1.6            | 1.8            | 1.2            | 0.3            | 0.3            | 0.2            | +              |
| Biodiesel (BD100)         | +            | +            | 1.0          | 2.0          | 3.0          | 2.3          | 4.4          | 14.6         | 41.4           | 50.2           | 39.1           | 46.4           | 39.4           | 149.4          | 180.3          | 310.3          | 333.2          |
| NEVs                      | +            | 7.5          | 49.8         | 66.4         | 119.4        | 158.7        | 121.2        | 127.2        | 149.4          | 63.6           | 64.2           | 46.8           | 51.8           | 58.1           | 48.4           | 52.7           | 58.3           |
| Electric Vehicle          | +            | 0.2          | 1.5          | 1.9          | 3.5          | 4.6          | 2.5          | 1.9          | 2.3            | 1.5            | 2.3            | 1.5            | 0.9            | 169.3          | 533.5          | 1,484.1        | 2,771.9        |
| SI PHEV - Electricity     | +            | +            | +            | +            | +            | +            | +            | +            | +              | +              | +              | +              | +              | 22.6           | 137.2          | 276.5          | 442.2          |
| Fuel Cell Hydrogen        | +            | +            | +            | +            | +            | +            | +            | +            | +              | 0.3            | 0.2            | 0.5            | 0.2            | 0.1            | 0.1            | 0.1            | 0.1            |
| <b>Light-Duty Trucks</b>  | <b>77.4</b>  | <b>93.3</b>  | <b>180.9</b> | <b>225.0</b> | <b>274.4</b> | <b>302.9</b> | <b>271.8</b> | <b>333.1</b> | <b>491.8</b>   | <b>556.3</b>   | <b>458.8</b>   | <b>511.5</b>   | <b>463.9</b>   | <b>1,223.3</b> | <b>1,341.2</b> | <b>2,036.1</b> | <b>2,086.2</b> |
| Ethanol-Flex Fuel ICE     | +            | 0.3          | 23.4         | 28.0         | 33.9         | 38.1         | 45.4         | 54.5         | 62.8           | 77.0           | 89.6           | 102.7          | 130.9          | 144.2          | 191.8          | 227.2          | 278.7          |
| CNG ICE                   | +            | 0.1          | 5.6          | 9.1          | 12.2         | 14.3         | 15.9         | 14.5         | 15.2           | 13.7           | 10.8           | 10.3           | 9.2            | 9.7            | 9.9            | 9.5            | 10.2           |
| CNG Bi-fuel               | +            | 0.4          | 47.2         | 76.6         | 103.4        | 121.0        | 84.4         | 72.8         | 68.8           | 61.3           | 26.3           | 22.0           | 20.6           | 19.7           | 16.0           | 17.4           | 17.2           |
| LPG ICE                   | 22.4         | 26.5         | 27.6         | 27.7         | 28.5         | 29.3         | 23.7         | 32.6         | 28.9           | 23.3           | 11.6           | 13.5           | 10.8           | 10.6           | 6.5            | 6.9            | 5.4            |
| LPG Bi-fuel               | 55.0         | 65.1         | 67.7         | 68.3         | 70.3         | 72.4         | 71.4         | 67.6         | 54.8           | 32.0           | 24.8           | 28.9           | 25.1           | 13.2           | 5.3            | 6.4            | 2.9            |
| LNG                       | +            | +            | 0.1          | 0.2          | 0.2          | 0.2          | 0.2          | 0.3          | 0.2            | 0.2            | 0.3            | 0.2            | +              | +              | +              | +              | +              |
| Biodiesel (BD100)         | +            | +            | 4.1          | 8.1          | 13.5         | 11.2         | 24.0         | 85.3         | 253.9          | 341.1          | 287.9          | 326.6          | 260.2          | 1,020.9        | 1,107.2        | 1,752.0        | 1,741.4        |
| Electric Vehicle          | +            | 0.8          | 5.3          | 7.0          | 12.5         | 16.5         | 6.8          | 5.6          | 7.1            | 7.7            | 7.5            | 7.3            | 7.0            | 4.6            | 4.4            | 16.4           | 30.1           |
| Fuel Cell Hydrogen        | +            | +            | +            | +            | +            | +            | +            | +            | +              | 0.1            | 0.1            | 0.2            | 0.1            | 0.3            | 0.2            | 0.2            | 0.2            |
| <b>Medium Duty Trucks</b> | <b>324.5</b> | <b>317.4</b> | <b>308.6</b> | <b>327.8</b> | <b>353.9</b> | <b>364.6</b> | <b>265.7</b> | <b>352.8</b> | <b>543.5</b>   | <b>640.9</b>   | <b>580.1</b>   | <b>607.0</b>   | <b>457.1</b>   | <b>1,414.9</b> | <b>1,475.6</b> | <b>2,335.2</b> | <b>2,360.6</b> |
| CNG ICE                   | +            | +            | 1.1          | 1.8          | 2.4          | 2.9          | 2.9          | 4.1          | 2.8            | 5.8            | 8.0            | 6.9            | 6.6            | 9.0            | 10.6           | 11.1           | 13.1           |
| CNG Bi-fuel               | +            | 0.1          | 10.5         | 17.2         | 23.3         | 27.4         | 14.6         | 15.5         | 13.0           | 11.4           | 9.7            | 8.0            | 7.5            | 7.3            | 8.3            | 8.7            | 10.2           |
| LPG ICE                   | 271.5        | 265.4        | 242.4        | 246.1        | 254.4        | 262.8        | 166.6        | 155.6        | 82.1           | 61.3           | 46.5           | 41.5           | 36.5           | 34.1           | 32.3           | 29.6           | 29.2           |
| LPG Bi-fuel               | 53.0         | 51.8         | 47.3         | 48.1         | 49.7         | 51.3         | 41.6         | 43.8         | 23.6           | 10.3           | 16.5           | 8.4            | 10.3           | 9.3            | 12.6           | 13.7           | 16.2           |
| LNG                       | +            | +            | +            | +            | +            | +            | +            | +            | +              | +              | +              | +              | +              | +              | +              | 0.1            | 0.1            |
| Biodiesel (BD100)         | +            | +            | 7.3          | 14.5         | 24.0         | 20.2         | 39.8         | 133.7        | 422.0          | 552.0          | 499.4          | 542.3          | 396.2          | 1,355.2        | 1,411.8        | 2,271.9        | 2,291.8        |
| <b>Heavy-Duty Trucks</b>  | <b>130.4</b> | <b>127.6</b> | <b>137.1</b> | <b>159.1</b> | <b>186.5</b> | <b>181.1</b> | <b>217.7</b> | <b>439.5</b> | <b>1,046.6</b> | <b>1,388.9</b> | <b>1,180.1</b> | <b>1,227.1</b> | <b>1,023.1</b> | <b>3,368.5</b> | <b>3,368.0</b> | <b>5,320.4</b> | <b>5,300.8</b> |
| Neat Ethanol ICE          | +            | +            | +            | +            | +            | 1.3          | 1.6          | 1.9          | 2.2            | 2.7            | 3.1            | 3.5            | 4.5            | 7.0            | 11.3           | 15.7           | 22.2           |
| CNG ICE                   | +            | +            | 0.9          | 1.5          | 2.0          | 2.4          | 1.9          | 1.9          | 2.7            | 2.9            | 2.7            | 3.4            | 3.6            | 3.6            | 4.1            | 5.0            | 6.1            |
| LPG ICE                   | 122.1        | 119.5        | 109.1        | 110.7        | 114.4        | 118.1        | 106.4        | 87.2         | 76.6           | 65.8           | 56.2           | 49.6           | 41.0           | 43.1           | 28.0           | 27.6           | 20.6           |
| LPG Bi-fuel               | 8.3          | 8.1          | 7.4          | 7.5          | 7.8          | 8.0          | 6.4          | 5.4          | 4.8            | 4.6            | 4.6            | 5.4            | 5.6            | 8.3            | 6.5            | 7.1            | 5.9            |
| LNG                       | +            | +            | +            | +            | +            | +            | +            | 0.1          | 0.1            | +              | 0.1            | +              | +              | +              | +              | +              | +              |
| Biodiesel (BD100)         | +            | +            | 19.7         | 39.3         | 62.3         | 51.2         | 101.4        | 343.0        | 960.2          | 1,312.8        | 1,113.4        | 1,165.1        | 968.4          | 3,306.4        | 3,318.2        | 5,265.1        | 5,246.0        |
| <b>Buses</b>              | <b>24.6</b>  | <b>46.3</b>  | <b>150.7</b> | <b>225.9</b> | <b>299.2</b> | <b>349.1</b> | <b>496.6</b> | <b>524.6</b> | <b>629.8</b>   | <b>625.6</b>   | <b>656.3</b>   | <b>685.3</b>   | <b>695.5</b>   | <b>779.6</b>   | <b>770.3</b>   | <b>833.7</b>   | <b>804.8</b>   |
| Neat Methanol ICE         | 7.8          | 12.7         | +            | +            | +            | +            | +            | +            | +              | +              | +              | +              | +              | +              | +              | +              | +              |
| Neat Ethanol ICE          | +            | 5.9          | 0.1          | +            | +            | +            | +            | +            | +              | +              | +              | +              | +              | +              | 0.1            | 0.1            | 0.1            |
| CNG ICE                   | +            | 1.1          | 104.1        | 170.3        | 231.7        | 271.8        | 392.8        | 413.5        | 481.7          | 509.8          | 546.2          | 581.7          | 605.4          | 637.1          | 628.3          | 650.1          | 623.7          |
| LPG ICE                   | 16.4         | 15.8         | 14.4         | 14.6         | 15.2         | 15.6         | 17.5         | 11.7         | 13.3           | 12.3           | 13.4           | 9.0            | 8.1            | 4.8            | 4.7            | 5.0            | 5.2            |

|                    |              |              |              |                |                |                |                |                |                |                |                |                |                |                |                |                 |                 |
|--------------------|--------------|--------------|--------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------------|-----------------|
| LNG                | 0.4          | 8.9          | 23.2         | 28.5           | 30.1           | 33.7           | 58.9           | 61.4           | 66.8           | 40.2           | 39.8           | 36.0           | 36.8           | 39.5           | 41.1           | 29.4            | 27.5            |
| Biodiesel (BD100)  | +            | +            | 0.8          | 1.6            | 2.6            | 2.0            | 4.1            | 13.2           | 38.9           | 53.3           | 46.6           | 51.7           | 38.1           | 90.1           | 90.7           | 143.5           | 142.8           |
| Electric           | +            | 1.8          | 8.2          | 10.9           | 19.6           | 26.0           | 23.2           | 24.8           | 29.0           | 9.9            | 10.2           | 6.8            | 7.0            | 7.7            | 5.1            | 5.3             | 5.2             |
| Fuel Cell Hydrogen | +            | +            | +            | +              | +              | +              | +              | +              | 0.1            | 0.1            | 0.1            | 0.1            | 0.2            | 0.3            | 0.3            | 0.3             | 0.3             |
| <b>Total VMT</b>   | <b>560.8</b> | <b>646.0</b> | <b>893.4</b> | <b>1,087.9</b> | <b>1,336.9</b> | <b>1,465.2</b> | <b>1,501.2</b> | <b>1,904.7</b> | <b>3,008.0</b> | <b>3,437.3</b> | <b>3,094.9</b> | <b>3,248.2</b> | <b>2,875.1</b> | <b>7,324.2</b> | <b>8,020.9</b> | <b>12,840.0</b> | <b>14,367.6</b> |

+ Does not exceed 0.05 million vehicle miles traveled

Note: Throughout the rest of this Inventory, medium-duty trucks are grouped with heavy-duty trucks; they are reported separately here because these two categories may run on a slightly different range of fuel types.

In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data.

EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated for the 2014

Inventory and apply to the 1990-2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes.

Source: Derived from Browning (2015).

**Table A-99: Age Distribution by Vehicle/Fuel Type for On-Road Vehicles,<sup>a</sup> 2014**

| Vehicle Age  | LDGV          | LDGT          | HDGV          | LDDV          | LDDT          | HDDV          | MC            |
|--------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| 0            | 7.1%          | 7.9%          | 6.2%          | 14.7%         | 8.6%          | 6.0%          | 7.2%          |
| 1            | 6.8%          | 7.5%          | 5.6%          | 14.1%         | 8.1%          | 5.4%          | 6.9%          |
| 2            | 6.6%          | 7.1%          | 5.1%          | 13.6%         | 7.8%          | 5.0%          | 6.5%          |
| 3            | 4.1%          | 4.8%          | 2.9%          | 8.4%          | 5.3%          | 3.1%          | 5.2%          |
| 4            | 4.6%          | 4.2%          | 1.9%          | 8.1%          | 3.1%          | 2.1%          | 4.7%          |
| 5            | 4.2%          | 3.2%          | 1.7%          | 5.4%          | 2.8%          | 2.5%          | 4.9%          |
| 6            | 5.3%          | 5.4%          | 3.4%          | 0.5%          | 6.9%          | 3.8%          | 8.6%          |
| 7            | 5.7%          | 5.7%          | 3.1%          | 0.3%          | 6.1%          | 7.4%          | 7.7%          |
| 8            | 5.3%          | 5.7%          | 4.4%          | 6.4%          | 7.8%          | 6.4%          | 7.3%          |
| 9            | 5.2%          | 5.8%          | 3.4%          | 4.4%          | 6.6%          | 5.9%          | 6.4%          |
| 10           | 4.8%          | 5.8%          | 4.3%          | 2.6%          | 5.9%          | 4.1%          | 5.4%          |
| 11           | 5.0%          | 5.2%          | 3.7%          | 3.4%          | 5.3%          | 3.6%          | 4.6%          |
| 12           | 4.9%          | 4.9%          | 3.7%          | 3.7%          | 4.5%          | 2.9%          | 4.1%          |
| 13           | 4.6%          | 4.3%          | 3.1%          | 2.4%          | 5.1%          | 3.8%          | 3.4%          |
| 14           | 4.5%          | 4.0%          | 6.0%          | 2.0%          | 2.7%          | 5.9%          | 2.8%          |
| 15           | 3.5%          | 3.4%          | 5.8%          | 1.1%          | 3.9%          | 4.7%          | 2.1%          |
| 16           | 2.8%          | 2.7%          | 2.4%          | 1.1%          | 1.4%          | 3.2%          | 1.8%          |
| 17           | 2.6%          | 2.3%          | 4.5%          | 0.4%          | 1.8%          | 3.0%          | 1.7%          |
| 18           | 2.1%          | 1.7%          | 2.7%          | 0.4%          | 1.4%          | 2.7%          | 1.5%          |
| 19           | 2.1%          | 1.6%          | 3.7%          | 0.3%          | 1.0%          | 3.3%          | 1.1%          |
| 20           | 1.6%          | 1.4%          | 2.9%          | 0.0%          | 0.6%          | 2.5%          | 1.3%          |
| 21           | 1.4%          | 1.0%          | 2.3%          | 0.2%          | 0.6%          | 1.8%          | 1.1%          |
| 22           | 1.1%          | 0.8%          | 1.8%          | 0.2%          | 0.6%          | 1.3%          | 0.9%          |
| 23           | 0.9%          | 0.6%          | 1.5%          | 0.5%          | 0.3%          | 1.2%          | 0.7%          |
| 24           | 0.8%          | 0.6%          | 2.0%          | 0.1%          | 0.3%          | 1.5%          | 0.6%          |
| 25           | 0.6%          | 0.6%          | 2.4%          | 0.1%          | 0.2%          | 1.5%          | 0.4%          |
| 26           | 0.5%          | 0.5%          | 2.0%          | 0.0%          | 0.2%          | 1.3%          | 0.4%          |
| 27           | 0.4%          | 0.4%          | 1.8%          | 0.9%          | 0.1%          | 1.1%          | 0.3%          |
| 28           | 0.3%          | 0.4%          | 1.8%          | 0.5%          | 0.3%          | 0.8%          | 0.3%          |
| 29           | 0.3%          | 0.3%          | 1.4%          | 1.6%          | 0.2%          | 0.8%          | 0.2%          |
| 30           | 0.3%          | 0.2%          | 2.4%          | 2.5%          | 0.2%          | 1.3%          | 0.2%          |
| <b>Total</b> | <b>100.0%</b> |

Source: EPA (2015b).

Note: This year's Inventory includes updated vehicle population data based on the MOVES 2014a Model.

<sup>a</sup> The following abbreviations correspond to vehicle types: LDGV (light-duty gasoline vehicles), LDGT (light-duty gasoline trucks), HDGV (heavy-duty gasoline vehicles), LDDV (light-duty diesel vehicles), LDDT (light-duty diesel trucks), HDDV (heavy-duty diesel vehicles), and MC (motorcycles).**Table A-100: Annual Average Vehicle Mileage Accumulation per Vehicle<sup>a</sup> (miles)**

| Vehicle Age | LDGV   | LDGT   | HDGV   | LDDV   | LDDT   | HDDV   | MC <sup>b</sup> |
|-------------|--------|--------|--------|--------|--------|--------|-----------------|
| 0           | 13,809 | 15,607 | 19,166 | 13,809 | 15,608 | 41,091 | 7,566           |
| 1           | 13,547 | 15,314 | 19,169 | 13,547 | 15,314 | 40,875 | 4,040           |
| 2           | 13,263 | 14,983 | 19,173 | 13,263 | 14,983 | 41,583 | 3,057           |
| 3           | 12,960 | 14,621 | 17,828 | 12,960 | 14,621 | 48,627 | 2,527           |
| 4           | 12,641 | 14,231 | 15,436 | 12,641 | 14,231 | 47,399 | 2,187           |
| 5           | 12,307 | 13,816 | 14,949 | 12,307 | 13,816 | 48,996 | 1,944           |
| 6           | 11,959 | 13,380 | 15,707 | 11,959 | 13,380 | 34,924 | 1,763           |
| 7           | 11,601 | 12,928 | 13,175 | 11,601 | 12,928 | 46,171 | 1,619           |
| 8           | 11,234 | 12,463 | 13,437 | 11,234 | 12,463 | 38,948 | 1,498           |
| 9           | 10,861 | 11,990 | 11,436 | 10,861 | 11,990 | 36,966 | 1,400           |
| 10          | 10,483 | 11,511 | 11,274 | 10,483 | 11,511 | 30,576 | 1,316           |
| 11          | 10,104 | 11,031 | 10,074 | 10,104 | 11,031 | 29,579 | 1,241           |
| 12          | 9,725  | 10,555 | 9,275  | 9,725  | 10,555 | 25,374 | 1,180           |
| 13          | 9,347  | 10,087 | 7,987  | 9,347  | 10,087 | 23,342 | 1,120           |
| 14          | 8,974  | 9,630  | 8,500  | 8,974  | 9,630  | 21,710 | 1,067           |
| 15          | 8,608  | 9,187  | 7,310  | 8,608  | 9,187  | 18,601 | 1,021           |
| 16          | 8,250  | 8,764  | 6,062  | 8,250  | 8,764  | 16,648 | 984             |

|    |       |       |       |       |       |        |     |
|----|-------|-------|-------|-------|-------|--------|-----|
| 17 | 7,902 | 8,362 | 5,478 | 7,902 | 8,362 | 12,487 | 946 |
| 18 | 7,567 | 7,989 | 5,183 | 7,567 | 7,989 | 13,300 | 908 |
| 19 | 7,247 | 7,648 | 5,149 | 7,247 | 7,648 | 11,161 | 878 |
| 20 | 6,945 | 7,340 | 4,657 | 6,945 | 7,340 | 9,880  | 847 |
| 21 | 6,662 | 7,072 | 4,650 | 6,662 | 7,072 | 9,275  | 825 |
| 22 | 6,400 | 6,845 | 3,963 | 6,400 | 6,845 | 8,528  | 802 |
| 23 | 6,161 | 6,668 | 3,920 | 6,161 | 6,668 | 7,821  | 757 |
| 24 | 5,949 | 6,540 | 3,693 | 5,949 | 6,540 | 6,747  | 711 |
| 25 | 5,764 | 6,465 | 3,484 | 5,764 | 6,465 | 5,858  | 666 |
| 26 | 5,609 | 6,451 | 3,157 | 5,609 | 6,451 | 5,308  | 613 |
| 27 | 5,486 | 6,451 | 2,920 | 5,486 | 6,451 | 5,090  | 567 |
| 28 | 5,397 | 6,451 | 2,736 | 5,397 | 6,451 | 4,296  | 537 |
| 29 | 5,345 | 6,451 | 2,724 | 5,345 | 6,451 | 3,475  | 499 |
| 30 | 5,345 | 6,451 | 2,408 | 5,345 | 6,451 | 2,598  | 462 |

Source: EPA (2015b).

<sup>a</sup> The following abbreviations correspond to vehicle types: LDGV (light-duty gasoline vehicles), LDGT (light-duty gasoline trucks), HDGV (heavy-duty gasoline vehicles), LDDV (light-duty diesel vehicles), LDDT (light-duty diesel trucks), HDDV (heavy-duty diesel vehicles), and MC (motorcycles).

<sup>b</sup> Because of a lack of data, all motorcycles over 12 years old are considered to have the same emissions and travel characteristics, and therefore are presented in aggregate.

**Table A-101: VMT Distribution by Vehicle Age and Vehicle/Fuel Type, <sup>a</sup> 2014**

| Vehicle Age  | LDGV           | LDGT           | HDGV           | LDDV           | LDDT           | HDDV           | MC             |
|--------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| 0            | 9.18%          | 10.20%         | 11.71%         | 17.18%         | 10.81%         | 8.56%          | 24.81%         |
| 1            | 8.61%          | 9.46%          | 10.55%         | 16.11%         | 10.01%         | 7.64%          | 12.70%         |
| 2            | 8.14%          | 8.84%          | 9.70%          | 15.25%         | 9.35%          | 7.27%          | 9.04%          |
| 3            | 4.92%          | 5.79%          | 5.12%          | 9.21%          | 6.27%          | 5.32%          | 5.99%          |
| 4            | 5.39%          | 4.98%          | 2.96%          | 8.67%          | 3.57%          | 3.43%          | 4.68%          |
| 5            | 4.76%          | 3.63%          | 2.57%          | 5.57%          | 3.12%          | 4.34%          | 4.31%          |
| 6            | 5.85%          | 6.00%          | 5.20%          | 0.49%          | 7.40%          | 4.63%          | 6.93%          |
| 7            | 6.17%          | 6.05%          | 4.02%          | 0.33%          | 6.38%          | 11.95%         | 5.66%          |
| 8            | 5.52%          | 5.85%          | 5.79%          | 6.04%          | 7.78%          | 8.69%          | 4.96%          |
| 9            | 5.29%          | 5.77%          | 3.89%          | 4.01%          | 6.40%          | 7.54%          | 4.06%          |
| 10           | 4.73%          | 5.48%          | 4.76%          | 2.32%          | 5.43%          | 4.36%          | 3.24%          |
| 11           | 4.70%          | 4.74%          | 3.70%          | 2.91%          | 4.72%          | 3.73%          | 2.62%          |
| 12           | 4.45%          | 4.32%          | 3.41%          | 3.03%          | 3.86%          | 2.58%          | 2.19%          |
| 13           | 3.96%          | 3.62%          | 2.42%          | 1.86%          | 4.17%          | 3.12%          | 1.76%          |
| 14           | 3.75%          | 3.21%          | 5.04%          | 1.52%          | 2.11%          | 4.48%          | 1.34%          |
| 15           | 2.82%          | 2.60%          | 4.16%          | 0.84%          | 2.87%          | 3.05%          | 0.97%          |
| 16           | 2.18%          | 1.95%          | 1.46%          | 0.76%          | 1.02%          | 1.83%          | 0.80%          |
| 17           | 1.89%          | 1.61%          | 2.43%          | 0.27%          | 1.22%          | 1.32%          | 0.73%          |
| 18           | 1.47%          | 1.12%          | 1.37%          | 0.29%          | 0.89%          | 1.25%          | 0.62%          |
| 19           | 1.41%          | 1.04%          | 1.90%          | 0.21%          | 0.63%          | 1.30%          | 0.45%          |
| 20           | 1.04%          | 0.85%          | 1.34%          | 0.02%          | 0.35%          | 0.86%          | 0.50%          |
| 21           | 0.84%          | 0.59%          | 1.07%          | 0.09%          | 0.36%          | 0.60%          | 0.40%          |
| 22           | 0.67%          | 0.43%          | 0.71%          | 0.12%          | 0.32%          | 0.38%          | 0.32%          |
| 23           | 0.54%          | 0.35%          | 0.56%          | 0.24%          | 0.18%          | 0.34%          | 0.24%          |
| 24           | 0.44%          | 0.31%          | 0.73%          | 0.07%          | 0.14%          | 0.36%          | 0.18%          |
| 25           | 0.35%          | 0.31%          | 0.82%          | 0.05%          | 0.13%          | 0.30%          | 0.13%          |
| 26           | 0.26%          | 0.27%          | 0.61%          | 0.01%          | 0.10%          | 0.24%          | 0.10%          |
| 27           | 0.21%          | 0.20%          | 0.53%          | 0.43%          | 0.04%          | 0.19%          | 0.09%          |
| 28           | 0.17%          | 0.20%          | 0.50%          | 0.24%          | 0.15%          | 0.12%          | 0.07%          |
| 29           | 0.13%          | 0.13%          | 0.38%          | 0.72%          | 0.11%          | 0.09%          | 0.06%          |
| 30           | 0.15%          | 0.11%          | 0.58%          | 1.12%          | 0.11%          | 0.11%          | 0.05%          |
| <b>Total</b> | <b>100.00%</b> |

Model that affects this distribution.

<sup>a</sup> The following abbreviations correspond to vehicle types: LDGV (light-duty gasoline vehicles), LDGT (light-duty gasoline trucks), HDGV (heavy-duty gasoline vehicles), LDDV (light-duty diesel vehicles), LDDT (light-duty diesel trucks), HDDV (heavy-duty diesel vehicles), and MC (motorcycles).

Note: Estimated by weighting data in by data in Table A-100. This year's Inventory includes updated vehicle population data based on the MOVES 2014a

**Table A-102: Fuel Consumption for Off-Road Sources by Fuel Type (million gallons)**

| Vehicle Type/Year                                | 1990          | 1995          | 2000          | 2001          | 2002          | 2003          | 2004          | 2005          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|--|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Aircraft<sup>a</sup></b>                      | <b>19,560</b> | <b>18,320</b> | <b>20,304</b> | <b>19,745</b> | <b>19,284</b> | <b>18,640</b> | <b>19,407</b> | <b>19,714</b> | <b>18,973</b> | <b>18,670</b> | <b>17,984</b> | <b>16,030</b> | <b>15,762</b> | <b>15,262</b> | <b>14,914</b> | <b>15,274</b> | <b>15,418</b> |
| Aviation Gasoline                                | 374           | 329           | 302           | 291           | 281           | 251           | 260           | 294           | 278           | 263           | 235           | 221           | 225           | 225           | 209           | 186           | 181           |
| Jet Fuel   | 19,186        | 17,991        | 20,002        | 19,454        | 19,004        | 18,389        | 19,147        | 19,420        | 18,695        | 18,407        | 17,749        | 15,809        | 15,537        | 15,036        | 14,705        | 15,088        | 15,237        |
| Commercial Aviation <sup>b</sup>                 | 11,569        | 12,136        | 14,672        | 13,121        | 12,774        | 12,943        | 13,147        | 13,976        | 14,426        | 14,708        | 13,400        | 12,588        | 11,931        | 12,067        | 11,932        | 12,031        | 12,131        |
| <b>Ships and Other Boats</b>                     | <b>4,507</b>  | <b>5,789</b>  | <b>6,407</b>  | <b>4,393</b>  | <b>4,810</b>  | <b>4,066</b>  | <b>4,277</b>  | <b>4,858</b>  | <b>5,120</b>  | <b>5,723</b>  | <b>4,860</b>  | <b>4,289</b>  | <b>5,740</b>  | <b>5,915</b>  | <b>5,340</b>  | <b>5,293</b>  | <b>4,329</b>  |
| Diesel   | 1,043         | 1,546         | 1,750         | 1,630         | 1,592         | 1,711         | 1,347         | 1,470         | 1,409         | 1,489         | 1,470         | 1,480         | 1,446         | 1,727         | 1,475         | 1,499         | 1,370         |
| Gasoline   | 1,403         | 1,597         | 1,629         | 1,632         | 1,630         | 1,625         | 1,616         | 1,607         | 1,597         | 1,587         | 1,577         | 1,568         | 1,556         | 1,545         | 1,535         | 1,528         | 1,522         |
| Residual   | 2,061         | 2,646         | 3,028         | 1,131         | 1,588         | 730           | 1,313         | 1,781         | 2,115         | 2,647         | 1,812         | 1,241         | 2,738         | 2,643         | 2,330         | 2,265         | 1,437         |
| <b>Construction/Mining Equipment<sup>c</sup></b> | <b>4,160</b>  | <b>4,835</b>  | <b>5,523</b>  | <b>5,984</b>  | <b>6,156</b>  | <b>6,339</b>  | <b>6,522</b>  | <b>6,617</b>  | <b>6,755</b>  | <b>6,785</b>  | <b>6,939</b>  | <b>7,066</b>  | <b>7,312</b>  | <b>7,418</b>  | <b>7,586</b>  | <b>8,187</b>  | <b>7,949</b>  |
| Diesel   | 3,674         | 4,387         | 5,181         | 5,329         | 5,477         | 5,625         | 5,774         | 5,922         | 6,069         | 6,216         | 6,363         | 6,511         | 6,658         | 6,806         | 6,954         | 7,102         | 7,250         |
| Gasoline   | 486           | 448           | 342           | 654           | 678           | 714           | 749           | 695           | 686           | 569           | 575           | 556           | 655           | 612           | 632           | 1,085         | 698           |
| <b>Agricultural Equipment<sup>d</sup></b>        | <b>3,134</b>  | <b>3,698</b>  | <b>3,929</b>  | <b>4,163</b>  | <b>4,277</b>  | <b>4,383</b>  | <b>4,708</b>  | <b>4,776</b>  | <b>5,011</b>  | <b>4,926</b>  | <b>4,582</b>  | <b>4,708</b>  | <b>4,807</b>  | <b>4,998</b>  | <b>5,157</b>  | <b>5,021</b>  | <b>5,094</b>  |
| Diesel   | 2,321         | 2,772         | 3,277         | 3,361         | 3,445         | 3,530         | 3,614         | 3,699         | 3,782         | 3,865         | 3,948         | 4,032         | 4,115         | 4,199         | 4,282         | 4,366         | 4,450         |
| Gasoline   | 813           | 927           | 652           | 802           | 832           | 853           | 1,094         | 1,078         | 1,229         | 1,061         | 634           | 676           | 692           | 799           | 875           | 655           | 644           |
| <b>Rail</b>                                      | <b>3,461</b>  | <b>3,864</b>  | <b>4,106</b>  | <b>4,119</b>  | <b>4,089</b>  | <b>4,176</b>  | <b>4,407</b>  | <b>4,446</b>  | <b>4,665</b>  | <b>4,539</b>  | <b>4,216</b>  | <b>3,535</b>  | <b>3,807</b>  | <b>3,999</b>  | <b>3,921</b>  | <b>4,025</b>  | <b>4,180</b>  |
| Diesel   | 3,461         | 3,864         | 4,106         | 4,119         | 4,089         | 4,176         | 4,407         | 4,446         | 4,665         | 4,539         | 4,216         | 3,535         | 3,807         | 3,999         | 3,921         | 4,025         | 4,180         |
| <b>Other<sup>e</sup></b>                         | <b>5,916</b>  | <b>6,525</b>  | <b>6,798</b>  | <b>7,630</b>  | <b>7,813</b>  | <b>8,022</b>  | <b>8,236</b>  | <b>8,255</b>  | <b>8,370</b>  | <b>8,229</b>  | <b>8,360</b>  | <b>8,455</b>  | <b>8,804</b>  | <b>8,768</b>  | <b>8,703</b>  | <b>8,800</b>  | <b>8,952</b>  |
| Diesel   | 1,423         | 1,720         | 2,050         | 2,114         | 2,181         | 2,247         | 2,313         | 2,380         | 2,446         | 2,512         | 2,579         | 2,645         | 2,711         | 2,778         | 2,844         | 2,910         | 2,977         |
| Gasoline   | 4,493         | 4,805         | 4,748         | 5,515         | 5,632         | 5,775         | 5,922         | 5,875         | 5,924         | 5,717         | 5,782         | 5,810         | 6,093         | 5,990         | 5,859         | 5,890         | 5,975         |
| <b>Total</b>                                     | <b>40,738</b> | <b>43,031</b> | <b>47,067</b> | <b>46,032</b> | <b>46,429</b> | <b>45,627</b> | <b>47,557</b> | <b>48,666</b> | <b>48,894</b> | <b>48,872</b> | <b>46,941</b> | <b>44,083</b> | <b>46,233</b> | <b>46,359</b> | <b>45,622</b> | <b>46,600</b> | <b>45,922</b> |

<sup>a</sup> For aircraft, this is aviation gasoline. For all other categories, this is motor gasoline.

<sup>b</sup> Commercial aviation, as modeled in FAA's AEDT, consists of passenger aircraft, cargo, and other chartered flights.

<sup>c</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.

<sup>d</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.

<sup>e</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

Note: In 2015, EPA incorporated the NONROAD2008 model into MOVES2014a. This year's inventory uses the NONROAD component of MOVES2014a for years 1999 through 2014. This update resulted in small changes (<2%) to the 1999-2013 time series for NONROAD fuel consumption due to differences in the gasoline and diesel default fuel densities used within the model iterations.

Sources: AAR (2008 through 2014), APTA (2007 through 2013), BEA (1991 through 2014), Benson (2002 through 2004), DHS (2008), DOC (1991 through 2014), DESC (2014), DOE (1993 through 2013), DOT (1991 through 2014), EIA (2002), EIA (2007b), EIA (2016), EIA (2007 through 2014), EIA (1991 through 2014), EPA (2015d), FAA (2016), Gaffney (2007), and Whorton (2006 through 2013).

**Table A-103: Control Technology Assignments for Gasoline Passenger Cars (Percent of VMT)**

| Model Years | Non-catalyst | Oxidation | EPA Tier 0 | EPA Tier 1 | LEV | EPA Tier 2 |
|-------------|--------------|-----------|------------|------------|-----|------------|
| 1973-1974   | 100%         | -         | -          | -          | -   | -          |
| 1975        | 20%          | 80%       | -          | -          | -   | -          |
| 1976-1977   | 15%          | 85%       | -          | -          | -   | -          |
| 1978-1979   | 10%          | 90%       | -          | -          | -   | -          |
| 1980        | 5%           | 88%       | 7%         | -          | -   | -          |
| 1981        | -            | 15%       | 85%        | -          | -   | -          |
| 1982        | -            | 14%       | 86%        | -          | -   | -          |
| 1983        | -            | 12%       | 88%        | -          | -   | -          |
| 1984-1993   | -            | -         | 100%       | -          | -   | -          |
| 1994        | -            | -         | 60%        | 40%        | -   | -          |
| 1995        | -            | -         | 20%        | 80%        | -   | -          |
| 1996        | -            | -         | 1%         | 97%        | 2%  | -          |
| 1997        | -            | -         | 0.5%       | 96.5%      | 3%  | -          |
| 1998        | -            | -         | <1%        | 87%        | 13% | -          |
| 1999        | -            | -         | <1%        | 67%        | 33% | -          |
| 2000        | -            | -         | -          | 44%        | 56% | -          |
| 2001        | -            | -         | -          | 3%         | 97% | -          |
| 2002        | -            | -         | -          | 1%         | 99% | -          |
| 2003        | -            | -         | -          | <1%        | 87% | 13%        |
| 2004        | -            | -         | -          | <1%        | 41% | 59%        |
| 2005        | -            | -         | -          | -          | 38% | 62%        |
| 2006        | -            | -         | -          | -          | 18% | 82%        |
| 2007        | -            | -         | -          | -          | 4%  | 96%        |
| 2008        | -            | -         | -          | -          | 2%  | 98%        |
| 2009-14     | -            | -         | -          | -          | -   | 100%       |

Sources: EPA (1998), EPA (2007a), and EPA (2007b).

- Not applicable.

Note: Detailed descriptions of emissions control technologies are provided in the following section of this Annex.

**Table A-104: Control Technology Assignments for Gasoline Light-Duty Trucks (Percent of VMT)<sup>a</sup>**

| Model Years | Non-catalyst | Oxidation | EPA Tier 0 | EPA Tier 1 | LEV <sup>b</sup> | EPA Tier 2 |
|-------------|--------------|-----------|------------|------------|------------------|------------|
| 1973-1974   | 100%         | -         | -          | -          | -                | -          |
| 1975        | 30%          | 70%       | -          | -          | -                | -          |
| 1976        | 20%          | 80%       | -          | -          | -                | -          |
| 1977-1978   | 25%          | 75%       | -          | -          | -                | -          |
| 1979-1980   | 20%          | 80%       | -          | -          | -                | -          |
| 1981        | -            | 95%       | 5%         | -          | -                | -          |
| 1982        | -            | 90%       | 10%        | -          | -                | -          |
| 1983        | -            | 80%       | 20%        | -          | -                | -          |
| 1984        | -            | 70%       | 30%        | -          | -                | -          |
| 1985        | -            | 60%       | 40%        | -          | -                | -          |
| 1986        | -            | 50%       | 50%        | -          | -                | -          |
| 1987-1993   | -            | 5%        | 95%        | -          | -                | -          |
| 1994        | -            | -         | 60%        | 40%        | -                | -          |
| 1995        | -            | -         | 20%        | 80%        | -                | -          |
| 1996        | -            | -         | -          | 100%       | -                | -          |
| 1997        | -            | -         | -          | 100%       | -                | -          |
| 1998        | -            | -         | -          | 80%        | 20%              | -          |
| 1999        | -            | -         | -          | 57%        | 43%              | -          |
| 2000        | -            | -         | -          | 65%        | 35%              | -          |
| 2001        | -            | -         | -          | 1%         | 99%              | -          |
| 2002        | -            | -         | -          | 10%        | 90%              | -          |
| 2003        | -            | -         | -          | <1%        | 53%              | 47%        |
| 2004        | -            | -         | -          | -          | 72%              | 28%        |
| 2005        | -            | -         | -          | -          | 38%              | 62%        |
| 2006        | -            | -         | -          | -          | 25%              | 75%        |
| 2007        | -            | -         | -          | -          | 14%              | 86%        |
| 2008-2014   | -            | -         | -          | -          | -                | 100%       |

Sources: EPA (1998), EPA (2007a), and EPA (2007b).

- Not applicable.

<sup>a</sup> Detailed descriptions of emissions control technologies are provided in the following section of this Annex.

<sup>b</sup> The proportion of LEVs as a whole has decreased since 2001, as carmakers have been able to achieve greater emission reductions with certain types of LEVs, such as ULEVs. Because ULEVs emit about half the emissions of LEVs, a carmaker can reduce the total number of LEVs they need to build to meet a specified emission average for all of their vehicles in a given model year.

**Table A-105: Control Technology Assignments for Gasoline Heavy-Duty Vehicles (Percent of VMT)<sup>a</sup>**

| Model Years | Uncontrolled | Non-catalyst | Oxidation | EPA Tier 0 | EPA Tier 1 | LEV <sup>b</sup> | EPA Tier 2 |
|-------------|--------------|--------------|-----------|------------|------------|------------------|------------|
| ≤1981       | 100%         | -            | -         | -          | -          | -                | -          |
| 1982-1984   | 95%          | -            | 5%        | -          | -          | -                | -          |
| 1985-1986   | -            | 95%          | 5%        | -          | -          | -                | -          |
| 1987        | -            | 70%          | 15%       | 15%        | -          | -                | -          |
| 1988-1989   | -            | 60%          | 25%       | 15%        | -          | -                | -          |
| 1990-1995   | -            | 45%          | 30%       | 25%        | -          | -                | -          |
| 1996        | -            | -            | 25%       | 10%        | 65%        | -                | -          |
| 1997        | -            | -            | 10%       | 5%         | 85%        | -                | -          |
| 1998        | -            | -            | -         | -          | 96%        | 4%               | -          |
| 1999        | -            | -            | -         | -          | 78%        | 22%              | -          |
| 2000        | -            | -            | -         | -          | 54%        | 46%              | -          |
| 2001        | -            | -            | -         | -          | 64%        | 36%              | -          |
| 2002        | -            | -            | -         | -          | 69%        | 31%              | -          |
| 2003        | -            | -            | -         | -          | 65%        | 30%              | 5%         |
| 2004        | -            | -            | -         | -          | 5%         | 37%              | 59%        |
| 2005        | -            | -            | -         | -          | -          | 23%              | 77%        |
| 2006        | -            | -            | -         | -          | -          | 20%              | 80%        |
| 2007        | -            | -            | -         | -          | -          | 10%              | 90%        |
| 2008-2014   | -            | -            | -         | -          | -          | 0%               | 100%       |

Sources: EPA (1998), EPA (2007a), and EPA (2007b).

- Not applicable.

<sup>a</sup> Detailed descriptions of emissions control technologies are provided in the following section of this Annex.

<sup>b</sup> The proportion of LEVs as a whole has decreased since 2000, as carmakers have been able to achieve greater emission reductions with certain types of LEVs, such as ULEVs. Because ULEVs emit about half the emissions of LEVs, a manufacturer can reduce the total number of LEVs they need to build to meet a specified emission average for all of their vehicles in a given model year.

**Table A-106: Control Technology Assignments for Diesel On-Road Vehicles and Motorcycles**

| Vehicle Type/Control Technology                       | Model Years |
|---|-------------|
| <b>Diesel Passenger Cars and Light-Duty Trucks</b>    |             |
| Uncontrolled  | 1960-1982   |
| Moderate control                                      | 1983-1995   |
| Advanced control                                      | 1996-2014   |
| <b>Diesel Medium- and Heavy-Duty Trucks and Buses</b> |             |
| Uncontrolled  | 1960-1990   |
| Moderate control                                      | 1991-2003   |
| Advanced control                                      | 2004-2006   |
| Aftertreatment  | 2007-2014   |
| <b>Motorcycles</b>                                    |             |
| Uncontrolled  | 1960-1995   |
| Non-catalyst controls                                 | 1996-2014   |

Note: Detailed descriptions of emissions control technologies are provided in the following section of this Annex.

Source: EPA (1998) and Browning (2005).

**Table A-107: Emission Factors for CH<sub>4</sub> and N<sub>2</sub>O for On-Road Vehicles**

| Vehicle Type/Control Technology | N <sub>2</sub> O<br>(g/mi) | CH <sub>4</sub><br>(g/mi) |
|---------------------------------|----------------------------|---------------------------|
| <b>Gasoline Passenger Cars</b>  |                            |                           |
| EPA Tier 2                      | 0.0036                     | 0.0173                    |
| Low Emission Vehicles           | 0.0150                     | 0.0105                    |
| EPA Tier 1 <sup>a</sup>         | 0.0429                     | 0.0271                    |
| EPA Tier 0 <sup>a</sup>         | 0.0647                     | 0.0704                    |
| Oxidation Catalyst              | 0.0504                     | 0.1355                    |
| Non-Catalyst Control            | 0.0197                     | 0.1696                    |

|   |        |        |
|---|--------|--------|
| Uncontrolled  | 0.0197 | 0.1780 |
| <b>Gasoline Light-Duty Trucks</b>                     |        |        |
| EPA Tier 2  | 0.0066 | 0.0163 |
| Low Emission Vehicles                                 | 0.0157 | 0.0148 |
| EPA Tier 1 <sup>a</sup>                               | 0.0871 | 0.0452 |
| EPA Tier 0 <sup>a</sup>                               | 0.1056 | 0.0776 |
| Oxidation Catalyst                                    | 0.0639 | 0.1516 |
| Non-Catalyst Control                                  | 0.0218 | 0.1908 |
| Uncontrolled  | 0.0220 | 0.2024 |
| <b>Gasoline Heavy-Duty Vehicles</b>                   |        |        |
| EPA Tier 2  | 0.0134 | 0.0333 |
| Low Emission Vehicles                                 | 0.0320 | 0.0303 |
| EPA Tier 1 <sup>a</sup>                               | 0.1750 | 0.0655 |
| EPA Tier 0 <sup>a</sup>                               | 0.2135 | 0.2630 |
| Oxidation Catalyst                                    | 0.1317 | 0.2356 |
| Non-Catalyst Control                                  | 0.0473 | 0.4181 |
| Uncontrolled  | 0.0497 | 0.4604 |
| <b>Diesel Passenger Cars</b>                          |        |        |
| Advanced  | 0.0010 | 0.0005 |
| Moderate  | 0.0010 | 0.0005 |
| Uncontrolled  | 0.0012 | 0.0006 |
| <b>Diesel Light-Duty Trucks</b>                       |        |        |
| Advanced  | 0.0015 | 0.0010 |
| Moderate  | 0.0014 | 0.0009 |
| Uncontrolled  | 0.0017 | 0.0011 |
| <b>Diesel Medium- and Heavy-Duty Trucks and Buses</b> |        |        |
| Aftertreatment  | 0.0048 | 0.0051 |
| Advanced  | 0.0048 | 0.0051 |
| Moderate  | 0.0048 | 0.0051 |
| Uncontrolled  | 0.0048 | 0.0051 |
| <b>Motorcycles</b>                                    |        |        |
| Non-Catalyst Control                                  | 0.0069 | 0.0672 |
| Uncontrolled  | 0.0087 | 0.0899 |

Source: ICF (2006b and 2004).

<sup>a</sup> The categories "EPA Tier 0" and "EPA Tier 1" were substituted for the early three-way catalyst and advanced three-way catalyst categories, respectively, as defined in the 2006 IPCC Guidelines. Detailed descriptions of emissions control technologies are provided at the end of this Annex.

**Table A-108: Emission Factors for CH<sub>4</sub> and N<sub>2</sub>O for Alternative Fuel Vehicles (g/mi)**

|                                      | N <sub>2</sub> O | CH <sub>4</sub> |
|--------------------------------------|------------------|-----------------|
| <b>Light Duty Vehicles</b>           |                  |                 |
| Methanol                             | 0.067            | 0.018           |
| CNG                                  | 0.050            | 0.737           |
| LPG                                  | 0.067            | 0.037           |
| Ethanol                              | 0.067            | 0.055           |
| Biodiesel (BD20)                     | 0.001            | 0.0005          |
| <b>Medium- and Heavy-Duty Trucks</b> |                  |                 |
| Methanol                             | 0.175            | 0.066           |
| CNG                                  | 0.175            | 1.966           |
| LNG                                  | 0.175            | 1.966           |
| LPG                                  | 0.175            | 0.066           |
| Ethanol                              | 0.175            | 0.197           |
| Biodiesel (BD20)                     | 0.005            | 0.005           |
| <b>Buses</b>                         |                  |                 |
| Methanol                             | 0.175            | 0.066           |
| CNG                                  | 0.175            | 1.966           |
| Ethanol                              | 0.175            | 0.197           |
| Biodiesel (BD20)                     | 0.005            | 0.005           |

Source: Developed by ICF (2006a) using ANL (2006) and Lipman and Delucchi (2002).

**Table A-109: Emission Factors for CH<sub>4</sub> and N<sub>2</sub>O Emissions from Non-Road Mobile Combustion (g/kg fuel)**

| Vehicle Type/Fuel Type                           | N <sub>2</sub> O | CH <sub>4</sub> |
|--|------------------|-----------------|
| <b>Ships and Boats</b>                           |                  |                 |
| Residual   | 0.16             | 0.03            |
| Gasoline   | 0.08             | 0.23            |
| Diesel   | 0.14             | 0.02            |
| <b>Rail</b>                                      |                  |                 |
| Diesel   | 0.08             | 0.25            |
| <b>Agricultural Equipment<sup>a</sup></b>        |                  |                 |
| Gasoline   | 0.08             | 0.45            |
| Diesel   | 0.08             | 0.45            |
| <b>Construction/Mining Equipment<sup>b</sup></b> |                  |                 |
| Gasoline   | 0.08             | 0.18            |
| Diesel   | 0.08             | 0.18            |
| <b>Other Non-Road</b>                            |                  |                 |
| All "Other" Categories <sup>c</sup>              | 0.08             | 0.18            |
| <b>Aircraft</b>                                  |                  |                 |
| Jet Fuel <sup>d</sup>                            | 0.10             | 0.00            |
| Aviation Gasoline                                | 0.04             | 2.64            |

Source: IPCC (2006) and ICF (2009).

<sup>a</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.

<sup>b</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.

<sup>c</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

<sup>d</sup> Emissions of CH<sub>4</sub> from jet fuels have been zeroed out across the time series. Recent research indicates that modern aircraft jet engines are typically net consumers of methane (Santoni et al. 2011). Methane is emitted at low power and idle operation, but at higher power modes aircraft engines consumer methane. Over the range of engine operating modes, aircraft engines are net consumers of methane on average. Based on this data, methane emissions factors for jet aircraft were changed to zero in this year's Inventory to reflect the latest emissions testing data.

**Table A-110: NO<sub>x</sub> Emissions from Mobile Combustion (kt)**

| Fuel Type/Vehicle Type                      | 1990          | 1995          | 2000          | 2001         | 2002          | 2003          | 2004          | 2005          | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|---|---------------|---------------|---------------|--------------|---------------|---------------|---------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Gasoline On-Road</b>                     | <b>5,746</b>  | <b>4,560</b>  | <b>3,812</b>  | <b>3,715</b> | <b>4,940</b>  | <b>4,621</b>  | <b>4,303</b>  | <b>3,984</b>  | <b>3,819</b> | <b>3,654</b> | <b>3,317</b> | <b>2,966</b> | <b>2,724</b> | <b>2,805</b> | <b>2,585</b> | <b>2,365</b> | <b>2,145</b> |
| Passenger Cars                              | 3,847         | 2,752         | 2,084         | 2,027        | 2,695         | 2,521         | 2,347         | 2,174         | 2,083        | 1,993        | 1,810        | 1,618        | 1,486        | 1,530        | 1,410        | 1,290        | 1,170        |
| Light-Duty Trucks                           | 1,364         | 1,325         | 1,303         | 1,285        | 1,708         | 1,598         | 1,488         | 1,378         | 1,321        | 1,264        | 1,147        | 1,026        | 942          | 970          | 894          | 818          | 742          |
| Medium- and Heavy-Duty Trucks and Buses     | 515           | 469           | 411           | 390          | 518           | 485           | 452           | 418           | 401          | 383          | 348          | 311          | 286          | 294          | 271          | 248          | 225          |
| Motorcycles                                 | 20            | 14            | 13            | 14           | 18            | 17            | 16            | 15            | 14           | 13           | 12           | 11           | 10           | 10           | 10           | 9            | 8            |
| <b>Diesel On-Road</b>                       | <b>2,956</b>  | <b>3,493</b>  | <b>3,803</b>  | <b>3,338</b> | <b>4,438</b>  | <b>4,152</b>  | <b>3,866</b>  | <b>3,580</b>  | <b>3,431</b> | <b>3,283</b> | <b>2,980</b> | <b>2,665</b> | <b>2,448</b> | <b>2,520</b> | <b>2,323</b> | <b>2,125</b> | <b>1,927</b> |
| Passenger Cars                              | 39            | 19            | 7             | 6            | 8             | 7             | 7             | 6             | 6            | 6            | 5            | 5            | 4            | 4            | 4            | 4            | 3            |
| Light-Duty Trucks                           | 20            | 12            | 6             | 5            | 7             | 7             | 6             | 6             | 6            | 5            | 5            | 4            | 4            | 4            | 4            | 3            | 3            |
| Medium- and Heavy-Duty Trucks and Buses     | 2,897         | 3,462         | 3,791         | 3,326        | 4,423         | 4,138         | 3,853         | 3,568         | 3,420        | 3,272        | 2,970        | 2,656        | 2,439        | 2,512        | 2,315        | 2,118        | 1,921        |
| <b>Alternative Fuel On-Road<sup>a</sup></b> | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>    | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    |
| <b>Non-Road</b>                             | <b>2,160</b>  | <b>2,483</b>  | <b>2,584</b>  | <b>2,643</b> | <b>3,107</b>  | <b>2,981</b>  | <b>2,856</b>  | <b>2,731</b>  | <b>2,490</b> | <b>2,249</b> | <b>2,226</b> | <b>2,166</b> | <b>2,118</b> | <b>1,968</b> | <b>1,881</b> | <b>1,793</b> | <b>1,705</b> |
| Ships and Boats                             | 402           | 488           | 506           | 544          | 643           | 617           | 591           | 565           | 515          | 465          | 460          | 448          | 438          | 407          | 389          | 371          | 353          |
| Rail  | 338           | 433           | 451           | 485          | 574           | 550           | 527           | 504           | 460          | 415          | 411          | 400          | 391          | 363          | 347          | 331          | 315          |
| Aircraft <sup>b</sup>                       | 25            | 31            | 40            | 39           | 46            | 45            | 43            | 41            | 37           | 34           | 33           | 32           | 32           | 29           | 28           | 27           | 25           |
| Agricultural Equipment <sup>c</sup>         | 437           | 478           | 484           | 480          | 562           | 539           | 516           | 494           | 450          | 407          | 402          | 392          | 383          | 356          | 340          | 324          | 308          |
| Construction/Mining Equipment <sup>d</sup>  | 641           | 697           | 697           | 690          | 807           | 774           | 742           | 709           | 647          | 584          | 578          | 563          | 550          | 511          | 488          | 466          | 443          |
| Other <sup>e</sup>                          | 318           | 357           | 407           | 406          | 476           | 456           | 437           | 418           | 381          | 344          | 341          | 332          | 324          | 301          | 288          | 274          | 261          |
| <b>Total</b>                                | <b>10,862</b> | <b>10,536</b> | <b>10,199</b> | <b>9,696</b> | <b>12,485</b> | <b>11,755</b> | <b>11,025</b> | <b>10,295</b> | <b>9,740</b> | <b>9,186</b> | <b>8,523</b> | <b>7,797</b> | <b>7,290</b> | <b>7,294</b> | <b>6,788</b> | <b>6,283</b> | <b>5,777</b> |

IE - Included Elsewhere

<sup>a</sup> NO<sub>x</sub> emissions from alternative fuel on-road vehicles are included under gasoline and diesel on-road.<sup>b</sup> Aircraft estimates include only emissions related to LTO cycles, and therefore do not include cruise altitude emissions.<sup>c</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.<sup>d</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.<sup>e</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

Note: The source of this data is the National Emissions Inventory. Updates to estimates from MOVES2014a is a change that affects the emissions time series.

Note: Totals may not sum due to independent rounding.

**Table A-111: CO Emissions from Mobile Combustion (kt)**

| Fuel Type/Vehicle Type                      | 1990           | 1995          | 2000          | 2001          | 2002          | 2003          | 2004          | 2005          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|---|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Gasoline On-Road</b>                     | <b>98,328</b>  | <b>74,673</b> | <b>60,657</b> | <b>56,716</b> | <b>46,115</b> | <b>43,498</b> | <b>40,882</b> | <b>38,265</b> | <b>35,781</b> | <b>33,298</b> | <b>29,626</b> | <b>24,515</b> | <b>25,235</b> | <b>24,442</b> | <b>22,925</b> | <b>21,408</b> | <b>19,890</b> |
| Passenger Cars                              | 60,757         | 42,065        | 32,867        | 31,600        | 25,693        | 24,235        | 22,777        | 21,319        | 19,936        | 18,552        | 16,506        | 13,659        | 14,060        | 13,618        | 12,773        | 11,927        | 11,082        |
| Light-Duty Trucks                           | 29,237         | 27,048        | 24,532        | 22,574        | 18,355        | 17,313        | 16,272        | 15,230        | 14,242        | 13,253        | 11,792        | 9,758         | 10,044        | 9,729         | 9,125         | 8,521         | 7,917         |
| Medium- and Heavy-Duty Trucks and Buses     | 8,093          | 5,404         | 3,104         | 2,411         | 1,960         | 1,849         | 1,738         | 1,627         | 1,521         | 1,416         | 1,259         | 1,042         | 1,073         | 1,039         | 975           | 910           | 846           |
| Motorcycles                                 | 240            | 155           | 154           | 131           | 107           | 101           | 95            | 89            | 83            | 77            | 69            | 57            | 58            | 57            | 53            | 50            | 46            |
| <b>Diesel On-Road</b>                       | <b>1,696</b>   | <b>1,424</b>  | <b>1,088</b>  | <b>869</b>    | <b>707</b>    | <b>667</b>    | <b>626</b>    | <b>586</b>    | <b>548</b>    | <b>510</b>    | <b>454</b>    | <b>376</b>    | <b>387</b>    | <b>375</b>    | <b>351</b>    | <b>328</b>    | <b>305</b>    |
| Passenger Cars                              | 35             | 18            | 7             | 6             | 5             | 4             | 4             | 4             | 4             | 3             | 3             | 3             | 3             | 3             | 2             | 2             | 2             |
| Light-Duty Trucks                           | 22             | 16            | 6             | 5             | 4             | 4             | 4             | 4             | 3             | 3             | 3             | 2             | 2             | 2             | 2             | 2             | 2             |
| Medium- and Heavy-Duty Trucks and Buses     | 1,639          | 1,391         | 1,075         | 858           | 698           | 658           | 618           | 579           | 541           | 504           | 448           | 371           | 382           | 370           | 347           | 324           | 301           |
| <b>Alternative Fuel On-Road<sup>a</sup></b> | <b>IE</b>      | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     | <b>IE</b>     |
| <b>Non-Road</b>                             | <b>19,337</b>  | <b>21,533</b> | <b>21,814</b> | <b>22,266</b> | <b>20,414</b> | <b>20,197</b> | <b>19,980</b> | <b>19,763</b> | <b>18,382</b> | <b>17,001</b> | <b>16,137</b> | <b>14,365</b> | <b>13,853</b> | <b>13,488</b> | <b>13,214</b> | <b>12,940</b> | <b>12,666</b> |
| Ships and Boats                             | 1,559          | 1,781         | 1,825         | 1,831         | 1,679         | 1,661         | 1,643         | 1,626         | 1,512         | 1,398         | 1,327         | 1,182         | 1,140         | 1,109         | 1,087         | 1,064         | 1,042         |
| Rail  | 85             | 93            | 90            | 90            | 82            | 81            | 81            | 80            | 74            | 69            | 65            | 58            | 56            | 54            | 53            | 52            | 51            |
| Aircraft <sup>b</sup>                       | 217            | 224           | 245           | 233           | 214           | 212           | 210           | 207           | 193           | 178           | 169           | 151           | 145           | 141           | 139           | 136           | 133           |
| Agricultural Equipment <sup>c</sup>         | 581            | 628           | 626           | 621           | 569           | 563           | 557           | 551           | 513           | 474           | 450           | 401           | 386           | 376           | 369           | 361           | 353           |
| Construction/Mining Equipment <sup>d</sup>  | 1,090          | 1,132         | 1,047         | 1,041         | 955           | 944           | 934           | 924           | 860           | 795           | 755           | 672           | 648           | 631           | 618           | 605           | 592           |
| Other <sup>e</sup>                          | 15,805         | 17,676        | 17,981        | 18,449        | 16,914        | 16,735        | 16,555        | 16,375        | 15,231        | 14,087        | 13,371        | 11,903        | 11,479        | 11,176        | 10,949        | 10,722        | 10,494        |
| <b>Total</b>                                | <b>119,360</b> | <b>97,630</b> | <b>83,559</b> | <b>79,851</b> | <b>67,235</b> | <b>64,362</b> | <b>61,488</b> | <b>58,615</b> | <b>54,712</b> | <b>50,809</b> | <b>46,217</b> | <b>39,256</b> | <b>39,475</b> | <b>38,305</b> | <b>36,491</b> | <b>34,676</b> | <b>32,861</b> |

IE - Included Elsewhere

<sup>a</sup> CO emissions from alternative fuel on-road vehicles are included under gasoline and diesel on-road.<sup>b</sup> Aircraft estimates include only emissions related to LTO cycles, and therefore do not include cruise altitude emissions.<sup>c</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.<sup>d</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.<sup>e</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

Notes: The source of this data is the National Emissions Inventory. Updates to estimates from MOVES2014a is a change that affects the emissions time series. Totals may not sum due to independent rounding.

**Table A-112: NMVOCs Emissions from Mobile Combustion (kt)**

| Fuel Type/Vehicle Type                      | 1990          | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|---|---------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Gasoline On-Road</b>                     | <b>8,110</b>  | <b>5,819</b> | <b>4,615</b> | <b>4,285</b> | <b>3,473</b> | <b>3,308</b> | <b>3,144</b> | <b>2,979</b> | <b>2,997</b> | <b>3,015</b> | <b>2,641</b> | <b>2,384</b> | <b>2,393</b> | <b>2,485</b> | <b>2,279</b> | <b>2,074</b> | <b>1,868</b> |
| Passenger Cars                              | 5,120         | 3,394        | 2,610        | 2,393        | 1,939        | 1,847        | 1,756        | 1,664        | 1,674        | 1,684        | 1,475        | 1,332        | 1,336        | 1,388        | 1,273        | 1,158        | 1,043        |
| Light-Duty Trucks                           | 2,374         | 2,019        | 1,750        | 1,664        | 1,348        | 1,285        | 1,221        | 1,157        | 1,164        | 1,171        | 1,025        | 926          | 929          | 965          | 885          | 805          | 725          |
| Medium- and Heavy-Duty Trucks and Buses     | 575           | 382          | 232          | 206          | 167          | 159          | 151          | 143          | 144          | 145          | 127          | 115          | 115          | 120          | 110          | 100          | 90           |
| Motorcycles                                 | 42            | 24           | 23           | 22           | 18           | 17           | 16           | 15           | 15           | 15           | 14           | 12           | 12           | 13           | 12           | 11           | 10           |
| <b>Diesel On-Road</b>                       | <b>406</b>    | <b>304</b>   | <b>216</b>   | <b>207</b>   | <b>168</b>   | <b>160</b>   | <b>152</b>   | <b>144</b>   | <b>145</b>   | <b>146</b>   | <b>128</b>   | <b>115</b>   | <b>116</b>   | <b>120</b>   | <b>110</b>   | <b>100</b>   | <b>90</b>    |
| Passenger Cars                              | 16            | 8            | 3            | 3            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 1            | 1            |
| Light-Duty Trucks                           | 14            | 9            | 4            | 4            | 3            | 3            | 3            | 3            | 3            | 3            | 2            | 2            | 2            | 2            | 2            | 2            | 2            |
| Medium- and Heavy-Duty Trucks and Buses     | 377           | 286          | 209          | 201          | 163          | 155          | 147          | 140          | 140          | 141          | 124          | 112          | 112          | 116          | 107          | 97           | 88           |
| <b>Alternative Fuel On-Road<sup>a</sup></b> | <b>IE</b>     | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    | <b>IE</b>    |
| <b>Non-Road</b>                             | <b>2,415</b>  | <b>2,622</b> | <b>2,398</b> | <b>2,379</b> | <b>2,800</b> | <b>2,733</b> | <b>2,667</b> | <b>2,600</b> | <b>2,491</b> | <b>2,383</b> | <b>2,310</b> | <b>2,150</b> | <b>2,082</b> | <b>1,957</b> | <b>1,863</b> | <b>1,768</b> | <b>1,674</b> |
| Ships and Boats                             | 608           | 739          | 744          | 730          | 859          | 839          | 818          | 798          | 764          | 731          | 709          | 660          | 639          | 600          | 572          | 543          | 514          |
| Rail  | 33            | 36           | 35           | 35           | 42           | 41           | 40           | 39           | 37           | 35           | 34           | 32           | 31           | 29           | 28           | 26           | 25           |
| Aircraft <sup>b</sup>                       | 28            | 28           | 24           | 19           | 23           | 22           | 22           | 21           | 20           | 19           | 19           | 17           | 17           | 16           | 15           | 14           | 14           |
| Agricultural Equipment <sup>c</sup>         | 85            | 86           | 76           | 72           | 85           | 83           | 81           | 79           | 76           | 73           | 70           | 65           | 63           | 60           | 57           | 54           | 51           |
| Construction/Mining Equipment <sup>d</sup>  | 149           | 152          | 130          | 125          | 147          | 144          | 140          | 137          | 131          | 125          | 121          | 113          | 109          | 103          | 98           | 93           | 88           |
| Other <sup>e</sup>                          | 1,512         | 1,580        | 1,390        | 1,397        | 1,644        | 1,605        | 1,566        | 1,527        | 1,463        | 1,399        | 1,356        | 1,263        | 1,223        | 1,149        | 1,094        | 1,038        | 983          |
| <b>Total</b>                                | <b>10,932</b> | <b>8,745</b> | <b>7,230</b> | <b>6,872</b> | <b>6,440</b> | <b>6,201</b> | <b>5,962</b> | <b>5,724</b> | <b>5,634</b> | <b>5,544</b> | <b>5,078</b> | <b>4,650</b> | <b>4,591</b> | <b>4,562</b> | <b>4,252</b> | <b>3,942</b> | <b>3,632</b> |

IE - Included Elsewhere

<sup>a</sup> NMVOC emissions from alternative fuel on-road vehicles are included under gasoline and diesel on-road.<sup>b</sup> Aircraft estimates include only emissions related to LTO cycles, and therefore do not include cruise altitude emissions.<sup>c</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.<sup>d</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.<sup>e</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

Notes: The source of this data is the National Emissions Inventory. Updates to estimates from MOVES2014a is a change that affects the emissions time series. Totals may not sum due to independent rounding.

## Definitions of Emission Control Technologies and Standards

The N<sub>2</sub>O and CH<sub>4</sub> emission factors used depend on the emission standards in place and the corresponding level of control technology for each vehicle type. Table A-103 through Table A-106 show the years in which these technologies or standards were in place and the penetration level for each vehicle type. These categories are defined below and were compiled from EPA (1993, 1994a, 1994b, 1998, 1999a) and IPCC/UNEP/OECD/IEA (1997).

### **Uncontrolled**

Vehicles manufactured prior to the implementation of pollution control technologies are designated as uncontrolled. Gasoline passenger cars and light-duty trucks (pre-1973), gasoline heavy-duty vehicles (pre-1984), diesel vehicles (pre-1983), and motorcycles (pre-1996) are assumed to have no control technologies in place.

### **Gasoline Emission Controls**

Below are the control technologies and emissions standards applicable to gasoline vehicles.

#### ***Non-catalyst***

These emission controls were common in gasoline passenger cars and light-duty gasoline trucks during model years (1973-1974) but phased out thereafter, in heavy-duty gasoline vehicles beginning in the mid-1980s, and in motorcycles beginning in 1996. This technology reduces hydrocarbon (HC) and carbon monoxide (CO) emissions through adjustments to ignition timing and air-fuel ratio, air injection into the exhaust manifold, and exhaust gas recirculation (EGR) valves, which also helps meet vehicle NO<sub>x</sub> standards.

#### ***Oxidation Catalyst***

This control technology designation represents the introduction of the catalytic converter, and was the most common technology in gasoline passenger cars and light-duty gasoline trucks made from 1975 to 1980 (cars) and 1975 to 1985 (trucks). This technology was also used in some heavy-duty gasoline vehicles between 1982 and 1997. The two-way catalytic converter oxidizes HC and CO, significantly reducing emissions over 80 percent beyond non-catalyst-system capacity. One reason unleaded gasoline was introduced in 1975 was due to the fact that oxidation catalysts cannot function properly with leaded gasoline.

#### ***EPA Tier 0***

This emission standard from the Clean Air Act was met through the implementation of early "three-way" catalysts, therefore this technology was used in gasoline passenger cars and light-duty gasoline trucks sold beginning in the early 1980s, and remained common until 1994. This more sophisticated emission control system improves the efficiency of the catalyst by converting CO and HC to CO<sub>2</sub> and H<sub>2</sub>O, reducing NO<sub>x</sub> to nitrogen and oxygen, and using an on-board diagnostic computer and oxygen sensor. In addition, this type of catalyst includes a fuel metering system (carburetor or fuel injection) with electronic "trim" (also known as a "closed-loop system"). New cars with three-way catalysts met the Clean Air Act's amended standards (enacted in 1977) of reducing HC to 0.41 g/mile by 1980, CO to 3.4 g/mile by 1981 and NO<sub>x</sub> to 1.0 g/mile by 1981.

#### ***EPA Tier 1***

This emission standard created through the 1990 amendments to the Clean Air Act limited passenger car NO<sub>x</sub> emissions to 0.4 g/mi, and HC emissions to 0.25 g/mi. These bounds respectively amounted to a 60 and 40 percent reduction from the EPA Tier 0 standard set in 1981. For light-duty trucks, this standard set emissions at 0.4 to 1.1 g/mi for NO<sub>x</sub>, and 0.25 to 0.39 g/mi for HCs, depending on the weight of the truck. Emission reductions were met through the use of more advanced emission control systems, and applied to light-duty gasoline vehicles beginning in 1994. These advanced emission control systems included advanced three-way catalysts, electronically controlled fuel injection and ignition timing, EGR, and air injection.

#### ***EPA Tier 2***

This emission standard was specified in the 1990 amendments to the Clean Air Act, limiting passenger car NO<sub>x</sub> emissions to 0.07 g/mi on average and aligning emissions standards for passenger cars and light-duty trucks. Manufacturers can meet this average emission level by producing vehicles in 11 emission "Bins," the three highest of which expire in 2006.

These new emission levels represent a 77 to 95 percent reduction in emissions from the EPA Tier 1 standard set in 1994. Emission reductions were met through the use of more advanced emission control systems and lower sulfur fuels and are applied to vehicles beginning in 2004. These advanced emission control systems include improved combustion, advanced three-way catalysts, electronically controlled fuel injection and ignition timing, EGR, and air injection.

### ***Low Emission Vehicles (LEV)***

This emission standard requires a much higher emission control level than the Tier 1 standard. Applied to light-duty gasoline passenger cars and trucks beginning in small numbers in the mid-1990s, LEV includes multi-port fuel injection with adaptive learning, an advanced computer diagnostics systems and advanced and close coupled catalysts with secondary air injection. LEVs as defined here include transitional low-emission vehicles (TLEVs), low emission vehicles, ultra-low emission vehicles (ULEVs) and super ultra-low emission vehicles (SULEVs). In this analysis, all categories of LEVs are treated the same due to the fact that there are very limited CH<sub>4</sub> or N<sub>2</sub>O emission factor data for LEVs to distinguish among the different types of vehicles. Zero emission vehicles (ZEVs) are incorporated into the alternative fuel and advanced technology vehicle assessments.

### **Diesel Emission Controls**

Below are the three levels of emissions control for diesel vehicles.

#### ***Moderate control***

Improved injection timing technology and combustion system design for light- and heavy-duty diesel vehicles (generally in place in model years 1983 to 1995) are considered moderate control technologies. These controls were implemented to meet emission standards for diesel trucks and buses adopted by the EPA in 1985 to be met in 1991 and 1994.

#### ***Advanced control***

EGR and modern electronic control of the fuel injection system are designated as advanced control technologies. These technologies provide diesel vehicles with the level of emission control necessary to comply with standards in place from 1996 through 2006.

#### ***Aftertreatment***

Use of diesel particulate filters (DPFs), oxidation catalysts and NO<sub>x</sub> absorbers or selective catalytic reduction (SCR) systems are designated as aftertreatment control. These technologies provide diesel vehicles with a level of emission control necessary to comply with standards in place from 2007 on.

## **Supplemental Information on GHG Emissions from Transportation and Other Mobile Sources**

This section of this Annex includes supplemental information on the contribution of transportation and other mobile sources to U.S. greenhouse gas emissions. In the main body of the Inventory report, emission estimates are generally presented by greenhouse gas, with separate discussions of the methodologies used to estimate CO<sub>2</sub>, N<sub>2</sub>O, CH<sub>4</sub>, and HFC emissions. Although the inventory is not required to provide detail beyond what is contained in the body of this report, the IPCC allows presentation of additional data and detail on emission sources. The purpose of this sub-annex, within the Annex that details the calculation methods and data used for non-CO<sub>2</sub> calculations, is to provide all transportation estimates presented throughout the report in one place.

This section of this Annex reports total greenhouse gas emissions from transportation and other (non-transportation) mobile sources in CO<sub>2</sub> equivalents, with information on the contribution by greenhouse gas and by mode, vehicle type, and fuel type. In order to calculate these figures, additional analyses were conducted to develop estimates of CO<sub>2</sub> from non-transportation mobile sources (e.g., agricultural equipment, construction/mining equipment, recreational vehicles), and to provide more detailed breakdowns of emissions by source.

### **Estimation of CO<sub>2</sub> from Non-Transportation Mobile Sources**

The estimates of N<sub>2</sub>O and CH<sub>4</sub> from fuel combustion presented in the Energy chapter of the Inventory include both transportation sources and other mobile sources. Other mobile sources include construction/mining equipment, agricultural equipment, vehicles used off-road, and other sources that have utility associated with their movement but do not have a primary purpose of transporting people or goods (e.g., snowmobiles, riding lawnmowers, etc.). Estimates of CO<sub>2</sub> from non-transportation mobile sources, based on EIA fuel consumption estimates, are included in the agricultural, industrial, and commercial sectors. In order to provide comparable information on transportation and mobile sources, Table A-113 provides estimates of CO<sub>2</sub> from these other mobile sources, developed from EPA's NONROAD model and FHWA's *Highway*

*Statistics.* These other mobile source estimates were developed using the same fuel consumption data utilized in developing the N<sub>2</sub>O and CH<sub>4</sub> estimates.

**Table A-113: CO<sub>2</sub> Emissions from Non-Transportation Mobile Sources (MMT CO<sub>2</sub> Eq.)**

| Fuel Type/Vehicle Type                     | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Agricultural Equipment <sup>a</sup>        | 31.0         | 36.6         | 39.3         | 41.5         | 42.7         | 43.7         | 46.7         | 47.4         | 49.6         | 49.0         | 46.0         | 47.2         | 48.2         | 49.9         | 51.4         | 50.4         | 51.2         |
| Construction/Mining Equipment <sup>b</sup> | 42.0         | 48.9         | 56.2         | 60.4         | 62.2         | 64.0         | 65.8         | 66.9         | 68.3         | 68.8         | 70.3         | 71.6         | 73.9         | 75.1         | 76.8         | 82.2         | 80.4         |
| Other Sources <sup>c</sup>                 | 54.5         | 59.9         | 62.6         | 70.1         | 71.8         | 73.7         | 75.7         | 75.9         | 77.1         | 75.9         | 76.6         | 77.3         | 80.2         | 79.9         | 79.4         | 80.3         | 81.7         |
| <b>Total</b>                               | <b>127.6</b> | <b>145.4</b> | <b>158.1</b> | <b>172.0</b> | <b>176.7</b> | <b>181.3</b> | <b>188.2</b> | <b>190.2</b> | <b>195.1</b> | <b>193.7</b> | <b>192.9</b> | <b>196.1</b> | <b>202.3</b> | <b>204.9</b> | <b>207.6</b> | <b>212.9</b> | <b>213.3</b> |

<sup>a</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.

<sup>b</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.

<sup>c</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

Note: In 2015, EPA incorporated the NONROAD2008 model into MOVES2014a. This year's inventory uses the NONROAD component of MOVES2014a for years 1999 through 2014. This update resulted in small changes (<2%) to the 1999-2013 time series for NONROAD fuel consumption due to differences in the gasoline and diesel default fuel densities used within the model iterations.

### Estimation of HFC Emissions from Transportation Sources

In addition to CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> emissions, transportation sources also result in emissions of HFCs. HFCs are emitted to the atmosphere during equipment manufacture and operation (as a result of component failure, leaks, and purges), as well as at servicing and disposal events. There are three categories of transportation-related HFC emissions; Mobile air-conditioning represents the emissions from air conditioning units in passenger cars and light-duty trucks; Comfort Cooling represents the emissions from air conditioning units in passenger trains and buses; and Refrigerated Transport represents the emissions from units used to cool freight during transportation.

Table A-114 below presents these HFC emissions. Table A-115 presents all transportation and mobile source greenhouse gas emissions, including HFC emissions.

**Table A-114: HFC Emissions from Transportation Sources (MMT CO<sub>2</sub>Eq.)**

| Vehicle Type                                | 1990 | 1995        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|---|------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>Mobile AC</b>                            | -    | <b>18.9</b> | <b>53.5</b> | <b>59.1</b> | <b>61.3</b> | <b>62.7</b> | <b>64.3</b> | <b>65.0</b> | <b>65.6</b> | <b>66.0</b> | <b>66.3</b> | <b>65.2</b> | <b>61.7</b> | <b>55.7</b> | <b>49.9</b> | <b>44.0</b> | <b>40.9</b> |
| Passenger Cars                              | -    | 11.2        | 28.1        | 30.7        | 31.5        | 31.6        | 31.8        | 31.7        | 31.7        | 31.5        | 31.2        | 29.9        | 27.5        | 23.9        | 20.6        | 17.3        | 16.0        |
| Light-Duty Trucks                           | -    | 7.8         | 25.4        | 28.4        | 29.8        | 31.1        | 32.4        | 33.3        | 33.9        | 34.5        | 35.1        | 35.2        | 34.2        | 31.7        | 29.3        | 26.7        | 25.0        |
| <b>Comfort Cooling for Trains and Buses</b> | -    | <b>+</b>    | <b>0.1</b>  | <b>0.2</b>  | <b>0.2</b>  | <b>0.2</b>  | <b>0.2</b>  | <b>0.3</b>  | <b>0.3</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.5</b>  | <b>0.5</b>  | <b>0.5</b>  | <b>0.5</b>  | <b>0.5</b>  | <b>0.5</b>  |
| School and Tour Buses                       | -    | +           | 0.1         | 0.2         | 0.2         | 0.2         | 0.2         | 0.3         | 0.3         | 0.3         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         |
| Transit Buses                               | -    | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | 0.1         |
| Rail  | -    | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           |
| <b>Refrigerated Transport</b>               | -    | <b>0.2</b>  | <b>0.8</b>  | <b>1.0</b>  | <b>1.2</b>  | <b>1.4</b>  | <b>1.6</b>  | <b>1.8</b>  | <b>2.0</b>  | <b>2.3</b>  | <b>2.6</b>  | <b>2.9</b>  | <b>3.5</b>  | <b>4.1</b>  | <b>4.7</b>  | <b>5.3</b>  | <b>5.8</b>  |
| Medium- and Heavy-Duty Trucks               | -    | 0.1         | 0.6         | 0.8         | 0.9         | 1.2         | 1.4         | 1.5         | 1.7         | 1.9         | 2.2         | 2.4         | 2.9         | 3.4         | 3.9         | 4.4         | 4.4         |
| Rail  | -    | 0.0         | 0.2         | 0.2         | 0.3         | 0.2         | 0.3         | 0.3         | 0.3         | 0.4         | 0.4         | 0.5         | 0.6         | 0.7         | 0.8         | 0.9         | 1.4         |
| Ships and Other Boats                       | -    | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| <b>Total</b>                                | -    | <b>19.1</b> | <b>54.5</b> | <b>60.3</b> | <b>62.7</b> | <b>64.3</b> | <b>66.1</b> | <b>67.1</b> | <b>67.9</b> | <b>68.7</b> | <b>69.3</b> | <b>68.5</b> | <b>65.6</b> | <b>60.2</b> | <b>55.1</b> | <b>49.8</b> | <b>47.2</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

- Unreported or zero

Note: Totals may not sum due to independent rounding.

## Contribution of Transportation and Mobile Sources to Greenhouse Gas Emissions, by Mode/Vehicle Type/Fuel Type

Table A-115 presents estimates of greenhouse gas emissions from an expanded analysis including all transportation and additional mobile sources, as well as emissions from electricity generation by the consuming category, in CO<sub>2</sub> equivalents. In total, transportation and non-transportation mobile sources emitted 2,029.7 MMT CO<sub>2</sub> Eq. in 2014, an increase of 21 percent from 1990. Transportation sources account for 1,814.5 MMT CO<sub>2</sub> Eq. while non-transportation mobile sources account for 215.3 MMT CO<sub>2</sub> Eq. These estimates include HFC emissions for mobile AC, comfort cooling for trains and buses, and refrigerated transport. These estimates were generated using the estimates of CO<sub>2</sub> emissions from transportation sources reported in the Carbon Dioxide Emissions from Fossil Fuel Combustion section, and CH<sub>4</sub> emissions and N<sub>2</sub>O emissions reported in the Mobile Combustion section of the Energy chapter; information on HFCs from mobile air conditioners, comfort cooling for trains and buses, and refrigerated transportation from the Substitutes for Ozone Depleting Substances section of the IPPU chapter; and estimates of CO<sub>2</sub> emitted from non-transportation mobile sources reported in Table A-111 above.

Although all emissions reported here are based on estimates reported throughout this Inventory, some additional calculations were performed in order to provide a detailed breakdown of emissions by mode and vehicle category. In the case of N<sub>2</sub>O and CH<sub>4</sub>, additional calculations were performed to develop emission estimates by type of aircraft and type of heavy-duty vehicle (i.e., medium- and heavy-duty trucks or buses) to match the level of detail for CO<sub>2</sub> emissions. N<sub>2</sub>O estimates for both jet fuel and aviation gasoline, and CH<sub>4</sub> estimates for aviation gasoline were developed for individual aircraft types by multiplying the emissions estimates for each fuel type (jet fuel and aviation gasoline) by the portion of fuel used by each aircraft type (from FAA 2016 and DESC 2015). Emissions of CH<sub>4</sub> from jet fuels are no longer considered to be emitted from aircraft gas turbine engines burning jet fuel A at higher power settings. This update applies to the entire time series.<sup>54</sup> Recent research indicates that modern aircraft jet engines are typically net consumers of methane (Santoni et al 2011). Methane is emitted at low power and idle operation, but at higher power modes aircraft engines consume methane. Over the range of engine operating modes, aircraft engines are net consumers of methane on average. Based on this data, CH<sub>4</sub> emission factors for jet aircraft were reported as zero to reflect the latest emissions testing data.

Similarly, N<sub>2</sub>O and CH<sub>4</sub> estimates were developed for medium- and heavy-duty trucks and buses by multiplying the emission estimates for heavy-duty vehicles for each fuel type (gasoline, diesel) from the Mobile Combustion section in the Energy chapter, by the portion of fuel used by each vehicle type (from DOE 1993 through 2015). Carbon dioxide emissions from non-transportation mobile sources are calculated using data from EPA's NONROAD component of MOVES2014a (EPA 2015d). Otherwise, the table and figure are drawn directly from emission estimates presented elsewhere in the Inventory, and are dependent on the methodologies presented in Annex 2.1 (for CO<sub>2</sub>), Chapter 4, and Annex 3.8 (for HFCs), and earlier in this Annex (for CH<sub>4</sub> and N<sub>2</sub>O).

Transportation sources include on-road vehicles, aircraft, boats and ships, rail, and pipelines (note: pipelines are a transportation source but are stationary, not mobile sources). In addition, transportation-related greenhouse gas emissions also include HFC released from mobile air-conditioners and refrigerated transport, and the release of CO<sub>2</sub> from lubricants (such as motor oil) used in transportation. Together, transportation sources were responsible for 1,814.5 MMT CO<sub>2</sub> Eq. in 2014.

On-road vehicles were responsible for about 75 percent of all transportation and non-transportation mobile greenhouse gas emissions in 2014. Although passenger cars make up the largest component of on-road vehicle greenhouse gas emissions, medium- and heavy-duty trucks have been the primary sources of growth in on-road vehicle emissions. Between 1990 and 2014, greenhouse gas emissions from passenger cars increased by 16 percent, while emissions from light-duty trucks increased by one percent.<sup>55</sup> Meanwhile, greenhouse gas emissions from medium- and heavy-duty trucks increased 76 percent between 1990 and 2014, reflecting the increased volume of total freight movement and an increasing share transported by trucks.

---

<sup>54</sup> Recommended Best Practice for Quantifying Speciated Organic Gas Emissions from Aircraft Equipped with Turbofan, Turbojet and Turboprop Engines," EPA-420-R-09-901, May 27, 2009 (see <<http://www.epa.gov/otaq/regs/nonroad/aviation/420r09901.pdf>>).

<sup>55</sup> In 2011 FHWA changed how they defined vehicle types for the purposes of reporting VMT for the years 2007-2010. The old approach to vehicle classification was based on body type and split passenger vehicles into "Passenger Cars" and "Other 2 Axle 4-Tire Vehicles". The new approach is a vehicle classification system based on wheelbase. Vehicles with a wheelbase less than or equal to 121 inches are counted as "Light-duty Vehicles—Short Wheelbase". Passenger vehicles with a wheelbase greater than 121 inches are counted as "Light-duty Vehicles - Long Wheelbase". This change in vehicle classification has moved some smaller trucks and sport utility vehicles from the light truck category to the passenger vehicle category in this Inventory. These changes are reflected in a large drop in light-truck emissions between 2006 and 2007.

Greenhouse gas emissions from aircraft decreased 20 percent between 1990 and 2014. Emissions from military aircraft decreased 56 percent between 1990 and 2014. Commercial aircraft emissions rose 27 percent between 1990 and 2007 then dropped 18 percent from 2007 to 2014, a change of approximately 5 percent between 1990 and 2014.

Non-transportation mobile sources, such as construction/mining equipment, agricultural equipment, and industrial/commercial equipment, emitted approximately 215.3 MMT CO<sub>2</sub> Eq. in 2014. Together, these sources emitted more greenhouse gases than ships and boats, and rail combined. Emissions from non-transportation mobile sources increased rapidly, growing approximately 67 percent between 1990 and 2014. Methane and N<sub>2</sub>O emissions from these sources are included in the “Mobile Combustion” section and CO<sub>2</sub> emissions are included in the relevant economic sectors.

#### **Contribution of Transportation and Mobile Sources to Greenhouse Gas Emissions, by Gas**

Table A-116 presents estimates of greenhouse gas emissions from transportation and other mobile sources broken down by greenhouse gas. As this table shows, CO<sub>2</sub> accounts for the vast majority of transportation greenhouse gas emissions (approximately 97 percent in 2014). Emissions of CO<sub>2</sub> from transportation and mobile sources increased by 327.8 MMT CO<sub>2</sub> Eq. between 1990 and 2014. In contrast, the combined emissions of CH<sub>4</sub> and N<sub>2</sub>O decreased by 28.5 MMT CO<sub>2</sub> Eq. over the same period, due largely to the introduction of control technologies designed to reduce criteria pollutant emissions.<sup>56</sup> Meanwhile, HFC emissions from mobile air-conditioners and refrigerated transport increased from virtually no emissions in 1990 to 47.2 MMT CO<sub>2</sub> Eq. in 2014 as these chemicals were phased in as substitutes for ozone depleting substances. It should be noted, however, that the ozone depleting substances that HFCs replaced are also powerful greenhouse gases, but are not included in national greenhouse gas inventories per UNFCCC reporting requirements.

#### **Greenhouse Gas Emissions from Freight and Passenger Transportation**

Table A-117 and

Table A-118 present greenhouse gas estimates from transportation, broken down into the passenger and freight categories. Passenger modes include light-duty vehicles, buses, passenger rail, aircraft (general aviation and commercial aircraft), recreational boats, and mobile airconditioners, and are illustrated in Table A-117. Freight modes include medium- and heavy-duty trucks, freight rail, refrigerated transport, waterborne freight vessels, pipelines, and commercial aircraft and are illustrated in

Table A-118. Commercial aircraft do carry some freight, in addition to passengers, and emissions have been split between passenger and freight transportation. The amount of commercial aircraft emissions to allocate to the passenger and freight categories was calculated using BTS data on freight shipped by commercial aircraft, and the total number of passengers enplaned. Each passenger was considered to weigh an average of 150 pounds, with a luggage weight of 50 pounds. The total freight weight and total passenger weight carried were used to determine percent shares which were used to split the total commercial aircraft emission estimates. The remaining transportation and mobile emissions were from sources not considered to be either freight or passenger modes (e.g., construction/mining and agricultural equipment, lubricants).

The estimates in these tables are derived from the estimates presented in Table A-115. In addition, estimates of fuel consumption from DOE (1993 through 2015) were used to allocate rail emissions between passenger and freight categories.

In 2014, passenger transportation modes emitted 1,261.5 MMT CO<sub>2</sub> Eq., while freight transportation modes emitted 518.3 MMT CO<sub>2</sub> Eq. Between 1990 and 2014, the percentage growth of greenhouse gas emissions from freight sources was 47 percent, while emissions from passenger sources grew by 9 percent. This difference in growth is due largely to the rapid increase in emissions associated with medium- and heavy-duty trucks.

---

<sup>56</sup> The decline in CFC emissions is not captured in the official transportation estimates.

**Table A-115: Total U.S. Greenhouse Gas Emissions from Transportation and Mobile Sources (MMT CO<sub>2</sub> Eq.)**

| Mode / Vehicle Type / Fuel Type               | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           | Percent Change 1990-2014 |
|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------------------|
| <b>Transportation Total<sup>a</sup></b>       | <b>1,554.4</b> | <b>1,698.4</b> | <b>1,926.7</b> | <b>1,911.3</b> | <b>1,951.4</b> | <b>1,941.2</b> | <b>1,986.0</b> | <b>2,004.4</b> | <b>1,998.1</b> | <b>1,999.3</b> | <b>1,902.0</b> | <b>1,823.6</b> | <b>1,832.0</b> | <b>1,803.9</b> | <b>1,784.3</b> | <b>1,794.0</b> | <b>1,814.5</b> | <b>17%</b>               |
| <b>On-Road Vehicles</b>                       | <b>1,233.5</b> | <b>1,370.5</b> | <b>1,572.8</b> | <b>1,583.1</b> | <b>1,621.4</b> | <b>1,630.6</b> | <b>1,663.8</b> | <b>1,672.4</b> | <b>1,668.5</b> | <b>1,664.6</b> | <b>1,586.5</b> | <b>1,540.9</b> | <b>1,541.7</b> | <b>1,515.0</b> | <b>1,504.6</b> | <b>1,504.3</b> | <b>1,531.1</b> | <b>24%</b>               |
| <b>Passenger Cars</b>                         | <b>656.6</b>   | <b>646.7</b>   | <b>697.3</b>   | <b>704.1</b>   | <b>716.9</b>   | <b>694.2</b>   | <b>690.2</b>   | <b>708.9</b>   | <b>683.0</b>   | <b>843.5</b>   | <b>802.3</b>   | <b>792.8</b>   | <b>783.6</b>   | <b>774.3</b>   | <b>767.9</b>   | <b>763.2</b>   | <b>762.5</b>   | <b>16%</b>               |
| Gasoline <sup>b</sup>                         | 648.7          | 627.6          | 665.5          | 669.7          | 681.7          | 658.4          | 654.1          | 673.0          | 647.2          | 807.9          | 767.4          | 759.3          | 752.3          | 746.3          | 743.2          | 741.8          | 742.4          | 14%                      |
| Diesel <sup>b, c</sup>                        | 7.9            | 7.8            | 3.7            | 3.7            | 3.7            | 4.2            | 4.3            | 4.2            | 4.1            | 4.1            | 3.7            | 3.6            | 3.7            | 4.1            | 4.1            | 4.1            | 4.1            | -48%                     |
| AFVs <sup>d</sup>                             | +              | 0.1            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 5,608%                   |
| HFCs from Mobile AC                           | +              | 11.2           | 28.1           | 30.7           | 31.5           | 31.6           | 31.8           | 31.7           | 31.7           | 31.5           | 31.2           | 29.9           | 27.5           | 23.9           | 20.6           | 17.3           | 16.0           | NA                       |
| <b>Light-Duty Trucks</b>                      | <b>335.6</b>   | <b>436.8</b>   | <b>515.0</b>   | <b>520.8</b>   | <b>532.7</b>   | <b>567.4</b>   | <b>589.0</b>   | <b>551.5</b>   | <b>564.0</b>   | <b>367.2</b>   | <b>348.6</b>   | <b>351.5</b>   | <b>348.9</b>   | <b>332.0</b>   | <b>326.0</b>   | <b>323.4</b>   | <b>338.1</b>   | <b>1%</b>                |
| Gasoline <sup>b</sup>                         | 323.5          | 413.6          | 469.0          | 471.0          | 480.4          | 508.3          | 526.7          | 491.1          | 502.2          | 318.1          | 299.7          | 303.1          | 300.9          | 285.8          | 282.1          | 281.9          | 297.4          | -8%                      |
| Diesel <sup>b, c</sup>                        | 11.5           | 14.9           | 20.1           | 20.8           | 21.9           | 27.2           | 28.9           | 25.8           | 26.7           | 13.5           | 12.1           | 12.0           | 12.5           | 13.0           | 12.9           | 12.9           | 13.9           | 21%                      |
| AFVs <sup>d</sup>                             | 0.6            | 0.5            | 0.5            | 0.6            | 0.6            | 0.8            | 0.9            | 1.3            | 1.2            | 1.0            | 1.8            | 1.2            | 1.3            | 1.5            | 1.6            | 1.9            | 1.9            | 234%                     |
| HFCs from Mobile AC                           | +              | 7.8            | 25.4           | 28.4           | 29.8           | 31.1           | 32.4           | 33.3           | 33.9           | 34.5           | 35.1           | 35.2           | 34.2           | 31.7           | 29.3           | 26.7           | 25.0           | NA                       |
| <b>Medium- and Heavy-Duty Trucks</b>          | <b>231.1</b>   | <b>275.9</b>   | <b>347.5</b>   | <b>346.1</b>   | <b>360.0</b>   | <b>356.6</b>   | <b>367.7</b>   | <b>398.2</b>   | <b>407.4</b>   | <b>432.0</b>   | <b>414.2</b>   | <b>376.3</b>   | <b>389.7</b>   | <b>388.4</b>   | <b>388.7</b>   | <b>395.7</b>   | <b>407.4</b>   | <b>76%</b>               |
| Gasoline <sup>b</sup>                         | 39.5           | 36.8           | 37.0           | 36.1           | 36.6           | 31.6           | 31.8           | 35.7           | 36.1           | 47.1           | 47.3           | 43.4           | 43.2           | 39.6           | 39.3           | 40.1           | 40.5           | 2%                       |
| Diesel <sup>b, c</sup>                        | 190.7          | 238.4          | 309.5          | 309.0          | 322.2          | 323.5          | 334.3          | 360.5          | 369.1          | 382.6          | 364.0          | 329.9          | 343.1          | 344.7          | 344.8          | 350.4          | 361.7          | 90%                      |
| AFVs <sup>d</sup>                             | 0.9            | 0.6            | 0.3            | 0.3            | 0.3            | 0.4            | 0.3            | 0.4            | 0.5            | 0.4            | 0.7            | 0.5            | 0.6            | 0.6            | 0.7            | 0.8            | 0.8            | -8%                      |
| HFCs from Refrigerated Transport <sup>e</sup> | +              | 0.1            | 0.6            | 0.8            | 0.9            | 1.2            | 1.4            | 1.5            | 1.7            | 1.9            | 2.2            | 2.4            | 2.9            | 3.4            | 3.9            | 4.4            | 4.4            | NA                       |
| <b>Buses</b>                                  | <b>8.4</b>     | <b>9.2</b>     | <b>11.2</b>    | <b>10.3</b>    | <b>10.0</b>    | <b>10.8</b>    | <b>15.1</b>    | <b>12.1</b>    | <b>12.2</b>    | <b>17.6</b>    | <b>16.9</b>    | <b>16.0</b>    | <b>15.8</b>    | <b>16.8</b>    | <b>17.8</b>    | <b>18.0</b>    | <b>19.1</b>    | <b>129%</b>              |
| Gasoline <sup>b</sup>                         | 0.4            | 0.4            | 0.4            | 0.4            | 0.3            | 0.3            | 0.5            | 0.4            | 0.4            | 0.7            | 0.7            | 0.8            | 0.7            | 0.7            | 0.8            | 0.9            | 0.9            | 154%                     |
| Diesel <sup>b, c</sup>                        | 8.0            | 8.7            | 10.2           | 9.3            | 8.8            | 9.5            | 13.5           | 10.6           | 10.6           | 15.5           | 14.6           | 13.6           | 13.5           | 14.4           | 15.4           | 15.5           | 16.6           | 107%                     |
| AFVs <sup>d</sup>                             | +              | 0.1            | 0.5            | 0.5            | 0.6            | 0.8            | 0.9            | 0.9            | 0.9            | 1.0            | 1.2            | 1.2            | 1.1            | 1.2            | 1.1            | 1.2            | 1.2            | 88,269%                  |
| HFCs from Comfort Cooling                     | +              | 0.0            | 0.1            | 0.2            | 0.2            | 0.2            | 0.2            | 0.3            | 0.3            | 0.4            | 0.4            | 0.4            | 0.4            | 0.4            | 0.4            | 0.4            | 0.4            | NA                       |
| <b>Motorcycles</b>                            | <b>1.8</b>     | <b>1.8</b>     | <b>1.9</b>     | <b>1.7</b>     | <b>1.7</b>     | <b>1.7</b>     | <b>1.8</b>     | <b>1.7</b>     | <b>1.9</b>     | <b>4.3</b>     | <b>4.4</b>     | <b>4.2</b>     | <b>3.7</b>     | <b>3.6</b>     | <b>4.2</b>     | <b>4.0</b>     | <b>3.9</b>     | <b>118%</b>              |
| Gasoline <sup>b</sup>                         | 1.8            | 1.8            | 1.9            | 1.7            | 1.7            | 1.7            | 1.8            | 1.7            | 1.9            | 4.3            | 4.4            | 4.2            | 3.7            | 3.6            | 4.2            | 4.0            | 3.9            | 118%                     |
| <b>Aircraft General Aviation</b>              | <b>189.2</b>   | <b>176.7</b>   | <b>199.4</b>   | <b>193.9</b>   | <b>189.4</b>   | <b>183.1</b>   | <b>190.6</b>   | <b>193.6</b>   | <b>186.3</b>   | <b>183.4</b>   | <b>176.7</b>   | <b>157.4</b>   | <b>154.8</b>   | <b>149.9</b>   | <b>146.5</b>   | <b>150.1</b>   | <b>151.5</b>   | <b>-20%</b>              |
| <b>Aircraft</b>                               | <b>42.9</b>    | <b>35.8</b>    | <b>35.9</b>    | <b>43.7</b>    | <b>45.1</b>    | <b>36.9</b>    | <b>41.9</b>    | <b>40.1</b>    | <b>30.1</b>    | <b>24.4</b>    | <b>30.5</b>    | <b>21.2</b>    | <b>26.7</b>    | <b>22.5</b>    | <b>19.9</b>    | <b>23.6</b>    | <b>19.7</b>    | <b>-54%</b>              |

|  |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |             |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|-------------|
| Jet Fuel   | 39.8         | 33.0         | 33.4         | 41.2         | 42.7         | 34.7         | 39.7         | 37.6         | 27.7         | 22.2         | 28.5         | 19.4         | 24.8         | 20.6         | 18.2         | 22.0         | 18.2         | -54%        |
| Aviation Gasoline                                      | 3.2          | 2.8          | 2.6          | 2.5          | 2.4          | 2.1          | 2.2          | 2.5          | 2.4          | 2.2          | 2.0          | 1.9          | 1.9          | 1.9          | 1.8          | 1.6          | 1.5          | -52%        |
| <b>Commercial Aircraft</b>                             | <b>110.9</b> | <b>116.3</b> | <b>140.6</b> | <b>125.8</b> | <b>122.4</b> | <b>124.0</b> | <b>126.0</b> | <b>134.0</b> | <b>138.3</b> | <b>141.0</b> | <b>128.4</b> | <b>120.6</b> | <b>114.4</b> | <b>115.7</b> | <b>114.3</b> | <b>115.4</b> | <b>116.3</b> | <b>5%</b>   |
| Jet Fuel   | 110.9        | 116.3        | 140.6        | 125.8        | 122.4        | 124.0        | 126.0        | 134.0        | 138.3        | 141.0        | 128.4        | 120.6        | 114.4        | 115.7        | 114.3        | 115.4        | 116.3        | 5%          |
| <b>Military Aircraft</b>                               | <b>35.3</b>  | <b>24.5</b>  | <b>22.9</b>  | <b>24.5</b>  | <b>21.9</b>  | <b>22.2</b>  | <b>22.7</b>  | <b>19.5</b>  | <b>18.0</b>  | <b>18.0</b>  | <b>17.7</b>  | <b>15.5</b>  | <b>13.7</b>  | <b>11.7</b>  | <b>12.2</b>  | <b>11.1</b>  | <b>15.5</b>  | <b>-56%</b> |
| Jet Fuel   | 35.3         | 24.5         | 22.9         | 24.5         | 21.9         | 22.2         | 22.7         | 19.5         | 18.0         | 18.0         | 17.7         | 15.5         | 13.7         | 11.7         | 12.2         | 11.1         | 15.5         | -56%        |
| <b>Ships and Boats<sup>g</sup></b>                     | <b>44.9</b>  | <b>58.5</b>  | <b>61.1</b>  | <b>42.4</b>  | <b>47.2</b>  | <b>37.1</b>  | <b>39.8</b>  | <b>44.9</b>  | <b>48.0</b>  | <b>54.7</b>  | <b>45.3</b>  | <b>38.6</b>  | <b>44.7</b>  | <b>46.4</b>  | <b>40.1</b>  | <b>39.4</b>  | <b>28.6</b>  | <b>-36%</b> |
| Gasoline   | 12.4         | 14.1         | 10.2         | 14.4         | 14.4         | 14.2         | 14.1         | 14.0         | 13.8         | 13.6         | 13.2         | 13.0         | 12.7         | 12.6         | 12.5         | 12.4         | 12.3         | 0%          |
| Distillate Fuel <sup>c</sup>                           | 9.6          | 14.9         | 17.1         | 15.8         | 15.4         | 15.3         | 11.5         | 11.4         | 10.8         | 11.6         | 11.4         | 11.5         | 11.1         | 14.0         | 11.4         | 11.5         | 10.2         | 6%          |
| Residual Fuel <sup>h</sup>                             | 22.9         | 29.5         | 33.8         | 12.2         | 17.4         | 7.6          | 14.2         | 19.6         | 23.4         | 29.4         | 20.7         | 14.2         | 20.9         | 19.8         | 16.2         | 15.5         | 6.0          | -75%        |
| HFCs from Refrigerated Transport <sup>e</sup>          | +            | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | NA          |
| <b>Rail</b>  | <b>39.0</b>  | <b>43.1</b>  | <b>46.1</b>  | <b>46.5</b>  | <b>46.1</b>  | <b>47.8</b>  | <b>50.5</b>  | <b>51.1</b>  | <b>53.0</b>  | <b>52.2</b>  | <b>48.5</b>  | <b>41.3</b>  | <b>44.2</b>  | <b>45.9</b>  | <b>44.6</b>  | <b>45.5</b>  | <b>47.6</b>  | <b>22%</b>  |
| Distillate Fuel <sup>c,i</sup>                         | 35.8         | 40.0         | 42.5         | 42.6         | 42.3         | 43.2         | 45.6         | 46.0         | 48.1         | 46.7         | 43.3         | 36.3         | 39.0         | 40.8         | 39.9         | 40.5         | 42.1         | 17%         |
| Electricity  | 3.1          | 3.1          | 3.5          | 3.7          | 3.5          | 4.3          | 4.6          | 4.8          | 4.6          | 5.1          | 4.7          | 4.5          | 4.5          | 4.3          | 3.9          | 4.1          | 4.1          | 34%         |
| Other Emissions from Rail Electricity Use <sup>j</sup> | 0.1          | 0.1          | +            | +            | +            | +            | 0.1          | 0.1          | +            | 0.1          | +            | +            | +            | +            | +            | +            | +            | -30%        |
| HFCs from Comfort Cooling                              | +            | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | NA          |
| HFCs from Refrigerated Transport <sup>e</sup>          | +            | 0.0          | 0.2          | 0.2          | 0.3          | 0.2          | 0.3          | 0.3          | 0.3          | 0.4          | 0.4          | 0.5          | 0.6          | 0.7          | 0.8          | 0.9          | 1.4          | NA          |
| <b>Pipelines<sup>k</sup></b>                           | <b>36.0</b>  | <b>38.2</b>  | <b>35.2</b>  | <b>34.4</b>  | <b>36.4</b>  | <b>32.5</b>  | <b>31.1</b>  | <b>32.2</b>  | <b>32.3</b>  | <b>34.2</b>  | <b>35.6</b>  | <b>36.7</b>  | <b>37.1</b>  | <b>37.8</b>  | <b>40.3</b>  | <b>45.9</b>  | <b>46.5</b>  | <b>29%</b>  |
| Natural Gas  | 36.0         | 38.2         | 35.2         | 34.4         | 36.4         | 32.5         | 31.1         | 32.2         | 32.3         | 34.2         | 35.6         | 36.7         | 37.1         | 37.8         | 40.3         | 45.9         | 46.5         | 29%         |
| <b>Other Transportation</b>                            | <b>11.8</b>  | <b>11.3</b>  | <b>12.1</b>  | <b>11.1</b>  | <b>10.9</b>  | <b>10.1</b>  | <b>10.2</b>  | <b>10.2</b>  | <b>9.9</b>   | <b>10.2</b>  | <b>9.5</b>   | <b>8.5</b>   | <b>9.5</b>   | <b>9.0</b>   | <b>8.3</b>   | <b>8.8</b>   | <b>9.1</b>   | <b>-23%</b> |
| Lubricants   | 11.8         | 11.3         | 12.1         | 11.1         | 10.9         | 10.1         | 10.2         | 10.2         | 9.9          | 10.2         | 9.5          | 8.5          | 9.5          | 9.0          | 8.3          | 8.8          | 9.1          | -23%        |
| <b>Non-Transportation Mobile Total</b>                 | <b>128.8</b> | <b>146.8</b> | <b>159.6</b> | <b>173.6</b> | <b>178.3</b> | <b>183.0</b> | <b>190.0</b> | <b>191.9</b> | <b>196.9</b> | <b>195.5</b> | <b>194.7</b> | <b>198.0</b> | <b>204.2</b> | <b>206.8</b> | <b>209.6</b> | <b>214.9</b> | <b>215.3</b> | <b>67%</b>  |
| <b>Agricultural Equipment<sup>l</sup></b>              | <b>31.4</b>  | <b>37.0</b>  | <b>39.8</b>  | <b>42.0</b>  | <b>43.1</b>  | <b>44.2</b>  | <b>47.2</b>  | <b>47.9</b>  | <b>50.2</b>  | <b>49.5</b>  | <b>46.5</b>  | <b>47.7</b>  | <b>48.7</b>  | <b>50.5</b>  | <b>52.0</b>  | <b>51.0</b>  | <b>51.7</b>  | <b>65%</b>  |
| Gasoline   | 7.3          | 8.3          | 5.8          | 7.1          | 7.4          | 7.6          | 9.7          | 9.6          | 11.0         | 9.4          | 5.6          | 5.9          | 6.0          | 7.0          | 7.6          | 5.7          | 5.6          | -23%        |
| Diesel   | 24.1         | 28.7         | 34.0         | 34.9         | 35.7         | 36.6         | 37.5         | 38.4         | 39.2         | 40.1         | 40.9         | 41.8         | 42.7         | 43.5         | 44.4         | 45.3         | 46.1         | 92%         |
| <b>Construction/ Mining Equipment<sup>m</sup></b>      | <b>42.4</b>  | <b>49.4</b>  | <b>56.6</b>  | <b>61.0</b>  | <b>62.7</b>  | <b>64.5</b>  | <b>66.4</b>  | <b>67.4</b>  | <b>68.9</b>  | <b>69.4</b>  | <b>70.9</b>  | <b>72.2</b>  | <b>74.6</b>  | <b>75.7</b>  | <b>77.4</b>  | <b>82.9</b>  | <b>81.1</b>  | <b>91%</b>  |
| Gasoline   | 4.4          | 4.0          | 3.0          | 5.8          | 6.0          | 6.3          | 6.6          | 6.2          | 6.1          | 5.1          | 5.1          | 4.9          | 5.7          | 5.3          | 5.5          | 9.4          | 6.1          | 39%         |
| Diesel   | 38.0         | 45.4         | 53.6         | 55.1         | 56.7         | 58.2         | 59.7         | 61.3         | 62.8         | 64.3         | 65.8         | 67.4         | 68.9         | 70.4         | 72.0         | 73.5         | 75.0         | 97%         |

|   |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |            |
|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|------------|
| <b>Other Equipment<sup>a</sup></b>                        | <b>55.0</b>    | <b>60.4</b>    | <b>63.2</b>    | <b>70.7</b>    | <b>72.5</b>    | <b>74.3</b>    | <b>76.4</b>    | <b>76.6</b>    | <b>77.8</b>    | <b>76.6</b>    | <b>77.3</b>    | <b>78.0</b>    | <b>80.9</b>    | <b>80.6</b>    | <b>80.1</b>    | <b>81.0</b>    | <b>82.5</b>    | <b>50%</b> |
| Gasoline  | 40.3           | 42.6           | 42.0           | 48.8           | 49.9           | 51.1           | 52.4           | 51.9           | 52.5           | 50.6           | 50.6           | 50.6           | 52.9           | 51.9           | 50.7           | 50.9           | 51.6           | 28%        |
| Diesel  | 14.7           | 17.8           | 21.2           | 21.9           | 22.6           | 23.3           | 23.9           | 24.6           | 25.3           | 26.0           | 26.7           | 27.4           | 28.1           | 28.7           | 29.4           | 30.1           | 30.8           | 109%       |
| <b>Transportation and Non-Transportation Mobile Total</b> | <b>1,683.1</b> | <b>1,845.2</b> | <b>2,086.4</b> | <b>2,085.0</b> | <b>2,129.7</b> | <b>2,124.2</b> | <b>2,176.0</b> | <b>2,196.4</b> | <b>2,195.0</b> | <b>2,194.8</b> | <b>2,096.7</b> | <b>2,021.5</b> | <b>2,036.2</b> | <b>2,010.8</b> | <b>1,993.9</b> | <b>2,008.9</b> | <b>2,029.7</b> | <b>21%</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

NA - Not Applicable, as there were no HFC emissions allocated to the transport sector in 1990, and thus a growth rate cannot be calculated.

<sup>a</sup> Not including emissions from international bunker fuels.

<sup>b</sup> Gasoline and diesel highway vehicle fuel consumption estimates are based on data from FHWA Highway Statistics Table VM-1 and MF-27 (FHWA 1996 through 2015). These fuel consumption estimates are combined with estimates of fuel shares by vehicle type from DOE's TEDB Annex Tables A.1 through A.6 (DOE 1993 through 2015). TEDB data for 2014 has not been published yet, therefore 2013 data is as a proxy.

<sup>c</sup> Updates to the distillate fuel oil heat content data from EIA for years 1993 through present resulted in changes to the time series for energy consumption and emissions compared to the previous Inventory.

<sup>d</sup> In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data. EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated in this year's inventory and apply to the 1990-2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes.

<sup>e</sup> Updated Commodity Flow Survey data was used to allocate refrigerated transport emissions to the trucking, rail and marine sectors in 2014.

<sup>f</sup> Updates to the jet fuel heat content used in the mobile N<sub>2</sub>O emissions estimates for years 1990 through present resulted in small changes to the time series emissions compared to the previous Inventory.

<sup>g</sup> Fluctuations in emission estimates reflect data collection problems. Note that CH<sub>4</sub> and N<sub>2</sub>O from U.S. Territories are included in this value, but not CO<sub>2</sub> emissions from U.S. Territories, which are estimated separately in the section on U.S. Territories.

<sup>h</sup> Domestic residual fuel for ships and boats is estimated by taking the total amount of residual fuel and subtracting out an estimate of international bunker fuel use.

<sup>i</sup> Class II and Class III diesel consumption data for 2014 is not available yet, therefore 2013 data is used as a proxy.

<sup>j</sup> Other emissions from electricity generation are a result of waste incineration (as the majority of municipal solid waste is combusted in "trash-to-steam" electricity generation plants), electrical transmission and distribution, and a portion of Other Process Uses of Carbonates (from pollution control equipment installed in electricity generation plants).

<sup>k</sup> Includes only CO<sub>2</sub> from natural gas used to power natural gas pipelines; does not include emissions from electricity use or non-CO<sub>2</sub> gases.

<sup>l</sup> Includes equipment, such as tractors and combines, as well as fuel consumption from trucks that are used off-road in agriculture.

<sup>m</sup> Includes equipment, such as cranes, dumpers, and excavators, as well as fuel consumption from trucks that are used off-road in construction.

<sup>n</sup> "Other" includes snowmobiles and other recreational equipment, logging equipment, lawn and garden equipment, railroad equipment, airport equipment, commercial equipment, and industrial equipment, as well as fuel consumption from trucks that are used off-road for commercial/industrial purposes.

Notes: Increases to CH<sub>4</sub> and N<sub>2</sub>O emissions from mobile combustion relative to previous Inventories are largely due to updates made to the Motor Vehicle Emissions Simulator (MOVES2014a) model that is used to estimate on-road gasoline vehicle distribution and mileage across the time series. See Section 3.1 "CH<sub>4</sub> and N<sub>2</sub>O from Mobile Combustion" for more detail. In 2015, EPA incorporated the NONROAD2008 model into MOVES2014a. This year's inventory uses the NONROAD component of MOVES2014a for years 1999 through 2014. This update resulted in small changes (<2%) to the 1999-2013 time series for NONROAD fuel consumption due to differences in the gasoline and diesel default fuel densities used within the model iterations.

**Table A-116: Transportation and Mobile Source Emissions by Gas (MMT CO<sub>2</sub> Eq.)**

|                          | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           | Percent Change 1990-2014 |
|--------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------------------|
| CO <sub>2</sub>          | 1,636.2        | 1,769.6        | 1,979.1        | 1,976.1        | 2,021.7        | 2,018.0        | 2,070.1        | 2,092.1        | 2,092.1        | 2,095.1        | 1,998.9        | 1,926.1        | 1,944.5        | 1,925.8        | 1,916.5        | 1,938.7        | 1,964.1        | 20%                      |
| N <sub>2</sub> O         | 41.2           | 51.2           | 49.0           | 45.0           | 42.1           | 38.9           | 36.8           | 34.4           | 32.2           | 28.4           | 26.0           | 24.5           | 23.6           | 22.4           | 20.0           | 18.2           | 16.3           | -60%                     |
| CH <sub>4</sub>          | 5.6            | 5.2            | 3.7            | 3.5            | 3.2            | 3.0            | 2.9            | 2.7            | 2.7            | 2.5            | 2.4            | 2.3            | 2.3            | 2.2            | 2.2            | 2.1            | 2.0            | -64%                     |
| HFC                      | -              | 19.1           | 54.5           | 60.3           | 62.7           | 64.3           | 66.1           | 67.1           | 67.9           | 68.7           | 69.3           | 68.5           | 65.6           | 60.2           | 55.1           | 49.8           | 47.2           | NA                       |
| <b>Total<sup>a</sup></b> | <b>1,683.1</b> | <b>1,845.1</b> | <b>2,086.3</b> | <b>2,084.9</b> | <b>2,129.7</b> | <b>2,124.2</b> | <b>2,175.9</b> | <b>2,196.3</b> | <b>2,194.9</b> | <b>2,194.7</b> | <b>2,096.7</b> | <b>2,021.5</b> | <b>2,036.1</b> | <b>2,010.7</b> | <b>1,993.8</b> | <b>2,008.8</b> | <b>2,029.6</b> | <b>21%</b>               |

- Unreported or zero


NA - Not applicable, as there were no HFC emissions allocated to the transport sector in 1990, and thus a growth rate cannot be calculated.

<sup>a</sup> Total excludes other emissions from electricity generation and CH<sub>4</sub> and N<sub>2</sub>O emissions from electric rail.

Note: The current Inventory includes updated vehicle population data based on the MOVES 2014a Model.

Gasoline and diesel highway vehicle mileage are based on data from FHWA Highway Statistics Table VM-1 (FHWA 1996 through 2015). These mileage consumption estimates are combined with estimates of fuel shares by vehicle type from DOE's TEDB Annex Tables A.1 through A.6 (DOE 1993 through 2015). TEDB data for 2014 has not been published yet, therefore 2013 data is used as a proxy.

**Figure A-4: Domestic Greenhouse Gas Emissions by Mode and Vehicle Type, 1990 to 2014 (MMT CO<sub>2</sub> Eq.)**



**Table A-17: Greenhouse Gas Emissions from Passenger Transportation (MMT CO<sub>2</sub> Eq.)**

| Vehicle Type                        | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           | Percent Change 1990-2014 |
|-------------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|--------------------------|
| <b>On-Road Vehicles<sup>a</sup></b> | <b>1,002.4</b> | <b>1,094.6</b> | <b>1,225.3</b> | <b>1,236.9</b> | <b>1,261.4</b> | <b>1,274.0</b> | <b>1,296.0</b> | <b>1,274.2</b> | <b>1,261.1</b> | <b>1,232.6</b> | <b>1,172.3</b> | <b>1,164.6</b> | <b>1,151.9</b> | <b>1,126.6</b> | <b>1,115.9</b> | <b>1,108.6</b> | <b>1,123.7</b> | <b>12%</b>               |
| Passenger Cars                      | 656.6          | 646.7          | 697.3          | 704.1          | 716.9          | 694.2          | 690.2          | 708.9          | 683.0          | 843.5          | 802.3          | 792.8          | 783.6          | 774.3          | 767.9          | 763.2          | 762.5          | 16%                      |
| Light-Duty Trucks                   | 335.6          | 436.8          | 515.0          | 520.8          | 532.7          | 567.4          | 589.0          | 551.5          | 564.0          | 367.2          | 348.6          | 351.5          | 348.9          | 332.0          | 326.0          | 323.4          | 338.1          | 1%                       |
| Buses                               | 8.4            | 9.2            | 11.2           | 10.3           | 10.0           | 10.8           | 15.1           | 12.1           | 12.2           | 17.6           | 16.9           | 16.0           | 15.8           | 16.8           | 17.8           | 18.0           | 19.1           | 129%                     |
| Motorcycles                         | 1.8            | 1.8            | 1.9            | 1.7            | 1.7            | 1.7            | 1.8            | 1.7            | 1.9            | 4.3            | 4.4            | 4.2            | 3.7            | 3.6            | 4.2            | 4.0            | 3.9            | 118%                     |
| <b>Aircraft</b>                     | <b>134.6</b>   | <b>132.0</b>   | <b>152.2</b>   | <b>147.7</b>   | <b>146.3</b>   | <b>139.4</b>   | <b>146.8</b>   | <b>152.7</b>   | <b>146.6</b>   | <b>144.9</b>   | <b>140.9</b>   | <b>125.2</b>   | <b>124.8</b>   | <b>122.1</b>   | <b>118.5</b>   | <b>123.1</b>   | <b>119.7</b>   | <b>-11%</b>              |
| General Aviation                    | 42.9           | 35.8           | 35.9           | 43.7           | 45.1           | 36.9           | 41.9           | 40.1           | 30.1           | 24.4           | 30.5           | 21.2           | 26.7           | 22.5           | 19.9           | 23.6           | 19.7           | -54%                     |
| Commercial Aircraft                 | 91.7           | 96.2           | 116.3          | 104.0          | 101.2          | 102.6          | 104.9          | 112.6          | 116.5          | 120.4          | 110.4          | 103.9          | 98.0           | 99.6           | 98.6           | 99.5           | 100.0          | 9%                       |

|                    |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |           |
|--------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|-----------|
| Recreational Boats | 14.3           | 16.4           | 13.0           | 17.2           | 17.3           | 17.2           | 17.2           | 17.2           | 17.1           | 17.0           | 16.6           | 16.5           | 16.3           | 16.2           | 16.2           | 12.4           | 12.3           | -14%      |
| Passenger Rail     | 4.4            | 4.5            | 5.2            | 5.4            | 5.1            | 5.8            | 6.0            | 6.2            | 6.0            | 6.6            | 6.2            | 6.1            | 6.2            | 5.9            | 5.5            | 5.8            | 5.8            | 33%       |
| <b>Total</b>       | <b>1,155.7</b> | <b>1,247.5</b> | <b>1,395.7</b> | <b>1,407.2</b> | <b>1,430.1</b> | <b>1,436.5</b> | <b>1,466.1</b> | <b>1,450.3</b> | <b>1,430.8</b> | <b>1,401.0</b> | <b>1,336.0</b> | <b>1,312.4</b> | <b>1,299.1</b> | <b>1,270.9</b> | <b>1,256.1</b> | <b>1,249.8</b> | <b>1,261.5</b> | <b>9%</b> |

<sup>a</sup>The current Inventory includes updated vehicle population data based on the MOVES 2014a Model.

Notes: Data from DOE (1993 through 2015) were used to disaggregate emissions from rail and buses. Emissions from HFCs have been included in these estimates. Updates to the distillate fuel oil heat content data from EIA for years 1993 through present resulted in changes to the time series for energy consumption and emissions compared to the previous Inventory. In 2015, EPA incorporated the NONROAD2008 model into MOVES2014a. This year's inventory uses the NONROAD component of MOVES2014a for years 1999 through 2014. This update resulted in small changes (<2%) to the 1999-2013 time series for NONROAD fuel consumption due to differences in the gasoline and diesel default fuel densities used within the model iterations. In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data. EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated in this year's inventory and apply to the 1990-2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes

**Table A-118: Greenhouse Gas Emissions from Domestic Freight Transportation (MMT CO<sub>2</sub> Eq.)**

| By Mode                | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         | Percent Change 1990-2014 |
|------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------------------|
| Trucking <sup>a</sup>  | 231.1        | 275.9        | 347.5        | 346.1        | 360.0        | 356.6        | 367.7        | 398.2        | 407.4        | 432.0        | 414.2        | 376.3        | 389.7        | 388.4        | 388.7        | 395.7        | 407.4        | 76%                      |
| Freight Rail           | 34.5         | 38.6         | 40.9         | 41.1         | 40.9         | 41.9         | 44.4         | 44.8         | 47.0         | 45.6         | 42.2         | 35.1         | 38.0         | 39.9         | 39.0         | 39.7         | 41.8         | 21%                      |
| Ships and Other        |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |                          |
| Boats                  | 30.6         | 42.1         | 48.1         | 25.1         | 29.9         | 19.8         | 22.6         | 27.8         | 31.0         | 37.7         | 28.7         | 22.2         | 28.5         | 30.2         | 23.9         | 15.7         | 6.3          | -80%                     |
| Pipelines <sup>b</sup> | 36.0         | 38.2         | 35.2         | 34.4         | 36.4         | 32.5         | 31.1         | 32.2         | 32.3         | 34.2         | 35.6         | 36.7         | 37.1         | 37.8         | 40.3         | 45.9         | 46.5         | 29%                      |
| Commercial             |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |                          |
| Aircraft               | 19.2         | 20.1         | 24.3         | 21.8         | 21.2         | 21.5         | 21.1         | 21.4         | 21.8         | 20.5         | 18.0         | 16.7         | 16.3         | 16.0         | 15.8         | 15.9         | 16.2         | -15%                     |
| <b>Total</b>           | <b>351.4</b> | <b>415.0</b> | <b>496.1</b> | <b>468.5</b> | <b>488.4</b> | <b>472.3</b> | <b>487.0</b> | <b>524.3</b> | <b>539.4</b> | <b>570.0</b> | <b>538.7</b> | <b>487.0</b> | <b>509.6</b> | <b>512.3</b> | <b>507.7</b> | <b>513.0</b> | <b>518.3</b> | <b>47%</b>               |

<sup>a</sup> The current Inventory includes updated vehicle population data based on the MOVES 2014a Model.

<sup>b</sup> Pipelines reflect CO<sub>2</sub> emissions from natural gas powered pipelines transporting natural gas

Note: Data from DOE (1993 through 2015) were used to disaggregate emissions from rail and buses. Emissions from HFCs have been included in these estimates.

Note: In 2015, EPA incorporated the NONROAD2008 model into MOVES2014a. This year's inventory uses the NONROAD component of MOVES2014a for years 1999 through 2014. This update resulted in small changes (<2 percent) to the 1999 to 2013 time series for NONROAD fuel consumption due to differences in the gasoline and diesel default fuel densities used within the model iterations.

Note: In 2015, EIA changed its methods for estimating AFV fuel consumption. These methodological changes included how vehicle counts are estimated, moving from estimates based on modeling to one that is based on survey data. EIA now publishes data about fuel use and number of vehicles for only four types of AFV fleets: federal government, state government, transit agencies, and fuel providers. These changes were first incorporated in this year's inventory and apply to the 1990 to 2014 time period. This resulted in large reductions in AFV VMT, thus leading to a shift in VMT to conventional on-road vehicle classes.

## References

- AAR (2008 through 2015) *Railroad Facts*. Policy and Economics Department, Association of American Railroads, Washington, D.C. Obtained from Clyde Crimmel at AAR.
- ANL (2006) Argonne National Laboratory (2006) GREET model Version 1.7. June 2006.
- ANL (2015) *The Greenhouse Gases, Regulated Emissions, and Energy Use in Transportation Model (GREET\_1\_2015)*. Argonne National Laboratory. October 2015. Available at <<https://greet.es.anl.gov>>.
- APTA (2007 through 2015) *Public Transportation Fact Book*. American Public Transportation Association, Washington, D.C. Available online at: <<http://www.apta.com/resources/statistics/Pages/transitstats.aspx>>.
- APTA (2006) *Commuter Rail National Totals*. American Public Transportation Association, Washington, D.C. Available online at: <<http://www.apta.com/research/stats/rail/crsun.cfm>>.
- Benson, D. (2002 through 2004) Unpublished data. Upper Great Plains Transportation Institute, North Dakota State University and American Short Line & Regional Railroad Association.
- Browning, L. (2015) "Methodology for Highway Vehicle Alternative Fuel GHG Estimates". Technical Memo, December 2015.
- Browning, L. (2005) Personal communication with Lou Browning, Emission control technologies for diesel highway vehicles specialist, ICF International.
- DHS (2008) Email Communication. Elissa Kay, Department of Homeland Security and Joe Aamidor, ICF International. January 11, 2008.
- DLA Energy (2015) Unpublished data from the Defense Fuels Automated Management System (DFAMS). Defense Energy Support Center, Defense Logistics Agency, U.S. Department of Defense. Washington, D.C.
- DOE (1993 through 2015) *Transportation Energy Data Book*. Office of Transportation Technologies, Center for Transportation Analysis, Energy Division, Oak Ridge National Laboratory. ORNL-6978.
- EDTA (2015) *Electric Drive Sales Dashboard*. Electric Drive Transportation Association, Washington, D.C. Available at: <<http://electricdrive.org/index.php?ht=d/sp/i/20952/pid/20952>>.
- EIA (2015) *Natural Gas Annual 2014*. Energy Information Administration, U.S. Department of Energy. Washington, D.C. DOE/EIA-0131(06).
- EIA (2016) *Monthly Energy Review, February 2016*, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2016/02).
- EIA (1991 through 2015) *Fuel Oil and Kerosene Sales*. Energy Information Administration, U.S. Department of Energy. Washington, D.C. Available online at: <<http://www.eia.gov/petroleum/fueloilkerosene/>>.
- EPA (2015d). *Motor Vehicle Emissions Simulator (Moves) 2014*. Office of Transportation and Air Quality, U.S. Environmental Protection Agency. Available online at: <<http://www.epa.gov/otaq/models/moves/index.htm>>.
- EPA (2015e) Annual Certification Test Results Report. Office of Transportation and Air Quality, U.S. Environmental Protection Agency. Available online at: <<http://www.epa.gov/otaq/crttst.htm>>.
- EPA (2015f) Confidential Engine Family Sales Data Submitted To EPA By Manufacturers. Office of Transportation and Air Quality, U.S. Environmental Protection Agency.
- EPA (2015g) "1970 - 2014 Average annual emissions, all criteria pollutants in MS Excel." *National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data*. Office of Air Quality Planning and Standards. Available online at: <<http://www.epa.gov/ttn/chieftrends/index.html>>.
- EPA (2000) *Mobile6 Vehicle Emission Modeling Software*. Office of Mobile Sources, U.S. Environmental Protection Agency, Ann Arbor, Michigan.
- EPA (1999b) Regulatory Announcement: EPA's Program for Cleaner Vehicles and Cleaner Gasoline. Office of Mobile Sources. December 1999. EPA420-F-99-051. Available online at: <<http://www.epa.gov/otaq/regs/ld-hwy/tier-2/frm/f99051.pdf>>.
- EPA (1998b) *Emissions of Nitrous Oxide from Highway Mobile Sources: Comments on the Draft Inventory of U.S. Greenhouse Gas Emissions and Sinks, 1990–1996*. Office of Mobile Sources, Assessment and Modeling Division, U.S. Environmental Protection Agency. August 1998. EPA420-R-98-009.
- FAA (2016) Personal Communication between FAA and Leif Hockstad for aviation emissions estimates from the Aviation Environmental Design Tool (AEDT). February 2016.

- FHWA (1996 through 2015) *Highway Statistics*. Federal Highway Administration, U.S. Department of Transportation, Washington, D.C. Report FHWA-PL-96-023-annual. Available online at: <<http://www.fhwa.dot.gov/policy/ohpi/hss/hsspubs.htm>>.
- Gaffney, J. (2007) Email Communication. John Gaffney, American Public Transportation Association and Joe Aamidor, ICF International. December 17, 2007.
- ICF (2006a) *Revised Gasoline Vehicle EFs for LEV and Tier 2 Emission Levels*. Memorandum from ICF International to John Davies, Office of Transportation and Air Quality, U.S. Environmental Protection Agency. November 2006.
- ICF (2006b) *Revisions to Alternative Fuel Vehicle (AFV) Emission Factors for the U.S. Greenhouse Gas Inventory*. Memorandum from ICF International to John Davies, Office of Transportation and Air Quality, U.S. Environmental Protection Agency. November 2006.
- ICF (2004) *Update of Methane and Nitrous Oxide Emission Factors for On-Highway Vehicles*. Final Report to U.S. Environmental Protection Agency. February 2004.
- Lipman, T. and M. Delucchi (2002) "Emissions of Nitrous Oxide and Methane from Conventional and Alternative Fuel Motor Vehicles." *Climate Change*, 53:477-516.
- Santoni, G., B. Lee, E. Wood, S. Herndon, R. Miake-Lye, S. Wofsy, J. McManus, D. Nelson, M. Zahniser (2011) Aircraft emissions of methane and nitrous oxide during the alternative aviation fuel experiment. *Environ Sci Technol*. 2011 Aug 15; 45(16):7075-82.
- Whorton, D. (2006 through 2014) Personal communication, Class II and III Rail energy consumption, American Short Line and Regional Railroad Association.

### 3.3. Methodology for Estimating Emissions from Commercial Aircraft Jet Fuel Consumption

**IPCC Tier 3B Method:** Commercial aircraft jet fuel burn and carbon dioxide (CO<sub>2</sub>) emissions estimates were developed by the U.S. Federal Aviation Administration (FAA) using radar-informed data from the FAA Enhanced Traffic Management System (ETMS) for 2000 through 2014 as modeled with the Aviation Environmental Design Tool (AEDT). This bottom-up approach is built from modeling dynamic aircraft performance for each flight occurring within an individual calendar year. The analysis incorporates data on the aircraft type, date, flight identifier, departure time, arrival time, departure airport, arrival airport, ground delay at each airport, and real-world flight trajectories. To generate results for a given flight within AEDT, the radar-informed aircraft data is correlated with engine and aircraft performance data to calculate fuel burn and exhaust emissions. Information on exhaust emissions for in-production aircraft engines comes from the International Civil Aviation Organization (ICAO) Aircraft Engine Emissions Databank (EDB). This bottom-up approach is in accordance with the Tier 3B method from the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006).

**International Bunkers:** The IPCC guidelines define international aviation (International Bunkers) as emissions from flights that depart from one country and arrive in a different country. Bunker fuel emissions estimates for commercial aircraft were developed for this report for 2000 through 2014 using the same radar-informed data modeled with AEDT. Since this process builds estimates from flight-specific information, the emissions estimates for commercial aircraft can include emissions associated with the U.S. Territories (i.e., American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands). However, to allow for the alignment of emissions estimates for commercial aircraft with other data that is provided without the U.S. Territories, this annex includes emissions estimates for commercial aircraft both with and without the U.S. Territories included.

**Time Series and Analysis Update:** The FAA incrementally improves the consistency, robustness, and fidelity of the CO<sub>2</sub> emissions modeling for commercial aircraft, which is the basis of the Tier 3B inventories presented in this report. While the FAA does not anticipate significant changes to the AEDT model in the future, recommended improvements are limited by budget and time constraints, as well as data availability. For instance, previous reports included reported annual CO<sub>2</sub> emission estimates for 2000 through 2005 that were modeled using the FAA's System for assessing Aviation's Global Emissions (SAGE). That tool and its capabilities were significantly improved after it was incorporated and evolved into AEDT. For this report, the AEDT model was used to generate annual CO<sub>2</sub> emission estimates for 2000, 2005, 2010, 2011, 2012, 2013 and 2014 only. The reported annual CO<sub>2</sub> emissions values for 2001 through 2004 were estimated from the previously reported SAGE data. Likewise, CO<sub>2</sub> emissions values for 2006 through 2009 were estimated by interpolation to preserve trends from past reports.

Commercial aircraft radar data sets are not available for years prior to 2000. Instead, the FAA applied a Tier3B methodology by developing Official Airline Guide (OAG) schedule-informed estimates modeled with AEDT and great circle trajectories for 1990, 2000 and 2010. The ratios between the OAG schedule-informed and the radar-informed inventories for the years 2000 and 2010 were applied to the 1990 OAG scheduled-informed inventory to generate the best possible CO<sub>2</sub> inventory estimate for commercial aircraft in 1990. The resultant 1990 CO<sub>2</sub> inventory served as the reference for generating the additional 1991-1999 emissions estimates, which were established using previously available trends.

**Notes on the 1990 CO<sub>2</sub> Emissions Inventory for Commercial Aircraft:** There are uncertainties associated with the modeled 1990 data that do not exist for the modeled 2000 to 2014 data. Radar-based data is not available for 1990. The OAG schedule information generally includes fewer carriers than radar information, and this will result in a different fleet mix, and in turn, different CO<sub>2</sub> emissions than would be quantified using a radar-based data set. For this reason, the FAA adjusted the OAG-informed schedule for 1990 with a ratio based on radar-informed information. In addition, radar trajectories are also generally longer than great circle trajectories. While the 1990 fuel burn data was adjusted to address these differences, it inherently adds greater uncertainty to the revised 1990 commercial aircraft CO<sub>2</sub> emissions as compared to data from 2000 forward. Also, the revised 1990 CO<sub>2</sub> emissions inventory now reflects only commercial aircraft jet fuel consumption, while previous reports may have aggregated jet fuel sales data from non-commercial aircraft into this category. Thus, it would be inappropriate to compare 1990 to future years for other than qualitative purposes.

The 1990 commercial aircraft CO<sub>2</sub> emissions estimate is approximately 5 percent lower than the 2014 CO<sub>2</sub> emissions estimate. It is important to note that the distance flown increased by more than 40 percent over this 25-year period

and that fuel burn and aviation activity trends over the past two decades indicate significant improvements in commercial aviation's ability to provide increased service levels while using less fuel.<sup>57</sup>

**Methane Emissions:** Contributions of methane (CH<sub>4</sub>) emissions from commercial aircraft are reported as zero. Years of scientific measurement campaigns conducted at the exhaust exit plane of commercial aircraft gas turbine engines have repeatedly indicated that CH<sub>4</sub> emissions are consumed over the full mission flight envelope (Santoni et al. 2011). As a result, the U.S. EPA published that "...methane is no longer considered to be an emission from aircraft gas turbine engines burning Jet A at higher power settings and is, in fact, consumed in net at these higher powers."<sup>58</sup> In accordance with the following statements in the 2006 IPCC Guidelines (IPCC 2006), the FAA does not calculate CH<sub>4</sub> emissions for either the domestic or international bunker commercial aircraft jet fuel emissions inventories. "*Methane (CH<sub>4</sub>) may be emitted by gas turbines during idle and by older technology engines, but recent data suggest that little or no CH<sub>4</sub> is emitted by modern engines.*" "*Current scientific understanding does not allow other gases (e.g., N<sub>2</sub>O and CH<sub>4</sub>) to be included in calculation of cruise emissions.*" (IPCC 1999).

**Results:** For each inventory calendar year the graph and table below include four jet fuel burn values. These values are comprised of domestic and international fuel burn totals for the U.S. 50 States and the U.S. 50 States + Territories. Data are presented for domestic defined as jet fuel burn from any commercial aircraft flight departing and landing in the U.S. 50 States and for the U.S. 50 States + Territories. The data presented as international is respective of the two different domestic definitions, and represents flights departing from the specified domestic area and landing anywhere in the world outside of that area.

Note that the graph and table present less fuel burn for the international U.S. 50 States + Territories than for the international U.S. 50 States. This is because the flights between the 50 states and U.S. Territories are "international" when only the 50 states are defined as domestic, but they are "domestic" for the U.S. 50 States + Territories definition.


---

<sup>57</sup> Additional information on the AEDT modeling process is available at:

[http://www.faa.gov/about/office\\_org/headquarters\\_offices/apl/research/models/](http://www.faa.gov/about/office_org/headquarters_offices/apl/research/models/)

<sup>58</sup> Recommended Best Practice for Quantifying Speciated Organic Gas Emissions from Aircraft Equipped with Turbofan, Turbojet and Turboprop Engines, EPA-420-R-09-901, May 27, 2009. See <<http://www.epa.gov/otaq/aviation.html>>.

**Figure A-5: Commercial Aviation Fuel Burn for the United States and Territories**



Note: Hollow markers are estimates from data generated by prior tools and methods. 1990 is estimated using non-radar methods.

**Table A-119: Commercial Aviation Fuel Burn for the United States and Territories**

| Year  | Region  | Distance Flown (nmi) | Fuel Burn (M Gallon) | Fuel Burn (Tbtu) | Fuel Burn (Kg) | CO <sub>2</sub> (MMT) |
|-------|---|----------------------|----------------------|------------------|----------------|-----------------------|
| 1990  | Domestic U.S. 50 States and U.S. Territories      | 4,057,195,988        | 11,568               | 1,562            | 34,820,800,463 | 109.9                 |
|       | International U.S. 50 States and U.S. Territories | 599,486,893          | 3,155                | 426              | 9,497,397,919  | 30.0                  |
|       | Domestic U.S. 50 States                           | 3,984,482,217        | 11,287               | 1,524            | 33,972,832,399 | 107.2                 |
|       | International U.S. 50 States                      | 617,671,849          | 3,228                | 436              | 9,714,974,766  | 30.7                  |
| 1995* | Domestic U.S. 50 States and U.S. Territories      | N/A                  | 12,136               | 1,638            | 36,528,990,675 | 115.2                 |
| 1996* | Domestic U.S. 50 States and U.S. Territories      | N/A                  | 12,492               | 1,686            | 37,600,624,534 | 118.6                 |
| 1997* | Domestic U.S. 50 States and U.S. Territories      | N/A                  | 12,937               | 1,747            | 38,940,896,854 | 122.9                 |
| 1998* | Domestic U.S. 50 States and U.S. Territories      | N/A                  | 12,601               | 1,701            | 37,930,582,643 | 119.7                 |
| 1999* | Domestic U.S. 50 States and U.S. Territories      | N/A                  | 13,726               | 1,853            | 41,314,843,250 | 130.3                 |
| 2000  | Domestic U.S. 50 States and U.S. Territories      | 5,994,679,944        | 14,672               | 1,981            | 44,161,841,348 | 139.3                 |
|       | International U.S. 50 States and U.S. Territories | 1,309,565,963        | 6,040                | 815              | 18,181,535,058 | 57.4                  |
|       | Domestic U.S. 50 States                           | 5,891,481,028        | 14,349               | 1,937            | 43,191,000,202 | 136.3                 |
|       | International U.S. 50 States                      | 1,331,784,289        | 6,117                | 826              | 18,412,169,613 | 58.1                  |
| 2001* | Domestic U.S. 50 States and U.S. Territories      | 5,360,977,447        | 13,121               | 1,771            | 39,493,457,147 | 124.6                 |
|       | International U.S. 50 States and U.S. Territories | 1,171,130,679        | 5,402                | 729              | 16,259,550,186 | 51.3                  |
|       | Domestic U.S. 50 States                           | 5,268,687,772        | 12,832               | 1,732            | 38,625,244,409 | 121.9                 |
|       | International U.S. 50 States                      | 1,191,000,288        | 5,470                | 739              | 16,465,804,174 | 51.9                  |
| 2002* | Domestic U.S. 50 States and U.S. Territories      | 5,219,345,344        | 12,774               | 1,725            | 38,450,076,259 | 121.3                 |
|       | International U.S. 50 States and U.S. Territories | 1,140,190,481        | 5,259                | 710              | 15,829,987,794 | 49.9                  |

|       |   |               |        |       |                |       |
|-------|---|---------------|--------|-------|----------------|-------|
|       | Domestic U.S. 50 States                           | 5,129,493,877 | 12,493 | 1,687 | 37,604,800,905 | 118.6 |
|       | International U.S. 50 States                      | 1,159,535,153 | 5,326  | 719   | 16,030,792,741 | 50.6  |
| 2003* | Domestic U.S. 50 States and U.S. Territories      | 5,288,138,079 | 12,942 | 1,747 | 38,956,861,262 | 122.9 |
|       | International U.S. 50 States and U.S. Territories | 1,155,218,577 | 5,328  | 719   | 16,038,632,384 | 50.6  |
|       | Domestic U.S. 50 States                           | 5,197,102,340 | 12,658 | 1,709 | 38,100,444,893 | 120.2 |
|       | International U.S. 50 States                      | 1,174,818,219 | 5,396  | 728   | 16,242,084,008 | 51.2  |
| 2004* | Domestic U.S. 50 States and U.S. Territories      | 5,371,498,689 | 13,146 | 1,775 | 39,570,965,441 | 124.8 |
|       | International U.S. 50 States and U.S. Territories | 1,173,429,093 | 5,412  | 731   | 16,291,460,535 | 51.4  |
|       | Domestic U.S. 50 States                           | 5,279,027,890 | 12,857 | 1,736 | 38,701,048,784 | 122.1 |
|       | International U.S. 50 States                      | 1,193,337,698 | 5,481  | 740   | 16,498,119,309 | 52.1  |
| 2005  | Domestic U.S. 50 States and U.S. Territories      | 6,476,007,697 | 13,976 | 1,887 | 42,067,562,737 | 132.7 |
|       | International U.S. 50 States and U.S. Territories | 1,373,543,928 | 5,858  | 791   | 17,633,508,081 | 55.6  |
|       | Domestic U.S. 50 States                           | 6,370,544,998 | 13,654 | 1,843 | 41,098,359,387 | 129.7 |
|       | International U.S. 50 States                      | 1,397,051,323 | 5,936  | 801   | 17,868,972,965 | 56.4  |
| 2006* | Domestic U.S. 50 States and U.S. Territories      | 5,894,323,482 | 14,426 | 1,948 | 43,422,531,461 | 137.0 |
|       | International U.S. 50 States and U.S. Territories | 1,287,642,623 | 5,939  | 802   | 17,877,159,421 | 56.4  |
|       | Domestic U.S. 50 States                           | 5,792,852,211 | 14,109 | 1,905 | 42,467,943,091 | 134.0 |
|       | International U.S. 50 States                      | 1,309,488,994 | 6,015  | 812   | 18,103,932,940 | 57.1  |
| 2007* | Domestic U.S. 50 States and U.S. Territories      | 6,009,247,818 | 14,707 | 1,986 | 44,269,160,525 | 139.7 |
|       | International U.S. 50 States and U.S. Territories | 1,312,748,383 | 6,055  | 817   | 18,225,718,619 | 57.5  |
|       | Domestic U.S. 50 States                           | 5,905,798,114 | 14,384 | 1,942 | 43,295,960,105 | 136.6 |
|       | International U.S. 50 States                      | 1,335,020,703 | 6,132  | 828   | 18,456,913,646 | 58.2  |
| 2008* | Domestic U.S. 50 States and U.S. Territories      | 5,475,092,456 | 13,400 | 1,809 | 40,334,124,033 | 127.3 |
|       | International U.S. 50 States and U.S. Territories | 1,196,059,638 | 5,517  | 745   | 16,605,654,741 | 52.4  |
|       | Domestic U.S. 50 States                           | 5,380,838,282 | 13,105 | 1,769 | 39,447,430,318 | 124.5 |
|       | International U.S. 50 States                      | 1,216,352,196 | 5,587  | 754   | 16,816,299,099 | 53.1  |
| 2009* | Domestic U.S. 50 States and U.S. Territories      | 5,143,268,671 | 12,588 | 1,699 | 37,889,631,668 | 119.5 |
|       | International U.S. 50 States and U.S. Territories | 1,123,571,175 | 5,182  | 700   | 15,599,251,424 | 49.2  |
|       | Domestic U.S. 50 States                           | 5,054,726,871 | 12,311 | 1,662 | 37,056,676,966 | 116.9 |
|       | International U.S. 50 States                      | 1,142,633,881 | 5,248  | 709   | 15,797,129,457 | 49.8  |
| 2010  | Domestic U.S. 50 States and U.S. Territories      | 5,652,264,576 | 11,931 | 1,611 | 35,912,723,830 | 113.3 |
|       | International U.S. 50 States and U.S. Territories | 1,474,839,733 | 6,044  | 816   | 18,192,953,916 | 57.4  |
|       | Domestic U.S. 50 States                           | 5,554,043,585 | 11,667 | 1,575 | 35,116,863,245 | 110.8 |
|       | International U.S. 50 States                      | 1,497,606,695 | 6,113  | 825   | 18,398,996,825 | 58.0  |
| 2011  | Domestic U.S. 50 States and U.S. Territories      | 5,767,378,664 | 12,067 | 1,629 | 36,321,170,730 | 114.6 |
|       | International U.S. 50 States and U.S. Territories | 1,576,982,962 | 6,496  | 877   | 19,551,631,939 | 61.7  |
|       | Domestic U.S. 50 States                           | 5,673,689,481 | 11,823 | 1,596 | 35,588,754,827 | 112.3 |
|       | International U.S. 50 States                      | 1,596,797,398 | 6,554  | 885   | 19,727,043,614 | 62.2  |
| 2012  | Domestic U.S. 50 States and U.S. Territories      | 5,735,605,432 | 11,932 | 1,611 | 35,915,745,616 | 113.3 |
|       | International U.S. 50 States and U.S. Territories | 1,619,012,587 | 6,464  | 873   | 19,457,378,739 | 61.4  |
|       | Domestic U.S. 50 States                           | 5,636,910,529 | 11,672 | 1,576 | 35,132,961,140 | 110.8 |
|       | International U.S. 50 States                      | 1,637,917,110 | 6,507  | 879   | 19,587,140,347 | 61.8  |
| 2013  | Domestic U.S. 50 States and U.S. Territories      | 5,808,034,123 | 12,031 | 1,624 | 36,212,974,471 | 114.3 |
|       | International U.S. 50 States and U.S. Territories | 1,641,151,400 | 6,611  | 892   | 19,898,871,458 | 62.8  |
|       | Domestic U.S. 50 States                           | 5,708,807,315 | 11,780 | 1,590 | 35,458,690,595 | 111.9 |
|       | International U.S. 50 States                      | 1,661,167,498 | 6,657  | 899   | 20,036,865,038 | 63.2  |
| 2014  | Domestic U.S. 50 States and U.S. Territories      | 5,825,999,388 | 12,131 | 1,638 | 36,514,970,659 | 115.2 |
|       | International U.S. 50 States and U.S. Territories | 1,724,559,209 | 6,980  | 942   | 21,008,818,741 | 66.3  |
|       | Domestic U.S. 50 States                           | 5,725,819,482 | 11,882 | 1,604 | 35,764,791,774 | 112.8 |
|       | International U.S. 50 States                      | 1,745,315,059 | 7,027  | 949   | 21,152,418,387 | 66.7  |

\*Estimates for these years were derived from previously reported tools and methods

### 3.4. Methodology for Estimating CH<sub>4</sub> Emissions from Coal Mining

The methodology for estimating CH<sub>4</sub> emissions from coal mining consists of two steps:

- **Estimate emissions from underground mines.** These emissions have two sources: ventilation systems and degasification systems. They are estimated on a mine-by-mine basis, then summed to determine total CH<sub>4</sub> liberated. The CH<sub>4</sub> recovered and used is then subtracted from this total, resulting in an estimate of net emissions to the atmosphere.
- **Estimate CH<sub>4</sub> emissions from surface mines and post-mining activities.** This step does not use mine-specific data; rather, it consists of multiplying coal-basin-specific coal production by coal-basin-specific gas content and an emission factor.

#### Step 1: Estimate CH<sub>4</sub> Liberated and CH<sub>4</sub> Emitted from Underground Mines

Underground mines generate CH<sub>4</sub> from ventilation systems and from degasification systems. Some mines recover and use the generated CH<sub>4</sub>, thereby reducing emissions to the atmosphere. Total CH<sub>4</sub> emitted from underground mines equals the CH<sub>4</sub> liberated from ventilation systems, plus the CH<sub>4</sub> liberated from degasification systems, minus CH<sub>4</sub> recovered and used.

##### Step 1.1: Estimate CH<sub>4</sub> Liberated from Ventilation Systems

All coal mines with detectable CH<sub>4</sub> emissions use ventilation systems to ensure that CH<sub>4</sub> levels remain within safe concentrations. Many coal mines do not have detectable levels of CH<sub>4</sub>; others emit several million cubic feet per day (MMCFD) from their ventilation systems. On a quarterly basis, the U.S. Mine Safety and Health Administration (MSHA) measures CH<sub>4</sub> emissions levels at underground mines. MSHA maintains a database of measurement data from all underground mines with detectable levels of CH<sub>4</sub> in their ventilation air (MSHA 2015).<sup>59</sup> Based on the four quarterly measurements, MSHA estimates average daily CH<sub>4</sub> liberated at each of these underground mines.

For 1990 through 1999, average daily CH<sub>4</sub> emissions from MSHA were multiplied by the number of days in the year (i.e., coal mine assumed in operation for all four quarters) to determine the annual emissions for each mine. For 2000 through 2014, the average daily CH<sub>4</sub> emissions were multiplied by the number of days corresponding to the number of quarters the mine vent was operating. For example, if the mine vent was operational in one out of the four quarters, the average daily CH<sub>4</sub> emissions were multiplied by 92 days. Total ventilation emissions for a particular year were estimated by summing emissions from individual mines.

Since 2011, the nation's "gassiest" underground coal mines—those that liberate more than 36,500,000 actual cubic feet of CH<sub>4</sub> per year (about 14,700 MT CO<sub>2</sub> eq.)—have been required to report to the EPA's GHGRP (EPA 2015).<sup>60</sup> Mines that report to the GHGRP must report quarterly measurements of CH<sub>4</sub> emissions from ventilation systems to EPA; they have the option of recording their own measurements, or using the measurements taken by MSHA as part of that agency's quarterly safety inspections of all mines in the U.S. with detectable CH<sub>4</sub> concentrations.<sup>61</sup>

Since 2013, ventilation emission estimates have been calculated based on both GHGRP data submitted by underground mines that recorded their own measurements, and on quarterly measurement data obtained directly from MSHA for the remaining mines (not MSHA data reported by the mines to the GHGRP).<sup>62</sup> The quarterly measurements are used to determine the average daily emissions rate for the reporting year quarter. The CH<sub>4</sub> liberated from ventilation systems was estimated by summing the emissions from the GHGRP self-reported mines and emissions based on MSHA quarterly measurements for the remaining mines.

---

<sup>59</sup> MSHA records coal mine methane readings with concentrations of greater than 50 ppm (parts per million) methane. Readings below this threshold are considered non-detectable.

<sup>60</sup> Underground coal mines report to EPA under Subpart FF of the GHGRP. In 2014, 128 underground coal mines reported to the program.

<sup>61</sup> MSHA records coal mine CH<sub>4</sub> readings with concentrations of greater than 50 ppm (parts per million) CH<sub>4</sub>. Readings below this threshold are considered non-detectable.

<sup>62</sup> EPA has determined that certain mines are having difficulty interpreting the MSHA data so that they report them correctly to the GHGRP. EPA is working with these mines to correct their GHGRP reports, and in the meantime is relying on data obtained directly from MSHA for purposes of the national inventory.

**Table A-120: Mine-Specific Data Used to Estimate Ventilation Emissions**

| Year | Individual Mine Data Used  |
|------|--|
| 1990 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 1991 | 1990 Emissions Factors Used Instead of Mine-Specific Data  |
| 1992 | 1990 Emissions Factors Used Instead of Mine-Specific Data  |
| 1993 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 1994 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 1995 | All Mines Emitting at Least 0.5 MMCFD (Assumed to Account for 94.1% of Total)*                     |
| 1996 | All Mines Emitting at Least 0.5 MMCFD (Assumed to Account for 94.1% of Total)*                     |
| 1997 | All Mines with Detectable Emissions (Assumed to Account for 100% of Total)                         |
| 1998 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 1999 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2000 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2001 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2002 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2003 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2004 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2005 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2006 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 97.8% of Total)*                     |
| 2007 | All Mines with Detectable Emissions (Assumed to Account for 100% of Total)                         |
| 2008 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 98.96% of Total)**                   |
| 2009 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 98.96% of Total)**                   |
| 2010 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 98.96% of Total)**                   |
| 2011 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 98.96% of Total)**                   |
| 2012 | All Mines Emitting at Least 0.1 MMCFD (Assumed to Account for 98.96% of Total)**                   |
| 2013 | All Mines with Detectable Emissions and GHGRP reported data (Assumed to account for 100% of Total) |
| 2014 | All Mines with Detectable Emissions and GHGRP reported data (Assumed to account for 100% of Total) |

\* Factor derived from a complete set of individual mine data collected for 1997.

\*\* Factor derived from a complete set of individual mine data collected for 2007.

### Step 1.2: Estimate CH<sub>4</sub> Liberated from Degasification Systems

Coal mines use several types of degasification systems to remove CH<sub>4</sub>, including pre-mining vertical and horizontal wells (to recover CH<sub>4</sub> before mining) and post-mining vertical wells and horizontal boreholes (to recover CH<sub>4</sub> during mining of the coal seam). Post-mining gob wells and cross-measure boreholes recover CH<sub>4</sub> from the overburden (i.e., gob area) after mining of the seam (primarily in longwall mines).

Twenty-five mines employed degasification systems in 2014, and the CH<sub>4</sub> liberated through these systems was reported to the EPA's GHGRP (EPA 2015). Sixteen of these mines reported CH<sub>4</sub> recovery and use projects, and the other nine reported emitting CH<sub>4</sub> from degasification systems to the atmosphere. Several of the mines venting CH<sub>4</sub> from degasification systems use a small portion of the gas to fuel gob well blowers or compressors in remote locations where electricity is not available. However, this CH<sub>4</sub> use is not considered to be a formal recovery and use project.

Degasification information reported to the GHGRP by underground coal mines is the primary source of data used to develop estimates of CH<sub>4</sub> liberated from degasification systems. Data reported to the GHGRP were used to estimate CH<sub>4</sub> liberated from degasification systems at 20 of the 25 mines that used degasification systems in 2014.

Degasification volumes for the life of any pre-mining wells are attributed to the mine as emissions in the year in which the well is mined through.<sup>63</sup> The GHGRP does not require gas production from virgin coal seams (coalbed methane) to be reported by coal mines under subpart FF. Most pre-mining wells drilled from the surface are considered coalbed methane wells and are reported under another subpart of the program (subpart W, "Petroleum and Natural Gas Systems"). As a result, for the 10 mines with degasification systems that include pre-mining wells, GHGRP information was supplemented with historical data from state gas well production databases (GSA 2016; WVGES 2015), as well as with mine-specific information regarding the dates on which pre-mining wells are mined through (JWR 2010; El Paso 2009). For pre-mining wells, the cumulative CH<sub>4</sub> production from the well is totaled using gas sales data, and considered liberated from the mine's degasification system the year in which the well is mined through.

<sup>63</sup> A well is "mined through" when coal mining development or the working face intersects the borehole or well.

EPA's GHGRP reports with CH<sub>4</sub> liberated from degasification systems are reviewed for errors in reporting. For some mines, GHGRP data are corrected for the inventory based on expert judgment. Common errors include reporting CH<sub>4</sub> liberated as CH<sub>4</sub> destroyed and vice versa. Other errors include reporting CH<sub>4</sub> destroyed without reporting any CH<sub>4</sub> liberated by degasification systems. In the rare cases where GHGRP data are inaccurate and gas sales data unavailable, estimates of CH<sub>4</sub> liberated are based on historical CH<sub>4</sub> liberation rates.

### **Step 1.3: Estimate CH<sub>4</sub> Recovered from Ventilation and Degasification Systems, and Utilized or Destroyed (Emissions Avoided)**

Of the 16 active coal mines with operational CH<sub>4</sub> recovery and use projects in 2014, 14 sold the recovered CH<sub>4</sub> to a pipeline, including one that also used CH<sub>4</sub> to fuel a thermal coal dryer. Uses at other mines include electrical power generation (one mine) and heating mine ventilation air (one mine).

Ten of the 16 mines deployed degasification systems in 2014; for those mines, estimates of CH<sub>4</sub> recovered from the systems were exclusively based on GHGRP data. Based on weekly measurements of gas flow and CH<sub>4</sub> concentrations, the GHGRP summary data for degasification destruction at each mine were added together to estimate the CH<sub>4</sub> recovered and used from degasification systems.

Of the 10 mines with degasification systems in 2014, four intersected pre-mining wells in 2014. GHGRP and supplemental data were used to estimate CH<sub>4</sub> recovered and used at two of these mines, while supplemental data alone were used at the other two mines, that reported as a single entity to the GHGRP. Supplemental information was used for these four mines because estimating CH<sub>4</sub> recovery and use from pre-mining wells requires additional data (not reported under subpart FF of the GHGRP; see discussion in step 1.2 above) to account for the emissions avoided. The supplemental data came from state gas production databases (GSA 2016; WVGES 2015), as well as mine-specific information on the timing of mined-through pre-mining wells (JWR 2010; El Paso 2009). For pre-mining wells, the cumulative CH<sub>4</sub> production from the wells was totaled using gas sales data, and considered to be CH<sub>4</sub> recovered and used from the mine's degasification system the year in which the well is mined through.

EPA's GHGRP reports with CH<sub>4</sub> recovered and used from degasification systems are reviewed for errors in reporting. For some mines, GHGRP data are corrected for the inventory based on expert judgment (see further discussion in Step 1.2). In 2014, GHGRP information was not used to estimate CH<sub>4</sub> recovered and used at two mines because of a lack of mine-provided information used in prior years and GHGRP reporting discrepancies.

In 2014, one mine destroyed a portion of its CH<sub>4</sub> emissions from ventilation systems using thermal oxidation technology. The amount of CH<sub>4</sub> recovered and destroyed by the project was determined through publicly available emission reduction project information (CAR 2015).

## **Step 2: Estimate CH<sub>4</sub> Emitted from Surface Mines and Post-Mining Activities**

Mine-specific data were not available for estimating CH<sub>4</sub> emissions from surface coal mines or for post-mining activities. For surface mines, basin-specific coal production obtained from the Energy Information Administration's *Annual Coal Report* was multiplied by basin-specific gas contents and a 150 percent emission factor (to account for CH<sub>4</sub> from over- and under-burden) to estimate CH<sub>4</sub> emissions (see King 1994; Saghafi 2013). For post-mining activities, basin-specific coal production was multiplied by basin-specific gas contents and a mid-range 32.5 percent emission factor accounting for CH<sub>4</sub> desorption during coal transportation and storage (Creedy 1993). Basin-specific *in situ* gas content data were compiled from AAPG (1984) and USBM (1986). Beginning in 2006, revised data on *in situ* CH<sub>4</sub> content and emissions factors have been used (EPA 1996, 2005).

### **Step 2.1: Define the Geographic Resolution of the Analysis and Collect Coal Production Data**

The first step in estimating CH<sub>4</sub> emissions from surface mining and post-mining activities was to define the geographic resolution of the analysis and to collect coal production data at that level of resolution. The analysis was conducted by coal basin as defined in Table A-121, which presents coal basin definitions by basin and by state.

The Energy Information Administration's *Annual Coal Report* (EIA 2015) includes state- and county-specific underground and surface coal production by year. To calculate production by basin, the state level data were grouped into coal basins using the basin definitions listed in Table A-121. For two states—West Virginia and Kentucky—county-level production data were used for the basin assignments because coal production occurred in geologically distinct coal basins within these states. Table A-122 presents the coal production data aggregated by basin.

### Step 2.2: Estimate Emissions Factors for Each Emissions Type

Emission factors for surface-mined coal were developed from the *in situ* CH<sub>4</sub> content of the surface coal in each basin. Based on analyses conducted in Canada and Australia on coals similar to those present in the U.S. (King 1994; Saghafi 2013), the surface mining emission factor used was conservatively estimated to be 150 percent of the *in situ* CH<sub>4</sub> content of the basin. Furthermore, the post-mining emission factors used were estimated to be 25 to 40 percent of the average *in situ* CH<sub>4</sub> content in the basin. For this analysis, the post-mining emission factor was determined to be 32.5 percent of the *in situ* CH<sub>4</sub> content in the basin. Table A-123 presents the average *in situ* content for each basin, along with the resulting emission factor estimates.

### Step 2.3: Estimate CH<sub>4</sub> Emitted

The total amount of CH<sub>4</sub> emitted from surface mines and post-mining activities was calculated by multiplying the coal production in each basin by the appropriate emission factors.

Table A-121 lists each of the major coal mine basins in the United States and the states in which they are located. As shown in Figure A-6, several coal basins span several states. Table A-122 shows annual underground, surface, and total coal production (in short tons) for each coal basin. Table A-123 shows the surface, post-surface, and post-underground emission factors used for estimating CH<sub>4</sub> emissions for each of the categories. Table A-124 presents annual estimates of CH<sub>4</sub> emissions for ventilation and degasification systems, and CH<sub>4</sub> used and emitted by underground coal mines. Table A-125 presents annual estimates of total CH<sub>4</sub> emissions from underground, post-underground, surface, and post-surface activities. Table A-126 provides the total net CH<sub>4</sub> emissions by state.


**Table A-121: Coal Basin Definitions by Basin and by State**

| Basin                        | States   |
|------------------------------|--|
| Northern Appalachian Basin   | Maryland, Ohio, Pennsylvania, West Virginia North            |
| Central Appalachian Basin    | Kentucky East, Tennessee, Virginia, West Virginia South      |
| Warrior Basin                | Alabama, Mississippi   |
| Illinois Basin               | Illinois, Indiana, Kentucky West                             |
| South West and Rockies Basin | Arizona, California, Colorado, New Mexico, Utah              |
| North Great Plains Basin     | Montana, North Dakota, Wyoming                               |
| West Interior Basin          | Arkansas, Iowa, Kansas, Louisiana, Missouri, Oklahoma, Texas |
| Northwest Basin              | Alaska, Washington   |
| State                        | Basin  |
| Alabama                      | Warrior Basin  |
| Alaska                       | Northwest Basin  |
| Arizona                      | South West and Rockies Basin                                 |
| Arkansas                     | West Interior Basin  |
| California                   | South West and Rockies Basin                                 |
| Colorado                     | South West and Rockies Basin                                 |
| Illinois                     | Illinois Basin   |
| Indiana                      | Illinois Basin   |
| Iowa                         | West Interior Basin  |
| Kansas                       | West Interior Basin  |
| Kentucky (east)              | Central Appalachian Basin                                    |
| Kentucky (west)              | Illinois Basin   |
| Louisiana                    | West Interior Basin  |
| Maryland                     | Northern Appalachian Basin                                   |
| Mississippi                  | Warrior Basin  |
| Missouri                     | West Interior Basin  |
| Montana                      | North Great Plains Basin                                     |
| New Mexico                   | South West and Rockies Basin                                 |
| North Dakota                 | North Great Plains Basin                                     |
| Ohio                         | Northern Appalachian Basin                                   |
| Oklahoma                     | West Interior Basin  |
| Pennsylvania                 | Northern Appalachian Basin                                   |
| Tennessee                    | Central Appalachian Basin                                    |
| Texas                        | West Interior Basin  |
| Utah                         | South West and Rockies Basin                                 |
| Virginia                     | Central Appalachian Basin                                    |
| Washington                   | Northwest Basin  |
| West Virginia South          | Central Appalachian Basin                                    |

West Virginia North  
Wyoming

Northern Appalachian Basin  
North Great Plains Basin

**Figure A-6: Locations of U.S. Coal Basins**



**Table A-122: Annual Coal Production (Thousand Short Tons)**

| Basin                              | 1990             | 2005             | 2008             | 2009             | 2010             | 2011             | 2012             | 2013           | 2014           |
|------------------------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|----------------|----------------|
| <b>Underground Coal Production</b> | <b>423,556</b>   | <b>368,611</b>   | <b>357,074</b>   | <b>332,061</b>   | <b>337,155</b>   | <b>345,607</b>   | <b>342,387</b>   | <b>341,216</b> | <b>354,705</b> |
| N. Appalachia                      | 103,865          | 111,151          | 105,228          | 99,629           | 103,109          | 105,752          | 103,408          | 104,198        | 116,700        |
| Cent. Appalachia                   | 198,412          | 123,083          | 114,998          | 98,689           | 96,354           | 94,034           | 78,067           | 70,440         | 64,219         |
| Warrior                            | 17,531           | 13,295           | 12,281           | 11,505           | 12,513           | 10,879           | 12,570           | 13,391         | 12,516         |
| Illinois                           | 69,167           | 59,180           | 64,609           | 67,186           | 72,178           | 81,089           | 92,500           | 98,331         | 105,211        |
| S. West/Rockies                    | 32,754           | 60,865           | 55,781           | 50,416           | 44,368           | 45,139           | 45,052           | 41,232         | 44,302         |
| N. Great Plains                    | 1,722            | 572              | 3,669            | 4,248            | 8,208            | 8,179            | 10,345           | 13,126         | 11,272         |
| West Interior                      | 105              | 465              | 508              | 388              | 425              | 535              | 445              | 498            | 485            |
| Northwest                          | 0                | 0                | 0                | 0                | 0                | 0                | 0                | 0              | 0              |
| <b>Surface Coal Production</b>     | <b>602,753</b>   | <b>762,191</b>   | <b>813,321</b>   | <b>740,175</b>   | <b>764,709</b>   | <b>754,871</b>   | <b>672,748</b>   | <b>640,740</b> | <b>643,721</b> |
| N. Appalachia                      | 60,761           | 28,873           | 30,413           | 26,552           | 26,082           | 26,382           | 21,411           | 19,339         | 17,300         |
| Cent. Appalachia                   | 94,343           | 112,222          | 118,962          | 97,778           | 89,788           | 90,778           | 69,721           | 57,173         | 52,399         |
| Warrior                            | 11,413           | 11,599           | 11,172           | 10,731           | 11,406           | 10,939           | 9,705            | 8,695          | 7,584          |
| Illinois                           | 72,000           | 33,702           | 34,266           | 34,837           | 32,911           | 34,943           | 34,771           | 33,798         | 31,969         |
| S. West/Rockies                    | 43,863           | 42,756           | 34,283           | 32,167           | 28,889           | 31,432           | 30,475           | 28,968         | 27,564         |
| N. Great Plains                    | 249,356          | 474,056          | 538,387          | 496,290          | 507,995          | 502,734          | 455,320          | 444,740        | 458,112        |
| West Interior                      | 64,310           | 52,263           | 44,361           | 39,960           | 46,136           | 55,514           | 49,293           | 46,477         | 47,201         |
| Northwest                          | 6,707            | 6,720            | 1,477            | 1,860            | 2,151            | 2,149            | 2,052            | 1,550          | 1,502          |
| <b>Total Coal Production</b>       | <b>1,026,309</b> | <b>1,130,802</b> | <b>1,170,395</b> | <b>1,072,236</b> | <b>1,101,864</b> | <b>1,100,478</b> | <b>1,015,135</b> | <b>981,956</b> | <b>998,426</b> |
| N. Appalachia                      | 164,626          | 140,024          | 135,641          | 126,181          | 129,191          | 132,134          | 124,819          | 123,537        | 134,000        |
| Cent. Appalachia                   | 292,755          | 235,305          | 233,960          | 196,467          | 186,142          | 184,812          | 147,788          | 127,613        | 116,618        |
| Warrior                            | 28,944           | 24,894           | 23,453           | 22,236           | 23,919           | 21,818           | 22,275           | 22,086         | 20,100         |
| Illinois                           | 141,167          | 92,882           | 98,875           | 102,023          | 105,089          | 116,032          | 127,271          | 132,129        | 137,180        |
| S. West/Rockies                    | 76,617           | 103,621          | 90,064           | 82,583           | 73,257           | 76,571           | 75,527           | 70,200         | 71,956         |
| N. Great Plains                    | 251,078          | 474,628          | 542,056          | 500,538          | 516,203          | 510,913          | 465,665          | 457,866        | 469,384        |
| West Interior                      | 64,415           | 52,728           | 44,869           | 40,348           | 46,561           | 56,049           | 49,738           | 46,975         | 47,686         |
| Northwest                          | 6,707            | 6,720            | 1,477            | 1,860            | 2,151            | 2,149            | 2,052            | 1,550          | 1,502          |

Source for 1990–2014 data: EIA (1990 through 2014), *Annual Coal Report*. Table 1. U.S. Department of Energy. Source for 2014 data: spreadsheet for the 2014 *Annual Coal Report*.

Note: Totals may not sum due to independent rounding.

**Table A-123: Coal Underground, Surface, and Post-Mining CH<sub>4</sub> Emission Factors (ft<sup>3</sup> per Short Ton)**

| Basin  | Surface Average<br>In Situ Content | Underground Average<br>In Situ Content | Surface Mine<br>Factors | Post-Mining<br>Surface Factors | Post Mining<br>Underground |
|--|------------------------------------|--|-------------------------|--------------------------------|----------------------------|
| Northern Appalachia                          | 59.5                               | 138.4                                  | 89.3                    | 19.3                           | 45.0                       |
| Central Appalachia (WV)                      | 24.9                               | 136.8                                  | 37.4                    | 8.1                            | 44.5                       |
| Central Appalachia (VA)                      | 24.9                               | 399.1                                  | 37.4                    | 8.1                            | 129.7                      |
| Central Appalachia (E KY)                    | 24.9                               | 61.4                                   | 37.4                    | 8.1                            | 20.0                       |
| Warrior                                      | 30.7                               | 266.7                                  | 46.1                    | 10.0                           | 86.7                       |
| Illinois                                     | 34.3                               | 64.3                                   | 51.5                    | 11.1                           | 20.9                       |
| Rockies (Piceance Basin)                     | 33.1                               | 196.4                                  | 49.7                    | 10.8                           | 63.8                       |
| Rockies (Uinta Basin)                        | 16.0                               | 99.4                                   | 24.0                    | 5.2                            | 32.3                       |
| Rockies (San Juan Basin)                     | 7.3                                | 104.8                                  | 11.0                    | 2.4                            | 34.1                       |
| Rockies (Green River Basin)                  | 33.1                               | 247.2                                  | 49.7                    | 10.8                           | 80.3                       |
| Rockies (Raton Basin)                        | 33.1                               | 127.9                                  | 49.7                    | 10.8                           | 41.6                       |
| N. Great Plains (WY, MT)                     | 20.0                               | 15.8                                   | 30.0                    | 6.5                            | 5.1                        |
| N. Great Plains (ND)                         | 5.6                                | 15.8                                   | 8.4                     | 1.8                            | 5.1                        |
| West Interior (Forest City, Cherokee Basins) | 34.3                               | 64.3                                   | 51.5                    | 11.1                           | 20.9                       |
| West Interior (Arkoma Basin)                 | 74.5                               | 331.2                                  | 111.8                   | 24.2                           | 107.6                      |
| West Interior (Gulf Coast Basin)             | 11.0                               | 127.9                                  | 16.5                    | 3.6                            | 41.6                       |
| Northwest (AK)                               | 16.0                               | 160.0                                  | 24.0                    | 1.8                            | 52.0                       |
| Northwest (WA)                               | 16.0                               | 47.3                                   | 24.0                    | 5.2                            | 15.4                       |

Sources: 1986 USBM Circular 9067, *Results of the Direct Method Determination of the Gas Contents of U.S. Coal Basins*; U.S. DOE Report DOE/METC/83-76, *Methane Recovery from Coalbeds: A Potential Energy Source*; 1986–1988 Gas Research Institute Topical Report, *A Geologic Assessment of Natural Gas from Coal Seams*; 2005 U.S. EPA Draft Report, *Surface Mines Emissions Assessment*.

**Table A-124: Underground Coal Mining CH<sub>4</sub> Emissions (Billion Cubic Feet)**

| Activity                         | 1990       | 2005      | 2008       | 2009       | 2010       | 2011       | 2012      | 2013      | 2014       |
|----------------------------------|------------|-----------|------------|------------|------------|------------|-----------|-----------|------------|
| Ventilation Output               | 112        | 75        | 100        | 114        | 117        | 97         | 90        | 89        | 89         |
| Adjustment Factor for Mine Data* | 98%        | 98%       | 99%        | 99%        | 99%        | 99%        | 99%       | 100%      | 100%       |
| Adjusted Ventilation Output      | 114        | 77        | 101        | 115        | 118        | 98         | 91        | 89        | 89         |
| Degasification System Liberated  | 54         | 48        | 49         | 49         | 58         | 48         | 45        | 48        | 48         |
| Total Underground Liberated      | 168        | 124       | 150        | 163        | 177        | 147        | 137       | 137       | 136        |
| Recovered & Used                 | (14)       | (37)      | (40)       | (40)       | (49)       | (42)       | (38)      | (41)      | (34)       |
| <b>Total</b>                     | <b>154</b> | <b>87</b> | <b>110</b> | <b>123</b> | <b>128</b> | <b>104</b> | <b>98</b> | <b>96</b> | <b>102</b> |

\* Refer to Table A-120.

Note: Totals may not sum due to independent rounding.

**Table A-125: Total Coal Mining CH<sub>4</sub> Emissions (Billion Cubic Feet)**

| Activity                     | 1990       | 2005       | 2008       | 2009       | 2010       | 2011       | 2012       | 2013       | 2014       |
|------------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Underground Mining           | 154        | 87         | 110        | 123        | 128        | 104        | 98         | 96         | 102        |
| Surface Mining               | 22         | 25         | 27         | 24         | 24         | 24         | 21         | 20         | 20         |
| Post-Mining<br>(Underground) | 19         | 16         | 15         | 14         | 14         | 14         | 14         | 14         | 14         |
| Post-Mining (Surface)        | 5          | 5          | 6          | 5          | 5          | 5          | 5          | 4          | 4          |
| <b>Total</b>                 | <b>200</b> | <b>132</b> | <b>157</b> | <b>166</b> | <b>171</b> | <b>148</b> | <b>138</b> | <b>134</b> | <b>140</b> |

Note: Totals may not sum due to independent rounding.

**Table A-126: Total Coal Mining CH<sub>4</sub> Emissions by State (Million Cubic Feet)**

| State      | 1990   | 2005   | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   |
|------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Alabama    | 32,097 | 15,789 | 20,992 | 22,119 | 21,377 | 18,530 | 18,129 | 17,486 | 16,301 |
| Alaska     | 50     | 42     | 43     | 54     | 63     | 63     | 60     | 45     | 44     |
| Arizona    | 151    | 161    | 107    | 100    | 103    | 108    | 100    | 101    | 107    |
| Arkansas   | 5      | +      | 237    | 119    | 130    | 348    | 391    | 214    | 176    |
| California | 1      | 0      | 0      | 0      | 0      | 0      | 0      | 0      | 0      |
| Colorado   | 10,187 | 13,441 | 12,871 | 13,999 | 16,470 | 11,187 | 9,305  | 4,838  | 4,038  |
| Illinois   | 10,180 | 6,488  | 7,568  | 7,231  | 8,622  | 7,579  | 9,763  | 8,920  | 9,217  |
| Indiana    | 2,232  | 3,303  | 5,047  | 5,763  | 5,938  | 6,203  | 7,374  | 6,427  | 7,159  |
| Iowa       | 24     | 0      | 0      | 0      | 0      | 0      | 0      | 0      | 0      |
| Kansas     | 45     | 11     | 14     | 12     | 8      | 2      | 1      | 1      | 4      |

| State         | 1990           | 2005           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|---------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Kentucky      | 10,018         | 6,898          | 9,986          | 12,035         | 12,303         | 10,592         | 7,993          | 8,098          | 8,219          |
| Louisiana     | 64             | 84             | 77             | 73             | 79             | 168            | 80             | 56             | 52             |
| Maryland      | 474            | 361            | 263            | 219            | 238            | 263            | 197            | 166            | 169            |
| Mississippi   | 0              | 199            | 159            | 193            | 224            | 154            | 165            | 200            | 209            |
| Missouri      | 166            | 3              | 15             | 28             | 29             | 29             | 26             | 26             | 23             |
| Montana       | 1,373          | 1,468          | 1,629          | 1,417          | 1,495          | 1,445          | 1,160          | 1,269          | 1,379          |
| New Mexico    | 363            | 2,926          | 3,411          | 3,836          | 3,956          | 4,187          | 2,148          | 2,845          | 2,219          |
| North Dakota  | 299            | 306            | 303            | 306            | 296            | 289            | 281            | 282            | 298            |
| Ohio          | 4,406          | 3,120          | 3,686          | 4,443          | 3,614          | 3,909          | 3,389          | 3,182          | 3,267          |
| Oklahoma      | 226            | 825            | 932            | 624            | 436            | 360            | 499            | 282            | 112            |
| Pennsylvania  | 21,864         | 17,904         | 20,684         | 22,939         | 23,372         | 17,708         | 17,773         | 20,953         | 19,803         |
| Tennessee     | 276            | 115            | 86             | 69             | 67             | 60             | 35             | 31             | 22             |
| Texas         | 1,119          | 922            | 783            | 704            | 823            | 922            | 887            | 854            | 876            |
| Utah          | 3,587          | 4,787          | 5,524          | 5,449          | 5,628          | 3,651          | 3,624          | 2,733          | 1,605          |
| Virginia      | 46,041         | 8,649          | 9,223          | 8,042          | 9,061          | 8,526          | 6,516          | 8,141          | 12,680         |
| Washington    | 146            | 154            | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| West Virginia | 48,335         | 29,745         | 36,421         | 40,452         | 40,638         | 35,709         | 33,608         | 32,998         | 38,023         |
| Wyoming       | 6,671          | 14,745         | 16,959         | 15,627         | 16,032         | 15,916         | 14,507         | 14,025         | 14,339         |
| <b>Total</b>  | <b>200,399</b> | <b>132,481</b> | <b>157,112</b> | <b>165,854</b> | <b>171,000</b> | <b>147,908</b> | <b>138,012</b> | <b>134,173</b> | <b>140,343</b> |

+ Does not exceed 0.5 million cubic feet.

Note: The emission estimates provided above are inclusive of emissions from underground mines, surface mines and post-mining activities. The following states have neither underground nor surface mining and thus report no emissions as a result of coal mining: Connecticut, Delaware, Florida, Georgia, Hawaii, Idaho, Maine, Massachusetts, Michigan, Minnesota, Nebraska, Nevada, New Hampshire, New Jersey, New York, North Carolina, Oregon, Rhode Island, South Carolina, South Dakota, Vermont, and Wisconsin.

## References

- AAPG (1984) Coalbed Methane Resources of the United States. AAPG Studies in Geology Series #17.
- CAR (2015) Project Database. Climate Action Reserve. Available online at <<http://www.climateactionreserve.org/>>.
- Consol (2015) Ruby Canyon Summary 2014. CONSOL Energy Inc. excel spreadsheet.
- Creedy, D.P. (1993) Chemosphere. Vol. 26, pp. 419-440.
- EIA (2015) Annual Coal Report 1991-2014 (Formerly called Coal Industry Annual). Table 1. Energy Information Administration, U.S. Department of Energy, Washington, DC .
- EIA (2014) Annual Coal Report 1991-2013 (Formerly called Coal Industry Annual). Table 1. Energy Information Administration, U.S. Department of Energy, Washington, DC .
- El Paso (2009) Shoal Creek Mine Plan, El Paso Exploration & Production.
- EPA (2015) Greenhouse Gas Reporting Program (GHGRP) 2014 Envirofacts. Subpart FF: Underground Coal Mines. Available online at <<http://www.epa.gov/ghgreporting/ghgdata/reported/coalmines.html>>.
- EPA (2005) Surface Mines Emissions Assessment. Draft. U.S. Environmental Protection Agency.
- EPA (2004) Methane Emissions Estimates & Methodology for Abandoned Coal Mines in the U.S. Draft Final Report. Washington, DC. April - June 2004.
- EPA (1996) Evaluation and Analysis of Gas Content and Coal Properties of Major Coal Bearing Regions of the United States. U.S. Environmental Protection Agency. EPA/600/R-96-065.
- Geological Survey of Alabama State Oil and Gas Board (GSA) (2016) Well Records Database. Available online at <<http://www.gsa.state.al.us/ogb/database.aspx>>.
- IEA (2015) Key World Energy Statistics. Coal Production, International Energy Agency.
- IPCC (2011) Use of Models and Facility-Level Data in Greenhouse Gas Inventories. Report of IPCC Expert Meeting on Use of Models and Measurements in Greenhouse Gas Inventories 9-11 August 2010, Sydney, Australia. Eds: Eggleston H.S., Srivastava N., Tanabe K., Baasansuren J., Fukuda M. IGES.

- King, B. (1994) Management of Methane Emissions from Coal Mines: Environmental, Engineering, Economic and Institutional Implication of Options, Neil and Gunter Ltd., Halifax, March 1994.
- JWR (2014) Wells Intercepted in 2013. Jim Walter Resources excel spreadsheet.
- JWR (2010) No. 4 & 7 Mines General Area Maps. Walter Energy: Jim Walter Resources.
- MSHA (2015) Data Transparency at MSHA. Mine Safety and Health Administration. Available online at <<http://www.msha.gov/>>.
- Mutmansky, Jan M., and Yanbei Wang (2000) Analysis of Potential Errors in Determination of Coal Mine Annual Methane Emissions. Department of Energy and Geo-Environmental Engineering, Pennsylvania State University. University Park, PA.
- Saghafi, Abouna (2013) Estimation of fugitive emissions from open cut coal mining and measurable gas content, 13th Coal Operators' Conference, University of Wollongong, The Australian Institute of Mining and Metallurgy & Mine Managers Association of Australia, 2013, 306-313.
- USBM (1986) Results of the Direct Method Determination of the Gas Contents of U.S. Coal Basins. Circular 9067, U.S. Bureau of Mines.
- West Virginia Geological & Economic Survey (WVGES) (2015) Oil & Gas Production Data. Available online at <<http://www.wygs.wvnet.edu/www/datastat/datastat.htm>>.

### 3.5. Methodology for Estimating CH<sub>4</sub> and CO<sub>2</sub> Emissions from Petroleum Systems

The methodology for estimating CH<sub>4</sub> and non-combustion CO<sub>2</sub> emissions from the production, refineries and transportation segments of petroleum systems is generally based on EPA's Greenhouse Gas Reporting Program (GHGRP) data for 2014; the 1999 EPA report, Estimates of Methane Emissions from the U.S. Oil Industry (EPA 1999); and the study, Methane Emissions from the U.S. Petroleum Industry (EPA/GRI 1996). Fifty-seven activities that emit CH<sub>4</sub> and thirty-two activities that emit non-combustion CO<sub>2</sub> from petroleum systems were examined from these reports and the GHGRP data. Most of the activities are part of crude oil production field operations, which accounted for approximately 99 percent of total oil industry CH<sub>4</sub> emissions. Crude transportation and refining accounted for the remaining CH<sub>4</sub> emissions of approximately 0.3 and 0.9 percent, respectively. Non-combustion CO<sub>2</sub> emissions were analyzed for production operations and asphalt blowing, flaring, and process vents in refining operations. Non-combustion CO<sub>2</sub> emissions from transportation operations are not included because they are negligible. The following steps were taken to estimate CH<sub>4</sub> and CO<sub>2</sub> emissions from petroleum systems.

#### Step 1: Calculate Potential Methane (or Net Emissions)

##### Activity Data

Activity data were taken from the following source: DrillingInfo (2015), the Energy Information Administration annual and monthly reports (EIA 1990 through 2015), (EIA 1995 through 2015a, 2015b), "Methane Emissions from the Natural Gas Industry by the Gas Research Institute and EPA" (EPA/GRI 1996a-d), "Estimates of Methane Emissions from the U.S. Oil Industry" (EPA 1999), consensus of industry peer review panels, BOEMRE and BOEM reports (BOEMRE 2004, BOEM 2011), analysis of BOEMRE data (EPA 2005, BOEMRE 2004), the Oil & Gas Journal (OGJ 2015), the Interstate Oil and Gas Compact Commission (IOGCC 2012), the United States Army Corps of Engineers, (1995-2015), and the GHGRP (2010-2014).

For many sources, complete activity data were not available for all years of the time series. In such cases, one of three approaches was employed. Where appropriate, the activity data were calculated from related statistics using ratios developed based on EPA 1996, and/or GHGRP data. For major equipment, pneumatic controllers, and chemical injection pumps, GHGRP subpart W data were used to develop activity factors (i.e., count per well) that are applied to calculated activity in recent years; to populate earlier years of the time series, linear interpolation is used to connect GHGRP-based estimates with existing estimates in years 1990 to 1995. For hydraulically fractured oil well completions, EPA developed activity data specific to each year of the time series using the date of completion or first reported production available from a data set licensed by DrillingInfo, Inc. For more information on the DrillingInfo data processing, please see Annex 3.6 Methodology for Estimating CH<sub>4</sub> and CO<sub>2</sub> from Natural Gas Systems. In other cases, the activity data were held constant from 1990 through 2014 based on EPA (1999). Lastly, the previous year's data were used when data for the current year were unavailable. For offshore production, the number of platforms in shallow water and the number of platforms in deep water are used as activity data and are taken from Bureau of Ocean Energy Management (BOEM) (formerly Bureau of Ocean Energy Management, Regulation, and Enforcement [BOEMRE]) datasets (BOEM 2011a,b,c).

For petroleum refining activities, 2010 to 2014 emissions were directly obtained from EPA's GHGRP. All refineries have been required to report CH<sub>4</sub> and CO<sub>2</sub> emissions for all major activities since 2010. The national totals of these emissions for each activity were used for the 2010 to 2014 emissions. The national emission totals for each activity were divided by refinery feed rates for those four Inventory years to develop average activity-specific emission factors, which were used to estimate national emissions for each refinery activity from 1990 to 2009 based on national refinery feed rates for each year (EPA 2015d).

The activity data for the total crude transported in the transportation sector is not available. In this case, all the crude oil that was transported was assumed to go to refineries. Therefore, the activity data for the refining sector (i.e., refinery feed in 1000 bbl/year) was used also for the transportation sector. In the few cases where no data were located, oil industry data based on expert judgment was used. In the case of non-combustion CO<sub>2</sub> emission sources, the activity factors are the same as for CH<sub>4</sub> emission sources. In some instances, where 2014 data are not yet available 2013 or prior data has been used as proxy.

##### Potential methane factors and emission factors

The CH<sub>4</sub> emission factors for the majority of the activities are taken from the 1999 EPA draft report, which contained the most recent and comprehensive determination of CH<sub>4</sub> emission factors for the 57 CH<sub>4</sub>-emitting activities in the oil industry at that time. Since the time of this study, practices and technologies have changed. While this study still represents best available data in many cases, using these emission factors alone to represent actual emissions without adjusting for emissions controls would in many cases overestimate emissions. For this reason, "potential methane" is

calculated using the data, and then recent data on voluntary and regulatory emission reduction activities (Step 3) is deducted to calculate actual emissions.

For certain CH<sub>4</sub> emissions sources, new data and information allows for net emissions to be calculated directly: Oil well completions with hydraulic fracturing, pneumatic controllers, chemical injection pumps, and refineries. For these sources, EPA used emissions factors that directly reflect recent practice and the use of control technologies. The controlled and uncontrolled emission factors for hydraulically fractured (HF) well completions were developed using data analyzed for the 2015 NSPS OOOOa proposal. The gas engine emission factor is taken from the EPA/GRI 1996b study. For pneumatic controllers, separate estimates are developed for low bleed, high bleed, and intermittent controllers. Chemical injection pumps estimate are calculated with an emission factor developed with GHGRP data, which is based on the previous GRI/EPA factor but takes into account operating hours. The refineries emission factors were also developed from GHGRP data.

Other emission factors that are not based on the 1999 report include oil tank venting and offshore platforms. The oil tank venting emission factor is taken from the API E&P Tank Calc weighted average for API gravity less than 45 API degrees with the distribution of gravities taken from a sample of production data from the HPDI database. Offshore emissions from shallow water and deep water oil platforms are taken from analysis of the Gulf-wide Offshore Activity Data System (GOADS) report (EPA 2015; BOEM 2014). The emission factors were assumed to be representative of emissions from each source type over the period 1990 through 2014, and are used for each year throughout this period.

In general, the CO<sub>2</sub> emission factors were derived from the corresponding source CH<sub>4</sub> emission factors. The amount of CO<sub>2</sub> in the crude oil stream changes as it passes through various equipment in petroleum production operations. As a result, four distinct stages/streams with varying CO<sub>2</sub> contents exist. The four streams that are used to estimate the emissions factors are the associated gas stream separated from crude oil, hydrocarbons flashed out from crude oil (such as in storage tanks), whole crude oil itself when it leaks downstream, and gas emissions from offshore oil platforms. The standard approach used to estimate CO<sub>2</sub> emission factors was to use the existing CH<sub>4</sub> emissions factors and multiply them by a conversion factor, which is the ratio of CO<sub>2</sub> content to methane content for the particular stream. Ratios of CO<sub>2</sub> to CH<sub>4</sub> volume in emissions are presented in Table A-131. The exceptions are the emissions factor for storage tanks, which are estimated using API E&P Tank Calc simulation runs of tank emissions for crude oil of different gravities less than 45 API degrees; emission factors for shallow water and deep water platforms, which are estimated from analysis of the *2011 Gulf-Wide Emission Inventory Study* (BOEM 2014) and the emissions estimates for refineries, which are estimated using the data from U.S. EPA's GHGRP.

## **Step 2: Compile Reductions Data**

The methane emissions calculated in Step 1 above generally represent expected emissions from an activity in the absence of emissions controls (with the exceptions of pneumatic controllers, completions with hydraulic fracturing, etc. as noted above), and do not take into account any use of technologies or practices that reduce emissions. To take into account use of such technologies, data were collected on voluntary reductions. Voluntary reductions included in the Petroleum Sector calculations were those reported to Natural Gas STAR for the following activities: Artificial lift: gas lift, Artificial lift: use compression, Artificial lift: use pumping unit, Consolidate crude oil prod and water storage tanks, Lower heater-treater temperature, Re-inject gas for enhanced oil recovery, Re-inject gas into crude, and Route casing head gas to VRU or compressor.

Industry partners report CH<sub>4</sub> emission reductions by project to the Natural Gas STAR Program. The reductions from the implementation of specific technologies and practices are calculated by the reporting partners using actual measurement data or equipment-specific emission factors. The reductions undergo quality assurance and quality control checks to identify errors, inconsistencies, or irregular data before being incorporated into the Inventory. The technology-specific approach for pneumatic controllers allows net emissions to be calculated directly. Pneumatic controller-related reductions were removed from the Gas STAR reduction data set before it was used to adjust the petroleum segment estimates.

## **Step 3: Calculate Net Methane and Carbon Dioxide Emissions for Each Activity for Each Year**

Annual CH<sub>4</sub> emissions from each of the 57 petroleum system activities and CO<sub>2</sub> emissions from the 32 petroleum system activities were estimated by multiplying the activity data for each year by the corresponding emission factor, except for petroleum refineries segment. Emissions from refineries were obtained directly from the GHGRP data for 2010 through 2014; these four years of data were used to develop emission factors and activity data that are applied for the remainder of the time-series (i.e., 1990 through 2009). These annual emissions for each activity were then summed to estimate the total annual CH<sub>4</sub> and CO<sub>2</sub> emissions, respectively. As a final step, the relevant Natural Gas STAR reductions data is summed for each year and deducted from the total emissions to estimate net CH<sub>4</sub> emissions for the Inventory.

Table A-127, Table A-128, Table A-129, and Table A-132 provide 2014 activity data, emission factors, and emission estimates and Table A-130 and Table A-133 provide a summary of emission estimates for the years 1990, 1995, 2000, and 2005 through 2014. Table A-131 provides the CO<sub>2</sub> content in natural gas for equipment in different crude streams to estimate CO<sub>2</sub> emission factors using CH<sub>4</sub> emission factors.

The tables provide references for emission factors and activity data in footnotes (the lettered footnotes). The tables also provide information on which method was used for supplying activity data for 2014 (the numbered footnotes).

Key to table notations on methods for supplying activity data for 2014 for all tables:

1. Ratios relating other factors for which activity data are available.
2. Activity data for 2014 available from source.
3. Activity data were held constant from 1990 through 2014 based on EPA (1999).
4. 2009, 2010, 2011, 2012, or 2013 activity data are used to determine some or all of the 2014 activity data.

**Table A-127: 2014 CH<sub>4</sub> Emissions from Petroleum Production Field Operations**

| Activity/Equipment  | 2014 EPA Inventory Values                                   |   | Emissions<br>(Bcf/yr) | Emissions<br>(kt/yr) |
|---|---|---|-----------------------|----------------------|
|   | Emission Factor   | Activity Data   |                       |                      |
| <b>Vented Emissions</b>   |   |   | <b>128.872</b>        | <b>2,478.3</b>       |
| Oil Tanks   | 7.4 scf of CH <sub>4</sub> /bbl crude <sup>a</sup>          | 2,782 MMbbl/yr (non-stripper wells) <sup>b,c,d,1</sup>  | 20.575                | 395.7                |
| Pneumatic controllers, High Bleed                                     | 622 scfd CH <sub>4</sub> /controller <sup>r</sup>           | 43,211 No. of high-bleed controllers <sup>r,2</sup>     | 9.806                 | 188.6**              |
| Pneumatic controllers, Low Bleed                                      | 23 scfd CH <sub>4</sub> /controller <sup>r</sup>            | 300,940 No. of low-bleed controllers <sup>r,2</sup>     | 2.510                 | 48.3**               |
| Pneumatic controllers, Int Bleed                                      | 218 scfd CH <sub>4</sub> /controller <sup>r</sup>           | 868,079 No. of int-bleed controllers <sup>r,2</sup>     | 69.173                | 1,330.2**            |
| Chemical Injection Pumps  | 216 scfd CH <sub>4</sub> /pump <sup>r</sup>                 | 127,484 No. of pumps <sup>r,2</sup>                     | 10.030                | 192.9                |
| Vessel Blowdowns  | 78 scfy CH <sub>4</sub> /vessel <sup>h</sup>                | 502,754 No. of vessels <sup>c,g,i,q,1</sup>             | 0.039                 | 0.8                  |
| Compressor Blowdowns  | 3,775 scf/yr of CH <sub>4</sub> /compressor <sup>h</sup>    | 2,967 No. of compressors <sup>c,g,i,q,1</sup>           | 0.011                 | 0.2                  |
| Compressor Starts   | 8,443 scf/yr of CH <sub>4</sub> /compressor <sup>h</sup>    | 2,967 No. of compressors <sup>c,g,i,1</sup>             | 0.025                 | 0.5                  |
| Stripper wells  | 2,345 scf/yr of CH <sub>4</sub> /stripper well <sup>f</sup> | 327,674 No. of stripper wells vented <sup>f,d,4</sup>   | 0.768                 | 14.8                 |
| Well Completion Venting   | 733 scf/completion <sup>h</sup>                             | 15,753 Oil well completions <sup>s,4</sup>              | 0.012                 | 0.2                  |
| Well Workovers  | 96 scf CH <sub>4</sub> /workover <sup>i</sup>               | 67,370 Oil well workovers <sup>g,i,1,4</sup>            | 0.006                 | 0.1                  |
| HF Well Completions, Uncontrolled                                     | 351,146 scf/HF completions <sup>s</sup>                     | 17,522 HF oil well completions <sup>t,2</sup>           | 6.153                 | 118.3                |
| HF Well Completions, Controlled                                       | 17,557 scf/HF completion <sup>s</sup>                       | 1,319 HF oil well completions <sup>t,2</sup>            | 0.023                 | 0.4                  |
| Pipeline Pigging  | 2.4 scfd of CH <sub>4</sub> /pig station <sup>i</sup>       | 0 No. of crude pig stations <sup>e,3</sup>              | 0.000                 | 0.0                  |
| Offshore Platforms, Shallow water Oil, fugitive, vented and combusted | 16,552 scfd CH <sub>4</sub> /platform <sup>q</sup>          | 1,447 No. of shallow water oil platforms <sup>l,4</sup> | 8.739                 | 168.1                |
| Offshore Platforms, Deepwater oil, fugitive, vented and combusted     | 93,836 scfd CH <sub>4</sub> /platform <sup>q</sup>          | 29 No. of deep water oil platforms <sup>l,4</sup>       | 1.001                 | 19.3                 |
| <b>Fugitive Emissions</b>   |   |   | <b>6.479</b>          | <b>124.6</b>         |
| Oil Wellheads (heavy crude)   | 0.13 scfd/well <sup>e,m</sup>                               | 40,227 No. of hvy. Crude wells <sup>d,i,t,1,4</sup>     | 0.002                 | 0.0*                 |
| Oil Wellheads (light crude)   | 17 scfd/well <sup>e,m</sup>                                 | 530,367 No. of lt. crude wells <sup>d,i,t,1,4</sup>     | 3.219                 | 61.9                 |
| Separators (heavy crude)  | 0.15 scfd CH <sub>4</sub> /separator <sup>e,m</sup>         | 38,520 No. of hvy. Crude seps. <sup>r,2</sup>           | 0.002                 | 0.0*                 |
| Separators (light crude)  | 14 scfd CH <sub>4</sub> /separator <sup>e,m</sup>           | 350,573 No. of lt. crude seps. <sup>r,2</sup>           | 1.773                 | 34.1                 |
| Heater/Treaters (light crude)   | 19 scfd CH <sub>4</sub> /heater <sup>e,m</sup>              | 113,662 No. of heater treaters <sup>r,2</sup>           | 0.796                 | 15.3                 |
| Headers (heavy crude)   | 0.08 scfd CH <sub>4</sub> /header <sup>e,m</sup>            | 35,647 No. of hvy. Crude hdrs. <sup>r,2</sup>           | 0.001                 | 0.0*                 |
| Headers (light crude)   | 11 scfd CH <sub>4</sub> /header <sup>e,m</sup>              | 110,509 No. of lt. crude hdrs. <sup>r,2</sup>           | 0.438                 | 8.4                  |
| Floating Roof Tanks   | 338,306 tank/yr <sup>m,n</sup>                              | 24 No. of floating roof tanks <sup>e,3</sup>            | 0.008                 | 0.2                  |
| Compressors   | 100 scfd CH <sub>4</sub> /compressor <sup>e</sup>           | 2,967 No. of compressors <sup>c,i,t,1</sup>             | 0.108                 | 2.1                  |
| Large Compressors   | 16,360 scfd CH <sub>4</sub> /compressor <sup>e</sup>        | 0 No. of large comprs. <sup>e,3</sup>                   | 0.000                 | 0.0                  |
| Sales Areas   | 41 scf CH <sub>4</sub> /loading <sup>e</sup>                | 2,648,868 Loadings/year <sup>c,1</sup>                  | 0.107                 | 2.1                  |
| Pipelines   | NE scfd of CH <sub>4</sub> /mile of pipeline                | 14,571 Miles of gathering line <sup>e,2</sup>           | NE                    | NE                   |
| Well Drilling   | NE scfd of CH <sub>4</sub> /oil well drilled                | 17,774 No. of oil wells drilled <sup>c,2</sup>          | NE                    | NE                   |
| Battery Pumps   | 0.24 scfd of CH <sub>4</sub> /pump <sup>m</sup>             | 269,480 No. of battery pumps <sup>g,e,1</sup>           | 0.024                 | 0.5                  |
| <b>Combustion Emissions</b>   |   |   | <b>6.196</b>          | <b>119.2</b>         |
| Gas Engines   | 0.24 scf CH <sub>4</sub> /HP-hr <sup>h</sup>                | 18,694 MMHP-hr <sup>c,g,i,1</sup>                       | 4.487                 | 86.3                 |

| Activity/Equipment               | 2014 EPA Inventory Values                            |  |  | Emissions<br>(Bcf/yr) | Emissions<br>(kt/yr) |
|----------------------------------|--|--|--|-----------------------|----------------------|
|                                  | Emission Factor                                      | Activity Data                              |  |                       |                      |
| Heaters                          | 0.52 scf CH <sub>4</sub> /bbl <sup>n</sup>           | 3,179 MMbbl/yr <sup>c,2</sup>              |  | 1.657                 | 31.9                 |
| Well Drilling                    | 2,453 scf CH <sub>4</sub> /well drilled <sup>m</sup> | 17,774 Oil wells drilled <sup>c,4</sup>    |  | 0.044                 | 0.8                  |
| Flares                           | 20 scf CH <sub>4</sub> /Mcf flared <sup>i</sup>      | 452,651 Mcf flared/yr <sup>b,c,d,1,4</sup> |  | 0.009                 | 0.2                  |
| <b>Process Upset Emissions</b>   |  |  |  | <b>0.157</b>          | <b>3.0</b>           |
| Pressure Relief Valves           | 35 scf/yr/PR valve <sup>h</sup>                      | 265,554 No. of PR valves <sup>c,e,1</sup>  |  | 0.009                 | 0.2                  |
| Well Blowouts Onshore            | 2.5 MMscf/blowout <sup>f</sup>                       | 59 No. of blowouts/yr <sup>c,e,1</sup>     |  | 0.148                 | 2.8                  |
| <b>Voluntary Reductions</b>      |  |  |  | <b>(1.605)</b>        | <b>(30.9)</b>        |
| <b>Total Potential Emissions</b> |  |  |  | <b>141.704</b>        | <b>2,725.1</b>       |
| <b>Total Net Emissions</b>       |  |  |  | <b>140.100</b>        | <b>2,694.2</b>       |

- Zero Emissions

\* Emissions are not actually 0, but too small to show at this level of precision.

<sup>a</sup> TankCALC

<sup>b</sup> EPA / ICF International (1999)

<sup>c</sup> Energy Information Administration (EIA) Monthly Energy Review

<sup>d</sup> Interstate Oil & Gas Compact Commission (IOGCC) Marginal Wells Report

<sup>e</sup> Consensus of Industrial Review Panel

<sup>f</sup> Expert Judgment

<sup>g</sup> EIA Annual Energy Review

<sup>h</sup> Gas Research Institute (GRI) / EPA (1996)

<sup>i</sup> Radian (1999)

<sup>j</sup> Canadian Association of Petroleum Producers (CAPP) (1992)

<sup>k</sup> Adapted from the Minerals Management Service (MMS) Gulfwide Offshore Activities Data System (GOADS) by ICF (2005)

<sup>l</sup> Bureau of Ocean Energy Management (BOEM)

<sup>m</sup> American Petroleum Institute (API) (1996)

<sup>n</sup> EPA, AP 42 Compilation of Air Pollutant Emission Factors

<sup>o</sup> Oil and Gas Journal (OGJ) Petroleum Economics Issue

<sup>p</sup> BOEM 2011 Gulf-wide Emissions Inventory Study (2014)

<sup>r</sup> GHGRP data

<sup>s</sup> DrillingInfo data from analysis supporting the NSPS OOOOa proposal

<sup>t</sup> DrillingInfo data

**Table A-128: 2014 CH<sub>4</sub> Emissions from Petroleum Transportation**

| Activity/Equipment          | Emission  |       | Activity   | Emissions<br>(Bcf/yr) | Emissions<br>(kt/yr) |
|-----------------------------|---|-------|--|-----------------------|----------------------|
|                             | Factor  | Units |  |                       |                      |
| <b>Vented Emissions</b>     |   |       |  | <b>0.370</b>          | <b>7.1</b>           |
| Tanks                       | 0.021 scf CH <sub>4</sub> /yr/bbl of crude delivered to refineries <sup>a</sup> |       | 5,785 MMbbl crude feed/yr <sup>b,2</sup>         | 0.119                 | 2.3                  |
| Truck Loading               | 0.520 scf CH <sub>4</sub> /yr/bbl of crude transported by truck <sup>c</sup>    |       | 152 MMbbl crude trans. By truck <sup>d,2</sup>   | 0.079                 | 1.5                  |
| Marine Loading              | 2.544 scf CH <sub>4</sub> /1000 gal crude marine loadings <sup>c</sup>          |       | 33,954,617 1,000 gal/yr loaded <sup>e,1,4</sup>  | 0.086                 | 1.7                  |
| Rail Loading                | 0.520 scf CH <sub>4</sub> /yr/bbl of crude transported by rail <sup>c</sup>     |       | 133 MMbbl Crude by rail/yr <sup>d,2</sup>        | 0.069                 | 1.3                  |
| Pump Station Maintenance    | 36.80 scf CH <sub>4</sub> /station/yr <sup>f</sup>                              |       | 564 No. of pump stations <sup>g,1</sup>          | 0.000*                | 0.0*                 |
| Pipeline Pigging            | 39 scfd of CH <sub>4</sub> /pig station <sup>h</sup>                            |       | 1,128 No. of pig stations <sup>g,1</sup>         | 0.016                 | 0.3                  |
| <b>Fugitive Emissions</b>   |   |       |  | <b>0.050</b>          | <b>1.0</b>           |
| Pump Stations               | 25 scf CH <sub>4</sub> /mile/yr <sup>f</sup>                                    |       | 56,375 No. of miles of crude p/yr <sup>g,2</sup> | 0.001                 | 0.0*                 |
| Pipelines                   | NE scf CH <sub>4</sub> /bbl crude transported by pipeline <sup>f</sup>          |       | 9,289 MMbbl crude piped <sup>g,2</sup>           | NE                    | NE                   |
| Floating Roof Tanks         | 58,965 scf CH <sub>4</sub> /floating roof tank/yr <sup>i</sup>                  |       | 824 No. of floating roof tanks <sup>3</sup>      | 0.049                 | 0.9                  |
| <b>Combustion Emissions</b> |   |       |  | <b>NE</b>             | <b>NE</b>            |
| Pump Engine Drivers         | 0.24 scf CH <sub>4</sub> /hp-hr <sup>i</sup>                                    |       | NE No. of hp-hrs                                 | NE                    | NE                   |
| Heaters                     | 0.521 scf CH <sub>4</sub> /bbl burned <sup>k</sup>                              |       | NE No. of bbl Burned                             | NE                    | NE                   |
| <b>Total</b>                |   |       |  | <b>0.420</b>          | <b>8.1</b>           |

\* Emissions are not actually 0, but too small to show at this level of precision.

NE - Not estimated for lack of data

<sup>a</sup> API (1992)

<sup>b</sup> Energy Information Administration (EIA) Petroleum Supply Annual, Volume 1.

<sup>c</sup> EPA, AP 42 Compilation of Air Pollutant Emission Factors

<sup>d</sup> EIA Refinery Capacity Report

<sup>e</sup> EIA Monthly Energy Review  
<sup>f</sup> Radian (1996)  
<sup>g</sup> OGJ Petroleum Economics Issue  
<sup>h</sup> CAPP (1992)  
<sup>i</sup> API TANK  
<sup>j</sup> GRI / EPA (1996)  
<sup>k</sup> EPA / ICF International (1999)

**Table A-129: 2014 CH<sub>4</sub> Emissions from Petroleum Refining**

| Activity/Equipment                    | 2014 EPA Inventory Values          |   |                    |                   |
|---------------------------------------|------------------------------------|---|--------------------|-------------------|
|                                       | Emission Factor                    | Activity Factor                                 | Emissions (Bcf/yr) | Emissions (kt/yr) |
| <b>Vented Emissions</b>               |                                    |   | <b>0.373</b>       | <b>7.2</b>        |
| Uncontrolled Blowdowns                | 0.000971 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.292              | 5.6               |
| Asphalt Blowing                       | 0.000049 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.015              | 0.3               |
| Process Vents                         | 0.000215 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.065              | 1.2               |
| CEMS                                  | 0.000006 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.002              | 0.0*              |
| <b>Fugitive Emissions</b>             |                                    |   | <b>0.234</b>       | <b>4.5</b>        |
| Equipment Leaks                       | 0.000457 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.137              | 2.6               |
| Storage Tanks                         | 0.000237 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.071              | 1.4               |
| Wastewater Treating                   | 0.00798 lb VOC/bbl <sup>b,c</sup>  | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.011              | 0.2               |
| Cooling Towers                        | 0.010 lb VOC/bbl <sup>b,c</sup>    | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.014              | 0.3               |
| Loading Operations                    | 0.000002 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.001              | 0.0*              |
| <b>Combustion Emissions</b>           |                                    |   | <b>0.614</b>       | <b>11.8</b>       |
| Catalytic Cracking, Coking, Reforming | 0.000248 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.075              | 1.4               |
| Flares                                | 0.001611 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.485              | 9.3               |
| Delay Cokers                          | 0.000178 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.054              | 1.0               |
| Coke Calcining                        | 0.000005 MT CH <sub>4</sub> /Mbbld | 5,784,637 Mbbbl/year refinery feed <sup>a</sup> | 0.002              | 0.0*              |
| <b>Total</b>                          |                                    |   | <b>1.221</b>       | <b>23.5</b>       |

\* Emissions are not actually 0, but too small to show at this level of precision.

<sup>a</sup> EIA Petroleum Supply Annual, Volume 1.

<sup>b</sup> Radian (1996)

<sup>c</sup> Assuming methane is 1% of total hydrocarbons (AP-42)

<sup>d</sup> GHGRP data

Note: The methodology for year 2014 is to use GHGRP emissions data as-reported. The emission factors in this table were developed for this table by dividing 2014 emissions by 2014 refinery feed rate.

**Table A-130: Summary of CH<sub>4</sub> Emissions from Petroleum Systems (kt)**

| Activity                           | 1990         | 1995         | 2000         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Production Field Operations</b> | <b>1,519</b> | <b>1,549</b> | <b>1,767</b> | <b>2,056</b> | <b>2,103</b> | <b>2,120</b> | <b>2,193</b> | <b>2,263</b> | <b>2,347</b> | <b>2,586</b> | <b>2,725</b> |
| Pneumatic controller venting       | 761          | 832          | 1,044        | 1,262        | 1,286        | 1,308        | 1,328        | 1,346        | 1,332        | 1,509        | 1,567        |
| Tank venting                       | 250          | 226          | 214          | 193          | 185          | 202          | 210          | 220          | 278          | 330          | 396          |
| Combustion & process upsets        | 115          | 105          | 95           | 92           | 95           | 95           | 98           | 101          | 108          | 114          | 122          |
| Misc. venting & fugitives          | 334          | 332          | 365          | 459          | 482          | 461          | 502          | 540          | 570          | 573          | 578          |
| Wellhead fugitives                 | 59           | 54           | 48           | 50           | 55           | 53           | 54           | 56           | 59           | 60           | 62           |
| <b>Crude Oil Transportation</b>    | <b>7</b>     | <b>6</b>     | <b>5</b>     | <b>5</b>     | <b>5</b>     | <b>5</b>     | <b>5</b>     | <b>5</b>     | <b>6</b>     | <b>7</b>     | <b>8</b>     |
| <b>Refining</b>                    | <b>24</b>    | <b>25</b>    | <b>27</b>    | <b>27</b>    | <b>26</b>    | <b>25</b>    | <b>26</b>    | <b>28</b>    | <b>27</b>    | <b>26</b>    | <b>23</b>    |
| <b>Voluntary Reductions</b>        | <b>(0)</b>   | <b>(1)</b>   | <b>(17)</b>  | <b>(64)</b>  | <b>(77)</b>  | <b>(67)</b>  | <b>(60)</b>  | <b>(45)</b>  | <b>(45)</b>  | <b>(31)</b>  | <b>(31)</b>  |
| <b>Total Potential Emissions</b>   | <b>1,550</b> | <b>1,580</b> | <b>1,799</b> | <b>2,088</b> | <b>2,134</b> | <b>2,150</b> | <b>2,224</b> | <b>2,296</b> | <b>2,380</b> | <b>2,619</b> | <b>2,757</b> |
| <b>Total Net Emissions</b>         | <b>1,550</b> | <b>1,578</b> | <b>1,781</b> | <b>2,024</b> | <b>2,058</b> | <b>2,083</b> | <b>2,163</b> | <b>2,251</b> | <b>2,335</b> | <b>2,588</b> | <b>2,726</b> |

Note: Totals may not sum due to independent rounding. Parentheses indicate emissions reductions.

**Table A-131: Ratios of CO<sub>2</sub> to CH<sub>4</sub> Volume in Emissions from Petroleum Production Field Operations**

|   | Whole Crude,<br>Post-Separator | Associated Gas | Tank Flash Gas | Offshore |
|---|--------------------------------|----------------|----------------|----------|
| Ratio %CO <sub>2</sub> / %CH <sub>4</sub> | 0.052                          | 0.020          | 0.017          | 0.004    |

**Table A-132: 2014 CO<sub>2</sub> Emissions from Petroleum Production Field Operations and Petroleum Refining**

| Activity/Equipment  | 2014 EPA Inventory Values   |  |                       |                      |
|---|---|--|-----------------------|----------------------|
|   | Emission Factor   | Activity Factor  | Emissions<br>(Bcf/yr) | Emissions<br>(kt/yr) |
| <b>Vented Emissions</b>   |   |  | <b>11.974</b>         | <b>633.2</b>         |
| Oil Tanks   | 3.528 scf of CO <sub>2</sub> /bbl crude <sup>a</sup>                                    | 2,782 MMbbl/yr (non stripper wells) <sup>b,c,d,1,4</sup> | 9.817                 | 519.1                |
| Pneumatic controllers, High Bleed                                     | 12.615 scfd CO <sub>2</sub> /controller <sup>q</sup>                                    | 43,211 No. of high-bleed controllers <sup>q,2</sup>      | 0.199                 | 10.5                 |
| Pneumatic controllers, Low Bleed                                      | 0.464 scfd CO <sub>2</sub> /controller <sup>q</sup>                                     | 300,940 No. of low-bleed controllers <sup>q,2</sup>      | 0.051                 | 2.7                  |
| Pneumatic controllers, Int Bleed                                      | 4.430 scfd CO <sub>2</sub> /controller <sup>q</sup>                                     | 868,079 No. of int-bleed controllers <sup>q,2</sup>      | 1.404                 | 74.2                 |
| Chemical Injection Pumps  | 4.374 scfd CO <sub>2</sub> /pump <sup>q</sup>   | 127,484 No. of pumps <sup>q,2</sup>                      | 0.204                 | 10.8                 |
| Vessel Blowdowns  | 1.583 scfy CO <sub>2</sub> /vessel <sup>h</sup><br>scf/yr of                            | 502,754 No. of vessels <sup>c,g,i,1</sup>                | 0.001                 | 0.0*                 |
| Compressor Blowdowns  | 77 CO <sub>2</sub> /compressor <sup>h</sup><br>scf/yr of                                | 2,967 No. of compressors <sup>c,g,i,1</sup>              | 0.000*                | 0.0*                 |
| Compressor Starts   | 171 CO <sub>2</sub> /compressor <sup>h</sup><br>scf/yr of CO <sub>2</sub> /stripper     | 2,967 No. of compressors <sup>c,g,i,1</sup>              | 0.001                 | 0.0*                 |
| Stripper wells  | 48 well <sup>f</sup>  | 327,674 No. of stripper wells vented <sup>f,1,4</sup>    | 0.016                 | 0.8                  |
| Well Completion Venting   | 14.87 scf/completion <sup>h</sup>   | 15,753 Oil well completions <sup>c,2</sup>               | 0.000*                | 0.0*                 |
| Well Workovers  | 1.95 scf CO <sub>2</sub> /workover <sup>i</sup>   | 67,370 Oil well workovers <sup>g,i,1</sup>               | 0.000*                | 0.0*                 |
| HF Well Completions, Uncontrolled                                     | 7,125.06 scf/HF completion <sup>f</sup>   | 17,522 HF oil well completions <sup>s,1</sup>            | 0.125                 | 6.6                  |
| HF Well Completions, Controlled                                       | 356.25 scf/HF completion <sup>f</sup>   | 1,319 HF oil well completions <sup>s,1</sup>             | 0.000*                | 0.0*                 |
| Pipeline Pigging  | NE scfd of CO <sub>2</sub> /pig station   | 0 No. of crude pig stations                              | NE                    | NE                   |
| Offshore Platforms, Shallow water Oil, fugitive, vented and combusted | 276 scfd CO <sub>2</sub> /platform <sup>k</sup>   | 1,447 No. of shallow water oil platforms <sup>l,4</sup>  | 0.146                 | 7.7                  |
| Offshore Platforms, Deepwater oil, fugitive, vented and combusted     | 1,100 scfd CO <sub>2</sub> /platform <sup>k</sup>                                       | 29 No. of deep water oil platforms <sup>l,4</sup>        | 0.012                 | 0.6                  |
| <b>Fugitive Emissions</b>   |   |  | <b>0.133</b>          | <b>7.0</b>           |
| Oil Wellheads (heavy crude)   | 0.003 scfd/well <sup>e,m</sup>  | 40,227 No. of hvy. crude wells <sup>d,g,i,1,4</sup>      | 0.000*                | 0.0*                 |
| Oil Wellheads (light crude)   | 0.337 scfd/well <sup>e,m</sup>  | 530,367 No. of lt. crude wells <sup>d,g,i,1,4</sup>      | 0.065                 | 3.5                  |
| Separators (heavy crude)  | 0.003 scfd CO <sub>2</sub> /separator <sup>e,m</sup>                                    | 38,520 No. of hvy. crude seps. <sup>q,2</sup>            | 0.000*                | 0.0*                 |
| Separators (light crude)  | 0.281 scfd CO <sub>2</sub> /separator <sup>e,m</sup>                                    | 350,573 No. of lt. crude seps. <sup>q,2</sup>            | 0.036                 | 1.9                  |
| Heater/Treaters (light crude)   | 0.319 scfd CO <sub>2</sub> /heater <sup>e,m</sup>                                       | 113,662 No. of heater treaters <sup>q,2</sup>            | 0.013                 | 0.7                  |
| Headers (heavy crude)   | 0.002 scfd CO <sub>2</sub> /header <sup>e,m</sup>                                       | 35,647 No. of hvy. crude hdrs. <sup>q,2</sup>            | 0.000*                | 0.0*                 |
| Headers (light crude)   | 0.220 scfd CO <sub>2</sub> /header <sup>e,m</sup><br>scf CO <sub>2</sub> /floating roof | 110,509 No. of lt. crude hdrs. <sup>q,2</sup>            | 0.009                 | 0.5                  |
| Floating Roof Tanks   | 17,490 tank/yr <sup>m,n</sup>   | 24 No. of floating roof tanks <sup>e,3</sup>             | 0.000*                | 0.0*                 |
| Compressors   | 2.029 scfd CO <sub>2</sub> /compressor <sup>e</sup>                                     | 2,967 No. of compressors <sup>c,g,i,1</sup>              | 0.002                 | 0.1                  |
| Large Compressors   | 332 scfd CO <sub>2</sub> /compressor <sup>e</sup>                                       | - No. of largecompr. <sup>e,3</sup>                      | 0.000                 | 0.0                  |
| Sales Areas   | 2.096 scf CO <sub>2</sub> /loading <sup>e</sup><br>scfd of CO <sub>2</sub> /mile of     | 2,648,868 Loadings/year <sup>c,1</sup>                   | 0.006                 | 0.3                  |
| Pipelines   | NE pipeline   | 14,571 Miles of gathering line <sup>o,2</sup>            | NE                    | NE                   |

| Activity/Equipment                    | 2014 EPA Inventory Values                        |   |                    |                   |
|---------------------------------------|--|---|--------------------|-------------------|
|                                       | Emission Factor                                  | Activity Factor                                 | Emissions (Bcf/yr) | Emissions (kt/yr) |
| Well Drilling                         | scfd of CO <sub>2</sub> /oil well<br>NE drilled  | 17,774 No. of oil wells drilled <sup>c,2</sup>  | NE                 | NE                |
| Battery Pumps                         | 0.012 scfd of CO <sub>2</sub> /pump <sup>m</sup> | 269,480 No. of battery pumps <sup>g,e,1</sup>   | 0.001              | 0.1               |
| <b>Process Upset Emissions</b>        |  |   | <b>0.003</b>       | <b>0.2</b>        |
| Pressure Relief Valves                | 1.794 scf/yr/PR valve <sup>h</sup>               | 265,554 No. of PR valves <sup>c,e,1</sup>       | 0.000*             | 0.0*              |
| Well Blowouts Onshore                 | 0.051 MMscf/blowout <sup>e</sup>                 | 59 No. of blowouts/yr <sup>c,e,1</sup>          | 0.003              | 0.2               |
| <b>Refining Emissions<sup>1</sup></b> |  |   | <b>55.341</b>      | <b>2,926.7</b>    |
| Asphalt Blowing                       | 0.020 MT CO <sub>2</sub> /Mbbbl <sup>q</sup>     | 5,784,637 Mbbbl/yearrefineryfeed <sup>p</sup>   | 2.187              | 115.6             |
| Flaring                               | 0.477 MT CO <sub>2</sub> /Mbbbl <sup>q</sup>     | 5,784,637 Mbbbl/year refinery feed <sup>p</sup> | 52.196             | 2,760.4           |
| Process Vents                         | 0.009 MT CO <sub>2</sub> /Mbbbl <sup>q</sup>     | 5,784,637 Mbbbl/year refinery feed <sup>p</sup> | 0.958              | 50.7              |
| <b>Total</b>                          |  |   | <b>67.451</b>      | <b>3,567.1</b>    |

\* Emissions are not actually 0, but too small to show at this level of precision.

NE - Not estimated for lack of data

<sup>a</sup> TankCALC

<sup>b</sup> EPA / ICF International (1999)

<sup>c</sup> EIA Monthly Energy Review

<sup>d</sup> IOGCC Marginal Wells Report

<sup>e</sup> Consensus of Industrial Review Panel

<sup>f</sup> Expert Judgment

<sup>g</sup> EIA Annual Energy Review

<sup>h</sup> GRI / EPA (1996)

<sup>i</sup> Radian (1996)

<sup>j</sup> CAPP (1992)

<sup>k</sup> Adapted from the GOADS 2011 Study by ERG (2015)

<sup>l</sup> BOEM

<sup>m</sup> API (1996)

<sup>n</sup> EPA, AP 42 Compilation of Air Pollutant Emission Factors

<sup>o</sup> OGJ Petroleum Economics Issue

<sup>p</sup> EIA Petroleum Supply Annual, Volume 1

<sup>q</sup> GHGRP data

<sup>r</sup> Calculated using CH<sub>4</sub> emission factor and CO<sub>2</sub> content of gas.

<sup>s</sup> DrillingInfo Data

<sup>1</sup> The methodology for year 2014 is to use GHGRP emissions data as-reported (rather than an EFxAF approach, per se). The emission factors in this table were populated by dividing 2014 emissions by 2014 refinery feed rate.

Note: Energy use CO<sub>2</sub> emissions not estimated to avoid double counting with fossil fuel combustion

**Table A-133: Summary of CO<sub>2</sub> Emissions from Petroleum Systems (kt)**

| Activity                           | 1990         | 1995         | 2000         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Production Field Operations</b> | <b>391</b>   | <b>362</b>   | <b>360</b>   | <b>350</b>   | <b>343</b>   | <b>365</b>   | <b>379</b>   | <b>395</b>   | <b>473</b>   | <b>550</b>   | <b>640</b>   |
| Pneumatic controller venting       | 42           | 46           | 58           | 70           | 72           | 73           | 74           | 75           | 74           | 84           | 87           |
| Tank venting                       | 328          | 296          | 281          | 253          | 243          | 265          | 276          | 288          | 365          | 432          | 519          |
| Misc. venting & fugitives          | 17           | 17           | 18           | 23           | 25           | 24           | 26           | 28           | 30           | 30           | 30           |
| Wellhead fugitives                 | 3            | 3            | 3            | 3            | 3            | 3            | 3            | 3            | 3            | 3            | 3            |
| <b>Refining</b>                    | <b>3,162</b> | <b>3,295</b> | <b>3,562</b> | <b>3,574</b> | <b>3,463</b> | <b>3,380</b> | <b>3,775</b> | <b>3,797</b> | <b>3,404</b> | <b>3,143</b> | <b>2,927</b> |
| Asphalt Blowing                    | 95           | 99           | 107          | 107          | 104          | 101          | 97           | 84           | 117          | 125          | 116          |
| Flaring                            | 2,993        | 3,119        | 3,372        | 3,383        | 3,279        | 3,200        | 3,626        | 3,630        | 3,228        | 2,882        | 2,760        |
| Process Vents                      | 74           | 77           | 83           | 84           | 81           | 79           | 52           | 83           | 58           | 137          | 51           |
| <b>Total</b>                       | <b>3,553</b> | <b>3,657</b> | <b>3,923</b> | <b>3,924</b> | <b>3,806</b> | <b>3,745</b> | <b>4,154</b> | <b>4,192</b> | <b>3,876</b> | <b>3,693</b> | <b>3,567</b> |

## References

- Allen et al. (2014) Methane Emissions from Process Equipment at Natural Gas Production Sites in the United States: Pneumatic Controllers. ES&T. December 9, 2014. Available online at: <<http://pubs.acs.org/doi/pdf/10.1021/es5040156>>.
- API (2009) Compendium of Greenhouse Gas Emissions Methodologies for the Oil and Gas Industry. American Petroleum Institute. Austin, TX, August 2009.
- BOEM (2011a) OCS Platform Activity. Bureau of Ocean Energy Management, U.S. Department of Interior. Available online at: <[http://www.boem.gov/uploadedFiles/BOEM/Newsroom/Offshore\\_Stats\\_and\\_Facts/Gulf\\_of\\_Mexico\\_Region/OCSPlatformActivity.pdf](http://www.boem.gov/uploadedFiles/BOEM/Newsroom/Offshore_Stats_and_Facts/Gulf_of_Mexico_Region/OCSPlatformActivity.pdf)>.
- BOEM (2011b) Platform Information and Data. Bureau of Ocean Energy Management, U.S. Department of Interior. Available online at: <[https://www.data.boem.gov/homepg/data\\_center/platform/platform.asp](https://www.data.boem.gov/homepg/data_center/platform/platform.asp)>.
- BOEM (2011c) Pacific OCS Region. Bureau of Ocean Energy Management, U.S. Department of Interior. Available online at: <[http://www.data.boem.gov/homepg/data\\_center/platform/PacificFreePlat.asp](http://www.data.boem.gov/homepg/data_center/platform/PacificFreePlat.asp)>.
- BOEM (2014) Year 2011 Gulfwide Emission Inventory Study. Bureau of Ocean Energy Management, U.S. Department of Interior. OCS Study BOEM 2014-666. Available online at: <<http://www.data.boem.gov/PI/PDFImages/ESPIS/5/5440.pdf>>
- DrillingInfo (2015) March 2015 Download. DI Desktop® DrillingInfo, Inc.
- EIA (1990 through 2015) Refinery Capacity Report. Energy Information Administration, U.S. Department of Energy. Washington, DC. Available online at: <<http://www.eia.gov/petroleum/refinerycapacity/>>.
- EIA (1995 through 2015a) Monthly Energy Review. Energy Information Administration, U.S. Department of Energy. Washington, DC. Available online at: <<http://www.eia.gov/totalenergy/data/monthly/index.cfm>>.
- EIA (1995 through 2015b) Petroleum Supply Annual. Volume 1. U.S Department of Energy Washington, DC. Available online at: <<http://www.eia.gov/petroleum/supply/annual/volume1/>>.
- EPA (2016a) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas and Petroleum Production Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2016b) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas Gathering and Boosting Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2016c) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas Transmission and Storage Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2016d) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas Distribution Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2015a) Greenhouse Gas Reporting Program. Environmental Protection Agency. Data reported as of August 16, 2015.
- EPA (2015b) Background Technical Support Document for the Proposed New Source Performance Standards 40 CFR Part 60, subpart OOOOa. Available online at: <https://www.regulations.gov/#!documentDetail;D=EPA-HQ-OAR-2010-0505-5021>
- EPA (2015c) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2013: Update to Offshore Oil and Gas Platforms Emissions Estimate. Available online at: <http://www.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>.
- EPA (2015d) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2013: Update to Refineries Emissions Estimate. Available online at: <<http://www.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.

EPA (2005) Incorporating the Mineral Management Service Gulfwide Offshore Activities Data System (GOADS) 2000 data into the methane emissions inventories. Prepared by ICF International. U.S. Environmental Protection Agency. 2005.

EPA (1999a) Estimates of Methane Emissions from the U.S. Oil Industry (Draft Report). Prepared by ICF International. Office of Air and Radiation, U.S. Environmental Protection Agency. October 1999.

EPA (1999b) Methane Emissions from the U.S. Petroleum Industry. Prepared by Radian International. U.S. Environmental Protection Agency. February 1999.

EPA/GRI (1996a) Methane Emissions from the Natural Gas Industry, V7: Blow and Purge Activities. Prepared by Radian. U.S. Environmental Protection Agency. April 1996.

EPA/GRI (1996b) Methane Emissions from the Natural Gas Industry, V11: Compressor Driver Exhaust. Prepared by Radian. U.S. Environmental Protection Agency. April 1996.

EPA/GRI (1996c) Methane Emissions from the Natural Gas Industry, V12: Pneumatic Devices. Prepared by Radian. U.S. Environmental Protection Agency. April 1996.

EPA/GRI (1996d) Methane Emissions from the Natural Gas Industry, V13: Chemical Injection Pumps. Prepared by Radian. U.S. Environmental Protection Agency. April 1996.

HPDI (2011) Production and Permit Data, October 2009.

IOGCC (2012) Marginal Wells: fuel for economic growth 2012 Report. Interstate Oil & Gas Compact Commission. Available online at: < [http://iogcc.ok.gov/Websites/iogcc/images/MarginalWell/MW\\_2012\\_report.pdf](http://iogcc.ok.gov/Websites/iogcc/images/MarginalWell/MW_2012_report.pdf) >.

IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change. [H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.)]. Hayama, Kanagawa, Japan.

IPCC (2007) Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. [Pachauri, R.K and Reisinger, A. (eds.)]; IPCC, Geneva, Switzerland.

OGJ (2015) Oil and Gas Journal 1990-2014. Oil Pipelines, Vol 113, Issue 9, September 7, 2015.

United States Army Corps of Engineers (1995 through 2015) Waterborne Commerce of the United States, Part 5: National Summaries. U.S. Army Corps of Engineers. Washington, DC, July 20, 2015. Available online at: <<http://www.navigationdatacenter.us/wcsc/pdf/wcusnatl3.pdf>>.

### 3.6. Methodology for Estimating CH<sub>4</sub> and CO<sub>2</sub> Emissions from Natural Gas Systems

As described in the main body text on Natural Gas Systems, the Inventory methodology involves the calculation of CH<sub>4</sub> and CO<sub>2</sub> emissions for over 100 emissions sources, and then the summation of emissions for each natural gas sector stage.

#### Step 1: Calculate Potential Methane (or Net Emissions)

##### Potential methane factors and emission factors

The primary basis for potential CH<sub>4</sub> factors and emission factors for non-combustion-related CO<sub>2</sub> emissions from the U.S. natural gas industry is a detailed study by the Gas Research Institute and EPA (EPA/GRI 1996). The EPA/GRI study developed over 80 CH<sub>4</sub> emission factors to characterize emissions from the various components within the operating stages of the U.S. natural gas system. Since the time of this study, practices and technologies have changed. While this study still represents best available data in many cases, using these emission factors alone to represent actual emissions without adjusting for emissions controls would in many cases overestimate emissions. For this reason, “potential methane” is calculated using the data, and then recent data on voluntary and regulatory emission reduction activities (Step 3) is deducted to calculate actual emissions. See Section 3.7 of the main document on Natural Gas Systems for more information.

For certain CH<sub>4</sub> emissions sources, new data and information allows for net emissions to be calculated directly: gas well completions and workovers with hydraulic fracturing, liquids unloading, condensate storage tanks, centrifugal compressors, pneumatic controllers, chemical injection pumps, transmission and storage station fugitives and compressors, M&R stations, and distribution pipeline leaks. For these sources, EPA used emissions factors that directly reflect the use of control technologies. For gas well completions and workovers with hydraulic fracturing, separate emissions estimates were developed for hydraulically fractured completions and workovers that vent, flared hydraulic fracturing completions and workovers, hydraulic fracturing completions and workovers with reduced emissions completions (RECs), and hydraulic fracturing completions and workovers with RECs that flare. For liquids unloading, separate emissions estimates were developed for wells with plunger lifts and wells without plunger lifts. Likewise, for condensate tanks, emissions estimates were developed for tanks with and without control devices. For pneumatic controllers, separate estimates are developed for low bleed, high bleed, and intermittent controllers. Chemical injection pumps estimates are calculated with an emission factor developed with GHGRP data, which is based on the previous GRI/EPA factor but takes into account operating hours.

For potential CH<sub>4</sub> factors and emission factors used in the Inventory, see Table A-134 to Table A-138. Methane compositions from GTI 2001 are adjusted year to year using gross production for National Energy Modeling System (NEMS) oil and gas supply module regions from the EIA. These adjusted region-specific annual CH<sub>4</sub> compositions are presented in Table A-139 (for general sources), Table A-140 (for gas wells without hydraulic fracturing), and Table A-141 (for gas wells with hydraulic fracturing). Therefore, emission factors may vary from year to year due to slight changes in the CH<sub>4</sub> composition between each NEMS oil and gas supply module region.

##### *1990-2014 Inventory updates to emission factors*

Summary information for emission factors for sources with updates in this year’s inventory is below. The details are presented in four memoranda addressing production, gathering and boosting, transportation and storage, and distribution (See “Revisions to Natural Gas and Petroleum Production Emissions”, “Revisions to Natural Gas Gathering and Boosting Emissions”, “Revisions to Natural Gas Transmission and Storage Emissions”, and “Revisions to Natural Gas Distribution Emissions” [EPA 2016a through 2016d]).<sup>64</sup>

For the production segment, bleed type-specific (i.e., continuous high bleed, continuous low bleed, and intermittent bleed) emission factors for pneumatic controllers are based on GHGRP data (used for the full time series). Emission factors for gathering and boosting stations based on Marchese et al. are used for the full time series.

For the transmission and storage segment, station-level emission factors for transmission and storage stations and factors for compressors are based on the Zimmerle et al. study and used for 2011 to 2014. Emission factors for transmission and storage pneumatic controllers are from GHGRP and are used for the full time series.

For the distribution segment, emission factors for M&R stations and for emissions per leak for pipeline leaks are from the Lamb et al. study (used for 2011 to 2014). Emission factors for residential customer meters and

<sup>64</sup> See <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.

commercial/industrial customer meters are from the 2009 GTI study and 2011 Clearstone study and are used for the full time series.

In order to create time series consistency for emission factors between earlier years' estimates (1990 to 1992) that generally rely on data from GRI/EPA 1996 and the most recent years' estimates (2011 to 2014) that were calculated using data from studies or GHGRP, linear interpolation between the data endpoints of 1992 (GRI/EPA) and 2011 (GHGRP) was typically used for calculations. Exceptions to the use of linear interpolation include gathering and boosting stations and residential and commercial/industry customer meters.

#### Activity Data

Activity data were taken from the following sources: DrillingInfo, Inc. (DrillingInfo 2015); American Gas Association (AGA 1991–1998); Bureau of Ocean Energy Management, Regulation and Enforcement (previous Minerals and Management Service) (BOEMRE 2011a, 2011b, 2011c, 2011d); Monthly Energy Review (EIA 2012f, 2012g, 2012h, 2011a, 2011b, 2011c, 2011d); Natural Gas Liquids Reserves Report (EIA 2005); Natural Gas Monthly (EIA 2012c, 2012d, 2012e, 2013a, 2013b, 2013c); the Natural Gas STAR Program annual emissions savings (EPA 2012a, 2013c); Oil and Gas Journal (OGJ 1997–2015); Pipeline and Hazardous Materials Safety Administration (PHMSA 2015); Federal Energy Regulatory Commission (FERC 2014); GHGRP data for natural gas systems (40 CFR 98, subpart W); and other Energy Information Administration publications (EIA 2001, 2004, 2010, 2011, 2012i, 2015); (EPA 1999); Wyoming Oil and Gas Conservation Commission (Wyoming 2015); and the Alabama State Oil and Gas Board (Alabama 2015). Activity data are presented in Table A-134 through Table A-138.

For a few sources, recent direct activity data were not available. For these sources, either 2013 data were used as proxy for 2014 data or a set of industry activity data drivers was developed and was used to update activity data. Drivers include statistics on gas production, number of wells, system throughput, miles of various kinds of pipe, and other statistics that characterize the changes in the U.S. natural gas system infrastructure and operations. The key activity drivers are presented in Table A-142.

EPA used DI Desktop, a production database maintained by DrillingInfo, Inc. (DrillingInfo 2015), covering U.S. oil and natural gas wells to populate activity data for non-associated gas wells, oil wells (in petroleum systems), gas wells with hydraulic fracturing, and completions with hydraulic fracturing. EPA queried DI Desktop for relevant data on an individual well basis—including location, natural gas and liquids (i.e., oil and condensate) production by year, drill type (e.g., horizontal or vertical), and date of completion or first production. Non-associated gas wells were classified as any well within DI Desktop that had non-zero gas production in a given year, and with a gas-to-oil ratio (GOR) of greater than 100 mcf/bbl in that year. Oil wells were classified as any well that had non-zero liquids production in a given year, and with a GOR of less than or equal to 100 mcf/bbl in that year. Gas wells with hydraulic fracturing were assumed to be the subset of the non-associated gas wells that were horizontally drilled and/or located in an unconventional formation (i.e., shale, tight sands, or coalbed). Unconventional formations were identified based on well basin, reservoir, and field data reported in DI Desktop referenced against a formation type crosswalk developed by EIA (EIA 2012a).

For 1990 through 2010, gas well completions with hydraulic fracturing were identified as a subset of the gas wells with hydraulic fracturing that had a date of completion or first production in the specified year. To calculate workovers for 1990 through 2010, EPA applied a refracture rate of 1 percent (i.e., 1 percent of all wells with hydraulic fracturing are assumed to be refractured in a given year) to the total counts of wells with hydraulic fracturing from the DrillingInfo data. For 2011 through 2014, EPA used GHGRP data for the total number of well completions and workovers. The GHGRP data represents a subset of the national completions and workovers, due to the reporting threshold, and therefore using this data without scaling it up to national level results in an underestimate. However, because EPA's GHGRP counts of completions and workovers were higher than national counts of completions and workovers, obtained using DI Desktop data, EPA directly used the GHGRP data for completions and workovers for 2011 through 2014.

EPA calculated the percentage of gas well completions and workovers with hydraulic fracturing in the each of the four control categories using 2011 through 2014 Subpart W data. EPA assumed 0 percent RECs use from 1990 through 2000, used GHGRP RECs percentage for 2011 through 2014, and then used linear interpolation between the 2000 and 2011 percentages. For flaring, EPA used an assumption of 10 percent (the average of the percent of completions and workovers that were flared in 2011 through 2013 GHGRP data) flaring from 1990 through 2010 to recognize that some flaring has occurred over that time period. For 2011 through 2014, EPA used the GHGRP data on flaring.

Summary information for activity data for sources with updates in this year's inventory is below. The details are presented in four memoranda addressing production, gathering and boosting, transportation and storage, and distribution (see "Revisions to Natural Gas and Petroleum Production Emissions", "Revisions to Natural Gas Gathering and Boosting

Emissions”, “Revisions to Natural Gas Transmission and Storage Emissions”, and “Revisions to Natural Gas Distribution Emissions” [EPA 2016a through 2016d)].<sup>65</sup>

For the production segment, GHGRP-based activity factors (i.e., counts per gas well) were used for the years 2011 to 2014 for in-line heaters, separators, dehydrators, compressors, meters/piping, pneumatic pumps, and pneumatic controllers. In addition, for years 2011 to 2014, bleed type-specific (i.e., continuous high bleed, continuous low bleed, and intermittent bleed) activity data for pneumatic controllers were developed from GHGRP.

For the transmission and storage segment, activity factors (transmission stations; storage stations per storage field) are based on Zimmerle et al. study and applied for 2011 to 2014, reciprocating and centrifugal compressor activity data and compressor seal types (i.e., wet versus dry seals) are based on GHGRP and Zimmerle et al. study data for 2011-2014, and activity factors (i.e., controllers per station) for transmission and storage pneumatic controllers based on GHGRP and Zimmerle et al. study for 2011 to 2014.

For the distribution segment, activity factors for M&R stations for 2011 to 2014 are based on GHGRP and Lamb et al. study. Activity data for meters are from EIA for the full time series. Activity data for pipeline blowdowns and mishaps/dig-ins are from PHMSA for the full time series.

For sources where new activity data was determined to be applicable for only recent years of the time series, in order to create time series consistency for activity data between earlier years’ estimates (1990 to 1992) that generally rely on data from GRI/EPA 1996 and the most recent years’ estimates (2011 to 2014) that were calculated using data from studies or GHGRP, linear interpolation between the data endpoints of 1992 (GRI/EPA) and 2011 (GHGRP) was typically used for calculations. Exceptions to the use of linear interpolation include gathering and boosting stations and residential and commercial/industry customer meters.

## **Step 2: Compile Reductions Data**

The emissions calculated in Step 1 above represent expected emissions from an activity in the absence of emissions controls (with the exceptions of emission sources that use a “net” approach such as pneumatic controllers, gas well completions and workovers with hydraulic fracturing, liquids unloading, centrifugal compressors, condensate tanks, distribution M&R stations, distribution pipelines, etc. as noted above), and do not take into account any use of technologies or practices that reduce emissions. To take into account use of such technologies, data were collected on voluntary and regulatory reductions. Voluntary reductions included in the Inventory were those reported to Gas STAR for activities such as replacing a high bleed pneumatic controllers with a low bleed controller and replacing wet seals with dry seals at reciprocating compressors. Regulatory actions reducing emissions include National Emission Standards for Hazardous Air Pollutants (NESHAP) regulations for dehydrator vents and condensate tanks.

### Voluntary reductions

Industry partners report CH<sub>4</sub> emission reductions by project to the Natural Gas STAR Program. The reductions from the implementation of specific technologies and practices (e.g., vapor recovery units,) are calculated by the reporting partners using actual measurement data or equipment-specific emission factors. Natural Gas STAR Partners do not report reductions when they are required due to regulation. The reductions undergo quality assurance and quality control checks to identify errors, inconsistencies, or irregular data before being incorporated into the Inventory. In general, the Inventory uses aggregated Gas STAR reductions by natural gas system stage (i.e., production, processing, transmission and storage, and distribution). However, aggregate emissions reductions data by Gas STAR technology are provided for several sources, as shown in Table A-143. For those sources, EPA has also used data on potential emissions, and the Gas STAR data on reductions, to calculate net emissions, as shown in Table A-149 (along with other sources calculated with net emission factors). Many of the activities reported to Gas STAR are cross-cutting and apply to more than one emissions source and therefore cannot be assigned to one emissions source, but instead are included in the “other” category. For Inventory sources with emission factors that already take into account the use of control technologies (i.e., “net” emission sources) Natural Gas STAR reported reductions for those activities are not incorporated into the Inventory, as this would double count reductions. CH<sub>4</sub> emission reductions from the Natural Gas STAR Program are summarized in Table A-143.

For sources where net factor approaches are used, the use of data from the GHGRP and recent studies to revise certain emission factors as discussed above obviated the need to apply Gas STAR reductions data for these sources (i.e., the calculated emissions were already net emissions, instead of potential emissions).

---

<sup>65</sup> See <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.

There are significant Gas STAR reductions in the production segment that are not classified as applicable to specific emission sources (“other voluntary reductions” are 18 MMT CO<sub>2</sub> eq. of CH<sub>4</sub> in year 2014). As many sources in production are now calculated with net factor approaches, to address potential double-counting of reductions, a scaling factor was applied to the “other voluntary reductions” to reduce this reported amount based an estimate of the fraction of those reductions that occur in the sources that are now calculated using net emissions approaches. This fraction was developed by dividing the net emissions from sources with net approaches, by the total production segment emissions (without deducting the Gas STAR reductions). The result for 2014, is that around 50 percent of the reductions were estimated to occur in sources for which net emissions are now calculated, which yields an adjusted other reduction number of 9 MMT CO<sub>2</sub> Eq.

#### Federal regulations.

The 1990 Clean Air Act (CAA) sets limits on the amount of hazardous air pollutants (HAPs) that can be emitted in the United States. The NESHAP regulations set the standards to limit emissions of HAPs. The emission sources are required to use the Maximum Achievable Control Technology (MACT), giving the operators flexibility to choose the type of control measure(s) to implement. In regards to the oil and natural gas industry, the NESHAP regulation addresses HAPs from the oil and natural gas production sectors and the natural gas transmission and storage sectors of the industry. Though the regulation deals specifically with HAPs reductions, methane emissions are also incidentally reduced.

The NESHAP regulation requires that glycol dehydration unit vents and storage tanks that have HAP emissions and exceed a gas throughput and liquids throughput threshold, respectively, be connected to a closed loop emission control system that reduces emissions by 95 percent. Also, gas processing plants exceeding the threshold natural gas throughput limit are required to routinely implement Leak Detection and Repair (LDAR) programs. The emissions reductions achieved as a result of NESHAP regulations were calculated using data provided in the Federal Register Background Information Document (BID) for this regulation. The BID provides the levels of control measures in place before the enactment of regulation. The emissions reductions were estimated by analyzing the portion of the industry without control measures already in place that would be impacted by the regulation. CH<sub>4</sub> emission reductions from federal regulations, such as NESHAP, are summarized in Table A-144. In addition to the NESHAP applicable to natural gas, the Inventory reflects the 2012 New Source Performance Standards (NSPS) subpart OOOO for oil and gas, through the use of a net factor approach that captures shifts to lower emitting technologies. By separating gas well completions and workovers with hydraulic fracturing into four categories and developing control technology-specific methane emission factors for each category, EPA is implicitly accounting for subpart OOOO reductions from hydraulically fractured gas wells. This is also the case for high bleed pneumatic controllers in the production segment.

#### **Step 3: Calculate Net Methane and Carbon Dioxide Emissions for Each Activity for Each Year**

Annual CH<sub>4</sub> emissions and CO<sub>2</sub> emissions for each source were estimated by multiplying the activity data for each year by the corresponding emission factor. These annual emissions for each activity were then summed to estimate the total annual CH<sub>4</sub> and CO<sub>2</sub> emissions, respectively. As a final step, the relevant Natural Gas STAR reductions data from each segment is summed for each year and deducted from the total emissions to estimate net CH<sub>4</sub> emissions for the Inventory.

The same procedure for estimating CH<sub>4</sub> emissions holds true for estimating non-energy related CO<sub>2</sub> emissions, except the emission estimates are not adjusted for reductions due to the Natural Gas STAR program or regulations.

Produced natural gas is composed of primarily CH<sub>4</sub>, but as shown in Table A-150, the natural gas contains, in some cases, as much as 8 percent CO<sub>2</sub>. The same vented and fugitive natural gas that led to CH<sub>4</sub> emissions also contains a certain volume of CO<sub>2</sub>. Accordingly, the CO<sub>2</sub> emissions for each sector can be estimated using the same activity data for these vented and fugitive sources. The emission factors used to estimate CH<sub>4</sub> were also used to calculate non-combustion CO<sub>2</sub> emissions. The Gas Technology Institute’s (GTI, formerly GRI) Unconventional Natural Gas and Gas Composition Databases (GTI 2001) were used to adapt the CH<sub>4</sub> emission factors into non-combustion related CO<sub>2</sub> emission factors. Additional information about CO<sub>2</sub> content in transmission quality natural gas was obtained from numerous U.S. transmission companies to help further develop the non-combustion CO<sub>2</sub> emission factors. For the CO<sub>2</sub> content used to develop CO<sub>2</sub> emission factors from CH<sub>4</sub> potential factors, see Table A-150. The detailed source emission estimates for CH<sub>4</sub> and CO<sub>2</sub> from the production sector are presented in Table A-145 and Table A-154, respectively.

In the processing sector, the CO<sub>2</sub> content of the natural gas remains the same as the CO<sub>2</sub> content in the production sector for the equipment upstream of the acid gas removal unit because produced natural gas is usually only minimally treated after being produced and then transported to natural gas processing plants via gathering pipelines. The CO<sub>2</sub> content in gas for the remaining equipment that is downstream of the acid gas removal is the same as in pipeline quality gas. The EPA/GRI study estimates the average CH<sub>4</sub> content of natural gas in the processing sector to be 87 percent CH<sub>4</sub>. Consequently, the processing sector CO<sub>2</sub> emission factors were developed using CH<sub>4</sub> emission factors, proportioned to

reflect the CO<sub>2</sub> content of either produced natural gas or pipeline quality gas using the same methodology as the production sector. The detailed source emission estimates for CH<sub>4</sub> and CO<sub>2</sub> from the processing sector are presented in Table A-146 and Table A-152, respectively.

For the transmission sector, CO<sub>2</sub> content in natural gas transmission pipelines was estimated for the top 20 transmission pipeline companies in the United States (separate analyses identified the top 20 companies based on gas throughput and total pipeline miles). The weighted average CO<sub>2</sub> content in the transmission pipeline quality gas in both cases—total gas throughput and total miles of pipeline—was estimated to be about 1 percent. To estimate the CO<sub>2</sub> emissions for the transmission sector, the CH<sub>4</sub> emission factors were proportioned from the 93.4 percent CH<sub>4</sub> reported in EPA/GRI (1996) to reflect the 1 percent CO<sub>2</sub> content found in transmission quality natural gas. The detailed source emissions estimates for CH<sub>4</sub> and CO<sub>2</sub> for the transmission sector are presented in Table A-147 and Table A-153, respectively.

The natural gas in the distribution sector of the system has the same characteristics as the natural gas in the transmission sector. The CH<sub>4</sub> content (93.4 percent) and CO<sub>2</sub> content (1 percent) are identical to transmission segment contents due to the absence of any further treatment between sector boundaries. Thus, the CH<sub>4</sub> emissions factors were converted to CO<sub>2</sub> emission factors using the same methodology as discussed for the transmission sector. The detailed source emission estimates for CH<sub>4</sub> and CO<sub>2</sub> for the distribution sector are presented in Table A-148 and Table A-154, respectively.

Three exceptions to this methodology are CO<sub>2</sub> emissions from flares, CO<sub>2</sub> from acid gas removal units, and CO<sub>2</sub> from condensate tanks. In the case of flare emissions, a direct CO<sub>2</sub> emission factor from EIA (1996) was used. This emission factor was applied to the portion of offshore gas that is not vented and all of the gas reported as vented and flared onshore by EIA, including associated gas. The amount of CO<sub>2</sub> emissions from an acid gas unit in a processing plant is equal to the difference in CO<sub>2</sub> concentrations between produced natural gas and pipeline quality gas applied to the throughput of the plant. This methodology was applied to the national gas throughput using national average CO<sub>2</sub> concentrations in produced gas (3.45 percent) and transmission quality gas (1 percent). Data were unavailable to use annual values for CO<sub>2</sub> concentration. For condensate tanks, a series of E&P Tank (EPA 1999) simulations provide the total CO<sub>2</sub> vented per barrel of condensate throughput from fixed roof tank flash gas for condensate gravities of API 45 degree and higher. The ratios of emissions to throughput were used to estimate the CO<sub>2</sub> emission factor for condensate passing through fixed roof tanks.

Table A-134 through Table A-138 display the 2012 activity data, CH<sub>4</sub> emission factors, and calculated potential CH<sub>4</sub> emissions for each stage.

The tables provide references for emission factors and activity data in footnotes (the lettered footnotes). The tables also provide information on which method was used for supplying activity data for 2014 (the numbered footnotes).

Key to table notations on methods for supplying activity data for 2014 for all tables:

1. Ratios relating other factors for which activity data are available.
2. Activity data for 2014 available from source.
3. Activity data were held constant from 1990 through 2014 based on EPA (1999).

**Table A-134: 2014 Data and Calculated CH<sub>4</sub> Potential Emissions (Mg) for the Natural Gas Production Stage**

| Activity   | 2014 EPA Inventory Values                           |  |   |
|--|---|--|---|
|  | National Activity Data                              | National Emission Factor or Range of Regional Values (Potential) <sup>aa</sup> | Calculated Potential (Mg) <sup>bb</sup> |
| <b>Gas Wells</b>   |   |  |   |
| Associated Gas Wells   | 503,873 wells <sup>a,1,cc</sup>                     | NA <sup>dd</sup>   | 0.0                                     |
| Non-associated Gas Wells (less fractured wells)                      | 205,363 wells <sup>a,1</sup>                        | 7.43-42.49 scfd/well <sup>b</sup>  | 17,754.5                                |
| Gas Wells with Hydraulic Fracturing                                  | 250,777 wells <sup>a,1</sup>                        | 7.59-42.49 scfd/well <sup>b</sup>  | 35,085.1                                |
| <b>Well Pad Equipment</b>  |   |  |   |
| Heaters  | 99,038 heaters <sup>c,d,2</sup>                     | 14.87-67.29 scfd/heater <sup>b</sup>   | 23,953.4                                |
| Separators   | 306,377 separators <sup>c,d,2</sup>                 | 0.94-142.27 scfd/separator <sup>b</sup>  | 118,591.2                               |
| Dehydrators  | 17,126 dehydrators <sup>c,d,2</sup>                 | 23.18-106.25 scfd/dehydrator <sup>b</sup>                                      | 8,417.5                                 |
| Meters/Piping  | 523,885 meters <sup>c,d,2</sup>                     | 9.43-61.68 scfd/meter <sup>b</sup>   | 107,173.2                               |
| Compressors  | 48,518 compressors <sup>c,d,2</sup>                 | 263.85-312.19 scfd/compressor <sup>b</sup>                                     | 96,170.6                                |
| <b>Gathering and Boosting</b>  |   |  |   |
| Gathering and Boosting Stations*                                     | 4,999 stations <sup>e,2</sup>                       | 53,066 scfd CH <sub>4</sub> /station <sup>e</sup>                              | 1,864,870.3                             |
| Pipeline Leaks   | 431,051 miles <sup>f,2</sup>                        | 52.38-61.97 scfd/mile <sup>b</sup>   | 169,701.4                               |
| <b>Drilling, Well Completion, and Well Workover</b>                  |   |  |   |
| Gas Well Completions without Hydraulic Fracturing                    |   |  |   |
| Fracturing   | 767 completions/year <sup>g</sup>                   | 707.23-854.65 scf/completion <sup>b</sup>                                      | 11.3                                    |
| Gas Well Workovers without Hydraulic Fracturing                      | 8,933 workovers/year <sup>a,2</sup>                 | 2,367.7-2,861.3 scf/workover <sup>b</sup>                                      | 445.8                                   |
| Hydraulic Fracturing Completions and Workovers that vent*            | 1,791 completions and workovers/year <sup>c</sup>   | MT/(completion or workover) <sup>h</sup>                                       | 65,940.7                                |
| Flared Hydraulic Fracturing Completions and Workovers*               | 548 completions and workovers/year <sup>c</sup>     | MT/(completion or workover) <sup>h</sup>                                       | 2,690.6                                 |
| Hydraulic Fracturing Completions and Workovers with RECs*            | 1,043 completions and workovers/year <sup>c</sup>   | MT/(completion or workover) <sup>h</sup>                                       | 3,379.6                                 |
| Hydraulic Fracturing Completions and Workovers with RECs that flare* | 1,979 completions and workovers/year <sup>c</sup>   | MT/(completion or workover) <sup>h</sup>                                       | 9,653.4                                 |
| Well Drilling  | 18,837 wells <sup>i,1</sup>                         | 2,505.9-2,965.0 scf/well <sup>i</sup>  | 971.6                                   |
| <b>Normal Operations</b>   |   |  |   |
| Pneumatic Device Vents*  | 834,919 controllers <sup>c,d,2</sup>                | 176.74-209.12 scfd/device <sup>c,d</sup>                                       | 1,105,119.0                             |
| Pneumatic Device Vents - Low Bleed (LB)                              | 226,280 controllers <sup>c,d,2</sup>                | 22.52-26.64 scfd/device <sup>c,d</sup>   | Aggregated                              |
| Pneumatic Device Vents - High Bleed (HB)                             | 29,006 controllers <sup>c,d,2</sup>                 | 612.66-724.91 scfd/device <sup>c,d</sup>                                       | Aggregated                              |
| Pneumatic Device Vents - Intermittent Bleed (IB)                     | 579,633 controllers <sup>c,d,2</sup>                | 215.13-254.55 scfd/device <sup>c,d</sup>                                       | Aggregated                              |
| Chemical Injection Pumps*  | 83,249 active pumps <sup>c,d,2</sup>                | 208.89-252.30 scfd/pump <sup>c,d</sup>   | 128,876.5                               |
| Kimray Pumps   | 5,012,753 MMscf/yr <sup>b,2</sup>                   | 977.5-1,156.6 scf/MMscf <sup>b</sup>   | 100,857.2                               |
| Dehydrator Vents   | 5,625,985 MMscf/yr <sup>b,2</sup>                   | 271.58-321.34 scf/MMscf <sup>b</sup>   | 31,448.3                                |
| <b>Condensate Tank Vents</b>   |   |  |   |
| Condensate Tanks without Control Devices                             | 139 MMbbl/yr <sup>k,1</sup>                         | 21.87-302.75 scf/bbl <sup>l</sup>  | 253,092.6                               |
| Condensate Tanks with Control Devices*                               | 139 MMbbl/yr <sup>k,1</sup>                         | 4.37-60.55 scf/bbl <sup>l</sup>  | 50,618.5                                |
| <b>Compressor Exhaust Vented</b>                                     |   |  |   |
| Gas Engines  | 51,648 MMHPhr <sup>b,2</sup>                        | 0.237-0.280 scf/HPhr <sup>b</sup>  | 249,756.3                               |
| <b>Well Clean Ups</b>  |   |  |   |
| Liquids Unloading with Plunger Lifts*                                | 22,477 venting wells <sup>a,m,2</sup>               | 2,856-1,137,406 scfy/venting well <sup>m</sup>                                 | 112,568.8                               |
| Liquids Unloading without Plunger Lifts*                             | 37,912 venting wells <sup>a,m,2</sup>               | 77,891-2,002,960 scfy/venting well <sup>m</sup>                                | 148,075.1                               |
| <b>Blowdowns</b>   |   |  |   |
| Vessel Blowdowns   | 422,542 vessels <sup>b,2</sup>                      | 76.86-90.94 scfy/vessel <sup>b</sup>   | 668.5                                   |
| Pipeline Blowdowns   | 431,051 miles (gathering) <sup>b,2</sup>            | 304.49-360.28 scfy/mile <sup>b</sup>   | 2,702.9                                 |
| Compressor Blowdowns   | 48,518 compressors <sup>b,2</sup>                   | 3,719-4,400 scfy/compressor <sup>b</sup>                                       | 3,713.8                                 |
| Compressor Starts  | 48,518 compressors <sup>b,2</sup>                   | 8,320-9,844 scfy/compressor <sup>b</sup>                                       | 8,308.4                                 |
| <b>Upsets</b>  |   |  |   |
| Pressure Relief Valves   | 1,015,507 PRV <sup>b,2</sup>                        | 33.50-39.64 scfy/PRV <sup>b</sup>  | 700.3                                   |
| Mishaps  | 107,763 miles <sup>f,2</sup>                        | 659.24-780.03 scf/mile <sup>b</sup>  | 1,463.0                                 |
| <b>Produced Water from Coal Bed Methane Wells</b>                    |   |  |   |
| Black Warrior  | 5,480 wells <sup>o</sup>                            | 0.0023 kt/well <sup>o,1</sup>  | 12,790.5                                |
| Powder River   | 20,596,530,150 gal produced water <sup>n</sup>      | kt/gal water<br>2.3E-09 drainage <sup>n,1</sup>                                | 47,627.3                                |
| <b>Offshore Platforms</b>  |   |  |   |
| Shallow Water Gas Platforms (Gulf of Mexico and Pacific)             | shallow water gas<br>1,973 platforms <sup>p,3</sup> | 8,899 scfd/platform <sup>q</sup>   | 123,460.0                               |

|   |   |                                   |          |
|---|---|-----------------------------------|----------|
| Deep Water Gas Platforms (Gulf of Mexico and Pacific) | deep water gas<br>41 platforms <sup>p,3</sup> | 93,836 scfd/platform <sup>q</sup> | 27,105.3 |
|---|---|-----------------------------------|----------|

|                                       |                |
|---------------------------------------|----------------|
| <b>Regulatory Reductions (kt)</b>     | <b>(91.4)</b>  |
| <b>Voluntary Reductions (kt)</b>      | <b>(483.2)</b> |
| <b>Total Reductions (kt)</b>          | <b>(574.6)</b> |
| <b>Total Potential Emissions (kt)</b> | <b>4,933.8</b> |
| <b>Total Net Emissions (kt)</b>       | <b>4,359.2</b> |

\*The values in this table are net emissions for these sources.

<sup>a</sup> DI Desktop (2015)

<sup>b</sup> EPA/GRI (1996), Methane Emissions from the Natural Gas Industry

<sup>c</sup> 2014 GHGRP – Subpart W data

<sup>d</sup> EPA (2016a)

<sup>e</sup> EPA (2016b)

<sup>f</sup> ICF (1996), Estimation of Activity Factors for the Natural Gas Exploration and Production Industry in the U.S.

<sup>g</sup> API/ICF Memorandum (1997)

<sup>h</sup> 2011-2014 GHGRP – Subpart W data

<sup>i</sup> EIA Monthly Energy Review

<sup>j</sup> Radian (1992), Global Emissions of Methane Sources

<sup>k</sup> EIA U.S. Crude Oil, Natural Gas, and Natural Gas Liquids Reserves Annual Report

<sup>l</sup> EP&P/API Tank Calc runs

<sup>m</sup> API/ANGA (2012), Characterizing Pivotal Sources of Methane Emission from Natural Gas Production – Summary and Analysis of API and ANGA Survey

<sup>n</sup> Wyoming Oil and Gas Conservation Commission (2015)

<sup>o</sup> Alabama State Oil and Gas Board (2015)

<sup>p</sup> Bureau of Ocean Energy Management, Regulation and Enforcement (2011)

<sup>q</sup> EPA (2015), Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2013: Revision to Offshore Platform Emissions

<sup>bb</sup> Totals may not sum due to independent rounding.

<sup>cc</sup> Emissions from oil wells that produce associated gas are estimated in the Petroleum Systems model. In the Natural Gas Systems model, the oil wells counts are used as a driver only.

<sup>dd</sup> NA = not applicable (i.e., this data is not applicable for the Natural Gas Systems model).

<sup>1</sup> Activity data for 2014 available from source.

<sup>2</sup> Ratios relating other factors for which activity data are available.

<sup>3</sup> 2013 activity data are used to determine some or all of the 2014 activity.

\*The values in this table are net emissions for these sources.

**Table A-135: U.S. Activity Data for Hydraulic Fracturing (HF) Completions and Workovers split by 4 categories**

| Activity  | 1990         | 2005          | 2009          | 2010          | 2011 <sup>a</sup> | 2012 <sup>a</sup> | 2013 <sup>a</sup> | 2014 <sup>a</sup> |
|---|--------------|---------------|---------------|---------------|-------------------|-------------------|-------------------|-------------------|
| HF Completions and Workovers that vent            | 3,920        | 10,303        | 5,865         | 5,373         | 4,640             | 3,109             | 1,677             | 1,791             |
| HF Completions and Workovers that flare           | 436          | 1,493         | 1,124         | 1,120         | 1,386             | 703               | 835               | 548               |
| HF Completions and Workovers with RECs            | 0            | 2,353         | 3,187         | 3,528         | 3,884             | 3,413             | 3,156             | 1,043             |
| HF Completions and Workovers with RECs that flare | 0            | 784           | 1,062         | 1,176         | 1,295             | 1,911             | 2,117             | 1,979             |
| <b>Total</b>                                      | <b>4,355</b> | <b>14,933</b> | <b>11,238</b> | <b>11,196</b> | <b>11,204</b>     | <b>9,136</b>      | <b>7,785</b>      | <b>5,361</b>      |

<sup>a</sup> 2011, 2012, 2013, 2014 GHGRP - Subpart W data

**Table A-136: 2014 Data and CH<sub>4</sub> Emissions (Mg) for the Natural Gas Processing Stage**

| Activity                              | 2014 EPA Inventory Values |                            |   |                              | Calculated Potential Emissions (Mg) |
|---------------------------------------|---------------------------|----------------------------|---|------------------------------|-------------------------------------|
|                                       | Activity Data             |                            | Emission Factor (Potential) <sup>aa</sup> |                              |                                     |
| <b>Normal Fugitives</b>               |                           |                            |   |                              |                                     |
| Plants                                | 668                       | plants <sup>a.1</sup>      | 7,906                                     | scfd/plant <sup>b</sup>      | 37,126.4                            |
| Reciprocating Compressors             | 6,020                     | compressors <sup>c.2</sup> | 11,196                                    | scfd/compressor <sup>b</sup> | 473,828.9                           |
| Centrifugal Compressors (wet seals)   | 665                       | compressors <sup>d.2</sup> | 51,370                                    | scfd/compressor <sup>d</sup> | 240,031.0                           |
| Centrifugal Compressors (dry seals)   | 306                       | compressors <sup>d.2</sup> | 25,189                                    | scfd/compressor <sup>d</sup> | 54,117.2                            |
| <b>Vented and Combusted</b>           |                           |                            |   |                              |                                     |
| Gas Engines                           | 43,251                    | MMHPhr <sup>c.2</sup>      | 0.24                                      | scf/HPhr <sup>b</sup>        | 199,922.8                           |
| Gas Turbines                          | 51,283                    | MMHPhr <sup>c.2</sup>      | 0.01                                      | scf/HPhr <sup>b</sup>        | 5,630.0                             |
| AGR Vents                             | 339                       | AGRunits <sup>b.2</sup>    | 6,083                                     | scfd/AGR <sup>b</sup>        | 14,477.9                            |
| Kimray Pumps                          | 1,566,830                 | MMscf/yr <sup>c.2</sup>    | 177.75                                    | scf/MMscf <sup>b</sup>       | 5,364.0                             |
| Dehydrator Vents                      | 14,115,586                | MMscf/yr <sup>c.2</sup>    | 121.55                                    | scf/MMscf <sup>b</sup>       | 33,045.3                            |
| Pneumatic Controllers                 | 668                       | gasplants <sup>a.1</sup>   | 164,721                                   | scfy/plant <sup>b</sup>      | 2,119.2                             |
| <b>Routine Maintenance</b>            |                           |                            |   |                              |                                     |
| Blowdowns/Venting                     | 668                       | gasplants <sup>a.1</sup>   | 4,060                                     | Mscfy/plant <sup>b</sup>     | 52,234.7                            |
| <b>Regulatory Reductions (kt)</b>     |                           |                            |   |                              | <b>(17.5)</b>                       |
| <b>Voluntary Reductions (kt)</b>      |                           |                            |   |                              | <b>(140.8)</b>                      |
| <b>Total Reductions (kt)</b>          |                           |                            |   |                              | <b>(158.3)</b>                      |
| <b>Total Potential Emissions (kt)</b> |                           |                            |   |                              | <b>1,117.9</b>                      |
| <b>Total Net Emissions (kt)</b>       |                           |                            |   |                              | <b>959.6</b>                        |

<sup>a</sup> Oil and Gas Journal<sup>b</sup> EPA/GRI (1996), Methane Emissions from the Natural Gas Industry<sup>c</sup> ICF (2008), Natural Gas Model Activity Factor Basis Change<sup>d</sup> ICF (2010), Emissions from Centrifugal Compressors<sup>aa</sup> Emission factors listed in this table are for potential emissions (unless otherwise indicated in a footnote). See detailed explanation of methodology above.<sup>1</sup> Activity data for 2014 available from source.<sup>2</sup> Ratios relating other factors for which activity data are available.

**Table A-137: 2014 Data and CH<sub>4</sub> Emissions (Mg) for the Natural Gas Transmission Stage**

| Activity                                    | 2014 EPA Inventory Values |                                      |   |                                   | Calculated Potential Emissions (Mg) |
|---|---------------------------|--------------------------------------|---|-----------------------------------|-------------------------------------|
|   | Activity Data             |                                      | Emission Factor (Potential) <sup>aa</sup> |                                   |                                     |
| <b>Fugitives</b>                            |                           |                                      |   |                                   |                                     |
| Pipeline Leaks                              | 301,748                   | miles <sup>a</sup>                   | 1.55                                      | scfd/mile <sup>b</sup>            | 3,296.3                             |
| <i>Compressor Stations (Transmission)*</i>  |                           |                                      |   |                                   |                                     |
| Station Total Emissions                     | 1,834                     | stations <sup>c,d,2</sup>            | 44,459                                    | scfd/station <sup>c,d,bb</sup>    | 573,179.2                           |
| Station + Compressor Fugitive Emissions     | 5,221                     | compressors <sup>c,d,2</sup>         | 9,104                                     | scfd/station <sup>c,d,bb</sup>    | 117,370.9                           |
| Reciprocating Compressor                    | 2,173                     | compressors <sup>c,d,2</sup>         | 9,246                                     | scfd/compressor <sup>c,d,bb</sup> | 339,361.9                           |
| Centrifugal Compressor (wet seals)          | 869                       | compressors <sup>c,d,2</sup>         | 9,673                                     | scfd/compressor <sup>c,d,bb</sup> | 59,092.2                            |
| Centrifugal Compressor (dry seals)          | 1,304                     | compressors <sup>c,d,2</sup>         | 6,259                                     | scfd/compressor <sup>c,d,bb</sup> | 57,354.2                            |
| <i>Compressor Stations (Storage)*</i>       |                           |                                      |   |                                   |                                     |
| Station Total Emissions                     | 356                       | stations <sup>c,d,2</sup>            | 52,604                                    | scfd/station <sup>c,d,bb</sup>    | 131,647.9                           |
| Station + Compressor Fugitive Emissions     | 356                       | stations <sup>c,d,2</sup>            | 10,100                                    | scfd/station <sup>c,d,bb</sup>    | 25,276.0                            |
| Reciprocating Compressor                    | 1,520                     | compressors <sup>c,d,2</sup>         | 9,957                                     | scfd/compressor <sup>c,d,bb</sup> | 106,371.9                           |
| Wells (Storage)                             | 19,522                    | wells <sup>b,2</sup>                 | 114.50                                    | scfd/well <sup>b</sup>            | 15,714.0                            |
| M&R (Trans. Co. Interconnect)               | 2,686                     | stations <sup>e,2</sup>              | 3,984                                     | scfd/station <sup>b</sup>         | 75,230.0                            |
| M&R (Farm Taps + Direct Sales)              | 79,646                    | stations <sup>e,2</sup>              | 31.20                                     | scfd/station <sup>b</sup>         | 17,468.9                            |
| <b>Normal Operation</b>                     |                           |                                      |   |                                   |                                     |
| Dehydrator vents (Transmission)             | 1,169,007                 | MMscf/yr <sup>b,2</sup>              | 93.72                                     | scf/MMscf <sup>b</sup>            | 2,110.1                             |
| Dehydrator vents (Storage)                  | 2,169,267                 | MMscf/yr <sup>b,2</sup>              | 117.18                                    | scf/MMscf <sup>b</sup>            | 4,895.8                             |
| <i>Compressor Exhaust</i>                   |                           |                                      |   |                                   |                                     |
| Engines (Transmission)                      | 53,295                    | MMHPhr <sup>b,2</sup>                | 0.24                                      | scf/HPhr <sup>b</sup>             | 246,351.2                           |
| Turbines (Transmission)                     | 12,717                    | MMHPhr <sup>b,2</sup>                | 0.01                                      | scf/HPhr <sup>b</sup>             | 1,396.1                             |
| Engines (Storage)                           | 5,339                     | MMHPhr <sup>b,2</sup>                | 0.24                                      | scf/HPhr <sup>b</sup>             | 24,677.0                            |
| Turbines (Storage)                          | 1,875                     | MMHPhr <sup>b,2</sup>                | 0.01                                      | scf/HPhr <sup>b</sup>             | 205.9                               |
| Generators (Engines)                        | 2,608                     | MMHPhr <sup>b,2</sup>                | 0.24                                      | scf/HPhr <sup>b</sup>             | 12,055.2                            |
| Generators (Turbines)                       | 31                        | MMHPhr <sup>b,2</sup>                | 0.01                                      | scf/HPhr <sup>b</sup>             | 3.4                                 |
| <i>Pneumatic Devices Trans + Stor*</i>      |                           |                                      |   |                                   |                                     |
| Pneumatic Devices Transmission (High Bleed) | 47,140                    | devices <sup>c,d,2</sup>             | 30,611                                    | scfy/device <sup>c,d,bb</sup>     | 27,792.1                            |
| (Intermittent Bleed)                        | 4,129                     | devices <sup>c,d,2</sup>             | 151,969                                   | scfy/device <sup>c,d,bb</sup>     | 12,085.2                            |
| (Low Bleed)                                 | 39,216                    | devices <sup>c,d,2</sup>             | 19,712                                    | scfy/device <sup>c,d,bb</sup>     | 14,888.7                            |
| Pneumatic Devices Storage (High Bleed)      | 3,795                     | devices <sup>c,d,2</sup>             | 11,196                                    | scfy/device <sup>c,d,bb</sup>     | 818.2                               |
| (Intermittent Bleed)                        | 23,964                    | devices <sup>c,d,2</sup>             | 63,622                                    | scfy/device <sup>c,d,bb</sup>     | 29,364.5                            |
| (Low Bleed)                                 | 8,379                     | devices <sup>c,d,2</sup>             | 147,983                                   | scfy/device <sup>c,d,bb</sup>     | 23,882.0                            |
|   | 13,482                    | devices <sup>c,d,2</sup>             | 19,333                                    | scfy/device <sup>c,d,bb</sup>     | 5,020.1                             |
|   | 2,103                     | devices <sup>c,d,2</sup>             | 11,414                                    | scfy/device <sup>c,d,bb</sup>     | 462.3                               |
| <b>Routine Maintenance/Upsets</b>           |                           |                                      |   |                                   |                                     |
| Pipeline venting                            | 301,748                   | miles <sup>a,1</sup>                 | 31.65                                     | Mscfy/mile <sup>b</sup>           | 183,939.2                           |
| <i>Station venting Trans + Storage</i>      |                           |                                      |   |                                   |                                     |
| Station Venting Transmission                | 1,834                     | compressor stations <sup>c,d,2</sup> | 4,359                                     | Mscfy/station <sup>b</sup>        | 153,965.5                           |
| Station Venting Storage                     | 356                       | compressor stations <sup>c,d,2</sup> | 4,359                                     | Mscfy/station <sup>b</sup>        | 29,887.7                            |
| <b>LNG Storage</b>                          |                           |                                      |   |                                   |                                     |
| LNG Stations                                | 70                        | stations <sup>f,g,3</sup>            | 21,507                                    | scfd/station <sup>b</sup>         | 10,622.8                            |
| LNG Reciprocating Compressors               | 270                       | compressors <sup>f,g,3</sup>         | 21,116                                    | scfd/compressor <sup>b</sup>      | 40,146.5                            |
| LNG Centrifugal Compressors                 | 64                        | compressors <sup>f,g,3</sup>         | 30,573                                    | scfd/compressor <sup>b</sup>      | 13,766.0                            |
| <i>LNG Compressor Exhaust</i>               |                           |                                      |   |                                   |                                     |
| LNG Engines                                 | 579                       | MMHPhr <sup>f,g,3</sup>              | 0.24                                      | scf/HPhr <sup>b</sup>             | 2,677.7                             |
| LNG Turbines                                | 113                       | MMHPhr <sup>f,g,3</sup>              | 0.01                                      | scf/HPhr <sup>b</sup>             | 12.4                                |
| LNG Station venting                         | 70                        | stations <sup>f,g,3</sup>            | 4,359                                     | Mscfy/station <sup>b</sup>        | 5,898.6                             |
| <b>LNG Import Terminals</b>                 |                           |                                      |   |                                   |                                     |
| LNG Stations                                | 8                         | stations <sup>f,g,3</sup>            | 21,507                                    | scfd/station <sup>b</sup>         | 1,270.0                             |
| LNG Reciprocating Compressors               | 41                        | compressors <sup>f,g,3</sup>         | 21,116                                    | scfd/compressor <sup>b</sup>      | 6,056.5                             |
| LNG Centrifugal Compressors                 | 7                         | compressors <sup>f,g,3</sup>         | 30,573                                    | scfd/compressor <sup>b</sup>      | 1,547.5                             |
| <i>LNG Compressor Exhaust</i>               |                           |                                      |   |                                   |                                     |
| LNG Engines                                 | 303                       | MMHPhr <sup>f,g,3</sup>              | 0.24                                      | scf/HPhr <sup>b</sup>             | 1,401.7                             |

|                                       |    |                           |       |                            |         |
|---------------------------------------|----|---------------------------|-------|----------------------------|---------|
| LNG Turbines                          | 60 | MMHPh <sup>f,g,3</sup>    | 0.01  | scf/HPh <sup>b</sup>       | 6.6     |
| LNG Station venting                   | 8  | stations <sup>f,g,3</sup> | 4,359 | Mscfy/station <sup>b</sup> | 705.2   |
| <b>Regulatory Reductions (kt)</b>     |    |                           |       |                            | -       |
| <b>Voluntary Reductions (kt)</b>      |    |                           |       |                            | (335.0) |
| <b>Total Reductions (kt)</b>          |    |                           |       |                            | (335.0) |
| <b>Total Potential Emissions (kt)</b> |    |                           |       |                            | 1,617.3 |
| <b>Total Net Emissions (kt)</b>       |    |                           |       |                            | 1,282.3 |

<sup>a</sup> Pipeline and Hazardous Materials Safety Administration (PHMSA), Office of Pipeline Safety (OPS) (2014)

<sup>b</sup> EPA/GRI (1996), Methane Emissions from the Natural Gas Industry

<sup>c</sup> EPA (2016c)

<sup>d</sup> 2014 GHGRP – Subpart W data

<sup>e</sup> ICF (2008), Natural Gas Model Activity Factor Basis Change

<sup>f</sup> ICF (1996), Estimation of Activity Factors for the Natural Gas Exploration and Production Industry in the U.S.

<sup>g</sup> EIA (2004), U.S. LNG Markets and Uses

<sup>1</sup> Activity data for 2014 available from source.

<sup>2</sup> Ratios relating other factors for which activity data are available.

<sup>3</sup> 2013 activity data are used to determine some or all of the 2014 activity (to be updated).

<sup>aa</sup> Emission factors listed in this table are for potential emissions (unless otherwise indicated in a footnote). See detailed explanation of methodology above.

<sup>bb</sup> Emission factors represent actual emissions and can be used to calculate emissions directly.

\*The values in this table are net emissions for these sources.

**Table A-138: 2014 Data and CH<sub>4</sub> Emissions (Mg) for the Natural Gas Distribution Stage**

| Activity                              | 2014 EPA Inventory Values |                             |   |                                | Calculated Potential Emissions (Mg) |
|---------------------------------------|---------------------------|-----------------------------|---|--------------------------------|-------------------------------------|
|                                       | Activity Data             |                             | Emission Factor (Potential) <sup>aa</sup> |                                |                                     |
| <b>Pipeline Leaks*</b>                |                           |                             |   |                                |                                     |
| Mains—Cast Iron                       | 29,359                    | miles <sup>a,1</sup>        | 60.09                                     | Mscf/mile-yr <sup>c,bb</sup>   | 33,976.2                            |
| Mains—Unprotected steel               | 58,520                    | miles <sup>a,1</sup>        | 44.72                                     | Mscf/mile-yr <sup>c,bb</sup>   | 50,404.7                            |
| Mains—Protected steel                 | 486,432                   | miles <sup>a,1</sup>        | 5.02                                      | Mscf/mile-yr <sup>c,bb</sup>   | 47,061.8                            |
| Mains—Plastic                         | 690,029                   | miles <sup>a,1</sup>        | 1.50                                      | Mscf/mile-yr <sup>c,bb</sup>   | 19,905.5                            |
| Services—Unprotected steel            | 3,432,641                 | services <sup>a,1</sup>     | 0.75                                      | Mscf/service <sup>c,bb</sup>   | 49,727.5                            |
| Services Protected steel              | 14,588,827                | services <sup>a,1</sup>     | 0.07                                      | Mscf/service <sup>c,bb</sup>   | 18,896.3                            |
| Services—Plastic                      | 46,755,197                | services <sup>a,1</sup>     | 0.01                                      | Mscf/service <sup>c,bb</sup>   | 12,296.3                            |
| Services—Copper                       | 935,595                   | services <sup>a,1</sup>     | 0.25                                      | Mscf/service <sup>b</sup>      | 4,582.8                             |
| <b>Meter/Regulator (City Gates)*</b>  |                           |                             |   |                                |                                     |
| M&R >300                              | 3,952                     | stations <sup>c,d,2</sup>   | 12.70                                     | scfh/station <sup>c,d,bb</sup> | 8,468.2                             |
| M&R 100-300                           | 14,423                    | stations <sup>c,d,2</sup>   | 5.90                                      | scfh/station <sup>c,d,bb</sup> | 14,356.7                            |
| M&R <100                              | 7,709                     | stations <sup>c,d,2</sup>   | 4.31                                      | scfh/station <sup>b</sup>      | 5,605.7                             |
| Reg >300                              | 4,321                     | stations <sup>c,d,2</sup>   | 5.15                                      | scfh/station <sup>c,d,bb</sup> | 3,754.4                             |
| R-Vault >300                          | 1,501                     | stations <sup>c,d,2</sup>   | 0.30                                      | scfh/station <sup>c,d,bb</sup> | 76.0                                |
| Reg 100-300                           | 13,071                    | stations <sup>c,d,2</sup>   | 0.85                                      | scfh/station <sup>c,d,bb</sup> | 1,874.6                             |
| R-Vault 100-300                       | 3,473                     | stations <sup>c,d,2</sup>   | 0.30                                      | scfh/station <sup>c,d,bb</sup> | 175.8                               |
| Reg 40-100                            | 39,225                    | stations <sup>c,d,2</sup>   | 0.97                                      | scfh/station <sup>c,d,bb</sup> | 6,419.5                             |
| R-Vault 40-100                        | 20,573                    | stations <sup>c,d,2</sup>   | 0.30                                      | scfh/station <sup>c,d,bb</sup> | 1,041.3                             |
| Reg <40                               | 16,633                    | stations <sup>c,d,2</sup>   | 0.13                                      | scfh/station <sup>b</sup>      | 373.2                               |
| <b>Customer Meters</b>                |                           |                             |   |                                |                                     |
| Residential                           | 53,339,363                | Outdoor meters <sup>e</sup> | 77.31                                     | scfy/meter <sup>b,c</sup>      | 79,424.1                            |
| Commercial/Industry                   | 5,611,121                 | meters <sup>e</sup>         | 505.40                                    | scfy/meter <sup>b,c</sup>      | 54,619.1                            |
| <b>Routine Maintenance</b>            |                           |                             |   |                                |                                     |
| Pressure Relief Valve Releases        | 1,264,340                 | milemain <sup>a,1</sup>     | 0.05                                      | Mscf/mile <sup>b</sup>         | 1,217.6                             |
| Pipeline Blowdown                     | 2,168,588                 | miles <sup>b,2</sup>        | 0.10                                      | Mscfy/mile <sup>b</sup>        | 4,260.2                             |
| <b>Upsets</b>                         |                           |                             |   |                                |                                     |
| Mishaps (Dig-ins)                     | 2,168,588                 | miles <sup>b,2</sup>        | 1.59                                      | Mscfy/mile <sup>b</sup>        | 66,409.5                            |
| <b>Regulatory Reductions (kt)</b>     |                           |                             |   |                                | -                                   |
| <b>Voluntary Reductions (kt)</b>      |                           |                             |   |                                | (40.8)                              |
| <b>Total Reductions (kt)</b>          |                           |                             |   |                                | (40.8)                              |
| <b>Total Potential Emissions (kt)</b> |                           |                             |   |                                | 484.9                               |
| <b>Total Net Emissions (kt)</b>       |                           |                             |   |                                | 444.1                               |

<sup>a</sup> Pipeline and Hazardous Materials Safety Administration (PHMSA), Office of Pipeline Safety (OPS) (2013)

<sup>b</sup> EPA/GRI (1996), Methane Emissions from the Natural Gas Industry

<sup>c</sup> EPA (2016d)

<sup>d</sup> 2014 GHGRP – Subpart W data

<sup>f</sup> EIA (2015d, 2015e, 2015f) – Number of Consumers (Residential, Commercial, and Industrial)

<sup>aa</sup> Emission factors listed in this table are for potential emissions (unless otherwise indicated in a footnote). See detailed explanation of methodology above.

<sup>bb</sup> Emission factors represent actual emissions and can be used to calculate emissions directly.

<sup>1</sup> Activity data for 2014 available from source.

<sup>2</sup> Ratios relating other factors for which activity data are available.

\*The values in this table are net emissions for these sources.

**Table A-139: U.S. Production Sector CH<sub>4</sub> Content in Natural Gas by NEMS Region (General Sources)**

| Year | U.S. Region |              |                |            |            |            |                 |
|------|-------------|--------------|----------------|------------|------------|------------|-----------------|
|      | North East  | Midcontinent | Rocky Mountain | South West | West Coast | Gulf Coast | Lower 48 States |
| 1990 | 84.0%       | 78.3%        | 67.0%          | 64.4%      | 75.3%      | 79.8%      | n/a             |
| 1991 | 83.8%       | 78.7%        | 69.1%          | 67.1%      | 78.1%      | 80.1%      | n/a             |
| 1992 | 83.5%       | 79.1%        | 71.4%          | 75.4%      | 80.8%      | 82.7%      | n/a             |
| 1993 | 82.9%       | 79.9%        | 73.4%          | 76.1%      | 83.6%      | 84.1%      | n/a             |
| 1994 | 82.0%       | 80.7%        | 75.5%          | 77.4%      | 86.4%      | 85.6%      | n/a             |
| 1995 | 81.5%       | 81.6%        | 77.6%          | 79.0%      | 89.1%      | 87.2%      | n/a             |
| 1996 | 81.2%       | 82.6%        | 80.5%          | 80.5%      | 91.9%      | 88.7%      | 84.2%           |
| 1997 | 80.3%       | 82.5%        | 80.4%          | 80.5%      | 91.9%      | 88.6%      | 84.1%           |
| 1998 | 81.0%       | 82.5%        | 80.5%          | 80.5%      | 91.9%      | 88.6%      | 84.2%           |
| 1999 | 80.5%       | 82.5%        | 80.4%          | 80.5%      | 91.9%      | 88.7%      | 84.2%           |
| 2000 | 80.8%       | 82.5%        | 80.2%          | 80.5%      | 91.9%      | 88.7%      | 84.0%           |
| 2001 | 80.3%       | 82.5%        | 79.5%          | 80.5%      | 91.9%      | 88.7%      | 83.8%           |
| 2002 | 80.4%       | 82.5%        | 79.3%          | 80.5%      | 91.9%      | 88.6%      | 83.5%           |
| 2003 | 76.4%       | 82.6%        | 79.1%          | 80.5%      | 91.9%      | 88.6%      | 83.2%           |
| 2004 | 80.4%       | 82.7%        | 79.0%          | 80.5%      | 91.9%      | 88.6%      | 83.4%           |
| 2005 | 80.1%       | 82.7%        | 79.0%          | 80.5%      | 91.9%      | 88.6%      | 83.4%           |
| 2006 | 79.5%       | 83.0%        | 78.9%          | 80.5%      | 91.9%      | 88.6%      | 83.4%           |
| 2007 | 85.8%       | 82.7%        | 77.5%          | 80.5%      | 91.9%      | 88.6%      | 83.9%           |
| 2008 | 86.0%       | 82.7%        | 77.7%          | 80.5%      | 91.9%      | 88.5%      | 83.9%           |
| 2009 | 85.1%       | 82.7%        | 77.5%          | 80.5%      | 91.9%      | 88.5%      | 83.6%           |
| 2010 | 84.3%       | 82.8%        | 77.4%          | 80.5%      | 91.9%      | 88.3%      | 83.4%           |
| 2011 | 85.2%       | 82.6%        | 77.5%          | 80.5%      | 91.9%      | 88.2%      | 83.3%           |
| 2012 | 84.8%       | 82.5%        | 78.2%          | 80.5%      | 91.9%      | 88.2%      | 83.0%           |
| 2013 | 84.2%       | 82.5%        | 77.9%          | 80.5%      | 91.9%      | 88.9%      | 83.0%           |
| 2014 | 84.0%       | 82.4%        | 77.7%          | 80.5%      | 91.9%      | 88.9%      | 82.7%           |

**Table A-140: U.S. Production Sector CH<sub>4</sub> Content in Natural Gas by NEMS Region (Gas Wells Without Hydraulic Fracturing)**

| Year | U.S. Region |              |                |            |            |            |                 |
|------|-------------|--------------|----------------|------------|------------|------------|-----------------|
|      | North East  | Midcontinent | Rocky Mountain | South West | West Coast | Gulf Coast | Lower 48 States |
| 1990 | 84.0%       | 78.3%        | 67.0%          | 64.4%      | 75.3%      | 79.8%      | n/a             |
| 1991 | 83.8%       | 78.7%        | 69.1%          | 67.1%      | 78.1%      | 80.1%      | n/a             |
| 1992 | 83.5%       | 79.1%        | 71.4%          | 75.4%      | 80.8%      | 82.7%      | n/a             |
| 1993 | 82.9%       | 79.9%        | 73.4%          | 76.1%      | 83.6%      | 84.1%      | n/a             |
| 1994 | 82.0%       | 80.7%        | 75.5%          | 77.4%      | 86.4%      | 85.6%      | n/a             |
| 1995 | 81.5%       | 81.6%        | 77.6%          | 79.0%      | 89.1%      | 87.2%      | n/a             |
| 1996 | 81.2%       | 82.5%        | 79.6%          | 80.5%      | 91.9%      | 88.6%      | 84.0%           |
| 1997 | 80.5%       | 82.5%        | 79.5%          | 80.5%      | 91.9%      | 88.6%      | 83.9%           |
| 1998 | 81.2%       | 82.5%        | 79.5%          | 80.5%      | 91.9%      | 88.6%      | 84.0%           |
| 1999 | 80.7%       | 82.5%        | 79.5%          | 80.5%      | 91.9%      | 88.7%      | 83.9%           |
| 2000 | 81.0%       | 82.5%        | 79.2%          | 80.5%      | 91.9%      | 88.7%      | 83.8%           |
| 2001 | 80.4%       | 82.5%        | 78.3%          | 80.5%      | 91.9%      | 88.6%      | 83.5%           |
| 2002 | 80.5%       | 82.5%        | 78.1%          | 80.5%      | 91.9%      | 88.6%      | 83.2%           |
| 2003 | 76.5%       | 82.6%        | 77.9%          | 80.5%      | 91.9%      | 88.6%      | 82.9%           |
| 2004 | 80.5%       | 82.6%        | 77.8%          | 80.5%      | 91.9%      | 88.6%      | 83.1%           |
| 2005 | 80.3%       | 82.7%        | 77.7%          | 80.5%      | 91.9%      | 88.6%      | 83.1%           |
| 2006 | 79.6%       | 83.0%        | 77.7%          | 80.5%      | 91.9%      | 88.6%      | 83.1%           |
| 2007 | 85.6%       | 82.7%        | 75.8%          | 80.5%      | 91.9%      | 88.6%      | 83.5%           |
| 2008 | 85.6%       | 82.7%        | 76.0%          | 80.5%      | 91.9%      | 88.5%      | 83.5%           |
| 2009 | 84.7%       | 82.7%        | 75.8%          | 80.5%      | 91.9%      | 88.5%      | 83.2%           |
| 2010 | 83.8%       | 82.8%        | 75.6%          | 80.5%      | 91.9%      | 88.3%      | 82.9%           |
| 2011 | 85.0%       | 82.6%        | 75.8%          | 80.5%      | 91.9%      | 88.2%      | 82.8%           |

|      |       |       |       |       |       |       |       |
|------|-------|-------|-------|-------|-------|-------|-------|
| 2012 | 84.4% | 82.5% | 76.7% | 80.5% | 91.9% | 88.2% | 82.5% |
| 2013 | 83.9% | 82.5% | 76.3% | 80.5% | 91.9% | 88.9% | 82.4% |
| 2014 | 83.6% | 82.4% | 76.0% | 80.5% | 91.9% | 88.9% | 82.1% |

**Table A-141: U.S. Production Sector CH<sub>4</sub> Content in Natural Gas by NEMS Region (Gas Wells With Hydraulic Fracturing)**

| Year | U.S. Region |              |                |            |            |            |                 |
|------|-------------|--------------|----------------|------------|------------|------------|-----------------|
|      | North East  | Midcontinent | Rocky Mountain | South West | West Coast | Gulf Coast | Lower 48 States |
| 1990 | 84.0%       | 78.3%        | 67.0%          | 64.4%      | 75.3%      | 79.8%      | n/a             |
| 1991 | 83.8%       | 78.7%        | 69.1%          | 67.1%      | 78.1%      | 80.1%      | n/a             |
| 1992 | 83.5%       | 79.1%        | 71.4%          | 75.4%      | 80.8%      | 82.7%      | n/a             |
| 1993 | 82.9%       | 79.9%        | 73.4%          | 76.1%      | 83.6%      | 84.1%      | n/a             |
| 1994 | 82.0%       | 80.7%        | 75.5%          | 77.4%      | 86.4%      | 85.6%      | n/a             |
| 1995 | 81.5%       | 81.6%        | 77.6%          | 79.0%      | 89.1%      | 87.2%      | n/a             |
| 1996 | 83.2%       | 92.6%        | 74.4%          | 80.5%      | 91.9%      | 88.7%      | 82.1%           |
| 1997 | 83.1%       | 92.6%        | 74.9%          | 80.5%      | 91.9%      | 88.6%      | 82.1%           |
| 1998 | 83.1%       | 92.6%        | 75.5%          | 80.5%      | 91.9%      | 88.6%      | 82.3%           |
| 1999 | 83.1%       | 92.6%        | 75.3%          | 80.5%      | 91.9%      | 88.7%      | 81.9%           |
| 2000 | 83.0%       | 92.6%        | 76.4%          | 80.5%      | 91.9%      | 88.7%      | 82.5%           |
| 2001 | 83.0%       | 92.6%        | 78.9%          | 80.5%      | 91.9%      | 88.7%      | 83.6%           |
| 2002 | 83.0%       | 92.6%        | 80.5%          | 80.5%      | 91.9%      | 88.6%      | 84.4%           |
| 2003 | 83.1%       | 92.6%        | 81.4%          | 80.5%      | 91.9%      | 88.6%      | 84.9%           |
| 2004 | 83.0%       | 92.6%        | 81.7%          | 80.5%      | 91.9%      | 88.6%      | 85.2%           |
| 2005 | 83.0%       | 92.6%        | 82.0%          | 80.5%      | 91.9%      | 88.6%      | 85.3%           |
| 2006 | 83.0%       | 92.6%        | 82.6%          | 80.5%      | 91.9%      | 88.6%      | 85.7%           |
| 2007 | 83.5%       | 92.6%        | 86.5%          | 80.5%      | 91.9%      | 88.6%      | 88.7%           |
| 2008 | 84.1%       | 92.6%        | 86.3%          | 80.5%      | 91.9%      | 88.5%      | 88.4%           |
| 2009 | 84.1%       | 92.6%        | 86.8%          | 80.5%      | 91.9%      | 88.5%      | 88.7%           |
| 2010 | 84.3%       | 92.6%        | 86.8%          | 80.5%      | 91.9%      | 88.3%      | 89.0%           |
| 2011 | 83.6%       | 92.6%        | 87.9%          | 80.5%      | 91.9%      | 88.2%      | 89.4%           |
| 2012 | 84.0%       | 92.6%        | 87.6%          | 80.5%      | 91.9%      | 88.2%      | 89.1%           |
| 2013 | 83.9%       | 92.6%        | 88.1%          | 80.5%      | 91.9%      | 88.9%      | 89.3%           |
| 2014 | 84.2%       | 92.6%        | 88.8%          | 80.5%      | 91.9%      | 88.9%      | 90.0%           |

**Table A-142: Key Activity Data Drivers**

| Variable  | Units                | 1990              | 2005              | 2009              | 2010              | 2011              | 2012              | 2013              | 2014              |
|---|----------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| <b>Transmission Pipelines Length</b>                              | <b>miles</b>         | <b>291,925</b>    | <b>300,468</b>    | <b>304,573</b>    | <b>304,805</b>    | <b>305,058</b>    | <b>303,341</b>    | <b>302,777</b>    | <b>301,748</b>    |
| <b>Wells</b>  |                      |                   |                   |                   |                   |                   |                   |                   |                   |
| NE—Associated Gas Wells <sup>a,1</sup>                            | # wells              | 28,397            | 32,769            | 37,141            | 41,541            | 42,634            | 45,314            | 43,781            | 45,349            |
| NE—Non-associated Gas Wells <sup>a,1</sup>                        | # wells              | 64,620            | 128,114           | 148,980           | 155,754           | 158,294           | 158,666           | 158,017           | 160,692           |
| MC—Associated Gas Wells <sup>a,1</sup>                            | # wells              | 50,789            | 43,686            | 46,561            | 46,189            | 46,053            | 48,547            | 50,142            | 46,830            |
| MC—Non-associated Gas Wells <sup>a,1</sup>                        | # wells              | 62,473            | 85,656            | 101,654           | 101,457           | 102,687           | 101,312           | 99,609            | 100,655           |
| RM—Associated Gas Wells <sup>a,1</sup>                            | # wells              | 23,419            | 29,425            | 37,769            | 39,888            | 43,427            | 47,547            | 50,523            | 53,964            |
| RM—Non-associated Gas Wells <sup>a,1</sup>                        | # wells              | 22,737            | 61,832            | 79,083            | 78,607            | 79,427            | 78,918            | 76,848            | 75,745            |
| SW—Associated Gas Wells <sup>a,1</sup>                            | # wells              | 308,254           | 220,818           | 227,313           | 232,151           | 236,087           | 244,444           | 247,386           | 253,329           |
| SW—Non-associated Gas Wells <sup>a,1</sup>                        | # wells              | 32,665            | 42,806            | 48,949            | 49,056            | 49,351            | 48,812            | 48,901            | 49,023            |
| WC—Associated Gas Wells <sup>a,1</sup>                            | # wells              | 16,627            | 18,131            | 28,600            | 28,095            | 28,467            | 28,756            | 29,578            | 30,071            |
| WC—Non-associated Gas Wells <sup>a,1</sup>                        | # wells              | 2,130             | 2,293             | 2,788             | 2,672             | 2,738             | 2,732             | 2,516             | 2,424             |
| GC—Associated Gas Wells <sup>a,1</sup>                            | # wells              | 141,216           | 62,003            | 60,856            | 60,263            | 61,432            | 66,369            | 69,685            | 74,330            |
| GC—Non-associated Gas Wells <sup>a,1</sup>                        | # wells              | 34,084            | 53,202            | 67,185            | 69,274            | 70,701            | 70,148            | 68,600            | 67,601            |
| <b>Platforms<sup>aa</sup></b>                                     |                      |                   |                   |                   |                   |                   |                   |                   |                   |
| Gulf of Mexico and Pacific OCS Off-shore Platforms <sup>b,2</sup> | # platforms          | 3,941             | 3,909             | 3,570             | 3,432             | 3,432             | 3,432             | 3,432             | 3,432             |
| GoM and Pacific OCS Deep Water Platforms <sup>b,2</sup>           | # platforms          | 17                | 59                | 68                | 70                | 70                | 70                | 70                | 70                |
| <b>Gas Plants<sup>c,1</sup></b>                                   | <b># gas plants</b>  | <b>761</b>        | <b>566</b>        | <b>579</b>        | <b>585</b>        | <b>606</b>        | <b>606</b>        | <b>650</b>        | <b>668</b>        |
| <b>Distribution Services</b>                                      | <b># of services</b> | <b>47,883,083</b> | <b>61,832,574</b> | <b>64,498,352</b> | <b>64,079,565</b> | <b>64,774,921</b> | <b>65,057,667</b> | <b>65,554,546</b> | <b>65,712,260</b> |
| Steel—Unprotected <sup>d,1</sup>                                  | # of services        | 7,633,526         | 5,507,356         | 5,221,143         | 4,226,682         | 4,142,891         | 3,917,081         | 3,669,003         | 3,432,641         |
| Steel—Protected <sup>d,1</sup>                                    | # of services        | 19,781,581        | 16,529,118        | 15,470,934        | 15,324,020        | 15,275,872        | 14,954,692        | 14,753,566        | 14,588,827        |
| Plastic <sup>d,1</sup>  | # of services        | 18,879,865        | 38,549,089        | 42,719,579        | 43,450,479        | 44,299,530        | 45,176,606        | 46,158,870        | 46,755,197        |
| Copper <sup>d,1</sup>   | # of services        | 1,588,111         | 1,247,011         | 1,086,696         | 1,078,384         | 1,056,628         | 1,009,288         | 973,107           | 935,595           |
| <b>Distribution Mains</b>   | <b>miles</b>         | <b>944,157</b>    | <b>1,162,560</b>  | <b>1,218,148</b>  | <b>1,228,099</b>  | <b>1,237,203</b>  | <b>1,245,694</b>  | <b>1,253,316</b>  | <b>1,264,340</b>  |
| Cast Iron <sup>d,1</sup>  | miles                | 58,292            | 39,645            | 35,623            | 34,592            | 33,669            | 32,406            | 30,904            | 29,359            |
| Steel—Unprotected <sup>d,1</sup>                                  | miles                | 108,941           | 72,458            | 67,758            | 67,443            | 64,981            | 63,703            | 60,619            | 58,520            |
| Steel—Protected <sup>d,1</sup>                                    | miles                | 465,538           | 490,156           | 488,352           | 488,718           | 488,810           | 487,646           | 486,771           | 486,432           |
| Plastic <sup>d,1</sup>  | miles                | 311,386           | 560,301           | 626,415           | 637,346           | 649,743           | 661,939           | 675,022           | 690,029           |

<sup>a</sup> DI Desktop (2015)

<sup>b</sup> Bureau of Ocean Energy Management, Regulation and Enforcement (2011)

<sup>c</sup> Oil and Gas Journal

<sup>d</sup> Pipeline and Hazardous Materials Safety Administration (PHMSA), Office of Pipeline Safety (OPS) (2014)

<sup>1</sup> Activity data for 2014 available from source.

<sup>2</sup> 2013 activity data are used to determine some or all of the 2014 activity (to be updated).

<sup>aa</sup> Number of platforms include both oil and gas platforms

**Table A-143: CH<sub>4</sub> Reductions Derived from the Natural Gas STAR Program (kt)**

| Process                             | 1990         | 2005           | 2009           | 2010             | 2011           | 2012           | 2013           | 2014           |
|-------------------------------------|--------------|----------------|----------------|------------------|----------------|----------------|----------------|----------------|
| <b>Production</b>                   | <b>(3.5)</b> | <b>(253.8)</b> | <b>(349.1)</b> | <b>(393.9)</b>   | <b>(427.6)</b> | <b>(421.4)</b> | <b>(446.7)</b> | <b>(483.2)</b> |
| Pipeline Leaks                      | (0.0)        | (2.4)          | -              | -                | -              | -              | -              | -              |
| Gas Engines                         | (0.0)        | (97.9)         | (123.1)        | (133.3)          | (137.2)        | (140.0)        | (140.5)        | (141.0)        |
| Compressor Starts                   | -            | (0.2)          | (0.4)          | (0.5)            | (0.5)          | (0.5)          | (0.5)          | (0.5)          |
| Other Production                    | (3.5)        | (155.7)        | (225.6)        | (260.0)          | (289.9)        | (280.9)        | (305.7)        | (341.7)        |
| <b>Processing</b>                   | <b>(1.5)</b> | <b>(155.5)</b> | <b>(137.6)</b> | <b>(214.3)</b>   | <b>(140.4)</b> | <b>(140.4)</b> | <b>(140.7)</b> | <b>(140.8)</b> |
| Fugitives Reciprocating Compressors | -            | -              | -              | (1.0)            | -              | -              | (0.1)          | (0.1)          |
| Gas Engines                         | -            | (1.1)          | (6.0)          | (6.1)            | (6.1)          | (6.1)          | (6.1)          | (6.1)          |
| AGR Vents                           | -            | -              | -              | -                | -              | -              | -              | -              |
| Dehydrator Vents                    | (1.3)        | (2.1)          | (9.3)          | (9.3)            | (9.3)          | (9.3)          | (9.3)          | (9.3)          |
| Other Processing                    | (0.2)        | (152.2)        | (122.2)        | (197.9)          | (125.0)        | (125.0)        | (125.2)        | (125.2)        |
| <b>Transmission and Storage</b>     | <b>-</b>     | <b>(496.2)</b> | <b>(379.0)</b> | <b>(433.2)</b>   | <b>(341.9)</b> | <b>(375.4)</b> | <b>(331.8)</b> | <b>(335.0)</b> |
| Engines (Transmission)              | -            | (83.2)         | (116.2)        | (119.4)          | (121.7)        | (124.0)        | (126.9)        | (129.8)        |
| Pipeline Vents (Transmission)       | -            | (124.9)        | (88.7)         | (128.2)          | (58.9)         | (100.1)        | (59.4)         | (59.4)         |
| Other Transmission                  | -            | (288.1)        | (174.1)        | (185.6)          | (161.3)        | (151.2)        | (145.5)        | (145.8)        |
| <b>Distribution</b>                 | <b>-</b>     | <b>(48.3)</b>  | <b>(63.9)</b>  | <b>(54.5)</b>    | <b>(58.0)</b>  | <b>(45.1)</b>  | <b>(40.5)</b>  | <b>(40.8)</b>  |
| Mishaps (Dig-ins)                   | -            | (0.3)          | (0.8)          | (0.8)            | (4.7)          | (0.7)          | (0.8)          | (0.8)          |
| Other Distribution                  | -            | (48.0)         | (63.1)         | (53.7)           | (53.3)         | (44.4)         | (39.6)         | (40.0)         |
| <b>Total</b>                        | <b>(5.0)</b> | <b>(953.8)</b> | <b>(929.5)</b> | <b>(1,095.9)</b> | <b>(967.9)</b> | <b>(982.4)</b> | <b>959.7)</b>  | <b>999.8)</b>  |

**Table A-144: CH<sub>4</sub> Reductions Derived from Regulations (kt)**

| Process                         | 1990         | 2005          | 2009          | 2010           | 2011          | 2012          | 2013           | 2014           |
|---------------------------------|--------------|---------------|---------------|----------------|---------------|---------------|----------------|----------------|
| <b>Production</b>               | <b>NA</b>    | <b>(45.9)</b> | <b>(55.1)</b> | <b>(94.3)</b>  | <b>(71.7)</b> | <b>(79.6)</b> | <b>(93.6)</b>  | <b>(91.4)</b>  |
| Dehydrator vents (NESHAP)       | NA           | (14.0)        | (10.9)        | (10.5)         | (10.3)        | (10.2)        | (10.1)         | (10.1)         |
| Condensate tanks (NESHAP)       | NA           | (31.9)        | (44.2)        | (83.8)         | (61.4)        | (69.4)        | (83.6)         | (81.3)         |
| <b>Processing</b>               | <b>(0.0)</b> | <b>(12.1)</b> | <b>(13.9)</b> | <b>(14.4)</b>  | <b>(15.5)</b> | <b>(16.3)</b> | <b>(16.4)</b>  | <b>(17.5)</b>  |
| Dehydrator vents (NESHAP)       | (0.0)        | (12.1)        | (13.9)        | (14.4)         | (15.5)        | (16.3)        | (16.4)         | (17.5)         |
| <b>Transmission and Storage</b> | <b>NA</b>    | <b>NA</b>     | <b>NA</b>     | <b>NA</b>      | <b>NA</b>     | <b>NA</b>     | <b>NA</b>      | <b>NA</b>      |
| <b>Distribution</b>             | <b>NA</b>    | <b>NA</b>     | <b>NA</b>     | <b>NA</b>      | <b>NA</b>     | <b>NA</b>     | <b>NA</b>      | <b>NA</b>      |
| <b>Total</b>                    | <b>(0.0)</b> | <b>(58.0)</b> | <b>(69.1)</b> | <b>(108.7)</b> | <b>(87.2)</b> | <b>(95.9)</b> | <b>(110.1)</b> | <b>(108.9)</b> |

NA Not applicable

Note: Totals may not sum due to independent rounding.

**Table A-145: National CH<sub>4</sub> Potential Emission Estimates from the Natural Gas Production Stage, and Reductions from the Natural Gas STAR Program and Regulations (kt)**

| Activity  | 1990  | 2005    | 2009    | 2010    | 2011    | 2012    | 2013    | 2014    |
|---|-------|---------|---------|---------|---------|---------|---------|---------|
| <b>Normal Fugitives</b>                         |       |         |         |         |         |         |         |         |
| <b>Gas Wells</b>                                |       |         |         |         |         |         |         |         |
| Associated Gas Wells                            | IE    | IE      | IE      | IE      | IE      | IE      | IE      | IE      |
| Non-associated Gas Wells (less fractured wells) | 12.8  | 17.1    | 18.6    | 18.4    | 18.5    | 18.1    | 17.8    | 17.8    |
| Gas Wells with Hydraulic Fracturing             | 7.8   | 25.3    | 34.3    | 34.7    | 35.5    | 35.5    | 35.0    | 35.1    |
| <b>Well Pad Equipment</b>                       |       |         |         |         |         |         |         |         |
| Heaters   | 13.0  | 21.0    | 23.8    | 24.4    | 24.6    | 24.5    | 24.1    | 24.0    |
| Separators                                      | 42.4  | 95.4    | 119.9   | 120.9   | 122.7   | 122.1   | 120.0   | 118.6   |
| Dehydrators                                     | 14.4  | 12.1    | 9.0     | 8.9     | 8.6     | 8.6     | 8.4     | 8.4     |
| Meters/Piping                                   | 49.4  | 90.2    | 107.4   | 108.8   | 110.4   | 109.8   | 108.0   | 107.2   |
| Compressors                                     | 33.9  | 74.4    | 92.4    | 96.2    | 98.1    | 97.4    | 96.0    | 96.2    |
| <b>Gathering and Boosting</b>                   |       |         |         |         |         |         |         |         |
| Gathering and Boosting Stations*                | 956.9 | 1,107.2 | 1,364.2 | 1,431.8 | 1,584.0 | 1,697.4 | 1,730.2 | 1,864.9 |
| Pipeline Leaks                                  | 89.8  | 144.9   | 169.7   | 171.1   | 173.2   | 172.4   | 170.2   | 169.7   |
| <b>Vented and Combusted</b>                     |       |         |         |         |         |         |         |         |
| Drilling, Well Completion, and Well Workover    |       |         |         |         |         |         |         |         |

|  |                |                |                |                |                |                |                |                |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Gas Well Completions without Hydraulic Fracturing                    | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            | 0.0            |
| Gas Well Workovers without Hydraulic Fracturing                      | 0.3            | 0.4            | 0.5            | 0.5            | 0.5            | 0.5            | 0.4            | 0.4            |
| Hydraulic Fracturing Completions and Workovers that vent*            | 144.3          | 379.4          | 216.0          | 197.8          | 170.9          | 114.5          | 61.7           | 65.9           |
| Flared Hydraulic Fracturing Completions and Workovers*               | 2.1            | 7.3            | 5.5            | 5.5            | 6.8            | 3.4            | 4.1            | 2.7            |
| Hydraulic Fracturing Completions and Workovers with RECs*            | 0.0            | 7.6            | 10.3           | 11.4           | 12.6           | 11.1           | 10.2           | 3.4            |
| Hydraulic Fracturing Completions and Workovers with RECs that flare* | 0.0            | 3.8            | 5.2            | 5.7            | 6.3            | 9.3            | 10.3           | 9.7            |
| Well Drilling  | 0.7            | 1.6            | 1.0            | 1.0            | 1.0            | 1.0            | 1.0            | 1.0            |
| <i>Produced Water from Coal Bed Methane</i>                          |                |                |                |                |                |                |                |                |
| Powder River   | 0.0            | 50.0           | 54.0           | 50.6           | 47.2           | 47.0           | 47.3           | 47.6           |
| Black Warrior  | 2.7            | 9.9            | 12.1           | 12.3           | 12.7           | 12.8           | 12.8           | 12.8           |
| <b>Normal Operations</b>   |                |                |                |                |                |                |                |                |
| Pneumatic Controller Vents*  | 556.3          | 1,079.3        | 1,261.7        | 1,249.3        | 1,229.7        | 1,245.3        | 1,259.8        | 1,105.1        |
| Low Bleed  | 0.0            | 23.0           | 40.0           | 43.5           | 47.0           | 40.7           | 24.5           | 38.3           |
| High Bleed   | 335.9          | 483.7          | 469.6          | 436.4          | 398.9          | 352.9          | 193.9          | 133.5          |
| Intermittent Bleed   | 220.5          | 572.5          | 752.0          | 769.4          | 783.9          | 851.8          | 1041.4         | 933.3          |
| Chemical Injection Pumps*  | 29.2           | 96.0           | 129.5          | 133.4          | 131.5          | 130.6          | 128.7          | 128.9          |
| Kimray Pumps   | 166.5          | 139.9          | 109.1          | 105.4          | 102.8          | 102.2          | 100.7          | 100.9          |
| Dehydrator Vents   | 51.9           | 43.6           | 34.0           | 32.9           | 32.1           | 31.9           | 31.4           | 31.4           |
| <b>Condensate Tank Vents</b>   |                |                |                |                |                |                |                |                |
| Condensate Tanks without Control Devices                             | 77.7           | 99.3           | 137.6          | 260.7          | 191.1          | 215.9          | 260.2          | 253.1          |
| Condensate Tanks with Control Devices*                               | 15.5           | 19.9           | 27.5           | 52.1           | 38.2           | 43.2           | 52.0           | 50.6           |
| <b>Compressor Exhaust Vented</b>                                     |                |                |                |                |                |                |                |                |
| Gas Engines  | 116.9          | 208.5          | 253.0          | 254.3          | 257.4          | 255.3          | 250.9          | 249.8          |
| <b>Well Clean Ups (Liquids Unloading)</b>                            |                |                |                |                |                |                |                |                |
| Liquids Unloading - Vent Using Plungers*                             | 0.0            | 69.2           | 106.6          | 113.2          | 115.0          | 114.1          | 112.1          | 112.6          |
| Liquids Unloading - Vent Without Using Plungers*                     | 805.9          | 636.9          | 276.2          | 149.0          | 151.6          | 151.0          | 148.4          | 148.1          |
| <b>Blowdowns</b>   |                |                |                |                |                |                |                |                |
| Vessel BD  | 0.3            | 0.5            | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            |
| Pipeline BD  | 1.4            | 2.3            | 2.7            | 2.7            | 2.8            | 2.7            | 2.7            | 2.7            |
| Compressor BD  | 1.3            | 2.9            | 3.6            | 3.7            | 3.8            | 3.8            | 3.7            | 3.7            |
| Compressor Starts  | 2.9            | 6.4            | 8.0            | 8.3            | 8.5            | 8.4            | 8.3            | 8.3            |
| <b>Upsets</b>  |                |                |                |                |                |                |                |                |
| Pressure Relief Valves   | 0.3            | 0.6            | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            |
| Mishaps  | 0.8            | 1.2            | 1.5            | 1.5            | 1.5            | 1.5            | 1.5            | 1.5            |
| <b>Offshore</b>  |                |                |                |                |                |                |                |                |
| Shallow water Gas Platforms (GoM and Pacific)                        | 134.8          | 149.3          | 129.1          | 123.5          | 123.5          | 123.5          | 123.5          | 123.5          |
| Deepwater Gas Platforms (GoM and Pacific)                            | 6.2            | 24.1           | 26.4           | 27.1           | 27.1           | 27.1           | 27.1           | 27.1           |
| <b>Regulatory Reductions (kt)</b>                                    | -              | (45.9)         | (55.1)         | (94.3)         | (71.7)         | (79.6)         | (93.6)         | (91.4)         |
| <b>Voluntary Reductions (kt)</b>                                     | (3.5)          | (256.2)        | (349.1)        | (393.9)        | (427.6)        | (421.4)        | (446.7)        | (483.2)        |
| <b>Total Reductions (kt)</b>   | (3.5)          | (302.1)        | (404.2)        | (488.2)        | (499.3)        | (501.0)        | (540.4)        | (574.6)        |
| <b>Total Potential Emissions (kt)</b>                                | <b>3,338.7</b> | <b>4,627.7</b> | <b>4,751.9</b> | <b>4,818.5</b> | <b>4,851.3</b> | <b>4,943.1</b> | <b>4,969.8</b> | <b>4,933.8</b> |
| <b>Total Net Emissions (kt)</b>                                      | <b>3,335.1</b> | <b>4,325.5</b> | <b>4,347.7</b> | <b>4,330.4</b> | <b>4,352.0</b> | <b>4,442.1</b> | <b>4,429.4</b> | <b>4,359.2</b> |

IE: Included Elsewhere. These emissions are included in the Petroleum Systems estimates.

\*The values in this table are net emissions for these sources.

Note: Totals may not sum due to independent rounding.

**Table A-146: Potential CH<sub>4</sub> Emission Estimates from the Natural Gas Processing Plants, and Reductions from the Natural Gas STAR Program and Regulations (kt)**

| Activity                | 1990 | 2005 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------------------|------|------|------|------|------|------|------|------|
| <b>Normal Fugitives</b> |      |      |      |      |      |      |      |      |
| Plants                  | 42.3 | 31.5 | 32.2 | 32.5 | 33.7 | 33.7 | 36.1 | 37.1 |

| Activity                              | 1990         | 2005         | 2009         | 2010         | 2011           | 2012           | 2013           | 2014           |
|---------------------------------------|--------------|--------------|--------------|--------------|----------------|----------------|----------------|----------------|
| Reciprocating Compressors             | 324.9        | 327.9        | 377.6        | 390.9        | 420.9          | 442.1          | 445.6          | 473.8          |
| Centrifugal Compressors (wet seals)   | 240.3        | 229.2        | 232.9        | 233.9        | 236.1          | 237.7          | 237.9          | 240.0          |
| Centrifugal Compressors (dry seals)   | -            | 6.5          | 22.7         | 27.1         | 36.8           | 43.8           | 44.9           | 54.1           |
| <b>Vented and Combusted</b>           |              |              |              |              |                |                |                |                |
| <i>Compressor Exhaust</i>             |              |              |              |              |                |                |                |                |
| Gas Engines                           | 137.1        | 138.3        | 159.3        | 164.9        | 177.6          | 186.5          | 188.0          | 199.9          |
| Gas Turbines                          | 3.9          | 3.9          | 4.5          | 4.6          | 5.0            | 5.3            | 5.3            | 5.6            |
| AGR Vents                             | 16.5         | 12.3         | 12.5         | 12.7         | 13.1           | 13.1           | 14.1           | 14.5           |
| Kimray Pumps                          | 3.7          | 3.7          | 4.3          | 4.4          | 4.8            | 5.0            | 5.0            | 5.4            |
| Dehydrator Vents                      | 22.7         | 22.9         | 26.3         | 27.3         | 29.4           | 30.8           | 31.1           | 33.0           |
| Pneumatic Controllers                 | 2.4          | 1.8          | 1.8          | 1.9          | 1.9            | 1.9            | 2.1            | 2.1            |
| <b>Routine Maintenance</b>            |              |              |              |              |                |                |                |                |
| Blowdowns/Venting                     | 59.5         | 44.3         | 45.3         | 45.7         | 47.4           | 47.4           | 50.8           | 52.2           |
| <b>Regulatory Reductions (kt)</b>     | -            | (12.1)       | (13.9)       | (14.4)       | (15.5)         | (16.3)         | (16.4)         | (17.5)         |
| <b>Voluntary Reductions (kt)</b>      | (1.5)        | (155.5)      | (137.6)      | (214.3)      | (140.4)        | (140.4)        | (140.7)        | (140.8)        |
| <b>Total Reductions (kt)</b>          | (1.5)        | (167.6)      | (151.5)      | (228.8)      | (155.9)        | (156.8)        | (157.2)        | (158.3)        |
| <b>Total Potential Emissions (kt)</b> | <b>853.2</b> | <b>822.2</b> | <b>919.5</b> | <b>946.0</b> | <b>1,006.6</b> | <b>1,047.3</b> | <b>1,060.9</b> | <b>1,117.9</b> |
| <b>Total Net Emissions (kt)</b>       | <b>851.8</b> | <b>654.6</b> | <b>768.0</b> | <b>717.2</b> | <b>850.7</b>   | <b>890.5</b>   | <b>903.7</b>   | <b>959.6</b>   |

Note: Totals may not sum due to independent rounding.

**Table A-147: Potential CH<sub>4</sub> Emission Estimates from the Natural Gas Transmission and Storage, and Reductions from the Natural Gas STAR Program and Regulations (kt)**

| Activity                                   | 1990    | 2005  | 2009  | 2010  | 2011  | 2012  | 2013  | 2014  |
|--|---------|-------|-------|-------|-------|-------|-------|-------|
| <b>Fugitives</b>                           |         |       |       |       |       |       |       |       |
| Pipeline Leaks                             | 3.2     | 3.3   | 3.3   | 3.3   | 3.3   | 3.3   | 3.3   | 3.3   |
| <b>Compressor Stations (Transmission)*</b> |         |       |       |       |       |       |       |       |
| Station Total Emissions                    | 1,098.4 | 667.5 | 549.6 | 521.1 | 493.1 | 496.0 | 536.0 | 573.2 |
| Station + Compressor Fugitive Emissions    | -       | -     | -     | -     | 94.8  | 101.6 | 109.7 | 117.4 |
| Reciprocating Compressor                   | -       | -     | -     | -     | 301.6 | 293.7 | 317.2 | 339.4 |
| Centrifugal Compressor (wet seals)         | -       | -     | -     | -     | 46.9  | 51.1  | 55.9  | 59.1  |
| Centrifugal Compressor (dry seals)         | -       | -     | -     | -     | 49.8  | 49.7  | 53.2  | 57.4  |
| <b>Compressor Stations (Storage)*</b>      |         |       |       |       |       |       |       |       |
| Station Total Emissions                    | 245.6   | 163.9 | 146.0 | 140.9 | 136.5 | 131.0 | 131.6 | 131.6 |
| Station + Compressor Fugitive Emissions    | -       | -     | -     | -     | -     | 25.1  | 25.3  | 25.3  |
| Reciprocating Compressor                   | -       | -     | -     | -     | -     | 105.8 | 106.4 | 106.4 |
| Wells (Storage)                            | 13.6    | 14.9  | 14.8  | 14.8  | 14.6  | 12.8  | 15.1  | 15.7  |
| M&R (Trans. Co. Interconnect)              | 72.8    | 74.9  | 75.9  | 76.0  | 76.1  | 75.6  | 75.5  | 75.2  |
| M&R (Farm Taps + Direct Sales)             | 16.9    | 17.4  | 17.6  | 17.6  | 17.7  | 17.6  | 17.5  | 17.5  |
| <b>Vented and Combusted</b>                |         |       |       |       |       |       |       |       |
| <b>Normal Operation</b>                    |         |       |       |       |       |       |       |       |
| Dehydrator vents (Transmission)            | 2.0     | 1.8   | 1.7   | 1.7   | 1.7   | 1.8   | 2.0   | 2.1   |
| Dehydrator vents (Storage)                 | 4.2     | 4.6   | 4.6   | 4.6   | 4.5   | 4.0   | 4.7   | 4.9   |
| <b>Compressor Exhaust</b>                  |         |       |       |       |       |       |       |       |
| Engines (Transmission)                     | 176.9   | 203.1 | 211.4 | 222.3 | 225.9 | 235.6 | 241.3 | 246.4 |
| Turbines (Transmission)                    | 1.0     | 1.2   | 1.2   | 1.3   | 1.3   | 1.3   | 1.4   | 1.4   |
| Engines (Storage)                          | 21.3    | 23.4  | 23.2  | 23.2  | 22.9  | 20.1  | 23.8  | 24.7  |
| Turbines (Storage)                         | 0.2     | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   | 0.2   |
| Generators (Engines)                       | 8.7     | 9.9   | 10.3  | 10.9  | 11.1  | 11.5  | 11.8  | 12.1  |
| Generators (Turbines)                      | 0.0     | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |
| <b>Pneumatic Devices Trans + Stor*</b>     |         |       |       |       |       |       |       |       |
| Pneumatic Devices Transmission             | 213.1   | 73.0  | 42.8  | 36.2  | 30.0  | 21.5  | 26.6  | 27.8  |
| (High Bleed)                               | -       | -     | -     | -     | 19.5  | 10.0  | 11.9  | 12.1  |
| (Intermittent Bleed)                       | -       | -     | -     | -     | 10.0  | 10.8  | 13.9  | 14.9  |
| (Low Bleed)                                | -       | -     | -     | -     | 0.5   | 0.6   | 0.7   | 0.8   |
| Pneumatic Devices Storage                  | 44.4    | 35.3  | 29.5  | 27.9  | 26.3  | 26.0  | 30.1  | 29.4  |
| (High Bleed)                               | -       | -     | -     | -     | 22.7  | 21.7  | 25.0  | 23.9  |
| (Intermittent Bleed)                       | -       | -     | -     | -     | 3.2   | 3.8   | 4.5   | 5.0   |
| (Low Bleed)                                | -       | -     | -     | -     | 0.4   | 0.4   | 0.6   | 0.5   |
| <b>Routine Maintenance/Upsets</b>          |         |       |       |       |       |       |       |       |
| Pipeline venting                           | 178.0   | 183.2 | 185.7 | 185.8 | 186.0 | 184.9 | 184.6 | 183.9 |
| <b>Station venting Trans + Storage</b>     |         |       |       |       |       |       |       |       |
| Station Venting Transmission               | 145.5   | 131.0 | 126.6 | 125.5 | 124.4 | 133.2 | 143.9 | 154.0 |

|                                       |         |         |         |         |         |         |         |         |
|---------------------------------------|---------|---------|---------|---------|---------|---------|---------|---------|
| Station Venting Storage               | 30.3    | 28.8    | 29.4    | 29.5    | 29.7    | 29.7    | 29.9    | 29.9    |
| <b>LNG Storage</b>                    |         |         |         |         |         |         |         |         |
| LNG Stations                          | 9.2     | 10.6    | 10.6    | 10.6    | 10.6    | 10.6    | 10.6    | 10.6    |
| LNG Reciprocating Compressors         | 34.5    | 40.1    | 40.1    | 40.1    | 40.1    | 40.1    | 40.1    | 40.1    |
| LNG Centrifugal Compressors           | 11.8    | 13.8    | 13.8    | 13.8    | 13.8    | 13.8    | 13.8    | 13.8    |
| <b>LNG Compressor Exhaust</b>         |         |         |         |         |         |         |         |         |
| LNG Engines                           | 2.6     | 2.7     | 2.7     | 2.7     | 2.7     | 2.7     | 2.7     | 2.7     |
| LNG Turbines                          | 0.0     | 0.0     | 0.0     | 0.0     | 0.0     | 0.0     | 0.0     | 0.0     |
| LNG Station venting                   | 5.1     | 5.9     | 5.9     | 5.9     | 5.9     | 5.9     | 5.9     | 5.9     |
| <b>LNG Import Terminals</b>           |         |         |         |         |         |         |         |         |
| LNG Stations                          | 0.2     | 0.4     | 1.2     | 1.2     | 1.2     | 1.2     | 1.2     | 1.3     |
| LNG Reciprocating Compressors         | 1.0     | 2.0     | 5.6     | 5.6     | 5.6     | 5.6     | 5.6     | 6.1     |
| LNG Centrifugal Compressors           | 0.3     | 0.5     | 1.4     | 1.4     | 1.4     | 1.4     | 1.4     | 1.5     |
| <b>LNG Compressor Exhaust</b>         |         |         |         |         |         |         |         |         |
| LNG Engines                           | 1.7     | 12.2    | 8.8     | 8.4     | 6.9     | 3.6     | 2.1     | 1.4     |
| LNG Turbines                          | 0.0     | 0.1     | 0.1     | 0.0     | 0.0     | 0.0     | 0.0     | 0.0     |
| LNG Station venting                   | 0.1     | 0.2     | 0.6     | 0.6     | 0.6     | 0.6     | 0.6     | 0.7     |
| <b>Regulatory Reductions (kt)</b>     | -       | -       | -       | -       | -       | -       | -       | -       |
| <b>Voluntary Reductions (kt)</b>      | -       | (496.2) | (379.0) | (433.2) | (341.9) | (375.4) | (331.8) | (335.0) |
| <b>Total Reductions (kt)</b>          | -       | (496.2) | (379.0) | (433.2) | (341.9) | (375.4) | (331.8) | (335.0) |
| <b>Total Potential Emissions (kt)</b> | 2,342.6 | 1,725.9 | 1,564.6 | 1,533.2 | 1,493.8 | 1,491.8 | 1,563.3 | 1,617.3 |
| <b>Total Net Emissions (kt)</b>       | 2,342.6 | 1,229.7 | 1,185.7 | 1,100.0 | 1,151.9 | 1,116.4 | 1,231.5 | 1,282.3 |

\*The values in this table are net emissions for these sources.

Note: Totals may not sum due to independent rounding.

**Table A-14B: Potential CH<sub>4</sub> Emission Estimates from the Natural Gas Distribution Stage, and Reductions from the Natural Gas STAR Program, and Regulations (kt)**

| Activity                              | 1990    | 2005   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   |
|---------------------------------------|---------|--------|--------|--------|--------|--------|--------|--------|
| <b>Normal Fugitives</b>               |         |        |        |        |        |        |        |        |
| <b>Pipeline Leaks*</b>                |         |        |        |        |        |        |        |        |
| Mains - Cast Iron                     | 268.0   | 88.9   | 54.1   | 46.3   | 39.0   | 37.5   | 35.8   | 34.0   |
| Mains - Unprotected steel             | 231.2   | 91.3   | 67.4   | 62.6   | 56.0   | 54.9   | 52.2   | 50.4   |
| Mains - Protected steel               | 27.5    | 41.6   | 45.3   | 46.3   | 47.3   | 47.2   | 47.1   | 47.1   |
| Mains - Plastic                       | 59.4    | 44.8   | 28.8   | 23.8   | 18.7   | 19.1   | 19.5   | 19.9   |
| Services - Unprotected steel          | 250.0   | 111.6  | 85.7   | 65.3   | 60.0   | 56.7   | 53.2   | 49.7   |
| Services Protected steel              | 67.2    | 32.4   | 23.5   | 21.5   | 19.8   | 19.4   | 19.1   | 18.9   |
| Services - Plastic                    | 3.4     | 9.1    | 10.9   | 11.2   | 11.7   | 11.9   | 12.1   | 12.3   |
| Services - Copper                     | 7.8     | 6.1    | 5.3    | 5.3    | 5.2    | 4.9    | 4.8    | 4.6    |
| <b>Meter/Regulator (City Gates)*</b>  |         |        |        |        |        |        |        |        |
| M&R >300                              | 140.6   | 64.9   | 20.2   | 14.4   | 9.4    | 9.4    | 10.1   | 8.5    |
| M&R 100-300                           | 272.8   | 123.9  | 37.4   | 25.9   | 15.9   | 15.9   | 17.2   | 14.4   |
| M&R <100                              | 6.6     | 8.3    | 5.6    | 5.6    | 6.2    | 6.2    | 6.7    | 5.6    |
| Reg >300                              | 138.4   | 59.3   | 15.8   | 9.8    | 4.2    | 4.2    | 4.5    | 3.8    |
| R-Vault >300                          | 0.7     | 0.4    | 0.2    | 0.2    | 0.1    | 0.1    | 0.1    | 0.1    |
| Reg 100-300                           | 104.7   | 43.9   | 11.1   | 6.5    | 2.1    | 2.1    | 2.2    | 1.9    |
| R-Vault 100-300                       | 0.2     | 0.4    | 0.3    | 0.3    | 0.2    | 0.2    | 0.2    | 0.2    |
| Reg 40-100                            | 8.1     | 9.8    | 6.5    | 6.5    | 7.1    | 7.1    | 7.7    | 6.4    |
| R-Vault 40-100                        | 0.6     | 2.0    | 1.6    | 1.7    | 1.4    | 1.5    | 1.3    | 1.0    |
| Reg <40                               | 0.4     | 0.6    | 0.4    | 0.4    | 0.4    | 0.4    | 0.4    | 0.4    |
| <b>Customer Meters</b>                |         |        |        |        |        |        |        |        |
| Residential                           | 58.4    | 75.0   | 77.2   | 77.4   | 77.9   | 78.4   | 78.9   | 79.4   |
| Commercial/Industry                   | 43.4    | 52.6   | 53.8   | 53.5   | 53.6   | 54.0   | 54.2   | 54.6   |
| <b>Vented</b>                         |         |        |        |        |        |        |        |        |
| <b>Routine Maintenance</b>            |         |        |        |        |        |        |        |        |
| Pressure Relief Valve Releases        | 0.9     | 1.1    | 1.2    | 1.2    | 1.2    | 1.2    | 1.2    | 1.2    |
| Pipeline Blowdown                     | 3.0     | 3.9    | 4.1    | 4.1    | 4.2    | 4.2    | 4.2    | 4.3    |
| <b>Upsets</b>                         |         |        |        |        |        |        |        |        |
| Mishaps (Dig-ins)                     | 47.4    | 60.1   | 63.9   | 64.4   | 64.9   | 65.5   | 65.8   | 66.4   |
| <b>Regulatory Reductions (kt)</b>     | -       | -      | -      | -      | -      | -      | -      | -      |
| <b>Voluntary Reductions (kt)</b>      | -       | (48.3) | (63.9) | (54.5) | (58.0) | (45.1) | (40.5) | (40.8) |
| <b>Total Reductions (kt)</b>          | -       | (48.3) | (63.9) | (54.5) | (58.0) | (45.1) | (40.5) | (40.8) |
| <b>Total Potential Emissions (kt)</b> | 1,740.7 | 931.9  | 620.2  | 554.1  | 506.5  | 501.9  | 498.5  | 484.9  |

|                                 |                |              |              |              |              |              |              |              |
|---------------------------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Total Net Emissions (kt)</b> | <b>1,740.7</b> | <b>883.6</b> | <b>556.3</b> | <b>499.6</b> | <b>448.5</b> | <b>456.7</b> | <b>458.0</b> | <b>444.1</b> |
|---------------------------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|

\*The values in this table are net emissions for these sources.

Note: Totals may not sum due to independent rounding.

**Table A-149: Net Emissions for Select Sources (kt)**

| Stage/Activity                                 | 1990           | 2005           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Production</b>                              | <b>3,335.1</b> | <b>4,325.5</b> | <b>4,347.7</b> | <b>4,330.4</b> | <b>4,352.0</b> | <b>4,442.1</b> | <b>4,429.4</b> | <b>4,359.2</b> |
| Gathering Stations                             | 956.9          | 1,107.2        | 1,364.2        | 1,431.8        | 1,584.0        | 1,697.4        | 1,730.2        | 1,864.9        |
| Hydraulic Fracturing Completions and Workovers | 146.5          | 398.2          | 237.0          | 220.5          | 196.6          | 138.3          | 86.4           | 81.7           |
| Liquids Unloading                              | 805.9          | 706.1          | 382.8          | 262.2          | 266.6          | 265.1          | 260.5          | 260.6          |
| Dehydrator Vents                               | 51.9           | 29.6           | 23.1           | 22.3           | 21.8           | 21.6           | 21.3           | 21.4           |
| Condensate Tanks                               | 93.2           | 87.3           | 121.0          | 229.1          | 167.9          | 189.8          | 228.6          | 222.4          |
| Pipeline Leaks                                 | 89.8           | 142.5          | 169.7          | 171.1          | 173.2          | 172.4          | 170.2          | 169.7          |
| Pneumatic Controller Vents                     | 556.3          | 1,079.3        | 1,261.7        | 1,249.3        | 1,229.7        | 1,245.3        | 1,259.8        | 1,105.1        |
| Chemical Injection Pumps                       | 29.2           | 96.0           | 129.5          | 133.4          | 131.5          | 130.6          | 128.7          | 128.9          |
| Gas Engines                                    | 116.9          | 110.6          | 130.0          | 120.9          | 120.2          | 115.3          | 110.4          | 108.8          |
| Compressor Starts                              | 2.9            | 6.2            | 7.6            | 7.8            | 8.0            | 7.9            | 7.8            | 7.8            |
| Other Production                               | 485.6          | 562.5          | 521.1          | 482.0          | 452.5          | 458.4          | 425.5          | 387.9          |
| <b>Processing</b>                              | <b>851.8</b>   | <b>654.6</b>   | <b>768.0</b>   | <b>717.2</b>   | <b>850.7</b>   | <b>890.5</b>   | <b>903.7</b>   | <b>959.6</b>   |
| Fugitives                                      |                |                |                |                |                |                |                |                |
| Reciprocating Compressors                      | 324.9          | 327.9          | 377.6          | 389.9          | 420.9          | 442.1          | 445.4          | 473.7          |
| Gas Engines                                    | 137.1          | 137.2          | 153.3          | 158.9          | 171.5          | 180.4          | 181.9          | 193.8          |
| AGR Vents                                      | 16.5           | 12.3           | 12.5           | 12.7           | 13.1           | 13.1           | 14.1           | 14.5           |
| Dehydrator Vents                               | 21.3           | 8.6            | 3.1            | 3.5            | 4.5            | 5.2            | 5.3            | 6.2            |
| Other Processing                               | 351.9          | 168.6          | 221.5          | 152.3          | 240.8          | 249.7          | 257.0          | 271.4          |
| <b>Transmission and Storage</b>                | <b>2,342.6</b> | <b>1,229.7</b> | <b>1,185.7</b> | <b>1,100.0</b> | <b>1,151.9</b> | <b>1,116.4</b> | <b>1,231.5</b> | <b>1,282.3</b> |
| Transmission Stations                          | 1098.4         | 667.5          | 549.6          | 521.1          | 493.1          | 496.0          | 536.0          | 573.2          |
| Station Fugitives                              | NA             | NA             | NA             | NA             | 94.8           | 101.6          | 109.7          | 117.4          |
| Reciprocating Compressors                      | NA             | NA             | NA             | NA             | 301.6          | 293.7          | 317.2          | 339.4          |
| Centrifugal Compressors (wet seals)            | NA             | NA             | NA             | NA             | 46.9           | 51.1           | 55.9           | 59.1           |
| Centrifugal compressors (dry seals)            | NA             | NA             | NA             | NA             | 49.8           | 49.7           | 53.2           | 57.4           |
| Transmission pneumatic controllers             | 213.1          | 73.0           | 42.8           | 36.2           | 30.0           | 21.5           | 26.6           | 27.8           |
| Storage stations                               | 245.6          | 163.9          | 146.0          | 140.9          | 136.5          | 131.0          | 131.6          | 131.6          |
| Station fugitives                              | NA             | NA             | NA             | NA             | NA             | 25.1           | 25.3           | 25.3           |
| Reciprocating compressors                      | NA             | NA             | NA             | NA             | NA             | 105.8          | 106.4          | 106.4          |
| Storage pneumatic controllers                  | 44.4           | 35.3           | 29.5           | 27.9           | 26.3           | 26.0           | 30.1           | 29.4           |
| Engines (Transmission)                         | 176.9          | 119.9          | 95.2           | 102.9          | 104.2          | 111.6          | 114.4          | 116.6          |
| Pipeline Vents (Transmission)                  | 178.0          | 58.3           | 97.0           | 57.6           | 127.0          | 84.8           | 125.2          | 124.6          |
| Other Transmission and Storage                 | 386.2          | 111.8          | 225.6          | 213.4          | 234.8          | 245.5          | 267.6          | 279.1          |
| <b>Distribution</b>                            | <b>1,740.7</b> | <b>883.6</b>   | <b>556.3</b>   | <b>499.6</b>   | <b>448.5</b>   | <b>456.7</b>   | <b>458.0</b>   | <b>444.1</b>   |
| <b>Pipeline Leaks</b>                          | <b>914.6</b>   | <b>425.8</b>   | <b>320.9</b>   | <b>282.4</b>   | <b>257.6</b>   | <b>251.6</b>   | <b>243.7</b>   | <b>236.9</b>   |
| <b>M&amp;R Stations</b>                        | <b>673.1</b>   | <b>313.4</b>   | <b>99.2</b>    | <b>71.2</b>    | <b>47.1</b>    | <b>47.1</b>    | <b>50.4</b>    | <b>42.1</b>    |
| Mishaps (Dig-ins)                              | 47.4           | 59.8           | 63.1           | 63.6           | 60.3           | 64.7           | 65.0           | 65.6           |
| Other Distribution                             | 105.6          | 84.6           | 73.1           | 82.4           | 83.5           | 93.3           | 98.9           | 99.5           |
| <b>Total</b>                                   | <b>8,270.2</b> | <b>7,093.5</b> | <b>6,857.6</b> | <b>6,647.1</b> | <b>6,803.1</b> | <b>6,905.8</b> | <b>7,022.7</b> | <b>7,045.2</b> |

Note: This table presents net emissions for each natural gas system stage, and also presents net emissions for select emissions sources for which disaggregated Gas STAR data and/or regulation reduction data can be matched to an Inventory source category, and sources for which emissions are calculated using net emission factors. In general, the Inventory uses aggregated Gas STAR reductions by natural gas system stage (i.e., production, processing, transmission and storage, and distribution). In some cases, emissions reductions reported to Gas STAR have been matched to potential emissions calculated in the Inventory, to provide a net emissions number for specific emissions sources. Net emission values presented here were calculated by deducting the voluntary reductions (Table A-143) and the regulatory reductions (Table A-144) from the potential emissions values in Table A-145 through Table A-148. Some reported Gas STAR reduction activities are cross-cutting and cover multiple Inventory sources. It is not possible to attribute those reductions to specific Inventory source categories, and they are included in the "Other" category.

**Table A-150: U.S. Production Sector CO<sub>2</sub> Content in Natural Gas by NEMS Region and Formation Type for all years**

| Formation Types   | U.S. Region |              |            |            |                |            |       | Lower-48 States |
|-------------------|-------------|--------------|------------|------------|----------------|------------|-------|-----------------|
|                   | North East  | Midcontinent | Gulf Coast | South West | Rocky Mountain | West Coast |       |                 |
| Conventional      | 0.92%       | 0.79%        | 2.17%      | 3.81%      | 7.95%          | 0.16%      | 3.41% |                 |
| Non-conventional* | 7.42%       | 0.31%        | 0.23%      | NA         | 0.64%          | NA         | 4.83% |                 |
| All types         | 3.04%       | 0.79%        | 2.17%      | 3.81%      | 7.58%          | 0.16%      | 3.45% |                 |

Source: GRI-01/0136 GTI's Gas Resource Database: Unconventional Natural Gas and Gas Composition Databases. Second Edition. August, 2001

\*In GTI, this refers to shale, coal bed methane, and tight geologic formations.

**Table A-151: CO<sub>2</sub> Emission Estimates from the Natural Gas Production Stage (kt)**

| Activity  | 1990 | 2005  | 2009  | 2010  | 2011  | 2012  | 2013  | 2014  |
|---|------|-------|-------|-------|-------|-------|-------|-------|
| <b>Normal Fugitives</b>                           |      |       |       |       |       |       |       |       |
| <b>Gas Wells</b>                                  |      |       |       |       |       |       |       |       |
| Associated Gas Wells                              | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |
| Non-associated Gas Wells (less fractured wells)   | 1.4  | 1.6   | 1.7   | 1.7   | 1.7   | 1.7   | 1.7   | 1.7   |
| Gas Wells with Hydraulic Fracturing               | 0.4  | 1.2   | 1.4   | 1.5   | 1.5   | 1.5   | 1.5   | 1.5   |
| <b>Well Pad Equipment</b>                         |      |       |       |       |       |       |       |       |
| Heaters   | 1.9  | 2.8   | 3.2   | 3.2   | 3.3   | 3.2   | 3.2   | 3.2   |
| Separators  | 6.0  | 14.3  | 18.4  | 18.5  | 18.7  | 18.5  | 18.2  | 18.0  |
| Dehydrators                                       | 1.4  | 1.2   | 1.0   | 1.0   | 0.9   | 0.9   | 0.9   | 0.9   |
| Meters/Piping                                     | 6.7  | 12.3  | 14.9  | 15.0  | 15.2  | 15.1  | 14.8  | 14.7  |
| Compressors                                       | 3.1  | 7.7   | 9.8   | 10.1  | 10.3  | 10.3  | 10.1  | 10.1  |
| <b>Gathering and Boosting</b>                     |      |       |       |       |       |       |       |       |
| Gathering and Boosting Stations                   | 88.7 | 135.4 | 167.1 | 171.6 | 182.6 | 192.6 | 194.5 | 207.5 |
| Pipeline Leaks                                    | 9.9  | 16.4  | 19.5  | 19.6  | 19.8  | 19.7  | 19.4  | 19.3  |
| <b>Vented and Combusted</b>                       |      |       |       |       |       |       |       |       |
| <b>Drilling and Well Completions</b>              |      |       |       |       |       |       |       |       |
| Gas Well Completions without Hydraulic Fracturing | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |
| Gas Well Completions with Hydraulic Fracturing    | 73.6 | 305.5 | 179.9 | 185.0 | 160.8 | 98.6  | 75.0  | 65.8  |
| Well Drilling                                     | 0.1  | 0.2   | 0.1   | 0.1   | 0.1   | 0.1   | 0.1   | 0.1   |
| <b>Produced Water from Coal Bed Methane</b>       |      |       |       |       |       |       |       |       |
| Powder River                                      | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |
| Black Warrior                                     | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |
| <b>Normal Operations</b>                          |      |       |       |       |       |       |       |       |
| Pneumatic Device Vents                            | 50.3 | 119.2 | 138.4 | 136.4 | 134.0 | 136.2 | 137.4 | 120.0 |
| Chemical Injection Pumps                          | 2.6  | 10.6  | 14.2  | 14.6  | 14.4  | 14.3  | 14.1  | 14.0  |
| Kimray Pumps                                      | 16.4 | 13.7  | 11.3  | 11.0  | 10.8  | 10.8  | 10.6  | 10.6  |
| Dehydrator Vents                                  | 5.1  | 4.3   | 3.5   | 3.4   | 3.4   | 3.4   | 3.3   | 3.3   |
| <b>Condensate Tank Vents</b>                      |      |       |       |       |       |       |       |       |
| Condensate Tanks without Control Devices          | 10.3 | 10.3  | 11.6  | 15.7  | 17.2  | 21.0  | 24.9  | 25.4  |
| Condensate Tanks with Control Devices             | 2.1  | 2.1   | 2.3   | 3.1   | 3.4   | 4.2   | 5.0   | 5.1   |
| <b>Compressor Exhaust Vented</b>                  |      |       |       |       |       |       |       |       |
| Gas Engines                                       | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |
| <b>Well Workovers and Clean Ups</b>               |      |       |       |       |       |       |       |       |
| Gas Well Workovers without Hydraulic Fracturing   | 0.0  | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   | 0.0   |

|   |                |                |                 |                 |                 |                 |                 |                 |
|---|----------------|----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Gas Well Workovers with Hydraulic Fracturing                | 15.3           | 44.3           | 57.1            | 58.3            | 44.1            | 26.1            | 26.1            | 26.1            |
| Well Clean Ups (LP Gas Wells) - Vent Using Plungers         | 0.0            | 8.4            | 13.1            | 13.9            | 14.1            | 14.1            | 13.8            | 13.8            |
| Well Clean Ups (LP Gas Wells) - Vent Without Using Plungers | 235.8          | 170.4          | 64.1            | 26.1            | 26.5            | 26.3            | 25.8            | 25.7            |
| <b>Blowdowns</b>  |                |                |                 |                 |                 |                 |                 |                 |
| Vessel BD   | 0.0            | 0.1            | 0.1             | 0.1             | 0.1             | 0.1             | 0.1             | 0.1             |
| Pipeline BD   | 0.2            | 0.3            | 0.3             | 0.3             | 0.3             | 0.3             | 0.3             | 0.3             |
| Compressor BD   | 0.1            | 0.3            | 0.4             | 0.4             | 0.4             | 0.4             | 0.4             | 0.4             |
| Compressor Starts   | 0.3            | 0.7            | 0.8             | 0.9             | 0.9             | 0.9             | 0.9             | 0.9             |
| <b>Upsets</b>   |                |                |                 |                 |                 |                 |                 |                 |
| Pressure Relief Valves                                      | 0.0            | 0.1            | 0.1             | 0.1             | 0.1             | 0.1             | 0.1             | 0.1             |
| Mishaps   | 0.1            | 0.1            | 0.2             | 0.2             | 0.2             | 0.2             | 0.2             | 0.2             |
| Flaring Emissions - Onshore                                 | 9,092.7        | 7,193.0        | 9,960.0         | 9,966.9         | 13,084.7        | 12,703.8        | 15,684.1        | 17,628.5        |
| <b>Offshore</b>   |                |                |                 |                 |                 |                 |                 |                 |
| Shallow water Gas Platforms (GoM and Pacific)               | 2.5            | 2.8            | 2.4             | 2.3             | 2.3             | 2.3             | 2.3             | 2.3             |
| Deepwater Gas Platforms (GoM and Pacific)                   | 0.1            | 0.3            | 0.3             | 0.3             | 0.3             | 0.3             | 0.3             | 0.3             |
| Flaring Emissions - Offshore                                | 230.4          | 180.7          | 360.3           | 359.3           | 373.7           | 357.1           | 360.3           | 365.2           |
| <b>Total Emissions (kt)</b>                                 | <b>9,857.4</b> | <b>8,260.0</b> | <b>11,057.7</b> | <b>11,040.8</b> | <b>14,145.8</b> | <b>13,684.0</b> | <b>16,649.3</b> | <b>18,585.0</b> |

<sup>a</sup> Energy use CO<sub>2</sub> emissions not estimated to avoid double counting. NE - Not Estimated.

<sup>b</sup> Emissions are not actually 0, but too small to show at this level of precision.

Note: Totals may not sum due to independent rounding.

**Table A-152: CO<sub>2</sub> Emission Estimates from the Natural Gas Processing Stage (kt)**

| Activity   | 1990            | 2005            | 2009            | 2010            | 2011            | 2012            | 2013            | 2014            |
|--|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| <b>Normal Fugitives</b>  |                 |                 |                 |                 |                 |                 |                 |                 |
| Plants – Before CO <sub>2</sub> removal                              | 2.6             | 1.9             | 1.9             | 2.0             | 2.0             | 2.0             | 2.2             | 2.2             |
| Plants – After CO <sub>2</sub> removal                               | 0.6             | 0.4             | 0.4             | 0.4             | 0.5             | 0.5             | 0.5             | 0.5             |
| Reciprocating Compressors – Before CO <sub>2</sub> removal           | 19.7            | 19.8            | 22.9            | 23.7            | 25.5            | 26.8            | 27.0            | 28.7            |
| Reciprocating Compressors – After CO <sub>2</sub> removal            | 4.4             | 4.4             | 5.1             | 5.3             | 5.7             | 5.9             | 6.0             | 6.4             |
| Centrifugal Compressors (wet seals) – Before CO <sub>2</sub> removal | 14.5            | 13.9            | 14.1            | 14.2            | 14.3            | 14.4            | 14.4            | 14.5            |
| Centrifugal Compressors (wet seals) – After CO <sub>2</sub> removal  | 3.2             | 3.1             | 3.1             | 3.1             | 3.2             | 3.2             | 3.2             | 3.2             |
| Centrifugal Compressors (dry seals) – Before CO <sub>2</sub> removal | -               | 0.4             | 1.4             | 1.6             | 2.2             | 2.6             | 2.7             | 3.3             |
| Centrifugal Compressors (dry seals) – After CO <sub>2</sub> removal  | -               | 0.1             | 0.3             | 0.4             | 0.5             | 0.6             | 0.6             | 0.7             |
| <b>Vented and Combusted</b>  |                 |                 |                 |                 |                 |                 |                 |                 |
| Compressor Exhaust   |                 |                 |                 |                 |                 |                 |                 |                 |
| Gas Engines <sup>a</sup>   | NE              |
| Gas Turbines <sup>a</sup>  | NE              |
| AGR Vents  | 27,708.2        | 21,694.3        | 21,130.5        | 21,286.7        | 21,403.6        | 21,403.6        | 21,690.3        | 23,643.5        |
| Kimray Pumps   | 0.4             | 0.4             | 0.5             | 0.5             | 0.5             | 0.5             | 0.5             | 0.6             |
| Dehydrator Vents   | 2.4             | 2.4             | 2.8             | 2.9             | 3.1             | 3.3             | 3.3             | 3.5             |
| Pneumatic Controllers  | 0.3             | 0.2             | 0.2             | 0.2             | 0.2             | 0.2             | 0.2             | 0.3             |
| <b>Routine Maintenance</b>   |                 |                 |                 |                 |                 |                 |                 |                 |
| Blowdowns/Venting  | 6.4             | 4.7             | 4.8             | 4.9             | 5.1             | 5.1             | 5.4             | 5.6             |
| <b>Total</b>   | <b>27,762.6</b> | <b>21,746.1</b> | <b>21,188.0</b> | <b>21,345.8</b> | <b>21,466.3</b> | <b>21,468.7</b> | <b>21,756.4</b> | <b>23,713.0</b> |

NE – Not Estimated.

<sup>a</sup> Energy use CO<sub>2</sub> emissions not estimated to avoid double counting.

Note: Totals may not sum due to independent rounding.

**Table A-153: CO<sub>2</sub> Emission Estimates from the Natural Gas Transmission and Storage Stage (kt)**

| Activity | 1990 | 2005 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 |
|----------|------|------|------|------|------|------|------|------|
|----------|------|------|------|------|------|------|------|------|

|   |             |             |             |             |             |             |             |             |
|---|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>Fugitives</b>                          |             |             |             |             |             |             |             |             |
| Pipeline Leaks                            | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         |
| <i>Compressor Stations (Transmission)</i> |             |             |             |             |             |             |             |             |
| Station                                   | 31.7        | 19.3        | 15.9        | 15.0        | 14.2        | 14.3        | 15.5        | 16.5        |
| Station + Compressor Fugitive Emissions   | -           | -           | -           | -           | 2.7         | 2.9         | 3.2         | 3.4         |
| Recip Compressor                          | -           | -           | -           | -           | 8.7         | 8.5         | 9.2         | 9.8         |
| Centrifugal Compressor (wet seals)        | -           | -           | -           | -           | 1.4         | 1.5         | 1.6         | 1.7         |
| Centrifugal Compressor (dry seals)        | -           | -           | -           | -           | 1.4         | 1.4         | 1.5         | 1.7         |
| <i>Compressor Stations (Storage)</i>      |             |             |             |             |             |             |             |             |
| Station                                   | 7.1         | 4.7         | 4.2         | 4.1         | 3.9         | 3.8         | 3.8         | 3.8         |
| Station + Compressor Fugitive Emissions   | -           | -           | -           | -           | -           | 0.7         | 0.7         | 0.7         |
| Recip Compressor                          | -           | -           | -           | -           | -           | 3.1         | 3.1         | 3.1         |
| Wells (Storage)                           | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.5         |
| M&R (Trans. Co. Interconnect)             | 2.1         | 2.2         | 2.2         | 2.2         | 2.2         | 2.2         | 2.2         | 2.2         |
| M&R (Farm Taps + Direct Sales)            | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         |
| <b>Vented and Combusted</b>               |             |             |             |             |             |             |             |             |
| <i>Normal Operation</i>                   |             |             |             |             |             |             |             |             |
| Dehydrator vents (Transmission)           | 0.1         | 0.1         | 0.1         | 0.0         | 0.0         | 0.1         | 0.1         | 0.1         |
| Dehydrator vents (Storage)                | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         |
| <i>Compressor Exhaust</i>                 |             |             |             |             |             |             |             |             |
| Engines (Transmission)                    | NE          |
| Turbines (Transmission)                   | NE          |
| Engines (Storage)                         | NE          |
| Turbines (Storage)                        | NE          |
| Generators (Engines)                      | NE          |
| Generators (Turbines)                     | NE          |
| <i>Pneumatic Devices Trans + Stor</i>     |             |             |             |             |             |             |             |             |
| Pneumatic Devices Trans                   | 6.1         | 2.1         | 1.2         | 1.0         | 0.9         | 0.6         | 0.8         | 0.8         |
| (High Bleed)                              | -           | -           | -           | -           | 0.6         | 0.3         | 0.3         | 0.3         |
| (Intermittent Bleed)                      | -           | -           | -           | -           | 0.3         | 0.3         | 0.4         | 0.4         |
| (Low Bleed)                               | -           | -           | -           | -           | 0.0         | 0.0         | 0.0         | 0.0         |
| Pneumatic Devices Storage                 | 1.3         | 1.0         | 0.9         | 0.8         | 0.8         | 0.7         | 0.9         | 0.8         |
| (High Bleed)                              | -           | -           | -           | -           | 0.7         | 0.6         | 0.7         | 0.7         |
| (Intermittent Bleed)                      | -           | -           | -           | -           | 0.1         | 0.1         | 0.1         | 0.1         |
| (Low Bleed)                               | -           | -           | -           | -           | 0.0         | 0.0         | 0.0         | 0.0         |
| <i>Routine Maintenance/Upsets</i>         |             |             |             |             |             |             |             |             |
| Pipeline venting                          | 5.1         | 5.3         | 5.4         | 5.4         | 5.4         | 5.3         | 5.3         | 5.3         |
| <i>Station venting Trans + Storage</i>    |             |             |             |             |             |             |             |             |
| Station Venting Transmission              | 4.2         | 3.8         | 3.7         | 3.6         | 3.6         | 3.8         | 4.2         | 4.4         |
| Station Venting Storage                   | 0.9         | 0.8         | 0.8         | 0.9         | 0.9         | 0.9         | 0.9         | 0.9         |
| <b>LNG Storage</b>                        |             |             |             |             |             |             |             |             |
| LNG Stations                              | 0.3         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         |
| LNG Reciprocating Compressors             | 1.2         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         |
| LNG Centrifugal Compressors               | 0.4         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         |
| <i>LNG Compressor Exhaust</i>             |             |             |             |             |             |             |             |             |
| LNG Engines                               | NE          |
| LNG Turbines                              | NE          |
| LNG Station venting                       | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         |
| <b>LNG Import Terminals</b>               |             |             |             |             |             |             |             |             |
| LNG Stations                              | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| LNG Reciprocating Compressors             | 0.0         | 0.1         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         |
| LNG Centrifugal Compressors               | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.1         |
| <i>LNG Compressor Exhaust</i>             |             |             |             |             |             |             |             |             |
| LNG Engines                               | NE          |
| LNG Turbines                              | NE          |
| LNG Station venting                       | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| <b>Total Emissions (kt)</b>               | <b>61.7</b> | <b>42.8</b> | <b>38.1</b> | <b>36.8</b> | <b>35.6</b> | <b>35.5</b> | <b>37.3</b> | <b>38.7</b> |

NE - Not Estimated

<sup>a</sup> Energy use CO<sub>2</sub> emissions not estimated to avoid double counting.

<sup>2</sup> Emissions are not actually 0, but too small to show at this level of precision.

Note: Totals may not sum due to independent rounding.

**Table A-154: CO<sub>2</sub> Emission Estimates from the Natural Gas Distribution Stage (kt)**

| Activity                            | 1990        | 2005        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|-------------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>Pipeline Leaks</b>               |             |             |             |             |             |             |             |             |
| Mains—Cast Iron                     | 7.7         | 2.6         | 1.6         | 1.3         | 1.1         | 1.1         | 1.0         | 1.0         |
| Mains—Unprotected steel             | 6.7         | 2.6         | 1.9         | 1.8         | 1.6         | 1.6         | 1.5         | 1.5         |
| Mains—Protected steel               | 0.8         | 1.2         | 1.3         | 1.3         | 1.4         | 1.4         | 1.4         | 1.4         |
| Mains—Plastic                       | 1.7         | 1.3         | 0.8         | 0.7         | 0.5         | 0.6         | 0.6         | 0.6         |
| <b>Total Pipeline Miles</b>         |             |             |             |             |             |             |             |             |
| Services—Unprotected steel          | 7.2         | 3.2         | 2.5         | 1.9         | 1.7         | 1.6         | 1.5         | 1.4         |
| Services Protected steel            | 1.9         | 0.9         | 0.7         | 0.6         | 0.6         | 0.6         | 0.6         | 0.5         |
| Services—Plastic                    | 0.1         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.4         | 0.4         |
| Services—Copper                     | 0.2         | 0.2         | 0.2         | 0.2         | 0.1         | 0.1         | 0.1         | 0.1         |
| <b>Meter/Regulator (City Gates)</b> |             |             |             |             |             |             |             |             |
| M&R >300                            | 4.1         | 1.9         | 0.6         | 0.4         | 0.3         | 0.3         | 0.3         | 0.2         |
| M&R 100-300                         | 7.9         | 3.6         | 1.1         | 0.7         | 0.5         | 0.5         | 0.5         | 0.4         |
| M&R <100                            | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         |
| Reg >300                            | 4.0         | 1.7         | 0.5         | 0.3         | 0.1         | 0.1         | 0.1         | 0.1         |
| R-Vault >300                        | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| Reg 100-300                         | 3.0         | 1.3         | 0.3         | 0.2         | 0.1         | 0.1         | 0.1         | 0.1         |
| R-Vault 100-300                     | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| Reg 40-100                          | 0.2         | 0.3         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         | 0.2         |
| R-Vault 40-100                      | 0.0         | 0.1         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| Reg <40                             | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| <b>Customer Meters</b>              |             |             |             |             |             |             |             |             |
| Residential                         | 1.7         | 2.2         | 2.2         | 2.2         | 2.2         | 2.3         | 2.3         | 2.3         |
| Commercial/Industry                 | 1.3         | 1.5         | 1.6         | 1.5         | 1.5         | 1.6         | 1.6         | 1.6         |
| <b>Routine Maintenance</b>          |             |             |             |             |             |             |             |             |
| Pressure Relief Valve Releases      | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         | 0.0         |
| Pipeline Blowdown                   | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         | 0.1         |
| <b>Upsets</b>                       |             |             |             |             |             |             |             |             |
| Mishaps (Dig-ins)                   | 1.4         | 1.7         | 1.8         | 1.9         | 1.9         | 1.9         | 1.9         | 1.9         |
| <b>Total</b>                        | <b>50.2</b> | <b>26.9</b> | <b>17.9</b> | <b>16.0</b> | <b>14.6</b> | <b>14.5</b> | <b>14.4</b> | <b>14.0</b> |

Note: Totals may not sum due to independent rounding.

## References

- AGA (1991 through 1998) Gas Facts. American Gas Association. Washington, DC.
- Alabama (2015) Alabama State Oil and Gas Board. Available online at <<http://www.ogb.state.al.us>>.
- API/ANGA (2012) *Characterizing Pivotal Sources of Methane Emissions from Natural Gas Production – Summary and Analysis of API and ANGA Survey Responses*. Final Report. American Petroleum Institute and America’s Natural Gas Alliance. September 21.
- BOEMRE (2004) Gulfwide Emission Inventory Study for the Regional Haze and Ozone Modeling Effort. OCS Study MMS 2004-072.
- BOEMRE (2011a) Gulf of Mexico Region Offshore Information. Bureau of Ocean Energy Management, Regulation and Enforcement, U.S. Department of Interior.
- BOEMRE (2011b) Pacific OCS Region Offshore Information. Bureau of Ocean Energy Management, Regulation and Enforcement, U.S. Department of Interior.
- BOEMRE (2011c) GOM and Pacific OCS Platform Activity. Bureau of Ocean Energy Management, Regulation and Enforcement, U.S. Department of Interior.
- BOEMRE (2011d) Pacific OCS Region. Bureau of Ocean Energy Management, Regulation and Enforcement, U.S. Department of Interior.
- DrillingInfo (2015) DI Desktop® December 2014 Download. DrillingInfo, Inc.
- EIA (2012a) Formation crosswalk. Energy Information Administration, U.S. Department of Energy, Washington, DC. Provided July 7.
- EIA (2012b) Lease Condensate Production, 1989-2011, Natural Gas Navigator. Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at <[http://www.eia.gov/dnav/ng/ng\\_prod\\_lc\\_sl\\_a.htm](http://www.eia.gov/dnav/ng/ng_prod_lc_sl_a.htm)>.
- EIA (2015a) “Table 1—Summary of natural gas supply and disposition in the United States 2008-2013.” Natural Gas Monthly, Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at <<http://www.eia.doe.gov>>.
- EIA (2015b) “Table 2—Natural Gas Consumption in the United States 2008-2013.” Natural Gas Monthly, Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at <<http://www.eia.doe.gov>>.
- EIA (2015c) “Table 7—Marketed production of natural gas in selected states and the Federal Gulf of Mexico, 2008-2013.” Natural Gas Monthly, Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <<http://www.eia.doe.gov>>.
- EIA (2015d) Number of Natural Gas Consumers – Residential. Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <[http://www.eia.gov/dnav/ng/ng\\_cons\\_num\\_a\\_EPG0\\_VN3\\_Count\\_a.htm](http://www.eia.gov/dnav/ng/ng_cons_num_a_EPG0_VN3_Count_a.htm)>.
- EIA (2015e) Number of Natural Gas Consumers – Commercial. Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <[http://www.eia.gov/dnav/ng/ng\\_cons\\_num\\_a\\_EPG0\\_VN5\\_Count\\_a.htm](http://www.eia.gov/dnav/ng/ng_cons_num_a_EPG0_VN5_Count_a.htm)>.
- EIA (2015f) Number of Natural Gas Consumers – Industrial. Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <[http://www.eia.gov/dnav/ng/ng\\_cons\\_num\\_a\\_EPG0\\_VN7\\_Count\\_a.htm](http://www.eia.gov/dnav/ng/ng_cons_num_a_EPG0_VN7_Count_a.htm)>.
- EIA (2014a) U.S. Imports by Country. Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <<http://www.eia.doe.gov>>.
- EIA (2014b) Natural Gas Gross Withdrawals and Production. Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <<http://www.eia.doe.gov>>.
- EIA (2005) “Table 5—U.S. Crude Oil, Natural Gas, and Natural Gas Liquids Reserves, 1977-2003.” Energy Information Administration, Department of Energy, Washington, DC.
- EIA (2004) *U.S. LNG Markets and Uses*. Energy Information Administration, U.S. Department of Energy, Washington, DC. June 2004. Available online at: <[http://www.eia.doe.gov/pub/oil\\_gas/natural\\_gas/feature\\_articles/2004/lng/lng2004.pdf](http://www.eia.doe.gov/pub/oil_gas/natural_gas/feature_articles/2004/lng/lng2004.pdf)>.

- EIA (2001) "Documentation of the Oil and Gas Supply Module (OGSM)." Energy Information Administration, U.S. Department of Energy, Washington, DC. Available online at: <[http://tonto.eia.doe.gov/FTP/ROOT/modeldoc/m063\(2001\).pdf](http://tonto.eia.doe.gov/FTP/ROOT/modeldoc/m063(2001).pdf)>.
- EIA (1996) "Emissions of Greenhouse Gases in the United States" Carbon Dioxide Emissions. Energy Information Administration, U.S. Department of Energy, Washington, DC.
- EPA (2016a) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas and Petroleum Production Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2016b) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas Gathering and Boosting Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2016c) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas Transmission and Storage Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2016d) Inventory of U.S. Greenhouse Gas Emissions and Sinks 1990-2014: Revisions to Natural Gas Distribution Emissions. Available online at: <<https://www3.epa.gov/climatechange/ghgemissions/usinventoryreport/natural-gas-systems.html>>.
- EPA (2013a) *Oil and Natural Gas Sector: Standards of Performance for Crude Oil and Natural Gas Production, Transmission, and Distribution. Background Supplemental Technical Support Document for the Final New Source Performance Standards*. Environmental Protection Agency. September 2013.
- EPA (2013b) *Oil and Natural Gas Sector: New Source Performance Standards and National Emission Standards for Hazardous Air Pollutants Reviews*. Environmental Protection Agency, 40 CFR Parts 60 and 63, [EPA-HQ-OAR-2010-0505; FRL-9665-1], RIN 2060-AP76.
- EPA (2013c) Natural GasSTAR Reductions 1990-2012. Natural GasSTAR Program. September 2013.
- EPA (2013d) Updating GHG Inventory Estimate for Hydraulically Fractured Gas Well Completions and Workovers. Available online at: <http://www.epa.gov/climatechange/Downloads/ghgemissions/memo-update-emissions-for-hydraulically-workovers.pdf>.
- EPA (2015) *Greenhouse Gas Reporting Program- Subpart W – Petroleum and Natural Gas Systems*. Environmental Protection Agency. Data reported as of August 16, 2015.
- EPA (1999) *Estimates of Methane Emissions from the U.S. Oil Industry (Draft Report)*. Prepared by ICF-Kaiser, Office of Air and Radiation, U.S. Environmental Protection Agency. October 1999.
- EPA/GRI (1996) *Methane Emissions from the Natural Gas Industry*. Prepared by Harrison, M., T. Shires, J. Wessels, and R. Cowgill, eds., Radian International LLC for National Risk Management Research Laboratory, Air Pollution Prevention and Control Division, Research Triangle Park, NC. EPA-600/R-96-080a.
- FERC (2015) *North American LNG Terminals*. Federal Energy Regulatory Commission, Washington, DC.
- GTI (2001) Gas Resource Database: Unconventional Natural Gas and Gas Composition Databases. Second Edition. GRI-01/0136.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe, eds.; Institute for Global Environmental Strategies (IGES). Hayama, Kanagawa, Japan.
- OGJ (1997-2013) "Worldwide Gas Processing." *Oil & Gas Journal*, PennWell Corporation, Tulsa, OK. Available online at: <<http://www.ogj.com/>>.
- PHMSA (2015a) Transmission Annuals Data. Pipeline and Hazardous Materials Safety Administration, U.S. Department of Transportation, Washington, DC. Available online at: <<http://phmsa.dot.gov/pipeline/library/data-stats>>.
- PHMSA (2015b) Gas Distribution Annual Data. Pipeline and Hazardous Materials Safety Administration, U.S. Department of Transportation, Washington, DC. Available online at: <<http://phmsa.dot.gov/pipeline/library/data-stats>>.

Wyoming (2015) Wyoming Oil and Gas Conservation Commission. Available online at:  
<<http://wogcc.state.wy.us/coalbedchart.cfm>>.

### 3.7. Methodology for Estimating CO<sub>2</sub>, N<sub>2</sub>O, and CH<sub>4</sub> Emissions from the Incineration of Waste

Emissions of CO<sub>2</sub> from the incineration of waste include CO<sub>2</sub> generated by the incineration of plastics, synthetic rubber and synthetic fibers in municipal solid waste (MSW), and incineration of tires (which are composed in part of synthetic rubber and C black) in a variety of other combustion facilities (e.g., cement kilns). Incineration of waste also results in emissions of N<sub>2</sub>O and CH<sub>4</sub>. The methodology for calculating emissions from each of these waste incineration sources is described in this Annex.

#### CO<sub>2</sub> from Plastics Incineration

In the *Municipal Solid Waste Generation, Recycling, and Disposal in the United States: Facts and Figures* reports (EPA 1999 through 2003, 2005 through 2014) and *Advancing Sustainable Materials Management: Facts and Figures 2013 – Assessing Trends in Material Generation, Recycling and Disposal in the United States* (EPA 2015), the flows of plastics in the U.S. waste stream are reported for seven resin categories. For 2014, the quantity generated, recovered, and discarded for each resin is shown in Table A-155. The data set for 1990 through 2014 is incomplete, and several assumptions were employed to bridge the data gaps. The EPA reports do not provide estimates for individual materials landfilled and incinerated, although they do provide such an estimate for the waste stream as a whole. To estimate the quantity of plastics landfilled and incinerated, total discards were apportioned based on the proportions of landfilling and incineration for the entire U.S. waste stream for each year in the time series according to *Biocycle’s State of Garbage in America* (van Haaren et al. 2010), Themelis and Shin (in press) and Shin (2014). For those years when distribution by resin category was not reported (1990 through 1994), total values were apportioned according to 1995 (the closest year) distribution ratios. Generation and recovery figures for 2002 and 2004 were linearly interpolated between surrounding years’ data.

**Table A-155: 2014 Plastics in the Municipal Solid Waste Stream by Resin (kt)**

| Waste Pathway           | PET   | HDPE  | PVC  | LDPE/ |       | PP    | PS    | Other         | Total |
|-------------------------|-------|-------|------|-------|-------|-------|-------|---------------|-------|
|                         |       |       |      | LLDPE |       |       |       |               |       |
| Generation              | 4,246 | 5,062 | 816  | 6,768 | 6,713 | 2,059 | 3,837 | <b>29,502</b> |       |
| Recovery                | 844   | 517   | 0    | 426   | 36    | 27    | 871   | <b>2,722</b>  |       |
| Discard                 | 3,402 | 455   | 816  | 6,341 | 6,677 | 2,032 | 2,921 | <b>26,781</b> |       |
| Landfill                | 3,143 | 420   | 754  | 5,859 | 6,170 | 1,878 | 2,699 | <b>20,924</b> |       |
| Combustion              | 259   | 35    | 62   | 482   | 507   | 154   | 222   | <b>1,721</b>  |       |
| Recovery <sup>a</sup>   | 20%   | 10%   | 0%   | 6%    | 1%    | 1%    | 23%   | <b>9%</b>     |       |
| Discard <sup>a</sup>    | 80%   | 9%    | 100% | 94%   | 99%   | 99%   | 76%   | <b>91%</b>    |       |
| Landfill <sup>a</sup>   | 74%   | 8%    | 92%  | 87%   | 92%   | 91%   | 70%   | <b>71%</b>    |       |
| Combustion <sup>a</sup> | 6%    | 1%    | 8%   | 7%    | 8%    | 7%    | 6%    | <b>6%</b>     |       |

<sup>a</sup> As a percent of waste generation.

Note: Totals may not sum due to independent rounding. Abbreviations: PET (polyethylene terephthalate), HDPE (high density polyethylene), PVC (polyvinyl chloride), LDPE/LLDPE (linear low density polyethylene), PP (polypropylene), PS (polystyrene).

Fossil fuel-based CO<sub>2</sub> emissions were calculated as the product of plastic combusted, C content, and fraction oxidized (see Table A-156). The C content of each of the six types of plastics is listed, with the value for “other plastics” assumed equal to the weighted average of the six categories. The fraction oxidized was assumed to be 98 percent.

**Table A-156: 2014 Plastics Incinerated (kt), Carbon Content (%), Fraction Oxidized (%) and Carbon Incinerated (kt)**

| Factor                                    | PET        | HDPE       | PVC        | LDPE/      |            | PP         | PS         | Other        | Total |
|---|------------|------------|------------|------------|------------|------------|------------|--------------|-------|
|   |            |            |            | LLDPE      |            |            |            |              |       |
| Quantity Combusted                        | 259        | 35         | 62         | 482        | 507        | 154        | 222        | <b>1,721</b> |       |
| Carbon Content of Resin                   | 63%        | 86%        | 38%        | 86%        | 86%        | 92%        | 66%        | -            |       |
| Fraction Oxidized                         | 98%        | 98%        | 98%        | 98%        | 98%        | 98%        | 98%        | -            |       |
| Carbon in Resin Combusted                 | 158        | 29         | 23         | 405        | 426        | 140        | 143        | <b>1,325</b> |       |
| <b>Emissions (MMT CO<sub>2</sub> Eq.)</b> | <b>0.6</b> | <b>1.1</b> | <b>0.1</b> | <b>1.5</b> | <b>1.6</b> | <b>0.5</b> | <b>0.5</b> | <b>4.9</b>   |       |

<sup>a</sup> Weighted average of other plastics produced.

Note: Totals may not sum due to independent rounding.

#### CO<sub>2</sub> from Incineration of Synthetic Rubber and Carbon Black in Tires

Emissions from tire incineration require two pieces of information: the amount of tires incinerated and the C content of the tires. “2013 U.S. Scrap Tire Management Summary” (RMA 2014a) reports that 2,120 thousand of the 3,667 thousand tons of scrap tires generated in 2013 (approximately 58 percent of generation) were used for fuel purposes. Using

RMA's estimates of average tire composition and weight, the mass of synthetic rubber and C black in scrap tires was determined:

- Synthetic rubber in tires was estimated to be 90 percent C by weight, based on the weighted average C contents of the major elastomers used in new tire consumption.<sup>66</sup> Table A-157 shows consumption and C content of elastomers used for tires and other products in 2002, the most recent year for which data are available.
- C black is 100 percent C (Aslett Rubber Inc. n.d.).

Multiplying the mass of scrap tires incinerated by the total C content of the synthetic rubber, C black portions of scrap tires, and then by a 98 percent oxidation factor, yielded CO<sub>2</sub> emissions, as shown in Table A-158. The disposal rate of rubber in tires (0.3 MMT C/yr) is smaller than the consumption rate for tires based on summing the elastomers listed in Table A-155 (1.3 MMT/yr); this is due to the fact that much of the rubber is lost through tire wear during the product's lifetime and may also reflect the lag time between consumption and disposal of tires. Tire production and fuel use for 1990 through 2014 were taken from RMA 2006, RMA 2009, RMA 2011; RMA 2014a; where data were not reported, they were linearly interpolated between bracketing years' data or, for the ends of time series, set equal to the closest year with reported data.

In 2009, RMA changed the reporting of scrap tire data from millions of tires to thousands of short tons of scrap tire. As a result, the average weight and percent of the market of light duty and commercial scrap tires was used to convert the previous years from millions of tires to thousands of short tons (STMC 1990 through 1997; RMA 2002 through 2006, 2014b).

**Table A-157: Elastomers Consumed in 2002 (kt)**

| Elastomer                             | Consumed     | Carbon Content | Carbon Equivalent |
|---------------------------------------|--------------|----------------|-------------------|
| <b>Styrene butadiene rubber solid</b> | <b>768</b>   | <b>91%</b>     | <b>700</b>        |
| For Tires                             | 660          | 91%            | 602               |
| For Other Products*                   | 108          | 91%            | 98                |
| <b>Polybutadiene</b>                  | <b>583</b>   | <b>89%</b>     | <b>518</b>        |
| For Tires                             | 408          | 89%            | 363               |
| For Other Products                    | 175          | 89%            | 155               |
| <b>Ethylene Propylene</b>             | <b>301</b>   | <b>86%</b>     | <b>258</b>        |
| For Tires                             | 6            | 86%            | 5                 |
| For Other Products                    | 295          | 86%            | 253               |
| <b>Polychloroprene</b>                | <b>54</b>    | <b>59%</b>     | <b>32</b>         |
| For Tires                             | 0            | 59%            | 0                 |
| For Other Products                    | 54           | 59%            | 32                |
| <b>Nitrile butadiene rubber solid</b> | <b>84</b>    | <b>77%</b>     | <b>65</b>         |
| For Tires                             | 1            | 77%            | 1                 |
| For Other Products                    | 83           | 77%            | 64                |
| <b>Polyisoprene</b>                   | <b>58</b>    | <b>88%</b>     | <b>51</b>         |
| For Tires                             | 48           | 88%            | 42                |
| For Other Products                    | 10           | 88%            | 9                 |
| <b>Others</b>                         | <b>367</b>   | <b>88%</b>     | <b>323</b>        |
| For Tires                             | 184          | 88%            | 161               |
| For Other Products                    | 184          | 88%            | 161               |
| <b>Total</b>                          | <b>2,215</b> | <b>-</b>       | <b>1,950</b>      |
| <b>For Tires</b>                      | <b>1,307</b> | <b>-</b>       | <b>1,174</b>      |

\* Used to calculate C content of non-tire rubber products in municipal solid waste.

- Not applicable

Note: Totals may not sum due to independent rounding.

<sup>66</sup>The carbon content of tires (1,174 kt C) divided by the mass of rubber in tires (1,307 kt) equals 90 percent.

**Table A-158: Scrap Tire Constituents and CO<sub>2</sub> Emissions from Scrap Tire Incineration in 2014**

| Material         | Weight of Material (MMT) | Fraction Oxidized | Carbon Content | Emissions (MMT CO <sub>2</sub> Eq.) |
|------------------|--------------------------|-------------------|----------------|-------------------------------------|
| Synthetic Rubber | 0.3                      | 98%               | 90%            | 1.2                                 |
| Carbon Black     | 0.4                      | 98%               | 100%           | 1.4                                 |
| <b>Total</b>     | <b>0.7</b>               | <b>-</b>          | <b>-</b>       | <b>2.6</b>                          |

- Not applicable

### CO<sub>2</sub> from Incineration of Synthetic Rubber in Municipal Solid Waste

Similar to the methodology for scrap tires, CO<sub>2</sub> emissions from synthetic rubber in MSW were estimated by multiplying the amount of rubber incinerated by an average rubber C content. The amount of rubber discarded in the MSW stream was estimated from generation and recycling data<sup>67</sup> provided in the *Municipal Solid Waste Generation, Recycling, and Disposal in the United States: Facts and Figures* reports (EPA 1999 through 2003, 2005 through 2014), *Advancing Sustainable Materials Management: Facts and Figures 2013: Assessing Trends in Material Generation, Recycling and Disposal in the United States* (EPA 2015) and unpublished backup data (Schneider 2007). The reports divide rubber found in MSW into three product categories: other durables (not including tires), non-durables (which includes clothing and footwear and other non-durables), and containers and packaging. EPA (2015) did not report rubber found in the product category “containers and packaging;” however, containers and packaging from miscellaneous material types were reported for 2009 through 2014. As a result, EPA assumes that rubber containers and packaging are reported under the “miscellaneous” category; and therefore, the quantity reported for 2009 through 2014 were set equal to the quantity reported for 2008. Since there was negligible recovery for these product types, all the waste generated is considered to be discarded. Similar to the plastics method, discards were apportioned into landfilling and incineration based on their relative proportions, for each year, for the entire U.S. waste stream. The report aggregates rubber and leather in the MSW stream; an assumed synthetic rubber content of 70 percent was assigned to each product type, as shown in Table A-159.<sup>68</sup> A C content of 85 percent was assigned to synthetic rubber for all product types (based on the weighted average C content of rubber consumed for non-tire uses), and a 98 percent fraction oxidized was assumed.

**Table A-159: Rubber and Leather in Municipal Solid Waste in 2014**

| Product Type                    | Incinerated (kt) | Synthetic Rubber (%) | Carbon Content (%) | Fraction Oxidized (%) | Emissions (MMT CO <sub>2</sub> Eq.) |
|---------------------------------|------------------|----------------------|--------------------|-----------------------|-------------------------------------|
| <b>Durables (not Tires)</b>     | <b>261</b>       | <b>70%</b>           | <b>85%</b>         | <b>98%</b>            | <b>0.8</b>                          |
| <b>Non-Durables</b>             | <b>77</b>        | <b>-</b>             | <b>-</b>           | <b>-</b>              | <b>0.2</b>                          |
| Clothing and Footwear           | 59               | 70%                  | 85%                | 98%                   | 0.2                                 |
| Other Non-Durables              | 18               | 70%                  | 85%                | 98%                   | 0.1                                 |
| <b>Containers and Packaging</b> | <b>2</b>         | <b>70%</b>           | <b>85%</b>         | <b>98%</b>            | <b>0.0</b>                          |
| <b>Total</b>                    | <b>341</b>       | <b>-</b>             | <b>-</b>           | <b>-</b>              | <b>1.1</b>                          |

- Not Applicable.

### CO<sub>2</sub> from Incineration of Synthetic Fibers

Carbon dioxide emissions from synthetic fibers were estimated as the product of the amount of synthetic fiber discarded annually and the average C content of synthetic fiber. Fiber in the MSW stream was estimated from data provided in the *Municipal Solid Waste Generation, Recycling, and Disposal in the United States: Facts and Figures* reports (EPA 1999 through 2003, 2005 through 2015) for textiles. Production data for the synthetic fibers was based on data from the American Chemical Society (FEB 2009). The amount of synthetic fiber in MSW was estimated by subtracting (a) the amount recovered from (b) the waste generated (see Table A-160). As with the other materials in the MSW stream, discards were apportioned based on the annually variable proportions of landfilling and incineration for the entire U.S. waste stream, as found in van Haaren et al. (2010), Themelis and Shin (in press), and Shin (2014). It was assumed that approximately 55 percent of the fiber was synthetic in origin, based on information received from the Fiber Economics Bureau (DeZan 2000). An average C content of 70 percent was assigned to synthetic fiber using the production-weighted average of the C contents of the four major fiber types (polyester, nylon, olefin, and acrylic) produced in 1999 (see Table A-161). The equation relating CO<sub>2</sub> emissions to the amount of textiles combusted is shown below.

$$\text{CO}_2 \text{ Emissions from the Incineration of Synthetic Fibers} = \text{Annual Textile Incineration (kt)} \times (\text{Percent of Total Fiber that is Synthetic}) \times (\text{Average C Content of Synthetic Fiber}) \times (44 \text{ g CO}_2 / 12 \text{ g C})$$

<sup>67</sup> Discards = Generation minus recycling.

<sup>68</sup> As a sustainably harvested biogenic material, the incineration of leather is assumed to have no net CO<sub>2</sub> emissions.

**Table A-160: Synthetic Textiles in MSW (kt)**

| Year | Generation | Recovery | Discards | Incineration |
|------|------------|----------|----------|--------------|
| 1990 | 2,884      | 328      | 2,557    | 332          |
| 1995 | 3,674      | 447      | 3,227    | 442          |
| 1996 | 3,832      | 472      | 3,361    | 467          |
| 1997 | 4,090      | 526      | 3,564    | 458          |
| 1998 | 4,269      | 556      | 3,713    | 407          |
| 1999 | 4,498      | 611      | 3,887    | 406          |
| 2000 | 4,706      | 655      | 4,051    | 417          |
| 2001 | 4,870      | 715      | 4,155    | 432          |
| 2002 | 5,123      | 750      | 4,373    | 459          |
| 2003 | 5,297      | 774      | 4,522    | 472          |
| 2004 | 5,451      | 884      | 4,567    | 473          |
| 2005 | 5,714      | 913      | 4,800    | 480          |
| 2006 | 5,893      | 933      | 4,959    | 479          |
| 2007 | 6,041      | 953      | 5,088    | 470          |
| 2008 | 6,305      | 968      | 5,337    | 470          |
| 2009 | 6,424      | 978      | 5,446    | 458          |
| 2010 | 6,508      | 998      | 5,510    | 441          |
| 2011 | 6,513      | 1,003    | 5,510    | 419          |
| 2012 | 7,114      | 1,117    | 5,997    | 456          |
| 2013 | 7,496      | 894      | 6,602    | 502          |
| 2014 | 7,496      | 894      | 6,602    | 502          |

**Table A-161: Synthetic Fiber Production in 1999**

| Fiber        | Production (MMT) | Carbon Content |
|--------------|------------------|----------------|
| Polyester    | 1.8              | 63%            |
| Nylon        | 1.2              | 64%            |
| Olefin       | 1.4              | 86%            |
| Acrylic      | 0.1              | 68%            |
| <b>Total</b> | <b>4.5</b>       | <b>70%</b>     |

## N<sub>2</sub>O and CH<sub>4</sub> from Incineration of Waste

Estimates of N<sub>2</sub>O emissions from the incineration of waste in the United States are based on the methodology outlined in the EPA's Compilation of Air Pollutant Emission Factors (EPA 1995) and presented in the *Municipal Solid Waste Generation, Recycling, and Disposal in the United States: Facts and Figures* reports (EPA 1999 through 2003, 2005 through 2014), *Advancing Sustainable Materials Management: Facts and Figures 2013: Assessing Trends in Material Generation, Recycling and Disposal in the United States* (EPA 2015), and unpublished backup data (Schneider 2007). According to this methodology, emissions of N<sub>2</sub>O from waste incineration are the product of the mass of waste incinerated, an emission factor of N<sub>2</sub>O emitted per unit mass of waste incinerated, and an N<sub>2</sub>O emissions control removal efficiency. The mass of waste incinerated was derived from the results of the biannual national survey of Municipal Solid Waste (MSW) Generation and Disposition in the U.S., published in *BioCycle* (van Haaren et al. 2010), Themelis and Shin (in press), and Shin (2014). For waste incineration in the United States, an emission factor of 50 g N<sub>2</sub>O/metric ton MSW based on the 2006 IPCC Guidelines and an estimated emissions control removal efficiency of zero percent were used (IPCC 2006). It was assumed that all MSW incinerators in the United States use continuously-fed stoker technology (Bahor 2009, ERC 2009).

Estimates of CH<sub>4</sub> emissions from the incineration of waste in the United States are based on the methodology outlined in IPCC's 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006). According to this methodology, emissions of CH<sub>4</sub> from waste incineration are the product of the mass of waste incinerated and an emission factor of CH<sub>4</sub> emitted per unit mass of waste incinerated. Similar to the N<sub>2</sub>O emissions methodology, the mass of waste incinerated was derived from the information published in *BioCycle* (van Haaren et al. 2010) for 1990 through 2008. Data for 2011 were derived from information forthcoming in Themelis and Shin (in press) and Shin (2014). For waste incineration in the United States, an emission factor of 0.20 kg CH<sub>4</sub>/kt MSW was used based on the 2006 IPCC Guidelines and assuming that all MSW incinerators in the United States use continuously-fed stoker technology (Bahor 2009; ERC 2009). No information was available on the mass of waste incinerated for 2012, 2013, or 2014, so these values were assumed to be equal to the 2011 value.

Despite the differences in methodology and data sources, the two series of references (EPA 2014; van Haaren, Rob, Themelis, N., and Goldstein, N. 2010) provide estimates of total solid waste incinerated that are relatively consistent (see Table A-162).

**Table A-162: U.S. Municipal Solid Waste Incinerated, as Reported by EPA and BioCycle (Metric Tons)**

| Year | EPA                     | BioCycle                |
|------|-------------------------|-------------------------|
| 1990 | 28,939,680              | 30,632,057              |
| 1995 | 32,241,888              | 29,639,040              |
| 2000 | 30,599,856              | 25,974,978              |
| 2001 | 30,481,920              | 25,942,036 <sup>a</sup> |
| 2002 | 30,255,120              | 25,802,917              |
| 2003 | 30,028,320              | 25,930,542 <sup>b</sup> |
| 2004 | 28,585,872              | 26,037,823              |
| 2005 | 28,685,664              | 25,973,520 <sup>c</sup> |
| 2006 | 28,985,040              | 25,853,401              |
| 2007 | 29,003,184              | 24,788,539 <sup>d</sup> |
| 2008 | 28,622,160              | 23,674,017              |
| 2009 | 26,317,872              | 22,714,122 <sup>e</sup> |
| 2010 | 26,544,672              | 21,741,734 <sup>e</sup> |
| 2011 | 26,544,672              | 20,756,870              |
| 2012 | 26,544,672              | 20,756,870 <sup>f</sup> |
| 2013 | 26,544,672 <sup>g</sup> | 20,756,870 <sup>f</sup> |
| 2014 | 26,544,672 <sup>g</sup> | 20,756,870 <sup>f</sup> |

<sup>a</sup> Interpolated between 2000 and 2002 values.

<sup>b</sup> Interpolated between 2002 and 2004 values.

<sup>c</sup> Interpolated between 2004 and 2006 values.

<sup>d</sup> Interpolated between 2006 and 2008 values

<sup>e</sup> Interpolated between 2011 and 2008 values

<sup>f</sup> Set equal to the 2011 value

<sup>g</sup> Set equal to the 2012 value.

## References

- ArSova, Ljupka, Rob van Haaren, Nora Goldstein, Scott M. Kaufman, and Nickolas J. Themelis (2008) "16th Annual BioCycle Nationwide Survey: The State of Garbage in America" Biocycle, JG Press, Emmaus, PA. December.
- Bahor, B (2009) Covanta Energy's public review comments re: *Draft Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2007*. Submitted via email on April 9, 2009 to Leif Hockstad, U.S. EPA.
- De Soete, G.G. (1993) "Nitrous Oxide from Combustion and Industry: Chemistry, Emissions and Control." In A. R. Van Amstel, (ed) Proc. of the International Workshop Methane and Nitrous Oxide: Methods in National Emission Inventories and Options for Control, Amersfoort, NL. February 3-5, 1993.
- DeZan, D. (2000) Personal Communication between Diane DeZan, Fiber Economics Bureau and Joe Casola, ICF Consulting. 4 August 2000.
- Energy Recovery Council (2009) "2007 Directory of Waste-to-Energy Plants in the United States." Accessed September 29, 2009.
- EPA (2015) Advancing Sustainable Materials Management: Facts and Figures 2013 – Assessing Trends in Material Generation, Recycling and Disposal in the United States. Office of Solid Waste and Emergency Response, U.S. Environmental Protection Agency. Washington, D.C. Available online at [http://www3.epa.gov/epawaste/nonhaz/municipal/pubs/2013\\_advncng\\_smm\\_rpt.pdf](http://www3.epa.gov/epawaste/nonhaz/municipal/pubs/2013_advncng_smm_rpt.pdf).
- EPA (2007, 2008, 2011, 2013, 2014) Municipal Solid Waste in the United States: Facts and Figures. Office of Solid Waste and Emergency Response, U.S. Environmental Protection Agency. Washington, DC. Available online at <http://www.epa.gov/osw/nonhaz/municipal/msw99.htm>.
- EPA (2006) Solid Waste Management and Greenhouse Gases: A Life-Cycle Assessment of Emissions and Sinks. Office of Solid Waste and Emergency Response, U.S. Environmental Protection Agency. Washington, DC.
- EPA (2000) Characterization of Municipal Solid Waste in the United States: Source Data on the 1999 Update. Office of Solid Waste, U.S. Environmental Protection Agency. Washington, DC. EPA530-F-00-024.
- Goldstein, N. and C. Madtes (2001) "13th Annual BioCycle Nationwide Survey: The State of Garbage in America." Biocycles, JG Press, Emmaus, PA. December 2001.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- Kaufman, et al. (2004) "14th Annual BioCycle Nationwide Survey: The State of Garbage in America 2004" Biocycle, JG Press, Emmaus, PA. January, 2004.
- RMA (2014a) "2013 U.S. Scrap Tire Management Summary." Rubber Manufacturers Association. November 2014. Available online at: [http://www.rma.org/download/scrap-tires/market-reports/US\\_STMarket2013.pdf](http://www.rma.org/download/scrap-tires/market-reports/US_STMarket2013.pdf). Accessed 17 November 2014.
- RMA (2014b) "Scrap Tire Markets: Facts and Figures – Scrap Tire Characteristics." Available online at [http://www.stref.orgwww.stref.org/scrap\\_tires/scrap\\_tire\\_markets/scrap\\_tire\\_characteristics/](http://www.stref.orgwww.stref.org/scrap_tires/scrap_tire_markets/scrap_tire_characteristics/). Accessed 17 November 2014.
- RMA (2012) "Rubber FAQs." Rubber Manufacturers Association. Available online at <http://www.rma.org/about-rma/rubber-faqs/> Accessed 19 November 2014.
- RMA (2011) "U.S. Scrap Tire Management Summary 2005-2009." Rubber Manufacturers Association. October 2011. Available online at: [http://www.rma.org/scrap\\_tires/scrap\\_tire\\_markets/2009\\_summary.pdf](http://www.rma.org/scrap_tires/scrap_tire_markets/2009_summary.pdf).
- Schneider, S. (2007) E-mail between Shelly Schneider of Franklin Associates (a division of ERG) and Sarah Shapiro of ICF International, January 10, 2007.
- Shin, D. (2014) Generation and Disposition of Municipal Solid Waste (MSW) in the United States—A National Survey. Thesis. Columbia University, Department of Earth and Environmental Engineering, January 3, 2014.
- Simmons, et al. (2006) "15th Nationwide Survey of Municipal Solid Waste Management in the United States: The State of Garbage in America" Biocycle, JG Press, Emmaus, PA. April 2006.

Themelis and Shin (in press) U.S. Survey of Generation and Disposition of Municipal Solid Waste. Waste Management. Submitted November 14, 2014.

van Haaren, Rob, Themelis, N., and Goldstein, N. (2010) "The State of Garbage in America." BioCycle, October 2010. Volume 51, Number 10, pg. 16-23.

### 3.8. Methodology for Estimating Emissions from International Bunker Fuels used by the U.S. Military

Bunker fuel emissions estimates for the Department of Defense (DoD) were developed using data generated by the Defense Logistics Agency Energy (DLA Energy) for aviation and naval fuels. DLA Energy prepared a special report based on data in the Fuels Automated System (FAS) for calendar year 2014 fuel sales in the Continental United States (CONUS).<sup>69</sup> The following steps outline the methodology used for estimating emissions from international bunker fuels used by the U.S. Military.

#### Step 1: Omit Extra-Territorial Fuel Deliveries

Beginning with the complete FAS data set for each year, the first step in quantifying DoD-related emissions from international bunker fuels was to identify data that would be representative of international bunker fuel consumption as defined by decisions of the UNFCCC (i.e., fuel sold to a vessel, aircraft, or installation within the United States or its territories and used in international maritime or aviation transport). Therefore, fuel data were categorized by the location of fuel delivery in order to identify and omit all international fuel transactions/deliveries (i.e., sales abroad).

#### Step 2: Allocate JP-8 between Aviation and Land-based Vehicles

As a result of DoD<sup>70</sup> and NATO<sup>71</sup> policies on implementing the Single Fuel For the Battlefield concept, DoD activities have been increasingly replacing diesel fuel with JP8 (a type of jet fuel) in compression ignition and turbine engines of land-based equipment. Based on this concept and examination of all data describing jet fuel used in land-based vehicles, it was determined that a portion of JP8 consumption should be attributed to ground vehicle use. Based on available Military Service data and expert judgment, a small fraction of the total JP8 use (i.e., between 1.78 and 2.7 times the quantity of diesel fuel used, depending on the Service) was reallocated from the aviation subtotal to a new land-based jet fuel category for 1997 and subsequent years. As a result of this reallocation, the JP8 use reported for aviation was reduced and the total fuel use for land-based equipment increased. DoD's total fuel use did not change.

Table A-163 displays DoD's consumption of transportation fuels, summarized by fuel type, that remain at the completion of Step 1, and reflects the adjustments for jet fuel used in land-based equipment, as described above.

#### Step 3: Omit Land-Based Fuels

Navy and Air Force land-based fuels (i.e., fuel not used by ships or aircraft) were omitted for the purpose of calculating international bunker fuels. The remaining fuels, listed below, were considered potential DoD international bunker fuels.

- **Aviation:** jet fuels (JP8, JP5, JP4, JAA, JA1, and JAB).
- **Marine:** naval distillate fuel (F76), marine gas oil (MGO), and intermediate fuel oil (IFO).

#### Step 4: Omit Fuel Transactions Received by Military Services that are not considered to be International Bunker Fuels

Only Navy and Air Force were deemed to be users of military international bunker fuels after sorting the data by Military Service and applying the following assumptions regarding fuel use by Service.

---

<sup>69</sup> FAS contains data for 1995 through 2014, but the dataset was not complete for years prior to 1995. Using DLA aviation and marine fuel procurement data, fuel quantities from 1990 to 1994 were estimated based on a back-calculation of the 1995 data in the legacy database, the Defense Fuels Automated Management System (DFAMS). The back-calculation was refined in 1999 to better account for the jet fuel conversion from JP4 to JP8 that occurred within DoD between 1992 and 1995.

<sup>70</sup> DoD Directive 4140.25-M-V1, Fuel Standardization and Cataloging, 2013; DoD Directive 4140.25, DoD Management Policy for Energy Commodities and Related Services, 2004.

<sup>71</sup> NATO Standard Agreement NATO STANAG 4362, Fuels for Future Ground Equipments Using Compression Ignition or Turbine Engines, 2012.

- Only fuel delivered to a ship, aircraft, or installation in the United States was considered a potential international bunker fuel. Fuel consumed in international aviation or marine transport was included in the bunker fuel estimate of the country where the ship or aircraft was fueled. Fuel consumed entirely within a country's borders was not considered a bunker fuel.
- Based on previous discussions with the Army staff, only an extremely small percentage of Army aviation emissions, and none of Army watercraft emissions, qualified as bunker fuel emissions. The magnitude of these emissions was judged to be insignificant when compared to Air Force and Navy emissions. Based on this research, Army bunker fuel emissions were assumed to be zero.
- Marine Corps aircraft operating while embarked consumed fuel that was reported as delivered to the Navy. Bunker fuel emissions from embarked Marine Corps aircraft were reported in the Navy bunker fuel estimates. Bunker fuel emissions from other Marine Corps operations and training were assumed to be zero.
- Bunker fuel emissions from other DoD and non-DoD activities (i.e., other federal agencies) that purchased fuel from DLA Energy were assumed to be zero.

### Step 5: Determine Bunker Fuel Percentages

It was necessary to determine what percent of the aviation and marine fuels were used as international bunker fuels. Military aviation bunkers include international operations (i.e., sorties that originate in the United States and end in a foreign country), operations conducted from naval vessels at sea, and operations conducted from U.S. installations principally over international water in direct support of military operations at sea (e.g., anti-submarine warfare flights). Methods for quantifying aviation and marine bunker fuel percentages are described below.

- **Aviation:** The Air Force Aviation bunker fuel percentage was determined to be 13.2 percent. A bunker fuel weighted average was calculated based on flying hours by major command. International flights were weighted by an adjustment factor to reflect the fact that they typically last longer than domestic flights. In addition, a fuel use correction factor was used to account for the fact that transport aircraft burn more fuel per hour of flight than most tactical aircraft. This percentage was multiplied by total annual Air Force aviation fuel delivered for U.S. activities, producing an estimate for international bunker fuel consumed by the Air Force.

The Naval Aviation bunker fuel percentage was calculated to be 40.4 percent by using flying hour data from Chief of Naval Operations Flying Hour Projection System Budget for fiscal year 1998 and estimates of bunker fuel percent of flights provided by the fleet. This Naval Aviation bunker fuel percentage was then multiplied by total annual Navy aviation fuel delivered for U.S. activities, yielding total Navy aviation bunker fuel consumed.

- **Marine:** For marine bunkers, fuels consumed while ships were underway were assumed to be bunker fuels. The Navy maritime bunker fuel percentage was determined to be 79 percent because the Navy reported that 79 percent of vessel operations were underway, while the remaining 21 percent of operations occurred in port (i.e., pierside) in the year 2000.<sup>72</sup>

Table A-164 and Table A-165 display DoD bunker fuel use totals for the Navy and Air Force.

### Step 6: Calculate Emissions from International Bunker Fuels

Bunker fuel totals were multiplied by appropriate emission factors to determine greenhouse gas emissions. CO<sub>2</sub> emissions from Aviation Bunkers and distillate Marine Bunkers are the total of military aviation and marine bunker fuels, respectively.

---

<sup>72</sup> Note that 79 percent is used because it is based on Navy data, but the percentage of time underway may vary from year-to-year depending on vessel operations. For example, for years prior to 2000, the bunker fuel percentage was 87 percent.

The rows labeled “U.S. Military” and “U.S. Military Naval Fuels” in the tables in the International Bunker Fuels section of the Energy chapter were based on the totals provided in Table A-164 and Table A-165, below. CO<sub>2</sub> emissions from aviation bunkers and distillate marine bunkers are presented in Table A-168, and are based on emissions from fuels tallied in Table A-164 and Table A-165.

**Table A-163: Transportation Fuels from Domestic Fuel Deliveries<sup>a</sup> (Million Gallons)**

| Vehicle Type/Fuel                        | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Aviation</b>                          | <b>4,598.4</b> | <b>3,099.9</b> | <b>2,664.4</b> | <b>2,900.6</b> | <b>2,609.8</b> | <b>2,615.0</b> | <b>2,703.1</b> | <b>2,338.1</b> | <b>2,092.0</b> | <b>2,081.0</b> | <b>2,067.8</b> | <b>1,814.5</b> | <b>1,663.9</b> | <b>1,405.0</b> | <b>1,449.7</b> | <b>1,336.4</b> | <b>1,796.2</b> |
| Total Jet Fuels                          | 4,598.4        | 3,099.9        | 2,664.4        | 2,900.6        | 2,609.6        | 2,614.9        | 2,703.1        | 2,338.0        | 2,091.9        | 2,080.9        | 2,067.7        | 1,814.3        | 1,663.7        | 1,404.8        | 1,449.5        | 1,336.2        | 1,795.9        |
| JP8                                      | 285.7          | 2,182.8        | 2,122.7        | 2,326.2        | 2,091.4        | 2,094.3        | 2,126.2        | 1,838.8        | 1,709.3        | 1,618.5        | 1,616.2        | 1,358.2        | 1,100.1        | 882.8          | 865.2          | 718.0          | 546.6          |
| JP5                                      | 1,025.4        | 691.2          | 472.1          | 503.2          | 442.2          | 409.1          | 433.7          | 421.6          | 325.5          | 376.1          | 362.2          | 361.2          | 399.3          | 372.3          | 362.5          | 316.4          | 311.0          |
| Other Jet Fuels                          | 3,287.3        | 225.9          | 69.6           | 71.2           | 76.1           | 111.4          | 143.2          | 77.6           | 57.0           | 86.3           | 89.2           | 94.8           | 164.3          | 149.7          | 221.8          | 301.7          | 938.3          |
| Aviation Gasoline                        | +              | +              | +              | +              | 0.1            | 0.1            | +              | 0.1            | 0.1            | 0.2            | 0.1            | 0.2            | 0.2            | 0.2            | 0.3            | 0.2            | 0.3            |
| <b>Marine</b>                            | <b>686.8</b>   | <b>438.9</b>   | <b>454.4</b>   | <b>418.4</b>   | <b>455.8</b>   | <b>609.1</b>   | <b>704.5</b>   | <b>604.9</b>   | <b>531.6</b>   | <b>572.8</b>   | <b>563.4</b>   | <b>485.8</b>   | <b>578.8</b>   | <b>489.9</b>   | <b>490.4</b>   | <b>390.4</b>   | <b>427.9</b>   |
| Middle Distillate (MGO)                  | +              | +              | 48.3           | 33.0           | 41.2           | 88.1           | 71.2           | 54.0           | 45.8           | 45.7           | 55.2           | 56.8           | 48.4           | 37.3           | 52.9           | 40.9           | 62.0           |
| Naval Distillate (F76)                   | 686.8          | 438.9          | 398.0          | 369.1          | 395.1          | 460.9          | 583.5          | 525.9          | 453.6          | 516.0          | 483.4          | 399.0          | 513.7          | 440.0          | 428.4          | 345.7          | 362.7          |
| Intermediate Fuel Oil (IFO) <sup>b</sup> | +              | +              | 8.1            | 16.3           | 19.5           | 60.2           | 49.9           | 25.0           | 32.2           | 11.1           | 24.9           | 30.0           | 16.7           | 12.5           | 9.1            | 3.8            | 3.2            |
| <b>Other<sup>c</sup></b>                 | <b>717.1</b>   | <b>310.9</b>   | <b>248.2</b>   | <b>109.8</b>   | <b>211.1</b>   | <b>221.2</b>   | <b>170.9</b>   | <b>205.6</b>   | <b>107.3</b>   | <b>169.0</b>   | <b>173.6</b>   | <b>206.8</b>   | <b>224.0</b>   | <b>208.6</b>   | <b>193.8</b>   | <b>180.6</b>   | <b>190.7</b>   |
| Diesel                                   | 93.0           | 119.9          | 126.6          | 26.6           | 57.7           | 60.8           | 46.4           | 56.8           | 30.6           | 47.3           | 49.1           | 58.3           | 64.1           | 60.9           | 57.9           | 54.9           | 57.5           |
| Gasoline                                 | 624.1          | 191.1          | 74.8           | 24.7           | 27.5           | 26.5           | 19.4           | 24.3           | 11.7           | 19.2           | 19.7           | 25.2           | 25.5           | 22.0           | 19.6           | 16.9           | 16.5           |
| Jet Fuel <sup>d</sup>                    | +              | +              | 46.7           | 58.4           | 125.9          | 133.9          | 105.1          | 124.4          | 65.0           | 102.6          | 104.8          | 123.3          | 134.4          | 125.6          | 116.2          | 108.8          | 116.7          |
| <b>Total (Including Bunkers)</b>         | <b>6,002.4</b> | <b>3,849.8</b> | <b>3,367.0</b> | <b>3,428.8</b> | <b>3,276.7</b> | <b>3,445.3</b> | <b>3,578.5</b> | <b>3,148.6</b> | <b>2,730.9</b> | <b>2,822.8</b> | <b>2,804.9</b> | <b>2,507.1</b> | <b>2,466.7</b> | <b>2,103.5</b> | <b>2,133.9</b> | <b>1,907.5</b> | <b>2,414.9</b> |

+ Indicates value does not exceed 0.05 million gallons.

<sup>a</sup> Includes fuel distributed in the United States and U.S. Territories.

<sup>b</sup> Intermediate fuel oil (IFO 180 and IFO 380) is a blend of distillate and residual fuels. IFO is used by the Military Sealift Command.

<sup>c</sup> Prior to 2001, gasoline and diesel fuel totals were estimated using data provided by the Military Services for 1990 and 1996. The 1991 through 1995 data points were interpolated from the Service inventory data. The 1997 through 1999 gasoline and diesel fuel data were initially extrapolated from the 1996 inventory data. Growth factors used for other diesel and gasoline were 5.2 and -21.1 percent, respectively. However, prior diesel fuel estimates from 1997 through 2000 were reduced according to the estimated consumption of jet fuel that is assumed to have replaced the diesel fuel consumption in land-based vehicles. Datasets for other diesel and gasoline consumed by the military in 2000 were estimated based on ground fuels consumption trends. This method produced a result that was more consistent with expected consumption for 2000. Since 2001, other gasoline and diesel fuel totals were generated by DLA Energy.

<sup>d</sup> The fraction of jet fuel consumed in land-based vehicles was estimated based on DLA Energy data as well as Military Service and expert judgment.

Note: Totals may not sum due to independent rounding.

**Table A-164: Total U.S. Military Aviation Bunker Fuel (Million Gallons)**

| Fuel Type/Service         | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|---------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Jet Fuels</b>          |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |
| <b>JP8</b>                | <b>56.7</b>  | <b>300.4</b> | <b>307.6</b> | <b>341.2</b> | <b>309.5</b> | <b>305.1</b> | <b>309.8</b> | <b>285.6</b> | <b>262.5</b> | <b>249.1</b> | <b>229.4</b> | <b>211.4</b> | <b>182.5</b> | <b>143.4</b> | <b>141.2</b> | <b>122.0</b> | <b>71.8</b>  |
| Navy                      | 56.7         | 38.3         | 53.4         | 73.8         | 86.6         | 76.3         | 79.2         | 70.9         | 64.7         | 62.7         | 59.2         | 55.4         | 60.8         | 47.1         | 50.4         | 48.9         | 19.8         |
| Air Force                 | +            | 262.2        | 254.2        | 267.4        | 222.9        | 228.7        | 230.6        | 214.7        | 197.8        | 186.5        | 170.3        | 156.0        | 121.7        | 96.2         | 90.8         | 73.0         | 52.0         |
| <b>JP5</b>                | <b>370.5</b> | <b>249.8</b> | <b>160.3</b> | <b>169.7</b> | <b>158.3</b> | <b>146.1</b> | <b>157.9</b> | <b>160.6</b> | <b>125.0</b> | <b>144.5</b> | <b>139.2</b> | <b>137.0</b> | <b>152.5</b> | <b>144.9</b> | <b>141.2</b> | <b>124.9</b> | <b>121.9</b> |
| Navy                      | 365.3        | 246.3        | 155.6        | 163.7        | 153.0        | 141.3        | 153.8        | 156.9        | 122.8        | 141.8        | 136.5        | 133.5        | 149.7        | 143.0        | 139.5        | 123.6        | 120.2        |
| Air Force                 | 5.3          | 3.5          | 4.7          | 6.1          | 5.3          | 4.9          | 4.1          | 3.7          | 2.3          | 2.7          | 2.6          | 3.5          | 2.8          | 1.8          | 1.7          | 1.3          | 1.6          |
| <b>JP4</b>                | <b>420.8</b> | <b>21.5</b>  | <b>0.0</b>   | <b>0.1</b>   | <b>0.0</b>   | <b>0.0</b>   | <b>0.0</b>   | <b>0.0</b>   |
| Navy                      | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            |
| Air Force                 | 420.8        | 21.5         | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | 0.1          | +            | +            | +            | +            |
| <b>JAA</b>                | <b>13.7</b>  | <b>9.2</b>   | <b>12.5</b>  | <b>12.6</b>  | <b>13.7</b>  | <b>21.7</b>  | <b>30.0</b>  | <b>15.5</b>  | <b>11.7</b>  | <b>15.6</b>  | <b>16.8</b>  | <b>18.1</b>  | <b>31.4</b>  | <b>31.1</b>  | <b>38.6</b>  | <b>46.5</b>  | <b>124.5</b> |
| Navy                      | 8.5          | 5.7          | 7.9          | 8.0          | 9.8          | 15.5         | 21.5         | 11.6         | 9.1          | 11.7         | 12.5         | 12.3         | 13.7         | 14.6         | 14.8         | 13.4         | 32.2         |
| Air Force                 | 5.3          | 3.5          | 4.5          | 4.6          | 3.8          | 6.2          | 8.6          | 3.9          | 2.6          | 3.9          | 4.3          | 5.9          | 17.7         | 16.5         | 23.8         | 33.1         | 92.3         |
| <b>JA1</b>                | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>0.1</b>   | <b>0.6</b>   | <b>0.2</b>   | <b>0.5</b>   | <b>0.5</b>   | <b>0.4</b>   | <b>1.1</b>   | <b>1.0</b>   | <b>0.6</b>   | <b>0.3</b>   | <b>(0.5)</b> | <b>(0.3)</b> | <b>0.6</b>   | <b>0.3</b>   |
| Navy                      | +            | +            | +            | +            | +            | +            | +            | +            | +            | 0.1          | 0.1          | 0.1          | 0.1          | (0.5)        | (0.3)        | 0.6          | +            |
| Air Force                 | +            | +            | +            | 0.1          | 0.6          | 0.2          | 0.5          | 0.5          | 0.4          | 1.0          | 0.8          | 0.5          | 0.1          | +            | +            | +            | 0.3          |
| <b>JAB</b>                | <b>+</b>     |
| Navy                      | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            |
| Air Force                 | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            |
| <b>Navy Subtotal</b>      | <b>430.5</b> | <b>290.2</b> | <b>216.9</b> | <b>245.5</b> | <b>249.4</b> | <b>233.1</b> | <b>254.4</b> | <b>239.4</b> | <b>196.6</b> | <b>216.3</b> | <b>208.3</b> | <b>201.3</b> | <b>224.4</b> | <b>204.3</b> | <b>204.5</b> | <b>186.5</b> | <b>172.3</b> |
| <b>Air Force Subtotal</b> | <b>431.3</b> | <b>290.7</b> | <b>263.5</b> | <b>278.1</b> | <b>232.7</b> | <b>239.9</b> | <b>243.7</b> | <b>222.9</b> | <b>203.1</b> | <b>194.0</b> | <b>178.1</b> | <b>165.9</b> | <b>142.4</b> | <b>114.5</b> | <b>116.3</b> | <b>107.4</b> | <b>146.2</b> |
| <b>Total</b>              | <b>861.8</b> | <b>580.9</b> | <b>480.4</b> | <b>523.6</b> | <b>482.1</b> | <b>473.0</b> | <b>498.1</b> | <b>462.3</b> | <b>399.7</b> | <b>410.3</b> | <b>386.3</b> | <b>367.2</b> | <b>366.7</b> | <b>318.8</b> | <b>320.8</b> | <b>293.9</b> | <b>318.5</b> |

+ Does not exceed 0.05 million gallons.

The negative values in this table represent returned products.

Note: Totals may not sum due to independent rounding.

**Table A-165: Total U.S. DoD Maritime Bunker Fuel (Million Gallons)**

| Marine Distillates | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|--------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Navy – MGO         | 0.0          | 0.0          | 23.8         | 22.5         | 27.1         | 63.7         | 56.2         | 38.0         | 33.0         | 31.6         | 40.9         | 39.9         | 32.9         | 25.5         | 36.5         | 32.3         | 43.3         |
| Navy – F76         | 522.4        | 333.8        | 298.6        | 282.6        | 305.6        | 347.8        | 434.4        | 413.1        | 355.9        | 404.1        | 376.9        | 311.4        | 402.2        | 346.6        | 337.9        | 273.1        | 286.2        |
| Navy – IFO         | 0.0          | 0.0          | 6.4          | 12.9         | 15.4         | 47.5         | 39.4         | 19.7         | 25.4         | 8.8          | 19.0         | 23.1         | 12.9         | 9.5          | 6.1          | 3.0          | 1.5          |
| <b>Total</b>       | <b>522.4</b> | <b>333.8</b> | <b>328.8</b> | <b>318.0</b> | <b>348.2</b> | <b>459.0</b> | <b>530.0</b> | <b>470.7</b> | <b>414.3</b> | <b>444.4</b> | <b>436.7</b> | <b>374.4</b> | <b>448.0</b> | <b>381.5</b> | <b>380.6</b> | <b>308.5</b> | <b>331.0</b> |

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 million gallons.

**Table A-166: Aviation and Marine Carbon Contents (MMT Carbon/QBtu) and Fraction Oxidized**

| Mode (Fuel)         | Carbon Content Coefficient | Fraction Oxidized |
|---------------------|----------------------------|-------------------|
| Aviation (Jet Fuel) | Variable                   | 1.00              |
| Marine (Distillate) | 20.17                      | 1.00              |
| Marine (Residual)   | 20.48                      | 1.00              |

Source: EPA (2010) and IPCC (2006).

**Table A-167: Annual Variable Carbon Content Coefficient for Jet Fuel (MMT Carbon/QBtu)**

| Fuel     | 1990  | 1995  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  | 2008  | 2009  | 2010  | 2011  | 2012  | 2013  | 2014  |
|----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Jet Fuel | 19.40 | 19.34 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 | 19.70 |

Source: EPA (2010)

**Table A-168: Total U.S. DoD CO<sub>2</sub> Emissions from Bunker Fuels (MMT CO<sub>2</sub> Eq.)**

| Mode         | 1990        | 1995       | 2000       | 2001       | 2002       | 2003       | 2004        | 2005       | 2006       | 2007       | 2008       | 2009       | 2010       | 2011       | 2012       | 2013       | 2014       |
|--------------|-------------|------------|------------|------------|------------|------------|-------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Aviation     | 8.1         | 5.5        | 4.7        | 5.1        | 4.7        | 4.6        | 4.8         | 4.5        | 3.9        | 4.0        | 3.8        | 3.6        | 3.6        | 3.1        | 3.1        | 2.9        | 3.1        |
| Marine       | 5.4         | 3.4        | 3.4        | 3.3        | 3.6        | 4.7        | 5.4         | 4.8        | 4.2        | 4.6        | 4.5        | 3.8        | 4.6        | 3.9        | 3.9        | 3.2        | 3.4        |
| <b>Total</b> | <b>13.4</b> | <b>9.0</b> | <b>8.0</b> | <b>8.3</b> | <b>8.3</b> | <b>9.3</b> | <b>10.3</b> | <b>9.3</b> | <b>8.1</b> | <b>8.5</b> | <b>8.2</b> | <b>7.4</b> | <b>8.2</b> | <b>7.0</b> | <b>7.0</b> | <b>6.0</b> | <b>6.5</b> |

Note: Totals may not sum due to independent rounding.

## References

DLA Energy (2015) Unpublished data from the Defense Fuels Automated Management System (DFAMS). Defense Energy Support Center, Defense Logistics Agency, U.S. Department of Defense. Washington, D.C.

IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.

### 3.9. Methodology for Estimating HFC and PFC Emissions from Substitution of Ozone Depleting Substances

Emissions of HFCs and PFCs from the substitution of ozone depleting substances (ODS) are developed using a country-specific modeling approach. The Vintaging Model was developed as a tool for estimating the annual chemical emissions from industrial sectors that have historically used ODS in their products. Under the terms of the Montreal Protocol and the United States Clean Air Act Amendments of 1990, the domestic U.S. consumption of ODS—chlorofluorocarbons (CFCs), halons, carbon tetrachloride, methyl chloroform, and hydrochlorofluorocarbons (HCFCs)—has been drastically reduced, forcing these industrial sectors to transition to more ozone friendly chemicals. As these industries have moved toward ODS alternatives such as hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs), the Vintaging Model has evolved into a tool for estimating the rise in consumption and emissions of these alternatives, and the decline of ODS consumption and emissions.

The Vintaging Model estimates emissions from five ODS substitute end-use sectors: refrigeration and air-conditioning, foams, aerosols, solvents, and fire-extinguishing. Within these sectors, there are more than 60 independently modeled end-uses. The model requires information on the market growth for each of the end-uses, a history of the market transition from ODS to alternatives, and the characteristics of each end-use such as market size or charge sizes and loss rates. As ODS are phased out, a percentage of the market share originally filled by the ODS is allocated to each of its substitutes.

The model, named for its method of tracking the emissions of annual “vintages” of new equipment that enter into service, is a “bottom-up” model. It models the consumption of chemicals based on estimates of the quantity of equipment or products sold, serviced, and retired each year, and the amount of the chemical required to manufacture and/or maintain the equipment. The Vintaging Model makes use of this market information to build an inventory of the in-use stocks of the equipment and ODS and ODS substitute in each of the end-uses. The simulation is considered to be a “business-as-usual” baseline case, and does not incorporate measures to reduce or eliminate the emissions of these gases other than those regulated by U.S. law or otherwise common in the industry. Emissions are estimated by applying annual leak rates, service emission rates, and disposal emission rates to each population of equipment. By aggregating the emission and consumption output from the different end-uses, the model produces estimates of total annual use and emissions of each chemical.

The Vintaging Model synthesizes data from a variety of sources, including data from the ODS Tracking System maintained by the Stratospheric Protection Division, the Greenhouse Gas Reporting Program maintained by the Climate Change Division, and information from submissions to EPA under the Significant New Alternatives Policy (SNAP) program. Published sources include documents prepared by the United Nations Environment Programme (UNEP) Technical Options Committees, reports from the Alternative Fluorocarbons Environmental Acceptability Study (AFEAS), and conference proceedings from the International Conferences on Ozone Protection Technologies and Earth Technologies Forums. EPA also coordinates extensively with numerous trade associations and individual companies. For example, the Alliance for Responsible Atmospheric Policy; the Air-Conditioning, Heating and Refrigeration Institute; the Association of Home Appliance Manufacturers; the American Automobile Manufacturers Association; and many of their member companies have provided valuable information over the years. In some instances the unpublished information that the EPA uses in the model is classified as Confidential Business Information (CBI). The annual emissions inventories of chemicals are aggregated in such a way that CBI cannot be inferred. Full public disclosure of the inputs to the Vintaging Model would jeopardize the security of the CBI that has been entrusted to the EPA.

The following sections discuss the emission equations used in the Vintaging Model for each broad end-use category. These equations are applied separately for each chemical used within each of the different end-uses. In the majority of these end-uses, more than one ODS substitute chemical is used.

In general, the modeled emissions are a function of the amount of chemical consumed in each end-use market. Estimates of the consumption of ODS alternatives can be inferred by determining the transition path of each regulated ODS used in the early 1990s. Using data gleaned from a variety of sources, assessments are made regarding which alternatives have been used, and what fraction of the ODS market in each end-use has been captured by a given alternative. By combining this with estimates of the total end-use market growth, a consumption value can be estimated for each chemical used within each end-use.

#### Methodology

The Vintaging Model estimates the use and emissions of ODS alternatives by taking the following steps:

1. *Gather historical data.* The Vintaging Model is populated with information on each end-use, taken from published sources and industry experts.

2. *Simulate the implementation of new, non-ODS technologies.* The Vintaging Model uses detailed characterizations of the existing uses of the ODS, as well as data on how the substitutes are replacing the ODS, to simulate the implementation of new technologies that enter the market in compliance with ODS phase-out policies. As part of this simulation, the ODS substitutes are introduced in each of the end-uses over time as seen historically and as needed to comply with the ODS phase-out and other regulations.

3. *Estimate emissions of the ODS substitutes.* The chemical use is estimated from the amount of substitutes that are required each year for the manufacture, installation, use, or servicing of products. The emissions are estimated from the emission profile for each vintage of equipment or product in each end-use. By aggregating the emissions from each vintage, a time profile of emissions from each end-use is developed.

Each set of end-uses is discussed in more detail in the following sections.

### Refrigeration and Air-Conditioning

For refrigeration and air conditioning products, emission calculations are split into two categories: emissions during equipment lifetime, which arise from annual leakage and service losses, and disposal emissions, which occur at the time of discard. Two separate steps are required to calculate the lifetime emissions from leakage and service, and the emissions resulting from disposal of the equipment. For any given year, these lifetime emissions (for existing equipment) and disposal emissions (from discarded equipment) are summed to calculate the total emissions from refrigeration and air-conditioning. As new technologies replace older ones, it is generally assumed that there are improvements in their leak, service, and disposal emission rates.

#### Step 1: Calculate lifetime emissions

Emissions from any piece of equipment include both the amount of chemical leaked during equipment operation and the amount emitted during service. Emissions from leakage and servicing can be expressed as follows:

$$Es_j = (l_a + l_s) \times \sum_{i=1}^k Qc_{j-i+1} \quad \text{for } i = 1 \rightarrow k$$

where:

|       |   |  |
|-------|---|--|
| $Es$  | = | Emissions from Equipment Serviced. Emissions in year $j$ from normal leakage and servicing (including recharging) of equipment.    |
| $l_a$ | = | Annual Leak Rate. Average annual leak rate during normal equipment operation (expressed as a percentage of total chemical charge). |
| $l_s$ | = | Service Leak Rate. Average leakage during equipment servicing (expressed as a percentage of total chemical charge).                |
| $Qc$  | = | Quantity of Chemical in New Equipment. Total amount of a specific chemical used to charge new equipment in a given year by weight. |
| $I$   | = | Counter, runs from 1 to lifetime ( $k$ ).  |
| $j$   | = | Year of emission.  |
| $k$   | = | Lifetime. The average lifetime of the equipment.   |

#### Step 2: Calculate disposal emissions

The disposal emission equations assume that a certain percentage of the chemical charge will be emitted to the atmosphere when that vintage is discarded. Disposal emissions are thus a function of the quantity of chemical contained in the retiring equipment fleet and the proportion of chemical released at disposal:

$$Ed_j = Qc_{j-k+1} \times [1 - (rm \times rc)]$$

where:

|      |   |  |
|------|---|--|
| $Ed$ | = | Emissions from Equipment Disposed. Emissions in year $j$ from the disposal of equipment.   |
| $Qc$ | = | Quantity of Chemical in New Equipment. Total amount of a specific chemical used to charge new equipment in year $j-k+1$ , by weight. |

|           |   |  |
|-----------|---|--|
| <i>rm</i> | = | Chemical Remaining. Amount of chemical remaining in equipment at the time of disposal (expressed as a percentage of total chemical charge).                      |
| <i>rc</i> | = | Chemical Recovery Rate. Amount of chemical that is recovered just prior to disposal (expressed as a percentage of chemical remaining at disposal ( <i>rm</i> )). |
| <i>j</i>  | = | Year of emission.  |
| <i>k</i>  | = | Lifetime. The average lifetime of the equipment.   |

**Step 3: Calculate total emissions**

Finally, lifetime and disposal emissions are summed to provide an estimate of total emissions.

$$E_j = Es_j + Ed_j$$

where:

|           |   |   |
|-----------|---|---|
| <i>E</i>  | = | Total Emissions. Emissions from refrigeration and air conditioning equipment in year <i>j</i> .                               |
| <i>Es</i> | = | Emissions from Equipment Serviced. Emissions in year <i>j</i> from leakage and servicing (including recharging) of equipment. |
| <i>Ed</i> | = | Emissions from Equipment Disposed. Emissions in year <i>j</i> from the disposal of equipment.                                 |
| <i>j</i>  | = | Year of emission.   |

**Assumptions**

The assumptions used by the Vintaging Model to trace the transition of each type of equipment away from ODS are presented in Table A-169, below. As new technologies replace older ones, it is generally assumed that there are improvements in their leak, service, and disposal emission rates. Additionally, the market for each equipment type is assumed to grow independently, according to annual growth rates.

**Table A-169: Refrigeration and Air-Conditioning Market Transition Assumptions**

| Initial Market Segment                             | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|--|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|  | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| <b>Centrifugal Chillers</b>                        |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11   | HCFC-123           | 1993       | 1993                                      | 45%                        | Unknown              |            |   |                            |                     |            |   |                            | 1.6%        |
|  | HCFC-22            | 1991       | 1993                                      | 16%                        | HFC-134a             | 2000       | 2010                                      | 100%                       | None                |            |   |                            |             |
|  | HFC-134a           | 1992       | 1993                                      | 39%                        | None                 |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HFC-134a           | 1992       | 1994                                      | 53%                        | None                 |            |   |                            |                     |            |   |                            | 1.5%        |
|  | HCFC-22            | 1991       | 1994                                      | 16%                        | HFC-134a             | 2000       | 2010                                      | 100%                       | None                |            |   |                            |             |
| R-500  | HCFC-123           | 1993       | 1994                                      | 31%                        | Unknown              |            |   |                            |                     |            |   |                            | 1.5%        |
|  | HFC-134a           | 1992       | 1994                                      | 53%                        | None                 |            |   |                            |                     |            |   |                            |             |
| CFC-114  | HCFC-22            | 1991       | 1994                                      | 16%                        | HFC-134a             | 2000       | 2010                                      | 100%                       | None                |            |   |                            | 1.4%        |
|  | HCFC-123           | 1993       | 1994                                      | 31%                        | Unknown              |            |   |                            |                     |            |   |                            |             |
|  | HFC-236fa          | 1993       | 1996                                      | 100%                       | HFC-134a             | 1998       | 2009                                      | 100%                       | None                |            |   |                            |             |
| <b>Cold Storage</b>                                |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HCFC-22            | 1990       | 1993                                      | 65%                        | R-404A               | 1996       | 2010                                      | 75%                        | None                |            |   |                            | 3.1%        |
|  |                    |            |   |                            | R-507                | 1996       | 2010                                      | 25%                        | None                |            |   |                            |             |
|  | R-404A             | 1994       | 1996                                      | 26%                        | None                 |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | R-507              | 1994       | 1996                                      | 9%                         | None                 |            |   |                            |                     |            |   |                            | 3.0%        |
|  | HCFC-22            | 1992       | 1993                                      | 100%                       | R-404A               | 1996       | 2009                                      | 8%                         | None                |            |   |                            |             |
|  |                    |            |   |                            | R-507                | 1996       | 2009                                      | 3%                         | None                |            |   |                            |             |
| R-502  |                    |            |   |                            | R-404A               | 2009       | 2010                                      | 68%                        | None                |            |   |                            | 2.6%        |
|  |                    |            |   |                            | R-507                | 2009       | 2010                                      | 23%                        | None                |            |   |                            |             |
|  | HCFC-22            | 1990       | 1993                                      | 40%                        | R-404A               | 1996       | 2010                                      | 38%                        | None                |            |   |                            |             |
|  |                    |            |   |                            | R-507                | 1996       | 2010                                      | 12%                        | None                |            |   |                            |             |
|  |                    |            |   |                            | Non-ODP/GWP          | 1996       | 2010                                      | 50%                        | None                |            |   |                            |             |
|  | R-404A             | 1993       | 1996                                      | 45%                        | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-507              | 1994       | 1996                                      | 15%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Commercial Unitary Air Conditioners (Large)</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HCFC-22            | 1992       | 1993                                      | 100%                       | R-410A               | 2001       | 2005                                      | 5%                         | None                |            |   |                            | 1.3%        |
|  |                    |            |   |                            | R-407C               | 2006       | 2009                                      | 1%                         | None                |            |   |                            |             |
|  |                    |            |   |                            | R-410A               | 2006       | 2009                                      | 9%                         | None                |            |   |                            |             |
|  |                    |            |   |                            | R-407C               | 2009       | 2010                                      | 5%                         | None                |            |   |                            |             |
|  |                    |            |   |                            | R-410A               | 2009       | 2010                                      | 81%                        | None                |            |   |                            |             |
| <b>Commercial Unitary Air Conditioners (Small)</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HCFC-22            | 1992       | 1993                                      | 100%                       | R-410A               | 1996       | 2000                                      | 3%                         | None                |            |   |                            | 1.3%        |
|  |                    |            |   |                            | R-410A               | 2001       | 2005                                      | 18%                        | None                |            |   |                            |             |
|  |                    |            |   |                            | R-410A               | 2006       | 2009                                      | 8%                         | None                |            |   |                            |             |
|  |                    |            |   |                            | R-410A               | 2009       | 2010                                      | 71%                        | None                |            |   |                            |             |

| Initial Market Segment                                 | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|--|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|  | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| <b>Dehumidifiers</b>                                   |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HFC-134a           | 1997       | 1997                                      | 89%                        | None                 |            |   |                            |                     |            |   |                            | 1.3%        |
|  | R-410A             | 2007       | 2010                                      | 11%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Ice Makers</b>                                      |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HFC-134a           | 1993       | 1995                                      | 25%                        | None                 |            |   |                            |                     |            |   |                            | 2.1%        |
|  | R-404A             | 1993       | 1995                                      | 75%                        |                      |            |   |                            |                     |            |   |                            |             |
| <b>Industrial Process Refrigeration</b>                |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11   | HCFC-123           | 1992       | 1994                                      | 70%                        | Unknown              |            |   |                            |                     |            |   |                            | 3.2%        |
|  | HFC-134a           | 1992       | 1994                                      | 15%                        | None                 |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HCFC-22            | 1991       | 1994                                      | 15%                        | HFC-134a             | 1995       | 2010                                      | 100%                       | None                |            |   |                            | 3.1%        |
|  | HCFC-22            | 1991       | 1994                                      | 10%                        | HFC-134a             | 1995       | 2010                                      | 15%                        |                     |            |   |                            |             |
|  |                    |            |   |                            | R-404A               | 1995       | 2010                                      | 50%                        |                     |            |   |                            |             |
|  |                    |            |   |                            | R-410A               | 1999       | 2010                                      | 20%                        |                     |            |   |                            |             |
|  |                    |            |   |                            | R-507                | 1995       | 2010                                      | 15%                        |                     |            |   |                            |             |
| HCFC-22  | HCFC-123           | 1992       | 1994                                      | 35%                        | Unknown              |            |   |                            | None                |            |   |                            | 3.0%        |
|  | HFC-134a           | 1992       | 1994                                      | 50%                        | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-401A             | 1995       | 1996                                      | 5%                         | HFC-134a             | 1997       | 2000                                      | 100%                       |                     |            |   |                            |             |
|  | HFC-134a           | 1995       | 2009                                      | 2%                         | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-404A             | 1995       | 2009                                      | 5%                         | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-410A             | 1999       | 2009                                      | 2%                         | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-507              | 1995       | 2009                                      | 2%                         | None                 |            |   |                            |                     |            |   |                            |             |
|  | HFC-134a           | 2009       | 2010                                      | 14%                        | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-404A             | 2009       | 2010                                      | 45%                        | None                 |            |   |                            |                     |            |   |                            |             |
|  | R-410A             | 2009       | 2010                                      | 18%                        | None                 |            |   |                            |                     |            |   |                            |             |
| R-507  | 2009               | 2010       | 14%                                       | None                       |                      |            |   |                            |                     |            |   |                            |             |
| <b>Mobile Air Conditioners (Passenger Cars)</b>        |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HFC-134a           | 1992       | 1994                                      | 100%                       | HFO-1234yf           | 2012       | 2015                                      | 1%                         | None                |            |   |                            | 0.3%        |
|  |                    |            |   |                            | HFO-1234yf           | 2016       | 2021                                      | 99%                        |                     |            |   |                            |             |
| <b>Mobile Air Conditioners (Light Duty Trucks)</b>     |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HFC-134a           | 1993       | 1994                                      | 100%                       | HFO-1234yf           | 2012       | 2015                                      | 1%                         | None                |            |   |                            | 1.4%        |
|  |                    |            |   |                            | HFO-1234yf           | 2016       | 2021                                      | 99%                        |                     |            |   |                            |             |
| <b>Mobile Air Conditioners (School and Tour Buses)</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HCFC-22            | 1994       | 1995                                      | 0.5%                       | HFC-134a             | 2006       | 2007                                      | 100%                       | None                |            |   |                            | 0.3%        |
|  | HFC-134a           | 1194       | 1997                                      | 99.5%                      | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Mobile Air Conditioners (Transit Buses)</b>         |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HFC-134a           | 1995       | 2009                                      | 100%                       | None                 |            |   |                            |                     |            |   |                            | 0.3%        |
| <b>Mobile Air Conditioners (Trains)</b>                |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HFC-134a           | 2002       | 2009                                      | 50%                        | None                 |            |   |                            |                     |            |   |                            | 0.3%        |

| Initial Market Segment                                   | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|--|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|  | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
|  | R-407C             | 2002       | 2009                                      | 50%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Packaged Terminal Air Conditioners and Heat Pumps</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | R-410A             | 2006       | 2009                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            | 3.0%        |
|  | R-410A             | 2009       | 2010                                      | 90%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Positive Displacement Chillers</b>                    |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HFC-134a           | 2000       | 2009                                      | 9%                         | R-407C               | 2010       | 2020                                      | 60%                        | None                |            |   |                            | 2.5%        |
|  | R-407C             | 2000       | 2009                                      | 1%                         | R-410A               | 2010       | 2020                                      | 40%                        | None                |            |   |                            |             |
|  | HFC-134a           | 2009       | 2010                                      | 81%                        | R-407C               | 2010       | 2020                                      | 60%                        | None                | 2010       | 2020                                      | 40%                        |             |
| CFC-12   | HCFC-22            | 1993       | 1993                                      | 100%                       | R-407C               | 2009       | 2010                                      | 9%                         | None                |            |   |                            | 2.5%        |
|  |                    |            |   |                            | HFC-134a             | 2000       | 2009                                      | 9%                         | R-407C              | 2010       | 2020                                      | 60%                        |             |
|  |                    |            |   |                            | R-407C               | 2000       | 2009                                      | 1%                         | R-410A              | 2010       | 2020                                      | 40%                        |             |
|  |                    |            |   |                            | HFC-134a             | 2009       | 2010                                      | 81%                        | R-407C              | 2010       | 2020                                      | 60%                        |             |
|  |                    |            |   |                            | R-407C               | 2009       | 2010                                      | 9%                         | R-410A              | 2010       | 2020                                      | 40%                        |             |
| <b>Refrigerated Appliances</b>                           |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | HFC-134a           | 1994       | 1995                                      | 100%                       | None                 |            |   |                            |                     |            |   |                            | 1.7%        |
| <b>Residential Unitary Air Conditioners</b>              |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22  | HCFC-22            | 2006       | 2006                                      | 70%                        | R-410A               | 2007       | 2010                                      | 29%                        | None                |            |   |                            | 1.3%        |
|  | R-410A             | 2000       | 2005                                      | 5%                         | R-410A               | 2010       | 2010                                      | 71%                        | None                |            |   |                            |             |
|  | R-410A             | 2000       | 2006                                      | 5%                         | R-410A               | 2006       | 2006                                      | 100%                       | None                |            |   |                            |             |
|  | R-410A             | 2006       | 2006                                      | 20%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Retail Food (Large; Technology Transition)</b>        |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| DX <sup>1</sup>  | DX                 | 2001       | 2006                                      | 67.5%                      | DX                   | 2006       | 2015                                      | 35%                        | None                |            |   |                            | 1.7%        |
|  |                    |            |   |                            | DR <sup>2</sup>      | 2000       | 2015                                      | 23%                        | None                |            |   |                            |             |
|  |                    |            |   |                            | SLS <sup>3</sup>     | 2000       | 2015                                      | 15%                        | None                |            |   |                            |             |
|  | DR                 | 2001       | 2006                                      | 22.5%                      | None                 |            |   |                            |                     |            |   |                            |             |
|  | SLS                | 2001       | 2006                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Retail Food (Large; Refrigerant Transition)</b>       |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12   | R-404A             | 1995       | 2000                                      | 17.5%                      | R-404A               | 2000       | 2000                                      | 3.3%                       | None                |            |   |                            | 1.7%        |
| R-502 <sup>4</sup>                                       |                    |            |   |                            | R-407A               | 2011       | 2015                                      | 63.3%                      | None                |            |   |                            |             |
|  | R-507              | 1995       | 2000                                      | 7.5%                       | R-507                | 2001       | 2005                                      | 70%                        | R-404A              | 2006       | 2010                                      | 29%                        |             |
|  |                    |            |   |                            | R-407A               | 2006       | 2010                                      | 30%                        | R-407A              | 2006       | 2010                                      | 71%                        |             |
|  | HCFC-22            | 1995       | 2000                                      | 75%                        | R-404A               | 2006       | 2010                                      | 13.3%                      | None                | 2011       | 2015                                      | 100%                       |             |

| Initial Market Segment                          | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|---|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|   | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
|   |                    |            |   |                            | R-407A               | 2001       | 2005                                      | 1.3%                       | None                |            |   |                            |             |
|   |                    |            |   |                            | R-404A               | 2001       | 2005                                      | 12%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | R-507                | 2001       | 2005                                      | 6.7%                       | R-407A              | 2011       | 2015                                      | 100%                       |             |
|   |                    |            |   |                            | R-404A               | 2006       | 2010                                      | 34%                        | R-407A              | 2011       | 2015                                      | 100%                       |             |
|   |                    |            |   |                            | R-404A               | 2006       | 2010                                      | 7.3%                       | None                |            |   |                            |             |
|   |                    |            |   |                            | R-407A               | 2006       | 2010                                      | 25.3%                      | None                |            |   |                            |             |
| <b>Retail Food (Large Condensing Units)</b>     |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22   | R-402A             | 1995       | 2005                                      | 5%                         | R-404A               | 2006       | 2006                                      | 100%                       | None                |            |   |                            | 1.5%        |
|   | R-404A             | 1995       | 2005                                      | 25%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-507              | 1995       | 2005                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-404A             | 2008       | 2010                                      | 45%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-507              | 2008       | 2010                                      | 15%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Retail Food (Small Condensing Units)</b>     |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22   | R-401A             | 1995       | 2005                                      | 6%                         | HFC-134a             | 2006       | 2006                                      | 100%                       | None                |            |   |                            | 1.6%        |
|   | R-402A             | 1995       | 2005                                      | 4%                         | HFC-134a             | 2006       | 2006                                      | 100%                       | None                |            |   |                            |             |
|   | HFC-134a           | 1993       | 2005                                      | 30%                        |                      |            |   |                            |                     |            |   |                            |             |
|   | R-404A             | 1995       | 2005                                      | 30%                        |                      |            |   |                            |                     |            |   |                            |             |
|   | R-404A             | 2008       | 2010                                      | 30%                        |                      |            |   |                            |                     |            |   |                            |             |
| <b>Retail Food (Small)</b>                      |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12  | HCFC-22            | 1990       | 1993                                      | 91%                        | HFC-134a             | 1993       | 1995                                      | 91%                        | CO <sub>2</sub>     | 2012       | 2015                                      | 1%                         | 2.2%        |
|   |                    |            |   |                            |                      |            |   |                            | Non-ODP/GWP         | 2012       | 2015                                      | 3.7%                       |             |
|   |                    |            |   |                            |                      |            |   |                            | CO <sub>2</sub>     | 2016       | 2016                                      | 11%                        |             |
|   |                    |            |   |                            |                      |            |   |                            | Non-ODP/GWP         | 2016       | 2016                                      | 17.3%                      |             |
|   |                    |            |   |                            | HFC-134a             | 2000       | 2009                                      | 9%                         |                     |            |   |                            |             |
|   | R-404A             | 1990       | 1993                                      | 9%                         | Non-ODP/GWP          | 2016       | 2016                                      | 30%                        |                     |            |   |                            |             |
| <b>Retail Food (Vending Machines)</b>           |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12  | HFC-134a           | 1995       | 1998                                      | 90%                        | CO <sub>2</sub>      | 2012       | 2012                                      | 1%                         | None                |            |   |                            | -0.03%      |
|   |                    |            |   |                            | CO <sub>2</sub>      | 2013       | 2017                                      | 39%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | Propane              | 2014       | 2014                                      | 1%                         | None                |            |   |                            |             |
|   |                    |            |   |                            | Propane              | 2015       | 2015                                      | 49%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | R-450A               | 2019       | 2019                                      | 5%                         | None                |            |   |                            |             |
|   |                    |            |   |                            | R-513A               | 2019       | 2019                                      | 5%                         | None                |            |   |                            |             |
|   | R-404A             | 1995       | 1998                                      | 10%                        | R-450A               | 2019       | 2019                                      | 50%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | R-513A               | 2019       | 2019                                      | 50%                        | None                |            |   |                            |             |
| <b>Transport Refrigeration (Road Transport)</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12  | HFC-134a           | 1993       | 1995                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            | 5.5%        |
|   | R-404A             | 1993       | 1995                                      | 60%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | HCFC-22            | 1993       | 1995                                      | 30%                        | R-410A               | 2000       | 2003                                      | 5%                         | CO <sub>2</sub>     | 2017       | 2021                                      | 5%                         |             |

| Initial Market Segment                                      | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|---|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|   | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
|   |                    |            |   |                            | R-404A               | 2006       | 2010                                      | 95%                        | CO <sub>2</sub>     | 2017       | 2021                                      | 5%                         |             |
| <b>Transport Refrigeration (Intermodal Containers)</b>      |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12  | HFC-134a           | 1993       | 1993                                      | 60%                        | CO <sub>2</sub>      | 2017       | 2021                                      | 5%                         | None                |            |   |                            | 7.3%        |
|   | R-404A             | 1993       | 1993                                      | 5%                         | CO <sub>2</sub>      | 2017       | 2021                                      | 5%                         | None                |            |   |                            |             |
|   | HCFC-22            | 1993       | 1993                                      | 35%                        | HFC-134a             | 2000       | 2010                                      | 10%                        | CO <sub>2</sub>     | 2017       | 2021                                      | 5%                         |             |
| <b>Transport Refrigeration (Merchant Fishing Transport)</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22   | HFC-134a           | 1993       | 1995                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            | 5.7%        |
|   | R-507              | 1994       | 1995                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-404A             | 1993       | 1995                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | HCFC-22            | 1993       | 1995                                      | 70%                        | R-407C               | 2000       | 2005                                      | 3%                         | R-410A              | 2005       | 2007                                      | 100%                       |             |
|   |                    |            |   |                            | R-507                | 2006       | 2010                                      | 49%                        | None                |            |   |                            |             |
|   |                    |            |   | R-404                      | 2006                 | 2010       | 49%                                       | None                       |                     |            |   |                            |             |
| <b>Transport Refrigeration (Reefer Ships)</b>               |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22   | HFC-134a           | 1993       | 1995                                      | 3.3%                       | None                 |            |   |                            |                     |            |   |                            | 4.2%        |
|   | R-507              | 1994       | 1995                                      | 3.3%                       | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-404A             | 1993       | 1995                                      | 3.3%                       | None                 |            |   |                            |                     |            |   |                            |             |
|   | HCFC-22            | 1993       | 1995                                      | 90%                        | HFC-134a             | 2006       | 2010                                      | 25%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | R-507                | 2006       | 2010                                      | 25%                        | None                |            |   |                            |             |
|   |                    |            |   | R-404A                     | 2006                 | 2010       | 25%                                       | None                       |                     |            |   |                            |             |
|   |                    |            |   | R-407C                     | 2006                 | 2010       | 25%                                       | None                       |                     |            |   |                            |             |
| <b>Transport Refrigeration (Vintage Rail Transport)</b>     |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12  | HCFC-22            | 1993       | 1995                                      | 100%                       | HFC-134a             | 1996       | 2000                                      | 100%                       | None                |            |   |                            | -100%       |
| <b>Transport Refrigeration (Modern Rail Transport)</b>      |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HFC-134a  | R-404A             | 1999       | 1999                                      | 50%                        | None                 |            |   |                            |                     |            |   |                            | 0.3%        |
|   | HFC-134A           | 2005       | 2005                                      | 50%                        | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Water-Source and Ground-Source Heat Pumps</b>            |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22   | R-407C             | 2000       | 2006                                      | 5%                         | None                 |            |   |                            |                     |            |   |                            | 1.3%        |
|   | R-410A             | 2000       | 2006                                      | 5%                         | None                 |            |   |                            |                     |            |   |                            |             |
|   | HFC-134a           | 2000       | 2009                                      | 2%                         | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-407C             | 2006       | 2009                                      | 2.5%                       | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-410A             | 2006       | 2009                                      | 4.5%                       | None                 |            |   |                            |                     |            |   |                            |             |
|   | HFC-134a           | 2009       | 2010                                      | 18%                        | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-407C             | 2009       | 2010                                      | 22.5%                      | None                 |            |   |                            |                     |            |   |                            |             |
|   | R-410A             | 2009       | 2010                                      | 40.5%                      | None                 |            |   |                            |                     |            |   |                            |             |
| <b>Window Units</b>   |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| HCFC-22   | R-410A             | 2008       | 2009                                      | 10%                        | None                 |            |   |                            |                     |            |   |                            | 4.0%        |
|   | R-410A             | 2009       | 2010                                      | 90%                        | None                 |            |   |                            |                     |            |   |                            |             |

<sup>1</sup> DX refers to direct expansion systems where the compressors are mounted together in a rack and share suction and discharge refrigeration lines that run throughout the store, feeding refrigerant to the display cases in the sales area.

<sup>2</sup> DR refers to distributed refrigeration systems that consist of multiple smaller units that are located close to the display cases that they serve such as on the roof above the cases, behind a nearby wall, or on top of or next to the case in the sales area.

<sup>3</sup> SLS refers to secondary loop systems wherein a secondary fluid such as glycol or carbon dioxide is cooled by the primary refrigerant in the machine room and then pumped throughout the store to remove heat from the display equipment.

<sup>4</sup> The CFC-12 large retail food market for new systems transitioned to R-502 from 1998 to 1990, and subsequently transitioned to HCFC-22 from 1990 to 1993. These transitions are not shown in the table in order to provide the HFC transitions in greater detail.

<sup>5</sup> HCFC-22 for new equipment after 2010 is assumed to be reclaimed material.

Table A-170 presents the average equipment lifetimes and annual HFC emission rates (for servicing and leaks) for each end-use assumed by the Vintaging Model.

**Table A-170: Refrigeration and Air-Conditioning Lifetime Assumptions**

| End-Use                          | Lifetime (Years) | HFC Emission Rates (%) |
|----------------------------------|------------------|------------------------|
| Centrifugal Chillers             | 20 – 27          | 2.0 – 10.9             |
| Cold Storage                     | 20 – 25          | 15.0                   |
| Commercial Unitary A/C           | 15               | 7.9 – 8.6              |
| Dehumidifiers                    | 11               | 0.5                    |
| Ice Makers                       | 8                | 3.0                    |
| Industrial Process Refrigeration | 25               | 3.6 – 12.3             |
| Mobile Air Conditioners          | 5 – 16           | 2.3 – 18.0             |
| Positive Displacement Chillers   | 20               | 0.5 – 1.5              |
| PTAC/PTHP                        | 12               | 3.9                    |
| Retail Food                      | 10 – 20          | 1.0 – 25               |
| Refrigerated Appliances          | 14               | 0.6                    |
| Residential Unitary A/C          | 15               | 11.8                   |
| Transport Refrigeration          | 9 – 40           | 19.4 – 36.4            |
| Water & Ground Source Heat Pumps | 20               | 3.9                    |
| Window Units                     | 12               | 0.6                    |

### Aerosols

ODSs, HFCs, and many other chemicals are used as propellant aerosols. Pressurized within a container, a nozzle releases the chemical, which allows the product within the can to also be released. Two types of aerosol products are modeled: metered dose inhalers (MDI) and consumer aerosols. In the United States, the use of CFCs in consumer aerosols was banned in 1978, and many products transitioned to hydrocarbons or “not-in-kind” technologies, such as solid deodorants and finger-pump hair sprays. However, MDIs can continue to use CFCs as propellants because their use has been deemed essential. Essential use exemptions granted to the United States under the Montreal Protocol for CFC use in MDIs are limited to the treatment of asthma and chronic obstructive pulmonary disease.

All HFCs and PFCs used in aerosols are assumed to be emitted in the year of manufacture. Since there is currently no aerosol recycling, it is assumed that all of the annual production of aerosol propellants is released to the atmosphere. The following equation describes the emissions from the aerosols sector.

$$E_j = Qc_j$$

where:

- $E$  = Emissions. Total emissions of a specific chemical in year  $j$  from use in aerosol products, by weight.
- $Qc$  = Quantity of Chemical. Total quantity of a specific chemical contained in aerosol products sold in year  $j$ , by weight.
- $j$  = Year of emission.

### Transition Assumptions

Transition assumptions and growth rates for those items that use ODSs or HFCs as propellants, including vital medical devices and specialty consumer products, are presented in Table A-171.

**Table A-171: Aerosol Product Transition Assumptions**

| Initial Market Segment              | Primary Substitute   |            |   |                            | Secondary Substitute |            |   |                            | Growth Rate |
|-------------------------------------|----------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|-------------|
|                                     | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| <b>MDIs</b>                         |                      |            |   |                            |                      |            |   |                            |             |
| CFC Mix <sup>a</sup>                | HFC-134a             | 1997       | 1997                                      | 6%                         | None                 |            |   |                            | 0.8%        |
|                                     | Non-ODP/GWP          | 1998       | 2007                                      | 7%                         | None                 |            |   |                            |             |
|                                     | CFC Mix <sup>a</sup> | 2000       | 2000                                      | 87%                        | HFC-134a             | 2002       | 2002                                      | 34%                        |             |
|                                     |                      |            |   |                            | HFC-134a             | 2003       | 2009                                      | 47%                        |             |
|                                     |                      |            |   |                            | HFC-227ea            | 2006       | 2009                                      | 5%                         |             |
|                                     |                      |            |   |                            | HFC-134a             | 2010       | 2011                                      | 6%                         |             |
|                                     |                      |            |   |                            | HFC-227ea            | 2010       | 2011                                      | 1%                         |             |
|                                     |                      |            |   |                            | HFC-134a             | 2011       | 2012                                      | 3%                         |             |
|                                     |                      |            |   |                            | HFC-227ea            | 2011       | 2012                                      | 0.3%                       |             |
|                                     |                      |            |   |                            | HFC-134a             | 2014       | 2014                                      | 3%                         |             |
|                                     |                      |            |   | HFC-227ea                  | 2014                 | 2014       | 0.3%                                      |                            |             |
| <b>Consumer Aerosols (Non-MDIs)</b> |                      |            |   |                            |                      |            |   |                            |             |
| NA <sup>b</sup>                     | HFC-152a             | 1990       | 1991                                      | 50%                        | None                 |            |   |                            | 2.0%        |
|                                     | HFC-134a             | 1995       | 1995                                      | 50%                        | HFC-152a             | 1997       | 1998                                      | 44%                        |             |
|                                     |                      |            |   |                            | HFC-152a             | 2001       | 2005                                      | 36%                        |             |

<sup>a</sup> CFC Mix consists of CFC-11, CFC-12 and CFC-114 and represents the weighted average of several CFCs consumed for essential use in MDIs from 1993 to 2008.

<sup>b</sup> Consumer Aerosols transitioned away from ODS prior to 1985, the year in which the Vintaging Model begins. The portion of the market that is now using HFC propellants is modeled.

**Solvents**

ODSs, HFCs, PFCs and other chemicals are used as solvents to clean items. For example, electronics may need to be cleaned after production to remove any manufacturing process oils or residues left. Solvents are applied by moving the item to be cleaned within a bath or stream of the solvent. Generally, most solvents are assumed to remain in the liquid phase and are not emitted as gas. Thus, emissions are considered “incomplete,” and are a fixed percentage of the amount of solvent consumed in a year. The remainder of the consumed solvent is assumed to be reused or disposed without being released to the atmosphere. The following equation calculates emissions from solvent applications.

$$E_j = l \times Qc_j$$

where:

- E* = Emissions. Total emissions of a specific chemical in year *j* from use in solvent applications, by weight.
- l* = Percent Leakage. The percentage of the total chemical that is leaked to the atmosphere, assumed to be 90 percent.
- Qc* = Quantity of Chemical. Total quantity of a specific chemical sold for use in solvent applications in the year *j*, by weight.
- j* = Year of emission.

**Transition Assumptions**

The transition assumptions and growth rates used within the Vintaging Model for electronics cleaning, metals cleaning, precision cleaning, and adhesives, coatings and inks, are presented in Table A-172.

**Table A-172: Solvent Market Transition Assumptions**

| Initial Market Segment           | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Growth Rate |
|----------------------------------|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|-------------|
|                                  | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| <b>Adhesives</b>                 |                    |            |   |                            |                      |            |   |                            |             |
| CH <sub>3</sub> CCl <sub>3</sub> | Non-ODP/GWP        | 1994       | 1995                                      | 100%                       | None                 |            |   |                            | 2.0%        |
| <b>Electronics</b>               |                    |            |   |                            |                      |            |   |                            |             |
| CFC-113                          | Semi-Aqueous       | 1994       | 1995                                      | 52%                        | None                 |            |   |                            | 2.0%        |
|                                  | HCFC-225ca/cb      | 1994       | 1995                                      | 0.2%                       | Unknown              |            |   |                            |             |
|                                  | HFC-43-10mee       | 1995       | 1996                                      | 0.7%                       | None                 |            |   |                            |             |
|                                  | HFE-7100           | 1994       | 1995                                      | 0.7%                       | None                 |            |   |                            |             |
|                                  | nPB                | 1992       | 1996                                      | 5%                         | None                 |            |   |                            |             |
|                                  | Methyl Siloxanes   | 1992       | 1996                                      | 0.8%                       | None                 |            |   |                            |             |
|                                  | No-Clean           | 1992       | 1996                                      | 40%                        | None                 |            |   |                            |             |
| CH <sub>3</sub> CCl <sub>3</sub> | Non-ODP/GWP        | 1996       | 1997                                      | 99.8%                      | None                 |            |   |                            | 2.0%        |
|                                  | PFC/PFPE           | 1996       | 1997                                      | 0.2%                       | Non-ODP/GWP          | 2000       | 2003                                      | 90%                        |             |
|                                  |                    |            |   |                            | Non-ODP/GWP          | 2005       | 2009                                      | 10%                        |             |
| <b>Metals</b>                    |                    |            |   |                            |                      |            |   |                            |             |
| CH <sub>3</sub> CCl <sub>3</sub> | Non-ODP/GWP        | 1992       | 1996                                      | 100%                       | None                 |            |   |                            | 2.0%        |
| CFC-113                          | Non-ODP/GWP        | 1992       | 1996                                      | 100%                       | None                 |            |   |                            | 2.0%        |
| CCl <sub>4</sub>                 | Non-ODP/GWP        | 1992       | 1996                                      | 100%                       | None                 |            |   |                            | 2.0%        |
| <b>Precision</b>                 |                    |            |   |                            |                      |            |   |                            |             |
| CH <sub>3</sub> CCl <sub>3</sub> | Non-ODP/GWP        | 1995       | 1996                                      | 99.3%                      | None                 |            |   |                            | 2.0%        |
|                                  | HFC-43-10mee       | 1995       | 1996                                      | 0.6%                       | None                 |            |   |                            |             |
|                                  | PFC/PFPE           | 1995       | 1996                                      | 0.1%                       | Non-ODP/GWP          | 2000       | 2003                                      | 90%                        |             |
|                                  |                    |            |   |                            | Non-ODP/GWP          | 2005       | 2009                                      | 10%                        |             |
| CFC-113                          | Non-ODP/GWP        | 1995       | 1996                                      | 96%                        | None                 |            |   |                            | 2.0%        |
|                                  | HCFC-225ca/cb      | 1995       | 1996                                      | 1%                         | Unknown              |            |   |                            |             |
|                                  | HFE-7100           | 1995       | 1996                                      | 3%                         | None                 |            |   |                            |             |

Non-ODP/GWP includes chemicals with zero ODP and low GWP, such as hydrocarbons and ammonia, as well as not-in-kind alternatives such as "no clean" technologies.

### Fire Extinguishing

ODSs, HFCs, PFCs and other chemicals are used as fire-extinguishing agents, in both hand-held "streaming" applications as well as in built-up "flooding" equipment similar to water sprinkler systems. Although these systems are generally built to be leak-tight, some leaks do occur and of course emissions occur when the agent is released. Total emissions from fire extinguishing are assumed, in aggregate, to equal a percentage of the total quantity of chemical in operation at a given time. For modeling purposes, it is assumed that fire extinguishing equipment leaks at a constant rate for an average equipment lifetime, as shown in the equation below. In streaming systems, non-halon emissions are assumed to be 3.5 percent of all chemical in use in each year, while in flooding systems 2.5 percent of the installed base of chemical is assumed to leak annually. Halon systems are assumed to leak at higher rates. The equation is applied for a single year, accounting for all fire protection equipment in operation in that year. Each fire protection agent is modeled separately. In the Vintaging Model, streaming applications have a 12-year lifetime and flooding applications have a 20-year lifetime.

$$E_j = r \times \sum Q_{C_{j+i}} \text{ for } i=1 \rightarrow k$$

where:

- E* = Emissions. Total emissions of a specific chemical in year *j* for streaming fire extinguishing equipment, by weight.
- r* = Percent Released. The percentage of the total chemical in operation that is released to the atmosphere.
- Q<sub>c</sub>* = Quantity of Chemical. Total amount of a specific chemical used in new fire extinguishing equipment in a given year, *j-i+1*, by weight.
- i* = Counter, runs from 1 to lifetime (*k*).

$j$  = Year of emission.

$k$  = Lifetime. The average lifetime of the equipment.

### Transition Assumptions

Transition assumptions and growth rates for these two fire extinguishing types are presented in Table A-173.

**Table A-173: Fire Extinguishing Market Transition Assumptions**

| Initial Market Segment  | Primary Substitute             |            |   |                            | Secondary Substitute |            |   |                            | Growth Rate |
|-------------------------|--------------------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|-------------|
|                         | Name of Substitute             | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| <b>Flooding Agents</b>  |                                |            |   |                            |                      |            |   |                            |             |
| Halon-1301              | Halon-1301*                    | 1994       | 1994                                      | 4%                         | Unknown              |            |   |                            | 2.2%        |
|                         | HFC-23                         | 1994       | 1999                                      | 0.2%                       | None                 |            |   |                            |             |
|                         | HFC-227ea                      | 1994       | 1999                                      | 18%                        | FK-5-1-12            | 2003       | 2010                                      | 10%                        |             |
|                         | Non-ODP/GWP                    | 1994       | 1994                                      | 46%                        | HFC-125              | 2001       | 2008                                      | 10%                        |             |
|                         | Non-ODP/GWP                    | 1995       | 2034                                      | 10%                        | FK-5-1-12            | 2003       | 2010                                      | 7%                         |             |
|                         | Non-ODP/GWP                    | 1998       | 2027                                      | 10%                        | None                 |            |   |                            |             |
|                         | C <sub>4</sub> F <sub>10</sub> | 1994       | 1999                                      | 1%                         | FK-5-1-12            | 2003       | 2003                                      | 100%                       |             |
|                         | HFC-125                        | 1997       | 2006                                      | 11%                        | None                 |            |   |                            |             |
| <b>Streaming Agents</b> |                                |            |   |                            |                      |            |   |                            |             |
| Halon-1211              | Halon-1211*                    | 1992       | 1992                                      | 5%                         | Unknown              |            |   |                            | 3.0%        |
|                         | HFC-236fa                      | 1997       | 1999                                      | 3%                         | None                 |            |   |                            |             |
|                         | Halotron                       | 1994       | 1995                                      | 0.1%                       | None                 |            |   |                            |             |
|                         | Halotron                       | 1996       | 2000                                      | 5.4%                       | Non-ODP/GWP          | 2020       | 2020                                      | 56%                        |             |
|                         | Non-ODP/GWP                    | 1993       | 1994                                      | 56%                        | None                 |            |   |                            |             |
|                         | Non-ODP/GWP                    | 1995       | 2024                                      | 20%                        | None                 |            |   |                            |             |
|                         | Non-ODP/GWP                    | 1999       | 2018                                      | 10%                        | None                 |            |   |                            |             |

\*Despite the 1994 consumption ban, a small percentage of new halon systems are assumed to continue to be built and filled with stockpiled or recovered supplies.

### Foam Blowing

ODSs, HFCs, and other chemicals are used to produce foams, including such items as the foam insulation panels around refrigerators, insulation sprayed on buildings, etc. The chemical is used to create pockets of gas within a substrate, increasing the insulating properties of the item. Foams are given emission profiles depending on the foam type (open cell or closed cell). Open cell foams are assumed to be 100 percent emissive in the year of manufacture. Closed cell foams are assumed to emit a portion of their total HFC content upon manufacture, a portion at a constant rate over the lifetime of the foam, a portion at disposal, and a portion after disposal; these portions vary by end-use.

#### Step 1: Calculate manufacturing emissions (open-cell and closed-cell foams)

Manufacturing emissions occur in the year of foam manufacture, and are calculated as presented in the following equation.

$$Em_j = lm \times Qc_j$$

where:

$Em_j$  = Emissions from manufacturing. Total emissions of a specific chemical in year  $j$  due to manufacturing losses, by weight.

$lm$  = Loss Rate. Percent of original blowing agent emitted during foam manufacture. For open-cell foams,  $lm$  is 100%.

$Qc$  = Quantity of Chemical. Total amount of a specific chemical used to manufacture closed-cell foams in a given year.

$j$  = Year of emission.

### Step 2: Calculate lifetime emissions (closed-cell foams)

Lifetime emissions occur annually from closed-cell foams throughout the lifetime of the foam, as calculated as presented in the following equation.

$$Eu_j = lu \times \sum Qc_{j+i} \text{ for } i=1 \rightarrow k$$

where:

- $Eu_j$  = Emissions from Lifetime Losses. Total emissions of a specific chemical in year j due to lifetime losses during use, by weight.
- $lu$  = Leak Rate. Percent of original blowing agent emitted each year during lifetime use.
- $Qc$  = Quantity of Chemical. Total amount of a specific chemical used to manufacture closed-cell foams in a given year.
- $i$  = Counter, runs from 1 to lifetime (k).
- $j$  = Year of emission.
- $k$  = Lifetime. The average lifetime of foam product.

### Step 3: Calculate disposal emissions (closed-cell foams)

Disposal emissions occur in the year the foam is disposed, and are calculated as presented in the following equation.

$$Ed_j = ld \times Qc_{j-k}$$

where:

- $Ed_j$  = Emissions from disposal. Total emissions of a specific chemical in year j at disposal, by weight.
- $ld$  = Loss Rate. Percent of original blowing agent emitted at disposal.
- $Qc$  = Quantity of Chemical. Total amount of a specific chemical used to manufacture closed-cell foams in a given year.
- $j$  = Year of emission.
- $k$  = Lifetime. The average lifetime of foam product.

### Step 4: Calculate post-disposal emissions (closed-cell foams)

Post-disposal emissions occur in the years after the foam is disposed; for example, emissions might occur while the disposed foam is in a landfill. Currently, the only foam type assumed to have post-disposal emissions is polyurethane foam used as domestic refrigerator and freezer insulation, which is expected to continue to emit for 26 years post-disposal, calculated as presented in the following equation.

$$Ep_j = lp \times \sum Qc_{j-m} \text{ for } m=k \rightarrow k + 26$$

where:

- $Ep_j$  = Emissions from post disposal. Total post-disposal emissions of a specific chemical in year j, by weight.
- $lp$  = Leak Rate. Percent of original blowing agent emitted post disposal.
- $Qc$  = Quantity of Chemical. Total amount of a specific chemical used to manufacture closed-cell foams in a given year.
- $k$  = Lifetime. The average lifetime of foam product.
- $m$  = Counter. Runs from lifetime (k) to (k+26).
- $j$  = Year of emission.

**Step 5: Calculate total emissions (open-cell and closed-cell foams)**

To calculate total emissions from foams in any given year, emissions from all foam stages must be summed, as presented in the following equation.

$$E_j = Em_j + Eu_j + Ed_j + Ep_j$$

where:

$E_j$  = Total Emissions. Total emissions of a specific chemical in year  $j$ , by weight.

$Em$  = Emissions from manufacturing. Total emissions of a specific chemical in year  $j$  due to manufacturing losses, by weight.

$Eu_j$  = Emissions from Lifetime Losses. Total emissions of a specific chemical in year  $j$  due to lifetime losses during use, by weight.

$Ed_j$  = Emissions from disposal. Total emissions of a specific chemical in year  $j$  at disposal, by weight.

$Ep_j$  = Emissions from post disposal. Total post-disposal emissions of a specific chemical in year  $j$ , by weight.

**Assumptions**

The Vintaging Model contains thirteen foam types, whose transition assumptions away from ODS and growth rates are presented in Table A-174. The emission profiles of these thirteen foam types are shown in Table A-175.

**Table A-174: Foam Blowing Market Transition Assumptions**

| Initial Market Segment  | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|---|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|   | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| <b>Commercial Refrigeration Foam</b>                          |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11  | HCFC-141b          | 1989       | 1996                                      | 40%                        | HFC-245fa            | 2002       | 2003                                      | 80%                        | None                |            |   |                            | 6.0%        |
|   |                    |            |   |                            | Non-ODP/GWP          | 2002       | 2003                                      | 20%                        | None                |            |   |                            |             |
|   | HCFC-142b          | 1989       | 1996                                      | 8%                         | Non-ODP/GWP          | 2009       | 2010                                      | 80%                        | None                |            |   |                            |             |
|   | HCFC-22            | 1989       | 1996                                      | 52%                        | HFC-245fa            | 2009       | 2010                                      | 20%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | Non-ODP/GWP          | 2009       | 2010                                      | 80%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | HFC-245fa            | 2009       | 2010                                      | 20%                        | None                |            |   |                            |             |
| <b>Flexible PU Foam: Integral Skin Foam</b>                   |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11  | HCFC-141b          | 1989       | 1990                                      | 100%                       | HFC-134a             | 1993       | 1996                                      | 25%                        | None                |            |   |                            | 2.0%        |
|   |                    |            |   |                            | HFC-134a             | 1994       | 1996                                      | 25%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | CO <sub>2</sub>      | 1993       | 1996                                      | 25%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | CO <sub>2</sub>      | 1994       | 1996                                      | 25%                        | None                |            |   |                            |             |
| <b>Flexible PU Foam: Slabstock Foam, Moulded Foam</b>         |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11  | Non-ODP/GWP        | 1992       | 1992                                      | 100%                       | None                 |            |   |                            |                     |            |   |                            | 2.0%        |
| <b>Phenolic Foam</b>  |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11  | HCFC-141b          | 1989       | 1990                                      | 100%                       | Non-ODP/GWP          | 1992       | 1992                                      | 100%                       | None                |            |   |                            | 2.0%        |
| <b>Polyolefin Foam</b>  |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-114   | HFC-152a           | 1989       | 1993                                      | 10%                        | Non-ODP/GWP          | 2005       | 2010                                      | 100%                       | None                |            |   |                            | 2.0%        |
|   | HCFC-142b          | 1989       | 1993                                      | 90%                        | Non-ODP/GWP          | 1994       | 1996                                      | 100%                       | None                |            |   |                            |             |
| <b>PU and PIR Rigid: Boardstock</b>                           |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11  | HCFC-141b          | 1993       | 1996                                      | 100%                       | Non-ODP/GWP          | 2000       | 2003                                      | 95%                        | None                |            |   |                            | 6.0%        |
|   |                    |            |   |                            | HC/HFC-245fa Blend   | 2000       | 2003                                      | 5%                         | None                |            |   |                            |             |
| <b>PU Rigid: Domestic Refrigerator and Freezer Insulation</b> |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-11  | HCFC-141b          | 1993       | 1995                                      | 100%                       | HFC-134a             | 1996       | 2001                                      | 7%                         | Non-ODP/GWP         | 2002       | 2003                                      | 100%                       | 0.8%        |
|   |                    |            |   |                            | HFC-245fa            | 2001       | 2003                                      | 50%                        | Non-ODP/GWP         | 2015       | 2029                                      | 100%                       |             |
|   |                    |            |   |                            | HFC-245fa            | 2006       | 2009                                      | 10%                        | Non-ODP/GWP         | 2015       | 2029                                      | 100%                       |             |
|   |                    |            |   |                            | Non-ODP/GWP          | 2002       | 2005                                      | 10%                        | None                |            |   |                            |             |
|   |                    |            |   |                            | Non-ODP/GWP          | 2006       | 2009                                      | 3%                         | None                |            |   |                            |             |
|   |                    |            |   |                            | Non-ODP/GWP          | 2009       | 2014                                      | 20%                        | None                |            |   |                            |             |
| <b>PU Rigid: One Component Foam</b>                           |                    |            |   |                            |                      |            |   |                            |                     |            |   |                            |             |
| CFC-12  | HCFC-142b/22 Blend | 1989       | 1996                                      | 70%                        | Non-ODP/GWP          | 2009       | 2010                                      | 80%                        | None                |            |   |                            | 4.0%        |
|   |                    |            |   |                            | HFC-134a             | 2009       | 2010                                      | 10%                        | None                |            |   |                            |             |

| Initial Market Segment   | Primary Substitute |            |   |                            | Secondary Substitute   |                              |   |                            | Tertiary Substitute  |              |   |                            | Growth Rate |
|--|--------------------|------------|---|----------------------------|--|------------------------------|---|----------------------------|--|--------------|---|----------------------------|-------------|
|  | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date                   | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date   | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
|  | HCFC-22            | 1989       | 1996                                      | 30%                        | HFC-152a<br>Non-ODP/GWP<br>HFC-134a<br>HFC-152a  | 2009<br>2009<br>2009<br>2009 | 2010<br>2010<br>2010<br>2010              | 10%<br>80%<br>10%<br>10%   | None<br>None<br>None<br>None   |              |   |                            |             |
| <b>PU Rigid: Other: Slabstock Foam</b>                         |                    |            |   |                            |  |                              |   |                            |  |              |   |                            |             |
| CFC-11   | HCFC-141b          | 1989       | 1996                                      | 100%                       | CO <sub>2</sub><br>Non-ODP/GWP<br>HCFC-22  | 1999<br>2001<br>2003         | 2003<br>2003<br>2003                      | 45%<br>45%<br>10%          | None<br>None<br>Non-ODP/GWP  | 2009         | 2010                                      | 100%                       | 2.0%        |
| <b>PU Rigid: Sandwich Panels: Continuous and Discontinuous</b> |                    |            |   |                            |  |                              |   |                            |  |              |   |                            |             |
| CFC-11   | HCFC-141b          | 1989       | 1996                                      | 82%                        | HCFC-22/Water Blend<br>HFC-245fa/CO <sub>2</sub> Blend<br>Non-ODP/GWP<br>HFC-134a<br>HFC-245fa/CO <sub>2</sub> Blend | 2001<br>2002<br>2001<br>2002 | 2003<br>2004<br>2004<br>2004              | 20%<br>20%<br>40%<br>20%   | HFC-245fa/CO <sub>2</sub> Blend<br>Non-ODP/GWP<br>None<br>None<br>None | 2009<br>2009 | 2010<br>2010                              | 50%<br>50%                 | 6.0%        |
|  | HCFC-22            | 1989       | 1996                                      | 18%                        | Non-ODP/GWP<br>CO <sub>2</sub><br>HFC-134a   | 2009<br>2009<br>2009         | 2010<br>2010<br>2010                      | 40%<br>20%<br>20%<br>20%   | None<br>None<br>None<br>None   |              |   |                            |             |
| <b>PU Rigid: Spray Foam</b>                                    |                    |            |   |                            |  |                              |   |                            |  |              |   |                            |             |
| CFC-11   | HCFC-141b          | 1989       | 1996                                      | 100%                       | HFC-245fa<br>HFC-245fa/CO <sub>2</sub> Blend<br>Non-ODP/GWP  | 2002<br>2002<br>2001         | 2003<br>2003<br>2003                      | 30%<br>60%<br>10%          | None<br>None<br>None   |              |   |                            | 6.0%        |
| <b>XPS: Boardstock Foam</b>                                    |                    |            |   |                            |  |                              |   |                            |  |              |   |                            |             |
| CFC-12   | HCFC-142b/22 Blend | 1989       | 1994                                      | 10%                        | HFC-134a<br>HFC-152a<br>CO <sub>2</sub><br>Non-ODP/GWP   | 2009<br>2009<br>2009<br>2009 | 2010<br>2010<br>2010<br>2010              | 70%<br>10%<br>10%<br>10%   | None<br>None<br>None<br>None   |              |   |                            | 2.5%        |
|  | HCFC-142b          | 1989       | 1994                                      | 90%                        | HFC-134a<br>HFC-152a<br>CO <sub>2</sub><br>Non-ODP/GWP   | 2009<br>2009<br>2009<br>2009 | 2010<br>2010<br>2010<br>2010              | 70%<br>10%<br>10%<br>10%   | None<br>None<br>None<br>None   |              |   |                            |             |
| <b>XPS: Sheet Foam</b>   |                    |            |   |                            |  |                              |   |                            |  |              |   |                            |             |
| CFC-12   | CO <sub>2</sub>    | 1989       | 1994                                      | 1%                         | None   |                              |   |                            |  |              |   |                            | 2.0%        |

| Initial Market Segment | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|------------------------|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|                        | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
|                        | Non-ODP/GWP        | 1989       | 1994                                      | 99%                        | CO <sub>2</sub>      | 1995       | 1999                                      | 9%                         | None                |            |   |                            |             |
|                        |                    |            |   |                            | HFC-152a             | 1995       | 1999                                      | 10%                        | None                |            |   |                            |             |

**Table A-175: Emission Profile for the Foam End-Uses**

| Foam End-Use  | Loss at Manufacturing (%) | Annual Leakage Rate (%) | Leakage Lifetime (years) | Loss at Disposal (%) | Total <sup>a</sup> (%) |
|---|---------------------------|-------------------------|--------------------------|----------------------|------------------------|
| Flexible PU Foam: Slabstock Foam, Moulded Foam                      | 100                       | 0                       | 1                        | 0                    | 100                    |
| Commercial Refrigeration  | 6                         | 0.25                    | 15                       | 90.25                | 100                    |
| Rigid PU: Spray Foam  | 15                        | 1.5                     | 56                       | 1                    | 100                    |
| Rigid PU: Slabstock and Other                                       | 37.5                      | 0.75                    | 15                       | 51.25                | 100                    |
| Phenolic Foam   | 23                        | 0.875                   | 32                       | 49                   | 100                    |
| Polyolefin Foam   | 95                        | 2.5                     | 2                        | 0                    | 100                    |
| Rigid PU: One Component Foam  | 100                       | 0                       | 1                        | 0                    | 100                    |
| XPS: Sheet Foam <sup>a</sup>  | 40                        | 2                       | 25                       | 0                    | 90                     |
| XPS: Boardstock Foam  | 25                        | 0.75                    | 50                       | 37.5                 | 100                    |
| Flexible PU Foam: Integral Skin Foam                                | 95                        | 2.5                     | 2                        | 0                    | 100                    |
| Rigid PU: Domestic Refrigerator and Freezer Insulation <sup>a</sup> | 4                         | 0.25                    | 14                       | 40.0                 | 47.5                   |
| PU and PIR Rigid: Boardstock  | 6                         | 1                       | 50                       | 44                   | 100                    |
| PU Sandwich Panels: Continuous and Discontinuous                    | 5.5                       | 0.5                     | 50                       | 69.5                 | 100                    |

PIR (Polyisocyanurate)

PU (Polyurethane)

XPS (Extruded Polystyrene)

<sup>a</sup> In general, total emissions from foam end-uses are assumed to be 100 percent, although work is underway to investigate that assumption. In the XPS Sheet/Insulation Board end-use, the source of emission rates and lifetimes did not yield 100 percent emission; it is unclear at this time whether that was intentional. In the Rigid PU Appliance Foam end-use, the source of emission rates and lifetimes did not yield 100 percent emission; the remainder is anticipated to be emitted at a rate of 2.0%/year post-disposal for the next 26 years.

### Sterilization

Sterilants kill microorganisms on medical equipment and devices. The principal ODS used in this sector was a blend of 12 percent ethylene oxide (EtO) and 88 percent CFC-12, known as "12/88." In that blend, ethylene oxide sterilizes the equipment and CFC-12 is a diluent solvent to form a non-flammable blend. The sterilization sector is modeled as a single end-use. For sterilization applications, all chemicals that are used in the equipment in any given year are assumed to be emitted in that year, as shown in the following equation.

$$E_j = Qc_j$$

where:

$E$  = Emissions. Total emissions of a specific chemical in year  $j$  from use in sterilization equipment, by weight.

$Qc$  = Quantity of Chemical. Total quantity of a specific chemical used in sterilization equipment in year  $j$ , by weight.

$j$  = Year of emission.

### Assumptions

The Vintaging Model contains one sterilization end-use, whose transition assumptions away from ODS and growth rates are presented in Table A-176.

**Table A-176: Sterilization Market Transition Assumptions**

| Initial Market Segment | Primary Substitute |            |   |                            | Secondary Substitute |            |   |                            | Tertiary Substitute |            |   |                            | Growth Rate |
|------------------------|--------------------|------------|---|----------------------------|----------------------|------------|---|----------------------------|---------------------|------------|---|----------------------------|-------------|
|                        | Name of Substitute | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute   | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration | Name of Substitute  | Start Date | Date of Full Penetration in New Equipment | Maximum Market Penetration |             |
| 12/88                  | EtO                | 1994       | 1995                                      | 95%                        | None                 |            |   |                            |                     |            |   |                            | 2.0%        |
|                        | Non-ODP/GWP        | 1994       | 1995                                      | 1%                         | None                 |            |   |                            |                     |            |   |                            |             |
|                        | HCFC/EtO Blends    | 1993       | 1994                                      | 4%                         | Non-ODP/GWP          | 2010       | 2010                                      | 100%                       | None                |            |   |                            |             |

## Model Output

By repeating these calculations for each year, the Vintaging Model creates annual profiles of use and emissions for ODS and ODS substitutes. The results can be shown for each year in two ways: 1) on a chemical-by-chemical basis, summed across the end-uses, or 2) on an end-use or sector basis. Values for use and emissions are calculated both in metric tons and in million metric tons of CO<sub>2</sub> equivalents (MMT CO<sub>2</sub> Eq.). The conversion of metric tons of chemical to MMT CO<sub>2</sub> Eq. is accomplished through a linear scaling of tonnage by the global warming potential (GWP) of each chemical.

Throughout its development, the Vintaging Model has undergone annual modifications. As new or more accurate information becomes available, the model is adjusted in such a way that both past and future emission estimates are often altered.

### Bank of ODS and ODS Substitutes

The bank of an ODS or an ODS substitute is “the cumulative difference between the chemical that has been consumed in an application or sub-application and that which has already been released” (IPCC 2006). For any given year, the bank is equal to the previous year’s bank, less the chemical in equipment disposed of during the year, plus chemical in new equipment entering the market during that year, less the amount emitted but not replaced, plus the amount added to replace chemical emitted prior to the given year, as shown in the following equation:

$$BC_j = BC_{j-1} - Qd_j + Qp_j + E_e - Q_r$$

where:

|        |   |  |
|--------|---|--|
| $BC_j$ | = | Bank of Chemical. Total bank of a specific chemical in year $j$ , by weight.   |
| $Qd_j$ | = | Quantity of Chemical in Equipment Disposed. Total quantity of a specific chemical in equipment disposed of in year $j$ , by weight.  |
| $Qp_j$ | = | Quantity of Chemical Penetrating the Market. Total quantity of a specific chemical that is entering the market in year $j$ , by weight.  |
| $E_e$  | = | Emissions of Chemical Not Replaced. Total quantity of a specific chemical that is emitted during year $j$ but is not replaced in that year. The Vintaging Model assumes all chemical emitted from refrigeration, air conditioning and fire extinguishing equipment is replaced in the year it is emitted, hence this term is zero for all sectors except foam blowing. |
| $Q_r$  | = | Chemical Replacing Previous Year’s Emissions. Total quantity of a specific chemical that is used to replace emissions that occurred prior to year $j$ . The Vintaging Model assumes all chemical emitted from refrigeration, air conditioning and fire extinguishing equipment is replaced in the year it is emitted, hence this term is zero for all sectors.         |
| $j$    | = | Year of emission.  |

Table A-177 provides the bank for ODS and ODS substitutes by chemical grouping in metric tons (MT) for 1990 to 2014.

**Table A-177. Banks of ODS and ODS Substitutes, 1990-2014 (MT)**

| Year | CFC     | HCFC      | HFC     |
|------|---------|-----------|---------|
| 1990 | 699,504 | 281,676   | 872     |
| 1995 | 792,162 | 495,808   | 48,449  |
| 2000 | 667,205 | 921,390   | 181,369 |
| 2001 | 639,001 | 990,934   | 209,791 |
| 2002 | 614,855 | 1,044,303 | 238,162 |
| 2003 | 590,932 | 1,081,093 | 272,302 |
| 2004 | 566,576 | 1,118,737 | 307,974 |
| 2005 | 543,858 | 1,159,977 | 345,872 |
| 2006 | 520,759 | 1,197,172 | 389,652 |
| 2007 | 500,425 | 1,225,447 | 434,729 |
| 2008 | 485,443 | 1,242,579 | 476,428 |
| 2009 | 478,491 | 1,235,393 | 522,965 |
| 2010 | 464,946 | 1,199,995 | 588,726 |
| 2011 | 451,334 | 1,155,064 | 653,741 |
| 2012 | 437,294 | 1,109,688 | 725,821 |
| 2013 | 423,535 | 1,060,341 | 799,683 |
| 2014 | 411,433 | 1,009,123 | 873,569 |

## References

IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.

### 3.10. Methodology for Estimating CH<sub>4</sub> Emissions from Enteric Fermentation

Methane emissions from enteric fermentation were estimated for seven livestock categories: cattle, horses, sheep, swine, goats, American bison, and the non-horse equines (mules and asses). Emissions from cattle represent the majority of U.S. emissions from enteric fermentation; consequently, a more detailed IPCC Tier 2 methodology was used to estimate emissions from cattle. The IPCC Tier 1 methodology was used to estimate emissions for the other types of livestock, including horses, goats, sheep, swine, American bison, and mules and asses.

#### Estimate Methane Emissions from Cattle

This section describes the process used to estimate CH<sub>4</sub> emissions from enteric fermentation from cattle using the Cattle Enteric Fermentation Model (CEFM). The CEFM was developed based on recommendations provided in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006) and uses information on population, energy requirements, digestible energy, and CH<sub>4</sub> conversion rates to estimate CH<sub>4</sub> emissions.<sup>73</sup> The emission methodology consists of the following three steps: (1) characterize the cattle population to account for animal population categories with different emission profiles; (2) characterize cattle diets to generate information needed to estimate emission factors; and (3) estimate emissions using these data and the IPCC Tier 2 equations.

#### Step 1: Characterize U.S. Cattle Population

The state-level cattle population estimates are based on data obtained from the U.S. Department of Agriculture's (USDA) National Agricultural Statistics Service Quick Stats database (USDA 2015). A summary of the annual average populations upon which all livestock-related emissions are based is provided in Table A-178. Cattle populations used in the Enteric Fermentation source category were estimated using the cattle transition matrix in the CEFM, which uses January 1 USDA population estimates and weight data to simulate the population of U.S. cattle from birth to slaughter, and results in an estimate of the number of animals in a particular cattle grouping while taking into account the monthly rate of weight gain, the average weight of the animals, and the death and calving rates. The use of supplemental USDA data and the cattle transition matrix in the CEFM results in cattle population estimates for this sector differing slightly from the January 1 or July 1 USDA point estimates and the cattle population data obtained from the Food and Agriculture Organization of the United Nations (FAO).

**Table A-178: Cattle Population Estimates from the CEFM Transition Matrix for 1990–2014 (1,000 head)**

| Livestock Type                       | 1990   | 1995   | 2000   | 2005   | 2010   | 2011   | 2012   | 2013   | 2014   |
|--------------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| <b>Dairy</b>                         |        |        |        |        |        |        |        |        |        |
| Dairy Calves (0–6 months)            | 5,369  | 5,091  | 4,951  | 4,628  | 4,666  | 4,709  | 4,770  | 4,758  | 4,727  |
| Dairy Cows                           | 10,015 | 9,482  | 9,183  | 9,004  | 9,087  | 9,156  | 9,236  | 9,221  | 9,208  |
| Dairy Replacements<br>(7–11 months)  | 1,214  | 1,216  | 1,196  | 1,257  | 1,351  | 1,362  | 1,348  | 1,341  | 1,356  |
| Dairy Replacements<br>(12–23 months) | 2,915  | 2,892  | 2,812  | 2,905  | 3,194  | 3,215  | 3,233  | 3,185  | 3,190  |
| <b>Beef</b>                          |        |        |        |        |        |        |        |        |        |
| Beef Calves (0–6 months)             | 16,909 | 18,177 | 17,431 | 16,918 | 16,067 | 15,817 | 15,288 | 14,859 | 14,946 |
| Bulls                                | 2,160  | 2,385  | 2,293  | 2,214  | 2,190  | 2,165  | 2,100  | 2,074  | 2,038  |
| Beef Cows                            | 32,455 | 35,190 | 33,575 | 32,674 | 31,440 | 30,913 | 30,282 | 29,631 | 29,085 |
| Beef Replacements<br>(7–11 months)   | 1,269  | 1,493  | 1,313  | 1,363  | 1,238  | 1,232  | 1,263  | 1,291  | 1,342  |
| Beef Replacements<br>(12–23 months)  | 2,967  | 3,637  | 3,097  | 3,171  | 3,050  | 2,889  | 2,968  | 3,041  | 3,113  |
| Steer Stockers                       | 10,321 | 11,716 | 8,724  | 8,185  | 8,234  | 7,568  | 7,173  | 7,457  | 7,411  |
| Heifer Stockers                      | 5,946  | 6,699  | 5,371  | 5,015  | 5,061  | 4,752  | 4,456  | 4,455  | 4,384  |
| Feedlot Cattle                       | 9,549  | 11,064 | 13,006 | 12,652 | 13,204 | 13,601 | 13,328 | 13,267 | 13,222 |

<sup>73</sup> Additional information on the Cattle Enteric Fermentation Model can be found in ICF (2006).

The population transition matrix in the CEFM simulates the U.S. cattle population over time and provides an estimate of the population age and weight structure by cattle type on a monthly basis.<sup>74</sup> Since cattle often do not remain in a single population type for an entire year (e.g., calves become stockers, stockers become feedlot animals), and emission profiles vary both between and within each cattle type, these monthly age groups are tracked in the enteric fermentation model to obtain more accurate emission estimates than would be available from annual point estimates of population (such as available from USDA statistics) and weight for each cattle type.

The transition matrix tracks both dairy and beef populations, and divides the populations into males and females, and subdivides the population further into specific cattle groupings for calves, replacements, stockers, feedlot, and mature animals. The matrix is based primarily on two types of data: population statistics and weight statistics (including target weights, slaughter weights, and weight gain). Using the weight data, the transition matrix simulates the growth of animals over time by month. The matrix also relies on supplementary data, such as feedlot placement statistics, slaughter statistics, death rates, and calving rates.

The basic method for tracking population of animals per category is based on the number of births (or graduates) into the monthly age group minus those animals that die or are slaughtered and those that graduate to the next category (such as stockers to feedlot placements).

Each stage in the cattle lifecycle was modeled to simulate the cattle population from birth to slaughter. This level of detail accounts for the variability in CH<sub>4</sub> emissions associated with each life stage. Given that a stage can last less than one year (e.g., calves are usually weaned between 4 and 6 months of age), each is modeled on a per-month basis. The type of cattle also influences CH<sub>4</sub> emissions (e.g., beef versus dairy). Consequently, there is an independent transition matrix for each of three separate lifecycle phases, 1) calves, 2) replacements and stockers, and 3) feedlot animals. In addition, the number of mature cows and bulls are tabulated for both dairy and beef stock. The transition matrix estimates total monthly populations for all cattle subtypes. These populations are then reallocated to the state level based on the percent of the cattle type reported in each state in the January 1 USDA data. Each lifecycle is discussed separately below, and the categories tracked are listed in Table A-179.

**Table A-179: Cattle Population Categories Used for Estimating CH<sub>4</sub> Emissions**

| Dairy Cattle        | Beef Cattle                           |
|---------------------|---------------------------------------|
| Calves              | Calves                                |
| Heifer Replacements | Heifer Replacements                   |
| Cows                | Heifer and Steer Stockers             |
|                     | Animals in Feedlots (Heifers & Steer) |
|                     | Cows                                  |
|                     | Bulls <sup>a</sup>                    |

<sup>a</sup>Bulls (beef and dairy) are accounted for in a single category.

The key variables tracked for each of these cattle population categories are as follows:

**Calves.** Although enteric emissions are only calculated for 4- to 6-month old calves, it is necessary to calculate populations from birth as emissions from manure management require total calf populations and the estimates of populations for older cattle rely on the available supply of calves from birth. The number of animals born on a monthly basis was used to initiate monthly cohorts and to determine population age structure. The number of calves born each month was obtained by multiplying annual births by the percentage of births per month. Annual birth information for each year was taken from USDA (2015). For dairy cows, the number of births is assumed to be distributed equally throughout the year (approximately 8.3 percent per month) while beef births are distributed according to Table A-180, based on approximations from the National Animal Health Monitoring System (NAHMS) (USDA/APHIS/VS 1998, 1994, 1993). To determine whether calves were born to dairy or beef cows, the dairy cow calving rate (USDA/APHIS/VS 2002, USDA/APHIS/VS 1996) was multiplied by the total dairy cow population to determine the number of births attributable to dairy cows, with the remainder assumed to be attributable to beef cows. Total annual calf births are obtained from USDA, and distributed into monthly cohorts by cattle type (beef or dairy). Calf growth is modeled by month, based on estimated monthly weight gain for each cohort (approximately 61 pounds per month). The total calf population is modified through time to account for veal calf slaughter at 4 months and a calf death loss of 0.35 percent annually (distributed across age cohorts up to 6 months of age). An example of a transition matrix for calves is shown in Table A-181. Note that 1- to 6-month old calves in January of each year have been tracked through the model based on births and death loss from the previous year.

<sup>74</sup> Mature animal populations are not assumed to have significant monthly fluctuations, and therefore the populations utilized are the January estimates downloaded from USDA (2015).

**Table A-180: Estimated Beef Cow Births by Month**

| Jan | Feb | Mar | Apr | May | Jun | Jul | Aug | Sep | Oct | Nov | Dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 7%  | 15% | 28% | 22% | 9%  | 3%  | 2%  | 2%  | 3%  | 4%  | 3%  | 3%  |

**Table A-181: Example of Monthly Average Populations from Calf Transition Matrix (1,000 head)**

| Age (month) | Jan   | Feb   | Mar   | Apr   | May   | Jun   | Jul   | Aug   | Sep   | Oct   | Nov   | Dec   |
|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 6           | 1,138 | 1,131 | 1,389 | 1,612 | 1,554 | 1,538 | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 |
| 5           | 1,131 | 1,389 | 1,612 | 1,554 | 1,538 | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 | 1,153 |
| 4           | 1,389 | 1,612 | 1,554 | 1,538 | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 | 1,153 | 1,144 |
| 3           | 1,612 | 1,554 | 1,538 | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 | 1,153 | 1,144 | 1,402 |
| 2           | 1,554 | 1,538 | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 | 1,153 | 1,144 | 1,402 | 1,625 |
| 1           | 1,538 | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 | 1,153 | 1,144 | 1,402 | 1,625 | 1,565 |
| 0           | 2,431 | 4,488 | 7,755 | 6,298 | 2,971 | 1,522 | 1,153 | 1,144 | 1,402 | 1,625 | 1,565 | 1,547 |

Note: The cohort starting at age 0 months on January 1 is tracked in order to illustrate how a single cohort moves through the transition matrix. Each month, the cohort reflects the decreases in population due to the estimated 0.35 percent annual death loss, and between months 4 and 5, a more significant loss is seen than in other months due to estimated veal slaughter.

**Replacements and Stockers.** At 7 months of age, calves “graduate” and are separated into the applicable cattle types. First the number of replacements required for beef and dairy cattle are calculated based on estimated death losses and population changes between beginning and end of year population estimates. Based on the USDA estimates for “replacement beef heifers” and “replacement dairy heifers,” the transition matrix for the replacements is back-calculated from the known animal totals from USDA, and the number of calves needed to fill that requirement for each month is subtracted from the known supply of female calves. All female calves remaining after those needed for beef and dairy replacements are removed and become “stockers” that can be placed in feedlots (along with all male calves). During the stocker phase animals are subtracted out of the transition matrix for placement into feedlots based on feedlot placement statistics from USDA (2015).

The data and calculations that occur for the stocker category include matrices that estimate the population of backgrounding heifers and steer, as well as a matrix for total combined stockers. The matrices start with the beginning of year populations in January and model the progression of each cohort. The age structure of the January population is based on estimated births by month from the previous two years, although in order to balance the population properly, an adjustment is added that slightly reduces population percentages in the older populations. The populations are modified through addition of graduating calves (added in month 7, bottom row of Table A-182) and subtraction through death loss and animals placed in feedlots. Eventually, an entire cohort population of stockers may reach zero, indicating that the complete cohort has been transitioned into feedlots. An example of the transition matrix for stockers is shown in Table A-182.

**Table A-182: Example of Monthly Average Populations from Stocker Transition Matrix (1,000 head)**

| Age (month) | Jan   | Feb   | Mar   | Apr   | May   | Jun   | Jul   | Aug   | Sep   | Oct   | Nov   | Dec   |
|-------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 23          | 185   | 180   | 104   | 37    | 15    | 9     | 8     | 8     | 6     | 3     | 1     | 0     |
| 22          | 320   | 146   | 49    | 19    | 12    | 9     | 9     | 9     | 6     | 3     | 17    | 181   |
| 21          | 260   | 69    | 25    | 14    | 11    | 11    | 11    | 8     | 6     | 68    | 218   | 313   |
| 20          | 123   | 35    | 19    | 14    | 14    | 13    | 10    | 8     | 133   | 331   | 387   | 254   |
| 19          | 63    | 27    | 19    | 17    | 16    | 13    | 10    | 196   | 472   | 615   | 318   | 120   |
| 18          | 48    | 27    | 23    | 20    | 16    | 13    | 241   | 610   | 900   | 514   | 149   | 61    |
| 17          | 47    | 33    | 27    | 19    | 15    | 295   | 709   | 1,179 | 759   | 237   | 129   | 47    |
| 16          | 58    | 38    | 26    | 19    | 363   | 828   | 1,380 | 1,000 | 348   | 340   | 47    | 46    |
| 15          | 67    | 36    | 25    | 452   | 977   | 1,619 | 1,172 | 456   | 603   | 47    | 46    | 57    |
| 14          | 65    | 36    | 599   | 1,172 | 1,921 | 1,378 | 534   | 862   | 47    | 46    | 57    | 66    |
| 13          | 64    | 845   | 1,478 | 2,309 | 1,639 | 629   | 1,117 | 47    | 46    | 57    | 66    | 63    |
| 12          | 982   | 1,602 | 2,556 | 1,858 | 755   | 1,512 | 214   | 46    | 57    | 66    | 63    | 63    |
| 11          | 1,814 | 2,770 | 2,056 | 855   | 1,872 | 277   | 138   | 76    | 89    | 81    | 80    | 1,016 |
| 10          | 3,133 | 2,255 | 945   | 2,241 | 385   | 189   | 184   | 231   | 209   | 185   | 1,135 | 2,445 |
| 9           | 2,545 | 1,062 | 2,502 | 484   | 335   | 341   | 420   | 372   | 371   | 1,292 | 2,786 | 5,299 |
| 8           | 1,200 | 2,951 | 664   | 482   | 557   | 759   | 658   | 649   | 1,503 | 3,247 | 5,984 | 4,877 |
| 7           | 3,381 | 800   | 794   | 956   | 1,160 | 1,109 | 1,100 | 1,876 | 3,666 | 6,504 | 5,243 | 2,353 |

Note: The cohort starting at age 7 months on January 1 is tracked in order to illustrate how a single cohort moves through the transition matrix. Each month, the cohort reflects the decreases in population due to the estimated 0.35 percent annual death loss and loss due to placement in feedlots (the latter resulting in the majority of the loss from the matrix).

In order to ensure a balanced population of both stockers and placements, additional data tables are utilized in the stocker matrix calculations. The tables summarize the placement data by weight class and month, and is based on the total number of animals within the population that are available to be placed in feedlots and the actual feedlot placement statistics provided by USDA (2015). In cases where there are discrepancies between the USDA estimated placements by weight class and the calculated animals available by weight, the model pulls available stockers from one higher weight category if available. If there are still not enough animals to fulfill requirements the model pulls animals from one lower weight category. In the current time series, this method was able to ensure that total placement data matched USDA estimates, and no shortfalls have occurred.

In addition, average weights were tracked for each monthly age group using starting weight and monthly weight gain estimates. Weight gain (i.e., pounds per month) was estimated based on weight gain needed to reach a set target weight, divided by the number of months remaining before target weight was achieved. Birth weight was assumed to be 88 pounds for both beef and dairy animals. Weaning weights were estimated at 515 pounds. Other reported target weights were available for 12-, 15-, 24-, and 36-month-old animals, depending on the animal type. Beef cow mature weight was taken from measurements provided by a major British Bos taurus breed (Enns 2008) and increased during the time series through 2007.<sup>75</sup> Bull mature weight was calculated as 1.5 times the beef cow mature weight (Doren et al. 1989). Beef replacement weight was calculated as 70 percent of mature weight at 15 months and 85 percent of mature weight at 24 months. As dairy weights are not a trait that is typically tracked, mature weight for dairy cows was estimated at 1,500 pounds for all years, based on a personal communication with Kris Johnson (2010) and an estimate from Holstein Association USA (2010).<sup>76</sup> Dairy replacement weight at 15 months was assumed to be 875 pounds and 1,300 pounds at 24 months. Live slaughter weights were estimated from dressed slaughter weight (USDA 2015) divided by 0.63. This ratio represents the dressed weight (i.e., weight of the carcass after removal of the internal organs), to the live weight (i.e., weight taken immediately before slaughter). The annual typical animal mass for each livestock type are presented in Table A-183.

Weight gain for stocker animals was based on monthly gain estimates from Johnson (1999) for 1989, and from average daily estimates from Lippke et al. (2000), Pinchack et al. (2004), Platter et al. (2003), and Skogerboe et al. (2000) for 2000. Interim years were calculated linearly, as shown in Table A-184, and weight gain was held constant starting in 2000. Table A-184 provides weight gains that vary by year in the CEFM.

<sup>75</sup> Mature beef weight is held constant after 2007 but future inventory submissions will incorporate known trends through 2007 and extrapolate to future years, as noted in the Planned Improvements section of 5.1 Enteric Fermentation.

<sup>76</sup> Mature dairy weight is based solely on Holstein weight, so could be higher than the national average. Future Inventory submissions will consider other dairy breeds, as noted in the Planned Improvements section of 5.1 Enteric Fermentation.

**Table A-183: Typical Animal Mass (lbs)**

| Year/Cattle Type | Calves | Dairy Cows <sup>a</sup> | Dairy Replacements <sup>b</sup> | Beef Cows <sup>a</sup> | Bulls <sup>a</sup> | Beef Replacements <sup>b</sup> | Steer Stockers <sup>b</sup> | Heifer Stockers <sup>b</sup> | Steer Feedlot <sup>b</sup> | Heifer Feedlot <sup>b</sup> |
|------------------|--------|-------------------------|---------------------------------|------------------------|--------------------|--------------------------------|-----------------------------|------------------------------|----------------------------|-----------------------------|
| 1990             | 269    | 1,500                   | 900                             | 1,221                  | 1,832              | 820                            | 692                         | 652                          | 923                        | 846                         |
| 1991             | 270    | 1,500                   | 898                             | 1,225                  | 1,838              | 822                            | 695                         | 656                          | 934                        | 856                         |
| 1992             | 269    | 1,500                   | 897                             | 1,263                  | 1,895              | 841                            | 714                         | 673                          | 984                        | 878                         |
| 1993             | 270    | 1,500                   | 899                             | 1,280                  | 1,920              | 852                            | 721                         | 683                          | 930                        | 864                         |
| 1994             | 270    | 1,500                   | 898                             | 1,280                  | 1,920              | 854                            | 721                         | 689                          | 944                        | 876                         |
| 1995             | 270    | 1,500                   | 898                             | 1,282                  | 1,923              | 858                            | 735                         | 701                          | 947                        | 880                         |
| 1996             | 269    | 1,500                   | 898                             | 1,285                  | 1,928              | 859                            | 739                         | 707                          | 940                        | 878                         |
| 1997             | 270    | 1,500                   | 900                             | 1,286                  | 1,929              | 861                            | 737                         | 708                          | 939                        | 877                         |
| 1998             | 270    | 1,500                   | 897                             | 1,296                  | 1,944              | 866                            | 736                         | 710                          | 957                        | 892                         |
| 1999             | 270    | 1,500                   | 899                             | 1,292                  | 1,938              | 862                            | 731                         | 709                          | 960                        | 895                         |
| 2000             | 270    | 1,500                   | 897                             | 1,272                  | 1,908              | 849                            | 720                         | 702                          | 961                        | 899                         |
| 2001             | 270    | 1,500                   | 898                             | 1,272                  | 1,908              | 850                            | 726                         | 707                          | 963                        | 901                         |
| 2002             | 270    | 1,500                   | 897                             | 1,276                  | 1,914              | 852                            | 726                         | 708                          | 982                        | 915                         |
| 2003             | 270    | 1,500                   | 900                             | 1,308                  | 1,962              | 872                            | 719                         | 702                          | 973                        | 905                         |
| 2004             | 270    | 1,500                   | 897                             | 1,323                  | 1,985              | 878                            | 719                         | 702                          | 967                        | 905                         |
| 2005             | 270    | 1,500                   | 895                             | 1,327                  | 1,991              | 880                            | 718                         | 706                          | 975                        | 917                         |
| 2006             | 270    | 1,500                   | 898                             | 1,341                  | 2,012              | 890                            | 725                         | 713                          | 984                        | 925                         |
| 2007             | 270    | 1,500                   | 897                             | 1,348                  | 2,022              | 895                            | 721                         | 707                          | 992                        | 928                         |
| 2008             | 270    | 1,500                   | 898                             | 1,348                  | 2,022              | 895                            | 721                         | 705                          | 1,000                      | 939                         |
| 2009             | 270    | 1,500                   | 896                             | 1,348                  | 2,022              | 895                            | 731                         | 715                          | 1,007                      | 948                         |
| 2010             | 270    | 1,500                   | 898                             | 1,348                  | 2,022              | 897                            | 727                         | 714                          | 997                        | 938                         |
| 2011             | 270    | 1,500                   | 898                             | 1,348                  | 2,022              | 892                            | 721                         | 713                          | 990                        | 932                         |
| 2012             | 270    | 1,500                   | 899                             | 1,348                  | 2,022              | 892                            | 714                         | 707                          | 1,004                      | 946                         |
| 2013             | 270    | 1,500                   | 899                             | 1,348                  | 2,022              | 893                            | 719                         | 710                          | 1,017                      | 958                         |
| 2014             | 270    | 1,500                   | 898                             | 1,348                  | 2,022              | 891                            | 722                         | 713                          | 1,023                      | 962                         |

<sup>a</sup> Input into the model.

<sup>b</sup> Annual average calculated in model based on age distribution.

**Table A-184: Weight Gains that Vary by Year (lbs)**

| Year/Cattle Type | Steer Stockers to 12 months (lbs/day) | Steer Stockers to 24 months (lbs/day) | Heifer Stockers to 12 months (lbs/day) | Heifer Stockers to 24 months (lbs/day) |
|------------------|---------------------------------------|---------------------------------------|--|--|
| 1990             | 1.53                                  | 1.23                                  | 1.23                                   | 1.08                                   |
| 1991             | 1.56                                  | 1.29                                  | 1.29                                   | 1.15                                   |
| 1992             | 1.59                                  | 1.35                                  | 1.35                                   | 1.23                                   |
| 1993             | 1.62                                  | 1.41                                  | 1.41                                   | 1.30                                   |
| 1994             | 1.65                                  | 1.47                                  | 1.47                                   | 1.38                                   |
| 1995             | 1.68                                  | 1.53                                  | 1.53                                   | 1.45                                   |
| 1996             | 1.71                                  | 1.59                                  | 1.59                                   | 1.53                                   |
| 1997             | 1.74                                  | 1.65                                  | 1.65                                   | 1.60                                   |
| 1998             | 1.77                                  | 1.71                                  | 1.71                                   | 1.68                                   |
| 1999             | 1.80                                  | 1.77                                  | 1.77                                   | 1.75                                   |
| 2000-onwards     | 1.83                                  | 1.83                                  | 1.83                                   | 1.83                                   |

Sources: Enns (2008), Johnson (1999), Lippke et al. (2000), NRC (1999), Pinchack et al. (2004), Platter et al. (2003), Skogerboe et al. (2000).

**Feedlot Animals.** Feedlot placement statistics from USDA provide data on the placement of animals from the stocker population into feedlots on a monthly basis by weight class. The model uses these data to shift a sufficient number of animals from the stocker cohorts into the feedlot populations to match the reported placement data. After animals are placed in feedlots they progress through two steps. First, animals spend 25 days on a step-up diet to become acclimated to the new feed type (e.g., more grain than forage, along with new dietary supplements), during this time weight gain is estimated to be 2.7 to 3 pounds per day (Johnson 1999). Animals are then switched to a finishing diet (concentrated, high energy) for a period of time before they are slaughtered. Weight gain during finishing diets is estimated to be 2.9 to 3.3 pounds per day (Johnson 1999). The length of time an animal spends in a feedlot depends on the start weight (i.e., placement weight), the rate of weight gain during the start-up and finishing phase of diet, and the target weight (as determined by

weights at slaughter). Additionally, animals remaining in feedlots at the end of the year are tracked for inclusion in the following year's emission and population counts. For 1990 to 1995, only the total placement data were available, therefore placements for each weight category (categories displayed in Table A-185) for those years are based on the average of monthly placements from the 1996 to 1998 reported figures. Placement data is available by weight class for all years from 1996 onward. Table A-185 provides a summary of the reported feedlot placement statistics for 2014.

**Table A-185: Feedlot Placements in the United States for 2014 (Number of animals placed/1,000 Head)**

| Weight Placed | When: | Jan          | Feb          | Mar          | Apr          | May          | Jun          | Jul          | Aug          | Sep          | Oct          | Nov          | Dec          |
|---------------|-------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| < 600 lbs     |       | 470          | 390          | 425          | 380          | 435          | 405          | 425          | 410          | 460          | 695          | 555          | 440          |
| 600 – 700 lbs |       | 435          | 330          | 290          | 250          | 290          | 245          | 260          | 285          | 350          | 570          | 435          | 365          |
| 700 – 800 lbs |       | 559          | 418          | 466          | 393          | 474          | 328          | 354          | 395          | 434          | 463          | 369          | 332          |
| > 800 lbs     |       | 550          | 520          | 620          | 600          | 710          | 490          | 520          | 635          | 770          | 640          | 435          | 400          |
| <b>Total</b>  |       | <b>2,014</b> | <b>1,658</b> | <b>1,801</b> | <b>1,623</b> | <b>1,909</b> | <b>1,468</b> | <b>1,559</b> | <b>1,725</b> | <b>2,014</b> | <b>2,368</b> | <b>1,794</b> | <b>1,537</b> |

Source: USDA (2015).

Note: Totals may not sum due to independent rounding.

**Mature Animals.** Energy requirements and hence, composition of diets, level of intake, and emissions for particular animals, are greatly influenced by whether the animal is pregnant or lactating. Information is therefore needed on the percentage of all mature animals that are pregnant each month, as well as milk production, to estimate CH<sub>4</sub> emissions. A weighted average percent of pregnant cows each month was estimated using information on births by month and average pregnancy term. For beef cattle, a weighted average total milk production per animal per month was estimated using information on typical lactation cycles and amounts (NRC 1999), and data on births by month. This process results in a range of weighted monthly lactation estimates expressed as pounds per animal per month. The monthly estimates for daily milk production by beef cows are shown in Table A-186. Annual estimates for dairy cows were taken from USDA milk production statistics. Dairy lactation estimates for 1990 through 2014 are shown in Table A-187. Beef and dairy cow and bull populations are assumed to remain relatively static throughout the year, as large fluctuations in population size are assumed to not occur. These estimates are taken from the USDA beginning and end of year population datasets.

**Table A-186: Estimates of Monthly Milk Production by Beef Cows (lbs/cow)**

|                                     | Jan | Feb | Mar | Apr  | May  | Jun  | Jul  | Aug | Sep | Oct | Nov | Dec |
|-------------------------------------|-----|-----|-----|------|------|------|------|-----|-----|-----|-----|-----|
| Beef Cow Milk Production (lbs/head) | 3.3 | 5.1 | 8.7 | 12.0 | 13.6 | 13.3 | 11.7 | 9.3 | 6.9 | 4.4 | 3.0 | 2.8 |

**Table A-187: Dairy Lactation Rates by State (lbs/ year/cow)**

| State/Year     | 1990   | 1995   | 2000   | 2005   | 2010   | 2011   | 2012   | 2013   | 2014   |
|----------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Alabama        | 12,214 | 14,176 | 13,920 | 14,000 | 14,182 | 14,300 | 13,000 | 13,000 | 13,625 |
| Alaska         | 13,300 | 17,000 | 14,500 | 12,273 | 11,833 | 13,800 | 14,250 | 10,667 | 11,667 |
| Arizona        | 17,500 | 19,735 | 21,820 | 22,679 | 23,452 | 23,473 | 23,979 | 23,626 | 24,347 |
| Arkansas       | 11,841 | 12,150 | 12,436 | 13,545 | 12,750 | 11,917 | 13,300 | 11,667 | 13,714 |
| California     | 18,456 | 19,573 | 21,130 | 21,404 | 23,025 | 23,438 | 23,457 | 23,178 | 23,785 |
| Colorado       | 17,182 | 18,687 | 21,618 | 22,577 | 23,664 | 23,430 | 24,158 | 24,292 | 24,951 |
| Connecticut    | 15,606 | 16,438 | 17,778 | 19,200 | 19,158 | 19,000 | 19,889 | 20,556 | 20,158 |
| Delaware       | 13,667 | 14,500 | 14,747 | 16,622 | 16,981 | 18,300 | 19,542 | 19,521 | 20,146 |
| Florida        | 14,033 | 14,698 | 15,688 | 16,591 | 18,711 | 19,067 | 19,024 | 19,374 | 20,382 |
| Georgia        | 12,973 | 15,550 | 16,284 | 17,259 | 17,658 | 18,354 | 19,138 | 19,600 | 20,790 |
| Hawaii         | 13,604 | 13,654 | 14,358 | 12,889 | 13,316 | 14,421 | 14,200 | 13,409 | 13,591 |
| Idaho          | 16,475 | 18,147 | 20,816 | 22,332 | 22,647 | 22,926 | 23,376 | 23,440 | 24,127 |
| Illinois       | 14,707 | 15,887 | 17,450 | 18,827 | 18,400 | 18,510 | 19,061 | 19,063 | 19,681 |
| Indiana        | 14,590 | 15,375 | 16,568 | 20,295 | 20,094 | 20,657 | 21,440 | 21,761 | 21,865 |
| Iowa           | 15,118 | 16,124 | 18,298 | 20,641 | 20,676 | 21,191 | 22,015 | 22,149 | 22,444 |
| Kansas         | 12,576 | 14,390 | 16,923 | 20,505 | 20,983 | 21,016 | 21,683 | 21,881 | 22,064 |
| Kentucky       | 10,947 | 12,469 | 12,841 | 12,896 | 14,769 | 14,342 | 15,135 | 15,070 | 15,905 |
| Louisiana      | 11,605 | 11,908 | 12,034 | 12,400 | 11,750 | 12,889 | 13,059 | 12,875 | 13,600 |
| Maine          | 14,619 | 16,025 | 17,128 | 18,030 | 18,344 | 18,688 | 18,576 | 19,548 | 20,000 |
| Maryland       | 13,461 | 14,725 | 16,083 | 16,099 | 18,537 | 18,654 | 19,196 | 19,440 | 19,740 |
| Massachusetts  | 14,871 | 16,000 | 17,091 | 17,059 | 17,286 | 16,923 | 18,250 | 17,692 | 17,923 |
| Michigan       | 15,394 | 17,071 | 19,017 | 21,635 | 23,277 | 23,164 | 23,976 | 24,116 | 24,638 |
| Minnesota      | 14,127 | 15,894 | 17,777 | 18,091 | 19,366 | 18,996 | 19,512 | 19,694 | 19,841 |
| Mississippi    | 12,081 | 12,909 | 15,028 | 15,280 | 13,118 | 14,571 | 14,214 | 13,286 | 14,462 |
| Missouri       | 13,632 | 14,158 | 14,662 | 16,026 | 14,596 | 14,611 | 14,979 | 14,663 | 15,539 |
| Montana        | 13,542 | 15,000 | 17,789 | 19,579 | 20,643 | 20,571 | 21,357 | 21,286 | 21,500 |
| Nebraska       | 13,866 | 14,797 | 16,513 | 17,950 | 19,797 | 20,579 | 21,179 | 21,574 | 22,130 |
| Nevada         | 16,400 | 18,128 | 19,000 | 21,680 | 23,714 | 22,966 | 22,931 | 22,034 | 23,793 |
| New Hampshire  | 15,100 | 16,300 | 17,333 | 18,875 | 19,600 | 20,429 | 19,643 | 20,923 | 20,143 |
| New Jersey     | 13,538 | 13,913 | 15,250 | 16,000 | 17,500 | 16,875 | 18,571 | 18,143 | 18,143 |
| New Mexico     | 18,815 | 18,969 | 20,944 | 21,192 | 24,551 | 24,854 | 24,694 | 24,944 | 25,093 |
| New York       | 14,658 | 16,501 | 17,378 | 18,639 | 20,807 | 21,046 | 21,623 | 22,070 | 22,330 |
| North Carolina | 15,220 | 16,314 | 16,746 | 18,741 | 19,682 | 20,089 | 20,435 | 20,326 | 20,891 |
| North Dakota   | 12,624 | 13,094 | 14,292 | 14,182 | 18,286 | 18,158 | 19,278 | 18,944 | 20,250 |
| Ohio           | 13,767 | 15,917 | 17,027 | 17,567 | 19,446 | 19,194 | 19,833 | 20,178 | 20,318 |
| Oklahoma       | 12,327 | 13,611 | 14,440 | 16,480 | 17,125 | 17,415 | 17,896 | 17,311 | 17,425 |
| Oregon         | 16,273 | 17,289 | 18,222 | 18,876 | 20,331 | 20,488 | 20,431 | 20,439 | 20,605 |
| Pennsylvania   | 14,726 | 16,492 | 18,081 | 18,722 | 19,847 | 19,495 | 19,549 | 19,797 | 20,157 |
| Rhode Island   | 14,250 | 14,773 | 15,667 | 17,000 | 17,727 | 17,909 | 16,636 | 19,000 | 19,000 |
| South Carolina | 12,771 | 14,481 | 16,087 | 16,000 | 17,875 | 17,438 | 17,250 | 16,500 | 16,375 |
| South Dakota   | 12,257 | 13,398 | 15,516 | 17,741 | 20,478 | 20,582 | 21,391 | 21,521 | 21,742 |
| Tennessee      | 11,825 | 13,740 | 14,789 | 15,743 | 16,346 | 16,200 | 16,100 | 15,938 | 16,304 |
| Texas          | 14,350 | 15,244 | 16,503 | 19,646 | 21,375 | 22,232 | 22,009 | 21,991 | 22,268 |
| Utah           | 15,838 | 16,739 | 17,573 | 18,875 | 21,898 | 22,161 | 22,863 | 22,432 | 22,968 |
| Vermont        | 14,528 | 16,210 | 17,199 | 18,469 | 18,537 | 18,940 | 19,316 | 19,448 | 20,197 |
| Virginia       | 14,213 | 15,116 | 15,833 | 16,990 | 18,095 | 17,906 | 17,990 | 18,337 | 19,140 |
| Washington     | 18,532 | 20,091 | 22,644 | 23,270 | 23,514 | 23,727 | 23,794 | 23,820 | 24,117 |
| West Virginia  | 11,250 | 12,667 | 15,588 | 14,923 | 15,700 | 15,700 | 15,400 | 15,200 | 15,556 |
| Wisconsin      | 13,973 | 15,397 | 17,306 | 18,500 | 20,630 | 20,599 | 21,436 | 21,693 | 21,869 |
| Wyoming        | 12,337 | 13,197 | 13,571 | 14,878 | 20,067 | 20,517 | 20,650 | 21,367 | 21,583 |

Source: USDA (2015).

## Step 2: Characterize U.S. Cattle Population Diets

To support development of digestible energy (DE, the percent of gross energy intake digested by the animal) and CH<sub>4</sub> conversion rate (Y<sub>m</sub>, the fraction of gross energy converted to CH<sub>4</sub>) values for each of the cattle population categories, data were collected on diets considered representative of different regions. For both grazing animals and animals being fed mixed rations, representative regional diets were estimated using information collected from state livestock specialists, the USDA, expert opinion, and other literature sources. The designated regions for this analysis for dairy cattle for all years and foraging beef cattle from 1990 through 2006 are shown in Table A-188. For foraging beef cattle from 2007 onwards, the regional designations were revised based on data available from the NAHMS 2007–2008 survey on cow-calf system management practices (USDA:APHIS:VS 2010) and are shown in and Table A-189. The data for each of the diets (e.g., proportions of different feed constituents, such as hay or grains) were used to determine feed chemical composition for use in estimating DE and Y<sub>m</sub> for each animal type.

**Table A-188: Regions used for Characterizing the Diets of Dairy Cattle (all years) and Foraging Cattle from 1990–2006**

| West       | California | Northern Great Plains | Midwestern | Northeast     | Southcentral | Southeast      |
|------------|------------|-----------------------|------------|---------------|--------------|----------------|
| Alaska     | California | Colorado              | Illinois   | Connecticut   | Arkansas     | Alabama        |
| Arizona    |            | Kansas                | Indiana    | Delaware      | Louisiana    | Florida        |
| Hawaii     |            | Montana               | Iowa       | Maine         | Oklahoma     | Georgia        |
| Idaho      |            | Nebraska              | Michigan   | Maryland      | Texas        | Kentucky       |
| Nevada     |            | North Dakota          | Minnesota  | Massachusetts |              | Mississippi    |
| New Mexico |            | South Dakota          | Missouri   | New Hampshire |              | North Carolina |
| Oregon     |            | Wyoming               | Ohio       | New Jersey    |              | South Carolina |
| Utah       |            |                       | Wisconsin  | New York      |              | Tennessee      |
| Washington |            |                       |            | Pennsylvania  |              | Virginia       |
|            |            |                       |            | Rhode Island  |              |                |
|            |            |                       |            | Vermont       |              |                |
|            |            |                       |            | West Virginia |              |                |

Source: USDA (1996).

**Table A-189: Regions used for Characterizing the Diets of Foraging Cattle from 2007–2014**

| West              | Central             | Northeast     | Southeast        |
|-------------------|---------------------|---------------|------------------|
| Alaska            | Illinois            | Connecticut   | Alabama          |
| Arizona           | Indiana             | Delaware      | <b>Arkansas</b>  |
| <b>California</b> | Iowa                | Maine         | Florida          |
| <b>Colorado</b>   | <b>Kansas</b>       | Maryland      | Georgia          |
| Hawaii            | Michigan            | Massachusetts | Kentucky         |
| Idaho             | Minnesota           | New Hampshire | <b>Louisiana</b> |
| <b>Montana</b>    | Missouri            | New Jersey    | Mississippi      |
| Nevada            | <b>Nebraska</b>     | New York      | North Carolina   |
| New Mexico        | <b>North Dakota</b> | Pennsylvania  | <b>Oklahoma</b>  |
| Oregon            | Ohio                | Rhode Island  | South Carolina   |
| Utah              | <b>South Dakota</b> | Vermont       | Tennessee        |
| Washington        | Wisconsin           | West Virginia | <b>Texas</b>     |
| <b>Wyoming</b>    |                     |               | Virginia         |

Source: Based on data from USDA:APHIS:VS (2010).

Note: States in **bold** represent a change in region from the 1990–2006 assessment.

DE and Y<sub>m</sub> vary by diet and animal type. The IPCC recommends Y<sub>m</sub> values of 3.0±1.0 percent for feedlot cattle and 6.5±1.0 percent for all other cattle (IPCC 2006). Given the availability of detailed diet information for different regions and animal types in the United States, DE and Y<sub>m</sub> values unique to the United States were developed for dairy and beef cattle. Digestible energy and Y<sub>m</sub> values were estimated across the time series for each cattle population category based on physiological modeling, published values, and/or expert opinion.

For dairy cows, ruminant digestion models were used to estimate Y<sub>m</sub>. The three major categories of input required by the models are animal description (e.g., cattle type, mature weight), animal performance (e.g., initial and final weight, age at start of period), and feed characteristics (e.g., chemical composition, habitat, grain or forage). Data used to simulate ruminant digestion is provided for a particular animal that is then used to represent a group of animals with similar characteristics. The Y<sub>m</sub> values were estimated for 1990 using the Donovan and Baldwin model (1999), which represents

physiological processes in the ruminant animals, as well as diet characteristics from USDA (1996). The Donovan and Baldwin model is able to account for differing diets (i.e., grain-based or forage-based), so that  $Y_m$  values for the variable feeding characteristics within the U.S. cattle population can be estimated. Subsequently, a literature review of dairy diets was conducted and nearly 250 diets were analyzed from 1990 through 2009 across 23 states—the review indicated highly variable diets, both temporally and spatially. Kebreab et al. (2008) conducted an evaluation of models and found that the COWPOLL model was the best model for estimating  $Y_m$  for dairy. The statistical analysis of the COWPOLL model showed a trend in predicting  $Y_m$ , and inventory team experts determined that the most comprehensive approach was to use the 1990 baseline from Donovan and Baldwin and then scale  $Y_m$  values for each of the diets beyond 1990 with the COWPOLL model. A function based on the national trend observed from the analysis of the dairy diets was used to calculate 1991 and beyond regional values based on the regional 1990  $Y_m$  values from Donovan and Baldwin. The resulting scaling factor (incorporating both Donovan and Baldwin (1999) and COWPOLL) is shown below:

$$Y_m = Y_m(1990) \text{EXP} \left( \frac{1.22}{(\text{Year} - 1980)} \right) / \text{EXP} \left( \frac{1.22}{(1990 - 1980)} \right)$$

DE values for dairy cows were estimated from the literature search based on the annual trends observed in the data collection effort. The regional variability observed in the literature search was not statistically significant, and therefore DE was not varied by region, but did vary over time, and was grouped by the following years 1990–1993, 1994–1998, 1999–2003, 2004–2006, 2007, and 2008 onwards.

Considerably less data was available for dairy heifers and dairy calves. Therefore, for dairy heifers assumptions were based on the relationship of the collected data in the literature on dairy heifers to the data on dairy cow diets. From this relationship, DE was estimated as the mature cow DE minus three percent, and  $Y_m$  was estimated as that of the mature dairy cow plus 0.1 percent.

To calculate the DE values for grazing beef cattle, diet composition assumptions were used to estimate weighted DE values for a combination of forage and supplemental diets. The forage portion makes up an estimated 85 to 95 percent of grazing beef cattle diets, and there is considerable variation of both forage type and quality across the United States. Currently there is no comprehensive survey of this data, so for this analysis two regional DE values were developed to account for the generally lower forage quality in the “West” region of the United States versus all other regions in Table A-188 (California, Northern Great Plains, Midwestern, Northeast, Southcentral, Southeast) and Table A-189 (Central, Northeast, and Southeast). For all non-western grazing cattle, the forage DE was an average of the estimated seasonal values for grass pasture diets for a calculated DE of 64.2 percent. For foraging cattle in the west, the forage DE was calculated as the seasonal average for grass pasture, meadow and range diets, for a calculated DE of 61.3 percent. The assumed specific components of each of the broad forage types, along with their corresponding DE value and the calculated regional DE values can be found in Table A-190. In addition, beef cattle are assumed to be fed a supplemental diet, consequently, two sets of supplemental diets were developed, one for 1990 through 2006 (Donovan 1999) and one for 2007 onwards (Preston 2010, Archibeque 2011, USDA:APHIS:VS 2010) as shown in Table A-191 and Table A-192 along with the percent of each total diet that is assumed to be made up of the supplemental portion. By weighting the calculated DE values from the forage and supplemental diets, the DE values for the composite diet were calculated.<sup>77</sup> These values are used for steer and heifer stockers and beef replacements. Finally, for mature beef cows and bulls, the DE value was adjusted downward by two percent to reflect the lower digestibility diets of mature cattle based on Johnson (2002).  $Y_m$  values for all grazing beef cattle were set at 6.5 percent based on Johnson (2002). The  $Y_m$  values and the resulting final weighted DE values by region for 2007 onwards are shown in Table A-193.

For feedlot animals, DE and  $Y_m$  are adjusted over time as diet compositions in actual feedlots are adjusted based on new and improved nutritional information and availability of feed types. Feedlot diets are assumed to not differ significantly by state, and therefore only a single set of national diet values is utilized for each year. The DE and  $Y_m$  values for 1990 were estimated by Dr. Don Johnson (1999). In the CEFM, the DE values for 1991 through 1999 were linearly extrapolated based on values for 1990 and 2000. DE and  $Y_m$  values from 2000 through the current year were estimated using the MOLLY model as described in Kebreab et al. (2008), based on a series of average diet feed compositions from Galyean and Gleghorn (2001) for 2000 through 2006 and Vasconcelos and Galyean (2007) for 2007 onwards. In addition, feedlot animals are assumed to spend the first 25 days in the feedlot on a “step-up” diet to become accustomed to the higher

<sup>77</sup> For example, the West has a forage DE of 61.3 which makes up 90 percent of the diet and a supplemented diet DE of 67.4 percent was used for 10 percent of the diet, for a total weighted DE of 61.9 percent, as shown in Table A-193

quality feedlot diets. The step-up DE and  $Y_m$  are calculated as the average of all state forage and feedlot diet DE and  $Y_m$  values.

For calves aged 4 through 6 months, a gradual weaning from milk is simulated, with calf diets at 4 months assumed to be 25 percent forage, increasing to 50 percent forage at age 5 months, and 75 percent forage at age 6 months. The portion of the diet allocated to milk results in zero emissions, as recommended by the IPCC (2006). For calves, the DE for the remainder of the diet is assumed to be similar to that of slightly older replacement heifers (both beef and dairy are calculated separately). The  $Y_m$  for beef calves is also assumed to be similar to that of beef replacement heifers (6.5 percent), as literature does not provide an alternative  $Y_m$  for use in beef calves. For dairy calves, the  $Y_m$  is assumed to be 7.8 percent at 4 months, 8.03 percent at 5 months, and 8.27 percent at 6 months based on estimates provided by Soliva (2006) for  $Y_m$  at 4 and 7 months of age and a linear interpolation for 5 and 6 months.

Table A-194 shows the regional DE and  $Y_m$  for U.S. cattle in each region for 2014.

**Table A-190: Feed Components and Digestible Energy Values Incorporated into Forage Diet Composition Estimates**

| Forage Type  | DE (% of GE) | Grass pasture - Spring | Grass pasture - Summer | Grass pasture - Fall | Range June | Range July | Range August | Range September | Range Winter | Meadow - Spring | Meadow - Fall |
|--|--------------|------------------------|------------------------|----------------------|------------|------------|--------------|-----------------|--------------|-----------------|---------------|
| Bahiagrass Paspalum notatum, fresh                         | 61.38        |                        |                        | x                    |            |            |              |                 |              |                 |               |
| Bermudagrass Cynodon dactylon, fresh                       | 66.29        |                        | x                      |                      |            |            |              |                 |              |                 |               |
| Bremudagrass, Coastal Cynodon dactylon, fresh              | 65.53        |                        | x                      |                      |            |            |              |                 |              |                 |               |
| Bluegrass, Canada Poa compressa, fresh, early vegetative   | 73.99        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Bluegrass, Kentucky Poa pratensis, fresh, early vegetative | 75.62        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Bluegrass, Kentucky Poa pratensis, fresh, mature           | 59.00        |                        | x                      | x                    |            |            |              |                 |              |                 |               |
| Bluestem Andropogon spp, fresh, early vegetative           | 73.17        |                        |                        |                      | x          |            |              |                 |              |                 |               |
| Bluestem Andropogon spp, fresh, mature                     | 56.82        |                        |                        |                      |            | x          | x            | x               | x            |                 | x             |
| Brome Bromus spp, fresh, early vegetative                  | 78.57        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Brome, Smooth Bromus inermis, fresh, early vegetative      | 75.71        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Brome, Smooth Bromus inermis, fresh, mature                | 57.58        |                        | x                      | x                    |            |            |              |                 | x            |                 |               |
| Buffalograss, Buchloe dactyloides, fresh                   | 64.02        |                        |                        |                      | x          | x          |              |                 |              |                 |               |
| Clover, Alsike Trifolium hybridum, fresh, early vegetative | 70.62        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Clover, Ladino Trifolium repens, fresh, early vegetative   | 73.22        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Clover, Red Trifolium pratense, fresh, early bloom         | 71.27        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Clover, Red Trifolium pratense, fresh, full bloom          | 67.44        |                        | x                      |                      | x          |            |              |                 |              |                 |               |

| Forage Type   | DE (% of GE) | Grass pasture - Spring | Grass pasture - Summer | Grass pasture - Fall | Range June | Range July | Range August | Range September | Range Winter | Meadow - Spring | Meadow - Fall |
|---|--------------|------------------------|------------------------|----------------------|------------|------------|--------------|-----------------|--------------|-----------------|---------------|
| Corn, Dent Yellow Zea mays indentata, aerial part without ears, without husks, sun-cured, (stover)(straw) | 55.28        |                        |                        | x                    |            |            |              |                 |              |                 |               |
| Dropseed, Sand Sporobolus cryptandrus, fresh, stem cured  | 64.69        |                        |                        |                      | x          | x          | x            |                 |              | x               |               |
| Fescue Festuca spp, hay, sun-cured, early vegetative  | 67.39        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Fescue Festuca spp, hay, sun-cured, early bloom   | 53.57        |                        |                        | x                    |            |            |              |                 |              |                 |               |
| Grama Bouteloua spp, fresh, early vegetative  | 67.02        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Grama Bouteloua spp, fresh, mature  | 63.38        |                        | x                      | x                    |            |            |              |                 |              | x               |               |
| Millet, Foxtail Setaria italica, fresh  | 68.20        | x                      |                        |                      | x          |            |              |                 |              |                 |               |
| Napiersgrass Pennisetum purpureum, fresh, late bloom  | 57.24        |                        | x                      | x                    |            |            |              |                 |              |                 |               |
| Needleandthread Stipa comata, fresh, stem cured   | 60.36        |                        |                        |                      |            | x          | x            | x               |              |                 |               |
| Orchardgrass Dactylis glomerata, fresh, early vegetative  | 75.54        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Orchardgrass Dactylis glomerata, fresh, midbloom  | 60.13        |                        | x                      |                      |            |            |              |                 |              |                 |               |
| Pearlmillet Pennisetum glaucum, fresh   | 68.04        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Prairie plants, Midwest, hay, sun-cured   | 55.53        |                        |                        | x                    |            |            |              |                 |              |                 | x             |
| Rape Brassica napus, fresh, early bloom   | 80.88        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Rye Secale cereale, fresh   | 71.83        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Ryegrass, Perennial Lolium perenne, fresh   | 73.68        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Saltgrass Distichlis spp, fresh, post ripe  | 58.06        |                        | x                      | x                    |            |            |              |                 |              |                 |               |
| Sorghum, Sudangrass Sorghum bicolor sudanense, fresh, early vegetative                                    | 73.27        | x                      |                        |                      |            |            |              |                 |              |                 |               |
| Squirreltail Stanion spp, fresh, stem-cured   | 62.00        |                        | x                      |                      |            | x          |              |                 |              |                 |               |
| Summercypress, Gray Kochia vestita, fresh, stem-cured   | 65.11        |                        |                        | x                    | x          | x          |              |                 |              |                 |               |

| Forage Type   | DE (% of GE) | Grass pasture - Spring | Grass pasture - Summer | Grass pasture - Fall | Range June   | Range July   | Range August | Range September | Range Winter | Meadow - Spring | Meadow - Fall |
|---|--------------|------------------------|------------------------|----------------------|--------------|--------------|--------------|-----------------|--------------|-----------------|---------------|
| Timothy Phleum pratense, fresh, late vegetative                   | 73.12        | x                      |                        |                      |              |              |              |                 |              |                 |               |
| Timothy Phleum pratense, fresh, midbloom                          | 66.87        |                        | x                      |                      |              |              |              |                 |              |                 |               |
| Trefoil, Birdsfoot Lotus corniculatus, fresh                      | 69.07        | x                      |                        |                      |              |              |              |                 |              |                 |               |
| Vetch Vicia spp, hay, sun-cured                                   | 59.44        |                        |                        | x                    |              |              |              |                 |              |                 |               |
| Wheat Triticum aestivum, straw                                    | 45.77        |                        |                        | x                    |              |              |              |                 |              |                 |               |
| Wheatgrass, Crested Agropyron desertorum, fresh, early vegetative | 79.78        | x                      |                        |                      |              |              |              |                 |              |                 |               |
| Wheatgrass, Crested Agropyron desertorum, fresh, full bloom       | 65.89        |                        | x                      |                      |              | x            |              |                 |              |                 |               |
| Wheatgrass, Crested Agropyron desertorum, fresh, post ripe        | 52.99        |                        |                        | x                    |              |              |              |                 | x            |                 | x             |
| Winterfat, Common Eurotia lanata, fresh, stem-cured               | 40.89        |                        |                        |                      |              |              |              |                 | x            |                 |               |
| <b>Weighted Average DE</b>  |              | <b>72.99</b>           | <b>62.45</b>           | <b>57.26</b>         | <b>67.11</b> | <b>62.70</b> | <b>60.62</b> | <b>58.59</b>    | <b>52.07</b> | <b>64.03</b>    | <b>55.11</b>  |
| <b>Forage Diet for West</b>                                       | <b>61.3</b>  | <b>10%</b>             | <b>10%</b>             | <b>10%</b>           | <b>10%</b>   | <b>10%</b>   | <b>10%</b>   | <b>10%</b>      | <b>10%</b>   | <b>10%</b>      | <b>10%</b>    |
| <b>Forage Diet for All Other Regions</b>                          | <b>64.2</b>  | <b>33.3%</b>           | <b>33.3%</b>           | <b>33.3%</b>         | <b>-</b>     | <b>-</b>     | <b>-</b>     | <b>-</b>        | <b>-</b>     | <b>-</b>        | <b>-</b>      |

Note that forages marked with an x indicate that the DE from that specific forage type is included in the general forage type for that column (e.g., grass pasture, range, meadow or meadow by month or season).  
Sources: Preston (2010) and Archibeque (2011).

**Table A-191: DE Values with Representative Regional Diets for the Supplemental Diet of Grazing Beef Cattle for 1990–2006**

| Feed             | Source of DE (NRC 1984)      | Unweighted DE (% of GE) | Northern Great Plains |      |        |              |           |         |           |
|------------------|------------------------------|-------------------------|-----------------------|------|--------|--------------|-----------|---------|-----------|
|                  |                              |                         | California *          | West | Plains | Southcentral | Northeast | Midwest | Southeast |
| Alfalfa Hay      | Table 8, feed #006           | 61.79                   | 65%                   | 30%  | 30%    | 29%          | 12%       | 30%     |           |
| Barley           |                              | 85.08                   | 10%                   | 15%  |        |              |           |         |           |
| Bermuda          | Table 8, feed #030           | 66.29                   |                       |      |        |              |           |         | 35%       |
| Bermuda Hay      | Table 8, feed #031           | 50.79                   |                       |      |        | 40%          |           |         |           |
| Corn             | Table 8, feed #089           | 88.85                   | 10%                   | 10%  | 25%    | 11%          | 13%       | 13%     |           |
| Corn Silage      | Table 8, feed #095           | 72.88                   |                       |      | 25%    |              | 20%       | 20%     |           |
| Cotton Seed Meal |                              |                         |                       |      |        | 7%           |           |         |           |
| Grass Hay        | Table 8, feed #126, 170, 274 | 58.37                   |                       | 40%  |        |              |           | 30%     |           |
| Orchard          | Table 8, feed #147           | 60.13                   |                       |      |        |              |           |         | 40%       |
| Soybean Meal     |                              |                         |                       |      |        |              |           |         |           |
| Supplement       |                              | 77.15                   |                       | 5%   | 5%     |              |           |         | 5%        |
| Sorghum          | Table 8, feed #211           | 84.23                   |                       |      |        |              |           |         | 20%       |
| Soybean Hulls    |                              | 66.86                   |                       |      |        |              |           | 7%      |           |
| Timothy Hay      | Table 8, feed #244           | 60.51                   |                       |      |        |              | 50%       |         |           |

|   |                    |             |             |             |             |             |             |             |
|---|--------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Whole Cotton Seed                         |                    | 75.75       | 5%          |             |             | 5%          |             |             |
| Wheat Middlings                           | Table 8, feed #257 | 68.09       |             | 15%         | 13%         |             |             |             |
| Wheat                                     | Table 8, feed #259 | 87.95       | 10%         |             |             |             |             |             |
| <b>Weighted Supplement DE (%)</b>         |                    | <b>70.1</b> | <b>67.4</b> | <b>73.0</b> | <b>62.0</b> | <b>67.6</b> | <b>66.9</b> | <b>68.0</b> |
| <b>Percent of Diet that is Supplement</b> |                    | <b>5%</b>   | <b>10%</b>  | <b>15%</b>  | <b>10%</b>  | <b>15%</b>  | <b>10%</b>  | <b>5%</b>   |

Source of representative regional diets: Donovan (1999).

\* Note that emissions are currently calculated on a state-by-state basis, but diets are applied by the regions shown in the table above.

**Table A-192: DE Values and Representative Regional Diets for the Supplemental Diet of Grazing Beef Cattle for 2007–2014**

| Feed                                       | Source of DE (NRC1984)                    | Unweighted DE (% of GE) | West <sup>a</sup> | Central <sup>a</sup> | Northeast <sup>a</sup> | Southeast <sup>a</sup> |
|--|---|-------------------------|-------------------|----------------------|------------------------|------------------------|
| Alfalfa Hay                                | Table 8, feed #006                        | 61.79                   | 65%               | 30%                  | 12%                    |                        |
| Bermuda                                    | Table 8, feed #030                        | 66.29                   |                   |                      |                        | 20%                    |
| Bermuda Hay                                | Table 8, feed #031                        | 50.79                   |                   |                      |                        | 20%                    |
| Corn                                       | Table 8, feed #089                        | 88.85                   | 10%               | 15%                  | 13%                    | 10%                    |
| Corn Silage                                | Table 8, feed #095                        | 72.88                   |                   | 35%                  | 20%                    |                        |
| Grass Hay                                  | Table 8, feed #126, 170, 274              | 58.37                   | 10%               |                      |                        |                        |
| Orchard                                    | Table 8, feed #147                        | 60.13                   |                   |                      |                        | 30%                    |
| Protein supplement (West)                  | Table 8, feed #082, 134, 225 <sup>b</sup> | 81.01                   | 10%               |                      |                        |                        |
| Protein Supplement (Central and Northeast) | Table 8, feed #082, 134, 225 <sup>b</sup> | 80.76                   |                   | 10%                  | 10%                    |                        |
| Protein Supplement (Southeast)             | Table 8, feed #082, 134, 101 <sup>b</sup> | 77.89                   |                   |                      |                        | 10%                    |
| Sorghum                                    | Table 8, feed #211                        | 84.23                   |                   | 5%                   |                        | 10%                    |
| Timothy Hay                                | Table 8, feed #244                        | 60.51                   |                   |                      | 45%                    |                        |
| Wheat Middlings                            | Table 8, feed #257                        | 68.09                   |                   | 5%                   |                        |                        |
| Wheat                                      | Table 8, feed #259                        | 87.95                   | 5%                |                      |                        |                        |
| <b>Weighted Supplement DE</b>              |   |                         | <b>67.4</b>       | <b>73.1</b>          | <b>68.9</b>            | <b>66.6</b>            |
| <b>Percent of Diet that is Supplement</b>  |   |                         | <b>10%</b>        | <b>15%</b>           | <b>5%</b>              | <b>15%</b>             |

<sup>a</sup> Note that emissions are currently calculated on a state-by-state basis, but diets are applied by the regions shown in the table above.

<sup>b</sup> Not in equal proportions.

Sources of representative regional diets: Donovan (1999), Preston (2010), Archibeque (2011), and USDA:APHIS:VS (2010).

**Table A-193: Foraging Animal DE (% of GE) and Y<sub>m</sub> Values for Each Region and Animal Type for 2007–2014**

| Animal Type          | Data                        | West <sup>a</sup> | Central | Northeast | Southeast |
|----------------------|-----------------------------|-------------------|---------|-----------|-----------|
| Beef Repl. Heifers   | DE <sup>b</sup>             | 61.9              | 65.6    | 64.5      | 64.6      |
|                      | Y <sub>m</sub> <sup>c</sup> | 6.5%              | 6.5%    | 6.5%      | 6.5%      |
| Beef Calves (4–6 mo) | DE                          | 61.9              | 65.6    | 64.5      | 64.6      |
|                      | Y <sub>m</sub>              | 6.5%              | 6.5%    | 6.5%      | 6.5%      |
| Steer Stockers       | DE                          | 61.9              | 65.6    | 64.5      | 64.6      |
|                      | Y <sub>m</sub>              | 6.5%              | 6.5%    | 6.5%      | 6.5%      |
| Heifer Stockers      | DE                          | 61.9              | 65.6    | 64.5      | 64.6      |
|                      | Y <sub>m</sub>              | 6.5%              | 6.5%    | 6.5%      | 6.5%      |
| Beef Cows            | DE                          | 59.9              | 63.6    | 62.5      | 62.6      |
|                      | Y <sub>m</sub>              | 6.5%              | 6.5%    | 6.5%      | 6.5%      |
| Bulls                | DE                          | 59.9              | 63.6    | 62.5      | 62.6      |
|                      | Y <sub>m</sub>              | 6.5%              | 6.5%    | 6.5%      | 6.5%      |

<sup>a</sup> Note that emissions are currently calculated on a state-by-state basis, but diets are applied by the regions shown in the table above. To see the regional designation per state, please see Table A-189.

<sup>b</sup> DE is the digestible energy in units of percent of GE (MJ/Day).

<sup>c</sup> Y<sub>m</sub> is the methane conversion rate, the fraction of GE in feed converted to methane.

**Table A-194: Regional DE (% of GE) and Y<sub>m</sub> Rates for Dairy and Feedlot Cattle by Animal Type for 2014**

| Animal Type           | Data            | Northern  |      |              |              |           |         |           |
|-----------------------|-----------------|---|------|--------------|--------------|-----------|---------|-----------|
|                       |                 | California <sup>a</sup>                             | West | Great Plains | Southcentral | Northeast | Midwest | Southeast |
| Dairy Repl. Heifers   | DE <sup>b</sup> | 63.7  | 63.7 | 63.7         | 63.7         | 63.7      | 63.7    | 63.7      |
|                       | Ym <sup>c</sup> | 6.0%  | 6.0% | 5.7%         | 6.5%         | 6.4%      | 5.7%    | 7.0%      |
| Dairy Calves (4–6 mo) | DE              | 63.7  | 63.7 | 63.7         | 63.7         | 63.7      | 63.7    | 63.7      |
|                       | Ym              | 7.8% (4 mo), 8.03% (5 mo), 8.27% (6 mo)-all regions |      |              |              |           |         |           |
| Dairy Cows            | DE              | 66.7  | 66.7 | 66.7         | 66.7         | 66.7      | 66.7    | 66.7      |
|                       | Ym              | 5.9%  | 5.9% | 5.6%         | 6.4%         | 6.3%      | 5.6%    | 6.9%      |
| Steer Feedlot         | DE              | 82.5  | 82.5 | 82.5         | 82.5         | 82.5      | 82.5    | 82.5      |
|                       | Ym              | 3.9%  | 3.9% | 3.9%         | 3.9%         | 3.9%      | 3.9%    | 3.9%      |
| Heifer Feedlot        | DE              | 82.5  | 82.5 | 82.5         | 82.5         | 82.5      | 82.5    | 82.5      |
|                       | Ym              | 3.9%  | 3.9% | 3.9%         | 3.9%         | 3.9%      | 3.9%    | 3.9%      |

<sup>a</sup> Note that emissions are currently calculated on a state-by-state basis, but diets are applied in Table A-188 by the regions shown in the table above. To see the regional designation for foraging cattle per state, please see Table A-188.

<sup>b</sup> DE is the digestible energy in units of percent of GE (MJ/Day).

<sup>c</sup> Y<sub>m</sub> is the methane conversion rate, the fraction of GE in feed converted to methane.

### Step 3: Estimate CH<sub>4</sub> Emissions from Cattle

Emissions by state were estimated in three steps: a) determine gross energy (GE) intake using the Tier 2 IPCC (2006) equations, b) determine an emission factor using the GE values, Y<sub>m</sub> and a conversion factor, and c) sum the daily emissions for each animal type. Finally, the state emissions were aggregated to obtain the national emissions estimate. The necessary data values for each state and animal type include:

- Body Weight (kg)
- Weight Gain (kg/day)
- Net Energy for Activity (C<sub>a</sub>, MJ/day)<sup>78</sup>
- Standard Reference Weight (kg)<sup>79</sup>
- Milk Production (kg/day)
- Milk Fat (percent of fat in milk = 4)
- Pregnancy (percent of population that is pregnant)
- DE (percent of GE intake digestible)
- Y<sub>m</sub> (the fraction of GE converted to CH<sub>4</sub>)
- Population

#### Step 3a: Determine Gross Energy, GE

As shown in the following equation, GE is derived based on the net energy estimates and the feed characteristics. Only variables relevant to each animal category are used (e.g., estimates for feedlot animals do not require the NE<sub>l</sub> factor). All net energy equations are provided in IPCC (2006).

$$GE = \left[ \frac{\left( \frac{NE_m + NE_a + NE_l + NE_{work} + NE_p}{REM} \right) + \left( \frac{NE_g}{REG} \right)}{\frac{DE\%}{100}} \right]$$

where,

GE = Gross energy (MJ/day)  
 NE<sub>m</sub> = Net energy required by the animal for maintenance (MJ/day)

<sup>78</sup> Zero for feedlot conditions, 0.17 for high quality confined pasture conditions, and 0.36 for extensive open range or hilly terrain grazing conditions. C<sub>a</sub> factor for dairy cows is weighted to account for the fraction of the population in the region that grazes during the year (IPCC 2006).

<sup>79</sup> Standard Reference Weight is the mature weight of a female animal of the animal type being estimated, used in the model to account for breed potential.

- NE<sub>a</sub> = Net energy for animal activity (MJ/day)
- NE<sub>l</sub> = Net energy for lactation (MJ/day)
- NE<sub>work</sub> = Net energy for work (MJ/day)
- NE<sub>p</sub> = Net energy required for pregnancy (MJ/day)
- REM = Ratio of net energy available in a diet for maintenance to digestible energy consumed
- NE<sub>g</sub> = Net energy needed for growth (MJ/day)
- REG = Ratio of net energy available for growth in a diet to digestible energy consumed
- DE = Digestible energy expressed as a percent of gross energy (percent)

**Step 3b: Determine Emission Factor**

The daily emission factor (DayEmit) was determined using the GE value and the methane conversion factor (Y<sub>m</sub>) for each category. This relationship is shown in the following equation:

$$DayEmit = \frac{GE \times Y_m}{55.65}$$

where,

- DayEmit = Emission factor (kg CH<sub>4</sub>/head/day)
- GE = Gross energy intake (MJ/head/day)
- Y<sub>m</sub> = CH<sub>4</sub> conversion rate, which is the fraction of GE in feed converted to CH<sub>4</sub> (%)
- 55.65 = A factor for the energy content of methane (MJ/kg CH<sub>4</sub>)

The daily emission factors were estimated for each animal type and state. Calculated annual national emission factors are shown by animal type in Table A-195.

**Table A-195: Calculated Annual National Emission Factors for Cattle by Animal Type (kg CH<sub>4</sub>/head/year)**

| Cattle Type                 | 1990 | 1995 | 2000 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-----------------------------|------|------|------|------|------|------|------|------|------|
| <b>Dairy</b>                |      |      |      |      |      |      |      |      |      |
| Calves                      | 12   | 12   | 12   | 12   | 12   | 12   | 12   | 12   | 12   |
| Cows                        | 124  | 125  | 132  | 133  | 142  | 142  | 144  | 144  | 145  |
| Replacements (7–11 months)  | 48   | 46   | 46   | 45   | 46   | 46   | 46   | 46   | 46   |
| Replacements (12–23 months) | 73   | 69   | 70   | 67   | 69   | 69   | 69   | 69   | 69   |
| <b>Beef</b>                 |      |      |      |      |      |      |      |      |      |
| Calves                      | 11   | 11   | 11   | 11   | 11   | 11   | 11   | 11   | 11   |
| Bulls                       | 91   | 94   | 94   | 97   | 98   | 98   | 98   | 98   | 98   |
| Cows                        | 89   | 92   | 91   | 94   | 95   | 95   | 95   | 95   | 95   |
| Replacements (7–11 months)  | 54   | 57   | 56   | 59   | 60   | 60   | 60   | 60   | 60   |
| Replacements (12–23 months) | 63   | 66   | 66   | 68   | 70   | 70   | 70   | 70   | 70   |
| Steer Stockers              | 55   | 57   | 58   | 58   | 58   | 58   | 58   | 58   | 58   |
| Heifer Stockers             | 52   | 56   | 60   | 60   | 60   | 60   | 60   | 60   | 60   |
| Feedlot Cattle              | 39   | 38   | 39   | 39   | 42   | 42   | 42   | 43   | 43   |

Note: To convert to a daily emission factor, the yearly emission factor can be divided by 365 (the number of days in a year).

For quality assurance purposes, U.S. emission factors for each animal type were compared to estimates provided by the other Annex I member countries of the United Nations Framework Convention on Climate Change (UNFCCC) (the most recently available summarized results for Annex I countries are through 2012 only). Results, presented in Table A-196, indicate that U.S. emission factors are comparable to those of other Annex I countries. Results in Table A-196 are presented along with Tier 1 emission factors provided by IPCC (2006). Throughout the time series, beef cattle in the United States generally emit more enteric CH<sub>4</sub> per head than other Annex I member countries, while dairy cattle in the United States generally emit comparable enteric CH<sub>4</sub> per head.

**Table A-196: Annex I Countries' Implied Emission Factors for Cattle by Year (kg CH<sub>4</sub>/head/year)<sup>80</sup>**

| Year   | Dairy Cattle                          |  | Beef Cattle                           |  |
|--|---------------------------------------|--|---------------------------------------|--|
|  | United States Implied Emission Factor | Mean of Implied Emission Factors for Annex I countries (excluding United States) | United States Implied Emission Factor | Mean of Implied Emission Factors for Annex I countries (excluding United States) |
| 1990   | 107                                   | 96   | 71                                    | 53   |
| 1991   | 107                                   | 97   | 71                                    | 53   |
| 1992   | 107                                   | 96   | 72                                    | 54   |
| 1993   | 106                                   | 97   | 72                                    | 54   |
| 1994   | 106                                   | 98   | 73                                    | 54   |
| 1995   | 106                                   | 98   | 72                                    | 54   |
| 1996   | 105                                   | 99   | 73                                    | 54   |
| 1997   | 106                                   | 100  | 73                                    | 54   |
| 1998   | 107                                   | 101  | 73                                    | 55   |
| 1999   | 110                                   | 102  | 72                                    | 55   |
| 2000   | 111                                   | 103  | 72                                    | 55   |
| 2001   | 110                                   | 104  | 73                                    | 55   |
| 2002   | 111                                   | 105  | 73                                    | 55   |
| 2003   | 111                                   | 106  | 73                                    | 55   |
| 2004   | 109                                   | 107  | 74                                    | 55   |
| 2005   | 110                                   | 109  | 74                                    | 55   |
| 2006   | 110                                   | 110  | 74                                    | 55   |
| 2007   | 114                                   | 111  | 75                                    | 55   |
| 2008   | 115                                   | 112  | 75                                    | 55   |
| 2009   | 115                                   | 112  | 75                                    | 56   |
| 2010   | 115                                   | 113  | 75                                    | 55   |
| 2011   | 116                                   | 113  | 75                                    | 55   |
| 2012   | 117                                   | 112  | 75                                    | 51   |
| 2013   | 117                                   | NA   | 75                                    | NA   |
| 2014   | 118                                   | NA   | 74                                    | NA   |
| Tier I EFs For North America, from IPCC (2006) |                                       | 121  |                                       | 53   |

**Step 3c: Estimate Total Emissions**

Emissions were summed for each month and for each state population category using the daily emission factor for a representative animal and the number of animals in the category. The following equation was used:

$$\text{Emissions}_{\text{state}} = \text{DayEmit}_{\text{state}} \times \text{Days/Month} \times \text{SubPop}_{\text{state}}$$

where,

|                            |   |  |
|----------------------------|---|--|
| Emissions <sub>state</sub> | = | Emissions for state during the month (kg CH <sub>4</sub> )                   |
| DayEmit <sub>state</sub>   | = | Emission factor for the subcategory and state (kg CH <sub>4</sub> /head/day) |
| Days/Month                 | = | Number of days in the month  |
| SubPop <sub>state</sub>    | = | Number of animals in the subcategory and state during the month              |

This process was repeated for each month, and the monthly totals for each state subcategory were summed to achieve an emission estimate for a state for the entire year and state estimates were summed to obtain the national total. The

<sup>80</sup> Excluding calves.

estimates for each of the 10 subcategories of cattle are listed in Table A-197. The emissions for each subcategory were then aggregated to estimate total emissions from beef cattle and dairy cattle for the entire year.

**Table A-197: CH<sub>4</sub> Emissions from Cattle (kt)**

| Cattle Type                 | 1990         | 1995         | 2000         | 2005         | 2010         | 2011         | 2012         | 2013         | 2014         |
|-----------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Dairy</b>                | <b>1,574</b> | <b>1,498</b> | <b>1,519</b> | <b>1,503</b> | <b>1,627</b> | <b>1,645</b> | <b>1,670</b> | <b>1,664</b> | <b>1,677</b> |
| Calves (4–6 months)         | 62           | 59           | 59           | 54           | 57           | 57           | 58           | 58           | 58           |
| Cows                        | 1,242        | 1,183        | 1,209        | 1,197        | 1,287        | 1,302        | 1,326        | 1,325        | 1,337        |
| Replacements (7–11 months)  | 58           | 56           | 55           | 56           | 62           | 63           | 62           | 61           | 62           |
| Replacements (12–23 months) | 212          | 201          | 196          | 196          | 222          | 223          | 224          | 220          | 220          |
| <b>Beef</b>                 | <b>4,763</b> | <b>5,419</b> | <b>5,070</b> | <b>5,007</b> | <b>4,984</b> | <b>4,873</b> | <b>4,763</b> | <b>4,722</b> | <b>4,667</b> |
| Calves (4–6 months)         | 182          | 193          | 186          | 179          | 169          | 166          | 161          | 157          | 158          |
| Bulls                       | 196          | 225          | 215          | 214          | 215          | 212          | 206          | 203          | 200          |
| Cows                        | 2,884        | 3,222        | 3,058        | 3,056        | 2,976        | 2,927        | 2,868        | 2,806        | 2,754        |
| Replacements (7–11 months)  | 69           | 85           | 74           | 80           | 75           | 74           | 76           | 78           | 81           |
| Replacements (12–23 months) | 188          | 241          | 204          | 217          | 213          | 202          | 208          | 213          | 217          |
| Steer Stockers              | 563          | 662          | 509          | 473          | 476          | 436          | 413          | 431          | 428          |
| Heifer Stockers             | 306          | 375          | 323          | 299          | 302          | 283          | 266          | 267          | 262          |
| Feedlot Cattle              | 375          | 416          | 502          | 488          | 560          | 573          | 565          | 568          | 567          |
| <b>Total</b>                | <b>6,338</b> | <b>6,917</b> | <b>6,589</b> | <b>6,510</b> | <b>6,612</b> | <b>6,518</b> | <b>6,433</b> | <b>6,386</b> | <b>6,344</b> |

Note: Totals may not sum due to independent rounding.

### Emission Estimates from Other Livestock

“Other livestock” include horses, sheep, swine, goats, American bison, and mules and asses. All livestock population data, except for American bison for years prior to 2002, were taken from the U.S. Department of Agriculture (USDA) National Agricultural Statistics Service (NASS) agricultural statistics database (USDA 2015) or earlier census data (USDA 1992, 1997). Annex 3.11 Methodology for Estimating CH<sub>4</sub> and N<sub>2</sub>O Emissions from Manure Management discusses the methods for obtaining annual average populations and disaggregating into state data where needed and provides the resulting population data for the other livestock that were used for estimating all livestock-related emissions (see Table A-199). For each animal category, the USDA publishes monthly, annual, or multi-year livestock population and production estimates. American bison estimates prior to 2002 were estimated using data from the National Bison Association (1999).

Methane emissions from sheep, goats, swine, horses, mules and asses were estimated by multiplying national population estimates by the default IPCC emission factor (IPCC 2006). For American bison the emission factor for buffalo (IPCC 2006) was used and adjusted based on the ratio of live weights of 300 kilogram for buffalo (IPCC 2006) and 1,130 pounds (513 kg) for American Bison (National Bison Association 2011) to the 0.75 power. This methodology for determining emission factors is recommended by IPCC (2006) for animals with similar digestive systems. Table A-198 shows the emission factors used for these other livestock. National enteric fermentation emissions from all livestock types are shown in Table A-199 and Table A-200. Enteric fermentation emissions from most livestock types, broken down by state, for 2014 are shown in Table A-201 and Table A-202. Livestock populations are shown in Table A-203.

**Table A-198: Emission Factors for Other Livestock (kg CH<sub>4</sub>/head/year)**

| Livestock Type  | Emission Factor |
|-----------------|-----------------|
| Swine           | 1.5             |
| Horses          | 18              |
| Sheep           | 8               |
| Goats           | 5               |
| American Bison  | 82.2            |
| Mules and Asses | 10.0            |

Source: IPCC (2006), except American Bison, as described in text.

**Table A-199: CH<sub>4</sub> Emissions from Enteric Fermentation (MMT CO<sub>2</sub> Eq.)**

| Livestock Type  | 1990         | 1995         | 2000         | 2005         | 2010         | 2011         | 2012         | 2013         | 2014         |
|-----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Beef Cattle     | 119.1        | 135.5        | 126.7        | 125.2        | 124.6        | 121.8        | 119.1        | 118.0        | 116.7        |
| Dairy Cattle    | 39.4         | 37.5         | 38.0         | 37.6         | 40.7         | 41.1         | 41.7         | 41.6         | 41.9         |
| Swine           | 2.0          | 2.2          | 2.2          | 2.3          | 2.4          | 2.5          | 2.5          | 2.5          | 2.4          |
| Horses          | 1.0          | 1.2          | 1.5          | 1.7          | 1.7          | 1.7          | 1.6          | 1.6          | 1.6          |
| Sheep           | 2.3          | 1.8          | 1.4          | 1.2          | 1.1          | 1.1          | 1.1          | 1.1          | 1.0          |
| Goats           | 0.3          | 0.3          | 0.3          | 0.4          | 0.4          | 0.3          | 0.3          | 0.3          | 0.3          |
| American Bison  | 0.1          | 0.2          | 0.4          | 0.4          | 0.4          | 0.3          | 0.3          | 0.3          | 0.3          |
| Mules and Asses | +            | +            | +            | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          |
| <b>Total</b>    | <b>164.2</b> | <b>178.7</b> | <b>170.6</b> | <b>168.9</b> | <b>171.1</b> | <b>168.7</b> | <b>166.3</b> | <b>164.5</b> | <b>164.3</b> |

Notes: Totals may not sum due to independent rounding.

+ Indicates emissions are less than 0.05 MMT CO<sub>2</sub> Eq.

**Table A-200: CH<sub>4</sub> Emissions from Enteric Fermentation (kt)**

| Livestock Type  | 1990         | 1995         | 2000         | 2005         | 2010         | 2011         | 2012         | 2013         | 2014         |
|-----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Beef Cattle     | 4,763        | 5,419        | 5,070        | 5,007        | 4,984        | 4,873        | 4,763        | 4,722        | 4,667        |
| Dairy Cattle    | 1,574        | 1,498        | 1,519        | 1,503        | 1,627        | 1,645        | 1,670        | 1,664        | 1,677        |
| Swine           | 81           | 88           | 88           | 92           | 97           | 98           | 100          | 98           | 96           |
| Horses          | 40           | 47           | 61           | 70           | 68           | 67           | 65           | 64           | 62           |
| Sheep           | 91           | 72           | 56           | 49           | 45           | 44           | 43           | 43           | 42           |
| Goats           | 13           | 12           | 12           | 14           | 14           | 14           | 13           | 13           | 12           |
| American Bison  | 4            | 9            | 16           | 17           | 15           | 14           | 13           | 13           | 12           |
| Mules and Asses | 1            | 1            | 1            | 2            | 3            | 3            | 3            | 3            | 3            |
| <b>Total</b>    | <b>6,566</b> | <b>7,146</b> | <b>6,824</b> | <b>6,755</b> | <b>6,853</b> | <b>6,757</b> | <b>6,670</b> | <b>6,619</b> | <b>6,572</b> |

Note: Totals may not sum due to independent rounding.

**Table A-201: CH<sub>4</sub> Emissions from Enteric Fermentation from Cattle (metric tons), by State, for 2014**

| State      | Dairy  |         | Dairy Repl.       | Dairy Repl.        | Bulls  | Beef   |           | Beef Repl.        | Beef Repl.         | Steer Stockers | Heifer Stockers | Feedlot | Total   |
|------------|--------|---------|-------------------|--------------------|--------|--------|-----------|-------------------|--------------------|----------------|-----------------|---------|---------|
|            | Calves | Cows    | Heif. 0-12 Months | Heif. 12-24 Months |        | Calves | Beef Cows | Heif. 0-12 Months | Heif. 12-24 Months |                |                 |         |         |
| Alabama    | 56     | 1,195   | 63                | 225                | 4,866  | 3,663  | 64,090    | 1,587             | 4,269              | 1,637          | 915             | 249     | 82,816  |
| Alaska     | 2      | 31      | 1                 | 5                  | 197    | 25     | 432       | 12                | 33                 | 12             | 3               | 1       | 755     |
| Arizona    | 1,202  | 29,450  | 1,008             | 3,581              | 2,078  | 1,034  | 17,883    | 467               | 1,249              | 8,956          | 668             | 11,844  | 79,420  |
| Arkansas   | 50     | 961     | 73                | 260                | 5,352  | 4,637  | 81,125    | 1,976             | 5,317              | 3,956          | 2,303           | 598     | 106,608 |
| California | 11,145 | 267,318 | 10,215            | 36,298             | 7,272  | 3,484  | 60,281    | 1,712             | 4,581              | 18,792         | 7,314           | 21,784  | 450,196 |
| Colorado   | 877    | 20,629  | 1,290             | 4,583              | 4,675  | 4,123  | 71,333    | 2,334             | 6,247              | 22,903         | 15,583          | 42,057  | 196,633 |
| Conn.      | 119    | 2,755   | 130               | 461                | 49     | 22     | 377       | 22                | 58                 | 55             | 15              | 7       | 4,069   |
| Delaware   | 29     | 681     | 37                | 133                | 29     | 15     | 264       | 7                 | 19                 | 55             | 18              | 7       | 1,296   |
| Florida    | 770    | 20,134  | 553               | 1,966              | 5,839  | 4,879  | 85,360    | 1,659             | 4,463              | 682            | 886             | 145     | 127,336 |
| Georgia    | 501    | 13,244  | 395               | 1,404              | 2,725  | 2,690  | 47,056    | 1,183             | 3,183              | 982            | 856             | 184     | 74,403  |
| Hawaii     | 14     | 245     | 14                | 48                 | 416    | 417    | 7,214     | 140               | 375                | 206            | 127             | 32      | 9,246   |
| Idaho      | 3,538  | 86,174  | 3,678             | 13,067             | 4,155  | 2,700  | 46,718    | 1,712             | 4,581              | 7,634          | 6,042           | 10,298  | 190,297 |
| Illinois   | 595    | 12,138  | 580               | 2,062              | 2,378  | 1,860  | 32,662    | 885               | 2,387              | 6,652          | 2,936           | 11,171  | 76,307  |
| Indiana    | 1,115  | 24,188  | 877               | 3,117              | 1,617  | 980    | 17,205    | 548               | 1,477              | 2,874          | 1,525           | 4,527   | 60,050  |
| Iowa       | 1,284  | 28,298  | 1,548             | 5,500              | 5,708  | 4,690  | 82,345    | 2,249             | 6,061              | 33,791         | 18,421          | 53,578  | 243,473 |
| Kansas     | 852    | 18,581  | 1,290             | 4,583              | 8,562  | 7,410  | 130,096   | 3,373             | 9,092              | 47,095         | 40,295          | 93,122  | 364,351 |
| Kentucky   | 426    | 9,740   | 711               | 2,528              | 6,812  | 5,336  | 93,359    | 2,164             | 5,822              | 5,457          | 3,690           | 771     | 136,815 |
| Louisiana  | 94     | 1,795   | 73                | 260                | 2,919  | 2,421  | 42,350    | 1,212             | 3,260              | 655            | 620             | 121     | 55,780  |
| Maine      | 188    | 4,331   | 245               | 870                | 146    | 59     | 1,038     | 43                | 117                | 137            | 59              | 20      | 7,254   |
| Maryland   | 313    | 7,164   | 360               | 1,280              | 390    | 205    | 3,586     | 137               | 370                | 411            | 222             | 433     | 14,872  |
| Mass.      | 75     | 1,629   | 108               | 384                | 98     | 32     | 566       | 29                | 78                 | 55             | 15              | 7       | 3,075   |
| Michigan   | 2,386  | 55,699  | 2,115             | 7,516              | 1,427  | 624    | 10,949    | 408               | 1,099              | 4,656          | 1,353           | 6,535   | 94,766  |
| Minn.      | 2,880  | 59,049  | 3,612             | 12,833             | 3,330  | 1,782  | 31,282    | 1,124             | 3,031              | 13,038         | 4,893           | 16,316  | 153,168 |
| Miss.      | 81     | 1,776   | 95                | 337                | 3,698  | 2,566  | 44,891    | 1,313             | 3,532              | 1,692          | 1,270           | 276     | 61,526  |
| Missouri   | 564    | 10,114  | 645               | 2,292              | 9,513  | 9,538  | 167,450   | 4,287             | 11,554             | 10,111         | 6,188           | 3,288   | 235,544 |
| Montana    | 88     | 1,883   | 116               | 412                | 10,389 | 8,571  | 148,292   | 6,691             | 17,908             | 6,460          | 6,710           | 2,271   | 209,790 |
| Nebraska   | 332    | 7,254   | 258               | 917                | 9,038  | 9,470  | 166,254   | 5,622             | 15,153             | 61,197         | 36,842          | 105,842 | 418,178 |
| Nevada     | 182    | 4,385   | 123               | 436                | 1,351  | 1,341  | 23,208    | 560               | 1,499              | 1,292          | 986             | 173     | 35,535  |
| N. Hamp.   | 85     | 1,957   | 101               | 358                | 49     | 16     | 283       | 14                | 39                 | 27             | 15              | 4       | 2,948   |
| N. Jersey  | 44     | 956     | 43                | 154                | 98     | 38     | 661       | 14                | 39                 | 55             | 30              | 8       | 2,139   |
| N. Mexico  | 2,022  | 50,489  | 1,634             | 5,808              | 3,636  | 2,363  | 40,891    | 1,089             | 2,915              | 2,643          | 2,226           | 453     | 116,170 |

| State     | Dairy Calves | Dairy Cows | Dairy Repl.       | Dairy Repl.        | Bulls  | Beef Calves | Beef Cows | Beef Repl.        | Beef Repl.         | Steer Stockers | Heifer Stockers | Feedlot | Total   |
|-----------|--------------|------------|-------------------|--------------------|--------|-------------|-----------|-------------------|--------------------|----------------|-----------------|---------|---------|
|           |              |            | Heif. 0-12 Months | Heif. 12-24 Months |        |             |           | Heif. 0-12 Months | Heif. 12-24 Months |                |                 |         |         |
| New York  | 3,851        | 94,740     | 5,116             | 18,177             | 1,757  | 567         | 9,909     | 651               | 1,752              | 876            | 1,185           | 1,012   | 139,591 |
| N. Car.   | 282          | 7,471      | 316               | 1,123              | 2,919  | 1,910       | 33,410    | 1,039             | 2,794              | 1,201          | 709             | 187     | 53,360  |
| N. Dakota | 106          | 2,208      | 129               | 458                | 5,232  | 4,837       | 84,921    | 2,389             | 6,440              | 7,184          | 6,044           | 1,754   | 121,704 |
| Ohio      | 1,672        | 34,747     | 1,677             | 5,958              | 2,378  | 1,536       | 26,958    | 773               | 2,084              | 5,188          | 1,727           | 6,926   | 91,623  |
| Oklahoma  | 282          | 6,112      | 292               | 1,038              | 12,164 | 9,656       | 168,931   | 4,688             | 12,614             | 24,829         | 10,038          | 12,140  | 262,784 |
| Oregon    | 776          | 17,197     | 817               | 2,904              | 3,740  | 2,996       | 51,842    | 1,634             | 4,373              | 4,698          | 3,816           | 3,163   | 97,957  |
| Penn      | 3,318        | 76,855     | 4,539             | 16,129             | 2,440  | 863         | 15,100    | 723               | 1,946              | 5,063          | 2,073           | 4,837   | 133,887 |
| R.Island  | 6            | 126        | 7                 | 26                 | 10     | 8           | 142       | 7                 | 19                 | 14             | 6               | 2       | 372     |
| S. Car.   | 100          | 2,326      | 95                | 337                | 1,362  | 909         | 15,905    | 433               | 1,164              | 327            | 354             | 63      | 23,376  |
| S. Dakota | 595          | 12,866     | 645               | 2,292              | 9,038  | 8,569       | 150,429   | 4,779             | 12,880             | 18,359         | 14,103          | 16,525  | 251,079 |
| Tenn.     | 288          | 6,673      | 474               | 1,685              | 5,839  | 4,648       | 81,313    | 1,875             | 5,045              | 3,683          | 2,510           | 130     | 114,163 |
| Texas     | 2,755        | 68,751     | 3,215             | 11,423             | 27,247 | 21,033      | 367,978   | 9,520             | 25,615             | 64,665         | 41,924          | 108,395 | 752,521 |
| Utah      | 595          | 14,057     | 627               | 2,226              | 2,389  | 1,974       | 34,159    | 1,089             | 2,915              | 2,496          | 2,194           | 1,125   | 65,847  |
| Vermont   | 826          | 19,163     | 807               | 2,867              | 293    | 65          | 1,132     | 65                | 175                | 109            | 163             | 27      | 25,694  |
| Virginia  | 582          | 14,695     | 632               | 2,247              | 3,892  | 3,427       | 59,949    | 1,659             | 4,463              | 4,638          | 2,067           | 891     | 99,143  |
| Wash.     | 1,665        | 40,560     | 1,703             | 6,050              | 1,974  | 1,243       | 21,500    | 778               | 2,082              | 4,698          | 4,134           | 8,876   | 95,263  |
| W. Virg.  | 56           | 1,133      | 72                | 256                | 1,366  | 1,031       | 18,025    | 506               | 1,362              | 1,232          | 711             | 208     | 25,959  |
| Wisconsin | 7,952        | 172,592    | 8,771             | 31,166             | 2,854  | 1,310       | 23,001    | 984               | 2,652              | 10,111         | 2,015           | 11,255  | 274,662 |
| Wyoming   | 38           | 809        | 52                | 183                | 4,155  | 4,030       | 69,725    | 2,723             | 7,288              | 4,404          | 3,371           | 3,205   | 99,984  |

**Table A-202: CH<sub>4</sub> Emissions from Enteric Fermentation from Other Livestock (metric tons), by State, for 2014**

| State          | Swine  | Horses | Sheep | Goats | American Bison | Mules and Asses | Total  |
|----------------|--------|--------|-------|-------|----------------|-----------------|--------|
| Alabama        | 158    | 978    | 97    | 208   | 17             | 116             | 1,574  |
| Alaska         | 2      | 23     | 97    | 3     | 143            | 1               | 269    |
| Arizona        | 203    | 1,833  | 1,200 | 417   | 0              | 35              | 3,688  |
| Arkansas       | 165    | 971    | 97    | 190   | 14             | 84              | 1,520  |
| California     | 165    | 2,291  | 4,720 | 719   | 88             | 65              | 8,048  |
| Colorado       | 1,043  | 1,924  | 2,920 | 145   | 726            | 63              | 6,821  |
| Connecticut    | 4      | 356    | 59    | 22    | 10             | 10              | 460    |
| Delaware       | 6      | 127    | 97    | 7     | 8              | 1               | 245    |
| Florida        | 26     | 2,182  | 97    | 249   | 7              | 96              | 2,656  |
| Georgia        | 230    | 1,208  | 97    | 334   | 17             | 88              | 1,973  |
| Hawaii         | 14     | 82     | 97    | 73    | 5              | 4               | 274    |
| Idaho          | 46     | 1,015  | 2,000 | 91    | 346            | 39              | 3,538  |
| Illinois       | 6,713  | 1,008  | 448   | 153   | 38             | 35              | 8,395  |
| Indiana        | 5,306  | 1,870  | 400   | 176   | 89             | 54              | 7,895  |
| Iowa           | 30,263 | 1,049  | 1,240 | 282   | 117            | 44              | 32,994 |
| Kansas         | 2,603  | 1,240  | 600   | 197   | 433            | 37              | 5,110  |
| Kentucky       | 488    | 2,311  | 392   | 252   | 107            | 130             | 3,679  |
| Louisiana      | 12     | 1,071  | 97    | 88    | 3              | 73              | 1,344  |
| Maine          | 7      | 215    | 59    | 33    | 18             | 4               | 335    |
| Maryland       | 32     | 501    | 97    | 41    | 25             | 12              | 708    |
| Massachusetts  | 14     | 364    | 59    | 44    | 6              | 5               | 492    |
| Michigan       | 1,568  | 1,489  | 648   | 134   | 99             | 41              | 3,979  |
| Minnesota      | 11,625 | 1,024  | 1,080 | 162   | 150            | 30              | 14,071 |
| Mississippi    | 833    | 1,009  | 97    | 110   | <1             | 88              | 2,137  |
| Missouri       | 3,938  | 1,882  | 664   | 533   | 112            | 96              | 7,224  |
| Montana        | 264    | 1,710  | 1,760 | 48    | 1,209          | 47              | 5,038  |
| Nebraska       | 4,613  | 1,149  | 608   | 112   | 2,077          | 38              | 8,596  |
| Nevada         | 3      | 434    | 640   | 126   | 4              | 6               | 1,213  |
| New Hampshire  | 5      | 158    | 59    | 26    | 26             | 3               | 277    |
| New Jersey     | 15     | 480    | 97    | 37    | 17             | 9               | 654    |
| New Mexico     | 2      | 892    | 648   | 146   | 435            | 18              | 2,142  |
| New York       | 105    | 1,660  | 600   | 175   | 54             | 37              | 2,631  |
| North Carolina | 12,225 | 1,121  | 216   | 268   | 15             | 93              | 13,937 |
| North Dakota   | 209    | 819    | 528   | 25    | 578            | 12              | 2,170  |
| Ohio           | 3,101  | 2,018  | 936   | 222   | 53             | 71              | 6,401  |
| Oklahoma       | 3,000  | 2,813  | 472   | 373   | 774            | 136             | 7,568  |
| Oregon         | 14     | 1,131  | 1,560 | 156   | 122            | 33              | 3,106  |
| Pennsylvania   | 1,691  | 2,184  | 752   | 233   | 62             | 95              | 5,016  |

| <b>State</b>   | <b>Swine</b> | <b>Horses</b> | <b>Sheep</b> | <b>Goats</b> | <b>American Bison</b> | <b>Mules and Asses</b> | <b>Total</b> |
|----------------|--------------|---------------|--------------|--------------|-----------------------|------------------------|--------------|
| Rhode Island   | 2            | 36            | 59           | 5            | 0                     | 1                      | 103          |
| South Carolina | 375          | 1,009         | 97           | 184          | 10                    | 57                     | 1,732        |
| South Dakota   | 1,853        | 1,231         | 2,160        | 94           | 2,598                 | 15                     | 7,952        |
| Tennessee      | 300          | 1,410         | 312          | 380          | 10                    | 143                    | 2,555        |
| Texas          | 1,039        | 6,815         | 5,840        | 3,872        | 310                   | 632                    | 18,508       |
| Utah           | 1,035        | 1,056         | 2,240        | 68           | 84                    | 32                     | 4,515        |
| Vermont        | 5            | 198           | 59           | 61           | 7                     | 13                     | 343          |
| Virginia       | 405          | 1,538         | 664          | 230          | 82                    | 70                     | 2,989        |
| Washington     | 46           | 982           | 440          | 124          | 60                    | 35                     | 1,688        |
| West Virginia  | 6            | 395           | 256          | 76           | 0                     | 29                     | 762          |
| Wisconsin      | 450          | 1,743         | 664          | 316          | 287                   | 58                     | 3,518        |
| Wyoming        | 126          | 1,247         | 2,840        | 48           | 688                   | 27                     | 4,975        |

## References

- Archibeque, S. (2011) Personal Communication. Shawn Archibeque, Colorado State University, Fort Collins, Colorado and staff at ICF International.
- Crutzen, P.J., I. Aselmann, and W. Seiler (1986) Methane Production by Domestic Animals, Wild Ruminants, Other Herbivores, Fauna, and Humans. *Tellus*, 38B:271-284.
- Donovan, K. (1999) Personal Communication. Kacey Donovan, University of California at Davis and staff at ICF International.
- Doren, P.E., J. F. Baker, C. R. Long and T. C. Cartwright (1989) Estimating Parameters of Growth Curves of Bulls, *J Animal Science* 67:1432-1445.
- Enns, M. (2008) Personal Communication. Dr. Mark Enns, Colorado State University and staff at ICF International.
- Galyean and Gleghorn (2001) Summary of the 2000 Texas Tech University Consulting Nutritionist Survey. Texas Tech University. Available online at: <[http://www.depts.ttu.edu/afs/burnett\\_center/progress\\_reports/bc12.pdf](http://www.depts.ttu.edu/afs/burnett_center/progress_reports/bc12.pdf)>. June 2009.
- Holstein Association (2010) *History of the Holstein Breed* (website). Available online at: <[http://www.holsteinusa.com/holstein\\_breed/breedhistory.html](http://www.holsteinusa.com/holstein_breed/breedhistory.html)>. Accessed September 2010.
- ICF (2006) *Cattle Enteric Fermentation Model: Model Documentation*. Prepared by ICF International for the Environmental Protection Agency. June 2006.
- ICF (2003) *Uncertainty Analysis of 2001 Inventory Estimates of Methane Emissions from Livestock Enteric Fermentation in the U.S.* Memorandum from ICF International to the Environmental Protection Agency. May 2003.
- IPCC (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.). Cambridge University Press. Cambridge, United Kingdom 996 pp.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- Johnson, D. (2002) Personal Communication. Don Johnson, Colorado State University, Fort Collins, and ICF International.
- Johnson, D. (1999) Personal Communication. Don Johnson, Colorado State University, Fort Collins, and David Conneely, ICF International.
- Johnson, K. (2010) Personal Communication. Kris Johnson, Washington State University, Pullman, and ICF International.
- Kebreab E., K. A. Johnson, S. L. Archibeque, D. Pape, and T. Wirth (2008) Model for estimating enteric methane emissions from United States dairy and feedlot cattle. *J. Anim. Sci.* 86: 2738-2748.
- Lippke, H., T. D. Forbes, and W. C. Ellis. (2000) Effect of supplements on growth and forage intake by stocker steers grazing wheat pasture. *J. Anim. Sci.* 78:1625-1635
- National Bison Association (2011) Handling & Carcass Info (on website). Available online at: <<http://www.bisoncentral.com/about-bison/handling-and-carcass-info>>. Accessed August 16, 2011.
- National Bison Association (1999) Total Bison Population—1999. Report provided during personal email communication with Dave Carter, Executive Director, National Bison Association July 19, 2011.
- NRC (1999) *1996 Beef NRC: Appendix Table 22*. National Research Council.
- NRC (1984) *Nutrient requirements for beef cattle (6th Ed.)*. National Academy Press, Washington, DC.
- Pinchak, W.E., D. R. Tolleson, M. McCloy, L. J. Hunt, R. J. Gill, R. J. Ansley, and S. J. Bevers (2004) Morbidity effects on productivity and profitability of stocker cattle grazing in the southern plains. *J. Anim. Sci.* 82:2773-2779.
- Platter, W. J., J. D. Tatum, K. E. Belk, J. A. Scanga, and G. C. Smith (2003) Effects of repetitive use of hormonal implants on beef carcass quality, tenderness, and consumer ratings of beef palatability. *J. Anim. Sci.* 81:984-996.

- Preston, R.L. (2010) What's The Feed Composition Value of That Cattle Feed? Beef Magazine, March 1, 2010. Available at: <<http://beefmagazine.com/nutrition/feed-composition-tables/feed-composition-value-cattle--0301>>.
- Skogerboe, T. L., L. Thompson, J. M. Cunningham, A. C. Brake, V. K. Karle (2000) The effectiveness of a single dose of doramectin pour-on in the control of gastrointestinal nematodes in yearling stocker cattle. *Vet. Parasitology* 87:173-181.
- Soliva, C.R. (2006) Report to the attention of IPCC about the data set and calculation method used to estimate methane formation from enteric fermentation of agricultural livestock population and manure management in Swiss agriculture. On behalf of the Federal Office for the Environment (FOEN), Berne, Switzerland,
- USDA (2015) *Quick Stats: Agricultural Statistics Database*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. Available online at: <<http://quickstats.nass.usda.gov/>>. Accessed August 3, 2015.
- USDA (2007) *Census of Agriculture: 2007 Census Report*. United States Department of Agriculture. Available online at: <<http://www.agcensus.usda.gov/Publications/2007/index.asp>>.
- USDA (2002) *Census of Agriculture: 2002 Census Report*. United States Department of Agriculture. Available online at: <<http://www.agcensus.usda.gov/Publications/2002/index.asp>>.
- USDA (1997) *Census of Agriculture: 1997 Census Report*. United States Department of Agriculture. Available online at: <<http://www.agcensus.usda.gov/Publications/1997/index.asp>>. Accessed July 18, 2011.
- USDA (1996) *Beef Cow/Calf Health and Productivity Audit (CHAPA): Forage Analyses from Cow/Calf Herds in 18 States*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. Available online at: <<http://www.aphis.usda.gov/vs/ceah/cahm>>. March 1996.
- USDA (1992) *Census of Agriculture: 1992 Census Report*. United States Department of Agriculture. Available online at: <<http://www.agcensus.usda.gov/Publications/1992/index.asp>>. Accessed July 18, 2011.
- USDA:APHIS:VS (2010) *Beef 2007–08, Part V: Reference of Beef Cow-calf Management Practices in the United States, 2007–08*. USDA–APHIS–VS, CEAH. Fort Collins, CO.
- USDA:APHIS:VS (2002) *Reference of 2002 Dairy Management Practices*. USDA–APHIS–VS, CEAH. Fort Collins, CO. Available online at: <<http://www.aphis.usda.gov/vs/ceah/cahm>>.
- USDA:APHIS:VS (1998) *Beef '97, Parts I-IV*. USDA–APHIS–VS, CEAH. Fort Collins, CO. Available online at: <[http://www.aphis.usda.gov/animal\\_health/nahms/beefcowcalf/index.shtml#beef97](http://www.aphis.usda.gov/animal_health/nahms/beefcowcalf/index.shtml#beef97)>
- USDA:APHIS:VS (1996) *Reference of 1996 Dairy Management Practices*. USDA–APHIS–VS, CEAH. Fort Collins, CO. Available online at: <<http://www.aphis.usda.gov/vs/ceah/cahm>>.
- USDA:APHIS:VS (1994) *Beef Cow/Calf Health and Productivity Audit*. USDA–APHIS–VS, CEAH. Fort Collins, CO. Available online at: <<http://www.aphis.usda.gov/vs/ceah/cahm>>.
- USDA:APHIS:VS (1993) *Beef Cow/Calf Health and Productivity Audit*. USDA–APHIS–VS, CEAH. Fort Collins, CO. August 1993. Available online at: <<http://www.aphis.usda.gov/vs/ceah/cahm>>.
- Vasconcelos and Galyean (2007) Nutritional recommendations of feedlot consulting nutritionists: The 2007 Texas Tech University Study. *J. Anim. Sci.* 85:2772-2781.

### 3.11. Methodology for Estimating CH<sub>4</sub> and N<sub>2</sub>O Emissions from Manure Management

The following steps were used to estimate methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) emissions from the management of livestock manure.<sup>81</sup>

#### Step 1: Livestock Population Characterization Data

Annual animal population data for 1990 through 2014 for all livestock types, except American bison, goats, horses, mules and asses were obtained from the USDA National Agricultural Statistics Service (NASS). The population data used in the emissions calculations for cattle, swine, and sheep were downloaded from the USDA NASS Quick Stats Database (USDA 2015a). Poultry population data were obtained from USDA NASS reports (USDA 1995a, 1995b, 1998, 1999, 2004a, 2004b, 2009a, 2009b, 2009c, 2009d, 2010a, 2010b, 2011a, 2011b, 2012a, 2012b, 2013a, 2013b, 2014b, 2014c, 2015b and 2015c). Goat population data for 1992, 1997, 2002, 2007, and 2012 were obtained from the Census of Agriculture (USDA 2014a), as were horse, mule and ass population data for 1987, 1992, 1997, 2002, 2007, and 2012, and American bison population for 2002, 2007, and 2012. American bison population data for 1990-1999 were obtained from the National Bison Association (1999). Additional data sources used and adjustments to these data sets are described below.

*Cattle:* For all cattle groups (cows, heifers, steers, bulls, and calves), the USDA data provide cattle inventories from January (for each state) and July (as a U. S. total only) of each year. Cattle inventories change over the course of the year, sometimes significantly, as new calves are born and as cattle are moved into feedlots and subsequently slaughtered; therefore, to develop the best estimate for the annual animal population, the populations and the individual characteristics, such as weight and weight gain, pregnancy, and lactation of each animal type were tracked in the Cattle Enteric Fermentation Model (CEFM—see section 5.1 Enteric Fermentation). For animals that have relatively static populations throughout the year, such as mature cows and bulls, the January 1 values were used. For animals that have fluctuating populations throughout the year, such as calves and growing heifers and steer, the populations are modeled based on a transition matrix that uses annual population data from USDA along with USDA data on animal births, placement into feedlots, and slaughter statistics.

*Swine:* The USDA provides quarterly data for each swine subcategory: breeding, market under 50 pounds (under 23 kg), market 50 to 119 pounds (23 to 54 kg), market 120 to 179 pounds (54 to 81 kg), and market 180 pounds and over (greater than 82 kg). The average of the quarterly data was used in the emission calculations. For states where only December inventory is reported, the December data were used directly.

*Sheep:* The USDA provides total state-level data annually for lambs and sheep. Population distribution data for lamb and sheep on feed are not available after 1993 (USDA 1994). The number of lamb and sheep on feed for 1994 through 2014 were calculated using the average of the percent of lamb and sheep on feed from 1990 through 1993. In addition, all of the sheep and lamb “on feed” are not necessarily on “feedlots;” they may be on pasture/crop residue supplemented by feed. Data for those animals on feed that are in feedlots versus pasture/crop residue were provided only for lamb in 1993. To calculate the populations of sheep and lamb in feedlots for all years, it was assumed that the percentage of sheep and lamb on feed that are in feedlots versus pasture/crop residue is the same as that for lambs in 1993 (Anderson 2000).

*Goats:* Annual goat population data by state were available for 1992, 1997, 2002, 2007, and 2012 (USDA 2014a). The data for 1992 were used for 1990 through 1992. Data for 1993 through 1996, 1998 through 2001, 2003 through 2006, 2008 through 2011, and 2013 through 2014 were extrapolated based on the 1992, 1997, 2002, 2007, and 2012 Census data.

*Horses:* Annual horse population data by state were available for 1987, 1992, 1997, 2002, 2007, and 2012 (USDA 2014a). Data for 1990 through 1991, 1993 through 1996, 1998 through 2001, 2003 through 2006, 2008 through 2011, and 2013 through 2014 were extrapolated based on the 1987, 1992, 1997, 2002, 2007, and 2012 Census data.

*Mules and Asses:* Annual mule and ass (burro and donkey) population data by state were available for 1987, 1992, 1997, 2002, 2007, and 2012 (USDA 2014a). Data for 1990 through 1991, 1993 through 1996, 1998 through 2001, 2003

---

<sup>81</sup> Note that direct N<sub>2</sub>O emissions from dung and urine spread onto fields either directly as daily spread or after it is removed from manure management systems (e.g., lagoon, pit, etc.) and from livestock dung and urine deposited on pasture, range, or paddock lands are accounted for and discussed in the Agricultural Soil Management source category within the Agriculture sector. Indirect N<sub>2</sub>O emissions dung and urine spread onto fields after it is removed from manure management systems (e.g., lagoon, pit, etc.) and from livestock dung and urine deposited on pasture, range, or paddock lands are also included in the Agricultural Soil Management source category.

through 2006, 2008 through 2011, and 2013 through 2014 were extrapolated based on the 1987, 1992, 1997, 2002, 2007, and 2012 Census data.

*American Bison:* Annual American bison population data by state were available for 2002, 2007, and 2012 (USDA 2014a). Data for 1990 through 1999 were obtained from the Bison Association (1999). Data for 2000, 2001, 2003 through 2006, 2008 through 2011, and 2013 through 2014 were extrapolated based on the Bison Association and 2002 2007, and 2012 Census data.

*Poultry:* The USDA provides population data for hens (one year old or older), pullets (hens younger than one year old), other chickens, and production (slaughter) data for broilers and turkeys (USDA 1995a, 1995b, 1998, 1999, 2004a, 2004b, 2009b, 2009c, 2009d, 2009e, 2010a, 2010b, 2011a, 2011b, 2012a, 2012b, 2013a, 2013b, 2014b, 2014c, 2015b and 2015c). All poultry population data were adjusted to account for states that report non-disclosed populations to USDA NASS. The combined populations of the states reporting non-disclosed populations are reported as “other” states. State populations for the non-disclosed states were estimated by equally distributing the population attributed to “other” states to each of the non-disclosed states.

Because only production data are available for boilers and turkeys, population data are calculated by dividing the number of animals produced by the number of production cycles per year, or the turnover rate. Based on personal communications with John Lange, an agricultural statistician with USDA NASS, the broiler turnover rate ranges from 3.4 to 5.5 over the course of the inventory (Lange 2000). For turkeys, the turnover rate ranges from 2.4 to 3.0. A summary of the livestock population characterization data used to calculate CH<sub>4</sub> and N<sub>2</sub>O emissions is presented in Table A-203.

## Step 2: Waste Characteristics Data

Methane and N<sub>2</sub>O emissions calculations are based on the following animal characteristics for each relevant livestock population:

- Volatile solids (VS) excretion rate;
- Maximum methane producing capacity (B<sub>0</sub>) for U.S. animal waste;
- Nitrogen excretion rate (Nex); and
- Typical animal mass (TAM).

Table A-204 presents a summary of the waste characteristics used in the emissions estimates. Published sources were reviewed for U.S.-specific livestock waste characterization data that would be consistent with the animal population data discussed in Step 1. The USDA’s *Agricultural Waste Management Field Handbook* (AWMFH; USDA 1996, 2008) is one of the primary sources of waste characteristics for non-cattle animal groups. Data from the 1996 and 2008 USDA AWMFH were used to estimate VS and Nex for most non-cattle animal groups across the time series of the inventory, as shown in Table A-205 (ERG 2010b and 2010c). The 1996 AWMFH data were based on measured values from U.S. farms; the 2008 AWMFH data were developed using the calculation method created by the American Society of Agricultural and Biological Engineers, which is based on U.S. animal dietary intake and performance measures. Since the values from each of the two AWMFHs result from different estimation methods and reflect changes in animal genetics and nutrition over time, both data sources were used to create a time series across the Inventory as neither value would be appropriate to use across the entire span of Inventory years. Although the AWMFH values are lower than the IPCC values, these values are more appropriate for U.S. systems because they have been calculated using U.S.-specific data. Animal-specific notes about VS and Nex are presented below:

- *Swine:* The VS and Nex data for breeding swine are from a combination of the types of animals that make up this animal group, namely gestating and farrowing swine and boars. It is assumed that a group of breeding swine is typically broken out as 80 percent gestating sows, 15 percent farrowing swine, and 5 percent boars (Safley 2000).
- *Poultry:* Due to the change in USDA reporting of hens and pullets in 2005, new nitrogen and VS excretion rates were calculated for the combined population of hens and pullets; a weighted average rate was calculated based on hen and pullet population data from 1990 to 2004.
- *Goats, Sheep, Horses, Mules and Asses:* In cases where data were not available in the USDA documents, data from the American Society of Agricultural Engineers, Standard D384.1 (ASAE 1998) or the *2006 IPCC Guidelines* were used as a supplement.

The method for calculating VS excretion and Nex for cattle (including American bison, beef and dairy cows, bulls, heifers, and steers) is based on the relationship between animal performance characteristics such as diet, lactation, and weight gain and energy utilization. The method used is outlined by the *2006 IPCC Guidelines* Tier II methodology, and is modeled

using the CEFM as described in the enteric fermentation portion of the inventory (documented in Moffroid and Pape 2013) in order to take advantage of the detailed diet and animal performance data assembled as part of the Tier II analysis for cattle. For American bison, VS and Nex were assumed to be the same as beef NOF bulls.

The VS content of manure is the fraction of the diet consumed by cattle that is not digested and thus excreted as fecal material; fecal material combined with urinary excretions constitutes manure. The CEFM uses the input of digestible energy (DE) and the energy requirements of cattle to estimate gross energy (GE) intake and enteric CH<sub>4</sub> emissions. GE and DE are used to calculate the indigestible energy per animal as gross energy minus digestible energy plus the amount of gross energy for urinary energy excretion per animal (2 or 4 percent). This value is then converted to VS production per animal using the typical conversion of dietary gross energy to dry organic matter of 18.45 MJ/kg, after subtracting out the ash content of manure. The current equation recommended by the 2006 IPCC Guidelines is:

$$\text{VS production (kg)} = [(GE - DE) + (UE \times GE)] \times \frac{1 - ASH}{18.45}$$

where,

|           |  |
|-----------|--|
| GE        | = Gross energy intake (MJ)   |
| DE        | = Digestible energy (MJ)   |
| (UE × GE) | = Urinary energy expressed as fraction of GE, assumed to be 0.04 except for feedlots which are reduced 0.02 as a result of the high grain content of their diet.                           |
| ASH       | = Ash content of manure calculated as a fraction of the dry matter feed intake (assumed to be 0.08).   |
| 18.45     | = Conversion factor for dietary GE per kg of dry matter (MJ per kg). This value is relatively constant across a wide range of forage and grain-based feeds commonly consumed by livestock. |

Total nitrogen ingestion in cattle is determined by dietary protein intake. When feed intake of protein exceeds the nutrient requirements of the animal, the excess nitrogen is excreted, primarily through the urine. To calculate the nitrogen excreted by each animal type, the CEFM utilizes the energy balance calculations recommended by the IPCC (2006) for gross energy and the energy required for growth along with inputs of weight gain, milk production, and the percent of crude protein in the diets. The total nitrogen excreted is measured in the CEFM as nitrogen consumed minus nitrogen retained by the animal for growth and in milk. The basic equation for calculating Nex is shown below, followed by the equations for each of the constituent parts.

$$N_{\text{excreted}} = N_{\text{consumed}} - (N_{\text{growth}} + N_{\text{milk}})$$

where,

|                       |   |
|-----------------------|---|
| N <sub>excreted</sub> | = Daily N excreted per animal, kg per animal per day.               |
| N <sub>consumed</sub> | = Daily N intake per animal, kg per animal per day                  |
| N <sub>growth</sub>   | = Nitrogen retained by the animal for growth, kg per animal per day |
| N <sub>milk</sub>     | = Nitrogen retained in milk, kg per animal per day                  |

The equation for N consumed is based on the 2006 IPCC Guidelines, and is estimated as:

$$N_{\text{consumed}} = \left[ \frac{GE}{18.45} * \left( \frac{CP\%}{6.25} \right) \right]$$

where,

|                       |  |
|-----------------------|--|
| N <sub>consumed</sub> | = Daily N intake per animal, kg per animal per day                           |
| GE                    | = Gross energy intake, as calculated in the CEFM, MJ per animal per day      |
| 18.45                 | = Conversion factor for dietary GE per kg of dry matter, MJ per kg.          |
| CP%                   | = Percent crude protein in diet, input into the CEFM                         |
| 6.25                  | = Conversion from kg of dietary protein to kg of dietary N, kg feed per kg N |

The portion of consumed N that is retained as product equals the nitrogen required for weight gain plus that in milk. The nitrogen retained in body weight gain by stockers, replacements, or feedlot animals is calculated using the net energy for growth (NEg), weight gain (WG), and other conversion factors and constants. The equation matches current 2006 IPCC Guidelines recommendations, and is as follows:

$$N_{growth} = \frac{\left\{ WG * \left[ 268 - \frac{(7.03 * NEg)}{WG} \right] \right\}}{\frac{1000}{6.25}}$$

where,

|              |   |
|--------------|---|
| $N_{growth}$ | = Nitrogen retained by the animal for growth, kg per animal per day                     |
| WG           | = Daily weight gain of the animal, as input into the CEFM transition matrix, kg per day |
| 268          | = Constant from 2006 IPCC Guidelines  |
| 7.03         | = Constant from 2006 IPCC Guidelines  |
| NEg          | = Net energy required for growth, as calculated in the CEFM, MJ per animal per day      |
| 1,000        | = Conversion from grams to kilograms  |
| 6.25         | = Conversion from kg of dietary protein to kg of dietary N, kg feed per kg N            |

The N content of milk produced also currently matches the 2006 IPCC Guidelines, and is calculated using milk production and percent protein, along with conversion factors. Milk N retained as product is calculated using the following equation:

$$N_{milk} = \frac{milk * \left( \frac{pr\%}{100} \right)}{6.38}$$

where,

|            |  |
|------------|--|
| $N_{milk}$ | = Nitrogen retained in milk, kg per animal per day   |
| milk       | = Milk production, kg per day  |
| pr%        | = Percent protein in milk, estimated from the fat content as $1.9 + 0.4 * \%Fat$<br>(Fat assumed to be 4%) |
| 100        | = Conversion from percent to value (e.g., 4% to 0.04)  |
| 6.38       | = Conversion from kg Protein to kg N   |

The VS and N equations above were used to calculate VS and Nex rates for each state, animal type (heifers and steer on feed, heifers and steer not on feed, bulls and American bison), and year. Table A-206 presents the state-specific VS and Nex production rates used for cattle in 2014.

### Step 3: Waste Management System Usage Data

Table A-207 summarizes 2014 manure distribution data among waste management systems (WMS) at beef feedlots, dairies, dairy heifer facilities, and swine, layer, broiler, and turkey operations. Manure from the remaining animal types (beef cattle not on feed, American bison, goats, horses, mules and asses and sheep) is managed on pasture, range, or paddocks, on drylot, or with solids storage systems. Additional information on the development of the manure distribution estimates for each animal type is presented below. Definitions of each WMS type are presented in Table A-208.

*Beef Cattle, Dairy Heifers and American Bison:* The beef feedlot and dairy heifer WMS data were developed using information from EPA's Office of Water's engineering cost analyses conducted to support the development of effluent limitations guidelines for Concentrated Animal Feeding Operations (EPA 2002b). Based on EPA site visits and state contacts supporting this work and additional personal communication with the national USDA office to estimate the percent of beef steers and heifers in feedlots (Milton 2000), feedlot manure is almost exclusively managed in drylots. Therefore, for these animal groups, the percent of manure deposited in drylots is assumed to be 100 percent. In addition, there is a

small amount of manure contained in runoff, which may or may not be collected in runoff ponds. Using EPA and USDA data and expert opinions (documented in ERG 2000a), the runoff from feedlots was calculated by region in *Calculations: Percent Distribution of Manure for Waste Management Systems* and was used to estimate the percentage of manure managed in runoff ponds in addition to drylots; this percentage ranges from 0.4 to 1.3 percent (ERG 2000a). The percentage of manure generating emissions from beef feedlots is therefore greater than 100 percent. The remaining population categories of beef cattle outside of feedlots are managed through pasture, range, or paddock systems, which are utilized for the majority of the population of beef cattle in the country. American bison WMS data were assumed to be the same as beef cattle NOF.

*Dairy Cows:* The WMS data for dairy cows were developed using data from the Census of Agriculture, EPA's Office of Water, USDA, and the expert sources noted below. Farm-size distribution data are reported in the 1992, 1997, 2002, and 2007 Census of Agriculture (USDA 2014a). It was assumed that the Census data provided for 1992 were the same as that for 1990 and 1991, and data provided for 2007 were the same as that for 2008 through 2014. Data for 1993 through 1996, 1998 through 2001, and 2003 through 2006, and 2008 through 2014 were extrapolated using the 1992, 1997, 2002, and 2007 data. The percent of waste by system was estimated using the USDA data broken out by geographic region and farm size.

Based on EPA site visits and the expert opinion of state contacts, manure from dairy cows at medium (200 through 700 head) and large (greater than 700 head) operations are managed using either flush systems or scrape/slurry systems. In addition, they may have a solids separator in place prior to their storage component. Estimates of the percent of farms that use each type of system (by geographic region) were developed by EPA's Office of Water, and were used to estimate the percent of waste managed in lagoons (flush systems), liquid/slurry systems (scrape systems), and solid storage (separated solids) (EPA 2002b).

Manure management system data for small (fewer than 200 head) dairies were obtained from USDA's Animal and Plant Health Inspection Service (APHIS)'s National Animal Health Monitoring System (Ott 2000). These data are based on a statistical sample of farms in the 20 U.S. states with the most dairy cows. Small operations are more likely to use liquid/slurry and solid storage management systems than anaerobic lagoon systems. The reported manure management systems were deep pit, liquid/slurry (includes slurry tank, slurry earth-basin, and aerated lagoon), anaerobic lagoon, and solid storage (includes manure pack, outside storage, and inside storage).

Data regarding the use of daily spread and pasture, range, or paddock systems for dairy cattle were obtained from personal communications with personnel from several organizations. These organizations include state NRCS offices, state extension services, state universities, USDA NASS, and other experts (Deal 2000, Johnson 2000, Miller 2000, Stettler 2000, Sweeten 2000, and Wright 2000). Contacts at Cornell University provided survey data on dairy manure management practices in New York (Poe et al. 1999). Census of Agriculture population data for 1992, 1997, 2002, and 2007 (USDA 2014a) were used in conjunction with the state data obtained from personal communications to determine regional percentages of total dairy cattle and dairy waste that are managed using these systems. These percentages were applied to the total annual dairy cow and heifer state population data for 1990 through 2014, which were obtained from the USDA NASS (USDA 2015a).

Of the dairies using systems other than daily spread and pasture, range, or paddock systems, some dairies reported using more than one type of manure management system. Due to limitations in how USDA APHIS collects the manure management data, the total percent of systems for a region and farm size is greater than 100 percent. However, manure is typically partitioned to use only one manure management system, rather than transferred between several different systems. Emissions estimates are only calculated for the final manure management system used for each portion of manure. To avoid double counting emissions, the reported percentages of systems in use were adjusted to equal a total of 100 percent using the same distribution of systems. For example, if USDA reported that 65 percent of dairies use deep pits to manage manure and 55 percent of dairies use anaerobic lagoons to manage manure, it was assumed that 54 percent (i.e., 65 percent divided by 120 percent) of the manure is managed with deep pits and 46 percent (i.e., 55 percent divided by 120 percent) of the manure is managed with anaerobic lagoons (ERG 2000a).

Finally, the percentage of manure managed with anaerobic digestion (AD) systems with methane capture and combustion was added to the WMS distributions. AD system data were obtained from EPA's AgSTAR Program's project database (EPA 2012). This database includes basic information for AD systems in the United States, based on publically available data and data submitted by farm operators, project developers, financiers, and others involved in the development of farm AD projects.

*Swine:* The distribution of manure managed in each WMS was estimated using data from a USDA APHIS report and EPA's Office of Water site visits (Bush 1998, ERG 2000a). The USDA APHIS data are based on a statistical sample of farms in the 16 U.S. states with the most hogs. For operations with less than 200 head, manure management system data

were obtained from USDA APHIS (Bush 1998), it was assumed that those operations use pasture, range, or paddock systems. For swine operations with greater than 200 head, the percent of waste managed in each system was estimated using the EPA and USDA data broken out by geographic region and farm size. Farm-size distribution data reported in the 1992, 1997, 2002, and 2007 Census of Agriculture (USDA 2014a) were used to determine the percentage of all swine utilizing the various manure management systems. It was assumed that the swine farm size data provided for 1992 were the same as that for 1990 and 1991, and data provided for 2007 were the same as that for 2008 through 2014. Data for 1993 through 1996, 1998 through 2001, and 2003 through 2006 were extrapolated using the 1992, 1997, 2002, and 2007 data. The manure management systems reported in the census were deep pit, liquid/slurry (includes above- and below-ground slurry), anaerobic lagoon, and solid storage (includes solids separated from liquids).

Some swine operations reported using more than one management system; therefore, the total percent of systems reported by USDA for a region and farm size was greater than 100 percent. Typically, this means that a portion of the manure at a swine operation is handled in one system (e.g., liquid system), and a separate portion of the manure is handled in another system (e.g., dry system). However, it is unlikely that the same manure is moved from one system to another, which could result in increased emissions, so reported systems data were normalized to 100 percent for incorporation into the WMS distribution, using the same method as described above for dairy operations. As with dairy, AD WMS were added to the WMS distribution based on data from EPA's AgSTAR database (EPA 2012).

*Sheep:* WMS data for sheep were obtained from USDA NASS sheep report for years 1990 through 1993 (USDA 1994). Data for 2001 are obtained from USDA APHIS's national sheep report (USDA, APHIS 2003). The USDA APHIS data are based on a statistical sampled of farms in the 22 U.S. states with the most sheep. The data for years 1994-2000 are calculated assuming a linear progression from 1993 to 2001. Due to lack of additional data, data for years 2002 and beyond are assumed to be the same as 2001. Based on expert opinion, it was assumed that all sheep manure not deposited in feedlots was deposited on pasture, range, or paddock lands (Anderson 2000).

*Goats, Horses, and Mules and Asses:* WMS data for 1990 to 2014 were obtained from Appendix H of *Global Methane Emissions from Livestock and Poultry Manure* (EPA 1992). This report presents state WMS usage in percentages for the major animal types in the United States, based on information obtained from extension service personnel in each state. It was assumed that all manure not deposited in pasture, range, or paddock lands was managed in dry systems. For mules and asses, the WMS was assumed to be the same as horses.

*Poultry—Hens (one year old or older), Pullets (hens less than one year old), and Other Chickens:* WMS data for 1992 were obtained from *Global Methane Emissions from Livestock and Poultry Manure* (EPA 1992). These data were also used to represent 1990 and 1991. The percentage of layer operations using a shallow pit flush house with anaerobic lagoon or high-rise house without bedding was obtained for 1999 from a United Egg Producers voluntary survey (UEP 1999). These data were augmented for key poultry states (AL, AR, CA, FL, GA, IA, IN, MN, MO, NC, NE, OH, PA, TX, and WA) with USDA data (USDA, APHIS 2000). It was assumed that the change in system usage between 1990 and 1999 is proportionally distributed among those years of the inventory. It was also assumed that system usage in 2000 through 2014 was equal to that estimated for 1999. Data collected for EPA's Office of Water, including information collected during site visits (EPA 2002b), were used to estimate the distribution of waste by management system and animal type. As with dairy and swine, using information about AD WMS from EPA's AgSTAR database (EPA 2012), AD was added to the WMS distribution for poultry operations.

*Poultry—Broilers and Turkeys:* The percentage of turkeys and broilers on pasture was obtained from the Office of Air and Radiation's *Global Methane Emissions from Livestock and Poultry Manure* (EPA 1992). It was assumed that one percent of poultry waste is deposited in pastures, ranges, and paddocks (EPA 1992). The remainder of waste is assumed to be deposited in operations with bedding management. As with dairy, swine, and other poultry, AD systems were added to the WMS distributions based on information from EPA's AgSTAR database (EPA 2012).

#### **Step 4: Emission Factor Calculations**

Methane conversion factors (MCFs) and N<sub>2</sub>O emission factors (EFs) used in the emission calculations were determined using the methodologies presented below.

##### ***Methane Conversion Factors (MCFs)***

Climate-based IPCC default MCFs (IPCC 2006) were used for all dry systems; these factors are presented in Table A-209. A U.S.-specific methodology was used to develop MCFs for all lagoon and liquid systems.

For animal waste managed in dry systems, the appropriate IPCC default MCF was applied based on annual average temperature data. The average county and state temperature data were obtained from the National Climate Data Center

(NOAA 2014) and each state and year in the inventory was assigned a climate classification of cool, temperate or warm. Although there are some specific locations in the United States that may be included in the warm climate category, no aggregated state-level annual average temperatures are included in this category. In addition, some counties in a particular state may be included in the cool climate category, although the aggregated state-level annual average temperature may be included in the temperate category. Although considering the temperatures at a state level instead of a county level may be causing some specific locations to be classified into an inappropriate climate category, using the state level annual average temperature provides an estimate that is appropriate for calculating the national average.

For anaerobic lagoons and other liquid systems a climate-based approach based on the van't Hoff-Arrhenius equation was developed to estimate MCFs that reflects the seasonal changes in temperatures, and also accounts for long-term retention time. This approach is consistent with the latest guidelines from IPCC (IPCC 2006). The van't Hoff-Arrhenius equation, with a base temperature of 30°C, is shown in the following equation (Safley and Westerman 1990):

$$f = \exp\left[\frac{E(T_2 - T_1)}{RT_1T_2}\right]$$

where,

|       |  |
|-------|--|
| $f$   | = van't Hoff-Arrhenius $f$ factor, the proportion of VS that are biologically available for conversion to CH <sub>4</sub> based on the temperature of the system |
| $T_1$ | = 303.15K  |
| $T_2$ | = Ambient temperature (K) for climate zone (in this case, a weighted value for each state)   |
| $E$   | = Activation energy constant (15,175 cal/mol)  |
| $R$   | = Ideal gas constant (1.987 cal/K mol)   |

For those animal populations using liquid manure management systems or manure runoff ponds (i.e., dairy cow, dairy heifer, layers, beef in feedlots, and swine) monthly average state temperatures were based on the counties where the specific animal population resides (i.e., the temperatures were weighted based on the percent of animals located in each county). County population data were calculated from state-level population data from NASS and county-state distribution data from the 1992, 1997, 2002, and 2007 Census data (USDA 2014a). County population distribution data for 1990 and 1991 were assumed to be the same as 1992; county population distribution data for 1993 through 1996 were extrapolated based on 1992 and 1997 data; county population data for 1998 through 2001 were extrapolated based on 1997 and 2002 data; county population data for 2003 through 2006 were extrapolated based on 2002 and 2007 data; and county population data for 2008 to 2014 were assumed to be the same as 2007.

Annual MCFs for liquid systems are calculated as follows for each animal type, state, and year of the inventory:

- The weighted-average temperature for a state is calculated using the county population estimates and average monthly temperature in each county. Monthly temperatures are used to calculate a monthly van't Hoff-Arrhenius  $f$  factor, using the equation presented above. A minimum temperature of 5°C is used for uncovered anaerobic lagoons and 7.5°C is used for liquid/slurry and deep pit systems due to the biological activity in the lagoon which keeps the temperature above freezing.
- Monthly production of VS added to the system is estimated based on the animal type, number of animals present, and the volatile solids excretion rate of the animals.
- For lagoon systems, the calculation of methane includes a management and design practices (MDP) factor. This factor, equal to 0.8, was developed based on model comparisons to empirical CH<sub>4</sub> measurement data from anaerobic lagoon systems in the United States (ERG 2001). The MDP factor represents management and design factors which cause a system to operate at a less than optimal level.
- For all systems other than anaerobic lagoons, the amount of VS available for conversion to CH<sub>4</sub> each month is assumed to be equal to the amount of VS produced during the month (from Step 3). For anaerobic lagoons, the amount of VS available also includes VS that may remain in the system from previous months.
- The amount of VS consumed during the month is equal to the amount available for conversion multiplied by the  $f$  factor.

- For anaerobic lagoons, the amount of VS carried over from one month to the next is equal to the amount available for conversion minus the amount consumed. Lagoons are also modeled to have a solids clean-out once per year, occurring in the month of October.
- The estimated amount of CH<sub>4</sub> generated during the month is equal to the monthly VS consumed multiplied by the maximum CH<sub>4</sub> potential of the waste (B<sub>o</sub>).

The annual MCF is then calculated as:

$$MCF_{\text{annual}} = \frac{\text{CH}_4 \text{ generated}_{\text{annual}}}{\text{VS produced}_{\text{annual}} \times B_o}$$

where,

MCF<sub>annual</sub> = Methane conversion factor  
 VS produced<sub>annual</sub> = Volatile solids excreted annually  
 B<sub>o</sub> = Maximum CH<sub>4</sub> producing potential of the waste

In order to account for the carry-over of VS from one year to the next, it is assumed that a portion of the VS from the previous year are available in the lagoon system in the next year. For example, the VS from October, November, and December of 2005 are available in the lagoon system starting January of 2006 in the MCF calculation for lagoons in 2006. Following this procedure, the resulting MCF for lagoons accounts for temperature variation throughout the year, residual VS in a system (carry-over), and management and design practices that may reduce the VS available for conversion to CH<sub>4</sub>. It is assumed that liquid-slurry systems have a retention time less than 30 days, so the liquid-slurry MCF calculation doesn't reflect the VS carry-over.

The liquid system MCFs are presented in Table A-210 by state, WMS, and animal group for 2014.

#### **Nitrous Oxide Emission Factors**

Direct N<sub>2</sub>O EFs for manure management systems (kg N<sub>2</sub>O-N/kg excreted N) were set equal to the most recent default IPCC factors (IPCC 2006), presented in Table A-211.

Indirect N<sub>2</sub>O EFs account for two fractions of nitrogen losses: volatilization of ammonia (NH<sub>3</sub>) and NO<sub>x</sub> (Frac<sub>gas</sub>) and runoff/leaching (Frac<sub>runoff/leach</sub>). IPCC default indirect N<sub>2</sub>O EFs were used to estimate indirect N<sub>2</sub>O emissions. These factors are 0.010 kg N<sub>2</sub>O-N/kg N for volatilization and 0.0075 kg N<sub>2</sub>O/kg N for runoff/leaching.

Country-specific estimates of N losses were developed for Frac<sub>gas</sub> and Frac<sub>runoff/leach</sub> for the United States. The vast majority of volatilization losses are NH<sub>3</sub>. Although there are also some small losses of NO<sub>x</sub>, no quantified estimates were available for use and those losses are believed to be small (about 1 percent) in comparison to the NH<sub>3</sub> losses. Therefore, Frac<sub>gas</sub> values were based on WMS-specific volatilization values estimated from U.S. EPA's *National Emission Inventory - Ammonia Emissions from Animal Agriculture Operations* (EPA 2005). To estimate Frac<sub>runoff/leach</sub>, data from EPA's Office of Water were used that estimate the amount of runoff from beef, dairy, and heifer operations in five geographic regions of the country (EPA 2002b). These estimates were used to develop U.S. runoff factors by animal type, WMS, and region. Nitrogen losses from leaching are believed to be small in comparison to the runoff losses and there are a lack of data to quantify these losses. Therefore, leaching losses were assumed to be zero and Frac<sub>runoff/leach</sub> was set equal to the runoff loss factor. Nitrogen losses from volatilization and runoff/leaching are presented in Table A-212.

#### **Step 5: CH<sub>4</sub> Emission Calculations**

To calculate CH<sub>4</sub> emissions for animals other than cattle, first the amount of VS excreted in manure that is managed in each WMS was estimated:

$$VS \text{ excreted}_{\text{State, Animal, WMS}} = \text{Population}_{\text{State, Animal}} \times \frac{TAM}{1000} \times VS \times WMS \times 365.25$$

where,

VS excreted<sub>State, Animal, WMS</sub> = Amount of VS excreted in manure managed in each WMS for each animal type (kg/yr)  
 Population<sub>State, Animal</sub> = Annual average state animal population by animal type (head)

|        |   |
|--------|---|
| TAM    | = Typical animal mass (kg)  |
| VS     | = Volatile solids production rate (kg VS/1000 kg animal mass/day)         |
| WMS    | = Distribution of manure by WMS for each animal type in a state (percent) |
| 365.25 | = Days per year   |

Using the CEFM VS data for cattle, the amount of VS excreted in manure that is managed in each WMS was estimated using the following equation:

$$\text{VS excreted}_{\text{State, Animal, WMS}} = \text{Population}_{\text{State, Animal}} \times \text{VS} \times \text{WMS}$$

where,

|   |  |
|---|--|
| VS excreted <sub>State, Animal, WMS</sub> | = Amount of VS excreted in manure managed in each WMS for each animal type (kg/yr) |
| Population <sub>State, Animal</sub>       | = Annual average state animal population by animal type (head)                     |
| VS  | = Volatile solids production rate (kg VS/animal/year)                              |
| WMS                                       | = Distribution of manure by WMS for each animal type in a state (percent)          |

For all animals, the estimated amount of VS excreted into a WMS was used to calculate CH<sub>4</sub> emissions using the following equation:

$$\text{CH}_4 = \sum_{\text{State, Animal, WMS}} (\text{VS excreted}_{\text{State, Animal, WMS}} \times B_o \times \text{MCF} \times 0.662)$$

where,

|                                   |  |
|-----------------------------------|--|
| CH <sub>4</sub>                   | = CH <sub>4</sub> emissions (kg CH <sub>4</sub> /yr)                                 |
| VS excreted <sub>WMS, State</sub> | = Amount of VS excreted in manure managed in each WMS (kg/yr)                        |
| B <sub>o</sub>                    | = Maximum CH <sub>4</sub> producing capacity (m <sup>3</sup> CH <sub>4</sub> /kg VS) |
| MCF <sub>animal, state, WMS</sub> | = MCF for the animal group, state and WMS (percent)                                  |
| 0.662                             | = Density of methane at 25° C (kg CH <sub>4</sub> /m <sup>3</sup> CH <sub>4</sub> )  |

A calculation was developed to estimate the amount of CH<sub>4</sub> emitted from AD systems utilizing CH<sub>4</sub> capture and combustion technology. First, AD systems were assumed to produce 90 percent of the maximum CH<sub>4</sub> producing capacity (B<sub>o</sub>) of the manure. This value is applied for all climate regions and AD system types. However, this is a conservative assumption as the actual amount of CH<sub>4</sub> produced by each AD system is very variable and will change based on operational and climate conditions and an assumption of 90 percent is likely overestimating CH<sub>4</sub> production from some systems and underestimating CH<sub>4</sub> production in other systems. The CH<sub>4</sub> production of AD systems is calculated using the equation below:

$$\text{CH}_4 \text{ Production}_{\text{AD}_{\text{ADSystem}}} = \text{Production}_{\text{AD}_{\text{ADSystem}}} \times \frac{\text{TAM}}{1000} \times \text{VS} \times B_o \times 0.662 \times 365.25 \times 0.90$$

where,

|  |   |
|--|---|
| CH <sub>4</sub> Production <sub>AD<sub>AD system</sub></sub> | = CH <sub>4</sub> production from a particular AD system, (kg/yr)                           |
| Population <sub>AD<sub>state</sub></sub>                     | = Number of animals on a particular AD system   |
| VS   | = Volatile solids production rate (kg VS/1000 kg animal mass-day)                           |
| TAM  | = Typical Animal Mass (kg/head)   |
| B <sub>o</sub>   | = Maximum CH <sub>4</sub> producing capacity (CH <sub>4</sub> m <sup>3</sup> /kg VS)        |
| 0.662  | = Density of CH <sub>4</sub> at 25° C (kg CH <sub>4</sub> /m <sup>3</sup> CH <sub>4</sub> ) |
| 365.25   | = Days/year   |
| 0.90   | = CH <sub>4</sub> production factor for AD systems  |

The total amount of CH<sub>4</sub> produced by AD is calculated only as a means to estimate the emissions from AD; i.e., only the estimated amount of CH<sub>4</sub> actually entering the atmosphere from AD is reported in the inventory. The emissions to the atmosphere from AD are a result of leakage from the system (e.g. from the cover, piping, tank, etc.) and incomplete

combustion and are calculated using the collection efficiency (CE) and destruction efficiency (DE) of the AD system. The three primary types of AD systems in the United States are covered lagoons, complete mix and plug flow systems. The CE of covered lagoon systems was assumed to be 75 percent, and the CE of complete mix and plug flow AD systems was assumed to be 99 percent (EPA 2008). The CH<sub>4</sub> DE from flaring or burning in an engine was assumed to be 98 percent; therefore, the amount of CH<sub>4</sub> that would not be flared or combusted was assumed to be 2 percent (EPA 2008). The amount of CH<sub>4</sub> produced by systems with AD was calculated with the following equation:

$$\text{CH}_4 \text{ Emissions AD} = \sum_{\text{State, Animal, AD Systems}} \left( \left[ \text{CH}_4 \text{ Production AD}_{\text{AD system}} \times \text{CE}_{\text{AD system}} \times (1 - \text{DE}) \right] + \left[ \text{CH}_4 \text{ Production AD}_{\text{AD system}} \times (1 - \text{CE}_{\text{AD system}}) \right] \right)$$

where,

|  |   |  |
|--|---|--|
| CH <sub>4</sub> Emissions AD                       | = | CH <sub>4</sub> emissions from AD systems, (kg/yr)               |
| CH <sub>4</sub> Production AD <sub>AD system</sub> | = | CH <sub>4</sub> production from a particular AD system, (kg/yr)  |
| CE <sub>AD system</sub>                            | = | Collection efficiency of the AD system, varies by AD system type |
| DE   | = | Destruction efficiency of the AD system, 0.98 for all systems    |

### Step 6: N<sub>2</sub>O Emission Calculations

Total N<sub>2</sub>O emissions from manure management systems were calculated by summing direct and indirect N<sub>2</sub>O emissions. The first step in estimating direct and indirect N<sub>2</sub>O emissions was calculating the amount of N excreted in manure and managed in each WMS. For calves and animals other than cattle the following equation was used:

$$\text{N excreted}_{\text{State, Animal, WMS}} = \text{Population}_{\text{State, Animal}} \times \text{WMS} \times \frac{\text{TAM}}{1000} \times \text{Nex} \times 365.25$$

where,

|  |   |   |
|--|---|---|
| N excreted <sub>State, Animal, WMS</sub> | = | Amount of N excreted in manure managed in each WMS for each animal type (kg/yr)             |
| Population <sub>state</sub>              | = | Annual average state animal population by animal type (head)                                |
| WMS                                      | = | Distribution of manure by waste management system for each animal type in a state (percent) |
| TAM                                      | = | Typical animal mass (kg)  |
| Nex                                      | = | Total Kjeldahl nitrogen excretion rate (kg N/1000 kg animal mass/day)                       |
| 365.25                                   | = | Days per year   |

Using the CEFM Nex data for cattle other than calves, the amount of N excreted was calculated using the following equation:

$$\text{N excreted}_{\text{State, Animal, WMS}} = \text{Population}_{\text{State, Animal}} \times \text{WMS} \times \text{Nex}$$

where,

|  |   |   |
|--|---|---|
| N excreted <sub>State, Animal, WMS</sub> | = | Amount of N excreted in manure managed in each WMS for each animal type (kg/yr)             |
| Population <sub>state</sub>              | = | Annual average state animal population by animal type (head)                                |
| WMS                                      | = | Distribution of manure by waste management system for each animal type in a state (percent) |
| Nex                                      | = | Total Kjeldahl N excretion rate (kg N/animal/year)  |

For all animals, direct N<sub>2</sub>O emissions were calculated as follows:

$$\text{Direct N}_2\text{O} = \sum_{\text{State, Animal, WMS}} \left( \text{N excreted}_{\text{State, Animal, WMS}} \times \text{EF}_{\text{WMS}} \times \frac{44}{28} \right)$$

where,

|                         |   |  |
|-------------------------|---|--|
| Direct N <sub>2</sub> O | = | Direct N <sub>2</sub> O emissions (kg N <sub>2</sub> O/yr) |
|-------------------------|---|--|

$N_{\text{excreted}}_{\text{State, Animal, WMS}}$  = Amount of N excreted in manure managed in each WMS for each animal type (kg/yr)  
 $EF_{\text{WMS}}$  = Direct N<sub>2</sub>O emission factor from IPCC guidelines (kg N<sub>2</sub>O-N /kg N)  
 $44/28$  = Conversion factor of N<sub>2</sub>O-N to N<sub>2</sub>O

Indirect N<sub>2</sub>O emissions were calculated for all animals with the following equation:

$$\text{Indirect N}_2\text{O} = \sum_{\text{State, Animal, WMS}} \left[ \begin{aligned} & \left[ N_{\text{excreted}}_{\text{State, Animal, WMS}} \times \frac{\text{Frac}_{\text{gas, WMS}}}{100} \times EF_{\text{volatilization}} \times \frac{44}{28} \right] + \\ & \left[ N_{\text{excreted}}_{\text{State, Animal, WMS}} \times \frac{\text{Frac}_{\text{runoff/leach, WMS}}}{100} \times EF_{\text{runoff/leach}} \times \frac{44}{28} \right] \end{aligned} \right]$$

where,

$\text{Indirect N}_2\text{O}$  = Indirect N<sub>2</sub>O emissions (kg N<sub>2</sub>O/yr)  
 $N_{\text{excreted}}_{\text{State, Animal, WMS}}$  = Amount of N excreted in manure managed in each WMS for each animal type (kg/yr)  
 $\text{Frac}_{\text{gas, WMS}}$  = Nitrogen lost through volatilization in each WMS  
 $\text{Frac}_{\text{runoff/leach, WMS}}$  = Nitrogen lost through runoff and leaching in each WMS (data were not available for leaching so the value reflects only runoff)  
 $EF_{\text{volatilization}}$  = Emission factor for volatilization (0.010 kg N<sub>2</sub>O-N/kg N)  
 $EF_{\text{runoff/leach}}$  = Emission factor for runoff/leaching (0.0075 kg N<sub>2</sub>O-N/kg N)  
 $44/28$  = Conversion factor of N<sub>2</sub>O-N to N<sub>2</sub>O

Emission estimates of CH<sub>4</sub> and N<sub>2</sub>O by animal type are presented for all years of the inventory in Table A-213 and Table A-214 respectively. Emission estimates for 2014 are presented by animal type and state in Table A-215 and Table A-216 respectively.

**Table A-203: Livestock Population (1,000 Head)**

| Animal Type              | 1990      | 1995      | 2001      | 2002      | 2003      | 2004      | 2005      | 2006      | 2007      | 2008      | 2009      | 2010      | 2011      | 2012      | 2013      | 2014      |
|--------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| Dairy Cattle             | 19,512    | 18,681    | 17,927    | 17,833    | 17,919    | 17,642    | 17,793    | 18,078    | 18,190    | 18,423    | 18,560    | 18,298    | 18,442    | 18,587    | 18,505    | 18,481    |
| Dairy Cows               | 10,015    | 9,482     | 9,172     | 9,106     | 9,142     | 8,988     | 9,004     | 9,104     | 9,145     | 9,257     | 9,333     | 9,087     | 9,156     | 9,236     | 9,221     | 9,208     |
| Dairy Heifer             | 4,129     | 4,108     | 4,045     | 4,060     | 4,073     | 4,033     | 4,162     | 4,294     | 4,343     | 4,401     | 4,437     | 4,545     | 4,577     | 4,581     | 4,525     | 4,546     |
| Dairy Calves             | 5,369     | 5,091     | 4,710     | 4,668     | 4,704     | 4,621     | 4,628     | 4,680     | 4,703     | 4,765     | 4,791     | 4,666     | 4,709     | 4,770     | 4,758     | 4,727     |
| Swine <sup>a</sup>       | 53,941    | 58,899    | 58,913    | 60,028    | 59,827    | 60,735    | 61,073    | 61,887    | 65,417    | 67,183    | 65,842    | 64,723    | 65,572    | 66,363    | 65,437    | 64,230    |
| Market <50 lb.           | 18,359    | 19,656    | 19,659    | 19,863    | 19,929    | 20,222    | 20,228    | 20,514    | 21,812    | 19,933    | 19,411    | 19,067    | 19,285    | 19,472    | 19,002    | 18,900    |
| Market 50-119 lb.        | 11,734    | 12,836    | 12,900    | 13,284    | 13,138    | 13,400    | 13,519    | 13,727    | 14,557    | 17,163    | 16,942    | 16,645    | 16,904    | 17,140    | 16,834    | 16,560    |
| Market 120-179 lb.       | 9,440     | 10,545    | 10,708    | 11,013    | 11,050    | 11,227    | 11,336    | 11,443    | 12,185    | 12,825    | 12,517    | 12,377    | 12,514    | 12,714    | 12,674    | 12,314    |
| Market >180 lb.          | 7,510     | 8,937     | 9,465     | 9,738     | 9,701     | 9,922     | 9,997     | 10,113    | 10,673    | 11,161    | 11,067    | 10,856    | 11,078    | 11,199    | 11,116    | 10,548    |
| Breeding                 | 6,899     | 6,926     | 6,181     | 6,129     | 6,011     | 5,963     | 5,993     | 6,090     | 6,190     | 6,102     | 5,905     | 5,778     | 5,791     | 5,839     | 5,812     | 5,908     |
| Beef Cattle <sup>b</sup> | 81,576    | 90,361    | 84,237    | 84,260    | 83,361    | 81,672    | 82,193    | 83,263    | 82,801    | 81,528    | 80,993    | 80,485    | 78,937    | 76,858    | 76,075    | 75,540    |
| Feedlot Steers           | 6,357     | 7,233     | 7,932     | 8,116     | 8,416     | 8,018     | 8,116     | 8,724     | 8,674     | 8,474     | 8,434     | 8,584     | 8,771     | 8,586     | 8,614     | 8,685     |
| Feedlot Heifers          | 3,192     | 3,831     | 4,569     | 4,557     | 4,676     | 4,521     | 4,536     | 4,801     | 4,730     | 4,585     | 4,493     | 4,620     | 4,830     | 4,742     | 4,653     | 4,537     |
| NOF Bulls                | 2,160     | 2,385     | 2,274     | 2,244     | 2,248     | 2,201     | 2,214     | 2,258     | 2,214     | 2,207     | 2,188     | 2,190     | 2,165     | 2,100     | 2,074     | 2,038     |
| Beef Calves              | 16,909    | 18,177    | 17,508    | 17,483    | 17,126    | 17,013    | 16,918    | 16,814    | 16,644    | 16,229    | 16,051    | 16,067    | 15,817    | 15,288    | 14,859    | 14,946    |
| NOF Heifers              | 10,182    | 11,829    | 9,832     | 9,843     | 9,564     | 9,321     | 9,550     | 9,716     | 9,592     | 9,356     | 9,473     | 9,350     | 8,874     | 8,687     | 8,787     | 8,838     |
| NOF Steers               | 10,321    | 11,716    | 8,724     | 8,883     | 8,347     | 8,067     | 8,185     | 8,248     | 8,302     | 8,243     | 8,560     | 8,234     | 7,568     | 7,173     | 7,457     | 7,411     |
| NOF Cows                 | 32,455    | 35,190    | 33,398    | 33,134    | 32,983    | 32,531    | 32,674    | 32,703    | 32,644    | 32,435    | 31,794    | 31,440    | 30,913    | 30,282    | 29,631    | 29,085    |
| Sheep                    | 11,358    | 8,989     | 6,908     | 6,623     | 6,321     | 6,065     | 6,135     | 6,200     | 6,120     | 5,950     | 5,747     | 5,620     | 5,470     | 5,375     | 5,335     | 5,245     |
| Sheep On Feed            | 1,180     | 1,771     | 3,256     | 3,143     | 3,049     | 2,923     | 2,971     | 3,026     | 3,000     | 2,911     | 2,806     | 2,778     | 2,687     | 2,666     | 2,626     | 2,593     |
| Sheep NOF                | 10,178    | 7,218     | 3,652     | 3,480     | 3,272     | 3,142     | 3,164     | 3,174     | 3,120     | 3,039     | 2,941     | 2,842     | 2,783     | 2,709     | 2,709     | 2,652     |
| Goats                    | 2,516     | 2,357     | 2,475     | 2,530     | 2,652     | 2,774     | 2,897     | 3,019     | 3,141     | 3,037     | 2,933     | 2,829     | 2,725     | 2,622     | 2,518     | 2,414     |
| Poultry <sup>c</sup>     | 1,537,074 | 1,826,977 | 2,060,398 | 2,097,691 | 2,085,268 | 2,130,877 | 2,150,410 | 2,154,236 | 2,166,936 | 2,175,990 | 2,088,828 | 2,104,335 | 2,095,951 | 2,168,697 | 2,106,502 | 2,111,689 |
| Hens >1 yr.              | 273,467   | 299,071   | 340,317   | 340,209   | 340,979   | 343,922   | 348,203   | 349,888   | 346,613   | 339,859   | 341,005   | 341,884   | 338,944   | 346,965   | 361,403   | 366,045   |
| Pullets                  | 73,167    | 81,369    | 95,656    | 95,289    | 100,346   | 101,429   | 96,809    | 96,596    | 103,816   | 99,458    | 102,301   | 105,738   | 102,233   | 104,460   | 106,646   | 106,602   |
| Chickens                 | 6,545     | 7,637     | 8,126     | 8,353     | 8,439     | 8,248     | 8,289     | 7,938     | 8,164     | 7,589     | 8,487     | 7,390     | 6,922     | 6,827     | 6,853     | 6,403     |
| Broilers                 | 1,066,209 | 1,331,940 | 1,525,413 | 1,562,015 | 1,544,155 | 1,589,209 | 1,613,091 | 1,612,327 | 1,619,400 | 1,638,055 | 1,554,582 | 1,567,927 | 1,565,018 | 1,625,945 | 1,551,600 | 1,553,473 |
| Turkeys                  | 117,685   | 106,960   | 90,887    | 91,826    | 91,349    | 88,069    | 84,018    | 87,487    | 88,943    | 91,029    | 82,453    | 81,396    | 82,833    | 84,500    | 80,000    | 79,167    |
| Horses                   | 2,212     | 2,632     | 3,519     | 3,644     | 3,721     | 3,798     | 3,875     | 3,952     | 4,029     | 3,947     | 3,866     | 3,784     | 3,703     | 3,621     | 3,540     | 3,458     |
| Mules and Asses          | 63        | 101       | 109       | 105       | 141       | 177       | 212       | 248       | 284       | 286       | 287       | 289       | 291       | 293       | 294       | 296       |
| American Bison           | 47        | 104       | 213       | 232       | 225       | 218       | 212       | 205       | 198       | 191       | 184       | 177       | 169       | 162       | 155       | 148       |

<sup>a</sup> Prior to 2008, the Market <50 lbs category was <60 lbs and the Market 50-119 lbs category was Market 60-119 lbs; USDA updated the categories to be more consistent with international animal categories.

<sup>b</sup> NOF - Not on Feed

<sup>c</sup> Pullets includes laying pullets, pullets younger than 3 months, and pullets older than 3 months.

Note: Totals may not sum due to independent rounding.

**Table A-204: Waste Characteristics Data**

| Animal Group              | Typical Animal Mass, TAM |              | Total Kjeldahl Nitrogen Excreted, Nex <sup>a</sup> |                                    | Maximum Methane Generation Potential, B <sub>0</sub> |                    | Volatile Solids Excreted, VS <sup>a</sup> |                                    |
|---------------------------|--------------------------|--------------|--|------------------------------------|--|--------------------|---|------------------------------------|
|                           | Value (kg)               | Source       | Value  | Source                             | Value (m <sup>3</sup> CH <sub>4</sub> /kg VS added)  | Source             | Value                                     | Source                             |
| Dairy Cows                | 680                      | CEFM         | Table A-206  | CEFM                               | 0.24   | Morris 1976        | Table A-206                               | CEFM                               |
| Dairy Heifers             | 406-408                  | CEFM         | Table A-206  | CEFM                               | 0.17   | Bryant et al. 1976 | Table A-206                               | CEFM                               |
| Feedlot Steers            | 419-457                  | CEFM         | Table A-206  | CEFM                               | 0.33   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| Feedlot Heifers           | 384-430                  | CEFM         | Table A-206  | CEFM                               | 0.33   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| NOF Bulls                 | 831-917                  | CEFM         | Table A-206  | CEFM                               | 0.17   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| NOF Calves                | 118                      | ERG 2003b    | Table A-205  | USDA 1996, 2008                    | 0.17   | Hashimoto 1981     | Table A-205                               | USDA 1996, 2008                    |
| NOF Heifers               | 296-407                  | CEFM         | Table A-206  | CEFM                               | 0.17   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| NOF Steers                | 314-335                  | CEFM         | Table A-206  | CEFM                               | 0.17   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| NOF Cows                  | 554-611                  | CEFM         | Table A-206  | CEFM                               | 0.17   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| American Bison            | 578.5                    | Meagher 1986 | Table A-206  | CEFM                               | 0.17   | Hashimoto 1981     | Table A-206                               | CEFM                               |
| Market Swine <50 lbs.     | 13                       | ERG 2010a    | Table A-205  | USDA 1996, 2008                    | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008                    |
| Market Swine <60 lbs.     | 16                       | Safley 2000  | Table A-205  | USDA 1996, 2008                    | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008                    |
| Market Swine 50-119 lbs.  | 39                       | ERG 2010a    | Table A-205  | USDA 1996, 2008                    | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008                    |
| Market Swine 60-119 lbs.  | 41                       | Safley 2000  | Table A-205  | USDA 1996, 2008                    | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008                    |
| Market Swine 120-179 lbs. | 68                       | Safley 2000  | Table A-205  | USDA 1996, 2008                    | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008                    |
| Market Swine >180 lbs.    | 91                       | Safley 2000  | Table A-205  | USDA 1996, 2008                    | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008                    |
| Breeding Swine            | 198                      | Safley 2000  | Table A-205  | USDA 1996, 2008<br>ASAE 1998, USDA | 0.48   | Hashimoto 1984     | Table A-205                               | USDA 1996, 2008<br>ASAE 1998, USDA |
| Feedlot Sheep             | 25                       | EPA 1992     | Table A-205  | 2008<br>ASAE 1998, USDA            | 0.36   | EPA 1992           | Table A-205                               | 2008<br>ASAE 1998, USDA            |
| NOF Sheep                 | 80                       | EPA 1992     | Table A-205  | 2008                               | 0.19   | EPA 1992           | Table A-205                               | 2008                               |
| Goats                     | 64                       | ASAE 1998    | Table A-205  | ASAE 1998<br>ASAE 1998, USDA       | 0.17   | EPA 1992           | Table A-205                               | ASAE 1998<br>ASAE 1998, USDA       |
| Horses                    | 450                      | ASAE 1998    | Table A-205  | 2008                               | 0.33   | EPA 1992           | Table A-205                               | 2008                               |
| Mules and Asses           | 130                      | IPCC 2006    | Table A-205  | IPCC 2006                          | 0.33   | EPA 1992           | Table A-205                               | IPCC 2006                          |
| Hens >= 1 yr              | 1.8                      | ASAE 1998    | Table A-205  | USDA 1996, 2008                    | 0.39   | Hill 1982          | Table A-205                               | USDA 1996, 2008                    |
| Pullets                   | 1.8                      | ASAE 1998    | Table A-205  | USDA 1996, 2008                    | 0.39   | Hill 1982          | Table A-205                               | USDA 1996, 2008                    |

|                |     |           |             |                 |      |           |             |                 |
|----------------|-----|-----------|-------------|-----------------|------|-----------|-------------|-----------------|
| Other Chickens | 1.8 | ASAE 1998 | Table A-205 | USDA 1996, 2008 | 0.39 | Hill 1982 | Table A-205 | USDA 1996, 2008 |
| Broilers       | 0.9 | ASAE 1998 | Table A-205 | USDA 1996, 2008 | 0.36 | Hill 1984 | Table A-205 | USDA 1996, 2008 |
| Turkeys        | 6.8 | ASAE 1998 | Table A-205 | USDA 1996, 2008 | 0.36 | Hill 1984 | Table A-205 | USDA 1996, 2008 |

<sup>a</sup> Nex and VS values vary by year; Table A-206 shows state-level values for 2014 only.

**Table A-205: Estimated Volatile Solids (VS) and Total Kjeldahl Nitrogen Excreted (Nex) Production Rates by year for Swine, Poultry, Sheep, Goats, Horses, Mules and Asses, and Cattle Calves (kg/day/1000 kg animal mass)**

| Animal Type                | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 |
|----------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| <b>VS</b>                  |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| Swine, Market <50 lbs.     | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  | 8.8  |
| Swine, Market 50-119 lbs.  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  |
| Swine, Market 120-179 lbs. | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  |
| Swine, Market >180 lbs.    | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  | 5.4  |
| Swine, Breeding            | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.6  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  | 2.7  |
| NOF Cattle Calves          | 6.4  | 6.4  | 6.4  | 6.4  | 6.4  | 6.4  | 6.4  | 6.5  | 6.6  | 6.7  | 6.8  | 6.9  | 7.1  | 7.2  | 7.3  | 7.4  | 7.5  | 7.6  | 7.7  | 7.7  | 7.7  | 7.7  | 7.7  | 7.7  | 7.7  |
| Sheep                      | 9.2  | 9.2  | 9.2  | 9.2  | 9.2  | 9.2  | 9.2  | 9.2  | 9.2  | 9.1  | 9.0  | 8.9  | 8.8  | 8.8  | 8.7  | 8.6  | 8.5  | 8.4  | 8.3  | 8.3  | 8.3  | 8.3  | 8.3  | 8.3  | 8.3  |
| Goats                      | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  | 9.5  |
| Hens >1yr.                 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 |
| Pullets                    | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.1 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 | 10.2 |
| Chickens                   | 10.8 | 10.8 | 10.8 | 10.8 | 10.8 | 10.8 | 10.8 | 10.8 | 10.8 | 10.9 | 10.9 | 10.9 | 10.9 | 10.9 | 10.9 | 11.0 | 11.0 | 11.0 | 11.0 | 11.0 | 11.0 | 11.0 | 11.0 | 11.0 | 11.0 |
| Broilers                   | 15.0 | 15.0 | 15.0 | 15.0 | 15.0 | 15.0 | 15.0 | 15.2 | 15.3 | 15.5 | 15.7 | 15.8 | 16.0 | 16.2 | 16.3 | 16.5 | 16.7 | 16.8 | 17.0 | 17.0 | 17.0 | 17.0 | 17.0 | 17.0 | 17.0 |
| Turkeys                    | 9.7  | 9.7  | 9.7  | 9.7  | 9.7  | 9.7  | 9.7  | 9.6  | 9.5  | 9.4  | 9.3  | 9.2  | 9.1  | 9.0  | 8.9  | 8.8  | 8.7  | 8.6  | 8.5  | 8.5  | 8.5  | 8.5  | 8.5  | 8.5  | 8.5  |
| Horses                     | 10.0 | 10.0 | 10.0 | 10.0 | 10.0 | 10.0 | 10.0 | 10.0 | 10.0 | 9.6  | 9.2  | 8.8  | 8.4  | 8.1  | 7.7  | 7.3  | 6.9  | 6.5  | 6.1  | 6.1  | 6.1  | 6.1  | 6.1  | 6.1  | 6.1  |
| Mules and Asses            | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  | 7.2  |
| <b>Nex</b>                 |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |      |
| Swine, Market <50 lbs.     | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.60 | 0.63 | 0.65 | 0.68 | 0.71 | 0.73 | 0.76 | 0.79 | 0.81 | 0.84 | 0.87 | 0.89 | 0.92 | 0.92 | 0.92 | 0.92 | 0.92 | 0.92 | 0.92 |

| Animal Type                   | 1990 | 1991 | 1992 | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 |
|-------------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Swine, Market<br>50-119 lbs.  | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.43 | 0.44 | 0.45 | 0.46 | 0.47 | 0.48 | 0.49 | 0.50 | 0.51 | 0.52 | 0.53 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 |
| Swine, Market<br>120-179 lbs. | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.43 | 0.44 | 0.45 | 0.46 | 0.47 | 0.48 | 0.49 | 0.50 | 0.51 | 0.52 | 0.53 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 |
| Swine, Market<br>>180 lbs.    | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.43 | 0.44 | 0.45 | 0.46 | 0.47 | 0.48 | 0.49 | 0.50 | 0.51 | 0.52 | 0.53 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 | 0.54 |
| Swine, Breeding               | 0.24 | 0.24 | 0.24 | 0.24 | 0.24 | 0.24 | 0.24 | 0.23 | 0.23 | 0.23 | 0.22 | 0.22 | 0.22 | 0.22 | 0.21 | 0.21 | 0.21 | 0.21 | 0.20 | 0.20 | 0.20 | 0.20 | 0.20 | 0.20 | 0.20 |
| NOF Cattle                    | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.31 | 0.33 | 0.34 | 0.35 | 0.36 | 0.38 | 0.39 | 0.40 | 0.41 | 0.43 | 0.44 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 |
| Calves                        | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.42 | 0.43 | 0.43 | 0.43 | 0.44 | 0.44 | 0.44 | 0.44 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 |
| Sheep                         | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 | 0.45 |
| Goats                         | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.71 | 0.72 | 0.73 | 0.73 | 0.74 | 0.75 | 0.76 | 0.77 | 0.77 | 0.78 | 0.79 | 0.79 | 0.79 | 0.79 | 0.79 | 0.79 | 0.79 |
| Hens >1yr.                    | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.70 | 0.71 | 0.72 | 0.73 | 0.73 | 0.74 | 0.75 | 0.76 | 0.77 | 0.77 | 0.78 | 0.79 | 0.79 | 0.79 | 0.79 | 0.79 | 0.79 | 0.79 |
| Pullets                       | 0.83 | 0.83 | 0.83 | 0.83 | 0.83 | 0.83 | 0.83 | 0.85 | 0.88 | 0.90 | 0.92 | 0.94 | 0.97 | 0.99 | 1.01 | 1.03 | 1.06 | 1.08 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 |
| Chickens                      | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.10 | 1.09 | 1.08 | 1.07 | 1.05 | 1.04 | 1.03 | 1.02 | 1.01 | 1.00 | 0.98 | 0.97 | 0.96 | 0.96 | 0.96 | 0.96 | 0.96 | 0.96 | 0.96 |
| Broilers                      | 0.74 | 0.74 | 0.74 | 0.74 | 0.74 | 0.74 | 0.74 | 0.73 | 0.72 | 0.71 | 0.70 | 0.69 | 0.68 | 0.67 | 0.66 | 0.65 | 0.64 | 0.63 | 0.63 | 0.63 | 0.63 | 0.63 | 0.63 | 0.63 | 0.63 |
| Turkeys                       | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.29 | 0.29 | 0.28 | 0.28 | 0.27 | 0.27 | 0.26 | 0.26 | 0.25 | 0.25 | 0.25 | 0.25 | 0.25 | 0.25 | 0.25 | 0.25 |
| Horses                        | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 |
| Mules and Asses               | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 | 0.30 |

**Table A-206: Estimated Volatile Solids (VS) and Total Kjeldahl Nitrogen Excreted (Nex) Production Rates by State for Cattle (other than Calves) and American Bison<sup>a</sup> for 2014 (kg/animal/year)**

| State          | Volatile Solids |               |              |                  |                |                 |               |               |                | Nitrogen Excreted |               |              |                  |                |                 |               |               |                |
|----------------|-----------------|---------------|--------------|------------------|----------------|-----------------|---------------|---------------|----------------|-------------------|---------------|--------------|------------------|----------------|-----------------|---------------|---------------|----------------|
|                | Dairy Cow       | Dairy Heifers | Beef NOF Cow | Beef NOF Heifers | Beef NOF Steer | Beef OF Heifers | Beef OF Steer | Beef NOF Bull | American Bison | Dairy Cow         | Dairy Heifers | Beef NOF Cow | Beef NOF Heifers | Beef NOF Steer | Beef OF Heifers | Beef OF Steer | Beef NOF Bull | American Bison |
| Alabama        | 2,171           | 1,252         | 1,664        | 1,100            | 971            | 682             | 663           | 1,721         | 1,721          | 132               | 69            | 73           | 51               | 42             | 55              | 56            | 83            | 83             |
| Alaska         | 1,971           | 1,252         | 1,891        | 1,272            | 1,116          | 682             | 663           | 1,956         | 1,956          | 121               | 69            | 59           | 42               | 33             | 55              | 56            | 69            | 69             |
| Arizona        | 2,918           | 1,252         | 1,891        | 1,244            | 1,116          | 682             | 663           | 1,956         | 1,956          | 162               | 69            | 59           | 40               | 33             | 55              | 56            | 69            | 69             |
| Arkansas       | 2,128           | 1,252         | 1,664        | 1,088            | 971            | 682             | 663           | 1,721         | 1,721          | 128               | 69            | 73           | 50               | 42             | 55              | 56            | 83            | 83             |
| California     | 2,857           | 1,252         | 1,891        | 1,211            | 1,116          | 682             | 663           | 1,956         | 1,956          | 159               | 69            | 59           | 39               | 33             | 55              | 56            | 69            | 69             |
| Colorado       | 2,963           | 1,252         | 1,891        | 1,198            | 1,116          | 682             | 663           | 1,956         | 1,956          | 164               | 69            | 59           | 38               | 33             | 55              | 56            | 69            | 69             |
| Connecticut    | 2,605           | 1,252         | 1,674        | 1,104            | 977            | 682             | 664           | 1,731         | 1,731          | 148               | 69            | 74           | 51               | 42             | 55              | 56            | 84            | 84             |
| Delaware       | 2,604           | 1,252         | 1,674        | 1,076            | 977            | 682             | 663           | 1,731         | 1,731          | 148               | 69            | 74           | 49               | 42             | 55              | 56            | 84            | 84             |
| Florida        | 2,676           | 1,252         | 1,664        | 1,101            | 971            | 682             | 662           | 1,721         | 1,721          | 153               | 69            | 73           | 51               | 42             | 55              | 56            | 83            | 83             |
| Georgia        | 2,707           | 1,252         | 1,664        | 1,097            | 971            | 682             | 664           | 1,721         | 1,721          | 155               | 69            | 73           | 50               | 42             | 55              | 57            | 83            | 83             |
| Hawaii         | 2,114           | 1,252         | 1,891        | 1,254            | 1,116          | 682             | 664           | 1,956         | 1,956          | 127               | 69            | 59           | 41               | 33             | 55              | 57            | 69            | 69             |
| Idaho          | 2,902           | 1,252         | 1,891        | 1,217            | 1,116          | 682             | 663           | 1,956         | 1,956          | 161               | 69            | 59           | 39               | 33             | 55              | 56            | 69            | 69             |
| Illinois       | 2,569           | 1,252         | 1,589        | 1,009            | 923            | 682             | 663           | 1,643         | 1,643          | 147               | 69            | 75           | 49               | 43             | 55              | 56            | 85            | 85             |
| Indiana        | 2,733           | 1,252         | 1,589        | 1,014            | 923            | 682             | 662           | 1,643         | 1,643          | 154               | 69            | 75           | 50               | 43             | 55              | 56            | 85            | 85             |
| Iowa           | 2,776           | 1,252         | 1,589        | 987              | 923            | 682             | 663           | 1,643         | 1,643          | 156               | 69            | 75           | 48               | 43             | 55              | 56            | 85            | 85             |
| Kansas         | 2,747           | 1,252         | 1,589        | 979              | 923            | 682             | 663           | 1,643         | 1,643          | 155               | 69            | 75           | 47               | 43             | 55              | 56            | 85            | 85             |
| Kentucky       | 2,342           | 1,252         | 1,664        | 1,079            | 971            | 682             | 663           | 1,721         | 1,721          | 139               | 69            | 73           | 49               | 42             | 55              | 56            | 83            | 83             |
| Louisiana      | 2,120           | 1,252         | 1,664        | 1,102            | 971            | 682             | 663           | 1,721         | 1,721          | 128               | 69            | 73           | 51               | 42             | 55              | 56            | 83            | 83             |
| Maine          | 2,593           | 1,252         | 1,674        | 1,091            | 977            | 682             | 664           | 1,731         | 1,731          | 148               | 69            | 74           | 50               | 42             | 55              | 57            | 84            | 84             |
| Maryland       | 2,574           | 1,252         | 1,674        | 1,087            | 977            | 682             | 663           | 1,731         | 1,731          | 147               | 69            | 74           | 50               | 42             | 55              | 56            | 84            | 84             |
| Massachusetts  | 2,438           | 1,252         | 1,674        | 1,108            | 977            | 682             | 664           | 1,731         | 1,731          | 141               | 69            | 74           | 52               | 42             | 55              | 57            | 84            | 84             |
| Michigan       | 2,940           | 1,252         | 1,589        | 1,009            | 923            | 682             | 663           | 1,643         | 1,643          | 163               | 69            | 75           | 49               | 43             | 55              | 56            | 85            | 85             |
| Minnesota      | 2,581           | 1,252         | 1,589        | 1,002            | 923            | 682             | 663           | 1,643         | 1,643          | 147               | 69            | 75           | 49               | 43             | 55              | 56            | 85            | 85             |
| Mississippi    | 2,234           | 1,252         | 1,664        | 1,092            | 971            | 682             | 662           | 1,721         | 1,721          | 134               | 69            | 73           | 50               | 42             | 55              | 56            | 83            | 83             |
| Missouri       | 2,260           | 1,252         | 1,589        | 1,030            | 923            | 682             | 663           | 1,643         | 1,643          | 134               | 69            | 75           | 51               | 43             | 55              | 56            | 85            | 85             |
| Montana        | 2,705           | 1,252         | 1,891        | 1,252            | 1,116          | 682             | 662           | 1,956         | 1,956          | 153               | 69            | 59           | 41               | 33             | 55              | 56            | 69            | 69             |
| Nebraska       | 2,752           | 1,252         | 1,589        | 992              | 923            | 682             | 663           | 1,643         | 1,643          | 155               | 69            | 75           | 48               | 43             | 55              | 56            | 85            | 85             |
| Nevada         | 2,877           | 1,252         | 1,891        | 1,238            | 1,116          | 682             | 663           | 1,956         | 1,956          | 160               | 69            | 59           | 40               | 33             | 55              | 56            | 69            | 69             |
| New Hampshire  | 2,604           | 1,252         | 1,674        | 1,097            | 977            | 682             | 664           | 1,731         | 1,731          | 148               | 69            | 74           | 51               | 42             | 55              | 57            | 84            | 84             |
| New Jersey     | 2,454           | 1,252         | 1,674        | 1,081            | 977            | 682             | 664           | 1,731         | 1,731          | 142               | 69            | 74           | 50               | 42             | 55              | 56            | 84            | 84             |
| New Mexico     | 2,974           | 1,252         | 1,891        | 1,234            | 1,116          | 682             | 664           | 1,956         | 1,956          | 164               | 69            | 59           | 40               | 33             | 55              | 57            | 69            | 69             |
| New York       | 2,767           | 1,252         | 1,674        | 1,084            | 977            | 682             | 663           | 1,731         | 1,731          | 155               | 69            | 74           | 50               | 42             | 55              | 56            | 84            | 84             |
| North Carolina | 2,714           | 1,252         | 1,664        | 1,098            | 971            | 682             | 663           | 1,721         | 1,721          | 155               | 69            | 73           | 50               | 42             | 55              | 56            | 83            | 83             |

| State          | Volatile Solids |               |              |                  |                |                 |               |               |                | Nitrogen Excreted |               |              |                  |                |                 |               |               |                |
|----------------|-----------------|---------------|--------------|------------------|----------------|-----------------|---------------|---------------|----------------|-------------------|---------------|--------------|------------------|----------------|-----------------|---------------|---------------|----------------|
|                | Dairy Cow       | Dairy Heifers | Beef NOF Cow | Beef NOF Heifers | Beef NOF Steer | Beef OF Heifers | Beef OF Steer | Beef NOF Bull | American Bison | Dairy Cow         | Dairy Heifers | Beef NOF Cow | Beef NOF Heifers | Beef NOF Steer | Beef OF Heifers | Beef OF Steer | Beef NOF Bull | American Bison |
| North Dakota   | 2,612           | 1,252         | 1,589        | 1,016            | 923            | 682             | 664           | 1,643         | 1,643          | 149               | 69            | 75           | 50               | 43             | 55              | 57            | 85            | 85             |
| Ohio           | 2,617           | 1,252         | 1,589        | 1,020            | 923            | 682             | 663           | 1,643         | 1,643          | 149               | 69            | 75           | 50               | 43             | 55              | 56            | 85            | 85             |
| Oklahoma       | 2,405           | 1,252         | 1,664        | 1,073            | 971            | 682             | 664           | 1,721         | 1,721          | 140               | 69            | 73           | 49               | 42             | 55              | 57            | 83            | 83             |
| Oregon         | 2,638           | 1,252         | 1,891        | 1,230            | 1,116          | 682             | 662           | 1,956         | 1,956          | 150               | 69            | 59           | 40               | 33             | 55              | 56            | 69            | 69             |
| Pennsylvania   | 2,605           | 1,252         | 1,674        | 1,072            | 977            | 682             | 663           | 1,731         | 1,731          | 148               | 69            | 74           | 49               | 42             | 55              | 56            | 84            | 84             |
| Rhode Island   | 2,519           | 1,252         | 1,674        | 1,101            | 977            | 682             | 662           | 1,731         | 1,731          | 145               | 69            | 74           | 51               | 42             | 54              | 56            | 84            | 84             |
| South Carolina | 2,377           | 1,252         | 1,664        | 1,095            | 971            | 682             | 662           | 1,721         | 1,721          | 140               | 69            | 73           | 50               | 42             | 55              | 56            | 83            | 83             |
| South Dakota   | 2,723           | 1,252         | 1,589        | 1,012            | 923            | 682             | 663           | 1,643         | 1,643          | 154               | 69            | 75           | 50               | 43             | 55              | 56            | 85            | 85             |
| Tennessee      | 2,372           | 1,252         | 1,664        | 1,085            | 971            | 682             | 663           | 1,721         | 1,721          | 140               | 69            | 73           | 50               | 42             | 55              | 56            | 83            | 83             |
| Texas          | 2,767           | 1,252         | 1,664        | 1,053            | 971            | 682             | 663           | 1,721         | 1,721          | 156               | 69            | 73           | 47               | 42             | 55              | 56            | 83            | 83             |
| Utah           | 2,815           | 1,252         | 1,891        | 1,234            | 1,116          | 682             | 663           | 1,956         | 1,956          | 157               | 69            | 59           | 40               | 33             | 55              | 56            | 69            | 69             |
| Vermont        | 2,608           | 1,252         | 1,674        | 1,075            | 977            | 682             | 664           | 1,731         | 1,731          | 149               | 69            | 74           | 49               | 42             | 55              | 57            | 84            | 84             |
| Virginia       | 2,584           | 1,252         | 1,664        | 1,086            | 971            | 682             | 663           | 1,721         | 1,721          | 149               | 69            | 73           | 50               | 42             | 55              | 56            | 83            | 83             |
| Washington     | 2,901           | 1,252         | 1,891        | 1,204            | 1,116          | 682             | 664           | 1,956         | 1,956          | 161               | 69            | 59           | 38               | 33             | 55              | 57            | 69            | 69             |
| West Virginia  | 2,261           | 1,252         | 1,674        | 1,090            | 977            | 682             | 662           | 1,731         | 1,731          | 134               | 69            | 74           | 50               | 42             | 55              | 56            | 84            | 84             |
| Wisconsin      | 2,733           | 1,252         | 1,589        | 1,022            | 923            | 682             | 663           | 1,643         | 1,643          | 154               | 69            | 75           | 50               | 43             | 55              | 56            | 85            | 85             |
| Wyoming        | 2,712           | 1,252         | 1,891        | 1,247            | 1,116          | 682             | 663           | 1,956         | 1,956          | 153               | 69            | 59           | 40               | 33             | 55              | 56            | 69            | 69             |

<sup>a</sup> Beef NOF Bull values were used for American bison Nex and VS.

Source: CEFM. NA: Not available; no population exists in this state.

**Table A-207: 2014 Manure Distribution Among Waste Management Systems by Operation (Percent)**

| State         | Beef Feedlots        |                            | Beef Not on Feed Operations | Dairy Cow Farms <sup>a</sup> |              |               |               |                  |          | Dairy Heifer Facilities   |                      |                            |                                      | Swine Operations <sup>a</sup> |               |               |                  |          | Layer Operations |                        | Broiler and Turkey Operations |                     |
|---------------|----------------------|----------------------------|-----------------------------|------------------------------|--------------|---------------|---------------|------------------|----------|---------------------------|----------------------|----------------------------|--------------------------------------|-------------------------------|---------------|---------------|------------------|----------|------------------|------------------------|-------------------------------|---------------------|
|               | Dry Lot <sup>b</sup> | Liquid/Slurry <sup>b</sup> | Pasture, Range, Paddock     | Pasture, Range, Paddock      | Daily Spread | Solid Storage | Liquid/Slurry | Anaerobic Lagoon | Deep Pit | Daily Spread <sup>b</sup> | Dry Lot <sup>b</sup> | Liquid/Slurry <sup>b</sup> | Pasture, Range, Paddock <sup>b</sup> | Pasture, Range, Paddock       | Solid Storage | Liquid/Slurry | Anaerobic Lagoon | Deep Pit | Anaerobic Lagoon | Poultry without Litter | Pasture, Range, Paddock       | Poultry with Litter |
| Alabama       | 100                  | 1                          | 100                         | 51                           | 16           | 7             | 10            | 16               | 0        | 17                        | 38                   | 0                          | 45                                   | 5                             | 4             | 7             | 54               | 31       | 42               | 58                     | 1                             | 99                  |
| Alaska        | 100                  | 1                          | 100                         | 5                            | 9            | 34            | 19            | 24               | 9        | 6                         | 90                   | 1                          | 4                                    | 64                            | 2             | 10            | 7                | 17       | 25               | 75                     | 1                             | 99                  |
| Arizona       | 100                  | 0                          | 100                         | 0                            | 10           | 9             | 19            | 61               | 0        | 10                        | 90                   | 0                          | 0                                    | 6                             | 3             | 6             | 55               | 29       | 60               | 40                     | 1                             | 99                  |
| Arkansas      | 100                  | 1                          | 100                         | 60                           | 14           | 10            | 7             | 9                | 0        | 15                        | 28                   | 0                          | 57                                   | 4                             | 4             | 13            | 45               | 35       | 0                | 100                    | 1                             | 99                  |
| California    | 100                  | 1                          | 100                         | 1                            | 11           | 9             | 20            | 59               | 0        | 11                        | 88                   | 1                          | 1                                    | 10                            | 3             | 7             | 50               | 29       | 12               | 88                     | 1                             | 99                  |
| Colorado      | 100                  | 0                          | 100                         | 1                            | 1            | 11            | 23            | 64               | 0        | 1                         | 98                   | 0                          | 1                                    | 1                             | 6             | 26            | 17               | 50       | 60               | 40                     | 1                             | 99                  |
| Connecticut   | 100                  | 1                          | 100                         | 6                            | 43           | 16            | 20            | 13               | 2        | 43                        | 51                   | 0                          | 6                                    | 78                            | 1             | 6             | 5                | 11       | 5                | 95                     | 1                             | 99                  |
| Delaware      | 100                  | 1                          | 100                         | 6                            | 44           | 19            | 19            | 10               | 2        | 44                        | 50                   | 0                          | 6                                    | 8                             | 5             | 25            | 17               | 46       | 5                | 95                     | 1                             | 99                  |
| Florida       | 100                  | 1                          | 100                         | 13                           | 22           | 7             | 15            | 43               | 0        | 22                        | 61                   | 1                          | 17                                   | 72                            | 1             | 8             | 6                | 13       | 42               | 58                     | 1                             | 99                  |
| Georgia       | 100                  | 1                          | 100                         | 37                           | 18           | 9             | 12            | 23               | 0        | 18                        | 42                   | 0                          | 40                                   | 4                             | 4             | 8             | 53               | 31       | 42               | 58                     | 1                             | 99                  |
| Hawaii        | 100                  | 1                          | 100                         | 10                           | 0            | 9             | 23            | 57               | 0        | 0                         | 99                   | 1                          | 1                                    | 31                            | 3             | 19            | 14               | 32       | 25               | 75                     | 1                             | 99                  |
| Idaho         | 100                  | 0                          | 100                         | 0                            | 0            | 11            | 23            | 65               | 0        | 1                         | 99                   | 0                          | 0                                    | 12                            | 5             | 23            | 15               | 44       | 60               | 40                     | 1                             | 99                  |
| Illinois      | 100                  | 1                          | 100                         | 4                            | 6            | 39            | 31            | 16               | 5        | 8                         | 87                   | 0                          | 5                                    | 1                             | 5             | 29            | 14               | 52       | 2                | 98                     | 1                             | 99                  |
| Indiana       | 100                  | 1                          | 100                         | 5                            | 8            | 29            | 31            | 24               | 3        | 13                        | 79                   | 0                          | 8                                    | 1                             | 5             | 28            | 14               | 52       | 0                | 100                    | 1                             | 99                  |
| Iowa          | 100                  | 1                          | 100                         | 4                            | 8            | 34            | 30            | 20               | 4        | 10                        | 83                   | 0                          | 6                                    | 1                             | 4             | 9             | 54               | 33       | 0                | 100                    | 1                             | 99                  |
| Kansas        | 100                  | 1                          | 100                         | 2                            | 3            | 21            | 37            | 36               | 2        | 5                         | 92                   | 0                          | 3                                    | 2                             | 5             | 28            | 13               | 52       | 2                | 98                     | 1                             | 99                  |
| Kentucky      | 100                  | 1                          | 100                         | 60                           | 14           | 14            | 7             | 3                | 2        | 14                        | 24                   | 0                          | 61                                   | 5                             | 4             | 10            | 48               | 33       | 5                | 95                     | 1                             | 99                  |
| Louisiana     | 100                  | 1                          | 100                         | 59                           | 15           | 10            | 7             | 9                | 1        | 14                        | 26                   | 0                          | 60                                   | 88                            | 1             | 3             | 3                | 6        | 60               | 40                     | 1                             | 99                  |
| Maine         | 100                  | 1                          | 100                         | 7                            | 45           | 20            | 17            | 10               | 2        | 45                        | 48                   | 0                          | 7                                    | 65                            | 2             | 10            | 7                | 16       | 5                | 95                     | 1                             | 99                  |
| Maryland      | 100                  | 1                          | 100                         | 7                            | 44           | 22            | 16            | 8                | 3        | 44                        | 49                   | 0                          | 7                                    | 7                             | 5             | 25            | 17               | 47       | 5                | 95                     | 1                             | 99                  |
| Massachusetts | 100                  | 1                          | 100                         | 7                            | 44           | 22            | 16            | 8                | 3        | 45                        | 47                   | 0                          | 7                                    | 56                            | 2             | 12            | 9                | 20       | 5                | 95                     | 1                             | 99                  |
| Michigan      | 100                  | 1                          | 100                         | 2                            | 4            | 24            | 38            | 29               | 3        | 6                         | 91                   | 0                          | 3                                    | 4                             | 5             | 26            | 17               | 48       | 2                | 98                     | 1                             | 99                  |
| Minnesota     | 100                  | 1                          | 100                         | 5                            | 8            | 39            | 28            | 17               | 4        | 10                        | 84                   | 0                          | 6                                    | 1                             | 5             | 26            | 18               | 50       | 0                | 100                    | 1                             | 99                  |
| Mississippi   | 100                  | 1                          | 100                         | 54                           | 15           | 10            | 8             | 12               | 0        | 15                        | 28                   | 0                          | 57                                   | 2                             | 4             | 6             | 58               | 31       | 60               | 40                     | 1                             | 99                  |
| Missouri      | 100                  | 1                          | 100                         | 7                            | 12           | 42            | 22            | 11               | 5        | 14                        | 77                   | 0                          | 8                                    | 2                             | 5             | 28            | 13               | 52       | 0                | 100                    | 1                             | 99                  |
| Montana       | 100                  | 0                          | 100                         | 2                            | 4            | 19            | 28            | 42               | 4        | 4                         | 93                   | 0                          | 3                                    | 3                             | 5             | 25            | 17               | 49       | 60               | 40                     | 1                             | 99                  |
| Nebraska      | 100                  | 1                          | 100                         | 2                            | 4            | 26            | 35            | 29               | 3        | 6                         | 90                   | 0                          | 4                                    | 1                             | 5             | 28            | 14               | 51       | 2                | 98                     | 1                             | 99                  |
| Nevada        | 100                  | 0                          | 100                         | 0                            | 0            | 10            | 24            | 65               | 0        | 0                         | 99                   | 0                          | 0                                    | 34                            | 3             | 18            | 14               | 31       | 0                | 100                    | 1                             | 99                  |
| New Hampshire | 100                  | 1                          | 100                         | 7                            | 44           | 19            | 18            | 10               | 2        | 44                        | 49                   | 0                          | 7                                    | 64                            | 2             | 10            | 8                | 17       | 5                | 95                     | 1                             | 99                  |
| New Jersey    | 100                  | 1                          | 100                         | 7                            | 45           | 25            | 13            | 6                | 3        | 45                        | 47                   | 0                          | 8                                    | 36                            | 3             | 18            | 14               | 30       | 5                | 95                     | 1                             | 99                  |
| New Mexico    | 100                  | 0                          | 100                         | 0                            | 10           | 9             | 19            | 61               | 0        | 10                        | 90                   | 0                          | 0                                    | 100                           | 0             | 0             | 0                | 0        | 60               | 40                     | 1                             | 99                  |

| State          | Beef Feedlots        |                            | Beef Not on Feed Operations | Dairy Cow Farms <sup>a</sup> |              |               |               |                  |          | Dairy Heifer Facilities   |                      |                            |                                      | Swine Operations <sup>a</sup> |               |               |                  |          | Layer Operations |                        | Broiler and Turkey Operations |                     |
|----------------|----------------------|----------------------------|-----------------------------|------------------------------|--------------|---------------|---------------|------------------|----------|---------------------------|----------------------|----------------------------|--------------------------------------|-------------------------------|---------------|---------------|------------------|----------|------------------|------------------------|-------------------------------|---------------------|
|                | Dry Lot <sup>b</sup> | Liquid/Slurry <sup>b</sup> | Pasture, Range, Paddock     | Pasture, Range, Paddock      | Daily Spread | Solid Storage | Liquid/Slurry | Anaerobic Lagoon | Deep Pit | Daily Spread <sup>b</sup> | Dry Lot <sup>b</sup> | Liquid/Slurry <sup>b</sup> | Pasture, Range, Paddock <sup>b</sup> | Pasture, Range, Paddock       | Solid Storage | Liquid/Slurry | Anaerobic Lagoon | Deep Pit | Anaerobic Lagoon | Poultry without Litter | Pasture, Range, Paddock       | Poultry with Litter |
| New York       | 100                  | 1                          | 100                         | 6                            | 44           | 17            | 18            | 13               | 2        | 45                        | 48                   | 0                          | 7                                    | 13                            | 5             | 23            | 15               | 44       | 5                | 95                     | 1                             | 99                  |
| North Carolina | 100                  | 1                          | 100                         | 46                           | 17           | 11            | 15            | 10               | 2        | 15                        | 31                   | 0                          | 54                                   | 0                             | 4             | 7             | 57               | 32       | 42               | 58                     | 1                             | 99                  |
| North Dakota   | 100                  | 1                          | 100                         | 7                            | 11           | 38            | 26            | 15               | 4        | 11                        | 83                   | 0                          | 6                                    | 5                             | 5             | 25            | 17               | 48       | 2                | 98                     | 1                             | 99                  |
| Ohio           | 100                  | 1                          | 100                         | 6                            | 11           | 38            | 26            | 15               | 4        | 14                        | 78                   | 0                          | 8                                    | 3                             | 5             | 28            | 14               | 51       | 0                | 100                    | 1                             | 99                  |
| Oklahoma       | 100                  | 0                          | 100                         | 0                            | 7            | 21            | 22            | 45               | 4        | 6                         | 94                   | 0                          | 0                                    | 1                             | 4             | 6             | 58               | 31       | 60               | 40                     | 1                             | 99                  |
| Oregon         | 100                  | 1                          | 100                         | 16                           | 0            | 11            | 22            | 50               | 1        | 0                         | 80                   | 1                          | 20                                   | 48                            | 2             | 14            | 11               | 24       | 25               | 75                     | 1                             | 99                  |
| Pennsylvania   | 100                  | 1                          | 100                         | 8                            | 46           | 24            | 12            | 6                | 2        | 47                        | 44                   | 0                          | 9                                    | 4                             | 5             | 26            | 18               | 48       | 0                | 100                    | 1                             | 99                  |
| Rhode Island   | 100                  | 1                          | 100                         | 9                            | 47           | 25            | 13            | 5                | 2        | 47                        | 44                   | 0                          | 9                                    | 72                            | 1             | 8             | 6                | 13       | 5                | 95                     | 1                             | 99                  |
| South Carolina | 100                  | 1                          | 100                         | 47                           | 17           | 8             | 11            | 18               | 0        | 15                        | 31                   | 0                          | 54                                   | 3                             | 4             | 7             | 55               | 31       | 60               | 40                     | 1                             | 99                  |
| South Dakota   | 100                  | 1                          | 100                         | 3                            | 4            | 24            | 36            | 31               | 2        | 8                         | 87                   | 0                          | 5                                    | 1                             | 5             | 26            | 17               | 50       | 2                | 98                     | 1                             | 99                  |
| Tennessee      | 100                  | 1                          | 100                         | 58                           | 15           | 12            | 9             | 4                | 2        | 15                        | 26                   | 0                          | 59                                   | 13                            | 3             | 11            | 41               | 32       | 5                | 95                     | 1                             | 99                  |
| Texas          | 100                  | 0                          | 100                         | 0                            | 9            | 11            | 22            | 58               | 1        | 8                         | 92                   | 0                          | 0                                    | 3                             | 4             | 6             | 57               | 30       | 12               | 88                     | 1                             | 99                  |
| Utah           | 100                  | 0                          | 100                         | 1                            | 1            | 15            | 26            | 56               | 2        | 1                         | 98                   | 0                          | 1                                    | 1                             | 6             | 26            | 17               | 51       | 60               | 40                     | 1                             | 99                  |
| Vermont        | 100                  | 1                          | 100                         | 6                            | 44           | 17            | 19            | 13               | 2        | 44                        | 49                   | 0                          | 7                                    | 63                            | 2             | 10            | 8                | 18       | 5                | 95                     | 1                             | 99                  |
| Virginia       | 100                  | 1                          | 100                         | 56                           | 15           | 11            | 10            | 5                | 2        | 15                        | 28                   | 0                          | 57                                   | 4                             | 4             | 7             | 54               | 31       | 5                | 95                     | 1                             | 99                  |
| Washington     | 100                  | 1                          | 100                         | 11                           | 0            | 11            | 22            | 56               | 1        | 0                         | 83                   | 1                          | 17                                   | 43                            | 3             | 15            | 11               | 28       | 12               | 88                     | 1                             | 99                  |
| West Virginia  | 100                  | 1                          | 100                         | 8                            | 46           | 23            | 14            | 7                | 2        | 45                        | 48                   | 0                          | 7                                    | 59                            | 2             | 11            | 7                | 20       | 5                | 95                     | 1                             | 99                  |
| Wisconsin      | 100                  | 1                          | 100                         | 5                            | 9            | 38            | 28            | 17               | 4        | 12                        | 82                   | 0                          | 7                                    | 13                            | 4             | 23            | 17               | 42       | 2                | 98                     | 1                             | 99                  |
| Wyoming        | 100                  | 0                          | 100                         | 4                            | 6            | 19            | 23            | 43               | 4        | 12                        | 81                   | 0                          | 7                                    | 4                             | 5             | 25            | 16               | 49       | 60               | 40                     | 1                             | 99                  |

<sup>a</sup> In the methane inventory for manure management, the percent of dairy cows and swine with AD systems is estimated using data from EPA's AgSTAR Program.

<sup>b</sup> Because manure from beef feedlots and dairy heifers may be managed for long periods of time in multiple systems (i.e., both drylot and runoff collection pond), the percent of manure that generates emissions is greater than 100 percent.

**Table A-208: Manure Management System Descriptions**

| Manure Management System | Description <sup>a</sup>   |
|--------------------------|--|
| Pasture, Range, Paddock  | The manure from pasture and range grazing animals is allowed to lie as is, and is not managed. Methane emissions are accounted for under Manure Management, but the N <sub>2</sub> O emissions from manure deposited on PRP are included under the Agricultural Soil Management category.  |
| Daily Spread             | Manure is routinely removed from a confinement facility and is applied to cropland or pasture within 24 hours of excretion. Methane and indirect N <sub>2</sub> O emissions are accounted for under Manure Management. Direct N <sub>2</sub> O emissions from land application are covered under the Agricultural Soil Management category.  |
| Solid Storage            | The storage of manure, typically for a period of several months, in unconfined piles or stacks. Manure is able to be stacked due to the presence of a sufficient amount of bedding material or loss of moisture by evaporation.  |
| Dry Lot                  | A paved or unpaved open confinement area without any significant vegetative cover where accumulating manure may be removed periodically. Dry lots are most typically found in dry climates but also are used in humid climates.  |
| Liquid/ Slurry           | Manure is stored as excreted or with some minimal addition of water to facilitate handling and is stored in either tanks or earthen ponds, usually for periods less than one year.   |
| Anaerobic Lagoon         | Uncovered anaerobic lagoons are designed and operated to combine waste stabilization and storage. Lagoon supernatant is usually used to remove manure from the associated confinement facilities to the lagoon. Anaerobic lagoons are designed with varying lengths of storage (up to a year or greater), depending on the climate region, the VS loading rate, and other operational factors. Anaerobic lagoons accumulate sludge over time, diminishing treatment capacity. Lagoons must be cleaned out once every 5 to 15 years, and the sludge is typically applied to agricultural lands. The water from the lagoon may be recycled as flush water or used to irrigate and fertilize fields. Lagoons are sometimes used in combination with a solids separator, typically for dairy waste. Solids separators help control the buildup of nondegradable material such as straw or other bedding materials. |
| Anaerobic Digester       | Animal excreta with or without straw are collected and anaerobically digested in a large containment vessel (complete mix or plug flow digester) or covered lagoon. Digesters are designed and operated for waste stabilization by the microbial reduction of complex organic compounds to CO <sub>2</sub> and CH <sub>4</sub> , which is captured and flared or used as a fuel.   |
| Deep Pit                 | Collection and storage of manure usually with little or no added water typically below a slatted floor in an enclosed animal confinement facility. Typical storage periods range from 5 to 12 months, after which manure is removed from the pit and transferred to a treatment system or applied to land.   |
| Poultry with Litter      | Enclosed poultry houses use bedding derived from wood shavings, rice hulls, chopped straw, peanut hulls, or other products, depending on availability. The bedding absorbs moisture and dilutes the manure produced by the birds. Litter is typically cleaned out completely once a year. These manure systems are typically used for all poultry breeder flocks and for the production of meat type chickens (broilers) and other fowl.   |
| Poultry without Litter   | In high-rise cages or scrape-out/belt systems, manure is excreted onto the floor below with no bedding to absorb moisture. The ventilation system dries the manure as it is stored. When designed and operated properly, this high-rise system is a form of passive windrow composting.  |

<sup>a</sup> Manure management system descriptions are based on the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Volume 4: Agriculture, Forestry and Other Land Use, Chapter 10: Emissions from Livestock and Manure Management, Tables 10.18 and 10.21) and the Development Document for the Final Revisions to the National Pollutant Discharge Elimination System Regulation and the Effluent Guidelines for Concentrated Animal Feeding Operations (EPA-821-R-03-001, December 2002).

**Table A-209: Methane Conversion Factors (percent) for Dry Systems**

| Waste Management System       | Cool Climate MCF | Temperate Climate MCF | Warm Climate MCF |
|-------------------------------|------------------|-----------------------|------------------|
| Aerobic Treatment             | 0                | 0                     | 0                |
| Anaerobic Digester            | 0                | 0                     | 0                |
| Cattle Deep Litter (<1 month) | 3                | 3                     | 30               |
| Cattle Deep Litter (>1 month) | 21               | 44                    | 76               |

| Waste Management System       | Cool Climate MCF | Temperate Climate MCF | Warm Climate MCF |
|-------------------------------|------------------|-----------------------|------------------|
| Composting - In Vessel        | 0.5              | 0.5                   | 0.5              |
| Composting - Static Pile      | 0.5              | 0.5                   | 0.5              |
| Composting-Extensive/ Passive | 0.5              | 1                     | 1.5              |
| Composting-Intensive          | 0.5              | 1                     | 1.5              |
| Daily Spread                  | 0.1              | 0.5                   | 1                |
| Dry Lot                       | 1                | 1.5                   | 5                |
| Fuel                          | 10               | 10                    | 10               |
| Pasture                       | 1                | 1.5                   | 2                |
| Poultry with bedding          | 1.5              | 1.5                   | 1.5              |
| Poultry without bedding       | 1.5              | 1.5                   | 1.5              |
| Solid Storage                 | 2                | 4                     | 5                |

Source: IPCC 2006

**Table A-210: Methane Conversion Factors by State for Liquid Systems for 2014 (percent)**

| State          | Dairy            |                            | Swine            |                            | Beef          | Poultry          |
|----------------|------------------|----------------------------|------------------|----------------------------|---------------|------------------|
|                | Anaerobic Lagoon | Liquid/Slurry and Deep Pit | Anaerobic Lagoon | Liquid/Slurry and Deep Pit | Liquid/Slurry | Anaerobic Lagoon |
| Alabama        | 75               | 37                         | 75               | 36                         | 38            | 75               |
| Alaska         | 47               | 15                         | 47               | 15                         | 15            | 47               |
| Arizona        | 78               | 57                         | 77               | 47                         | 52            | 74               |
| Arkansas       | 75               | 34                         | 76               | 37                         | 35            | 75               |
| California     | 73               | 32                         | 72               | 31                         | 41            | 74               |
| Colorado       | 65               | 22                         | 68               | 24                         | 24            | 65               |
| Connecticut    | 69               | 25                         | 69               | 26                         | 26            | 69               |
| Delaware       | 73               | 31                         | 73               | 31                         | 31            | 73               |
| Florida        | 79               | 55                         | 79               | 53                         | 53            | 79               |
| Georgia        | 76               | 39                         | 75               | 38                         | 37            | 75               |
| Hawaii         | 76               | 57                         | 76               | 57                         | 57            | 76               |
| Idaho          | 69               | 25                         | 66               | 22                         | 22            | 68               |
| Illinois       | 72               | 29                         | 72               | 28                         | 27            | 72               |
| Indiana        | 70               | 27                         | 71               | 27                         | 27            | 71               |
| Iowa           | 70               | 25                         | 70               | 26                         | 26            | 70               |
| Kansas         | 74               | 32                         | 74               | 32                         | 32            | 74               |
| Kentucky       | 73               | 31                         | 73               | 31                         | 30            | 73               |
| Louisiana      | 77               | 45                         | 77               | 46                         | 46            | 77               |
| Maine          | 63               | 21                         | 63               | 21                         | 21            | 64               |
| Maryland       | 72               | 30                         | 72               | 30                         | 31            | 73               |
| Massachusetts  | 67               | 24                         | 68               | 25                         | 25            | 68               |
| Michigan       | 67               | 23                         | 67               | 24                         | 24            | 67               |
| Minnesota      | 68               | 24                         | 69               | 24                         | 24            | 67               |
| Mississippi    | 76               | 40                         | 76               | 38                         | 41            | 76               |
| Missouri       | 73               | 30                         | 73               | 30                         | 30            | 74               |
| Montana        | 61               | 19                         | 64               | 21                         | 21            | 64               |
| Nebraska       | 72               | 27                         | 72               | 27                         | 27            | 72               |
| Nevada         | 70               | 26                         | 71               | 27                         | 25            | 70               |
| New Hampshire  | 64               | 22                         | 65               | 22                         | 22            | 65               |
| New Jersey     | 71               | 28                         | 71               | 29                         | 28            | 71               |
| New Mexico     | 73               | 31                         | 71               | 28                         | 30            | 70               |
| New York       | 65               | 23                         | 66               | 23                         | 23            | 66               |
| North Carolina | 73               | 31                         | 75               | 36                         | 30            | 73               |
| North Dakota   | 66               | 22                         | 66               | 22                         | 22            | 66               |
| Ohio           | 69               | 26                         | 70               | 27                         | 27            | 70               |
| Oklahoma       | 76               | 37                         | 75               | 35                         | 36            | 76               |
| Oregon         | 64               | 21                         | 63               | 21                         | 22            | 63               |
| Pennsylvania   | 69               | 26                         | 70               | 27                         | 27            | 70               |
| Rhode Island   | 69               | 26                         | 69               | 26                         | 26            | 69               |

| State          | Dairy            |                            | Swine            |                            | Beef          | Poultry          |
|----------------|------------------|----------------------------|------------------|----------------------------|---------------|------------------|
|                | Anaerobic Lagoon | Liquid/Slurry and Deep Pit | Anaerobic Lagoon | Liquid/Slurry and Deep Pit | Liquid/Slurry | Anaerobic Lagoon |
| South Carolina | 75               | 37                         | 76               | 38                         | 36            | 75               |
| South Dakota   | 69               | 24                         | 70               | 25                         | 25            | 70               |
| Tennessee      | 73               | 31                         | 74               | 32                         | 31            | 73               |
| Texas          | 76               | 41                         | 76               | 44                         | 38            | 77               |
| Utah           | 65               | 22                         | 69               | 25                         | 24            | 65               |
| Vermont        | 63               | 21                         | 63               | 21                         | 21            | 63               |
| Virginia       | 71               | 28                         | 72               | 31                         | 29            | 71               |
| Washington     | 64               | 21                         | 66               | 22                         | 23            | 65               |
| West Virginia  | 69               | 26                         | 70               | 27                         | 26            | 69               |
| Wisconsin      | 66               | 23                         | 68               | 24                         | 23            | 67               |
| Wyoming        | 63               | 20                         | 64               | 21                         | 22            | 64               |

Note: MCFs developed using Tier 2 methods described in IPCC 2006, Section 10.4.2.

**Table A-211: Direct Nitrous Oxide Emission Factors for 2014 (kg N<sub>2</sub>O-N/kg Kjdl N)**

| Waste Management System              | Direct N <sub>2</sub> O Emission Factor |
|--------------------------------------|---|
| Aerobic Treatment (forced aeration)  | 0.005                                   |
| Aerobic Treatment (natural aeration) | 0.01                                    |
| Anaerobic Digester                   | 0                                       |
| Anaerobic Lagoon                     | 0                                       |
| Cattle Deep Bed (active mix)         | 0.07                                    |
| Cattle Deep Bed (no mix)             | 0.01                                    |
| Composting_in vessel                 | 0.006                                   |
| Composting_intensive                 | 0.1                                     |
| Composting_passive                   | 0.01                                    |
| Composting_static                    | 0.006                                   |
| Daily Spread                         | 0                                       |
| Deep Pit                             | 0.002                                   |
| Dry Lot                              | 0.02                                    |
| Fuel                                 | 0                                       |
| Liquid/Slurry                        | 0.005                                   |
| Pasture                              | 0                                       |
| Poultry with bedding                 | 0.001                                   |
| Poultry without bedding              | 0.001                                   |
| Solid Storage                        | 0.005                                   |

Source: IPCC 2006

**Table A-212: Indirect Nitrous Oxide Loss Factors (percent)**

| Animal Type  | Waste Management System | Volatilization Nitrogen Loss | Runoff/Leaching Nitrogen Loss <sup>a</sup> |         |              |         |       |
|--------------|-------------------------|------------------------------|--|---------|--------------|---------|-------|
|              |                         |                              | Central                                    | Pacific | Mid-Atlantic | Midwest | South |
| Beef Cattle  | Dry Lot                 | 23                           | 1.1  | 3.9     | 3.6          | 1.9     | 4.3   |
| Beef Cattle  | Liquid/Slurry           | 26                           | 0  | 0       | 0            | 0       | 0     |
| Beef Cattle  | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Dairy Cattle | Anaerobic Lagoon        | 43                           | 0.2  | 0.8     | 0.7          | 0.4     | 0.9   |
| Dairy Cattle | Daily Spread            | 10                           | 0  | 0       | 0            | 0       | 0     |
| Dairy Cattle | Deep Pit                | 24                           | 0  | 0       | 0            | 0       | 0     |
| Dairy Cattle | Dry Lot                 | 15                           | 0.6  | 2       | 1.8          | 0.9     | 2.2   |
| Dairy Cattle | Liquid/Slurry           | 26                           | 0.2  | 0.8     | 0.7          | 0.4     | 0.9   |
| Dairy Cattle | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |

| Animal Type     | Waste Management System | Volatilization Nitrogen Loss | Runoff/Leaching Nitrogen Loss <sup>a</sup> |         |              |         |       |
|-----------------|-------------------------|------------------------------|--|---------|--------------|---------|-------|
|                 |                         |                              | Central                                    | Pacific | Mid-Atlantic | Midwest | South |
| Dairy Cattle    | Solid Storage           | 27                           | 0.2  | 0       | 0            | 0       | 0     |
| American Bison  | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Goats           | Dry Lot                 | 23                           | 1.1  | 3.9     | 3.6          | 1.9     | 4.3   |
| Goats           | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Horses          | Dry Lot                 | 23                           | 0  | 0       | 0            | 0       | 0     |
| Horses          | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Mules and Asses | Dry Lot                 | 23                           | 0  | 0       | 0            | 0       | 0     |
| Mules and Asses | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Poultry         | Anaerobic Lagoon        | 54                           | 0.2  | 0.8     | 0.7          | 0.4     | 0.9   |
| Poultry         | Liquid/Slurry           | 26                           | 0.2  | 0.8     | 0.7          | 0.4     | 0.9   |
| Poultry         | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Poultry         | Poultry with bedding    | 26                           | 0  | 0       | 0            | 0       | 0     |
| Poultry         | Poultry without bedding | 34                           | 0  | 0       | 0            | 0       | 0     |
| Poultry         | Solid Storage           | 8                            | 0  | 0       | 0            | 0       | 0     |
| Sheep           | Dry Lot                 | 23                           | 1.1  | 3.9     | 3.6          | 1.9     | 4.3   |
| Sheep           | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Swine           | Anaerobic Lagoon        | 58                           | 0.2  | 0.8     | 0.7          | 0.4     | 0.9   |
| Swine           | Deep Pit                | 34                           | 0  | 0       | 0            | 0       | 0     |
| Swine           | Liquid/Slurry           | 26                           | 0.2  | 0.8     | 0.7          | 0.4     | 0.9   |
| Swine           | Pasture                 | 0                            | 0  | 0       | 0            | 0       | 0     |
| Swine           | Solid Storage           | 45                           | 0  | 0       | 0            | 0       | 0     |

<sup>a</sup> Data for nitrogen losses due to leaching were not available, so the values represent only nitrogen losses due to runoff.  
Source: EPA 2002b, 2005.

**Table A-213: Methane Emissions from Livestock Manure Management (kt)<sup>a</sup>**

| Animal Type                | 1990       | 1991       | 1992       | 1993       | 1994       | 1995       | 1996       | 1997       | 1998       | 1999       | 2000       | 2001       | 2002       | 2003         | 2004       | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|----------------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Dairy Cattle</b>        | <b>590</b> | <b>607</b> | <b>591</b> | <b>614</b> | <b>656</b> | <b>685</b> | <b>695</b> | <b>725</b> | <b>771</b> | <b>841</b> | <b>889</b> | <b>951</b> | <b>985</b> | <b>1,036</b> | <b>988</b> | <b>1,057</b> | <b>1,091</b> | <b>1,212</b> | <b>1,230</b> | <b>1,218</b> | <b>1,217</b> | <b>1,245</b> | <b>1,306</b> | <b>1,271</b> | <b>1,289</b> |
| <i>Dairy Cows</i>          | 581        | 598        | 583        | 606        | 647        | 676        | 687        | 716        | 763        | 832        | 880        | 942        | 977        | 1,027        | 980        | 1,049        | 1,083        | 1,202        | 1,220        | 1,208        | 1,207        | 1,236        | 1,295        | 1,261        | 1,278        |
| <i>Dairy Heifer</i>        | 7          | 7          | 7          | 7          | 7          | 7          | 7          | 7          | 6          | 7          | 7          | 7          | 7          | 7            | 6          | 7            | 7            | 8            | 8            | 8            | 8            | 8            | 9            | 8            | 8            |
| <i>Dairy Calves</i>        | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2          | 2            | 2          | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            | 2            |
| <b>Swine</b>               | <b>622</b> | <b>674</b> | <b>637</b> | <b>678</b> | <b>740</b> | <b>763</b> | <b>729</b> | <b>782</b> | <b>891</b> | <b>849</b> | <b>835</b> | <b>854</b> | <b>877</b> | <b>859</b>   | <b>858</b> | <b>916</b>   | <b>902</b>   | <b>984</b>   | <b>936</b>   | <b>896</b>   | <b>945</b>   | <b>942</b>   | <b>972</b>   | <b>920</b>   | <b>896</b>   |
| Market Swine               | 483        | 522        | 499        | 533        | 584        | 608        | 581        | 625        | 720        | 693        | 681        | 697        | 719        | 705          | 707        | 755          | 742          | 816          | 779          | 748          | 790          | 788          | 814          | 771          | 744          |
| <i>Market &lt;50 lbs.</i>  | 102        | 110        | 103        | 108        | 119        | 121        | 116        | 125        | 140        | 133        | 131        | 134        | 137        | 135          | 135        | 142          | 141          | 155          | 109          | 103          | 110          | 109          | 113          | 105          | 104          |
| <i>Market 50-119 lbs.</i>  | 101        | 110        | 104        | 110        | 119        | 123        | 117        | 127        | 144        | 138        | 136        | 138        | 144        | 140          | 141        | 150          | 148          | 163          | 174          | 167          | 176          | 0            | 0            | 0            | 0            |
| <i>Market 120-179 lbs.</i> | 136        | 147        | 139        | 150        | 164        | 170        | 163        | 175        | 201        | 193        | 189        | 192        | 199        | 196          | 196        | 210          | 206          | 228          | 228          | 218          | 232          | 230          | 238          | 228          | 221          |
| <i>Market &gt;180 lbs.</i> | 144        | 156        | 152        | 164        | 182        | 193        | 184        | 198        | 234        | 229        | 225        | 232        | 240        | 234          | 235        | 252          | 247          | 270          | 268          | 259          | 272          | 274          | 281          | 266          | 250          |
| Breeding Swine             | 139        | 151        | 138        | 146        | 156        | 155        | 148        | 157        | 171        | 157        | 155        | 158        | 158        | 154          | 151        | 161          | 160          | 168          | 157          | 148          | 155          | 154          | 158          | 150          | 152          |
| <b>Beef Cattle</b>         | <b>126</b> | <b>126</b> | <b>129</b> | <b>130</b> | <b>135</b> | <b>139</b> | <b>136</b> | <b>134</b> | <b>137</b> | <b>137</b> | <b>131</b> | <b>134</b> | <b>131</b> | <b>131</b>   | <b>129</b> | <b>133</b>   | <b>137</b>   | <b>134</b>   | <b>130</b>   | <b>130</b>   | <b>132</b>   | <b>131</b>   | <b>128</b>   | <b>121</b>   | <b>120</b>   |
| <i>Feedlot Steers</i>      | 14         | 14         | 14         | 13         | 14         | 14         | 14         | 13         | 13         | 14         | 15         | 15         | 15         | 16           | 15         | 15           | 16           | 16           | 16           | 16           | 16           | 17           | 16           | 16           | 16           |
| <i>Feedlot Heifers</i>     | 7          | 7          | 7          | 7          | 8          | 8          | 8          | 8          | 8          | 8          | 9          | 9          | 9          | 9            | 9          | 9            | 9            | 9            | 9            | 9            | 9            | 9            | 9            | 9            | 9            |
| <i>NOF Bulls</i>           | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5          | 5            | 5          | 5            | 5            | 5            | 5            | 5            | 5            | 5            | 5            | 5            | 5            |
| <i>Beef Calves</i>         | 6          | 6          | 6          | 6          | 7          | 7          | 6          | 6          | 7          | 7          | 7          | 7          | 7          | 7            | 7          | 7            | 7            | 7            | 7            | 7            | 7            | 7            | 7            | 6            | 6            |
| <i>NOF Heifers</i>         | 12         | 12         | 13         | 14         | 14         | 15         | 15         | 14         | 15         | 14         | 13         | 13         | 13         | 13           | 12         | 13           | 13           | 13           | 13           | 13           | 13           | 12           | 12           | 12           | 12           |
| <i>NOF Steers</i>          | 12         | 12         | 13         | 14         | 13         | 14         | 14         | 13         | 13         | 12         | 11         | 11         | 11         | 10           | 10         | 10           | 11           | 10           | 10           | 11           | 10           | 10           | 9            | 9            | 9            |
| <i>NOF Cows</i>            | 69         | 69         | 70         | 71         | 74         | 76         | 75         | 74         | 76         | 76         | 71         | 73         | 71         | 71           | 71         | 73           | 75           | 73           | 70           | 70           | 71           | 71           | 69           | 64           | 63           |
| <b>Sheep</b>               | <b>7</b>   | <b>7</b>   | <b>7</b>   | <b>6</b>   | <b>6</b>   | <b>5</b>   | <b>5</b>   | <b>5</b>   | <b>5</b>   | <b>4</b>   | <b>4</b>   | <b>4</b>   | <b>4</b>   | <b>4</b>     | <b>3</b>   | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     | <b>3</b>     |
| <b>Goats</b>               | <b>1</b>     | <b>1</b>   | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     | <b>1</b>     |
| <b>Poultry</b>             | <b>131</b> | <b>131</b> | <b>127</b> | <b>131</b> | <b>131</b> | <b>128</b> | <b>125</b> | <b>128</b> | <b>130</b> | <b>126</b> | <b>127</b> | <b>131</b> | <b>129</b> | <b>130</b>   | <b>129</b> | <b>131</b>   | <b>134</b>   | <b>129</b>   | <b>128</b>   | <b>129</b>   | <b>127</b>   | <b>128</b>   | <b>128</b>   | <b>128</b>   | <b>130</b>   |
| <i>Hens &gt;1 yr.</i>      | 73         | 72         | 70         | 73         | 72         | 69         | 68         | 67         | 70         | 66         | 66         | 70         | 67         | 68           | 66         | 66           | 66           | 67           | 64           | 64           | 64           | 64           | 63           | 65           | 66           |
| <i>Total Pullets</i>       | 25         | 26         | 23         | 23         | 23         | 22         | 21         | 23         | 23         | 21         | 22         | 22         | 22         | 22           | 23         | 22           | 23           | 25           | 23           | 23           | 24           | 23           | 23           | 24           | 24           |
| <i>Chickens</i>            | 4          | 4          | 4          | 4          | 4          | 4          | 3          | 3          | 4          | 4          | 3          | 3          | 4          | 4            | 3          | 3            | 3            | 3            | 4            | 3            | 3            | 3            | 3            | 3            | 3            |
| <i>Broilers</i>            | 19         | 20         | 20         | 21         | 22         | 23         | 24         | 25         | 26         | 27         | 28         | 28         | 29         | 29           | 30         | 31           | 32           | 32           | 33           | 31           | 31           | 31           | 32           | 31           | 31           |
| <i>Turkeys</i>             | 10         | 10         | 10         | 10         | 9          | 9          | 9          | 9          | 8          | 7          | 7          | 7          | 7          | 7            | 7          | 7            | 7            | 7            | 7            | 6            | 6            | 6            | 6            | 6            | 6            |
| <b>Horses</b>              | <b>9</b>   | <b>9</b>   | <b>9</b>   | <b>9</b>   | <b>10</b>  | <b>11</b>  | <b>11</b>  | <b>12</b>  | <b>13</b>  | <b>13</b>  | <b>13</b>  | <b>13</b>  | <b>13</b>  | <b>13</b>    | <b>12</b>  | <b>12</b>    | <b>12</b>    | <b>11</b>    | <b>10</b>    | <b>10</b>    | <b>10</b>    | <b>10</b>    | <b>10</b>    | <b>9</b>     | <b>9</b>     |
| <b>Mules and Asses</b>     | <b>+</b>     | <b>+</b>   | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     |
| <b>American Bison</b>      | <b>+</b>     | <b>+</b>   | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     |

+ Does not exceed 0.5 kt.

<sup>a</sup> Accounts for CH<sub>4</sub> reductions due to capture and destruction of CH<sub>4</sub> at facilities using anaerobic digesters.

**Table A-214: Total (Direct and Indirect) Nitrous Oxide Emissions from Livestock Manure Management (kt)**

| Animal Type                | 1990        | 1991        | 1992        | 1993        | 1994        | 1995        | 1996        | 1997        | 1998        | 1999        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |            |
|----------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|------------|
| <b>Dairy Cattle</b>        | <b>17.7</b> | <b>17.6</b> | <b>17.7</b> | <b>17.9</b> | <b>18.0</b> | <b>18.2</b> | <b>18.3</b> | <b>18.4</b> | <b>18.5</b> | <b>18.1</b> | <b>18.4</b> | <b>18.7</b> | <b>18.9</b> | <b>19.1</b> | <b>18.2</b> | <b>18.7</b> | <b>19.3</b> | <b>19.3</b> | <b>19.0</b> | <b>19.2</b> | <b>19.3</b> | <b>19.5</b> | <b>19.7</b> | <b>19.6</b> | <b>19.7</b> |            |
| <i>Dairy Cows</i>          | 10.6        | 10.6        | 10.5        | 10.5        | 10.6        | 10.7        | 10.8        | 10.9        | 10.9        | 10.6        | 10.8        | 10.9        | 11.0        | 11.1        | 10.6        | 10.8        | 11.1        | 11.1        | 10.9        | 11.1        | 10.9        | 11.1        | 11.3        | 11.3        | 11.4        |            |
| <i>Dairy Heifer</i>        | 7.1         | 7.0         | 7.2         | 7.3         | 7.4         | 7.5         | 7.5         | 7.5         | 7.6         | 7.5         | 7.6         | 7.8         | 7.9         | 8.0         | 7.6         | 7.8         | 8.2         | 8.2         | 8.0         | 8.1         | 8.3         | 8.4         | 8.5         | 8.3         | 8.3         |            |
| <i>Dairy Calves</i>        | NA          |            |
| <b>Swine</b>               | <b>4.0</b>  | <b>4.2</b>  | <b>4.3</b>  | <b>4.4</b>  | <b>4.6</b>  | <b>4.5</b>  | <b>4.4</b>  | <b>4.7</b>  | <b>5.1</b>  | <b>5.0</b>  | <b>5.0</b>  | <b>5.1</b>  | <b>5.3</b>  | <b>5.4</b>  | <b>5.6</b>  | <b>5.7</b>  | <b>5.9</b>  | <b>6.3</b>  | <b>6.4</b>  | <b>6.3</b>  | <b>6.2</b>  | <b>6.3</b>  | <b>6.4</b>  | <b>6.3</b>  | <b>6.2</b>  |            |
| <i>Market Swine</i>        | 3.0         | 3.1         | 3.3         | 3.3         | 3.5         | 3.5         | 3.3         | 3.6         | 4.0         | 4.1         | 4.1         | 4.2         | 4.4         | 4.5         | 4.7         | 4.9         | 5.0         | 5.5         | 5.6         | 5.5         | 5.4         | 5.5         | 5.6         | 5.5         | 5.4         |            |
| <i>Market &lt;50 lbs.</i>  | 0.6         | 0.6         | 0.6         | 0.6         | 0.6         | 0.6         | 0.6         | 0.7         | 0.7         | 0.7         | 0.8         | 0.8         | 0.8         | 0.9         | 0.9         | 0.9         | 1.0         | 1.1         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         |            |
| <i>Market 50-119 lbs.</i>  | 0.6         | 0.7         | 0.7         | 0.7         | 0.7         | 0.7         | 0.7         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         | 0.9         | 0.9         | 0.9         | 1.0         | 1.0         | 1.1         | 1.3         | 1.2         | 1.2         | 1.2         | 1.3         | 1.2         | 1.2         |            |
| <i>Market 120-179 lbs.</i> | 0.9         | 0.9         | 0.9         | 1.0         | 1.0         | 1.0         | 1.0         | 1.0         | 1.1         | 1.1         | 1.1         | 1.2         | 1.2         | 1.3         | 1.3         | 1.4         | 1.4         | 1.5         | 1.6         | 1.6         | 1.6         | 1.6         | 1.6         | 1.6         | 1.6         |            |
| <i>Market &gt;180 lbs.</i> | 0.9         | 1.0         | 1.0         | 1.0         | 1.1         | 1.1         | 1.1         | 1.2         | 1.3         | 1.3         | 1.3         | 1.4         | 1.5         | 1.5         | 1.6         | 1.6         | 1.6         | 1.8         | 1.9         | 1.9         | 1.8         | 1.9         | 1.9         | 1.9         | 1.8         |            |
| <i>Breeding Swine</i>      | 1.0         | 1.1         | 1.1         | 1.1         | 1.1         | 1.1         | 1.0         | 1.1         | 1.1         | 1.0         | 0.9         | 0.9         | 0.9         | 0.9         | 0.9         | 0.9         | 0.9         | 0.9         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         |            |
| <b>Beef Cattle</b>         | <b>19.8</b> | <b>20.3</b> | <b>20.1</b> | <b>19.1</b> | <b>20.9</b> | <b>21.8</b> | <b>21.4</b> | <b>21.5</b> | <b>21.6</b> | <b>24.0</b> | <b>25.0</b> | <b>24.1</b> | <b>24.8</b> | <b>25.0</b> | <b>23.6</b> | <b>24.0</b> | <b>25.7</b> | <b>25.6</b> | <b>25.1</b> | <b>25.1</b> | <b>25.3</b> | <b>25.9</b> | <b>25.8</b> | <b>26.0</b> | <b>26.0</b> |            |
| <i>Feedlot Steers</i>      | 13.4        | 13.6        | 13.5        | 12.8        | 13.9        | 14.4        | 14.0        | 13.9        | 14.1        | 15.5        | 16.1        | 15.4        | 16.0        | 16.3        | 15.3        | 15.5        | 16.7        | 16.7        | 16.5        | 16.5        | 16.6        | 16.9        | 16.7        | 17.0        | 17.3        |            |
| <i>Feedlot Heifers</i>     | 6.4         | 6.6         | 6.6         | 6.3         | 7.0         | 7.4         | 7.4         | 7.6         | 7.6         | 8.5         | 8.9         | 8.6         | 8.7         | 8.8         | 8.4         | 8.5         | 9.0         | 8.9         | 8.7         | 8.6         | 8.7         | 9.1         | 9.0         | 9.0         | 8.8         |            |
| <b>Sheep</b>               | <b>0.4</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.6</b>  | <b>0.7</b>  | <b>0.8</b>  | <b>0.9</b>  | <b>0.9</b>  | <b>1.0</b>  | <b>1.1</b>  | <b>1.2</b>  | <b>1.2</b>  | <b>1.2</b>  | <b>1.1</b>  | <b>1.2</b>  | <b>1.2</b>  | <b>1.2</b>  | <b>1.1</b>  | <b>1.1</b>  | <b>1.1</b>  | <b>1.1</b>  | <b>1.0</b>  | <b>1.0</b>  | <b>1.0</b>  |            |
| <b>Goats</b>               | <b>0.1</b>  |            |
| <b>Poultry</b>             | <b>4.7</b>  | <b>4.8</b>  | <b>4.9</b>  | <b>5.0</b>  | <b>5.1</b>  | <b>5.1</b>  | <b>5.3</b>  | <b>5.3</b>  | <b>5.3</b>  | <b>5.3</b>  | <b>5.3</b>  | <b>5.3</b>  | <b>5.4</b>  | <b>5.3</b>  | <b>5.4</b>  | <b>5.4</b>  | <b>5.4</b>  | <b>5.4</b>  | <b>5.4</b>  | <b>5.4</b>  | <b>5.2</b>  | <b>5.2</b>  | <b>5.2</b>  | <b>5.3</b>  | <b>5.2</b>  | <b>5.2</b> |
| <i>Hens &gt;1 yr.</i>      | 1.0         | 1.0         | 1.0         | 1.0         | 1.0         | 1.0         | 1.0         | 1.1         | 1.1         | 1.1         | 1.1         | 1.2         | 1.2         | 1.2         | 1.2         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.3         | 1.4         |            |
| <i>Total Pullets</i>       | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         |            |
| <i>Chickens</i>            | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           |            |
| <i>Broilers</i>            | 2.2         | 2.3         | 2.4         | 2.5         | 2.6         | 2.7         | 2.8         | 2.8         | 2.9         | 2.9         | 2.9         | 2.9         | 3.0         | 2.9         | 2.9         | 3.0         | 2.9         | 2.9         | 2.9         | 2.9         | 2.7         | 2.8         | 2.8         | 2.9         | 2.7         |            |
| <i>Turkeys</i>             | 1.2         | 1.2         | 1.2         | 1.1         | 1.1         | 1.1         | 1.1         | 1.1         | 1.0         | 0.9         | 0.9         | 0.9         | 0.9         | 0.9         | 0.8         | 0.8         | 0.8         | 0.8         | 0.8         | 0.7         | 0.7         | 0.7         | 0.7         | 0.7         | 0.7         |            |
| <b>Horses</b>              | <b>0.3</b>  | <b>0.3</b>  | <b>0.3</b>  | <b>0.3</b>  | <b>0.3</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.4</b>  | <b>0.5</b>  | <b>0.4</b>  |            |
| <b>Mules and Asses</b>     | <b>+</b>    |            |
| <b>American Bison</b>      | <b>NA</b>   |            |

+ Does not exceed 0.5 kt.

NA (Not applicable)

Note: American bison are maintained entirely on unmanaged WMS; there are no American bison N<sub>2</sub>O emissions from managed systems.

**Table A-215: Methane Emissions by State from Livestock Manure Management for 2014 (kt)<sup>a</sup>**

| State          | Beef on Feedlots | Beef Not on Feed <sup>b</sup> | Dairy Cow | Dairy Heifer | Swine—Market | Swine—Breeding | Layer   | Broiler | Turkey | Sheep  | Goats  | Horses | Mules and Asses | American Bison |
|----------------|------------------|-------------------------------|-----------|--------------|--------------|----------------|---------|---------|--------|--------|--------|--------|-----------------|----------------|
| Alabama        | 0.0170           | 2.4961                        | 0.5233    | 0.0106       | 1.6284       | 0.4402         | 8.9934  | 3.8556  | 0.0211 | 0.0085 | 0.0156 | 0.1785 | 0.0131          | 0.0006         |
| Alaska         | 0.0001           | 0.0154                        | 0.0154    | 0.0002       | 0.0018       | 0.0015         | 0.1926  | +       | 0.0210 | 0.0057 | 0.0002 | 0.0029 | +               | 0.0038         |
| Arizona        | 0.6906           | 1.0848                        | 52.7906   | 0.1741       | 1.9359       | 0.7272         | 0.6745  | +       | 0.0211 | 0.1057 | 0.0313 | 0.3346 | 0.0039          | +              |
| Arkansas       | 0.0296           | 2.1448                        | 0.2667    | 0.0094       | 0.6193       | 1.9556         | 0.5434  | 3.5225  | 0.7503 | 0.0085 | 0.0143 | 0.1772 | 0.0094          | 0.0003         |
| California     | 1.5156           | 3.8144                        | 402.0657  | 1.9264       | 1.6039       | 0.1800         | 3.0921  | 0.1985  | 0.2751 | 0.4158 | 0.0539 | 0.4182 | 0.0073          | 0.0035         |
| Colorado       | 1.5724           | 2.7254                        | 31.0978   | 0.1501       | 4.0379       | 2.8453         | 4.0236  | +       | 0.0210 | 0.1715 | 0.0073 | 0.2341 | 0.0047          | 0.0194         |
| Connecticut    | 0.0003           | 0.0158                        | 1.0347    | 0.0153       | 0.0049       | 0.0036         | 0.2501  | +       | 0.0210 | 0.0034 | 0.0011 | 0.0433 | 0.0007          | 0.0002         |
| Delaware       | 0.0003           | 0.0092                        | 0.2790    | 0.0046       | 0.0269       | 0.0218         | 0.0754  | 0.8837  | 0.0210 | 0.0057 | 0.0003 | 0.0154 | 0.0001          | 0.0002         |
| Florida        | 0.0108           | 3.2376                        | 22.1675   | 0.1010       | 0.0635       | 0.0424         | 6.4911  | 0.2423  | 0.0211 | 0.0085 | 0.0187 | 0.3982 | 0.0108          | 0.0003         |
| Georgia        | 0.0124           | 1.8294                        | 7.6846    | 0.0667       | 1.9133       | 0.9216         | 15.6498 | 4.8098  | 0.0211 | 0.0085 | 0.0250 | 0.2205 | 0.0099          | 0.0006         |
| Hawaii         | 0.0024           | 0.2895                        | 0.4264    | 0.0029       | 0.0686       | 0.0639         | 0.3014  | +       | 0.0211 | 0.0085 | 0.0054 | 0.0149 | 0.0005          | 0.0002         |
| Idaho          | 0.3828           | 1.6793                        | 125.0109  | 0.4087       | 0.1619       | 0.0987         | 0.6159  | +       | 0.0210 | 0.1175 | 0.0046 | 0.1235 | 0.0029          | 0.0093         |
| Illinois       | 0.4437           | 1.0049                        | 8.1855    | 0.0725       | 40.2193      | 10.3785        | 0.2649  | 0.1978  | 0.0210 | 0.0263 | 0.0077 | 0.1227 | 0.0027          | 0.0008         |
| Indiana        | 0.1800           | 0.5545                        | 14.3566   | 0.1086       | 32.9344      | 5.5527         | 0.8942  | 0.1978  | 0.4736 | 0.0235 | 0.0088 | 0.2275 | 0.0040          | 0.0020         |
| Iowa           | 2.1095           | 3.0403                        | 20.9373   | 0.1904       | 293.3499     | 31.0226        | 1.9147  | 0.1978  | 0.2617 | 0.0728 | 0.0141 | 0.1277 | 0.0033          | 0.0026         |
| Kansas         | 3.7901           | 4.8366                        | 23.3681   | 0.1634       | 18.5299      | 3.8114         | 0.0440  | +       | 0.0210 | 0.0352 | 0.0099 | 0.1508 | 0.0028          | 0.0097         |
| Kentucky       | 0.0343           | 2.5216                        | 1.4173    | 0.0789       | 4.0954       | 1.2243         | 0.6210  | 1.1150  | 0.0210 | 0.0230 | 0.0126 | 0.2812 | 0.0097          | 0.0025         |
| Louisiana      | 0.0087           | 1.6486                        | 0.5940    | 0.0138       | 0.0130       | 0.0093         | 2.3022  | 0.1985  | 0.0211 | 0.0085 | 0.0066 | 0.1955 | 0.0082          | 0.0001         |
| Maine          | 0.0008           | 0.0383                        | 1.3343    | 0.0280       | 0.0102       | 0.0068         | 0.2830  | +       | 0.0210 | 0.0034 | 0.0017 | 0.0261 | 0.0003          | 0.0004         |
| Maryland       | 0.0193           | 0.1179                        | 2.4502    | 0.0436       | 0.1827       | 0.0640         | 0.3077  | 1.0419  | 0.0210 | 0.0057 | 0.0021 | 0.0610 | 0.0009          | 0.0006         |
| Massachusetts  | 0.0003           | 0.0201                        | 0.4851    | 0.0126       | 0.0303       | 0.0186         | 0.0127  | +       | 0.0210 | 0.0034 | 0.0022 | 0.0443 | 0.0004          | 0.0002         |
| Michigan       | 0.2545           | 0.4752                        | 48.2147   | 0.2582       | 8.7176       | 1.9544         | 0.7766  | 0.1978  | 0.1271 | 0.0381 | 0.0067 | 0.1812 | 0.0031          | 0.0022         |
| Minnesota      | 0.6363           | 1.2341                        | 31.5912   | 0.4408       | 63.2767      | 10.6950        | 0.3802  | 0.1694  | 1.1341 | 0.0634 | 0.0081 | 0.1246 | 0.0022          | 0.0034         |
| Mississippi    | 0.0191           | 1.8134                        | 0.5460    | 0.0161       | 8.7714       | 1.6819         | 7.6425  | 2.6413  | 0.0211 | 0.0085 | 0.0083 | 0.1841 | 0.0099          | +              |
| Missouri       | 0.1323           | 4.3993                        | 5.6464    | 0.0811       | 19.5727      | 7.7160         | 0.3223  | 1.0444  | 0.4237 | 0.0390 | 0.0266 | 0.2290 | 0.0072          | 0.0025         |
| Montana        | 0.0841           | 4.4117                        | 1.7741    | 0.0134       | 1.1634       | 0.3959         | 0.4143  | +       | 0.0210 | 0.1034 | 0.0024 | 0.2081 | 0.0035          | 0.0324         |
| Nebraska       | 4.1931           | 5.9583                        | 7.3544    | 0.0320       | 25.7524      | 7.9160         | 0.5798  | 0.1978  | 0.0210 | 0.0357 | 0.0056 | 0.1398 | 0.0029          | 0.0467         |
| Nevada         | 0.0065           | 0.6579                        | 6.8818    | 0.0137       | 0.0126       | 0.0073         | 0.0224  | +       | 0.0210 | 0.0376 | 0.0063 | 0.0528 | 0.0004          | 0.0001         |
| New Hampshire  | 0.0002           | 0.0116                        | 0.6205    | 0.0116       | 0.0103       | 0.0057         | 0.0693  | +       | 0.0210 | 0.0034 | 0.0013 | 0.0192 | 0.0002          | 0.0006         |
| New Jersey     | 0.0004           | 0.0204                        | 0.2564    | 0.0052       | 0.0671       | 0.0150         | 0.0744  | +       | 0.0210 | 0.0057 | 0.0018 | 0.0584 | 0.0007          | 0.0004         |
| New Mexico     | 0.0173           | 1.2629                        | 77.5423   | 0.1849       | 0.0002       | 0.0003         | 0.6320  | +       | 0.0210 | 0.0381 | 0.0073 | 0.1086 | 0.0014          | 0.0117         |
| New York       | 0.0425           | 0.4557                        | 31.2877   | 0.5901       | 0.4865       | 0.1467         | 0.5324  | 0.1978  | 0.0210 | 0.0352 | 0.0088 | 0.2020 | 0.0028          | 0.0013         |
| North Carolina | 0.0083           | 0.9086                        | 2.4938    | 0.0352       | 119.6529     | 30.7160        | 11.8916 | 2.8787  | 0.7104 | 0.0127 | 0.0134 | 0.1364 | 0.0070          | 0.0003         |
| North Dakota   | 0.0677           | 2.3399                        | 1.2266    | 0.0157       | 0.7321       | 0.6034         | 0.0416  | +       | 0.0210 | 0.0310 | 0.0012 | 0.0996 | 0.0009          | 0.0130         |
| Ohio           | 0.2740           | 0.8580                        | 19.7712   | 0.2071       | 17.8795      | 3.2873         | 1.0144  | 0.2737  | 0.1271 | 0.0550 | 0.0111 | 0.2455 | 0.0053          | 0.0012         |
| Oklahoma       | 0.4728           | 4.9499                        | 7.8066    | 0.0454       | 25.9088      | 15.5288        | 3.5335  | 0.7432  | 0.0210 | 0.0277 | 0.0186 | 0.3423 | 0.0102          | 0.0182         |

| State          | Beef on Feedlots | Beef Not on Feed <sup>b</sup> | Dairy Cow | Dairy Heifer | Swine—Market | Swine—Breeding | Layer  | Broiler | Turkey | Sheep  | Goats  | Horses | Mules and Asses | American Bison |
|----------------|------------------|-------------------------------|-----------|--------------|--------------|----------------|--------|---------|--------|--------|--------|--------|-----------------|----------------|
| Oregon         | 0.1394           | 1.5941                        | 17.9099   | 0.1025       | 0.0301       | 0.0195         | 0.8158 | 0.1978  | 0.0210 | 0.0916 | 0.0078 | 0.1376 | 0.0025          | 0.0033         |
| Pennsylvania   | 0.2089           | 0.6687                        | 16.7144   | 0.5361       | 10.2501      | 1.9201         | 0.7949 | 0.6563  | 0.1745 | 0.0442 | 0.0116 | 0.2657 | 0.0071          | 0.0015         |
| Rhode Island   | 0.0001           | 0.0044                        | 0.0290    | 0.0009       | 0.0032       | 0.0032         | 0.0729 | +       | 0.0210 | 0.0034 | 0.0002 | 0.0044 | 0.0001          | +              |
| South Carolina | 0.0042           | 0.6335                        | 1.1129    | 0.0158       | 4.7818       | 0.4586         | 4.5615 | 0.8445  | 0.0211 | 0.0085 | 0.0138 | 0.1841 | 0.0065          | 0.0004         |
| South Dakota   | 0.6473           | 4.3569                        | 12.3807   | 0.0790       | 10.1884      | 3.3233         | 0.1697 | +       | 0.1122 | 0.1269 | 0.0047 | 0.1498 | 0.0012          | 0.0584         |
| Tennessee      | 0.0058           | 2.1576                        | 1.2620    | 0.0527       | 2.5978       | 0.5478         | 0.2375 | 0.6538  | 0.0210 | 0.0183 | 0.0190 | 0.1716 | 0.0107          | 0.0002         |
| Texas          | 6.1094           | 17.0779                       | 100.3698  | 0.5029       | 10.2790      | 3.1653         | 4.5570 | 2.1495  | 0.0211 | 0.5145 | 0.2903 | 1.2439 | 0.0711          | 0.0110         |
| Utah           | 0.0420           | 1.0336                        | 18.1451   | 0.0691       | 5.3791       | 1.2998         | 3.5876 | +       | 0.0997 | 0.1316 | 0.0034 | 0.1285 | 0.0024          | 0.0023         |
| Vermont        | 0.0011           | 0.0658                        | 5.9194    | 0.0921       | 0.0084       | 0.0072         | 0.0121 | +       | 0.0210 | 0.0034 | 0.0030 | 0.0241 | 0.0010          | 0.0002         |
| Virginia       | 0.0391           | 1.6537                        | 2.8422    | 0.0689       | 4.5280       | 0.1581         | 0.3495 | 0.9485  | 0.4187 | 0.0390 | 0.0115 | 0.1871 | 0.0053          | 0.0019         |
| Washington     | 0.3929           | 0.8259                        | 47.0772   | 0.2127       | 0.1092       | 0.0651         | 1.3173 | 0.1978  | 0.0210 | 0.0258 | 0.0062 | 0.1195 | 0.0027          | 0.0016         |
| West Virginia  | 0.0089           | 0.4978                        | 0.3084    | 0.0085       | 0.0154       | 0.0088         | 0.1706 | 0.3450  | 0.0773 | 0.0150 | 0.0038 | 0.0481 | 0.0022          | +              |
| Wisconsin      | 0.4381           | 1.0851                        | 93.9587   | 1.0665       | 2.1879       | 0.6948         | 0.3439 | 0.1933  | 0.0210 | 0.0390 | 0.0158 | 0.2121 | 0.0043          | 0.0065         |
| Wyoming        | 0.1190           | 2.0593                        | 0.8553    | 0.0060       | 0.1890       | 0.3462         | 0.5771 | +       | 0.0210 | 0.1668 | 0.0024 | 0.1517 | 0.0020          | 0.0184         |

+ Does not exceed 0.00005 kt.

<sup>a</sup> Accounts for CH<sub>4</sub> reductions due to capture and destruction of CH<sub>4</sub> at facilities using anaerobic digesters.

<sup>b</sup> Beef Not on Feed includes calves.

**Table A-216: Nitrous Oxide Emissions by State from Livestock Manure Management for 2014 (kt)**

|                | Beef<br>Feedlot-<br>Heifer | Beef<br>Feedlot-<br>Steers | Dairy<br>Cow | Dairy<br>Heifer | Swine-<br>Market | Swine-<br>Breeding | Layer  | Broiler | Turkey | Sheep  | Goats  | Horses | Mules<br>and<br>Asses |
|----------------|----------------------------|----------------------------|--------------|-----------------|------------------|--------------------|--------|---------|--------|--------|--------|--------|-----------------------|
| Alabama        | 0.0039                     | 0.0077                     | 0.0040       | 0.0037          | 0.0085           | 0.0017             | 0.0650 | 0.3411  | 0.0024 | 0.0046 | 0.0012 | 0.0061 | 0.0005                |
| Alaska         | +                          | +                          | 0.0003       | 0.0002          | +                | +                  | 0.0033 | +       | 0.0024 | 0.0015 | +      | 0.0001 | +                     |
| Arizona        | 0.1828                     | 0.3594                     | 0.2417       | 0.1565          | 0.0091           | 0.0026             | 0.0035 | +       | 0.0024 | 0.0165 | 0.0025 | 0.0115 | 0.0001                |
| Arkansas       | 0.0094                     | 0.0184                     | 0.0029       | 0.0034          | 0.0036           | 0.0084             | 0.0766 | 0.3116  | 0.0869 | 0.0040 | 0.0011 | 0.0061 | 0.0003                |
| California     | 0.3396                     | 0.6690                     | 2.1973       | 1.5656          | 0.0090           | 0.0007             | 0.0651 | 0.0176  | 0.0319 | 0.0735 | 0.0043 | 0.0144 | 0.0003                |
| Colorado       | 0.6505                     | 1.2767                     | 0.1943       | 0.2294          | 0.0429           | 0.0223             | 0.0242 | +       | 0.0024 | 0.0402 | 0.0009 | 0.0121 | 0.0003                |
| Connecticut    | 0.0001                     | 0.0002                     | 0.0172       | 0.0112          | +                | +                  | 0.0109 | +       | 0.0024 | 0.0028 | 0.0001 | 0.0022 | +                     |
| Delaware       | 0.0001                     | 0.0002                     | 0.0042       | 0.0032          | 0.0002           | 0.0001             | 0.0031 | 0.0784  | 0.0024 | 0.0046 | +      | 0.0008 | +                     |
| Florida        | 0.0023                     | 0.0045                     | 0.1122       | 0.0512          | 0.0003           | 0.0002             | 0.0439 | 0.0214  | 0.0024 | 0.0046 | 0.0015 | 0.0137 | 0.0004                |
| Georgia        | 0.0029                     | 0.0057                     | 0.0552       | 0.0252          | 0.0100           | 0.0036             | 0.1128 | 0.4255  | 0.0024 | 0.0046 | 0.0020 | 0.0076 | 0.0004                |
| Hawaii         | 0.0005                     | 0.0010                     | 0.0022       | 0.0023          | 0.0004           | 0.0002             | 0.0033 | +       | 0.0024 | 0.0015 | 0.0004 | 0.0005 | +                     |
| Idaho          | 0.1593                     | 0.3126                     | 0.7753       | 0.6240          | 0.0018           | 0.0008             | 0.0035 | +       | 0.0024 | 0.0276 | 0.0005 | 0.0064 | 0.0002                |
| Illinois       | 0.1728                     | 0.3401                     | 0.1380       | 0.0920          | 0.3798           | 0.0716             | 0.0190 | 0.0176  | 0.0024 | 0.0184 | 0.0009 | 0.0063 | 0.0001                |
| Indiana        | 0.0695                     | 0.1377                     | 0.2528       | 0.1261          | 0.3206           | 0.0396             | 0.1242 | 0.0176  | 0.0551 | 0.0164 | 0.0010 | 0.0117 | 0.0002                |
| Iowa           | 0.8306                     | 1.6316                     | 0.3025       | 0.2352          | 1.7882           | 0.1387             | 0.2658 | 0.0176  | 0.0304 | 0.0508 | 0.0017 | 0.0066 | 0.0002                |
| Kansas         | 1.4446                     | 2.8361                     | 0.1995       | 0.2154          | 0.1617           | 0.0246             | 0.0031 | +       | 0.0024 | 0.0246 | 0.0012 | 0.0078 | 0.0001                |
| Kentucky       | 0.0120                     | 0.0236                     | 0.0296       | 0.0265          | 0.0240           | 0.0053             | 0.0259 | 0.0990  | 0.0024 | 0.0187 | 0.0015 | 0.0145 | 0.0005                |
| Louisiana      | 0.0019                     | 0.0037                     | 0.0057       | 0.0031          | 0.0001           | +                  | 0.0118 | 0.0176  | 0.0024 | 0.0040 | 0.0005 | 0.0067 | 0.0003                |
| Maine          | 0.0003                     | 0.0006                     | 0.0263       | 0.0200          | 0.0001           | 0.0001             | 0.0129 | +       | 0.0024 | 0.0028 | 0.0002 | 0.0013 | +                     |
| Maryland       | 0.0068                     | 0.0133                     | 0.0443       | 0.0300          | 0.0016           | 0.0004             | 0.0128 | 0.0925  | 0.0024 | 0.0046 | 0.0002 | 0.0031 | +                     |
| Massachusetts  | 0.0001                     | 0.0002                     | 0.0103       | 0.0087          | 0.0003           | 0.0001             | 0.0006 | +       | 0.0024 | 0.0028 | 0.0003 | 0.0023 | +                     |
| Michigan       | 0.1013                     | 0.1990                     | 0.5981       | 0.3485          | 0.0915           | 0.0153             | 0.0578 | 0.0176  | 0.0148 | 0.0266 | 0.0008 | 0.0093 | 0.0002                |
| Minnesota      | 0.2529                     | 0.4969                     | 0.6511       | 0.5530          | 0.6434           | 0.0801             | 0.0528 | 0.0150  | 0.1319 | 0.0443 | 0.0010 | 0.0064 | 0.0001                |
| Mississippi    | 0.0043                     | 0.0085                     | 0.0058       | 0.0041          | 0.0448           | 0.0062             | 0.0399 | 0.2337  | 0.0024 | 0.0046 | 0.0007 | 0.0063 | 0.0004                |
| Missouri       | 0.0510                     | 0.1002                     | 0.1086       | 0.0910          | 0.1885           | 0.0542             | 0.0449 | 0.0927  | 0.0493 | 0.0272 | 0.0032 | 0.0118 | 0.0004                |
| Montana        | 0.0349                     | 0.0689                     | 0.0191       | 0.0196          | 0.0134           | 0.0034             | 0.0025 | +       | 0.0024 | 0.0243 | 0.0003 | 0.0107 | 0.0002                |
| Nebraska       | 1.6420                     | 3.2236                     | 0.0792       | 0.0423          | 0.2495           | 0.0564             | 0.0417 | 0.0176  | 0.0024 | 0.0249 | 0.0007 | 0.0072 | 0.0002                |
| Nevada         | 0.0027                     | 0.0053                     | 0.0398       | 0.0209          | 0.0001           | +                  | 0.0031 | +       | 0.0024 | 0.0088 | 0.0007 | 0.0027 | +                     |
| New Hampshire  | 0.0001                     | 0.0001                     | 0.0120       | 0.0084          | 0.0001           | +                  | 0.0031 | +       | 0.0024 | 0.0028 | 0.0002 | 0.0010 | +                     |
| New Jersey     | 0.0001                     | 0.0003                     | 0.0059       | 0.0035          | 0.0006           | 0.0001             | 0.0031 | +       | 0.0024 | 0.0046 | 0.0002 | 0.0030 | +                     |
| New Mexico     | 0.0070                     | 0.0138                     | 0.4120       | 0.2537          | +                | +                  | 0.0035 | +       | 0.0024 | 0.0089 | 0.0009 | 0.0056 | 0.0001                |
| New York       | 0.0158                     | 0.0311                     | 0.5698       | 0.4185          | 0.0051           | 0.0011             | 0.0237 | 0.0176  | 0.0024 | 0.0286 | 0.0010 | 0.0104 | 0.0001                |
| North Carolina | 0.0029                     | 0.0057                     | 0.0288       | 0.0147          | 0.6303           | 0.1194             | 0.0875 | 0.2555  | 0.0826 | 0.0103 | 0.0016 | 0.0070 | 0.0004                |
| North Dakota   | 0.0275                     | 0.0535                     | 0.0229       | 0.0194          | 0.0079           | 0.0048             | 0.0031 | +       | 0.0024 | 0.0217 | 0.0001 | 0.0051 | +                     |
| Ohio           | 0.1073                     | 0.2109                     | 0.3649       | 0.2401          | 0.1756           | 0.0237             | 0.1407 | 0.0243  | 0.0148 | 0.0443 | 0.0013 | 0.0127 | 0.0003                |

|                |        |        |        |        |        |        |        |        |        |        |        |        |        |
|----------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Oklahoma       | 0.1892 | 0.3689 | 0.0547 | 0.0438 | 0.1336 | 0.0584 | 0.0184 | 0.0660 | 0.0024 | 0.0193 | 0.0022 | 0.0176 | 0.0005 |
| Oregon         | 0.0492 | 0.0971 | 0.1396 | 0.1126 | 0.0003 | 0.0002 | 0.0107 | 0.0176 | 0.0024 | 0.0243 | 0.0009 | 0.0071 | 0.0001 |
| Pennsylvania   | 0.0756 | 0.1484 | 0.4531 | 0.3435 | 0.0984 | 0.0138 | 0.1105 | 0.0583 | 0.0203 | 0.0358 | 0.0014 | 0.0137 | 0.0004 |
| Rhode Island   | +      | 0.0001 | 0.0008 | 0.0005 | +      | +      | 0.0031 | +      | 0.0024 | 0.0028 | +      | 0.0002 | +      |
| South Carolina | 0.0010 | 0.0019 | 0.0084 | 0.0044 | 0.0245 | 0.0017 | 0.0239 | 0.0747 | 0.0024 | 0.0046 | 0.0011 | 0.0063 | 0.0002 |
| South Dakota   | 0.2567 | 0.5034 | 0.1393 | 0.1025 | 0.1012 | 0.0243 | 0.0124 | +      | 0.0130 | 0.0886 | 0.0006 | 0.0077 | 0.0001 |
| Tennessee      | 0.0020 | 0.0040 | 0.0208 | 0.0190 | 0.0155 | 0.0024 | 0.0100 | 0.0580 | 0.0024 | 0.0149 | 0.0023 | 0.0088 | 0.0006 |
| Texas          | 1.6794 | 3.2913 | 0.5474 | 0.4745 | 0.0583 | 0.0132 | 0.0930 | 0.1902 | 0.0024 | 0.0805 | 0.0229 | 0.0427 | 0.0025 |
| Utah           | 0.0174 | 0.0342 | 0.1320 | 0.1051 | 0.0546 | 0.0108 | 0.0214 | +      | 0.0116 | 0.0309 | 0.0004 | 0.0066 | 0.0001 |
| Vermont        | 0.0004 | 0.0008 | 0.1176 | 0.0671 | 0.0001 | 0.0001 | 0.0006 | +      | 0.0024 | 0.0028 | 0.0004 | 0.0012 | 0.0001 |
| Virginia       | 0.0140 | 0.0273 | 0.0462 | 0.0267 | 0.0254 | 0.0007 | 0.0149 | 0.0842 | 0.0487 | 0.0316 | 0.0014 | 0.0096 | 0.0003 |
| Washington     | 0.1399 | 0.2730 | 0.3354 | 0.2451 | 0.0011 | 0.0005 | 0.0311 | 0.0176 | 0.0024 | 0.0068 | 0.0007 | 0.0062 | 0.0001 |
| West Virginia  | 0.0032 | 0.0064 | 0.0070 | 0.0059 | 0.0001 | 0.0001 | 0.0075 | 0.0306 | 0.0090 | 0.0122 | 0.0005 | 0.0025 | 0.0001 |
| Wisconsin      | 0.1743 | 0.3427 | 1.8468 | 1.3065 | 0.0220 | 0.0051 | 0.0256 | 0.0172 | 0.0024 | 0.0272 | 0.0019 | 0.0109 | 0.0002 |
| Wyoming        | 0.0493 | 0.0972 | 0.0077 | 0.0076 | 0.0034 | 0.0046 | 0.0035 | +      | 0.0024 | 0.0391 | 0.0003 | 0.0078 | 0.0001 |

+ Does not exceed 0.00005 kt.

## References

- Anderson, S. (2000) Personal Communication. Steve Anderson, Agricultural Statistician, National Agriculture Statistics Service, U.S. Department of Agriculture and Lee-Ann Tracy, ERG. Washington, D.C. May 31, 2000.
- ASAE (1998) *ASAE Standards 1998, 45<sup>th</sup> Edition*. American Society of Agricultural Engineers. St. Joseph, MI. Bryant, M.P., V.H. Varel, R.A. Frobish, and H.R. Isaacson (1976) In H.G. Schlegel (ed.); *Seminar on Microbial Energy Conversion*. E. Goltz KG. Göttingen, Germany.
- Bush, E. (1998) Personal communication with Eric Bush, Centers for Epidemiology and Animal Health, U.S. Department of Agriculture regarding *National Animal Health Monitoring System's (NAHMS) Swine '95 Study*.
- Deal, P. (2000) Personal Communication. Peter B. Deal, Rangeland Management Specialist, Florida Natural Resource Conservation Service and Lee-Ann Tracy, ERG. June 21, 2000.
- EPA (2012) AgSTAR Anaerobic Digester Database. Available online at: <<http://www.epa.gov/agstar/projects/index.html#database>>.
- EPA (2008) *Climate Leaders Greenhouse Gas Inventory Protocol Offset Project Methodology for Project Type Managing Manure with Biogas Recovery Systems*. Available online at: <[http://www.epa.gov/climateleaders/documents/resources/ClimateLeaders\\_DraftManureOffsetProtocol.pdf](http://www.epa.gov/climateleaders/documents/resources/ClimateLeaders_DraftManureOffsetProtocol.pdf)>.
- EPA (2006) *AgSTAR Digest*. Office of Air and Radiation, U.S. Environmental Protection Agency. Washington, D.C. Winter 2006. Available online at: <<http://www.epa.gov/agstar/pdf/2006digest.pdf>>. Retrieved July 2006.
- EPA (2005) *National Emission Inventory—Ammonia Emissions from Animal Agricultural Operations, Revised Draft Report*. U.S. Environmental Protection Agency. Washington, D.C. April 22, 2005. Available online at: <[ftp://ftp.epa.gov/EmisInventory/2002finalnei/documentation/nonpoint/nh3inventory\\_draft\\_042205.pdf](ftp://ftp.epa.gov/EmisInventory/2002finalnei/documentation/nonpoint/nh3inventory_draft_042205.pdf)>. Retrieved August 2007.
- EPA (2003) *AgSTAR Digest*. Office of Air and Radiation, U.S. Environmental Protection Agency. Washington, D.C. Winter 2003. Available online at: <<http://www.epa.gov/agstar/pdf/2003digest.pdf>>. Retrieved July 2006.
- EPA (2002a) *Development Document for the Final Revisions to the National Pollutant Discharge Elimination System (NPDES) Regulation and the Effluent Guidelines for Concentrated Animal Feeding Operations (CAFOS)*. U.S. Environmental Protection Agency. EPA-821-R-03-001. December 2002.
- EPA (2002b) *Cost Methodology for the Final Revisions to the National Pollutant Discharge Elimination System Regulation and the Effluent Guidelines for Concentrated Animal Feeding Operations*. U.S. Environmental Protection Agency. EPA-821-R-03-004. December 2002.
- EPA (2000) *AgSTAR Digest*. Office of Air and Radiation, U.S. Environmental Protection Agency. Washington, D.C. Spring 2000. Available online at: <<http://www.epa.gov/agstar/news-events/digest/2000digest.pdf>>.
- EPA (1992) *Global Methane Emissions from Livestock and Poultry Manure*, Office of Air and Radiation, U.S. Environmental Protection Agency. February 1992.
- ERG (2010a) "Typical Animal Mass Values for Inventory Swine Categories." Memorandum to EPA from ERG. July 19, 2010.
- ERG (2010b) Telecon with William Boyd of USDA NRCS and Cortney Itle of ERG Concerning Updated VS and Nex Rates. August 8, 2010.
- ERG (2010c) "Updating Current Inventory Manure Characteristics new USDA Agricultural Waste Management Field Handbook Values." Memorandum to EPA from ERG. August 13, 2010.
- ERG (2008) "Methodology for Improving Methane Emissions Estimates and Emission Reductions from Anaerobic Digestion System for the 1990-2007 Greenhouse Gas Inventory for Manure Management." Memorandum to EPA from ERG. August 18, 2008.
- ERG (2003a) "Methodology for Estimating Uncertainty for Manure Management Greenhouse Gas Inventory." Contract No. GS-10F-0036, Task Order 005. Memorandum to EPA from ERG, Lexington, MA. September 26, 2003.
- ERG (2003b) "Changes to Beef Calves and Beef Cows Typical Animal Mass in the Manure Management Greenhouse Gas Inventory." Memorandum to EPA from ERG, October 7, 2003.

- ERG (2001) *Summary of development of MDP Factor for methane conversion factor calculations*. ERG, Lexington, MA. September 2001.
- ERG (2000a) *Calculations: Percent Distribution of Manure for Waste Management Systems*. ERG, Lexington, MA. August 2000.
- ERG (2000b) *Discussion of Methodology for Estimating Animal Waste Characteristics* (Summary of B<sub>0</sub> Literature Review). ERG, Lexington, MA. June 2000.
- Garrett, W.N. and D.E. Johnson (1983) "Nutritional energetics of ruminants." *Journal of Animal Science*, 57(suppl.2):478-497.
- Groffman, P.M., R. Brumme, K. Butterbach-Bahl, K.E. Dobbie, A.R. Mosier, D. Ojima, H. Papen, W.J. Parton, K.A. Smith, and C. Wagner-Riddle (2000) "Evaluating annual nitrous oxide fluxes at the ecosystem scale." *Global Biogeochemical Cycles*, 14(4):1061-1070.
- Hashimoto, A.G. (1984) "Methane from Swine Manure: Effect of Temperature and Influent Substrate Composition on Kinetic Parameter (k)." *Agricultural Wastes*, 9:299-308.
- Hashimoto, A.G., V.H. Varel, and Y.R. Chen (1981) "Ultimate Methane Yield from Beef Cattle Manure; Effect of Temperature, Ration Constituents, Antibiotics and Manure Age." *Agricultural Wastes*, 3:241-256.
- Hill, D.T. (1984) "Methane Productivity of the Major Animal Types." *Transactions of the ASAE*, 27(2):530-540.
- Hill, D.T. (1982) "Design of Digestion Systems for Maximum Methane Production." *Transactions of the ASAE*, 25(1):226-230.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- Johnson, D. (2000) Personal Communication. Dan Johnson, State Water Management Engineer, California Natural Resource Conservation Service and Lee-Ann Tracy, ERG. June 23, 2000.
- Lange, J. (2000) Personal Communication. John Lange, Agricultural Statistician, U.S. Department of Agriculture, National Agriculture Statistics Service and Lee-Ann Tracy, ERG. Washington, D.C. May 8, 2000.
- Meagher, M. (1986). *Bison. Mammalian Species*. 266: 1-8.
- Miller, P. (2000) Personal Communication. Paul Miller, Iowa Natural Resource Conservation Service and Lee-Ann Tracy, ERG. June 12, 2000.
- Milton, B. (2000) Personal Communication. Bob Milton, Chief of Livestock Branch, U.S. Department of Agriculture, National Agriculture Statistics Service and Lee-Ann Tracy, ERG. May 1, 2000.
- Moffroid, K. and D. Pape. (2014) *1990-2013 Volatile Solids and Nitrogen Excretion Rates*. Dataset to EPA from ICF International. August 2014.
- Morris, G.R. (1976) *Anaerobic Fermentation of Animal Wastes: A Kinetic and Empirical Design Fermentation*. M.S. Thesis. Cornell University.
- National Bison Association (1999) Total Bison Population—1999. Report provided during personal email communication with Dave Carter, Executive Director, National Bison Association July 19, 2011.
- NOAA (2014) *National Climate Data Center (NCDC)*. Available online at: [<ftp://ftp.ncdc.noaa.gov/pub/data/cirs/climdiv/>](ftp://ftp.ncdc.noaa.gov/pub/data/cirs/climdiv/) (for all states except Alaska and Hawaii) and [<ftp://ftp.ncdc.noaa.gov/pub/data/g sod/2008/>](ftp://ftp.ncdc.noaa.gov/pub/data/g sod/2008/). (for Alaska and Hawaii). September 2014.
- Ott, S.L. (2000) *Dairy '96 Study*. Stephen L. Ott, Animal and Plant Health Inspection Service, U.S. Department of Agriculture. June 19, 2000.
- Poe, G., N. Bills, B. Bellows, P. Crosscombe, R. Koelsch, M. Kreher, and P. Wright (1999) *Staff Paper Documenting the Status of Dairy Manure Management in New York: Current Practices and Willingness to Participate in Voluntary Programs*. Department of Agricultural, Resource, and Managerial Economics; Cornell University, Ithaca, New York, September.

- Safley, L.M., Jr. (2000) Personal Communication. Deb Bartram, ERG and L.M. Safley, President, Agri-Waste Technology. June and October 2000.
- Safley, L.M., Jr. and P.W. Westerman (1990) "Psychrophilic anaerobic digestion of animal manure: proposed design methodology." *Biological Wastes*, 34:133-148.
- Stettler, D. (2000) Personal Communication. Don Stettler, Environmental Engineer, National Climate Center, Oregon Natural Resource Conservation Service and Lee-Ann Tracy, ERG. June 27, 2000.
- Sweeten, J. (2000) Personal Communication. John Sweeten, Texas A&M University and Indra Mitra, ERG. June 2000.
- UEP (1999) *Voluntary Survey Results—Estimated Percentage Participation/Activity*. Caged Layer Environmental Management Practices, Industry data submissions for EPA profile development, United Egg Producers and National Chicken Council. Received from John Thorne, Capitolink. June 2000.
- USDA (2015a) *Quick Stats: Agricultural Statistics Database*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. Available online at: <<http://quickstats.nass.usda.gov/>>.
- USDA (2015b) *Chicken and Eggs 2014 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2015. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2015c) *Poultry - Production and Value 2014 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2015. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2014a) *1987, 1992, 1997, 2002, 2007, and 2012 Census of Agriculture*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. Available online at: <<http://www.nass.usda.gov/census/>>. May 2014.
- USDA (2014b) *Chicken and Eggs 2013 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2014. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2014c) *Poultry - Production and Value 2013 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2014. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2013a) *Chicken and Eggs 2012 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2013. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2013b) *Poultry - Production and Value 2012 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2013. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2012a) *Chicken and Eggs 2011 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2012. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2012b) *Poultry - Production and Value 2011 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2012. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2011a) *Chicken and Eggs 2010 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2011. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2011b) *Poultry - Production and Value 2010 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2011. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.

- USDA (2010a) *Chicken and Eggs 2009 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2010. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2010b) *Poultry - Production and Value 2009 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2010. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2009a) *Chicken and Eggs 2008 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. February 2009. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2009b) *Poultry - Production and Value 2008 Summary*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2009. Available online at: <<http://www.nass.usda.gov/Publications/index.asp>>.
- USDA (2009c) *Chicken and Eggs – Final Estimates 2003-2007*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. March 2009. Available online at: <<http://usda.mannlib.cornell.edu/usda/nass/SB980/sb1024.pdf>>.
- USDA (2009d) *Poultry Production and Value—Final Estimates 2003-2007*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. May 2009. Available online at: <<http://usda.mannlib.cornell.edu/usda/nass/SB994/sb1028.pdf>>.
- USDA (2008) *Agricultural Waste Management Field Handbook, National Engineering Handbook (NEH)*, Part 651. Natural Resources Conservation Service, U.S. Department of Agriculture.
- USDA (2004a) *Chicken and Eggs—Final Estimates 1998-2003*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2004. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA (2004b) *Poultry Production and Value—Final Estimates 1998-2002*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. April 2004. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA (1999) *Poultry Production and Value—Final Estimates 1994-97*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. March 1999. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA (1998) *Chicken and Eggs—Final Estimates 1994-97*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. December 1998. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA (1996) *Agricultural Waste Management Field Handbook, National Engineering Handbook (NEH)*, Part 651. Natural Resources Conservation Service, U.S. Department of Agriculture. July 1996.
- USDA (1995a) *Poultry Production and Value—Final Estimates 1988-1993*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. March 1995. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA (1995b) *Chicken and Eggs—Final Estimates 1988-1993*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. December 1995. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA (1994) *Sheep and Goats—Final Estimates 1989-1993*. National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, D.C. January 31, 1994. Available online at: <<http://usda.mannlib.cornell.edu/reports/general/sb/>>.
- USDA, APHIS (2003) *Sheep 2001, Part I: Reference of Sheep Management in the United States, 2001 and Part IV: Baseline Reference of 2001 Sheep Feedlot Health and Management*. USDA-APHIS-VS. Fort Collins, CO. #N356.0702. <[http://www.aphis.usda.gov/animal\\_health/nahms/sheep/index.shtml#sheep2001](http://www.aphis.usda.gov/animal_health/nahms/sheep/index.shtml#sheep2001)>.
- USDA, APHIS (2000) *Layers '99—Part II: References of 1999 Table Egg Layer Management in the U.S.* USDA-APHIS-VS. Fort Collins, CO. <[http://www.aphis.usda.gov/animal\\_health/nahms/poultry/downloads/layers99/Layers99\\_dr\\_PartII.pdf](http://www.aphis.usda.gov/animal_health/nahms/poultry/downloads/layers99/Layers99_dr_PartII.pdf)>.

USDA, APHIS (1996) *Swine '95: Grower/Finisher Part II: Reference of 1995 U.S. Grower/Finisher Health & Management Practices*. USDA-APHIS-VS. Fort Collins, CO.  
<[http://www.aphis.usda.gov/animal\\_health/nahms/swine/downloads/swine95/Swine95\\_dr\\_PartII.pdf](http://www.aphis.usda.gov/animal_health/nahms/swine/downloads/swine95/Swine95_dr_PartII.pdf)>.

Wright, P. (2000) Personal Communication. Lee-Ann Tracy, ERG and Peter Wright, Cornell University, College of Agriculture and Life Sciences. June 23, 2000.

### 3.12. Methodology for Estimating N<sub>2</sub>O Emissions, CH<sub>4</sub> Emissions and Soil Organic C Stock Changes from Agricultural Lands (Cropland and Grassland)

Nitrous oxide (N<sub>2</sub>O) is produced in soils through the microbial processes of nitrification and denitrification<sup>82</sup>. Management influences these processes by modifying the availability of mineral nitrogen (N), which is a key control on the N<sub>2</sub>O emissions rates (Mosier et al. 1998). Emissions can occur directly in the soil where the N is made available or can be transported to another location following volatilization, leaching, or runoff, and then converted into N<sub>2</sub>O. Management practices influence soil organic C stocks in agricultural soils by modifying the natural processes of photosynthesis (i.e., crop and forage production) and microbial decomposition. CH<sub>4</sub> emissions from rice cultivation occur under flooded conditions through the process of methanogenesis. This sub-annex describes the methodologies used to calculate N<sub>2</sub>O emissions from agricultural soil management and annual carbon (C) stock changes from mineral and organic soils classified as *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland*, and *Land Converted to Grassland*<sup>83</sup>, and CH<sub>4</sub> emissions from *Rice Cultivation*. This annex provides the underlying methodologies for these three emission sources because there is considerable overlap in the methods with the majority of emissions estimated using the DAYCENT biogeochemical<sup>84</sup> simulation model.

A combination of Tier 1, 2 and 3 approaches are used to estimate direct and indirect N<sub>2</sub>O emissions and C stock changes in agricultural soils.

More specifically, the methodologies used to estimate soil N<sub>2</sub>O emissions include:

- 1) A Tier 3 method using the DAYCENT biogeochemical simulation model to estimate direct emissions from mineral soils that have less than 35 percent coarse fragments by volume and are used to produce alfalfa hay, barley, corn, cotton, dry beans, grass hay, grass-clover hay, oats, onions, peanuts, potatoes, rice, sorghum, soybeans, sugar beets, sunflowers, tomatoes, and wheat, as well as non-federal grasslands and land use change between grassland and cropland (with the crops listed above and less than 35 percent coarse fragments);
- 2) A combination of the Tier 3 and 1 methods to estimate indirect N<sub>2</sub>O emissions associated with management of cropland and grassland simulated with DAYCENT in Item 1;
- 3) A Tier 1 method to estimate direct and indirect N<sub>2</sub>O emissions from mineral soils that are not simulated with DAYCENT, including very gravelly, cobbly, or shaley soils (greater than 35 percent coarse fragments by volume); mineral soils with less than 35 percent coarse fragments that are used to produce crops that are not simulated by DAYCENT; and crops that are rotated with the crops that are not simulated with DAYCENT/Pasture/Range/Paddock (PRP) manure N deposited on federal grasslands; and
- 4) A Tier 1 method to estimate direct N<sub>2</sub>O emissions due to partial or complete drainage of organic soils in croplands and grasslands.

The methodologies used to estimate soil CH<sub>4</sub> emissions from rice cultivation include:

- 1) A Tier 3 method using the DAYCENT biogeochemical simulation model to estimate CH<sub>4</sub> emissions from mineral soils that have less than 35 percent coarse fragments by volume and rice grown continuously or in rotation with a crop listed in (1) for soil N<sub>2</sub>O emissions; and
- 2) A Tier 1 method to estimate CH<sub>4</sub> emissions from all other soils used to produce rice that are not estimated with the Tier 3 method, including rice grown on organic soils (i.e., *Histosols*), mineral soils with very gravelly, cobbly, or shaley soils (greater than 35 percent coarse fragments by volume), and rice grown in rotation with crops that are not simulated by DAYCENT.

---

<sup>82</sup> Nitrification and denitrification are driven by the activity of microorganisms in soils. Nitrification is the aerobic microbial oxidation of ammonium (NH<sub>4</sub><sup>+</sup>) to nitrate (NO<sub>3</sub><sup>-</sup>), and denitrification is the anaerobic microbial reduction of nitrate to N<sub>2</sub>. Nitrous oxide is a gaseous intermediate product in the reaction sequence of denitrification, which leaks from microbial cells into the soil and then into the atmosphere. Nitrous oxide is also produced during nitrification, although by a less well-understood mechanism (Nevison 2000).

<sup>83</sup> Soil C stock change methods for forestland are described in the *Forestland Remaining Forestland* section.

<sup>84</sup> Biogeochemical cycles are the flow of chemical elements and compounds between living organisms and the physical environment.

The methodologies used to estimate soil organic C stock changes include:

- 1) A Tier 3 method using the DAYCENT biogeochemical simulation model to estimate soil organic C stock changes in mineral soils as described in Item 1 for N<sub>2</sub>O emissions;
- 2) Tier 2 methods with country-specific stock change factors for estimating mineral soil organic C stock changes for mineral soils that are very gravelly, cobbly, or shaley (greater than 35 percent coarse fragments by volume), are used to produce crops or have land use changes to cropland and grassland (other than the conversions between cropland and grassland that are included in Item 1) that are not simulated with DAYCENT;
- 3) Tier 2 methods with country-specific stock change factors for estimating mineral soil organic C stock changes on federal lands;
- 4) Tier 2 methods with country-specific emission factors for estimating losses of C from organic soils that are partly or completely drained for agricultural production; and
- 5) Tier 2 methods for estimating additional changes in mineral soil C stocks due to sewage sludge additions to soils and enrollment changes in the Conservation Reserve Program (CRP) after 2010.

As described above, the Inventory uses a Tier 3 approach to estimate direct soil N<sub>2</sub>O emissions, CH<sub>4</sub> emissions from rice cultivation, and C stock changes for the majority of agricultural lands. This approach has the following advantages over the IPCC Tier 1 or 2 approaches:

- 1) It utilizes actual weather data at sub-county scales enabling quantification of inter-annual variability in N<sub>2</sub>O emissions and C stock changes at finer spatial scales, as opposed to a single emission factor for the entire country for soil N<sub>2</sub>O or broad climate region classification for soil C stock changes;
- 2) The model uses a more detailed characterization of spatially-mapped soil properties that influence soil C and N dynamics, as opposed to the broad soil taxonomic classifications of the IPCC methodology;
- 3) The simulation approach provides a more detailed representation of management influences and their interactions than are represented by a discrete factor-based approach in the Tier 1 and 2 methods; and
- 4) Soil N<sub>2</sub>O and CH<sub>4</sub> emissions, and C stock changes are estimated on a more continuous, daily basis as a function of the interaction of climate, soil, and land management, compared with the linear rate changes that are estimated with the Tier 1 and 2 methods.

The DAYCENT process-based simulation model (daily time-step version of the Century model) has been selected for the Tier 3 approach based on the following criteria:

- 1) The model has been developed in the U.S. and extensively tested and verified for U.S. conditions (e.g., Parton et al. 1987, 1993). In addition, the model has been widely used by researchers and agencies in many other parts of the world for simulating soil C dynamics at local, regional and national scales (e.g., Brazil, Canada, India, Jordan, Kenya, Mexico), soil N<sub>2</sub>O emissions (e.g., Canada, China, Ireland, New Zealand) (Abdalla et al. 2010; Li et al. 2005; Smith et al. 2008; Stehfest and Muller 2004; Cheng et al. 2014), and CH<sub>4</sub> emissions (Cheng et al. 2013).
- 2) The model is capable of simulating cropland, grassland, forest, and savanna ecosystems, and land-use transitions between these different land uses. It is, thus, well suited to model land-use change effects.
- 3) The model is designed to simulate management practices that influence soil C dynamics, CH<sub>4</sub> emissions and direct N<sub>2</sub>O emissions, with the exception of cultivated organic soils; cobbly, gravelly, or shaley soils; and crops that have not been parameterized for DAYCENT simulations (e.g., some vegetables, tobacco, perennial/horticultural crops, and crops that are rotated with these crops). For these latter cases, an IPCC Tier 2 method has been used for soil C stock changes and IPCC Tier 1 method for CH<sub>4</sub> and N<sub>2</sub>O emissions. The model can also be used estimate the amount of N leaching and runoff, as well as volatilization of N, which is subject to indirect N<sub>2</sub>O emissions.
- 4) Much of the data needed for the model is available from existing national databases. The exceptions are CRP enrollment after 2010, management of federal grasslands, and sewage sludge amendments to soils, which are not known at a sufficient resolution to use the Tier 3 model. Soil N<sub>2</sub>O emissions and C stock changes associated with these practices are addressed with a Tier 1 and 2 method, respectively.

Overall, the Tier 3 approach is used to estimate approximately 89 percent of direct soil N<sub>2</sub>O emissions 94 percent of the rice cultivation, and 88 percent of the land area associated with estimation of soil organic C stock changes under agricultural management in the United States.

### Tier 3 Method Description and Model Evaluation

The DAYCENT biogeochemical model (Parton et al. 1998; Del Grosso et al. 2001, 2011) simulates biogeochemical C and N fluxes between the atmosphere, vegetation, and soil; and provides a more complete estimation of soil C stock changes and N<sub>2</sub>O emissions than IPCC Tier 1 or 2 methods by more thoroughly accounting for the influence of environmental conditions. These conditions include soil characteristics, weather patterns, crop and forage characteristics, and management practices. The DAYCENT model utilizes the soil C modeling framework developed in the Century model (Parton et al. 1987, 1988, 1994; Metherell et al. 1993), but has been refined to simulate dynamics at a daily time-step. Carbon and N dynamics are linked in plant-soil systems through biogeochemical processes of microbial decomposition and plant production (McGill and Cole 1981). Coupling the three source categories (i.e., agricultural soil C, rice CH<sub>4</sub> and soil N<sub>2</sub>O) in a single inventory analysis ensures that there is a consistent treatment of the processes and interactions between C and N cycling in soils. For example, plant growth is controlled by nutrient availability, water, and temperature stress. Plant growth, along with residue management, determines C inputs to soils, which influence C stock changes, and removal of mineral N from the soil where plant growth influences the amount of N that can be converted into N<sub>2</sub>O. Nutrient supply is a function of external nutrient additions as well as litter and soil organic matter (SOM) decomposition rates, and increasing decomposition can lead to a reduction in soil organic C stocks due to microbial respiration, and greater N<sub>2</sub>O emissions by enhancing mineral N availability in soils.

Key processes simulated by DAYCENT include (1) plant growth; (2) organic matter formation and decomposition; (3) soil water and temperature regimes by layer; (4) nitrification and denitrification processes; and (5) methanogenesis (Figure A-7). Each of these submodels will be described separately below.


- 1) The plant-growth submodel simulates C assimilation through photosynthesis; N uptake; dry matter production; partitioning of C within the crop or forage; senescence; and mortality. The primary function of the growth submodel is to estimate the amount, type, and timing of organic matter inputs to soil, and to represent the influence of the plant on soil water, temperature, and N balance. Yield and removal of harvested biomass are also simulated. Separate submodels are designed to simulate herbaceous plants (i.e., agricultural crops and grasses) and woody vegetation (i.e., trees and scrub). Maximum daily net primary production (NPP) is estimated using the NASA-CASA production algorithm (Potter et al. 1993, 2007) and MODIS Enhanced Vegetation Index (EVI) products, MOD13Q1 and MYD13Q1, or an approximation of EVI data derived from the MODIS products (Gurung et al. 2009). The NASA-CASA production algorithm is only used for the following major crops: corn, soybeans, sorghum, cotton and wheat.<sup>85</sup> Other regions and crops are simulated with a single value for the maximum daily NPP, instead of the more dynamic NASA-CASA algorithm. The maximum daily NPP rate is modified by air temperature and available water (to capture temperature and moisture stress). If the NASA-CASA algorithm is not used in the simulation, then production is further subject to nutrient limitations (i.e., nitrogen). Model evaluation has shown that the NASA-CASA algorithm improves the precision of NPP estimates using the EVI products to inform the production model. The  $r^2$  is 83 percent for the NASA-CASA algorithm and 64 percent for the single parameter value approach. See Figure A-8.
- 2) Dynamics of soil organic C and N (Figure A-7) are simulated for the surface and belowground litter pools and soil organic matter in the top 20 cm of the soil profile; mineral N dynamics are simulated through the whole soil profile. Organic C and N stocks are represented by two plant litter pools (metabolic and structural) and three soil organic matter (SOM) pools (active, slow, and passive). The metabolic litter pool represents the easily decomposable constituents of plant residues, while the structural litter pool is composed of more recalcitrant, ligno-cellulose plant materials. The three SOM pools represent a gradient in decomposability, from active SOM (representing microbial biomass and associated metabolites) having a rapid turnover (months to years), to passive SOM (representing highly processed, humified, condensed decomposition products), which is highly recalcitrant, with mean residence times on the order of several hundred years. The slow pool represents decomposition products of intermediate stability, having a mean residence time on the order of decades and is the fraction that tends to change the most in response to changes in land use and management. Soil texture influences turnover rates of the slow and passive pools. The clay and silt-sized mineral fraction of the soil provides physical protection from microbial decomposition, leading to enhanced SOM stabilization in finely textured soils. Soil temperature and moisture, tillage disturbance, aeration, and other factors influence decomposition and loss of C from the soil organic matter pools.
- 3) The soil-water balance submodel calculates water balance components and changes in soil water availability, which influences both plant growth and decomposition/nutrient cycling processes. The moisture content of soils are simulated

---


<sup>85</sup> It is a planned improvement to estimate NPP for additional crops and grass forage with the NASA-CASA method in the future.

through a multi-layer profile based on precipitation, snow accumulation and melting, interception, soil and canopy evaporation, transpiration, soil water movement, runoff, and drainage.

**Figure A-7: DAYCENT Model Flow Diagram**



**Figure A-8: Modeled versus measured net primary production ( $\text{g C m}^{-2}$ )**



Part a) presents results of the NASA-CASA algorithm ( $r^2 = 83\%$ ) and part b) presents the results of a single parameter value for maximum net primary production ( $r^2 = 64\%$ ).

- 4) Soil mineral N dynamics are modeled based on N inputs from fertilizer inputs (synthetic and organic), residue N inputs, soil organic matter mineralization in addition to symbiotic and asymbiotic N fixation. Mineral N is available for plant and microbial uptake, and is largely controlled by the specified stoichiometric limits for these organisms (i.e., C:N ratios). Mineral and organic N losses are simulated with leaching and runoff, and nitrogen can be volatilized and lost from the soil through ammonia volatilization, nitrification and denitrification.  $\text{N}_2\text{O}$  emissions occur through nitrification and denitrification. Denitrification is a function of soil  $\text{NO}_3^-$  concentration, water filled pore space (WFPS), heterotrophic (i.e., microbial) respiration, and texture. Nitrification is controlled by soil ammonium ( $\text{NH}_4^+$ ) concentration, water filled pore space, temperature, and pH (See Box 2 for more information).
- 5) Methanogenesis is modeled under anaerobic conditions and is controlled by carbon substrate availability, temperature, and redox potential (Cheng et al. 2013). Carbon substrate supply is determined by decomposition of residues and soil organic matter, in addition to root exudation. The transport of  $\text{CH}_4$  to the atmosphere occurs through the rice plant and via ebullition (i.e., bubbles).  $\text{CH}_4$  can be oxidized (methanotrophy) as it moves through a flooded soil and the oxidation rates are higher as the plants mature and in soils with more clay (Sass et al. 1994).

The model allows for a variety of management options to be simulated, including specifying different crop types, crop sequences (e.g., rotation), tillage practices, fertilization, organic matter addition (e.g., manure amendments), harvest events (with variable residue removal), drainage, flooding, irrigation, burning, and grazing intensity. An input “schedule” file is used to simulate the timing of management activities and temporal trends; schedules can be organized into discrete time blocks to define a repeated sequence of events (e.g., a crop rotation or a frequency of disturbance such as a burning cycle for perennial grassland). Management options can be specified for any day of a year within a scheduling block, where management codes point to operation-specific parameter files (referred to as \*.100 files), which contain the information used

to simulate management effects with the model algorithms. User-specified management activities can be defined by adding to or editing the contents of the \*.100 files. Additional details of the model formulation are given in Parton et al. (1987, 1988, 1994, 1998), Del Grosso et al. (2001, 2011), Cheng et al. (2013) and Metherell et al. (1993), and archived copies of the model source code are available.

[BEGIN TEXT BOX]

## Box 2. DAYCENT Model Simulation of Nitrification and Denitrification

The DAYCENT model simulates the two biogeochemical processes, nitrification and denitrification, that result in N<sub>2</sub>O emissions from soils (Del Grosso et al. 2000, Parton et al. 2001). Nitrification is calculated for the top 15 cm of soil (where nitrification mostly occurs) while denitrification is calculated for the entire soil profile (accounting for denitrification near the surface and subsurface as nitrate leaches through the profile). The equations and key parameters controlling N<sub>2</sub>O emissions from nitrification and denitrification are described below.

Nitrification is controlled by soil ammonium (NH<sub>4</sub><sup>+</sup>) concentration, temperature (t), Water Filled Pore Space (WFPS) and pH according to the following equation:

$$\text{Nit} = \text{NH}_{4+} \times K_{\text{max}} \times F(t) \times F(\text{WFPS}) \times F(\text{pH})$$

where,

|                  |   |  |
|------------------|---|--|
| Nit              | = | the soil nitrification rate (g N/m <sup>2</sup> /day)  |
| NH <sub>4+</sub> | = | the model-derived soil ammonium concentration (g N/m <sup>2</sup> )                          |
| K <sub>max</sub> | = | the maximum fraction of NH <sub>4</sub> <sup>+</sup> nitrified (K <sub>max</sub> = 0.10/day) |
| F(t)             | = | the effect of soil temperature on nitrification (Figure A-9a)                                |
| F(WFPS)          | = | the effect of soil water content and soil texture on nitrification (Figure A-9b)             |
| F(pH)            | = | the effect of soil pH on nitrification (Figure A-9c)   |

The current parameterization used in the model assumes that 1.2 percent of nitrified N is converted to N<sub>2</sub>O.

The model assumes that denitrification rates are controlled by the availability of soil NO<sub>3</sub><sup>-</sup> (electron acceptor), labile C compounds (electron donor) and oxygen (competing electron acceptor). Heterotrophic soil respiration is used as a proxy for labile C availability, while oxygen availability is a function of soil physical properties that influence gas diffusivity, soil WFPS, and oxygen demand. The model selects the minimum of the NO<sub>3</sub><sup>-</sup> and CO<sub>2</sub> functions to establish a maximum potential denitrification rate. These rates vary for particular levels of electron acceptor and C substrate, and account for limitations of oxygen availability to estimate daily denitrification rates according to the following equation:

$$\text{Den} = \min[F(\text{CO}_2), F(\text{NO}_3)] \times F(\text{WFPS})$$

where,

|                     |   |   |
|---------------------|---|---|
| Den                 | = | the soil denitrification rate (μg N/g soil/day)                   |
| F(NO <sub>3</sub> ) | = | a function relating N gas flux to nitrate levels (Figure A-10a)   |
| F(CO <sub>2</sub> ) | = | a function relating N gas flux to soil respiration (Figure A-10b) |

$F(\text{WFPS})$  = a dimensionless multiplier (Figure A-10c).

The x inflection point of  $F(\text{WFPS})$  is a function of respiration and soil gas diffusivity at field capacity ( $D_{\text{FC}}$ ):

$$x \text{ inflection} = 0.90 - M(\text{CO}_2)$$

where,

$M$  = a multiplier that is a function of  $D_{\text{FC}}$ . In technical terms, the inflection point is the domain where either  $F(\text{WFPS})$  is not differentiable or its derivative is 0. In this case, the inflection point can be interpreted as the WFPS value at which denitrification reaches half of its maximum rate.

Respiration has a much stronger effect on the water curve in clay soils with low  $D_{\text{FC}}$  than in loam or sandy soils with high  $D_{\text{FC}}$  (Figure A-9b). The model assumes that microsites in fine-textured soils can become anaerobic at relatively low water contents when oxygen demand is high. After calculating total N gas flux, the ratio of  $\text{N}_2/\text{N}_2\text{O}$  is estimated so that total N gas emissions can be partitioned between  $\text{N}_2\text{O}$  and  $\text{N}_2$ :

$$R_{\text{N}_2/\text{N}_2\text{O}} = F_r(\text{NO}_3/\text{CO}_2) \times F_r(\text{WFPS}).$$

where,

$R_{\text{N}_2/\text{N}_2\text{O}}$  = the ratio of  $\text{N}_2/\text{N}_2\text{O}$

$F_r(\text{NO}_3/\text{CO}_2)$  = a function estimating the impact of the availability of electron donor relative to substrate

$F_r(\text{WFPS})$  = a multiplier to account for the effect of soil water on  $\text{N}_2:\text{N}_2\text{O}$ .

For  $F_r(\text{NO}_3/\text{CO}_2)$ , as the ratio of electron donor to substrate increases, a higher portion of N gas is assumed to be in the form of  $\text{N}_2\text{O}$ . For  $F_r(\text{WFPS})$ , as WFPS increases, a higher portion of N gas is assumed to be in the form of  $\text{N}_2$ .


[End Box]

Comparison of model results and plot level data show that DAYCENT reliably simulates soil organic matter levels (Ogle et al. 2007). The model was tested and shown to capture the general trends in C storage across 908 treatment observations from over 80 experimental sites (Figure A-11). Some bias and imprecision occur in predictions of soil organic C, which is reflected in the uncertainty associated with DAYCENT model results. Regardless, the Tier 3 approach has considerably less uncertainty than Tier 1 and 2 methods (Del Grosso et al. 2010; Figure A-12).


Similarly, DAYCENT model results have been compared to trace gas  $\text{N}_2\text{O}$  fluxes for a number of native and managed systems (Del Grosso et al. 2001, 2005, 2010) (Figure A-13). In general, the model simulates accurate emissions, but some bias and imprecision does occur in predictions, which is reflected in the uncertainty associated with DAYCENT model results. Comparisons with measured data showed that DAYCENT estimated  $\text{N}_2\text{O}$  emissions more accurately and precisely than the IPCC Tier 1 methodology (IPCC 2006) (See Agricultural Soil Management, QA/QC and Verification Section). The linear regression of simulated vs. measured emissions for DAYCENT had higher  $r^2$  values and a fitted line closer to a perfect 1:1 relationship between measured and modeled  $\text{N}_2\text{O}$  emissions compared to the IPCC Tier 1 approach (Del Grosso et al. 2005, 2008). This is not surprising, since DAYCENT includes site-specific factors (climate, soil properties, and previous management) that influence  $\text{N}_2\text{O}$  emissions. Furthermore, DAYCENT also simulated  $\text{NO}_3^-$  leaching (root mean square error = 20 percent) more accurately than IPCC Tier 1 methodology (root mean square error = 69 percent) (Del Grosso et al. 2005). Volatilization of N gases that contribute to indirect soil  $\text{N}_2\text{O}$  emissions is the only component that has not been thoroughly tested, which is due to a lack of measurement data. Overall, the Tier 3 approach has reduced uncertainties in the agricultural soil C stock changes and  $\text{N}_2\text{O}$  emissions compared to using lower Tier methods.

DAYCENT predictions of soil  $\text{CH}_4$  emissions have also been compared to experimental measurements from sites in California, Texas, Arkansas and Louisiana (Figure A-14). There are 10 experiments and 126 treatment observations. In general, the model estimates  $\text{CH}_4$  emissions in most states with no apparent bias, but there is a lack of precision, which is addressed in the uncertainty analysis. The exception is California where the model tends to over-estimate low emission rates, and this additional uncertainty is captured in the error propagation associated with the inventory analysis for California.


**Figure A-9: Effect of Soil Temperature (a) , Water-Filled Pore Space (b) , and pH (c) on Nitrification Rates**




**Figure A-10: Effect of Soil Nitrite Concentration (a), Heterotrophic Respiration Rates (b), and Water-Filled Pore Space (c) on Denitrification Rates**



**Figure A-11: Comparisons of Results from DAYCENT Model and Measurements of Soil Organic C Stocks**




**Figure A-12: Comparison of Estimated Soil Organic C Stock Changes and Uncertainties using Tier 1 (IPCC 2006), Tier 2 (Ogle et al. 2003, 2006) and Tier 3 Methods**




Source: Tier 1 (IPCC 2007), Tier 2 (Ogle et al. 2003, 2006), Tier 3 (Ogle et al. 2010).

**Figure A-13: Comparisons of Results from DAYCENT Model and Measurements of Soil Nitrous Oxide Emissions**



**Figure A-14: Comparisons of Results from DAYCENT Model and Measurements of Soil Methane Emissions**



## Inventory Compilation Steps

There are five steps involved in estimating soil organic C stock changes for *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland* and *Land Converted to Grassland*; direct N<sub>2</sub>O emissions from cropland and grassland soils; indirect N<sub>2</sub>O emissions from volatilization, leaching, and runoff from croplands and grasslands; and CH<sub>4</sub> emissions from rice cultivation. First, the activity data are derived from a combination of land-use, livestock, crop, and grassland management surveys, as well as expert knowledge. In the second, third, and fourth steps, soil organic C stock changes, direct and indirect N<sub>2</sub>O emissions are estimated using DAYCENT and/or the Tier 1 and 2 methods. In the fifth step, total emissions are computed by summing all components separately for soil organic C stock changes and N<sub>2</sub>O emissions. The remainder of this annex describes the methods underlying each step.

### Step 1: Derive Activity Data

The following describes how the activity data are derived to estimate soil organic C stock changes and direct and indirect N<sub>2</sub>O emissions. The activity data requirements include: (1) land base and history data, (2) crop-specific mineral N fertilizer rates,<sup>86</sup> (3) crop-specific manure amendment N rates and timing, (4) other N inputs, (5) tillage practices, (6) irrigation data, (7) Enhanced Vegetation Index (EVI), (8) daily weather data, and (9) edaphic characteristics.<sup>87</sup>

#### Step 1a: Activity Data for the Agricultural Land Base and Histories

The U.S. Department of Agriculture's 2010 National Resources Inventory (NRI) (USDA-NRCS 2013) provides the basis for identifying the U.S. agricultural land base on non-federal lands, and classifying parcels into *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland*, and *Land Converted to Grassland*. In 1998, the NRI program began collecting annual data, and data are currently available through 2012 (USDA-NRCS, 2015) although this Inventory only uses NRI data through 2010 because newer data were not made available in time to incorporate the additional years into this Inventory. Note that the Inventory does not include estimates of N<sub>2</sub>O emissions for federal grasslands (with the exception of soil N<sub>2</sub>O from PRP manure N, i.e., manure deposited directly onto pasture, range or paddock by grazing livestock) and a minor amount of croplands on federal lands.

The NRI has a stratified multi-stage sampling design, where primary sample units are stratified on the basis of county and township boundaries defined by the U.S. Public Land Survey (Nusser and Goebel 1997). Within a primary sample unit, typically a 160-acre (64.75 ha) square quarter-section, three sample points are selected according to a restricted randomization procedure. Each point in the survey is assigned an area weight (expansion factor) based on other known areas and land-use information (Nusser and Goebel 1997). In principle, the expansion factors represent the amount of area with the land use and land use change history that is the same as the point location. It is important to note that the NRI uses a sampling approach, and therefore there is some uncertainty associated with scaling the point data to a region or the country using the expansion factors. In general, those uncertainties decline at larger scales, such as states compared to smaller county units, because of a larger sample size. An extensive amount of soils, land-use, and land management data have been collected through the survey (Nusser et al. 1998).<sup>88</sup> Primary sources for data include aerial photography and remote sensing imagery as well as field visits and county office records.

The annual NRI data product provides crop data for most years between 1979 and 2010, with the exception of 1983, 1988, and 1993. These years are gap-filled using an automated set of rules so that cropping sequences are filled with the most likely crop type given the historical cropping pattern at each NRI point location. Grassland data are reported on 5-year increments prior to 1998, but it is assumed that the land use is also grassland between the years of data collection (see Easter et al. 2008 for more information).

NRI points are included in the land base for the agricultural soil C and N<sub>2</sub>O emissions inventories if they are identified as cropland or grassland<sup>89</sup> between 1990 and 2010 (Table A-217).<sup>90</sup> NRI does not provide land use data on federal

<sup>86</sup> No data are currently available at the national scale to distinguish the type of fertilizer applied or timing of applications rates. It is a planned improvement to address variation in these practices in future inventories.

<sup>87</sup> Edaphic characteristics include such factors as water content, acidity, aeration, and the availability of nutrients.

<sup>88</sup> In the current Inventory, NRI data only provide land-use and management statistics through 2010. More recent data will be incorporated in the future to extend the time series of land use and management data.

<sup>89</sup> Includes only non-federal lands because federal lands are not classified into land uses as part of the NRI survey (i.e., they are only designated as federal lands).

<sup>90</sup> Land use for 2011 to 2014 is assumed to be the same as 2010, but will be updated with newer NRI (i.e. USDA-NRCS 2015).

lands, therefore land use on federal lands are derived from the National Land Cover Database (NLCD) (Fry et al. 2011; Homer et al. 2007; Homer et al. 2015). Federal NRI points classified as cropland or grassland according to the NLCD are included in the agricultural land base. The NRI data are reconciled with the Forest Inventory and Analysis Dataset, and in this process, the time series for *Grassland Remaining Grassland* and *Land Converted to Grassland* is modified to account for differences in forest land area between the two national surveys (See Section 6.1 for more information on the U.S. land representation). Overall, 674,613 NRI survey points are included in the inventory (USDA-NRCS 2013).

For each year, land parcels are subdivided into *Cropland Remaining Cropland*, *Land Converted to Cropland*, *Grassland Remaining Grassland*, and *Land Converted to Grassland*. Land parcels under cropping management in a specific year are classified as *Cropland Remaining Cropland* if the parcel has been used as cropland for at least 20 years<sup>91</sup>. Similarly land parcels under grassland management in a specific year of the inventory are classified as *Grassland Remaining Grassland* if they have been designated as grassland for at least 20 years. Otherwise, land parcels are classified as *Land Converted to Cropland* or *Land Converted to Grassland* based on the most recent use in the inventory time period. Lands are retained in the land-use change categories (i.e., *Land Converted to Cropland* and *Land Converted to Grassland*) for 20 years as recommended by the 2006 IPCC Guidelines. Lands converted into Cropland and Grassland are further subdivided into the specific land use conversions (e.g., *Forest Land Converted to Cropland*).

**Table A-217: Total Land Areas for the Agricultural Soil C and N<sub>2</sub>O Inventory, Subdivided by Land Use Categories (Million Hectares)**

| Category                            | Land Areas (million ha) |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------------|-------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
|                                     | 1990                    | 1991          | 1992          | 1993          | 1994          | 1995          | 1996          | 1997          | 1998          | 1999          | 2000          | 2001          |
| <b>Mineral Soils</b>                | <b>447.61</b>           | <b>446.84</b> | <b>446.07</b> | <b>445.35</b> | <b>444.64</b> | <b>443.71</b> | <b>442.77</b> | <b>441.91</b> | <b>441.02</b> | <b>439.68</b> | <b>439.60</b> | <b>438.95</b> |
| <b>Croplands</b>                    | <b>173.95</b>           | <b>173.85</b> | <b>173.64</b> | <b>173.11</b> | <b>172.79</b> | <b>172.32</b> | <b>172.05</b> | <b>171.70</b> | <b>169.18</b> | <b>168.33</b> | <b>168.39</b> | <b>168.11</b> |
| Cropland Remaining Cropland         | 161.24                  | 160.80        | 160.28        | 158.53        | 156.83        | 156.20        | 155.60        | 155.03        | 151.39        | 150.27        | 150.76        | 150.86        |
| Grassland Converted to Cropland     | 12.07                   | 12.40         | 12.71         | 13.88         | 15.20         | 15.36         | 15.70         | 15.92         | 16.99         | 17.26         | 16.84         | 16.48         |
| Forest Converted to Cropland        | 0.24                    | 0.24          | 0.23          | 0.22          | 0.21          | 0.21          | 0.20          | 0.20          | 0.19          | 0.17          | 0.15          | 0.14          |
| Other Lands Converted to Cropland   | 0.19                    | 0.20          | 0.21          | 0.23          | 0.26          | 0.26          | 0.27          | 0.27          | 0.28          | 0.30          | 0.32          | 0.31          |
| Settlements Converted to Croplands  | 0.08                    | 0.08          | 0.08          | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          | 0.12          | 0.13          | 0.13          | 0.13          |
| Wetlands Converted to Croplands     | 0.13                    | 0.13          | 0.13          | 0.16          | 0.19          | 0.19          | 0.19          | 0.20          | 0.20          | 0.20          | 0.20          | 0.20          |
| <b>Grasslands</b>                   | <b>273.65</b>           | <b>272.99</b> | <b>272.43</b> | <b>272.24</b> | <b>271.86</b> | <b>271.39</b> | <b>270.71</b> | <b>270.20</b> | <b>271.84</b> | <b>271.35</b> | <b>271.21</b> | <b>270.83</b> |
| Grasslands Remaining Grasslands     | 263.76                  | 262.86        | 262.03        | 260.35        | 258.59        | 257.79        | 256.97        | 256.24        | 253.84        | 252.77        | 252.34        | 251.64        |
| Croplands Converted to Grasslands   | 8.76                    | 8.87          | 9.06          | 10.42         | 11.67         | 11.94         | 12.08         | 12.26         | 15.07         | 15.60         | 15.89         | 16.16         |
| Forest Converted to Grasslands      | 0.33                    | 0.33          | 0.33          | 0.33          | 0.33          | 0.36          | 0.35          | 0.34          | 1.34          | 1.34          | 1.32          | 1.31          |
| Other Lands Converted to Grasslands | 0.43                    | 0.45          | 0.50          | 0.59          | 0.70          | 0.74          | 0.74          | 0.78          | 0.93          | 0.95          | 0.99          | 1.05          |
| Settlements Converted to Grasslands | 0.06                    | 0.07          | 0.07          | 0.08          | 0.09          | 0.09          | 0.10          | 0.10          | 0.12          | 0.13          | 0.13          | 0.14          |
| Wetlands Converted to Grasslands    | 0.32                    | 0.41          | 0.45          | 0.46          | 0.48          | 0.47          | 0.48          | 0.48          | 0.54          | 0.55          | 0.55          | 0.55          |
| <b>Organic Soils</b>                | <b>1.35</b>             | <b>1.33</b>   | <b>1.32</b>   | <b>1.32</b>   | <b>1.33</b>   | <b>1.32</b>   | <b>1.31</b>   | <b>1.29</b>   | <b>1.30</b>   | <b>1.28</b>   | <b>1.28</b>   | <b>1.27</b>   |
| <b>Croplands</b>                    | <b>0.75</b>             | <b>0.75</b>   | <b>0.74</b>   | <b>0.73</b>   | <b>0.74</b>   | <b>0.74</b>   | <b>0.74</b>   | <b>0.73</b>   | <b>0.73</b>   | <b>0.74</b>   | <b>0.74</b>   | <b>0.75</b>   |
| Cropland Remaining Cropland         | 0.66                    | 0.66          | 0.65          | 0.64          | 0.63          | 0.63          | 0.63          | 0.63          | 0.61          | 0.61          | 0.62          | 0.61          |
| Grassland Converted to Cropland     | 0.07                    | 0.07          | 0.07          | 0.07          | 0.08          | 0.09          | 0.09          | 0.08          | 0.10          | 0.10          | 0.10          | 0.12          |
| Forest Converted to Cropland        | 0.00                    | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Other Lands Converted to Cropland   | 0.00                    | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Settlements Converted to Croplands  | 0.00                    | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Wetlands Converted to Croplands     | 0.01                    | 0.01          | 0.01          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          |
| <b>Grasslands</b>                   | <b>0.59</b>             | <b>0.58</b>   | <b>0.58</b>   | <b>0.59</b>   | <b>0.59</b>   | <b>0.58</b>   | <b>0.57</b>   | <b>0.56</b>   | <b>0.56</b>   | <b>0.55</b>   | <b>0.55</b>   | <b>0.52</b>   |
| Grasslands Remaining Grasslands     | 0.53                    | 0.53          | 0.52          | 0.51          | 0.50          | 0.49          | 0.49          | 0.48          | 0.47          | 0.46          | 0.45          | 0.41          |
| Croplands Converted to Grasslands   | 0.05                    | 0.05          | 0.05          | 0.06          | 0.07          | 0.07          | 0.07          | 0.06          | 0.08          | 0.07          | 0.07          | 0.08          |
| Forest Converted to Grasslands      | 0.00                    | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Other Lands Converted to Grasslands | 0.00                    | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Settlements Converted to Grasslands | 0.00                    | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Wetlands Converted to Grasslands    | 0.01                    | 0.01          | 0.01          | 0.01          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          | 0.02          |
| <b>Total</b>                        | <b>448.95</b>           | <b>448.17</b> | <b>447.39</b> | <b>446.67</b> | <b>445.97</b> | <b>445.03</b> | <b>444.08</b> | <b>443.20</b> | <b>442.31</b> | <b>440.96</b> | <b>440.89</b> | <b>440.22</b> |

| Category                    | 2002          | 2003          | 2004          | 2005          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|-----------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Mineral Soils</b>        | <b>438.34</b> | <b>437.49</b> | <b>436.72</b> | <b>435.66</b> | <b>434.83</b> | <b>434.06</b> | <b>433.27</b> | <b>432.59</b> | <b>431.88</b> | <b>431.60</b> | <b>431.31</b> | <b>431.31</b> | <b>431.31</b> |
| <b>Croplands</b>            | <b>167.84</b> | <b>167.22</b> | <b>165.74</b> | <b>165.28</b> | <b>164.79</b> | <b>164.47</b> | <b>164.03</b> | <b>163.58</b> | <b>162.98</b> | <b>162.98</b> | <b>162.98</b> | <b>162.98</b> | <b>162.98</b> |
| Cropland Remaining Cropland | 150.89        | 151.50        | 151.29        | 151.10        | 150.82        | 151.23        | 151.48        | 151.79        | 151.38        | 151.38        | 151.39        | 151.39        | 151.39        |

<sup>91</sup> NRI points are classified according to land-use history records starting in 1982 when the NRI survey began, and consequently the classifications are based on less than 20 years from 1990 to 1998.

|                                     |               |               |               |               |               |               |               |               |               |               |               |               |               |
|-------------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Grassland Converted to Cropland     | 16.21         | 15.02         | 13.81         | 13.59         | 13.39         | 12.71         | 12.05         | 11.33         | 11.16         | 11.16         | 11.16         | 11.16         | 11.16         |
| Forest Converted to Cropland        | 0.13          | 0.12          | 0.11          | 0.09          | 0.09          | 0.06          | 0.05          | 0.05          | 0.05          | 0.05          | 0.05          | 0.05          | 0.05          |
| Other Lands Converted to Cropland   | 0.30          | 0.28          | 0.26          | 0.24          | 0.25          | 0.24          | 0.23          | 0.22          | 0.21          | 0.21          | 0.21          | 0.21          | 0.21          |
| Settlements Converted to Croplands  | 0.13          | 0.11          | 0.10          | 0.10          | 0.10          | 0.10          | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          |
| Wetlands Converted to Croplands     | 0.19          | 0.18          | 0.16          | 0.15          | 0.15          | 0.14          | 0.12          | 0.10          | 0.10          | 0.10          | 0.10          | 0.10          | 0.10          |
| <b>Grasslands</b>                   | <b>270.49</b> | <b>270.27</b> | <b>270.99</b> | <b>270.38</b> | <b>270.04</b> | <b>269.59</b> | <b>269.24</b> | <b>269.00</b> | <b>268.90</b> | <b>268.62</b> | <b>268.33</b> | <b>268.33</b> | <b>268.33</b> |
| Grasslands Remaining Grasslands     | 251.09        | 251.24        | 251.18        | 250.46        | 250.10        | 250.07        | 250.31        | 250.33        | 250.18        | 249.90        | 249.62        | 249.62        | 249.62        |
| Croplands Converted to Grasslands   | 16.36         | 16.06         | 16.85         | 16.55         | 16.58         | 16.22         | 15.79         | 15.61         | 15.79         | 15.79         | 15.79         | 15.79         | 15.79         |
| Forest Converted to Grasslands      | 1.30          | 1.25          | 1.25          | 1.82          | 1.82          | 1.77          | 1.75          | 1.75          | 1.73          | 1.73          | 1.73          | 1.73          | 1.73          |
| Other Lands Converted to Grasslands | 1.06          | 1.04          | 1.03          | 0.93          | 0.92          | 0.91          | 0.89          | 0.87          | 0.82          | 0.82          | 0.82          | 0.82          | 0.82          |
| Settlements Converted to Grasslands | 0.14          | 0.14          | 0.14          | 0.13          | 0.13          | 0.13          | 0.13          | 0.12          | 0.12          | 0.12          | 0.12          | 0.12          | 0.12          |
| Wetlands Converted to Grasslands    | 0.54          | 0.54          | 0.54          | 0.49          | 0.49          | 0.48          | 0.39          | 0.31          | 0.26          | 0.26          | 0.26          | 0.26          | 0.26          |
| <b>Organic Soils</b>                | <b>1.26</b>   | <b>1.25</b>   | <b>1.24</b>   | <b>1.24</b>   | <b>1.24</b>   | <b>1.23</b>   | <b>1.23</b>   | <b>1.22</b>   | <b>1.22</b>   | <b>1.22</b>   | <b>1.21</b>   | <b>1.21</b>   | <b>1.21</b>   |
| <b>Croplands</b>                    | <b>0.75</b>   | <b>0.74</b>   | <b>0.74</b>   | <b>0.74</b>   | <b>0.73</b>   | <b>0.73</b>   | <b>0.72</b>   | <b>0.70</b>   | <b>0.70</b>   | <b>0.70</b>   | <b>0.70</b>   | <b>0.70</b>   | <b>0.70</b>   |
| Cropland Remaining Cropland         | 0.61          | 0.61          | 0.64          | 0.64          | 0.63          | 0.63          | 0.63          | 0.62          | 0.62          | 0.62          | 0.62          | 0.62          | 0.62          |
| Grassland Converted to Cropland     | 0.12          | 0.11          | 0.09          | 0.09          | 0.09          | 0.08          | 0.08          | 0.07          | 0.07          | 0.07          | 0.07          | 0.07          | 0.07          |
| Forest Converted to Cropland        | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Other Lands Converted to Cropland   | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Settlements Converted to Croplands  | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Wetlands Converted to Croplands     | 0.02          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          | 0.01          |
| <b>Grasslands</b>                   | <b>0.51</b>   | <b>0.50</b>   | <b>0.50</b>   | <b>0.50</b>   | <b>0.50</b>   | <b>0.50</b>   | <b>0.50</b>   | <b>0.52</b>   | <b>0.51</b>   | <b>0.51</b>   | <b>0.51</b>   | <b>0.51</b>   | <b>0.51</b>   |
| Grasslands Remaining Grasslands     | 0.41          | 0.40          | 0.39          | 0.39          | 0.39          | 0.39          | 0.39          | 0.39          | 0.39          | 0.38          | 0.38          | 0.38          | 0.38          |
| Croplands Converted to Grasslands   | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          | 0.09          | 0.10          | 0.10          | 0.10          | 0.10          | 0.10          | 0.10          |
| Forest Converted to Grasslands      | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Other Lands Converted to Grasslands | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Settlements Converted to Grasslands | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          | 0.00          |
| Wetlands Converted to Grasslands    | 0.01          | 0.01          | 0.01          | 0.02          | 0.02          | 0.02          | 0.02          | 0.03          | 0.03          | 0.03          | 0.03          | 0.03          | 0.03          |
| <b>Total</b>                        | <b>439.60</b> | <b>438.73</b> | <b>437.96</b> | <b>436.90</b> | <b>436.07</b> | <b>435.29</b> | <b>434.50</b> | <b>433.81</b> | <b>433.10</b> | <b>432.81</b> | <b>432.53</b> | <b>432.53</b> | <b>432.53</b> |

Note: The area estimates are not consistent with the land area values shown in the Representation of the U.S. Land Base chapter because the current inventory does not estimate emissions and removals for all managed lands. Specifically, grassland and cropland in Alaska are not included in the current inventory.

The Tier 3 method using the DAYCENT model is applied to estimate soil C stock changes and N<sub>2</sub>O emissions for most of the NRI points that occur on mineral soils. For the Tier 3 inventory, the actual crop and grassland histories are simulated with the DAYCENT model. Parcels of land that are not simulated with DAYCENT are allocated to the Tier 2 approach for estimating soil organic C stock change, and a Tier 1 method (IPCC 2006) to estimate soil N<sub>2</sub>O emissions (Table A-218) (Note: the Tier 1 method for soil N<sub>2</sub>O does not require land area data with the exception of emissions from drainage and cultivation of organic soils, so in practice it is only the amount of N input to mineral soils that is addressed by the Tier 1 method and not the actual land area).

The land base for the Tier 1 and 2 methods includes (1) land parcels occurring on organic soils; (2) land parcels that include non-agricultural uses such as forest and federal lands in one or more years of the inventory; (3) land parcels on mineral soils that are very gravelly, cobbly, or shaly (i.e., classified as soils that have greater than 35 percent of soil volume comprised of gravel, cobbles, or shale); or (4) land parcels that are used to produce some of the vegetable crops, perennial/horticultural crops, and tobacco, which are either grown continuously or in rotation with other crops. DAYCENT has not been fully tested or developed to simulate biogeochemical processes in soils used to produce some annual (e.g., tobacco), horticultural (e.g., flowers), or perennial (e.g., vineyards, orchards) crops and agricultural use of organic soils. In addition, DAYCENT has not been adequately tested for soils with a high gravel, cobble, or shale content.

**Table A-218: Total Land Area Estimated with Tier 2 and 3 Inventory Approaches (Million Hectares)**

| Year | Land Areas (million ha) |         |        |          |        |
|------|-------------------------|---------|--------|----------|--------|
|      | Tier 1/2                | Mineral |        | Organic  |        |
|      |                         | Tier 3  | Total  | Tier 1/2 | Total  |
| 1990 | 115.64                  | 331.96  | 447.61 | 1.35     | 448.95 |
| 1991 | 114.57                  | 332.26  | 446.84 | 1.33     | 448.17 |
| 1992 | 113.52                  | 332.55  | 446.07 | 1.32     | 447.39 |
| 1993 | 112.33                  | 333.01  | 445.35 | 1.32     | 446.67 |
| 1994 | 111.16                  | 333.48  | 444.64 | 1.33     | 445.97 |
| 1995 | 109.77                  | 333.94  | 443.71 | 1.32     | 445.03 |
| 1996 | 108.31                  | 334.46  | 442.77 | 1.31     | 444.08 |
| 1997 | 106.92                  | 334.98  | 441.91 | 1.29     | 443.20 |
| 1998 | 105.47                  | 335.55  | 441.02 | 1.30     | 442.31 |
| 1999 | 103.25                  | 336.43  | 439.68 | 1.28     | 440.96 |
| 2000 | 103.32                  | 336.28  | 439.60 | 1.28     | 440.89 |

|      |        |        |        |      |        |
|------|--------|--------|--------|------|--------|
| 2001 | 102.46 | 336.48 | 438.95 | 1.27 | 440.22 |
| 2002 | 101.66 | 336.68 | 438.34 | 1.26 | 439.60 |
| 2003 | 100.86 | 336.63 | 437.49 | 1.25 | 438.73 |
| 2004 | 100.29 | 336.44 | 436.72 | 1.24 | 437.96 |
| 2005 | 99.44  | 336.22 | 435.66 | 1.24 | 436.90 |
| 2006 | 98.76  | 336.07 | 434.83 | 1.24 | 436.07 |
| 2007 | 98.17  | 335.89 | 434.06 | 1.23 | 435.29 |
| 2008 | 97.69  | 335.58 | 433.27 | 1.23 | 434.50 |
| 2009 | 97.20  | 335.39 | 432.59 | 1.22 | 433.81 |
| 2010 | 96.78  | 335.10 | 431.88 | 1.22 | 433.10 |
| 2011 | 96.61  | 334.99 | 431.60 | 1.22 | 432.81 |
| 2012 | 96.44  | 334.87 | 431.31 | 1.21 | 432.53 |
| 2013 | 96.44  | 334.87 | 431.31 | 1.21 | 432.53 |
| 2014 | 96.44  | 334.87 | 431.31 | 1.21 | 432.53 |

NRI points on mineral soils are classified into specific crop categories, continuous pasture/rangeland, and other non-agricultural uses for the Tier 2 inventory analysis (Table A-219). NRI points are assigned to IPCC input categories (low, medium, high, and high with organic amendments) according to the classification provided in IPCC (2006). For croplands on federal lands, information on specific cropping systems is not available, so all croplands are assumed to be medium input. In addition, NRI differentiates between improved and unimproved grassland, where improvements include irrigation and interseeding of legumes. Grasslands on federal lands (as identified with the NLCD) are classified according to rangeland condition (nominal, moderately degraded and severely degraded) in areas where information is available. For lands managed for livestock grazing by the Bureau of Land Management (BLM), IPCC rangeland condition classes are interpreted at the state-level from the Rangeland Inventory, Monitoring and Evaluation Report (BLM 2014). In order to estimate uncertainties, probability distribution functions (PDFs) for the NRI land-use data are constructed as multivariate normal based on the total area estimates for each land-use/management category and associated covariance matrix. Through this approach, dependencies in land use are taken into account resulting from the likelihood that current use is correlated with past use. These dependencies occur because as some land use/management categories increase in area, the area of other land use/management categories will decline. The covariance matrix addresses these relationships.

**Table A-219: Total Land Areas by Land-Use and Management System for the Tier 2 Mineral Soil Organic C Approach (Million Hectares)**

| Land-Use/Management System         | Land Areas (million hectares) |               |               |               |               |               |               |               |               |               |               |               |  |
|------------------------------------|-------------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--|
|                                    | 1990                          | 1991          | 1992          | 1993          | 1994          | 1995          | 1996          | 1997          | 1998          | 1999          | 2000          | 2001          |  |
| <b>Cropland Systems</b>            | <b>21.65</b>                  | <b>21.35</b>  | <b>21.02</b>  | <b>20.55</b>  | <b>20.10</b>  | <b>19.68</b>  | <b>19.27</b>  | <b>18.88</b>  | <b>17.65</b>  | <b>16.91</b>  | <b>17.03</b>  | <b>16.82</b>  |  |
| Conservation Reserve Program       | 1.79                          | 2.02          | 2.06          | 1.90          | 1.77          | 1.68          | 1.55          | 1.51          | 1.13          | 0.96          | 1.02          | 1.01          |  |
| High Input Cropping Systems        | 2.25                          | 2.12          | 1.95          | 1.89          | 1.91          | 1.92          | 1.98          | 1.85          | 1.72          | 1.67          | 1.64          | 1.54          |  |
| Medium Input Cropping Systems      | 10.85                         | 10.54         | 10.24         | 10.05         | 9.84          | 9.55          | 9.41          | 9.23          | 8.44          | 8.21          | 8.25          | 8.21          |  |
| Low Input Cropping Systems         | 2.35                          | 2.32          | 2.42          | 2.39          | 2.32          | 2.45          | 2.43          | 2.42          | 2.32          | 2.23          | 2.19          | 2.13          |  |
| Hay with Legumes or Irrigation     | 1.08                          | 1.05          | 1.05          | 1.00          | 1.02          | 0.95          | 0.83          | 0.79          | 0.70          | 0.63          | 0.62          | 0.60          |  |
| Hay, Unimproved                    | 0.80                          | 0.80          | 0.80          | 0.80          | 0.71          | 0.64          | 0.60          | 0.63          | 0.69          | 0.54          | 0.54          | 0.51          |  |
| Perennial Croplands                | 2.43                          | 2.40          | 2.40          | 2.43          | 2.44          | 2.41          | 2.39          | 2.37          | 2.55          | 2.59          | 2.65          | 2.70          |  |
| Rice                               | 0.11                          | 0.10          | 0.11          | 0.09          | 0.10          | 0.08          | 0.09          | 0.08          | 0.10          | 0.08          | 0.12          | 0.11          |  |
| <b>Grassland Systems</b>           | <b>94.00</b>                  | <b>93.22</b>  | <b>92.50</b>  | <b>91.79</b>  | <b>91.06</b>  | <b>90.09</b>  | <b>89.03</b>  | <b>88.04</b>  | <b>87.82</b>  | <b>86.34</b>  | <b>86.29</b>  | <b>85.65</b>  |  |
| Pasture with Legumes or Irrigation | 4.14                          | 3.87          | 3.49          | 3.65          | 3.61          | 3.47          | 3.07          | 2.48          | 2.13          | 2.00          | 2.06          | 1.93          |  |
| Rangelands and Unimproved          |                               |               |               |               |               |               |               |               |               |               |               |               |  |
| Pasture                            | 89.86                         | 89.36         | 89.01         | 88.14         | 87.45         | 86.62         | 85.97         | 85.56         | 85.70         | 84.34         | 84.24         | 83.71         |  |
| <b>Total</b>                       | <b>115.64</b>                 | <b>114.57</b> | <b>113.52</b> | <b>112.33</b> | <b>111.16</b> | <b>109.77</b> | <b>108.31</b> | <b>106.92</b> | <b>105.47</b> | <b>103.25</b> | <b>103.32</b> | <b>102.46</b> |  |

| Land-Use/Management System     | Land Areas (million hectares) |              |              |              |              |              |              |              |              |              |              |              |              |
|--------------------------------|-------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
|                                | 2002                          | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
| <b>Cropland Systems</b>        | <b>16.59</b>                  | <b>16.36</b> | <b>15.96</b> | <b>15.74</b> | <b>15.54</b> | <b>15.38</b> | <b>15.24</b> | <b>15.08</b> | <b>14.94</b> | <b>14.94</b> | <b>14.94</b> | <b>14.94</b> | <b>14.94</b> |
| Conservation Reserve Program   | 0.94                          | 0.81         | 0.52         | 0.58         | 0.56         | 0.58         | 0.60         | 0.49         | 0.46         | 0.46         | 0.46         | 0.46         | 0.46         |
| High Input Cropping Systems    | 1.44                          | 1.40         | 1.38         | 1.36         | 1.32         | 1.26         | 1.25         | 1.28         | 1.24         | 1.24         | 1.24         | 1.24         | 1.24         |
| Medium Input Cropping Systems  | 8.16                          | 8.11         | 8.10         | 8.01         | 7.96         | 7.95         | 7.95         | 7.94         | 7.91         | 7.91         | 7.91         | 7.91         | 7.91         |
| Low Input Cropping Systems     | 2.01                          | 2.03         | 1.98         | 1.86         | 1.80         | 1.74         | 1.67         | 1.58         | 1.58         | 1.58         | 1.58         | 1.58         | 1.58         |
| Hay with Legumes or Irrigation | 0.66                          | 0.67         | 0.66         | 0.65         | 0.61         | 0.61         | 0.61         | 0.58         | 0.59         | 0.59         | 0.59         | 0.59         | 0.59         |
| Hay, Unimproved                | 0.54                          | 0.55         | 0.53         | 0.54         | 0.53         | 0.53         | 0.53         | 0.53         | 0.52         | 0.52         | 0.52         | 0.52         | 0.52         |
| Perennial Croplands            | 2.70                          | 2.68         | 2.68         | 2.65         | 2.66         | 2.62         | 2.54         | 2.57         | 2.54         | 2.54         | 2.54         | 2.54         | 2.54         |
| Rice                           | 0.12                          | 0.11         | 0.11         | 0.10         | 0.10         | 0.10         | 0.09         | 0.09         | 0.10         | 0.10         | 0.10         | 0.10         | 0.10         |
| <b>Grassland Systems</b>       | <b>85.06</b>                  | <b>84.50</b> | <b>84.33</b> | <b>83.69</b> | <b>83.22</b> | <b>82.78</b> | <b>82.46</b> | <b>82.12</b> | <b>81.84</b> | <b>81.67</b> | <b>81.50</b> | <b>81.50</b> | <b>81.50</b> |

|  |               |               |               |              |              |              |              |              |              |              |              |              |              |
|--|---------------|---------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Pasture with Legumes or Irrigation Rangelands and Unimproved Pasture | 1.83          | 1.80          | 1.79          | 1.73         | 1.64         | 1.58         | 1.53         | 1.49         | 1.42         | 1.42         | 1.42         | 1.42         | 1.42         |
| <b>Total</b>   | <b>101.66</b> | <b>100.86</b> | <b>100.29</b> | <b>99.44</b> | <b>98.77</b> | <b>98.17</b> | <b>97.69</b> | <b>97.20</b> | <b>96.78</b> | <b>96.61</b> | <b>96.44</b> | <b>96.44</b> | <b>96.44</b> |

Organic soils are categorized into land-use systems based on drainage (IPCC 2006). Undrained soils are treated as having no loss of organic C or soil N<sub>2</sub>O emissions. Drained soils are subdivided into those used for cultivated cropland, which are assumed to have high drainage and relatively large losses of C, and those used for managed pasture, which are assumed to have less drainage with smaller losses of C. N<sub>2</sub>O emissions are assumed to be similar for both drained croplands and grasslands. Overall, the area of organic soils drained for cropland and grassland has remained relatively stable since 1990 (see Table A-220).

**Table A-220: Total Land Areas for Drained Organic Soils By Land Management Category and Climate Region (Million Hectares)**

| IPCC Land-Use Category for Organic Soils | Land Areas (million ha) |             |             |             |             |             |             |             |             |             |             |             |             |             |
|--|-------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
|  | 1990                    | 1991        | 1992        | 1993        | 1994        | 1995        | 1996        | 1997        | 1998        | 1999        | 2000        | 2001        | 2002        | 2003        |
| <b>Cold Temperate</b>                    |                         |             |             |             |             |             |             |             |             |             |             |             |             |             |
| Cultivated Cropland (high drainage)      | 0.41                    | 0.40        | 0.41        | 0.40        | 0.40        | 0.40        | 0.40        | 0.41        | 0.41        | 0.41        | 0.40        | 0.39        | 0.39        | 0.39        |
| Managed Pasture (low drainage)           | 0.37                    | 0.37        | 0.36        | 0.37        | 0.37        | 0.36        | 0.36        | 0.34        | 0.35        | 0.34        | 0.34        | 0.33        | 0.33        | 0.33        |
| Undrained                                | 0.04                    | 0.05        | 0.05        | 0.04        | 0.04        | 0.04        | 0.04        | 0.03        | 0.02        | 0.03        | 0.04        | 0.03        | 0.02        | 0.02        |
| <b>Total</b>                             | <b>0.82</b>             | <b>0.82</b> | <b>0.81</b> | <b>0.81</b> | <b>0.81</b> | <b>0.80</b> | <b>0.80</b> | <b>0.78</b> | <b>0.78</b> | <b>0.77</b> | <b>0.77</b> | <b>0.75</b> | <b>0.74</b> | <b>0.73</b> |
| <b>Warm Temperate</b>                    |                         |             |             |             |             |             |             |             |             |             |             |             |             |             |
| Cultivated Cropland (high drainage)      | 0.09                    | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.09        | 0.09        |
| Managed Pasture (low drainage)           | 0.08                    | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        |
| Undrained                                | 0.01                    | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        |
| <b>Total</b>                             | <b>0.18</b>             | <b>0.17</b> | <b>0.16</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.16</b> | <b>0.16</b> | <b>0.16</b> | <b>0.17</b> | <b>0.16</b> | <b>0.17</b> | <b>0.17</b> |
| <b>Sub-Tropical</b>                      |                         |             |             |             |             |             |             |             |             |             |             |             |             |             |
| Cultivated Cropland (high drainage)      | 0.20                    | 0.20        | 0.20        | 0.20        | 0.21        | 0.21        | 0.21        | 0.21        | 0.21        | 0.20        | 0.20        | 0.25        | 0.24        | 0.24        |
| Managed Pasture (low drainage)           | 0.14                    | 0.14        | 0.14        | 0.14        | 0.14        | 0.14        | 0.13        | 0.13        | 0.13        | 0.12        | 0.13        | 0.10        | 0.10        | 0.09        |
| Undrained                                | 0.00                    | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.01        | 0.02        | 0.02        | 0.00        | 0.01        | 0.00        |
| <b>Total</b>                             | <b>0.34</b>             | <b>0.35</b> | <b>0.34</b> | <b>0.34</b> | <b>0.35</b> | <b>0.34</b> |

| IPCC Land-Use Category for Organic Soils | Land Areas (million ha) |             |             |             |             |             |             |             |             |             |             |
|--|-------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
|  | 2004                    | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
| <b>Cold Temperate</b>                    |                         |             |             |             |             |             |             |             |             |             |             |
| Cultivated Cropland (high drainage)      | 0.39                    | 0.39        | 0.39        | 0.39        | 0.38        | 0.38        | 0.39        | 0.39        | 0.39        | 0.39        | 0.39        |
| Managed Pasture (low drainage)           | 0.32                    | 0.32        | 0.32        | 0.32        | 0.33        | 0.33        | 0.32        | 0.32        | 0.32        | 0.32        | 0.32        |
| Undrained                                | 0.02                    | 0.02        | 0.02        | 0.02        | 0.03        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        |
| <b>Total</b>                             | <b>0.73</b>             | <b>0.73</b> | <b>0.73</b> | <b>0.73</b> | <b>0.73</b> | <b>0.73</b> | <b>0.72</b> | <b>0.72</b> | <b>0.72</b> | <b>0.72</b> | <b>0.72</b> |
| <b>Warm Temperate</b>                    |                         |             |             |             |             |             |             |             |             |             |             |
| Cultivated Cropland (high drainage)      | 0.09                    | 0.09        | 0.09        | 0.09        | 0.09        | 0.09        | 0.09        | 0.09        | 0.09        | 0.09        | 0.09        |
| Managed Pasture (low drainage)           | 0.08                    | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        | 0.08        |
| Undrained                                | 0.00                    | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        |
| <b>Total</b>                             | <b>0.17</b>             | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> | <b>0.17</b> |
| <b>Sub-Tropical</b>                      |                         |             |             |             |             |             |             |             |             |             |             |
| Cultivated Cropland (high drainage)      | 0.23                    | 0.23        | 0.23        | 0.21        | 0.20        | 0.21        | 0.21        | 0.21        | 0.21        | 0.21        | 0.21        |
| Managed Pasture (low drainage)           | 0.10                    | 0.10        | 0.10        | 0.10        | 0.10        | 0.11        | 0.11        | 0.11        | 0.11        | 0.11        | 0.11        |

|              |             |             |             |             |             |             |             |             |             |             |             |
|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Undrained    | 0.00        | 0.01        | 0.00        | 0.02        | 0.02        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        |
| <b>Total</b> | <b>0.34</b> | <b>0.34</b> | <b>0.33</b> | <b>0.33</b> | <b>0.33</b> | <b>0.32</b> | <b>0.32</b> | <b>0.32</b> | <b>0.32</b> | <b>0.32</b> | <b>0.32</b> |

The harvested rice cultivation area is estimated based on the NRI points classified as flooded rice (Table A-221). Ratoon crops occurs in the Southeast with a second season of rice during the year. Ratoon cropping occurs in Louisiana (LSU 2015 for years 2000 through 2013, 2015) and Texas (TAMU 2015 for years 1993 through 2014), averaging 32 percent and 48 percent of rice acres planted, respectively. Florida also has a large fraction of area with a ratoon crop (45 percent), and ratoon cropping occurs in Arkansas on relatively small fraction of fields estimated at about 1 percent. No data are available about ratoon crops in Missouri or Mississippi, and so amount of ratooning is assumed to be similar to Arkansas. Ratoon rice crops are not grown in California.

**Table A-221: Total Rice Harvested Area Estimated with Tier 1 and 3 Inventory Approaches (Million Hectares)**

| Year | Land Areas (Million Hectares) |        |       |
|------|-------------------------------|--------|-------|
|      | Tier 1                        | Tier 3 | Total |
| 1990 | 0.11                          | 1.53   | 1.64  |
| 1991 | 0.11                          | 1.61   | 1.72  |
| 1992 | 0.11                          | 1.66   | 1.78  |
| 1993 | 0.10                          | 1.64   | 1.74  |
| 1994 | 0.11                          | 1.52   | 1.63  |
| 1995 | 0.07                          | 1.59   | 1.66  |
| 1996 | 0.10                          | 1.55   | 1.65  |
| 1997 | 0.08                          | 1.50   | 1.58  |
| 1998 | 0.11                          | 1.49   | 1.60  |
| 1999 | 0.12                          | 1.51   | 1.63  |
| 2000 | 0.16                          | 1.55   | 1.71  |
| 2001 | 0.11                          | 1.56   | 1.66  |
| 2002 | 0.12                          | 1.72   | 1.83  |
| 2003 | 0.11                          | 1.49   | 1.59  |
| 2004 | 0.12                          | 1.52   | 1.64  |
| 2005 | 0.10                          | 1.75   | 1.85  |
| 2006 | 0.09                          | 1.35   | 1.45  |
| 2007 | 0.09                          | 1.35   | 1.44  |
| 2008 | 0.10                          | 1.30   | 1.40  |
| 2009 | 0.10                          | 1.54   | 1.63  |
| 2010 | 0.10                          | 1.52   | 1.62  |
| 2011 | 0.10                          | 1.52   | 1.62  |
| 2012 | 0.10                          | 1.52   | 1.62  |
| 2013 | 0.10                          | 1.52   | 1.62  |
| 2014 | 0.10                          | 1.52   | 1.62  |

Note: Land use data for 2011 through 2014 are based on the 2010 NRI data product.

**Step 1b: Obtain Management Activity Data for the Tier 3 Method to estimate Soil C Stock Changes, CH<sub>4</sub> and N<sub>2</sub>O Emissions from Mineral Soils**

*Synthetic N Fertilizer Application:* Data on N fertilizer rates are based primarily on the USDA–Economic Research Service Cropping Practices Survey through 1995 (USDA-ERS 1997), which became the Agricultural Resource Management Surveys (ARMS) in 1996 (USDA-ERS 2015)<sup>92</sup>. In these surveys, data on inorganic N fertilization rates are collected for crops simulated by DAYCENT (barley, corn, cotton, dry beans, hay, oats, onions, peanuts, potatoes, rice, sorghum, soybeans, sugar beets, sunflowers, tomatoes, and wheat) in the high production states and for a subset of low production states. These data are used to build a time series of fertilizer application rates for specific crops and states for the 1990 through 1999 time period and 2000 through 2014 time period. If only a single survey is available for a crop, as is the case with sorghum, the rates for the one survey are used for both time periods.

<sup>92</sup> Available online at <<http://www.ers.usda.gov/data-products/arms-farm-financial-and-crop-production-practices/arms-data.aspx>>

Mean fertilizer rates and standard deviations for irrigated and rainfed crops are produced for each state. If a state is not surveyed for a particular crop or if there are not enough data to produce a state-level estimate, then data are aggregated to USDA Farm Production Regions in order to estimate a mean and standard deviation for fertilization rates (Farm Production Regions are groups of states in the United States with similar agricultural commodities). If Farm Production Region data are not available, crop data are aggregated to the entire United States (all major states surveyed) to estimate a mean and standard deviation. Standard deviations for fertilizer rates are used to construct PDFs with log-normal densities in order to address uncertainties in application rates (see Step 2a for discussion of uncertainty methods). The survey summaries also present estimates for fraction of crop acres receiving fertilizer, and these fractions are used to determine if a crop is receiving fertilizer. Alfalfa hay and grass-clover hay are assumed to not be fertilized, but grass hay is fertilized according to rates from published farm enterprise budgets (NRIAI 2003). Total fertilizer application data are found in Table A-222.

Simulations are conducted for the period prior to 1990 in order to initialize the DAYCENT model (see Step 2a), and crop-specific regional fertilizer rates prior to 1990 are based largely on extrapolation/interpolation of fertilizer rates from the years with available data. For crops in some states, little or no data are available, and, therefore, a geographic regional mean is used to simulate N fertilization rates (e.g., no data are available for the State of Alabama during the 1970s and 1980s for corn fertilization rates; therefore, mean values from the southeastern United States are used to simulate fertilization to corn fields in this state).

*Managed Livestock Manure Amendments:*<sup>93</sup> County-level manure addition estimates have been derived from manure N addition rates developed by the USDA Natural Resources Conservation Service (NRCS) (Edmonds et al. 2003). Working with the farm-level crop and animal data from the 1997 Census of Agriculture, USDA-NRCS has coupled estimates of manure N produced with estimates of manure N recoverability by animal waste management system to produce county-level rates of manure N application to cropland and pasture. Edmonds et al. (2003) defined a hierarchy that included 24 crops, permanent pasture, and cropland used as pasture. They estimated the area amended with manure and application rates in 1997 for both manure-producing farms and manure-receiving farms within a county and for two scenarios—before implementation of Comprehensive Nutrient Management Plans (baseline) and after implementation (Edmonds et al. 2003). The goal of nutrient management plans is to apply manure nutrients at a rate meeting plant demand, thus limiting leaching losses of nutrients to groundwater and waterways.

For DAYCENT simulations, the rates for manure-producing farms and manure-receiving farms have been area-weighted and combined to produce a single county-level estimate for the amount of land amended with manure and the manure N application rate for each crop in each county. The estimates were based on the assumption that Comprehensive Nutrient Management Plans have not been fully implemented. This is a conservative assumption because it allows for higher leaching rates due to some over-application of manure to soils. In order to address uncertainty in these data, uniform probability distributions are constructed based on the proportion of land receiving manure versus the amount not receiving manure for each crop type and pasture. For example, if 20 percent of land producing corn in a county is amended with manure, randomly drawing a value equal to or greater than 0 and less than 20 would lead to a simulation with a manure amendment, while drawing a value greater than or equal to 20 and less than 100 would lead to no amendment in the simulation (see Step 2a for further discussion of uncertainty methods).

Edmonds et al. (2003) only provides manure application rate data for 1997, but the amount of managed manure available for soil application changes annually, so the area amended with manure is adjusted relative to 1997 to account for all the manure available for application in other years. Specifically, the manure N available for application in other years is divided by the manure N available in 1997. If the ratio is greater than 1, there is more manure N available in that county relative to the amount in 1997, and so it is assumed a larger area is amended with manure. In contrast, ratios less than one imply less area is amended with manure because there is a lower amount available in the year compared to 1997. The amendment area in each county for 1997 is multiplied by the ratio to reflect the impact of manure N availability on the area amended. The amount of managed manure N available for application to soils is calculated by determining the populations of livestock on feedlots or otherwise housed, requiring collection and management of the manure. The methods are described in the Manure Management section (Section 5.2) and annex (Annex 3.11). The total managed manure N applied to soils is found in Table A-223.

To estimate C inputs (associated with manure N application rates derived from Edmonds et al. (2003), carbon-nitrogen (C:N) ratios for livestock-specific manure types are adapted from the Agricultural Waste Management Field

---

<sup>93</sup> For purposes of the Inventory, total livestock manure is divided into two general categories: (1) managed manure, and (2) unmanaged manure. Managed manure includes manure that is stored in manure management systems such as drylots, pits and lagoons, as well as manure applied to soils through daily spread manure operations. Unmanaged manure encompasses all manure deposited on soils by animals on PRP.

Handbook (USDA 1996), On-Farm Composting Handbook (NRAES 1992), and recoverability factors provided by Edmonds et al (2003). The C:N ratios are applied to county-level estimates of manure N excreted by animal type and management system to produce a weighted county average C:N ratio for manure amendments. The average C:N ratio is used to determine the associated C input for crop amendments derived from Edmonds et al. (2003).

To account for the common practice of reducing inorganic N fertilizer inputs when manure is added to a cropland soil, crop-specific reduction factors are derived from mineral fertilization data for land amended with manure versus land not amended with manure in the ERS 1995 Cropping Practices Survey (USDA-ERS 1997). Mineral N fertilization rates are reduced for crops receiving manure N based on a fraction of the amount of manure N applied, depending on the crop and whether it is irrigated or rainfed. The reduction factors are randomly selected from PDFs with normal densities in order to address uncertainties in the dependence between manure amendments and mineral fertilizer application.

*PRP Manure N:* Another key source of N for grasslands is PRP manure N deposition (i.e., manure deposited by grazing livestock). The total amount of PRP manure N is estimated using methods described in the Manure Management section (Section 5.2) and annex (Annex 3.11). Nitrogen from PRP animal waste deposited on non-federal grasslands in a county is generated by multiplying the total PRP N (based on animal type and population data in a county) by the fraction of non-federal grassland area in the county. PRP manure N input rates for the Tier 3 DAYCENT simulations are estimated by dividing the total PRP manure N amount by the land area associated with non-federal grasslands in the county from the NRI survey data. The total PRP manure N added to soils is found in Table A-223.

*Residue N Inputs:* Crop residue N, fixation by legumes, and N residue inputs from senesced grass litter are included as sources of N to the soil, and are estimated in the DAYCENT simulations as a function of vegetation type, weather, and soil properties. That is, while the model accounts for the contribution of N from crop residues to the soil profile and subsequent N<sub>2</sub>O emissions, this source of mineral soil N is not “activity data” as it is not a model input. The simulated total N inputs of above- and below-ground residue N and fixed N that is not harvested and not burned (the DAYCENT simulations assumed that 3 percent of non-harvested above ground residues for crops are burned<sup>94</sup>) are provided in Table A-224.

*Other N Inputs:* Other N inputs are estimated within the DAYCENT simulation, and thus input data are not required, including mineralization from decomposition of soil organic matter and asymbiotic fixation of N from the atmosphere. Mineralization of soil organic matter will also include the effect of land use change on this process as recommended by the IPCC (2006). The influence of additional inputs of N are estimated in the simulations so that there is full accounting of all emissions from managed lands, as recommended by the IPCC (2006). The simulated N input from residues, soil organic matter mineralization and asymbiotic N fixation are provided in Table A-224.

*Tillage Practices:* Tillage practices are estimated for each cropping system based on data from the Conservation Technology Information Center<sup>95</sup> (CTIC 2004). CTIC compiles data on cropland area under five tillage classes by major crop species and year for each county. Because the surveys involve county-level aggregate area, they do not fully characterize tillage practices as they are applied within a management sequence (e.g., crop rotation). This is particularly true for area estimates of cropland under no-till, which include a relatively high proportion of “intermittent” no-till, where no-till in one year may be followed by tillage in a subsequent year. For example, a common practice in maize-soybean rotations is to use tillage in the maize crop while no-till is used for soybean, such that no-till practices are not continuous in time. Estimates of the area under continuous no-till are provided by experts at CTIC to account for intermittent tillage activity and its impact on soil C (Towery 2001).

Tillage practices are grouped into 3 categories: full, reduced, and no-tillage. Full tillage is defined as multiple tillage operations every year, including significant soil inversion (e.g., plowing, deep disking) and low surface residue coverage. This definition corresponds to the intensive tillage and “reduced” tillage systems as defined by CTIC (2004). No-till is defined as not disturbing the soil except through the use of fertilizer and seed drills and where no-till is applied to all crops in the rotation. Reduced tillage made up the remainder of the cultivated area, including mulch tillage and ridge tillage as defined by CTIC and intermittent no-till. The specific tillage implements and applications used for different crops, rotations, and regions to represent the three tillage classes are derived from the 1995 Cropping Practices Survey by the Economic Research Service (USDA-ERS 1997).

---

<sup>94</sup> Another improvement is to reconcile the amount of crop residues burned with the *Field Burning of Agricultural Residues* source category (Section 5.5).

<sup>95</sup> National scale tillage data are no longer collected by CTIC, and a new data source will be needed, which is a planned improvement.

Tillage data are further processed to construct PDFs. Transitions between tillage systems are based on observed county-level changes in the frequency distribution of the area under full, reduced, and no-till from the 1980s through 2004. Generally, the fraction of full tillage decreased during this time span, with concomitant increases in reduced till and no-till management. Transitions that are modeled and applied to NRI points occurring within a county are full tillage to reduced and no-till, and reduced tillage to no-till. The remaining amount of cropland is assumed to have no change in tillage (e.g., full tillage remained in full tillage). Transition matrices are constructed from CTIC data to represent tillage changes for three time periods, 1980 through 1989, 1990 through 1999, 2000 through 2007. Areas in each of the three tillage classes—full till (FT), reduced till (RT), no-till (NT)—in 1989 (the first year the CTIC data are available) are used for the first time period, data from 1997 are used for the second time period, and data from 2004 are used for the last time period. Percentage areas of cropland in each county are calculated for each possible transition (e.g., FT→FT, FT→RT, FT→NT, RT→RT, RT→NT) to obtain a probability for each tillage transition at an NRI point. It is assumed that there are no transitions for NT→FT or NT→NT after accounting for NT systems that have intermittent tillage. Uniform probability distributions are established for each tillage scenario in the county. For example, a particular crop rotation had 80 percent chance of remaining in full tillage over the two decades, a 15 percent chance of a transition from full to reduced tillage and a 5 percent chance of a transition from full to no-till. The uniform distribution is subdivided into three segments with random draws in the Monte Carlo simulation (discussed in Step 2b) leading to full tillage over the entire time period if the value is greater than or equal to 0 and less than 80, a transition from full to reduced till if the random draw is equal to or greater than 80 and less than 95, or a transition from full to no-till if the draw is greater than or equal to 95. See step 2b for additional discussion of the uncertainty analysis.

*Irrigation:* NRI (USDA-NRCS 2013) differentiates between irrigated and non-irrigated land, but does not provide more detailed information on the type and intensity of irrigation. Hence, irrigation is modeled by assuming that applied water to field capacity with intervals between irrigation events where the soils drain to about 60 percent of field capacity.

*Daily Weather Data:* Daily maximum/minimum temperature and precipitation data are based on gridded weather data from the PRISM Climate Group (2015). It is necessary to use computer-generated weather data because weather station data do not exist near all NRI points, and moreover weather station data are for a point in space. The PRISM product uses this information with interpolation algorithms to derive weather patterns for areas between these stations (Daly et al. 1998). PRISM weather data are available for the U.S. from 1981 through 2010 at a 4 km resolution. Each NRI point is assigned the PRISM weather data for the grid cell containing the point.

*Enhanced Vegetation Index:* The Enhanced Vegetation Index (EVI) from the MODIS vegetation products, (MOD13Q1 and MYD13Q1) is an input to DAYCENT for estimating net primary production using the NASA-CASA production algorithm (Potter et al. 1993, 2007). MODIS imagery is collected on a nominal 8 day-time frequency when combining the two products. A best approximation of the daily time series of EVI data is derived using a smoothing process based on the Savitzky-Golay Filter (Savitzky and Golay 1964) after pre-screening for outliers and for cloud-free, high quality data as identified in the MODIS data product quality layer. The NASA-CASA production algorithm is only used for the following crops: corn, soybeans, sorghum, cotton, wheat and other close-grown crops such as barley and oats.<sup>96</sup>

The MODIS EVI products have a 250 m spatial resolution, and some pixels in images have mixed land uses and crop types at this resolution, which is problematic for estimating NPP associated with a specific crop at a NRI point. Therefore, a threshold of 90 percent purity in an individual pixel is the cutoff for estimating NPP using the EVI data derived from the imagery (i.e., pixels with less than 90 percent purity for a crop are assumed to generate bias in the resulting NPP estimates). The USDA-NASS Crop Data Layer (CDL) (Johnson and Mueller 2010) is used to determine the purity levels of the EVI data. CDL data have a 30 to 58 m spatial resolution, depending on the year. The level of purity for individual pixels in the MODIS EVI products is determined by aggregating the crop cover data in CDL to the 250m resolution of the EVI data. In this step, the percent cover of individual crops is determined for the 250m EVI pixels. Pixels that did not meet a 90 percent purity level for any crop are eliminated from the dataset. CDL did not provide full coverage of crop maps for the conterminous United States until 2009 so it is not possible to evaluate purity for the entire cropland area prior to 2009. The nearest pixel with at least 90 percent purity for a crop is assigned to the NRI point based on a 10 km buffer surrounding the survey location. EVI data are not assigned to a point if there are no pixels with at least 90 percent purity within the 10 km buffer. In these cases, production is simulated with a single value for the maximum daily NPP, which is reduced if there is water, temperature or nutrient stress affecting the plants growth.

*Water Management for Rice Cultivation:* While rice crop production in the U.S. includes a minor amount of land with mid-season drainage or alternate wet-dry periods, the majority of rice growers use continuously flooded water management systems (Hardke 2015; UCCE 2015; Hollier 1999; Way et al. 2014). Therefore, continuous flooding is applied

---

<sup>96</sup> Additional crops and grassland will be used with the NASA-CASA method in the future, as a planned improvement.

to all rice cultivation areas in the inventory. Winter flooding is another key practice associated with water management in rice fields. Winter flooding occurs on 34 percent of rice fields in California (Miller et al. 2010; Fleskes et al. 2005), and approximately 21 percent of the fields in Arkansas (Wilson and Branson 2005 and 2006; Wilson and Runsick 2007 and 2008; Wilson et al. 2009 and 2010; Hardke and Wilson 2013 and 2014; Hardke 2015). No data are available on winter flooding for Texas, Louisiana, Florida, Missouri, or Mississippi. For these states, the average amount of flooding is assumed to be similar to Arkansas. In addition, the amount of winter flooding is assumed to be relatively constant over the Inventory time period.

*Organic Amendments for Rice Cultivation:* Rice straw is not typically harvested from fields in the U.S. The C input from rice straw is simulated directly within the DAYCENT model for the Tier 3 method. For the Tier 1 method, residue inputs are assumed to be left on the field for more than 30 days prior to cultivation and flooding for the next crop, with the exception of ratoon crops, which are assumed to have residues on the field for less than 30 days prior to the next crop. To estimate the amount of rice straw, crop yield data (except rice in Florida) are compiled from USDA NASS QuickStats (USDA 2015). Rice yield data for Florida are estimated separately because yield data are not collected by USDA. Total rice production for Florida is determined using NRI crop areas, and total yields are based on average primary and ratoon rice yields from Deren (2002). Relative proportions of ratoon crops are derived from information in several publications (Schueneman 1997, 1999, 2000, 2001; Deren 2002; Kirstein 2003, 2004, 2006; Cantens 2004, 2005; Gonzalez 2007 through 2014). The yields are multiplied by residue:crop product ratios from Strehler and Stützel (1987), to estimate rice straw input amounts for the Tier 1 method.

*Soil Properties:* Soil texture and natural drainage capacity (i.e., hydric vs. non-hydric soil characterization) are the main soil variables used as input to the DAYCENT model. Texture is one of the main controls on soil C turnover and stabilization in the DAYCENT model, which uses particle size fractions of sand (50-2,000  $\mu\text{m}$ ), silt (2-50  $\mu\text{m}$ ), and clay (<2  $\mu\text{m}$ ) as inputs. Hydric conditions are poorly-drained, and hence prone to have a high water table for part of the year in their native (pre-cultivation) condition. Non-hydric soils are moderately to well-drained.<sup>97</sup> Poorly drained soils can be subject to anaerobic (lack of oxygen) conditions if water inputs (precipitation and irrigation) exceed water losses from drainage and evapotranspiration. Depending on moisture conditions, hydric soils can range from being fully aerobic to completely anaerobic, varying over the year. Decomposition rates are modified according to a linear function that varies from 0.3 under completely anaerobic conditions to 1.0 under fully aerobic conditions (default parameters in DAYCENT).<sup>98</sup> Other soil characteristics needed in the simulation, such as field capacity and wilting-point water contents, are estimated from soil texture data using a standardized hydraulic properties calculator (Saxton et al. 1986). Soil input data are derived from Soil Survey Geographic Database (SSURGO) (Soil Survey Staff 2015). The data are based on field measurements collected as part of soil survey and mapping. Each NRI point is assigned the dominant soil component in the polygon containing the point from the SSURGO data product.

#### **Step 1c: Obtain Additional Management Activity Data for the Tier 1 Method to estimate Soil N<sub>2</sub>O Emissions from Mineral Soils**

*Synthetic N Fertilizer:* A process-of-elimination approach is used to estimate synthetic N fertilizer additions to crops in the Tier 1 method. The total amount of fertilizer used on-farms has been estimated by the USGS from 1990 through 2001 on a county scale from fertilizer sales data (Ruddy et al. 2006). For 2002 through 2014, county-level fertilizer used on-farms is adjusted based on annual fluctuations in total U.S. fertilizer sales (AAPFCO 1995 through 2014). Fertilizer application data are available for crops and grasslands simulated by DAYCENT (discussed in Step 1a section for Tier 3). Thus, the amount of N applied to crops in the Tier 1 method (i.e., not simulated by DAYCENT) is assumed to be the remainder of the fertilizer used on farms after subtracting the amount applied to crops and non-federal grasslands simulated by DAYCENT. The differences are aggregated to the state level, and PDFs are derived based on uncertainties in the amount of N applied to crops and non-federal grasslands for the Tier 3 method. Total fertilizer application to crops in the Tier 1 method is found in Table A-225.

*Managed Livestock Manure and Other Organic Amendments:* Manure N that is not applied to crops and grassland simulated by DAYCENT is assumed to be applied to other crops that are included in the Tier 1 method. Estimates of total national annual N additions from other commercial organic fertilizers are derived from organic fertilizer statistics (TVA 1991 through 1994; AAPFCO 1995 through 2014). Commercial organic fertilizers include dried blood, tankage, compost,

---

<sup>97</sup> Artificial drainage (e.g., ditch- or tile-drainage) is simulated as a management variable.

<sup>98</sup> Hydric soils are primarily subject to anaerobic conditions outside the plant growing season (i.e., in the absence of active plant water uptake). Soils that are water-logged during much of the year are typically classified as organic soils (e.g., peat), which are not simulated with the DAYCENT model.

and other; dried manure and sewage sludge that are used as commercial fertilizer are subtracted from totals to avoid double counting. The dried manure N is counted with the non-commercial manure applications, and sewage sludge is assumed to be applied only to grasslands. The organic fertilizer data, which are recorded in mass units of fertilizer, had to be converted to mass units of N by multiplying the consumption values by the average organic fertilizer N content of 0.5 percent (AAPFCO 2000). The fertilizer consumption data are recorded in “fertilizer year” totals, (i.e., July to June), but are converted to calendar year totals. This is done by assuming that approximately 35 percent of fertilizer usage occurred from July to December and 65 percent from January to June (TVA 1992b). Values for July to December are not available for calendar year 2013 so a “least squares line” statistical extrapolation using the previous 5 years of data is used to arrive at an approximate value. PDFs are derived for the organic fertilizer applications assuming a default  $\pm 50$  percent uncertainty. Annual consumption of other organic fertilizers is presented in Table A-226. The fate of manure N is summarized in Table A-223.

*PRP Manure N:* Soil N<sub>2</sub>O emissions from PRP manure N deposited on federal grasslands are estimated with a Tier 1 method. PRP manure N data are derived using methods described in the Manure Management section (Section 5.2) and Annex 3.11. PRP N deposited on federal grasslands is calculated using a process of elimination approach. The amount of PRP N generated by DAYCENT model simulations of non-federal grasslands was subtracted from total PRP N and this difference was assumed to be applied to federal grasslands. The total PRP manure N added to soils is found in Table A-223.

*Sewage Sludge Amendments:* Sewage sludge is generated from the treatment of raw sewage in public or private wastewater treatment works and is typically used as a soil amendment, or is sent to waste disposal facilities, such as landfills. In this Inventory, all sewage sludge that is amended to agricultural soils is assumed to be applied to grasslands. Estimates of the amounts of sewage sludge N applied to agricultural lands are derived from national data on sewage sludge generation, disposition, and N content. Total sewage sludge generation data for 1990 through 2012, in dry mass units, are obtained from AAPFCO (1995 through 2014). Values for 2013 and 2014 were not available so a “least squares line” statistical extrapolation using the previous 5 years of data was used to arrive at an approximate value. The total sludge generation estimates are then converted to units of N by applying an average N content of 69 percent (AAPFCO 2000), and disaggregated into use and disposal practices using historical data in EPA (1993) and NEBRA (2007). The use and disposal practices are agricultural land application, other land application, surface disposal, incineration, landfilling, ocean dumping (ended in 1992), and other disposal methods. The resulting estimates of sewage sludge N applied to agricultural land are used to estimate N<sub>2</sub>O emissions from agricultural soil management; the estimates of sewage sludge N applied to other land and surface-disposed are used in estimating N<sub>2</sub>O fluxes from soils in Settlements Remaining Settlements (see section 6.9 of the Land Use, Land-Use Change, and Forestry chapter). Sewage sludge disposal data are provided in Table A-227.

*Residue N Inputs:* Soil N<sub>2</sub>O emissions for residue N inputs from croplands that are not simulated by DAYCENT are estimated with a Tier 1 method. Annual crop production statistics for all major commodity and specialty crops are taken from U.S. Department of Agriculture crop production reports (USDA-NASS 2015). Total production for each crop is converted to tons of dry matter product using the residue dry matter fractions shown in Table A-228. Dry matter yield is then converted to tons of above- and below-ground biomass N. Above-ground biomass is calculated by using linear equations to estimate above-ground biomass given dry matter crop yields, and below-ground biomass is calculated by multiplying above-ground biomass by the below-to-above-ground biomass ratio. N inputs are estimated by multiplying above- and below-ground biomass by respective N concentrations and by the portion of cropland that was not simulated by DAYCENT. All ratios and equations used to calculate residue N inputs are from IPCC (2006) and Williams (2006). PDFs are derived assuming a  $\pm 50$  percent uncertainty in the yield estimates (USDA-NASS does not provide uncertainty), along with uncertainties provided by the IPCC (2006) for dry matter fractions, above-ground residue, ratio of below-ground to above-ground biomass, and residue N fractions. The resulting annual residue N inputs are presented in Table A-229.

#### **Step 1d: Obtain Additional Management Activity Data for the Tier 2 Method to estimate Soil C Stock Changes in Mineral Soils**

*Tillage Practices:* For the Tier 2 method that is used to estimate soil organic C stock changes, PDFs are constructed for the CTIC tillage data (CTIC 2004) as bivariate normal on a log-ratio scale to reflect negative dependence among tillage classes. This structure ensured that simulated tillage percentages are non-negative and summed to 100 percent. CTIC data do not differentiate between continuous and intermittent use of no-tillage, which is important for estimating SOC storage. Thus, regionally based estimates for continuous no-tillage (defined as 5 or more years of continuous use) are modified based on consultation with CTIC experts, as discussed in Step 1a (downward adjustment of total no-tillage area based on the amount of no-tillage that is rotated with more intensive tillage practices) (Towery 2001).

*Managed Livestock Manure Amendments:* USDA provides information on the amount of land amended with manure for 1997 based on manure production data and field-scale surveys detailing application rates that had been collected

in the *Census of Agriculture* (Edmonds et al. 2003). Similar to the DAYCENT model discussion in Step1b, the amount of land receiving manure is based on the estimates provided by Edmonds et al. (2003), as a proportion of crop and grassland amended with manure within individual climate regions. The resulting proportions are used to re-classify a portion of crop and grassland into a new management category. Specifically, a portion of medium input cropping systems is re-classified as high input, and a portion of the high input systems is re-classified as high input with amendment. In grassland systems, the estimated proportions for land amended with manure are used to re-classify a portion of nominally-managed grassland as improved, and a portion of improved grassland as improved with high input. These classification approaches are consistent with the IPCC inventory methodology (IPCC 2006). Uncertainties in the amount of land amended with manure are based on the sample variance at the climate region scale, assuming normal density PDFs (i.e., variance of the climate region estimates, which are derived from county-scale proportions).

**Sewage Sludge Amendments:** Sewage sludge is generated from the treatment of raw sewage in public or private wastewater treatment facilities and is typically used as a soil amendment or is sent for waste disposal to landfills. In this Inventory, all sewage sludge that is amended to agricultural soils is assumed to be applied to grasslands. See section on sewage sludge in Step 1c for more information about the methods used to derive sewage sludge N estimates. The total amount of sewage sludge N is given in Table A-227. Sewage sludge N is assumed to be applied at the assimilative capacity provided in Kellogg et al. (2000), which is the amount of nutrients taken up by a crop and removed at harvest, representing the recommended application rate for manure amendments. This capacity varies from year to year, because it is based on specific crop yields during the respective year (Kellogg et al. 2000). Total sewage sludge N available for application is divided by the assimilative capacity to estimate the total land area over which sewage sludge had been applied. The resulting estimates are used for the estimation of soil C stock change.

**CRP Enrollment after 2010:** The change in enrollment for the Conservation Reserve Program after 2010 is based on the amount of land under active contracts from 2011 through 2014 relative to 2010 (USDA-FSA 2014).

**Wetland Reserve:** Wetlands enrolled in the Conservation Reserve Program have been restored in the Northern Prairie Pothole Region through the Partners for Wildlife Program funded by the U.S. Fish and Wildlife Service (USFWS 2010). The area of restored wetlands is estimated from contract agreements (Euliss and Gleason 2002). While the contracts provide reasonable estimates of the amount of land restored in the region, they do not provide the information necessary to estimate uncertainty. Consequently, a  $\pm 50$  percent range is used to construct the PDFs for the uncertainty analysis.

**Table A-222: Synthetic Fertilizer N Added to Tier 3 Crops (kt N)**

|              | 1990  | 1995  | 2000   | 2001  | 2002  | 2003  | 2004   | 2005   | 2006  | 2007   | 2008   | 2009   | 2010  | 2011  | 2012  | 2013  | 2014  |
|--------------|-------|-------|--------|-------|-------|-------|--------|--------|-------|--------|--------|--------|-------|-------|-------|-------|-------|
| Fertilizer N | 9,809 | 9,529 | 10,063 | 9,877 | 9,996 | 9,918 | 10,113 | 10,168 | 9,863 | 10,613 | 10,085 | 10,117 | 9,598 | 9,598 | 9,599 | 9,599 | 9,599 |

**Table A-223: Fate of Livestock Manure Nitrogen (kt N)**

| Activity  | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Managed Manure N Applied to Tier 3 Cropland and Non-federal Grasslands <sup>a,b</sup> | 957          | 887          | 1,123        | 1,083        | 1,089        | 1,071        | 1,070        | 1,075        | 1,055        | 1,152        | 1,108        | 1,187        | 1,211        | 1,211        | 1,211        | 1,211        | 1,211        |
| Managed Manure N Applied to Tier 1 Cropland <sup>c</sup>                              | 1,453        | 1,645        | 1,487        | 1,521        | 1,553        | 1,585        | 1,511        | 1,545        | 1,647        | 1,576        | 1,599        | 1,499        | 1,469        | 1,498        | 1,526        | 1,519        | 1,519        |
| Managed Manure N Applied to Grasslands  | 53           | 55           | 72           | 72           | 71           | 72           | 68           | 68           | 68           | 67           | 68           | 71           | 72           | 72           | 72           | 72           | 72           |
| Pasture, Range, & Paddock Manure N  | 4,266        | 4,533        | 4,149        | 4,143        | 4,146        | 4,155        | 4,106        | 4,133        | 4,174        | 4,061        | 4,009        | 3,978        | 3,931        | 3,843        | 3,743        | 3,672        | 3,672        |
| <b>Total</b>  | <b>6,729</b> | <b>7,121</b> | <b>6,831</b> | <b>6,818</b> | <b>6,860</b> | <b>6,883</b> | <b>6,755</b> | <b>6,822</b> | <b>6,944</b> | <b>6,856</b> | <b>6,782</b> | <b>6,735</b> | <b>6,683</b> | <b>6,625</b> | <b>6,552</b> | <b>6,475</b> | <b>6,475</b> |

<sup>a</sup> Accounts for N volatilized and leached/runoff during treatment, storage and transport before soil application.

<sup>b</sup> Includes managed manure and daily spread manure amendments

<sup>c</sup> Totals may not sum exactly due to rounding.

**Table A-224: Crop Residue N and Other N Inputs to Tier 3 Crops as Simulated by DAYCENT (kt N)**

| Activity                             | 1990   | 1995   | 2000   | 2001   | 2002   | 2003   | 2004   | 2005   | 2006   | 2007   | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   |
|--------------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Residue N <sup>a</sup>               | 4,345  | 4,785  | 4,728  | 4,624  | 4,693  | 4,834  | 4,286  | 4,648  | 4,446  | 4,565  | 4,455  | 4,487  | 4,875  | 4,875  | 4,875  | 4,875  | 4,875  |
| Mineralization & Asymbiotic Fixation | 13,323 | 13,089 | 12,766 | 13,177 | 12,887 | 13,319 | 14,414 | 13,236 | 12,957 | 13,508 | 13,362 | 14,016 | 15,254 | 15,254 | 15,255 | 15,255 | 15,255 |

<sup>a</sup> Residue N inputs include unharvested fixed N from legumes as well as crop residue N.

**Table A-225: Synthetic Fertilizer N Added to Tier 1 Crops (kt N)**

| Activity     | 1990  | 1995  | 2000  | 2001  | 2002  | 2003  | 2004  | 2005  | 2006  | 2007  | 2008  | 2009  | 2010  | 2011  | 2012  | 2013  | 2014  |
|--------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Fertilizer N | 1,263 | 1,389 | 1,309 | 1,213 | 1,344 | 1,569 | 1,541 | 1,438 | 1,579 | 1,573 | 1,424 | 1,289 | 1,574 | 2,003 | 2,153 | 1,710 | 1,663 |

**Table A-226: Other Organic Commercial Fertilizer Consumption on Agricultural Lands (kt N)**

| Activity   | 1990 | 1995 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 |
|--|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Other Commercial Organic Fertilizer N <sup>a</sup> | 4    | 10   | 9    | 7    | 8    | 8    | 9    | 10   | 12   | 15   | 12   | 10   | 10   | 11   | 11   | 10   | 10   |

<sup>a</sup> Includes dried blood, tankage, compost, other. Excludes dried manure and sewage sludge used as commercial fertilizer to avoid double counting.

**Table A-227: Sewage Sludge Nitrogen by Disposal Practice (kt N)**

| Disposal Practice             | 1990      | 1995       | 2000       | 2001       | 2002       | 2003       | 2004       | 2005       | 2006       | 2007       | 2008       | 2009       | 2010       | 2011       | 2012       | 2013       | 2014       |
|-------------------------------|-----------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Applied to Agricultural Soils | 52        | 68         | 84         | 86         | 89         | 91         | 94         | 98         | 101        | 104        | 107        | 110        | 113        | 116        | 119        | 122        | 122        |
| Other Land Application        | 25        | 28         | 30         | 30         | 30         | 30         | 30         | 31         | 31         | 32         | 32         | 32         | 32         | 33         | 33         | 33         | 33         |
| Surface Disposal              | 20        | 16         | 10         | 9          | 8          | 6          | 5          | 5          | 4          | 4          | 3          | 3          | 3          | 2          | 2          | 2          | 2          |
| <b>Total</b>                  | <b>97</b> | <b>111</b> | <b>124</b> | <b>125</b> | <b>127</b> | <b>128</b> | <b>130</b> | <b>134</b> | <b>137</b> | <b>140</b> | <b>142</b> | <b>145</b> | <b>148</b> | <b>151</b> | <b>153</b> | <b>156</b> | <b>156</b> |

Note: Totals may not sum due to independent rounding.

**Table A-228: Key Assumptions for Crop Production in the Tier 1 Method**

| Crop             | Dry Matter Fraction of Harvested Product | Above-ground Residue |           | Ratio of Below-ground Residue to Above-ground Biomass | Residue N Fraction |              |
|------------------|--|----------------------|-----------|---|--------------------|--------------|
|                  |  | Slope                | Intercept |   | Above-ground       | Below-ground |
| Alfalfa          | 0.9                                      | 0.29                 | 0         | 0.4   | 0.027              | 0.019        |
| Artichokes       | 0.16                                     | 0.3                  | 0         | 0.2   | 0.006              | 0.009        |
| Asparagus        | 0.07                                     | 0.5                  | 0         | 0.2   | 0.006              | 0.009        |
| Barley           | 0.89                                     | 0.98                 | 0.59      | 0.22  | 0.007              | 0.014        |
| Beans            | 0.91                                     | 1.13                 | 0.85      | 0.19  | 0.008              | 0.008        |
| Beets            | 0.22                                     | 0.1                  | 1.06      | 0.2   | 0.019              | 0.014        |
| Broccoli         | 0.09                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Brussels sprouts | 0.14                                     | 0.1                  | 0         | 0.11  | 0.0093             | 0.0093       |
| Cabbage          | 0.08                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Cantaloup        | 0.06                                     | 1.77                 | 0         | 0.04  | 0.006              | 0.009        |
| Carrots          | 0.13                                     | 0.46                 | 0.02      | 0.15  | 0.019              | 0.014        |
| Cauliflower      | 0.08                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Celery           | 0.05                                     | 0.23                 | 0         | 0.11  | 0.006              | 0.009        |
| Collard greens   | 0.08                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Corn             | 0.87                                     | 1.03                 | 0.61      | 0.22  | 0.006              | 0.007        |
| Corn silage      | 0.3                                      | 0.3                  | 0         | 0.22  | 0.006              | 0.007        |
| Cotton           | 0.93                                     | 1.49                 | 4.41      | 0.13  | 0.012              | 0.007        |
| Cucumbers        | 0.04                                     | 1.77                 | 0         | 0.03  | 0.006              | 0.009        |
| Dry beans        | 0.9                                      | 0.36                 | 0.68      | 0.19  | 0.01               | 0.01         |
| Eggplant         | 0.08                                     | 1.77                 | 0         | 0.04  | 0.006              | 0.009        |
| Escarole         | 0.94                                     | 1.07                 | 1.54      | 0.2   | 0.016              | 0.014        |
| Flaxseed         | 0.88                                     | 1.09                 | 0.88      | 0.22  | 0.006              | 0.009        |
| Garlic           | 0.11                                     | 0.23                 | 0         | 0.15  | 0.019              | 0.014        |
| Hay              | 0.9                                      | 0.2075               | 0         | 0.505   | 0.018              | 0.01375      |
| Honeydew         | 0.06                                     | 1.77                 | 0         | 0.04  | 0.006              | 0.009        |
| Kale greens      | 0.08                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Lentils          | 0.91                                     | 1.13                 | 0.85      | 0.2   | 0.019              | 0.014        |
| Lettuce          | 0.04                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Millet           | 0.88                                     | 1.09                 | 0.88      | 0.22  | 0.006              | 0.009        |
| Mustard greens   | 0.08                                     | 0.1                  | 0         | 0.11  | 0.006              | 0.009        |
| Oats             | 0.89                                     | 0.91                 | 0.89      | 0.25  | 0.007              | 0.008        |
| Okra             | 0.1                                      | 1.4                  | 0         | 0.14  | 0.006              | 0.009        |
| Onions           | 0.12                                     | 0.23                 | 0         | 0.14  | 0.019              | 0.014        |
| Peanuts          | 0.94                                     | 1.07                 | 1.54      | 0.2   | 0.016              | 0.014        |
| Peas             | 0.91                                     | 1.13                 | 0.85      | 0.05  | 0.011              | 0.008        |

|                |       |      |      |      |       |       |
|----------------|-------|------|------|------|-------|-------|
| Peppermint Oil | 0.09  | 1    | 0    | 1.48 | 0.008 | 0.008 |
| Peppers        | 0.08  | 1.4  | 0    | 0.14 | 0.006 | 0.009 |
| Potatoes       | 0.22  | 0.1  | 1.06 | 0.2  | 0.019 | 0.014 |
| Pumpkins       | 0.1   | 1.77 | 0    | 0.04 | 0.006 | 0.009 |
| Radishes       | 0.05  | 1.21 | 0.46 | 0.15 | 0.019 | 0.014 |
| Rapeseed       | 0.88  | 1.09 | 0.88 | 0.22 | 0.006 | 0.009 |
| Rice           | 0.89  | 0.95 | 2.46 | 0.16 | 0.007 | 0.009 |
| Rye            | 0.88  | 1.09 | 0.88 | 0.22 | 0.006 | 0.009 |
| Safflower      | 0.88  | 1.09 | 0.88 | 0.22 | 0.006 | 0.009 |
| Sorghum        | 0.89  | 0.88 | 1.33 | 0.22 | 0.007 | 0.006 |
| Sorghum silage | 0.3   | 0.3  | 0    | 0.22 | 0.007 | 0.006 |
| Soybeans       | 0.91  | 0.93 | 1.35 | 0.19 | 0.008 | 0.008 |
| Spearmint Oil  | 0.09  | 1    | 0    | 1.48 | 0.008 | 0.008 |
| Spinach        | 0.08  | 0.9  | 0    | 0.11 | 0.006 | 0.009 |
| Squash         | 0.05  | 1.57 | 0    | 0.04 | 0.006 | 0.009 |
| Sugarcane      | 0.25  | 0.41 | 0    | 0.16 | 0.007 | 0.005 |
| Sunflower      | 0.88  | 1.09 | 0.88 | 0.22 | 0.006 | 0.009 |
| Sweet potatoes | 0.35  | 0.27 | 1.74 | 0.15 | 0.019 | 0.014 |
| Tobacco        | 0.87  | 0.3  | 0    | 0.4  | 0.008 | 0.018 |
| Tomatoes       | 0.05  | 0.59 | 0.57 | 0.19 | 0.006 | 0.009 |
| Turnip greens  | 0.08  | 0.1  | 0    | 0.11 | 0.006 | 0.009 |
| Watermelon     | 0.085 | 1.77 | 0    | 0.04 | 0.006 | 0.009 |
| Wheat          | 0.89  | 1.51 | 0.52 | 0.24 | 0.006 | 0.009 |

**Table A-229: Nitrogen in Crop Residues Retained on Soils Producing Crops not Simulated by DAYCENT (kt N)**

| Crop Type        | 1990    | 1995    | 2000    | 2001   | 2002   | 2003    | 2004    | 2005    | 2006    | 2007    |
|------------------|---------|---------|---------|--------|--------|---------|---------|---------|---------|---------|
| Alfalfa          | 68,294  | 61,587  | 41,929  | 34,943 | 34,573 | 34,231  | 35,115  | 34,279  | 30,769  | 30,213  |
| Artichokes       | 4       | 4       | 4       | 4      | 4      | 4       | 3       | 2       | 3       | 3       |
| Asparagus        | 5       | 5       | 5       | 5      | 4      | 4       | 5       | 2       | 2       | 2       |
| Barley           | 5,900   | 4,574   | 3,165   | 2,228  | 2,487  | 2,517   | 1,454   | 1,969   | 1,405   | 1,822   |
| Beans            | 1,888   | 1,881   | 1,836   | 1,697  | 1,706  | 1,623   | 1,832   | 1,020   | 1,233   | 1,170   |
| Beets            | 1,795   | 2,282   | 1,548   | 1,192  | 746    | 1,033   | 999     | 893     | 1,928   | 505     |
| Broccoli         | 18      | 18      | 18      | 17     | 16     | 17      | 18      | 10      | 12      | 12      |
| Brussels sprouts | 1       | 1       | 1       | 1      | 0      | 0       | 0       | 0       | 0       | 0       |
| Cabbage          | 22      | 22      | 21      | 21     | 19     | 17      | 20      | 11      | 13      | 13      |
| Cantaloup        | 103     | 102     | 91      | 95     | 89     | 90      | 92      | 50      | 57      | 58      |
| Carrots          | 403     | 401     | 345     | 345    | 298    | 324     | 335     | 195     | 208     | 200     |
| Cauliflower      | 6       | 6       | 6       | 5      | 5      | 5       | 5       | 3       | 4       | 4       |
| Celery           | 16      | 16      | 14      | 15     | 14     | 15      | 16      | 9       | 10      | 10      |
| Collard greens   | 0       | 0       | 1       | 1      | 0      | 0       | 0       | 0       | 0       | 0       |
| Corn             | 157,096 | 111,997 | 109,000 | 99,081 | 96,451 | 101,376 | 120,332 | 107,939 | 106,560 | 118,834 |
| Corn silage      | 9,419   | 8,109   | 6,779   | 6,513  | 4,855  | 6,431   | 6,867   | 5,223   | 5,754   | 5,751   |
| Cotton           | 40,634  | 52,686  | 52,456  | 57,126 | 51,494 | 50,597  | 48,554  | 55,583  | 52,681  | 38,979  |
| Cucumbers        | 72      | 72      | 63      | 61     | 60     | 59      | 62      | 32      | 36      | 38      |
| Dry beans        | 2,264   | 2,099   | 1,576   | 1,205  | 1,671  | 1,300   | 1,155   | 913     | 1,019   | 1,006   |
| Eggplant         | 5       | 5       | 9       | 10     | 0      | 0       | 0       | 0       | 0       | 0       |
| Escarole         | 91      | 90      | 128     | 129    | 0      | 0       | 0       | 0       | 0       | 0       |
| Flaxseed         | 1,234   | 732     | 2,064   | 2,055  | 1,925  | 1,079   | 694     | 819     | 846     | 481     |
| Garlic           | 29      | 29      | 25      | 27     | 25     | 28      | 25      | 13      | 14      | 13      |
| Hay              | 124,991 | 115,116 | 88,222  | 85,416 | 76,725 | 80,977  | 81,141  | 76,822  | 68,257  | 70,885  |
| Honeydew         | 24      | 24      | 21      | 20     | 20     | 21      | 21      | 10      | 12      | 12      |
| Kale greens      | 0       | 0       | 1       | 1      | 0      | 0       | 0       | 0       | 0       | 0       |
| Lentils          | 267     | 563     | 661     | 634    | 566    | 585     | 985     | 720     | 617     | 564     |
| Lettuce          | 40      | 40      | 39      | 39     | 38     | 41      | 44      | 23      | 29      | 26      |
| Millet           | 3,898   | 3,992   | 1,278   | 2,609  | 597    | 1,040   | 829     | 463     | 596     | 978     |
| Mustard greens   | 0       | 0       | 1       | 1      | 0      | 0       | 0       | 0       | 0       | 0       |
| Oats             | 3,848   | 2,151   | 1,287   | 847    | 1,203  | 1,579   | 1,528   | 1,487   | 1,745   | 1,303   |
| Okra             | 0       | 0       | 1       | 1      | 0      | 0       | 0       | 0       | 0       | 0       |
| Onions           | 378     | 376     | 354     | 344    | 323    | 349     | 423     | 210     | 247     | 261     |

|                |                |                |                |                |                |                |                |                |                |                |
|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Peanuts        | 13,885         | 11,663         | 9,748          | 10,315         | 8,565          | 11,914         | 14,400         | 13,586         | 13,023         | 13,702         |
| Peas           | 308            | 526            | 348            | 387            | 484            | 554            | 1,163          | 854            | 1,003          | 1,087          |
| Peppermint Oil | 1              | 2              | 1              | 1              | 1              | 1              | 1              | 1              | 1              | 1              |
| Peppers        | 113            | 113            | 112            | 109            | 99             | 110            | 122            | 68             | 78             | 74             |
| Potatoes       | 4,344          | 3,843          | 2,931          | 2,577          | 2,886          | 3,216          | 4,156          | 3,555          | 3,945          | 5,490          |
| Pumpkins       | 71             | 70             | 64             | 59             | 56             | 55             | 73             | 44             | 51             | 54             |
| Radishes       | 0              | 0              | 17             | 19             | 0              | 0              | 0              | 0              | 0              | 0              |
| Rapeseed       | 79             | 15             | 17             | 12             | 10             | 2              | 12             | 2              | 1              | 2              |
| Rice           | 6,719          | 4,601          | 9,910          | 7,407          | 8,515          | 7,120          | 8,577          | 7,541          | 7,033          | 6,454          |
| Rye            | 2,541          | 2,603          | 1,390          | 1,060          | 878            | 735            | 477            | 274            | 422            | 427            |
| Safflower      | 984            | 892            | 695            | 623            | 674            | 736            | 713            | 788            | 794            | 694            |
| Sorghum        | 7,944          | 5,731          | 4,118          | 4,517          | 3,458          | 4,036          | 3,253          | 1,867          | 1,505          | 2,000          |
| Sorghum silage | 873            | 504            | 387            | 185            | 162            | 117            | 104            | 217            | 14             | 38             |
| Soybeans       | 72,959         | 54,617         | 51,834         | 55,888         | 52,456         | 48,123         | 59,561         | 60,651         | 57,665         | 53,229         |
| Spearmint Oil  | 1              | 0              | 0              | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| Spinach        | 21             | 21             | 24             | 21             | 22             | 25             | 30             | 19             | 18             | 17             |
| Squash         | 29             | 29             | 27             | 24             | 26             | 22             | 23             | 14             | 17             | 13             |
| Sugarcane      | 19,061         | 12,147         | 15,376         | 15,019         | 15,472         | 15,441         | 17,283         | 16,033         | 18,923         | 16,296         |
| Sunflower      | 415            | 504            | 251            | 431            | 433            | 379            | 247            | 152            | 247            | 209            |
| Sweet potatoes | 437            | 424            | 409            | 424            | 352            | 431            | 462            | 254            | 305            | 328            |
| Tobacco        | 12,508         | 11,597         | 8,865          | 8,073          | 7,305          | 5,978          | 5,911          | 3,458          | 5,808          | 5,644          |
| Tomatoes       | 587            | 584            | 575            | 517            | 584            | 518            | 654            | 324            | 392            | 429            |
| Turnip greens  | 0              | 0              | 1              | 1              | 0              | 0              | 0              | 0              | 0              | 0              |
| Watermelon     | 251            | 250            | 221            | 242            | 222            | 221            | 226            | 129            | 164            | 149            |
| Wheat          | 40,306         | 30,994         | 26,666         | 22,252         | 21,399         | 26,664         | 22,829         | 23,199         | 17,851         | 21,863         |
| <b>Total</b>   | <b>607,182</b> | <b>510,710</b> | <b>446,941</b> | <b>426,852</b> | <b>399,973</b> | <b>411,740</b> | <b>442,832</b> | <b>421,735</b> | <b>403,316</b> | <b>401,341</b> |

| <b>Crop Type</b> | <b>2008</b> | <b>2009</b> | <b>2010</b> | <b>2011</b> | <b>2012</b> | <b>2013</b> | <b>2014</b> |
|------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Alfalfa          | 30,605      | 29,463      | 29,711      | 29,711      | 26,300      | 28,309      | 29,095      |
| Artichokes       | 3           | 4           | 3           | 4           | 3           | 3           | 3           |
| Asparagus        | 2           | 2           | 1           | 2           | 2           | 2           | 1           |
| Barley           | 1,781       | 2,443       | 2,622       | 2,495       | 2,426       | 2,565       | 2,610       |
| Beans            | 1,239       | 1,440       | 1,257       | 1,419       | 1,289       | 1,256       | 1,166       |
| Beets            | 1,238       | 1,030       | 1,339       | 1,198       | 1,397       | 1,364       | 1,328       |
| Broccoli         | 13          | 15          | 13          | 16          | 15          | 17          | 16          |
| Brussels sprouts | 0           | 0           | 0           | 0           | 0           | 0           | 0           |
| Cabbage          | 14          | 15          | 14          | 16          | 13          | 16          | 15          |
| Cantaloup        | 57          | 68          | 59          | 77          | 58          | 67          | 52          |
| Carrots          | 214         | 232         | 208         | 264         | 232         | 258         | 244         |
| Cauliflower      | 4           | 5           | 4           | 5           | 4           | 5           | 5           |
| Celery           | 11          | 13          | 12          | 15          | 13          | 12          | 12          |
| Collard greens   | 0           | 0           | 0           | 0           | 0           | 0           | 0           |
| Corn             | 111,272     | 117,718     | 104,642     | 101,834     | 86,596      | 108,517     | 116,268     |
| Corn silage      | 5,893       | 5,795       | 5,790       | 5,520       | 4,710       | 5,640       | 6,030       |
| Cotton           | 38,123      | 29,267      | 39,439      | 39,193      | 40,317      | 39,522      | 39,712      |
| Cucumbers        | 39          | 47          | 40          | 45          | 40          | 41          | 37          |
| Dry beans        | 1,047       | 1,261       | 1,409       | 1,152       | 1,485       | 1,234       | 1,406       |
| Eggplant         | 0           | 0           | 0           | 0           | 0           | 0           | 0           |
| Escarole         | 0           | 0           | 0           | 0           | 0           | 0           | 0           |
| Flaxseed         | 475         | 545         | 577         | 278         | 533         | 214         | 434         |
| Garlic           | 14          | 15          | 13          | 19          | 16          | 16          | 15          |
| Hay              | 70,656      | 69,663      | 66,882      | 64,680      | 58,899      | 64,128      | 67,431      |
| Honeydew         | 11          | 13          | 12          | 15          | 11          | 13          | 13          |
| Kale greens      | 0           | 0           | 0           | 0           | 0           | 0           | 0           |
| Lentils          | 437         | 1,087       | 1,458       | 1,099       | 1,024       | 972         | 631         |
| Lettuce          | 25          | 29          | 28          | 36          | 31          | 30          | 28          |

|                |                |                |                |                |                |                |                |
|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Millet         | 875            | 517            | 574            | 672            | 270            | 913            | 716            |
| Mustard greens | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| Oats           | 1,234          | 1,338          | 1,553          | 1,426          | 1,493          | 1,544          | 1,610          |
| Okra           | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| Onions         | 259            | 314            | 272            | 356            | 288            | 302            | 295            |
| Peanuts        | 13,798         | 15,525         | 11,364         | 11,550         | 13,620         | 13,092         | 12,895         |
| Peas           | 949            | 1,444          | 1,081          | 596            | 941            | 1,393          | 1,439          |
| Peppermint Oil | 1              | 1              | 1              | 1              | 1              | 1              | 1              |
| Peppers        | 78             | 100            | 82             | 109            | 91             | 91             | 91             |
| Potatoes       | 4,295          | 2,984          | 5,073          | 5,056          | 5,130          | 5,181          | 5,242          |
| Pumpkins       | 53             | 55             | 57             | 73             | 69             | 69             | 76             |
| Radishes       | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| Rapeseed       | 0              | 2              | 4              | 4              | 6              | 2              | 3              |
| Rice           | 6,910          | 6,964          | 6,813          | 7,074          | 7,414          | 7,592          | 7,498          |
| Rye            | 547            | 457            | 420            | 489            | 496            | 415            | 428            |
| Safflower      | 842            | 633            | 567            | 453            | 469            | 432            | 549            |
| Sorghum        | 1,602          | 1,225          | 1,132          | 919            | 866            | 985            | 1,081          |
| Sorghum silage | 102            | 207            | 38             | 31             | 36             | 43             | 39             |
| Soybeans       | 53,300         | 58,487         | 57,209         | 55,863         | 54,070         | 57,660         | 60,801         |
| Spearmint Oil  | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| Spinach        | 19             | 26             | 19             | 26             | 20             | 23             | 22             |
| Squash         | 15             | 19             | 15             | 21             | 19             | 17             | 15             |
| Sugarcane      | 13,743         | 13,828         | 11,800         | 12,669         | 13,471         | 12,754         | 13,207         |
| Sunflower      | 186            | 268            | 131            | 128            | 133            | 126            | 132            |
| Sweet potatoes | 350            | 433            | 470            | 686            | 563            | 557            | 622            |
| Tobacco        | 5,380          | 4,430          | 5,023          | 4,346          | 5,349          | 4,797          | 5,462          |
| Tomatoes       | 435            | 586            | 473            | 599            | 526            | 543            | 577            |
| Turnip greens  | 0              | 0              | 0              | 0              | 0              | 0              | 0              |
| Watermelon     | 168            | 196            | 187            | 210            | 177            | 190            | 157            |
| Wheat          | 26,921         | 27,396         | 25,467         | 24,219         | 25,510         | 25,960         | 24,262         |
| <b>Total</b>   | <b>395,233</b> | <b>397,604</b> | <b>385,359</b> | <b>376,667</b> | <b>356,441</b> | <b>388,884</b> | <b>403,771</b> |

### **Step 1e: Additional Activity Data for Indirect N<sub>2</sub>O Emissions**

A portion of the N that is applied as synthetic fertilizer, livestock manure, sewage sludge, and other organic amendments volatilizes as NH<sub>3</sub> and NO<sub>x</sub>. In turn, this N is returned to soils through atmospheric deposition, thereby increasing mineral N availability and enhancing N<sub>2</sub>O production. Additional N is lost from soils through leaching as water percolates through a soil profile and through runoff with overland water flow. N losses from leaching and runoff enter groundwater and waterways, from which a portion is emitted as N<sub>2</sub>O. However, N leaching is assumed to be an insignificant source of indirect N<sub>2</sub>O in cropland and grassland systems where the amount of precipitation plus irrigation does not exceed 80 percent of the potential evapotranspiration. These areas are typically semi-arid to arid, and nitrate leaching to groundwater is a relatively uncommon event; moreover IPCC (2006) recommends limiting the amount of nitrate leaching assumed to be a source of indirect N<sub>2</sub>O emissions based on precipitation, irrigation and potential evapotranspiration.

The activity data for synthetic fertilizer, livestock manure, other organic amendments, residue N inputs, sewage sludge N, and other N inputs are the same as those used in the calculation of direct emissions from agricultural mineral soils, and may be found in Table A-222 through Table A-227, and Table A-229.

Using the DAYCENT model, volatilization and leaching/surface run-off of N from soils is computed internally for crops and non-federal grasslands in the Tier 3 method. DAYCENT simulates the processes leading to these losses of N based on environmental conditions (i.e., weather patterns and soil characteristics), management impacts (e.g., plowing, irrigation, harvest), and soil N availability. Note that the DAYCENT model accounts for losses of N from all anthropogenic activity, not just the inputs of N from mineral fertilization and organic amendments, which are addressed in the Tier 1 methodology. Similarly, the N available for producing indirect emissions resulting from grassland management as well as deposited PRP manure is also estimated by DAYCENT. Estimated leaching losses of N from DAYCENT are not used in

the indirect N<sub>2</sub>O calculation if the amount of precipitation plus irrigation did not exceed 80 percent of the potential evapotranspiration. Volatilized losses of N are summed for each day in the annual cycle to provide an estimate of the amount of N subject to indirect N<sub>2</sub>O emissions. In addition, the daily losses of N through leaching and runoff in overland flow are summed for the annual cycle. Uncertainty in the estimates is derived from uncertainties in the activity data for the N inputs (i.e., fertilizer and organic amendments; see Step 1a for further information).

The Tier 1 method is used to estimate N losses from mineral soils due to volatilization and leaching/runoff for crops, sewage sludge applications, and PRP manure on federal grasslands, which is simulated by DAYCENT. To estimate volatilized losses, synthetic fertilizers, manure, sewage sludge, and other organic N inputs are multiplied by the fraction subject to gaseous losses using the respective default values of 0.1 kg N/kg N added as mineral fertilizers and 0.2 kg N/kg N added as manure (IPCC 2006). Uncertainty in the volatilized N ranges from 0.03-0.3 kg NH<sub>3</sub>-N+NO<sub>x</sub>-N/kg N for synthetic fertilizer and 0.05-0.5 kg NH<sub>3</sub>-N+NO<sub>x</sub>-N/kg N for organic amendments (IPCC 2006). Leaching/runoff losses of N are estimated by summing the N additions from synthetic and other organic fertilizers, manure, sewage sludge, and above- and below-ground crop residues, and then multiplying by the default fraction subject to leaching/runoff losses of 0.3 kg N/kg N applied, with an uncertainty from 0.1-0.8 kg NO<sub>3</sub>-N/kg N (IPCC 2006). However, N leaching is assumed to be an insignificant source of indirect N<sub>2</sub>O emissions if the amount of precipitation plus irrigation did not exceed 80 percent of the potential evapotranspiration. PDFs are derived for each of the N inputs in the same manner as direct N<sub>2</sub>O emissions, discussed in Steps 1a and 1c.

Volatilized N is summed for losses from croplands and grasslands. Similarly, the annual amounts of N lost from soil profiles through leaching and surface runoff are summed to obtain the total losses for this pathway.

## **Step 2: Estimate Soil Organic C Stock Changes, Direct N<sub>2</sub>O Emissions from Mineral Soils, and CH<sub>4</sub> Emissions from Rice Cultivation**

In this step, soil organic C stock changes, N<sub>2</sub>O emissions, and CH<sub>4</sub> emissions from rice cultivation are estimated for cropland and non-federal grasslands. Three methods are used to estimate soil organic C stock changes, direct N<sub>2</sub>O emissions from mineral soils, and CH<sub>4</sub> emissions from rice cultivation. The DAYCENT process-based model is used for the croplands and non-federal grasslands included in the Tier 3 method. A Tier 2 method is used to estimate soil organic C stock changes for crop histories that included crops that were not simulated by DAYCENT and land use change other than conversions between cropland and grassland. A Tier 1 methodology is used to estimate N<sub>2</sub>O emissions from crops that are not simulated by DAYCENT, PRP manure N deposition on federal grasslands, and CH<sub>4</sub> emissions from rice cultivation. Soil organic C stock changes and N<sub>2</sub>O emissions are not estimated for federal grasslands (other than the effect of PRP manure N), but are under evaluation as a planned improvement and may be estimated in future inventories.

### **Step 2a: Estimate Soil Organic C Stock Changes, N<sub>2</sub>O Emissions, and CH<sub>4</sub> emissions for Crops and Non-Federal Grassland with the Tier 3 DAYCENT Model**

Crops that are simulated with DAYCENT include alfalfa hay, barley, corn, cotton, dry beans, grass hay, grass-clover hay, oats, onions, peanuts, potatoes, rice, sorghum, soybeans, sugar beets, sunflowers, tomatoes, and wheat, which combined represent approximately 87 to 90 percent of total cropland in the U.S. The DAYCENT simulations also included all non-federal grasslands in the U.S.

The methodology description is divided into two sub-steps. First, the model is used to establish the initial conditions and C stocks for 1979, which is the last year before the NRI survey is initiated. In the second sub-step, DAYCENT is used to estimate changes in soil organic C stocks, direct N<sub>2</sub>O emissions, and CH<sub>4</sub> emissions from rice cultivation based on the land-use and management histories recorded in the NRI from 1990 through 2010 (USDA-NRCS 2013).

*Simulate Initial Conditions (Pre-NRI Conditions):* DAYCENT model initialization involves two steps, with the goal of estimating the most accurate stock for the pre-NRI history, and the distribution of organic C among the pools represented in the model (e.g., Structural, Metabolic, Active, Slow, and Passive). Each pool has a different turnover rate (representing the heterogeneous nature of soil organic matter), and the amount of C in each pool at any point in time influences the forward trajectory of the total soil organic C storage. There is currently no national set of soil C measurements that can be used for establishing initial conditions in the model. Sensitivity analysis of the soil organic C algorithms showed that the rate of change of soil organic matter is relatively insensitive to the *amount* of total soil organic C but is highly sensitive to the relative *distribution* of C among different pools (Parton et al. 1987). By simulating the historical land use prior to the inventory period, initial pool distributions are estimated in an unbiased way.

The first step involves running the model to a steady-state condition (e.g., equilibrium) under native vegetation, historical climate data based on the PRISM product (1981 through 2010), and the soil physical attributes for the NRI points.

Native vegetation is represented at the MLRA level for pre-settlement time periods in the United States. The model simulates 5,000 years in the pre-settlement era in order to achieve a steady-state condition.

The second step is to simulate the period of time from European settlement and expansion of agriculture to the beginning of the NRI survey, representing the influence of historic land-use change and management, particularly the conversion of native vegetation to agricultural uses. This encompasses a varying time period from land conversion (depending on historical settlement patterns) to 1979. The information on historical cropping practices used for DAYCENT simulations has been gathered from a variety of sources, ranging from the historical accounts of farming practices reported in the literature (e.g., Miner 1998) to national level databases (e.g., NASS 2004). A detailed description of the data sources and assumptions used in constructing the base history scenarios of agricultural practices can be found in Williams and Paustian (2005).

*NRI History Simulations:* After model initialization, DAYCENT is used to simulate the NRI land use and management histories from 1979 through 2010.<sup>99</sup> The simulations address the influence of soil management on direct N<sub>2</sub>O emissions, soil organic C stock changes and losses of N from the profile through leaching/runoff and volatilization. The NRI histories identify the land use and land use change histories for the NRI survey locations, as well as cropping patterns and irrigation history (see Step 1a for description of the NRI data). The input data for the model simulations also include the PRISM weather dataset and SSURGO soils data, synthetic N fertilizer rates, managed manure amendments to cropland and grassland, manure deposition on grasslands (i.e., PRP), tillage histories and EVI data (See Step 1b for description of the inputs). The total number of DAYCENT simulations is over 18 million with a 100 repeated simulations (i.e., iterations) for each NRI point location in a Monte Carlo Analysis. The simulation system incorporates a dedicated MySQL database server and a 30-node parallel processing computer cluster. Input/output operations are managed by a set of run executive programs written in PERL.

The simulations for the NRI history are integrated with the uncertainty analysis. Evaluating uncertainty is an integral part of the analysis and includes three components: (1) uncertainty in the main activity data inputs affecting soil C and N<sub>2</sub>O emissions (input uncertainty); (2) uncertainty in the model formulation and parameterization (structural uncertainty); and (3) uncertainty in the land-use and management system areas (scaling uncertainty) (Ogle et al. 2010; Del Grosso et al. 2010). For component 1, input uncertainty is evaluated for fertilization management, manure applications, and tillage, which are primary management activity data that are supplemental to the NRI observations and have significant influence on soil organic C dynamics, soil N<sub>2</sub>O and CH<sub>4</sub> emissions. As described in Step 1b, PDFs are derived from surveys at the county scale for the inputs in most cases. In addition, uncertainty is included for predictions of EVI data that are needed to fill-data gaps and extend the time series (see Enhanced Vegetation Index in Step 1b). To represent uncertainty in all of these inputs, a Monte-Carlo Analysis is used with 100 iterations for each NRI point; random draws are made from PDFs for fertilizer, manure application, tillage, and EVI predictions. As described above, an adjustment factor is also selected from PDFs with normal densities to represent the dependence between manure amendments and N fertilizer application rates.

The second component deals with uncertainty inherent in model formulation and parameterization. This component is the largest source of uncertainty in the Tier 3 model-based inventory analysis, accounting for more than 80 percent of the overall uncertainty in the final estimates (Ogle et al. 2010; Del Grosso et al. 2010). An empirically-based procedure is applied to develop a structural uncertainty estimator from the relationship between modeled results and field measurements from agricultural experiments (Ogle et al. 2007). For soil organic C, the DAYCENT model is evaluated with measurements from 84 long-term field experiments that have over 900 treatment observations, representing a variety of management conditions (e.g., variation in crop rotation, tillage, fertilization rates, and manure amendments). There are 27 experimental sites available with 93 treatment observations to evaluate structural uncertainty in the N<sub>2</sub>O emission predictions from DAYCENT (Del Grosso et al. 2010). There are 10 experiments with 126 treatment observations for CH<sub>4</sub> emissions from rice cultivation. The inputs to the model are essentially known in the simulations for the long-term experiments, and, therefore, the analysis is designed to evaluate uncertainties associated with the model structure (i.e., model algorithms and parameterization). USDA is developing a national soil monitoring network to evaluate the Inventory in the future (Spencer et al. 2011).

The relationship between modeled soil organic C stocks and field measurements are statistically analyzed using linear-mixed effect modeling techniques. Additional fixed effects are included in the mixed effect model if they explained

---

<sup>99</sup> The estimated soil C stock change in 2010 is currently assumed to represent the changes between 2011 and 2014. More recent data will be incorporated in the future to extend the time series of land use and management data.

significant variation in the relationship between modeled and measured stocks (i.e., if they met an alpha level of 0.05 for significance). Several variables are tested, including land-use class; type of tillage; cropping system; geographic location; climate; soil texture; time since the management change; original land cover (i.e., forest or grassland); grain harvest as predicted by the model compared to the experimental values; and variation in fertilizer and residue management. The final cropland model includes variables for modeled soil organic C inclusion of hay/pasture in cropping rotations, use of no-till, set-aside lands, organic matter amendments, and inclusion of bare fallow in the rotation, which are significant at an alpha level of 0.05. The final grassland model only included the model soil organic C. These fixed effects are used to make an adjustment to modeled values due to biases that are creating significant mismatches between the modeled and measured stocks. For soil N<sub>2</sub>O, simulated DAYCENT emissions are a highly significant predictor of the measurements, with a p-value of <0.01. Several other variables are considered in the statistical model to evaluate if DAYCENT exhibits bias under certain conditions related to climate, soil types, and management practices. Random effects are included in the model to capture the dependence in time series and data collected from the same site, which are needed to estimate appropriate standard deviations for parameter coefficients. For rice CH<sub>4</sub> emissions, simulated DAYCENT emissions are a significant predictor of measured emission, similar to the results for soil N<sub>2</sub>O emissions. Several other variables are tested including soil characteristics, geographic location (i.e., state), and management practices (e.g., with and without winter flooding). The only other significant variable is geographic location because the model does not predict emissions as accurately for California as other rice-producing states. Random effects are included to capture the dependence in time series and the data collected from the same site.

A Monte Carlo approach is used to apply the uncertainty estimator (Ogle et al. 2010). Parameter values for the statistical equation (i.e., fixed effects) are selected from their joint probability distribution, as well as random error associated with fine-scale estimates at NRI points, and the residual or unexplained error associated with the linear mixed-effect model. The estimate and associated management information is then used as input into the equation, and adjusted values are computed for each C stock, N<sub>2</sub>O and CH<sub>4</sub> emissions estimate. The variance of the adjusted estimates is computed from the 100 simulated values from the Monte Carlo analysis.

The third element is the uncertainty associated with scaling the DAYCENT results for each NRI point to the entire land base, using the expansion factors provided with the NRI survey dataset. The expansion factors represent the number of hectares associated with the land-use and management history for a particular point. This uncertainty is determined by computing the variances from a set of replicated weights for the expansion factor. For the land base that is simulated with the DAYCENT model, soil organic C stock changes are provided in Table A-230, soil N<sub>2</sub>O emissions are provided in Table A-231, and rice cultivation CH<sub>4</sub> emissions in Table A-232.

**Table A-230: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Land Base Simulated with the Tier 3 DAYCENT Model-Based Approach (MMT CO<sub>2</sub> Eq.)**

| Year | Cropland Remaining Cropland |                    | Land Converted to Cropland |                 | Grassland Remaining Grassland |                  | Land Converted to Grassland |                   |
|------|-----------------------------|--------------------|----------------------------|-----------------|-------------------------------|------------------|-----------------------------|-------------------|
|      | Estimate                    | 95% CI             | Estimate                   | 95% CI          | Estimate                      | 95% CI           | Estimate                    | 95% CI            |
| 1990 | (56.17)                     | (85.13) to (27.20) | 12.91                      | 3.82 to 21.99   | (17.87)                       | (57.42) to 21.68 | (3.81)                      | (8.28) to 0.67    |
| 1991 | (57.01)                     | (88.19) to (25.83) | 12.80                      | 3.55 to 22.05   | (8.44)                        | (41.12) to 24.25 | (4.72)                      | (9.42) to (0.01)  |
| 1992 | (58.33)                     | (92.79) to (23.88) | 13.63                      | 3.30 to 23.96   | (9.35)                        | (42.21) to 23.52 | (4.67)                      | (9.62) to 0.29    |
| 1993 | (37.09)                     | (70.77) to (3.40)  | 11.65                      | 1.84 to 21.46   | (8.68)                        | (46.74) to 29.38 | (5.12)                      | (11.21) to 0.96   |
| 1994 | (48.47)                     | (77.48) to (19.47) | 6.47                       | (4.12) to 17.06 | (24.18)                       | (66.44) to 18.08 | (5.42)                      | (11.13) to 0.28   |
| 1995 | (42.96)                     | (74.79) to (11.13) | 13.62                      | 3.64 to 23.61   | (7.36)                        | (40.08) to 25.37 | (5.61)                      | (11.06) to (0.16) |
| 1996 | (50.59)                     | (83.49) to (17.70) | 11.18                      | 0.04 to 22.33   | (16.46)                       | (54.41) to 21.49 | (5.96)                      | (11.88) to (0.04) |
| 1997 | (47.13)                     | (78.45) to (15.81) | 13.12                      | 2.94 to 23.30   | (9.94)                        | (53.81) to 33.94 | (6.64)                      | (12.54) to (0.74) |
| 1998 | (33.49)                     | (63.73) to (3.26)  | 7.63                       | (2.64) to 17.90 | (13.09)                       | (50.63) to 24.44 | (6.83)                      | (13.57) to (0.10) |
| 1999 | (46.54)                     | (80.91) to (12.17) | 8.52                       | (3.15) to 20.18 | (3.76)                        | (43.22) to 35.69 | (6.88)                      | (14.67) to 0.91   |
| 2000 | (51.06)                     | (79.03) to (23.10) | 8.41                       | (2.24) to 19.06 | (19.22)                       | (52.43) to 13.99 | (7.14)                      | (15.49) to 1.21   |
| 2001 | (50.85)                     | (83.65) to (18.06) | 8.16                       | (2.66) to 18.99 | (8.46)                        | (51.41) to 34.49 | (7.63)                      | (15.88) to 0.62   |
| 2002 | (42.21)                     | (72.67) to (11.75) | 8.62                       | (0.84) to 18.08 | (6.80)                        | (42.05) to 28.46 | (8.16)                      | (16.50) to 0.18   |
| 2003 | (32.95)                     | (64.45) to (1.45)  | 7.68                       | (2.02) to 17.37 | (5.15)                        | (45.88) to 35.58 | (7.31)                      | (16.01) to 1.39   |
| 2004 | (38.11)                     | (69.39) to (6.84)  | 6.93                       | (3.16) to 17.03 | (7.38)                        | (45.0) to 30.24  | (8.01)                      | (17.05) to 1.03   |
| 2005 | (37.19)                     | (69.31) to (5.07)  | 7.94                       | (1.98) to 17.86 | (5.26)                        | (44.04) to 33.52 | (7.23)                      | (16.46) to 2.0    |
| 2006 | (37.66)                     | (71.38) to (3.95)  | 7.94                       | (2.31) to 18.18 | (14.35)                       | (55.71) to 27.01 | (7.98)                      | (16.23) to 0.27   |
| 2007 | (37.76)                     | (69.09) to (6.44)  | 5.84                       | (3.94) to 15.62 | 1.26                          | (34.83) to 37.35 | (7.87)                      | (16.75) to 1.0    |
| 2008 | (29.74)                     | (63.12) to 3.64    | 6.79                       | (1.73) to 15.31 | (8.44)                        | (47.35) to 30.48 | (8.13)                      | (17.73) to 1.47   |
| 2009 | (22.33)                     | (51.73) to 7.07    | 5.77                       | (5.39) to 16.93 | (10.79)                       | (45.11) to 23.53 | (7.98)                      | (16.89) to 0.94   |
| 2010 | (22.03)                     | (56.14) to 12.07   | 6.65                       | (3.93) to 17.23 | (9.05)                        | (46.92) to 28.82 | (7.67)                      | (16.74) to 1.39   |
| 2011 | (36.74)                     | (68.95) to (4.53)  | 4.54                       | (5.34) to 14.42 | 1.06                          | (35.84) to 37.95 | (7.16)                      | (15.94) to 1.61   |
| 2012 | (36.75)                     | (68.95) to (4.54)  | 4.54                       | (5.34) to 14.42 | 1.08                          | (35.81) to 37.97 | (7.16)                      | (15.94) to 1.61   |

|      |         |                   |      |                 |      |                  |        |                 |
|------|---------|-------------------|------|-----------------|------|------------------|--------|-----------------|
| 2013 | (36.75) | (68.95) to (4.54) | 4.54 | (5.34) to 14.42 | 1.08 | (35.81) to 37.97 | (7.16) | (15.94) to 1.61 |
| 2014 | (36.75) | (68.95) to (4.54) | 4.54 | (5.34) to 14.42 | 1.08 | (35.81) to 37.97 | (7.16) | (15.94) to 1.61 |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

**Table A-231: Annual N<sub>2</sub>O Emissions (95% Confidence Interval) for the Land Base Simulated with the Tier 3 DAYCENT Model-Based Approach (MMT CO<sub>2</sub> Eq.)**

| Year | Tier 3 Cropland |                  | Non-Federal Grasslands |                |
|------|-----------------|------------------|------------------------|----------------|
|      | Estimate        | 95% CI           | Estimate               | 95% CI         |
| 1990 | 153.1           | 144.73 to 164.12 | 65.6                   | 61.13 to 72.06 |
| 1991 | 150.0           | 142.04 to 160.62 | 66.2                   | 61.67 to 72.46 |
| 1992 | 151.0           | 142.84 to 161.88 | 66.6                   | 62.37 to 72.78 |
| 1993 | 153.1           | 144.62 to 164.31 | 66.2                   | 61.92 to 72.23 |
| 1994 | 149.0           | 141.15 to 159.37 | 62.5                   | 58.41 to 68.26 |
| 1995 | 149.5           | 141.58 to 160.1  | 64.1                   | 59.93 to 69.93 |
| 1996 | 149.9           | 141.91 to 160.64 | 66.5                   | 62 to 72.93    |
| 1997 | 149.9           | 141.91 to 160.56 | 67.1                   | 62.78 to 73.07 |
| 1998 | 160.0           | 151.21 to 171.61 | 72.0                   | 67.3 to 78.85  |
| 1999 | 151.2           | 143.17 to 161.89 | 62.1                   | 58.28 to 67.58 |
| 2000 | 155.9           | 147.86 to 166.29 | 63.7                   | 59.41 to 70.08 |
| 2001 | 152.9           | 144.92 to 163.37 | 64.3                   | 60.03 to 70.38 |
| 2002 | 152.9           | 144.94 to 163.33 | 67.1                   | 62.54 to 73.77 |
| 2003 | 155.6           | 147.63 to 166.14 | 66.9                   | 62.5 to 73.33  |
| 2004 | 164.9           | 156.33 to 176.14 | 77.4                   | 72.13 to 85.08 |
| 2005 | 155.2           | 147.08 to 165.93 | 66.4                   | 62.24 to 72.41 |
| 2006 | 155.1           | 147.08 to 165.62 | 68.4                   | 63.93 to 74.82 |
| 2007 | 162.1           | 153.37 to 173.56 | 70.6                   | 65.83 to 77.54 |
| 2008 | 156.0           | 147.76 to 166.86 | 67.8                   | 63.45 to 74.11 |
| 2009 | 156.8           | 148.59 to 167.66 | 70.7                   | 66.05 to 77.44 |
| 2010 | 166.5           | 157.5 to 178.49  | 70.3                   | 65.83 to 76.87 |
| 2011 | 165.7           | 156.83 to 177.23 | 70.5                   | 65.81 to 77.27 |
| 2012 | 165.9           | 157.07 to 177.48 | 70.4                   | 65.71 to 77.16 |
| 2013 | 165.2           | 156.36 to 176.72 | 70.4                   | 65.71 to 77.16 |
| 2014 | 165.2           | 156.36 to 176.72 | 70.4                   | 65.71 to 77.16 |

**Table A-232: Annual CH<sub>4</sub> Emissions (95% Confidence Interval) for Rice Cultivation Simulated with the Tier 3 DAYCENT Model-Based Approach (MMT CO<sub>2</sub> Eq.)**

| Year | Estimate | 95% CI         |
|------|----------|----------------|
| 1990 | 12.08    | 10.55 to 13.61 |
| 1991 | 11.93    | 10.57 to 13.28 |
| 1992 | 11.91    | 10.18 to 13.64 |
| 1993 | 12.65    | 11.18 to 14.13 |
| 1994 | 9.32     | 8.37 to 10.27  |
| 1995 | 12.41    | 10.78 to 14.04 |
| 1996 | 11.45    | 10.02 to 12.87 |
| 1997 | 10.47    | 9.37 to 11.57  |
| 1998 | 12.39    | 9.94 to 14.85  |
| 1999 | 11.37    | 8.04 to 14.71  |
| 2000 | 12.83    | 10.21 to 15.45 |
| 2001 | 10.97    | 8.87 to 13.06  |
| 2002 | 14.01    | 9.98 to 18.03  |
| 2003 | 9.87     | 8.55 to 11.20  |
| 2004 | 9.70     | 7.78 to 11.63  |
| 2005 | 11.93    | 10.26 to 13.60 |
| 2006 | 9.25     | 7.74 to 10.76  |
| 2007 | 8.95     | 6.93 to 10.98  |
| 2008 | 7.84     | 6.21 to 9.46   |
| 2009 | 9.88     | 6.81 to 12.95  |
| 2010 | 10.71    | 8.78 to 12.64  |
| 2011 | 10.71    | 8.76 to 12.66  |
| 2012 | 10.72    | 8.79 to 12.65  |
| 2013 | 10.71    | 8.78 to 12.64  |

In DAYCENT, the model cannot distinguish among the original sources of N after the mineral N enters the soil pools, and therefore it is not possible to determine which management activity led to specific N<sub>2</sub>O emissions. This means, for example, that N<sub>2</sub>O emissions from applied synthetic fertilizer cannot be separated from emissions due to other N inputs, such as crop residues. It is desirable, however, to report emissions associated with specific N inputs. Thus, for each NRI point, the N inputs in a simulation are determined for anthropogenic practices discussed in IPCC (2006), including synthetic mineral N fertilization, organic amendments, and crop residue N added to soils (including N-fixing crops). The percentage of N input for anthropogenic practices is divided by the total N input, and this proportion is used to determine the amount of N<sub>2</sub>O emissions assigned to each of the practices.<sup>100</sup> For example, if 70 percent of the mineral N made available in the soil is due to mineral fertilization, then 70 percent of the N<sub>2</sub>O emissions are assigned to this practice. The remainder of soil N<sub>2</sub>O emissions is reported under “other N inputs,” which includes mineralization due to decomposition of soil organic matter and litter, as well as asymbiotic N fixation from the atmosphere. Asymbiotic N fixation by soil bacteria is a minor source of N, typically not exceeding 10 percent of total N inputs to agroecosystems. Mineralization of soil organic matter is a more significant source of N, but is still typically less than half of the amount of N made available in the cropland soils compared to application of synthetic fertilizers and manure amendments, along with symbiotic fixation. Mineralization of soil organic matter accounts for the majority of available N in grassland soils. Accounting for the influence of “other N inputs” is necessary in order to meet the recommendation for reporting all emissions from managed lands (IPCC 2006). While this method allows for attribution of N<sub>2</sub>O emissions to the individual N inputs to the soils, it is important to realize that sources such as synthetic fertilization may have a larger impact on N<sub>2</sub>O emissions than would be suggested by the associated level of N input for this source (Delgado et al. 2009). Further research will be needed to improve upon this attribution method, however. The results associated with subdividing the N<sub>2</sub>O emissions based on N inputs are provided in Table A-233 and Table A-234.

---

<sup>100</sup> This method is a simplification of reality to allow partitioning of N<sub>2</sub>O emissions, as it assumes that all N inputs have an identical chance of being converted to N<sub>2</sub>O. This is unlikely to be the case, but DAYCENT does not track N<sub>2</sub>O emissions by source of mineral N so this approximation is the only approach that can be used currently for partitioning N<sub>2</sub>O emissions by source of N input. Moreover, this approach is similar to the IPCC Tier 1 method (IPCC 2006), which uses the same direct emissions factor for most N sources (e.g., PRP). Further research and model development may allow for other approaches in the future.

**Table A-233: Direct N<sub>2</sub>O Emissions from Cropland Soils (MMT CO<sub>2</sub> Eq.)**

| N Source   | 1990         | 1991         | 1992         | 1993         | 1994         | 1995         | 1996         | 1997         | 1998         | 1999         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Total Mineral Soils</b>                             | <b>168.6</b> | <b>165.5</b> | <b>166.2</b> | <b>167.2</b> | <b>168.0</b> | <b>166.2</b> | <b>168.4</b> | <b>167.2</b> | <b>177.7</b> | <b>169.0</b> | <b>171.1</b> | <b>167.7</b> | <b>168.4</b> | <b>172.4</b> | <b>181.3</b> | <b>171.2</b> | <b>172.2</b> | <b>178.8</b> | <b>172.1</b> | <b>171.8</b> | <b>182.6</b> | <b>183.9</b> | <b>184.9</b> | <b>182.2</b> | <b>182.0</b> |
| <b>Tier 3 Cropland</b>                                 | <b>153.1</b> | <b>150.0</b> | <b>151.0</b> | <b>153.1</b> | <b>149.0</b> | <b>149.5</b> | <b>149.9</b> | <b>149.9</b> | <b>160.0</b> | <b>151.2</b> | <b>155.9</b> | <b>152.9</b> | <b>152.9</b> | <b>155.6</b> | <b>164.9</b> | <b>155.2</b> | <b>155.1</b> | <b>162.1</b> | <b>156.0</b> | <b>156.8</b> | <b>166.5</b> | <b>165.7</b> | <b>165.9</b> | <b>165.2</b> | <b>165.2</b> |
| Synthetic Fertilizer                                   | 53.3         | 53.0         | 55.2         | 53.4         | 54.3         | 50.8         | 53.8         | 53.3         | 53.5         | 50.7         | 55.6         | 52.9         | 54.0         | 53.8         | 56.6         | 54.7         | 54.9         | 58.2         | 55.0         | 53.9         | 51.9         | 51.6         | 51.7         | 51.5         | 51.5         |
| Managed Manure   | 5.0          | 4.9          | 5.3          | 5.1          | 5.0          | 4.5          | 4.9          | 4.9          | 5.1          | 5.8          | 6.0          | 5.6          | 5.7          | 5.6          | 5.8          | 5.6          | 5.6          | 6.1          | 5.8          | 6.2          | 6.5          | 6.4          | 6.4          | 6.4          | 6.4          |
| Residue N <sup>a</sup>                                 | 23.1         | 25.5         | 23.2         | 23.9         | 22.9         | 25.1         | 24.3         | 24.2         | 23.4         | 26.8         | 25.5         | 24.3         | 24.9         | 25.7         | 23.4         | 24.6         | 24.2         | 24.4         | 23.7         | 23.4         | 26.0         | 25.8         | 25.8         | 25.7         | 25.7         |
| Mineralization and Asymbiotic Fixation                 | 71.6         | 66.6         | 67.3         | 70.7         | 66.8         | 69.1         | 67.0         | 67.5         | 78.0         | 67.9         | 68.7         | 70.1         | 68.3         | 70.5         | 79.1         | 70.3         | 70.3         | 73.3         | 71.5         | 73.4         | 82.2         | 81.8         | 82.0         | 81.6         | 81.6         |
| <b>Tier 1 Cropland</b>                                 | <b>15.6</b>  | <b>15.4</b>  | <b>15.1</b>  | <b>14.2</b>  | <b>19.0</b>  | <b>16.6</b>  | <b>18.4</b>  | <b>17.3</b>  | <b>17.8</b>  | <b>17.7</b>  | <b>15.2</b>  | <b>14.8</b>  | <b>15.5</b>  | <b>16.7</b>  | <b>16.4</b>  | <b>16.0</b>  | <b>17.1</b>  | <b>16.7</b>  | <b>16.1</b>  | <b>15.0</b>  | <b>16.1</b>  | <b>18.2</b>  | <b>19.0</b>  | <b>17.0</b>  | <b>16.8</b>  |
| Synthetic Fertilizer                                   | 5.9          | 5.8          | 5.6          | 4.9          | 9.0          | 6.5          | 8.6          | 7.4          | 7.9          | 8.8          | 6.1          | 5.7          | 6.3          | 7.3          | 7.2          | 6.7          | 7.4          | 7.4          | 6.7          | 6.0          | 7.4          | 9.4          | 10.1         | 8.0          | 7.8          |
| Managed Manure and Other Organic Commercial Fertilizer | 6.8          | 7.0          | 6.9          | 6.8          | 7.3          | 7.8          | 7.5          | 7.6          | 7.6          | 6.9          | 7.0          | 7.2          | 7.3          | 7.5          | 7.1          | 7.3          | 7.8          | 7.5          | 7.5          | 7.1          | 6.9          | 7.1          | 7.2          | 7.2          | 7.2          |
| Residue N  | 2.8          | 2.6          | 2.7          | 2.4          | 2.7          | 2.4          | 2.4          | 2.3          | 2.2          | 2.1          | 2.1          | 2.0          | 1.9          | 1.9          | 2.1          | 2.0          | 1.9          | 1.9          | 1.9          | 1.9          | 1.8          | 1.8          | 1.7          | 1.8          | 1.9          |
| <b>Organic Soils</b>                                   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.1</b>   | <b>3.1</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.3</b>   | <b>3.3</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.1</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   |
| <b>Total*</b>  | <b>171.9</b> | <b>168.7</b> | <b>169.3</b> | <b>170.4</b> | <b>171.1</b> | <b>169.4</b> | <b>171.6</b> | <b>170.3</b> | <b>180.9</b> | <b>172.2</b> | <b>174.3</b> | <b>171.0</b> | <b>171.7</b> | <b>175.6</b> | <b>184.5</b> | <b>174.4</b> | <b>175.3</b> | <b>182.0</b> | <b>175.2</b> | <b>174.8</b> | <b>185.7</b> | <b>186.9</b> | <b>187.9</b> | <b>185.2</b> | <b>185.0</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.<sup>a</sup> Residue N inputs include unharvested fixed N from legumes as well as crop residue N.**Table A-234: Direct N<sub>2</sub>O Emissions from Grasslands (MMT CO<sub>2</sub> Eq.)**

| N Source                               | 1990        | 1991        | 1992        | 1993        | 1994        | 1995        | 1996        | 1997        | 1998        | 1999        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <b>Total Mineral Soils</b>             | <b>70.3</b> | <b>70.4</b> | <b>71.4</b> | <b>71.9</b> | <b>69.0</b> | <b>71.5</b> | <b>74.4</b> | <b>73.7</b> | <b>77.8</b> | <b>66.8</b> | <b>68.3</b> | <b>68.2</b> | <b>70.8</b> | <b>71.2</b> | <b>81.3</b> | <b>71.0</b> | <b>74.0</b> | <b>75.6</b> | <b>73.1</b> | <b>75.9</b> | <b>75.5</b> | <b>74.9</b> | <b>74.0</b> | <b>73.3</b> | <b>73.3</b> |
| <b>Tier 3</b>                          | <b>65.6</b> | <b>66.2</b> | <b>66.6</b> | <b>66.2</b> | <b>62.5</b> | <b>64.1</b> | <b>66.5</b> | <b>67.1</b> | <b>72.0</b> | <b>62.1</b> | <b>63.7</b> | <b>64.3</b> | <b>67.1</b> | <b>66.9</b> | <b>77.4</b> | <b>66.4</b> | <b>68.4</b> | <b>70.6</b> | <b>67.8</b> | <b>70.7</b> | <b>70.3</b> | <b>70.5</b> | <b>70.4</b> | <b>70.4</b> | <b>70.4</b> |
| Synthetic Fertilizer                   | 1.1         | 1.1         | 1.1         | 1.2         | 1.1         | 1.1         | 1.1         | 1.1         | 1.4         | 1.1         | 1.2         | 1.2         | 1.3         | 1.3         | 1.5         | 1.3         | 1.4         | 1.3         | 1.3         | 1.4         | 1.3         | 1.2         | 1.2         | 1.2         | 1.4         |
| PRP Manure                             | 8.9         | 8.8         | 8.8         | 8.5         | 8.5         | 8.2         | 8.5         | 8.3         | 9.3         | 8.0         | 8.4         | 8.5         | 8.9         | 8.6         | 9.6         | 8.2         | 8.5         | 8.4         | 8.0         | 8.3         | 7.9         | 8.0         | 8.0         | 8.0         | 8.0         |
| Managed Manure                         | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | +           | 0.2         |
| Residue N <sup>a</sup>                 | 19.7        | 20.1        | 20.3        | 20.5        | 18.3        | 20.3        | 19.8        | 20.4        | 20.2        | 20.3        | 19.3        | 19.8        | 20.1        | 20.2        | 20.9        | 21.0        | 20.3        | 21.5        | 20.7        | 20.3        | 21.8        | 21.7        | 21.7        | 21.7        | 21.6        |
| Mineralization and Asymbiotic Fixation | 35.8        | 36.1        | 36.3        | 35.8        | 34.4        | 34.4        | 36.9        | 37.1        | 41.1        | 32.5        | 34.7        | 34.6        | 36.7        | 36.6        | 45.3        | 35.8        | 38.0        | 39.3        | 37.7        | 40.5        | 39.1        | 39.4        | 39.4        | 39.4        | 39.2        |
| <b>Tier 1</b>                          | <b>4.7</b>  | <b>4.2</b>  | <b>4.8</b>  | <b>5.7</b>  | <b>6.6</b>  | <b>7.4</b>  | <b>7.8</b>  | <b>6.7</b>  | <b>5.8</b>  | <b>4.7</b>  | <b>4.5</b>  | <b>3.9</b>  | <b>3.7</b>  | <b>4.3</b>  | <b>3.8</b>  | <b>4.6</b>  | <b>5.6</b>  | <b>5.0</b>  | <b>5.4</b>  | <b>5.3</b>  | <b>4.4</b>  | <b>3.5</b>  | <b>2.9</b>  | <b>2.9</b>  | <b>2.9</b>  |
| PRP Manure                             | 4.5         | 4.0         | 4.5         | 5.4         | 6.2         | 7.1         | 7.5         | 6.3         | 5.4         | 4.3         | 4.1         | 3.5         | 3.3         | 3.9         | 3.4         | 4.1         | 5.1         | 4.5         | 4.9         | 4.8         | 4.6         | 3.8         | 3.0         | 2.3         | 2.3         |
| Sewage Sludge                          | 0.2         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.3         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.4         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.5         | 0.6         | 0.6         | 0.6         |
| <b>Organic Soils</b>                   | <b>2.9</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>2.9</b>  | <b>2.9</b>  | <b>2.9</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>3.0</b>  | <b>3.0</b>  | <b>2.9</b>  | <b>2.9</b>  | <b>2.9</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>2.8</b>  | <b>2.7</b>  | <b>2.7</b>  | <b>2.7</b>  | <b>2.7</b>  | <b>2.7</b>  | <b>2.7</b>  |
| <b>Total</b>                           | <b>73.2</b> | <b>73.2</b> | <b>74.2</b> | <b>74.7</b> | <b>71.9</b> | <b>74.3</b> | <b>77.2</b> | <b>76.5</b> | <b>80.6</b> | <b>69.6</b> | <b>71.1</b> | <b>71.1</b> | <b>73.8</b> | <b>74.1</b> | <b>84.1</b> | <b>73.9</b> | <b>76.8</b> | <b>78.4</b> | <b>75.9</b> | <b>78.7</b> | <b>78.1</b> | <b>77.6</b> | <b>76.6</b> | <b>76.0</b> | <b>76.0</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.<sup>a</sup> Residue N inputs include unharvested fixed N from legumes as well as crop residue N.

### **Step 2b: Soil N<sub>2</sub>O Emissions from Agricultural Lands on Mineral Soils Approximated with the Tier 1 Approach**

To estimate direct N<sub>2</sub>O emissions from N additions to crops in the Tier 1 method, the amount of N in applied synthetic fertilizer, manure and other commercial organic fertilizers (i.e., dried blood, tankage, compost, and other) is added to N inputs from crop residues, and the resulting annual totals are multiplied by the IPCC default emission factor of 0.01 kg N<sub>2</sub>O-N/kg N (IPCC 2006) (see Table A-233). The uncertainty is determined based on simple error propagation methods (IPCC 2006). The uncertainty in the default emission factor ranges from 0.3–3.0 kg N<sub>2</sub>O-N/kg N (IPCC 2006). For flooded rice soils, the IPCC default emission factor is 0.003 kg N<sub>2</sub>O-N/kg N and the uncertainty range is 0.000–0.006 kg N<sub>2</sub>O-N/kg N (IPCC 2006).<sup>101</sup> Uncertainties in the emission factor and fertilizer additions are combined with uncertainty in the equations used to calculate residue N additions from above- and below-ground biomass dry matter and N concentration to derive overall uncertainty.

The Tier 1 method is also used to estimate emissions from manure N deposited by livestock on federal lands (i.e., PRP manure N), and from sewage sludge application to grasslands. These two sources of N inputs to soils are multiplied by the IPCC (2006) default emission factors (0.01 kg N<sub>2</sub>O-N/kg N for sludge and horse, sheep, and goat manure, and 0.02 kg N<sub>2</sub>O-N/kg N for cattle, swine, and poultry manure) to estimate N<sub>2</sub>O emissions (Table A-234). The uncertainty is determined based on the Tier 1 error propagation methods provided by the IPCC (2006) with uncertainty in the default emission factor ranging from 0.007 to 0.06 kg N<sub>2</sub>O-N/kg N (IPCC 2006).

**Step 2c: Soil CH<sub>4</sub> Emissions from Agricultural Lands Approximated with the Tier 1 Approach** To estimate CH<sub>4</sub> emissions from rice cultivation for the Tier 1 method, an adjusted daily emission factor is calculated using the default baseline emission factor of 1.30 kg CH<sub>4</sub> ha<sup>-1</sup> d<sup>-1</sup> (ranging 0.8–2.2 kg CH<sub>4</sub> ha<sup>-1</sup> d<sup>-1</sup>) multiplied by a scaling factor for the cultivation water regime, pre-cultivation water regime and a scaling factor for organic amendments (IPCC 2006). The water regime during cultivation is continuously flooded for rice production in the United States and so the scaling factor is always 1 (ranging from 0.79 to 1.26). The pre-season water regime varies based on the proportion of land with winter flooding; land that does not have winter flooding is assigned a value of 0.68 (ranging from 0.58 to 0.80) and areas with winter flooding are assigned a value of 1 (ranging from 0.88 to 1.14). Organic amendments are estimated based on the amount of rice straw and multiplied by 1 (ranging 0.97 to 1.04) for straw incorporated greater than 30 days before cultivation, and by 0.29 (0.2 to 0.4) for straw incorporated greater than 30 days before cultivation. The adjusted daily emission factor is multiplied by the cultivation period and harvested area to estimate the total CH<sub>4</sub> emissions. The uncertainty is propagated through the calculation using an Approach 2 method with a Monte Carlo simulation (IPCC 2006), combining uncertainties associated with the calculation of the adjusted daily emission factor and the harvested areas derived from the USDA NRI survey data.

### **Step 2d: Soil Organic C Stock Changes in Agricultural Lands on Mineral Soils Approximated with the Tier 2 Approach**

Mineral soil organic C stock values are derived for crop rotations that were not simulated by DAYCENT and land converted from non-agricultural land uses to cropland or grassland from 1990–2010, based on the land-use and management activity data in conjunction with appropriate reference C stocks, land-use change, management, input, and wetland restoration factors. Each input to the inventory calculations for the Tier 2 approach has some level of uncertainty that is quantified in PDFs, including the land-use and management activity data, reference C stocks, and management factors. A Monte Carlo Analysis is used to quantify uncertainty in soil organic C stock changes for the inventory period based on uncertainty in the inputs. Input values are randomly selected from PDFs in an iterative process to estimate SOC change for 50,000 times and produce a 95 percent confidence interval for the inventory results.

*Derive Mineral Soil Organic C Stock Change Factors:* Stock change factors representative of U.S. conditions are estimated from published studies (Ogle et al. 2003; Ogle et al. 2006). The numerical factors quantify the impact of changing land use and management on SOC storage in mineral soils, including tillage practices, cropping rotation or intensification, and land conversions between cultivated and native conditions (including set-asides in the Conservation Reserve Program). Studies from the United States and Canada are used in this analysis under the assumption that they would best represent management impacts for the Inventory.

---

<sup>101</sup> Due to lack of data, uncertainties in managed manure N production, PRP manure N production, other commercial organic fertilizer amendments, indirect losses of N in the DAYCENT simulations, and sewage sludge amendments to soils are currently treated as certain; these sources of uncertainty will be included in future Inventories.

The IPCC inventory methodology for agricultural soils divides climate into eight distinct zones based upon average annual temperature, average annual precipitation, and the length of the dry season (IPCC 2006) (Table A-235). Seven of these climate zones occur in the conterminous United States and Hawaii (Eve et al. 2001).

**Table A-235: Characteristics of the IPCC Climate Zones that Occur in the United States**


| Climate Zone  | Annual Average Temperature (°C) | Average Annual Precipitation (mm) | Length of Dry Season (months) |
|---|---------------------------------|-----------------------------------|-------------------------------|
| Cold Temperate, Dry                                   | < 10                            | < Potential Evapotranspiration    | NA                            |
| Cold Temperate, Moist                                 | < 10                            | ≥ Potential Evapotranspiration    | NA                            |
| Warm Temperate, Dry                                   | 10 – 20                         | < 600                             | NA                            |
| Warm Temperate, Moist                                 | 10 – 20                         | ≥ Potential Evapotranspiration    | NA                            |
| Sub-Tropical, Dry <sup>a</sup>                        | > 20                            | < 1,000                           | Usually long                  |
| Sub-Tropical, Moist (w/short dry season) <sup>a</sup> | > 20                            | 1,000 – 2,000                     | < 5                           |

<sup>a</sup> The climate characteristics listed in the table for these zones are those that correspond to the tropical dry and tropical moist zones of the IPCC. They have been renamed “sub-tropical” here.

Mean precipitation and temperature (1950-2000) variables from the WorldClim data set (Hijmans et al. 2005)) and potential evapotranspiration data from the Consortium for Spatial Information (CGIAR-CSI) (Zomer et al. 2008; Zomer et al. 2007) are used to classify climate zones (Figure A-15). IPCC climate zones are assigned to NRI point locations.

Soils are classified into one of seven classes based upon texture, morphology, and ability to store organic matter (IPCC 2006). Six of the categories are mineral types and one is organic (i.e., *Histosol*). Reference C stocks, representing estimates from conventionally managed cropland, are computed for each of the mineral soil types across the various climate zones, based on pedon (i.e., soil) data from the National Soil Survey Characterization Database (NRCS 1997) (Table A-236). These stocks are used in conjunction with management factors to estimate the change in SOC stocks that result from management and land-use activity. PDFs, which represent the variability in the stock estimates, are constructed as normal densities based on the mean and variance from the pedon data. Pedon locations are clumped in various parts of the country, which reduces the statistical independence of individual pedon estimates. To account for this lack of independence, samples from each climate by soil zone are tested for spatial autocorrelation using the Moran’s I test, and variance terms are inflated by 10 percent for all zones with significant p-values.

**Figure A-15: IPCC Climate Zones**



**Table A-236: U.S. Soil Groupings Based on the IPCC Categories and Dominant Taxonomic Soil, and Reference Carbon Stocks (Metric Tons C/ha)**

| IPCC Inventory Soil Categories   | USDA Taxonomic Soil Orders  | Reference Carbon Stock in Climate Regions |                       |                     |                       |                   |                     |
|----------------------------------|---|---|-----------------------|---------------------|-----------------------|-------------------|---------------------|
|                                  |   | Cold Temperate, Dry                       | Cold Temperate, Moist | Warm Temperate, Dry | Warm Temperate, Moist | Sub-Tropical, Dry | Sub-Tropical, Moist |
| High Clay Activity Mineral Soils | Vertisols, Mollisols, Inceptisols, Aridisols, and high base status Alfisols               | 42 (n = 133)                              | 65 (n = 526)          | 37 (n = 203)        | 51 (n = 424)          | 42 (n = 26)       | 57 (n = 12)         |
| Low Clay Activity Mineral Soils  | Ultisols, Oxisols, acidic Alfisols, and many Entisols                                     | 45 (n = 37)                               | 52 (n = 113)          | 25 (n = 86)         | 40 (n = 300)          | 39 (n = 13)       | 47 (n = 7)          |
| Sandy Soils                      | Any soils with greater than 70 percent sand and less than 8 percent clay (often Entisols) | 24 (n = 5)                                | 40 (n = 43)           | 16 (n = 19)         | 30 (n = 102)          | 33 (n = 186)      | 50 (n = 18)         |
| Volcanic Soils                   | Andisols  | 124 (n = 12)                              | 114 (n = 2)           | 124 (n = 12)        | 124 (n = 12)          | 124 (n = 12)      | 128 (n = 9)         |
| Spodic Soils                     | Spodosols   | 86 (n=20)                                 | 74 (n = 13)           | 86 (n=20)           | 107 (n = 7)           | 86 (n=20)         | 86 (n=20)           |
| Aquic Soils                      | Soils with Aquic suborder   | 86 (n = 4)                                | 89 (n = 161)          | 48 (n = 26)         | 51 (n = 300)          | 63 (n = 503)      | 48 (n = 12)         |
| Organic Soils <sup>a</sup>       | Histosols   | NA  | NA                    | NA                  | NA                    | NA                | NA                  |

<sup>a</sup> C stocks are not needed for organic soils.

Notes: C stocks are for the top 30 cm of the soil profile, and are estimated from pedon data available in the National Soil Survey Characterization database (NRCS 1997); sample size provided in parentheses (i.e., 'n' values refer to sample size).

To estimate the land use, management and input factors, studies had to report SOC stocks (or information to compute stocks), depth of sampling, and the number of years since a management change to be included in the analysis. The data are analyzed using linear mixed-effect modeling, accounting for both fixed and random effects. Fixed effects included depth, number of years since a management change, climate, and the type of management change (e.g., reduced tillage vs. no-till). For depth increments, the data are not aggregated for the C stock measurements; each depth increment (e.g., 0-5 cm, 5-10 cm, and 10-30 cm) is included as a separate point in the dataset. Similarly, time series data are not aggregated in these datasets. Linear regression models assume that the underlying data are independent observations, but this is not the case with data from the same experimental site, or plot in a time series. These data are more related to each other than data from other sites (i.e., not independent). Consequently, random effects are needed to account for the dependence in time series data and the dependence among data points representing different depth increments from the same study. Factors are estimated for the effect of management practices at 20 years for the top 30 cm of the soil (Table A-237). Variance is calculated for each of the U.S. factor values, and used to construct PDFs with a normal density. In the IPCC method, specific factor values are given for improved grassland, high input cropland with organic amendments, and for wetland rice, each of which influences C stock changes in soils. Specifically, higher stocks are associated with increased productivity and C inputs (relative to native grassland) on improved grassland with both medium and high input.<sup>102</sup> Organic amendments in annual cropping systems also increase SOC stocks due to greater C inputs, while high SOC stocks in rice cultivation are associated with reduced decomposition due to periodic flooding. There are insufficient field studies to derive factor values for these systems from the published literature, and, thus, estimates from IPCC (2006) are used under the assumption that they would best approximate the impacts, given the lack of sufficient data to derive U.S.-specific factors. A measure of uncertainty is provided for these factors in IPCC (2006), which is used to construct PDFs.

<sup>102</sup> Improved grasslands are identified in the 2010 *National Resources Inventory* as grasslands that are irrigated or seeded with legumes, in addition to those reclassified as improved with manure amendments.

**Table A-237: Soil Organic Carbon Stock Change Factors for the United States and the IPCC Default Values Associated with Management Impacts on Mineral Soils**

|   | IPCC default | Warm Moist Climate | U.S. Factor      |                    |                  |
|---|--------------|--------------------|------------------|--------------------|------------------|
|   |              |                    | Warm Dry Climate | Cool Moist Climate | Cool Dry Climate |
| <b>Land-Use Change Factors</b>                    |              |                    |                  |                    |                  |
| Cultivated <sup>a</sup>                           | 1            | 1                  | 1                | 1                  | 1                |
| General Uncult. <sup>a,b</sup> (n=251)            | 1.4          | 1.42±0.06          | 1.37±0.05        | 1.24±0.06          | 1.20±0.06        |
| Set-Aside <sup>a</sup> (n=142)                    | 1.25         | 1.31±0.06          | 1.26±0.04        | 1.14±0.06          | 1.10±0.05        |
| <b>Improved Grassland Factors</b>                 |              |                    |                  |                    |                  |
| Medium Input                                      | 1.1          | 1.14±0.06          | 1.14±0.06        | 1.14±0.06          | 1.14±0.06        |
| High Input  | NA           | 1.11±0.04          | 1.11±0.04        | 1.11±0.04          | 1.11±0.04        |
| <b>Wetland Rice Production Factor<sup>b</sup></b> | 1.1          | 1.1                | 1.1              | 1.1                | 1.1              |
| <b>Tillage Factors</b>                            |              |                    |                  |                    |                  |
| Conv. Till  | 1            | 1                  | 1                | 1                  | 1                |
| Red. Till (n=93)                                  | 1.05         | 1.08±0.03          | 1.01±0.03        | 1.08±0.03          | 1.01±0.03        |
| No-till (n=212)                                   | 1.1          | 1.13±0.02          | 1.05±0.03        | 1.13±0.02          | 1.05±0.03        |
| <b>Cropland Input Factors</b>                     |              |                    |                  |                    |                  |
| Low (n=85)  | 0.9          | 0.94±0.01          | 0.94±0.01        | 0.94±0.01          | 0.94±0.01        |
| Medium  | 1            | 1                  | 1                | 1                  | 1                |
| High (n=22)                                       | 1.1          | 1.07±0.02          | 1.07±0.02        | 1.07±0.02          | 1.07±0.02        |
| High with amendment <sup>b</sup>                  | 1.2          | 1.38±0.06          | 1.34±0.08        | 1.38±0.06          | 1.34±0.08        |

<sup>a</sup> Factors in the IPCC documentation (IPCC 2006) are converted to represent changes in SOC storage from a cultivated condition rather than a native condition.

<sup>b</sup> U.S.-specific factors are not estimated for land improvements, rice production, or high input with amendment because of few studies addressing the impact of legume mixtures, irrigation, or manure applications for crop and grassland in the United States, or the impact of wetland rice production in the US. Factors provided in IPCC (2006) are used as the best estimates of these impacts.

Note: The "n" values refer to sample size.

Wetland restoration management also influences SOC storage in mineral soils, because restoration leads to higher water tables and inundation of the soil for at least part of the year. A stock change factor is estimated assessing the difference in SOC storage between restored and unrestored wetlands enrolled in the Conservation Reserve Program (Euliss and Gleason 2002), which represents an initial increase of C in the restored soils over the first 10 years (Table A-238). A PDF with a normal density is constructed from these data based on results from a linear regression model. Following the initial increase of C, natural erosion and deposition leads to additional accretion of C in these wetlands. The mass accumulation rate of organic C is estimated using annual sedimentation rates (cm/yr) in combination with percent organic C, and soil bulk density (g/cm<sup>3</sup>) (Euliss and Gleason 2002). Procedures for calculation of mass accumulation rate are described in Dean and Gorham (1998); the resulting rate and standard deviation are used to construct a PDF with a normal density (Table A-238).

**Table A-238: Rate and standard deviation for the Initial Increase and Subsequent Annual Mass Accumulation Rate (Mg C/ha-yr) in Soil Organic C Following Wetland Restoration of Conservation Reserve Program**

| Variable                                   | Value     |
|--|-----------|
| Factor (Initial Increase—First 10 Years)   | 1.22±0.18 |
| Mass Accumulation (After Initial 10 Years) | 0.79±0.05 |

Note: Mass accumulation rate represents additional gains in C for mineral soils after the first 10 years (Euliss and Gleason 2002).

*Estimate Annual Changes in Mineral Soil Organic C Stocks:* In accordance with IPCC methodology, annual changes in mineral soil C are calculated by subtracting the beginning stock from the ending stock and then dividing by 20.<sup>103</sup> For this analysis, stocks are estimated for each year and difference between years is the stock change. From the final distribution of 50,000 values, a 95 percent confidence interval is generated based on the simulated values at the 2.5 and 97.5 percentiles in the distribution (Ogle et al. 2003). Soil organic C stock changes are provided in Table A-239 through Table A-244.

<sup>103</sup> The difference in C stocks is divided by 20 because the stock change factors represent change over a 20-year time period.

### **Step 2e: Estimate Additional Changes in Soil Organic C Stocks Due to CRP Enrollment after 2010 and Sewage Sludge Amendments**

There are two additional land use and management activities in U.S. agricultural lands that are not estimated in Steps 2a and 2b. The first activity involves the application of sewage sludge to agricultural lands. Minimal data exist on where and how much sewage sludge is applied to U.S. agricultural soils, but national estimates of mineral soil land area receiving sewage sludge can be approximated based on sewage sludge N production data, and the assumption that amendments are applied at a rate equivalent to the assimilative capacity from Kellogg et al. (2000). It is assumed that sewage sludge for agricultural land application is applied to grassland because of the high heavy metal content and other pollutants found in human waste, which limits its application to crops. The impact of organic amendments on SOC is calculated as 0.38 metric tonnes C/ha-yr. This rate is based on the IPCC default method and country-specific factors (see Table A-237), by calculating the effect of converting nominal, medium-input grassland to high input improved grassland. The assumptions are that reference C stock are 50 metric tonnes C/ha, which represents a mid-range value of reference C stocks for the cropland soils in the United States,<sup>104</sup> that the land use factor for grassland of 1.4 and 1.11 for high input improved grassland are representative of typical conditions, and that the change in stocks are occurring over a 20 year (default value) time period (i.e.,  $[50 \times 1.4 \times 1.11 - 50 \times 1.4] / 20 = 0.38$ ). A nominal  $\pm 50$  percent uncertainty is attached to these estimates due to limited information on application and the rate of change in soil C stock change with sewage sludge amendments. The influence of sewage sludge on soil organic C stocks are provided in Table A-245.

The second activity is the change in enrollment for the Conservation Reserve Program after 2010 for mineral soils. Relative to the enrollment in 2010, the total area in the Conservation Reserve Program has decreased from 2011 to 2014 (USDA-FSA 2014). An average annual change in SOC of 0.5 metric tonnes C/ha-yr is used to estimate the effect of the enrollment changes. This rate is based on the IPCC default method and country-specific factors (see Table A-237) by estimating the impact of setting aside a medium input cropping system in the Conservation Reserve Program. The assumptions are that reference C stock are 50 metric tonnes C/ha, which represents a mid-range value for the dominant cropland soils in the United States, and the average country-specific factor is 1.2 for setting-aside cropland from production, with the change in stocks occurring over a 20 year (default value) time period equal to 0.5 (i.e.,  $[50 \times 1.2 - 50] / 20 = 0.5$ ). A nominal  $\pm 50$  percent uncertainty is attached to these estimates due to limited information about the enrollment trends at subregional scales, which creates uncertainty in the rate of soil C stock change (stock change factors for set-aside lands vary by climate region). Estimates are provided in Table A-252.

---

<sup>104</sup> Reference C stocks are based on cropland soils for the Tier 2 method applied in this Inventory.

**Table A-239: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Non-Federal Cropland Land Base Estimated with the Tier 2 Analysis using U.S. Factor Values (MMT CO<sub>2</sub> Eq./yr)**

| Non-Federal Croplands: | Cropland Remaining Cropland |                   | Grassland Converted to Cropland |                | Forest Converted to Cropland |                | Other Land Converted to Cropland |                | Settlements Converted to Cropland |                | Wetlands Converted to Cropland |                |
|------------------------|-----------------------------|-------------------|---------------------------------|----------------|------------------------------|----------------|----------------------------------|----------------|-----------------------------------|----------------|--------------------------------|----------------|
|                        | Year                        | Estimate          | 95% CI                          | Estimate       | 95% CI                       | Estimate       | 95% CI                           | Estimate       | 95% CI                            | Estimate       | 95% CI                         | Estimate       |
| <b>Mineral Soils</b>   |                             |                   |                                 |                |                              |                |                                  |                |                                   |                |                                |                |
| 1990                   | -6.13                       | (8.66) to (3.71)  | 1.31                            | (0.43) to 2.22 | 0.20                         | (0.07) to 0.34 | 0.16                             | (0.05) to 0.27 | 0.06                              | (0.02) to 0.11 | 0.11                           | (0.04) to 0.18 |
| 1991                   | -6.69                       | (9.39) to (4.11)  | 1.25                            | (0.43) to 2.23 | 0.18                         | (0.06) to 0.32 | 0.15                             | (0.05) to 0.26 | 0.06                              | (0.02) to 0.11 | 0.10                           | (0.03) to 0.18 |
| 1992                   | -6.69                       | (10.35) to (4.40) | 1.31                            | (0.42) to 2.32 | 0.18                         | (0.06) to 0.31 | 0.16                             | (0.05) to 0.28 | 0.07                              | (0.02) to 0.12 | 0.10                           | (0.03) to 0.18 |
| 1993                   | -7.05                       | (10.12) to (4.05) | 1.44                            | (0.41) to 2.50 | 0.17                         | (0.05) to 0.30 | 0.18                             | (0.05) to 0.32 | 0.07                              | (0.02) to 0.12 | 0.13                           | (0.04) to 0.22 |
| 1994                   | -6.91                       | (9.98) to (3.91)  | 1.44                            | (0.38) to 2.56 | 0.16                         | (0.04) to 0.28 | 0.20                             | (0.05) to 0.35 | 0.07                              | (0.02) to 0.13 | 0.14                           | (0.04) to 0.25 |
| 1995                   | -6.39                       | (9.26) to (3.59)  | 1.55                            | (0.38) to 2.68 | 0.17                         | (0.04) to 0.30 | 0.21                             | (0.05) to 0.36 | 0.07                              | (0.02) to 0.13 | 0.15                           | (0.04) to 0.27 |
| 1996                   | -6.04                       | (8.84) to (3.32)  | 1.77                            | (0.37) to 2.97 | 0.18                         | (0.04) to 0.30 | 0.24                             | (0.05) to 0.40 | 0.08                              | (0.02) to 0.14 | 0.17                           | (0.04) to 0.29 |
| 1997                   | -7.50                       | (11.01) to (4.13) | 1.46                            | (0.32) to 2.65 | 0.14                         | (0.03) to 0.26 | 0.20                             | (0.04) to 0.36 | 0.06                              | (0.01) to 0.12 | 0.14                           | (0.03) to 0.25 |
| 1998                   | -7.20                       | (10.72) to (3.83) | 1.50                            | (0.35) to 2.78 | 0.13                         | (0.03) to 0.24 | 0.20                             | (0.05) to 0.37 | 0.07                              | (0.02) to 0.13 | 0.14                           | (0.03) to 0.26 |
| 1999                   | -6.53                       | (9.82) to (3.38)  | 1.51                            | (0.27) to 2.80 | 0.12                         | (0.02) to 0.22 | 0.21                             | (0.04) to 0.39 | 0.07                              | (0.01) to 0.13 | 0.14                           | (0.02) to 0.26 |
| 2000                   | -6.96                       | (10.25) to (3.85) | 1.41                            | (0.27) to 2.63 | 0.10                         | (0.02) to 0.18 | 0.21                             | (0.04) to 0.39 | 0.07                              | (0.01) to 0.13 | 0.13                           | (0.03) to 0.25 |
| 2001                   | -6.78                       | (9.99) to (3.77)  | 1.32                            | (0.26) to 2.54 | 0.08                         | (0.02) to 0.16 | 0.19                             | (0.04) to 0.37 | 0.06                              | (0.01) to 0.12 | 0.12                           | (0.02) to 0.24 |
| 2002                   | -6.68                       | (9.74) to (3.85)  | 1.12                            | (0.24) to 2.25 | 0.07                         | (0.01) to 0.13 | 0.16                             | (0.03) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.10                           | (0.02) to 0.21 |
| 2003                   | -6.04                       | (8.92) to (3.36)  | 1.02                            | (0.23) to 2.07 | 0.06                         | (0.01) to 0.12 | 0.15                             | (0.03) to 0.30 | 0.05                              | (0.01) to 0.09 | 0.09                           | (0.02) to 0.19 |
| 2004                   | -5.43                       | (8.35) to (2.73)  | 1.14                            | (0.23) to 2.29 | 0.06                         | (0.01) to 0.12 | 0.16                             | (0.03) to 0.32 | 0.05                              | (0.01) to 0.10 | 0.10                           | (0.02) to 0.20 |
| 2005                   | -5.51                       | (8.26) to (2.97)  | 1.11                            | (0.24) to 2.24 | 0.06                         | (0.01) to 0.11 | 0.15                             | (0.03) to 0.30 | 0.05                              | (0.01) to 0.10 | 0.09                           | (0.02) to 0.19 |
| 2006                   | -4.58                       | (7.18) to (2.24)  | 1.34                            | (0.23) to 2.41 | 0.06                         | (0.01) to 0.12 | 0.19                             | (0.03) to 0.34 | 0.06                              | (0.01) to 0.11 | 0.11                           | (0.02) to 0.20 |
| 2007                   | -4.49                       | (6.96) to (2.29)  | 1.38                            | (0.21) to 2.41 | 0.04                         | (0.01) to 0.08 | 0.19                             | (0.03) to 0.33 | 0.07                              | (0.01) to 0.12 | 0.11                           | (0.02) to 0.19 |
| 2008                   | -3.85                       | (6.11) to (1.86)  | 1.37                            | (0.20) to 2.34 | 0.04                         | (0.01) to 0.08 | 0.19                             | (0.03) to 0.33 | 0.06                              | (0.01) to 0.11 | 0.10                           | (0.01) to 0.17 |
| 2009                   | -3.85                       | (6.09) to (1.89)  | 1.23                            | (0.18) to 2.18 | 0.04                         | (0.01) to 0.07 | 0.18                             | (0.03) to 0.31 | 0.05                              | (0.01) to 0.09 | 0.08                           | (0.01) to 0.15 |
| 2010                   | -3.90                       | (6.13) to (1.94)  | 1.24                            | (0.17) to 2.15 | 0.04                         | (0.01) to 0.07 | 0.18                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.08                           | (0.01) to 0.15 |
| 2011                   | -3.60                       | (5.71) to (1.80)  | 1.24                            | (0.17) to 2.16 | 0.04                         | (0.01) to 0.07 | 0.18                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.09                           | (0.01) to 0.15 |
| 2012                   | -3.26                       | (5.31) to (1.53)  | 1.25                            | (0.16) to 2.17 | 0.04                         | (0.01) to 0.07 | 0.18                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.09                           | (0.01) to 0.15 |
| 2013                   | -3.08                       | (5.10) to (1.37)  | 1.26                            | (0.16) to 2.18 | 0.04                         | (0.01) to 0.07 | 0.19                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.09                           | (0.01) to 0.15 |
| 2014                   | -3.11                       | (5.14) to (1.40)  | 1.27                            | (0.16) to 2.18 | 0.04                         | (0.01) to 0.07 | 0.19                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.09                           | (0.01) to 0.15 |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

**Table A-240: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Federal Cropland Land Base Estimated with the Tier 2 Analysis using U.S. Factor Values (MMT CO<sub>2</sub> Eq./yr)**

| Federal Croplands:   | Cropland Remaining Cropland |                  | Grassland Converted to Cropland |              | Forest Converted to Cropland |             | Other Land Converted to Cropland |            | Settlements Converted to Cropland |            | Wetlands Converted to Cropland |            |        |              |
|----------------------|-----------------------------|------------------|---------------------------------|--------------|------------------------------|-------------|----------------------------------|------------|-----------------------------------|------------|--------------------------------|------------|--------|--------------|
|                      | Year                        | Estimate         | 95% CI                          | Estimate     | 95% CI                       | Estimate    | 95% CI                           | Estimate   | 95% CI                            | Estimate   | 95% CI                         | Estimate   | 95% CI |              |
| <b>Mineral Soils</b> |                             |                  |                                 |              |                              |             |                                  |            |                                   |            |                                |            |        |              |
| 1990                 | -0.02                       | (0.06) to 0.01   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1991                 | -0.04                       | (0.09) to 0.01   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1992                 | -0.04                       | (0.15) to 0.03   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1993                 | -0.09                       | (0.19) to 0.01   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1994                 | -0.10                       | (0.23) to 0.03   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1995                 | -0.10                       | (0.23) to 0.03   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1996                 | -0.10                       | (0.23) to 0.03   | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1997                 | -0.23                       | (0.43) to (0.03) | 0.00                            | 0.0 to 0.0   | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.00   | 0.0 to 0.0   |
| 1998                 | -0.11                       | (0.20) to (0.02) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 1999                 | -0.10                       | (0.19) to (0.03) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2000                 | -0.10                       | (0.18) to (0.03) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2001                 | -0.10                       | (0.18) to (0.03) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2002                 | -0.09                       | (0.17) to (0.03) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2003                 | -0.10                       | (0.18) to (0.03) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2004                 | -0.11                       | (0.18) to (0.04) | 0.05                            | 0.03 to 0.08 | 0.01                         | 0.0 to 0.01 | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2005                 | -0.10                       | (0.17) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2006                 | -0.10                       | (0.17) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2007                 | -0.10                       | (0.17) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2008                 | -0.10                       | (0.17) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2009                 | -0.09                       | (0.15) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2010                 | -0.09                       | (0.14) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2011                 | -0.07                       | (0.12) to (0.04) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2012                 | -0.06                       | (0.10) to (0.03) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2013                 | -0.05                       | (0.08) to (0.03) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |
| 2014                 | -0.06                       | (0.09) to (0.03) | 0.06                            | 0.03 to 0.09 | 0.00                         | 0.0 to 0.0  | 0.00                             | 0.0 to 0.0 | 0.00                              | 0.0 to 0.0 | 0.00                           | 0.0 to 0.0 | 0.05   | 0.02 to 0.07 |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

**Table A-241: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Total Cropland Land Base Estimated with the Tier 2 Analysis using U.S. Factor Values (MMT CO<sub>2</sub> Eq./yr)**

| Total Croplands:     | Cropland Remaining Cropland |                   | Grassland Converted to Cropland |                | Forest Converted to Cropland |                | Other Land Converted to Cropland |                | Settlements Converted to Cropland |                | Wetlands Converted to Cropland |                |
|----------------------|-----------------------------|-------------------|---------------------------------|----------------|------------------------------|----------------|----------------------------------|----------------|-----------------------------------|----------------|--------------------------------|----------------|
|                      | Year                        | Estimate          | 95% CI                          | Estimate       | 95% CI                       | Estimate       | 95% CI                           | Estimate       | 95% CI                            | Estimate       | 95% CI                         | Estimate       |
| <b>Mineral Soils</b> |                             |                   |                                 |                |                              |                |                                  |                |                                   |                |                                |                |
| 1990                 | -6.15                       | (8.68) to (3.74)  | 1.31                            | (0.43) to 2.22 | 0.20                         | (0.07) to 0.34 | 0.16                             | (0.05) to 0.27 | 0.06                              | (0.02) to 0.11 | 0.11                           | (0.04) to 0.18 |
| 1991                 | -6.73                       | (9.43) to (4.14)  | 1.25                            | (0.43) to 2.23 | 0.18                         | (0.06) to 0.32 | 0.15                             | (0.05) to 0.26 | 0.06                              | (0.02) to 0.11 | 0.10                           | (0.03) to 0.18 |
| 1992                 | -6.73                       | (10.41) to (4.46) | 1.31                            | (0.42) to 2.32 | 0.18                         | (0.06) to 0.31 | 0.16                             | (0.05) to 0.28 | 0.07                              | (0.02) to 0.12 | 0.10                           | (0.03) to 0.18 |
| 1993                 | -7.14                       | (10.21) to (4.14) | 1.44                            | (0.41) to 2.50 | 0.17                         | (0.05) to 0.30 | 0.18                             | (0.05) to 0.32 | 0.07                              | (0.02) to 0.12 | 0.13                           | (0.04) to 0.22 |
| 1994                 | -7.01                       | (10.08) to (4.01) | 1.44                            | (0.38) to 2.56 | 0.16                         | (0.04) to 0.28 | 0.20                             | (0.05) to 0.35 | 0.07                              | (0.02) to 0.13 | 0.14                           | (0.04) to 0.25 |
| 1995                 | -6.49                       | (9.36) to (3.68)  | 1.55                            | (0.38) to 2.68 | 0.17                         | (0.04) to 0.30 | 0.21                             | (0.05) to 0.36 | 0.07                              | (0.02) to 0.13 | 0.15                           | (0.04) to 0.27 |
| 1996                 | -6.14                       | (8.94) to (3.41)  | 1.77                            | (0.37) to 2.97 | 0.18                         | (0.04) to 0.30 | 0.24                             | (0.05) to 0.40 | 0.08                              | (0.02) to 0.14 | 0.17                           | (0.04) to 0.29 |
| 1997                 | -7.72                       | (11.24) to (4.35) | 1.46                            | (0.32) to 2.65 | 0.14                         | (0.03) to 0.26 | 0.20                             | (0.04) to 0.36 | 0.06                              | (0.01) to 0.12 | 0.14                           | (0.03) to 0.25 |
| 1998                 | -7.30                       | (10.83) to (3.93) | 1.55                            | (0.30) to 2.83 | 0.14                         | (0.02) to 0.25 | 0.20                             | (0.05) to 0.37 | 0.07                              | (0.02) to 0.13 | 0.19                           | 0.01 to 0.31   |
| 1999                 | -6.63                       | (9.92) to (3.49)  | 1.56                            | (0.22) to 2.85 | 0.12                         | (0.01) to 0.22 | 0.21                             | (0.04) to 0.39 | 0.07                              | (0.01) to 0.13 | 0.19                           | 0.02 to 0.31   |
| 2000                 | -7.06                       | (10.35) to (3.95) | 1.46                            | (0.22) to 2.69 | 0.10                         | (0.01) to 0.19 | 0.21                             | (0.04) to 0.39 | 0.07                              | (0.01) to 0.13 | 0.18                           | 0.02 to 0.30   |
| 2001                 | -6.88                       | (10.09) to (3.87) | 1.37                            | (0.20) to 2.60 | 0.09                         | (0.01) to 0.17 | 0.19                             | (0.04) to 0.38 | 0.06                              | (0.01) to 0.12 | 0.17                           | 0.02 to 0.29   |
| 2002                 | -6.78                       | (9.84) to (3.94)  | 1.17                            | (0.19) to 2.30 | 0.07                         | (0.01) to 0.14 | 0.16                             | (0.03) to 0.32 | 0.06                              | (0.01) to 0.12 | 0.15                           | 0.02 to 0.26   |
| 2003                 | -6.15                       | (9.02) to (3.46)  | 1.07                            | (0.18) to 2.12 | 0.07                         | (0.01) to 0.13 | 0.15                             | (0.03) to 0.30 | 0.05                              | (0.01) to 0.09 | 0.14                           | 0.02 to 0.24   |
| 2004                 | -5.53                       | (8.46) to (2.83)  | 1.19                            | (0.17) to 2.34 | 0.07                         | (0.01) to 0.13 | 0.16                             | (0.03) to 0.32 | 0.05                              | (0.01) to 0.10 | 0.15                           | 0.03 to 0.25   |
| 2005                 | -5.61                       | (8.37) to (3.07)  | 1.17                            | (0.18) to 2.29 | 0.06                         | (0.01) to 0.11 | 0.15                             | (0.03) to 0.30 | 0.05                              | (0.01) to 0.10 | 0.14                           | 0.02 to 0.24   |
| 2006                 | -4.68                       | (7.28) to (2.34)  | 1.40                            | (0.18) to 2.46 | 0.07                         | (0.01) to 0.12 | 0.19                             | (0.03) to 0.34 | 0.06                              | (0.01) to 0.11 | 0.16                           | 0.02 to 0.25   |
| 2007                 | -4.59                       | (7.06) to (2.39)  | 1.44                            | (0.16) to 2.46 | 0.04                         | (0.01) to 0.08 | 0.19                             | (0.03) to 0.33 | 0.07                              | (0.01) to 0.12 | 0.15                           | 0.03 to 0.24   |
| 2008                 | -3.95                       | (6.21) to (1.96)  | 1.42                            | (0.14) to 2.39 | 0.04                         | (0.01) to 0.08 | 0.19                             | (0.03) to 0.33 | 0.06                              | (0.01) to 0.11 | 0.15                           | 0.03 to 0.22   |
| 2009                 | -3.95                       | (6.18) to (1.98)  | 1.29                            | (0.13) to 2.24 | 0.04                         | (0.01) to 0.07 | 0.18                             | (0.03) to 0.31 | 0.05                              | (0.01) to 0.09 | 0.13                           | 0.03 to 0.20   |
| 2010                 | -3.98                       | (6.22) to (2.02)  | 1.29                            | (0.11) to 2.21 | 0.04                         | 0.0 to 0.07    | 0.18                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.13                           | 0.03 to 0.20   |
| 2011                 | -3.68                       | (5.78) to (1.87)  | 1.30                            | (0.11) to 2.22 | 0.04                         | 0.0 to 0.07    | 0.18                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.13                           | 0.03 to 0.20   |
| 2012                 | -3.32                       | (5.37) to (1.59)  | 1.31                            | (0.11) to 2.23 | 0.04                         | 0.0 to 0.07    | 0.19                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.13                           | 0.03 to 0.20   |
| 2013                 | -3.13                       | (5.16) to (1.42)  | 1.32                            | (0.10) to 2.23 | 0.04                         | 0.0 to 0.07    | 0.19                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.13                           | 0.03 to 0.20   |
| 2014                 | -3.17                       | (5.20) to (1.45)  | 1.32                            | (0.10) to 2.24 | 0.04                         | 0.0 to 0.07    | 0.19                             | (0.02) to 0.32 | 0.06                              | (0.01) to 0.11 | 0.13                           | 0.03 to 0.20   |
| <b>Organic Soils</b> |                             |                   |                                 |                |                              |                |                                  |                |                                   |                |                                |                |
| 1990                 | 18.41                       | 40.41 to 3.19     | 1.90                            | 4.94 to 0.10   | 0.04                         | 0.18 to 0.11   | 0.01                             | 0.25 to 0.02   | 0.00                              | 0.06 to 0.61   | 0.30                           | 1.04 to 6.09   |
| 1991                 | 18.07                       | 39.89 to 3.20     | 1.91                            | 4.91 to 0.10   | 0.04                         | 0.18 to 0.11   | 0.01                             | 0.25 to 0.02   | 0.00                              | 0.06 to 0.62   | 0.30                           | 1.04 to 5.99   |
| 1992                 | 18.00                       | 39.96 to 3.18     | 1.90                            | 4.90 to 0.09   | 0.04                         | 0.17 to 0.04   | 0.00                             | 0.14 to 0.02   | 0.00                              | 0.06 to 0.62   | 0.30                           | 1.04 to 5.87   |
| 1993                 | 17.70                       | 39.55 to 3.13     | 1.86                            | 4.82 to 0.09   | 0.04                         | 0.16 to 0.09   | 0.00                             | 0.23 to 0.03   | 0.01                              | 0.07 to 0.74   | 0.39                           | 1.18 to 5.82   |
| 1994                 | 17.68                       | 39.57 to 3.37     | 2.03                            | 5.16 to 0.09   | 0.04                         | 0.16 to 0.09   | 0.00                             | 0.23 to 0.05   | 0.01                              | 0.10 to 0.87   | 0.50                           | 1.34 to 5.74   |
| 1995                 | 17.68                       | 39.81 to 3.71     | 2.20                            | 5.71 to 0.09   | 0.04                         | 0.16 to 0.09   | 0.00                             | 0.23 to 0.04   | 0.01                              | 0.09 to 0.90   | 0.53                           | 1.38 to 5.59   |
| 1996                 | 17.57                       | 39.49 to 3.74     | 2.27                            | 5.69 to 0.09   | 0.04                         | 0.16 to 0.09   | 0.00                             | 0.23 to 0.05   | 0.01                              | 0.10 to 0.91   | 0.54                           | 1.39 to 5.53   |
| 1997                 | 17.47                       | 39.47 to 3.78     | 2.29                            | 5.73 to 0.08   | 0.04                         | 0.15 to 0.09   | 0.00                             | 0.23 to 0.04   | 0.01                              | 0.09 to 0.91   | 0.54                           | 1.39 to 5.40   |
| 1998                 | 17.67                       | 39.35 to 4.28     | 1.68                            | 8.56 to 0.08   | 0.03                         | 0.14 to 0.06   | 0.00                             | 0.20 to 0.05   | 0.01                              | 0.11 to 0.97   | 0.50                           | 1.57 to 5.30   |

|      |       |               |      |               |      |              |      |              |      |              |      |              |
|------|-------|---------------|------|---------------|------|--------------|------|--------------|------|--------------|------|--------------|
| 1999 | 17.18 | 38.38 to 4.40 | 1.78 | 8.72 to 0.07  | 0.03 | 0.13 to 0.06 | 0.00 | 0.20 to 0.05 | 0.01 | 0.11 to 0.94 | 0.47 | 1.53 to 5.18 |
| 2000 | 17.13 | 38.39 to 4.07 | 1.66 | 8.27 to 0.07  | 0.02 | 0.12 to 0.06 | 0.00 | 0.20 to 0.05 | 0.01 | 0.11 to 0.77 | 0.42 | 1.22 to 5.15 |
| 2001 | 17.60 | 39.65 to 5.45 | 1.71 | 12.90 to 0.06 | 0.02 | 0.12 to 0.06 | 0.00 | 0.20 to 0.05 | 0.01 | 0.11 to 0.80 | 0.45 | 1.24 to 4.72 |
| 2002 | 17.84 | 39.86 to 5.29 | 1.62 | 12.67 to 0.05 | 0.01 | 0.10 to 0.06 | 0.00 | 0.20 to 0.06 | 0.01 | 0.12 to 0.70 | 0.37 | 1.12 to 4.64 |
| 2003 | 18.02 | 40.19 to 5.14 | 1.53 | 12.37 to 0.04 | 0.01 | 0.09 to 0.06 | 0.00 | 0.20 to 0.06 | 0.01 | 0.11 to 0.62 | 0.34 | 0.97 to 4.55 |
| 2004 | 17.78 | 43.90 to 4.22 | 1.30 | 10.74 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.05  | 0.01 | 0.09 to 0.51 | 0.23 | 0.88 to 4.49 |
| 2005 | 17.80 | 43.72 to 4.18 | 1.29 | 10.66 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.07  | 0.02 | 0.14 to 0.51 | 0.23 | 0.87 to 4.46 |
| 2006 | 17.66 | 43.26 to 4.16 | 1.24 | 10.72 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.07  | 0.02 | 0.14 to 0.51 | 0.23 | 0.88 to 4.43 |
| 2007 | 17.62 | 40.81 to 4.03 | 1.16 | 10.57 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.07  | 0.02 | 0.14 to 0.51 | 0.23 | 0.87 to 4.43 |
| 2008 | 17.42 | 40.10 to 3.88 | 1.06 | 10.37 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.06  | 0.01 | 0.12 to 0.46 | 0.23 | 0.75 to 4.44 |
| 2009 | 17.69 | 41.11 to 3.68 | 0.87 | 10.06 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.06  | 0.01 | 0.13 to 0.43 | 0.21 | 0.71 to 4.42 |
| 2010 | 17.72 | 40.96 to 3.81 | 0.87 | 10.25 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.07  | 0.02 | 0.13 to 0.43 | 0.21 | 0.71 to 4.38 |
| 2011 | 17.74 | 41.22 to 3.81 | 0.88 | 10.37 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.08  | 0.02 | 0.15 to 0.43 | 0.21 | 0.71 to 4.36 |
| 2012 | 17.69 | 40.91 to 3.78 | 0.88 | 10.20 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.08  | 0.02 | 0.15 to 0.43 | 0.21 | 0.71 to 4.33 |
| 2013 | 17.77 | 40.93 to 3.81 | 0.89 | 10.27 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.08  | 0.02 | 0.15 to 0.43 | 0.21 | 0.71 to 4.33 |
| 2014 | 17.79 | 40.98 to 3.82 | 0.86 | 10.25 to 0.02 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.08  | 0.02 | 0.15 to 0.43 | 0.21 | 0.71 to 4.33 |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

**Table A-242: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Non-Federal Grasslands Land Base Estimated with the Tier 2 Analysis using U.S. Factor Values (MMT CO<sub>2</sub> Eq./yr)**

| Non-Federal Grasslands: | Grassland Remaining Grassland |                | Cropland Converted to Grassland |                  | Forest Converted to Grassland |                  | Other Land Converted to Grassland |                  | Settlements Converted to Grassland |                  | Wetlands Converted to Grassland |                  |
|-------------------------|-------------------------------|----------------|---------------------------------|------------------|-------------------------------|------------------|-----------------------------------|------------------|------------------------------------|------------------|---------------------------------|------------------|
|                         | Year                          | Estimate       | 95% CI                          | Estimate         | 95% CI                        | Estimate         | 95% CI                            | Estimate         | 95% CI                             | Estimate         | 95% CI                          | Estimate         |
| <b>Mineral Soils</b>    |                               |                |                                 |                  |                               |                  |                                   |                  |                                    |                  |                                 |                  |
| 1990                    | -0.27                         | (0.60) to 0.01 | -3.14                           | (4.54) to (1.90) | -0.48                         | (0.69) to (0.29) | -0.62                             | (0.90) to (0.38) | -0.09                              | (0.13) to (0.05) | -0.47                           | (0.68) to (0.28) |
| 1991                    | -0.27                         | (0.60) to 0.0  | -3.07                           | (4.41) to (1.87) | -0.47                         | (0.68) to (0.29) | -0.64                             | (0.93) to (0.39) | -0.09                              | (0.13) to (0.06) | -0.59                           | (0.85) to (0.36) |
| 1992                    | -0.27                         | (0.58) to 0.01 | -2.97                           | (4.27) to (1.80) | -0.45                         | (0.64) to (0.27) | -0.68                             | (0.98) to (0.41) | -0.09                              | (0.14) to (0.06) | -0.61                           | (0.87) to (0.37) |
| 1993                    | -0.26                         | (0.57) to 0.0  | -3.30                           | (4.74) to (1.99) | -0.47                         | (0.67) to (0.28) | -0.83                             | (1.19) to (0.50) | -0.11                              | (0.16) to (0.07) | -0.64                           | (0.92) to (0.39) |
| 1994                    | -0.24                         | (0.53) to 0.0  | -3.41                           | (4.90) to (2.07) | -0.45                         | (0.65) to (0.27) | -0.95                             | (1.37) to (0.58) | -0.12                              | (0.17) to (0.07) | -0.65                           | (0.94) to (0.40) |
| 1995                    | -0.24                         | (0.52) to 0.0  | -3.25                           | (4.68) to (1.97) | -0.47                         | (0.68) to (0.29) | -0.98                             | (1.40) to (0.59) | -0.12                              | (0.17) to (0.07) | -0.62                           | (0.90) to (0.38) |
| 1996                    | -0.24                         | (0.51) to 0.0  | -3.05                           | (4.39) to (1.85) | -0.45                         | (0.65) to (0.27) | -0.94                             | (1.36) to (0.57) | -0.12                              | (0.18) to (0.07) | -0.61                           | (0.88) to (0.37) |
| 1997                    | -0.20                         | (0.44) to 0.0  | -2.90                           | (4.14) to (1.79) | -0.43                         | (0.61) to (0.26) | -0.99                             | (1.40) to (0.61) | -0.13                              | (0.18) to (0.08) | -0.61                           | (0.87) to (0.38) |
| 1998                    | -0.15                         | (0.48) to 0.16 | -3.19                           | (4.64) to (1.93) | -0.41                         | (0.60) to (0.25) | -1.07                             | (1.56) to (0.65) | -0.14                              | (0.21) to (0.09) | -0.60                           | (0.87) to (0.36) |
| 1999                    | -0.08                         | (0.37) to 0.19 | -2.87                           | (4.16) to (1.74) | -0.40                         | (0.57) to (0.24) | -1.07                             | (1.55) to (0.65) | -0.15                              | (0.22) to (0.09) | -0.59                           | (0.86) to (0.36) |
| 2000                    | -0.08                         | (0.37) to 0.20 | -3.06                           | (4.44) to (1.85) | -0.38                         | (0.55) to (0.23) | -1.16                             | (1.68) to (0.70) | -0.16                              | (0.23) to (0.10) | -0.61                           | (0.88) to (0.37) |
| 2001                    | -0.07                         | (0.34) to 0.20 | -2.89                           | (4.20) to (1.74) | -0.36                         | (0.53) to (0.22) | -1.19                             | (1.73) to (0.71) | -0.16                              | (0.24) to (0.10) | -0.59                           | (0.86) to (0.35) |
| 2002                    | -0.05                         | (0.32) to 0.20 | -2.64                           | (3.86) to (1.57) | -0.33                         | (0.49) to (0.20) | -1.12                             | (1.63) to (0.66) | -0.16                              | (0.24) to (0.10) | -0.54                           | (0.80) to (0.32) |
| 2003                    | -0.05                         | (0.31) to 0.21 | -2.40                           | (3.55) to (1.41) | -0.27                         | (0.39) to (0.16) | -1.04                             | (1.54) to (0.61) | -0.15                              | (0.22) to (0.09) | -0.52                           | (0.76) to (0.30) |

|      |       |                |       |                  |       |                  |       |                  |       |                  |       |                  |
|------|-------|----------------|-------|------------------|-------|------------------|-------|------------------|-------|------------------|-------|------------------|
| 2004 | -0.04 | (0.30) to 0.21 | -2.67 | (3.94) to (1.57) | -0.28 | (0.41) to (0.16) | -1.09 | (1.60) to (0.64) | -0.15 | (0.23) to (0.09) | -0.55 | (0.81) to (0.32) |
| 2005 | -0.06 | (0.32) to 0.19 | -2.46 | (3.63) to (1.45) | -0.28 | (0.41) to (0.16) | -1.08 | (1.60) to (0.64) | -0.15 | (0.23) to (0.09) | -0.53 | (0.79) to (0.31) |
| 2006 | -0.06 | (0.31) to 0.18 | -2.05 | (3.10) to (1.15) | -0.24 | (0.36) to (0.13) | -0.95 | (1.44) to (0.53) | -0.14 | (0.21) to (0.08) | -0.47 | (0.71) to (0.26) |
| 2007 | -0.04 | (0.28) to 0.20 | -1.73 | (2.66) to (0.95) | -0.18 | (0.27) to (0.10) | -0.89 | (1.36) to (0.49) | -0.13 | (0.20) to (0.07) | -0.43 | (0.66) to (0.24) |
| 2008 | -0.01 | (0.25) to 0.23 | -1.58 | (2.43) to (0.87) | -0.16 | (0.24) to (0.09) | -0.85 | (1.30) to (0.46) | -0.13 | (0.19) to (0.07) | -0.33 | (0.51) to (0.18) |
| 2009 | 0.00  | (0.24) to 0.24 | -1.52 | (2.34) to (0.83) | -0.16 | (0.24) to (0.09) | -0.84 | (1.29) to (0.46) | -0.12 | (0.19) to (0.07) | -0.27 | (0.41) to (0.15) |
| 2010 | 0.01  | (0.22) to 0.24 | -1.45 | (2.22) to (0.79) | -0.16 | (0.24) to (0.09) | -0.83 | (1.28) to (0.45) | -0.13 | (0.19) to (0.07) | -0.23 | (0.35) to (0.12) |
| 2011 | 0.01  | (0.21) to 0.24 | -1.44 | (2.21) to (0.78) | -0.16 | (0.24) to (0.08) | -0.83 | (1.28) to (0.45) | -0.13 | (0.19) to (0.07) | -0.22 | (0.34) to (0.12) |
| 2012 | 0.02  | (0.21) to 0.24 | -1.43 | (2.20) to (0.76) | -0.15 | (0.24) to (0.08) | -0.82 | (1.27) to (0.44) | -0.12 | (0.19) to (0.07) | -0.22 | (0.34) to (0.12) |
| 2013 | 0.02  | (0.21) to 0.25 | -1.42 | (2.19) to (0.76) | -0.15 | (0.24) to (0.08) | -0.82 | (1.26) to (0.44) | -0.12 | (0.19) to (0.07) | -0.22 | (0.34) to (0.12) |
| 2014 | 0.02  | (0.20) to 0.26 | -1.42 | (2.19) to (0.76) | -0.15 | (0.24) to (0.08) | -0.82 | (1.26) to (0.43) | -0.12 | (0.19) to (0.07) | -0.22 | (0.34) to (0.12) |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

**Table A-243: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Federal Grasslands Land Base Estimated with the Tier 2 Analysis using U.S. Factor Values (MMT CO<sub>2</sub> Eq./yr)**

| Federal Grasslands:  | Grassland Remaining Grassland |                 | Cropland Converted to Grassland |               | Forest Converted to Grassland |                | Other Land Converted to Grassland |                | Settlements Converted to Grassland |            | Wetlands Converted to Grassland |                |        |
|----------------------|-------------------------------|-----------------|---------------------------------|---------------|-------------------------------|----------------|-----------------------------------|----------------|------------------------------------|------------|---------------------------------|----------------|--------|
|                      | Year                          | Estimate        | 95% CI                          | Estimate      | 95% CI                        | Estimate       | 95% CI                            | Estimate       | 95% CI                             | Estimate   | 95% CI                          | Estimate       | 95% CI |
| <b>Mineral Soils</b> |                               |                 |                                 |               |                               |                |                                   |                |                                    |            |                                 |                |        |
| 1990                 | -0.21                         | (8.71) to 9.11  | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1991                 | -0.30                         | (9.01) to 8.48  | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1992                 | -0.30                         | (9.83) to 7.79  | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1993                 | -1.17                         | (10.81) to 7.36 | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1994                 | -1.46                         | (11.45) to 6.94 | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1995                 | -1.46                         | (11.63) to 6.81 | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1996                 | -0.81                         | (10.24) to 6.85 | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1997                 | -0.74                         | (10.05) to 7.06 | 0.00                            | 0.0 to 0.0    | 0.00                          | 0.0 to 0.0     | 0.00                              | 0.0 to 0.0     | 0.00                               | 0.0 to 0.0 | 0.00                            | 0.0 to 0.0     |        |
| 1998                 | -1.42                         | (12.70) to 6.88 | -0.02                           | (0.05) to 0.0 | -0.68                         | (1.55) to 0.13 | -0.04                             | (0.10) to 0.01 | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.01 |        |
| 1999                 | -1.27                         | (12.52) to 6.99 | -0.02                           | (0.05) to 0.0 | -0.69                         | (1.56) to 0.13 | -0.04                             | (0.10) to 0.01 | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.01 |        |
| 2000                 | -1.46                         | (11.84) to 6.24 | -0.02                           | (0.05) to 0.0 | -0.69                         | (1.55) to 0.12 | -0.04                             | (0.10) to 0.01 | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.01 |        |
| 2001                 | -1.45                         | (11.92) to 6.34 | -0.02                           | (0.05) to 0.0 | -0.68                         | (1.55) to 0.12 | -0.04                             | (0.10) to 0.01 | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.01 |        |
| 2002                 | -2.46                         | (13.81) to 6.87 | -0.02                           | (0.05) to 0.0 | -0.70                         | (1.50) to 0.05 | -0.05                             | (0.10) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.0  |        |
| 2003                 | -2.51                         | (13.94) to 7.52 | -0.02                           | (0.05) to 0.0 | -0.69                         | (1.49) to 0.05 | -0.04                             | (0.10) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.0  |        |
| 2004                 | -1.21                         | (10.83) to 8.60 | -0.02                           | (0.05) to 0.0 | -0.69                         | (1.49) to 0.06 | -0.04                             | (0.10) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.04                           | (0.09) to 0.0  |        |
| 2005                 | -1.27                         | (10.93) to 8.31 | 0.00                            | 0.0 to 0.0    | -0.71                         | (1.68) to 0.20 | -0.01                             | (0.02) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.02                           | (0.05) to 0.01 |        |
| 2006                 | -1.39                         | (11.27) to 8.31 | 0.00                            | 0.0 to 0.0    | -0.71                         | (1.68) to 0.19 | -0.01                             | (0.02) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.02                           | (0.05) to 0.01 |        |
| 2007                 | -1.49                         | (11.36) to 7.92 | 0.00                            | 0.0 to 0.0    | -0.71                         | (1.67) to 0.20 | -0.01                             | (0.02) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.02                           | (0.05) to 0.01 |        |
| 2008                 | -1.53                         | (11.51) to 8.04 | 0.00                            | 0.0 to 0.0    | -0.70                         | (1.67) to 0.20 | -0.01                             | (0.02) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.02                           | (0.05) to 0.01 |        |
| 2009                 | -1.30                         | (10.95) to 7.01 | 0.00                            | 0.0 to 0.0    | -0.69                         | (1.66) to 0.21 | -0.01                             | (0.02) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.02                           | (0.05) to 0.01 |        |
| 2010                 | -1.37                         | (10.93) to 7.36 | 0.00                            | 0.0 to 0.0    | -0.68                         | (1.65) to 0.22 | -0.01                             | (0.02) to 0.0  | 0.00                               | 0.0 to 0.0 | -0.02                           | (0.05) to 0.01 |        |

|      |       |                 |      |            |       |                |       |               |      |            |       |                |
|------|-------|-----------------|------|------------|-------|----------------|-------|---------------|------|------------|-------|----------------|
| 2011 | -1.00 | (10.18) to 7.87 | 0.00 | 0.0 to 0.0 | -0.67 | (1.64) to 0.23 | -0.01 | (0.02) to 0.0 | 0.00 | 0.0 to 0.0 | -0.02 | (0.05) to 0.01 |
| 2012 | -0.52 | (9.34) to 8.61  | 0.00 | 0.0 to 0.0 | -0.66 | (1.63) to 0.24 | -0.01 | (0.02) to 0.0 | 0.00 | 0.0 to 0.0 | -0.02 | (0.05) to 0.01 |
| 2013 | -0.28 | (8.76) to 9.05  | 0.00 | 0.0 to 0.0 | -0.66 | (1.63) to 0.25 | -0.01 | (0.02) to 0.0 | 0.00 | 0.0 to 0.0 | -0.02 | (0.05) to 0.01 |
| 2014 | -0.28 | (8.84) to 9.23  | 0.00 | 0.0 to 0.0 | -0.66 | (1.63) to 0.25 | -0.01 | (0.02) to 0.0 | 0.00 | 0.0 to 0.0 | -0.02 | (0.05) to 0.01 |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

**Table A-244: Annual Change in Soil Organic Carbon Stocks (95% Confidence Interval) for the Total Grassland Land Base Estimated with the Tier 2 Analysis using U.S. Factor Values (MMT CO<sub>2</sub> Eq./yr)**

| Total Grasslands:<br>Year | Grassland Remaining<br>Grassland |                 | Cropland Converted to<br>Grassland |                  | Forest Converted to<br>Grassland |                  | Other Land Converted to<br>Grassland |                  | Settlements Converted to<br>Grassland |                  | Wetlands Converted to<br>Grassland |                  |
|---------------------------|----------------------------------|-----------------|------------------------------------|------------------|----------------------------------|------------------|--------------------------------------|------------------|---------------------------------------|------------------|------------------------------------|------------------|
|                           | Estimate                         | 95% CI          | Estimate                           | 95% CI           | Estimate                         | 95% CI           | Estimate                             | 95% CI           | Estimate                              | 95% CI           | Estimate                           | 95% CI           |
| <b>Mineral Soils</b>      |                                  |                 |                                    |                  |                                  |                  |                                      |                  |                                       |                  |                                    |                  |
| 1990                      | -0.48                            | (8.98) to 8.84  | -3.14                              | (4.54) to (1.90) | -0.48                            | (0.69) to (0.29) | -0.62                                | (0.90) to (0.38) | -0.09                                 | (0.13) to (0.05) | -0.47                              | (0.68) to (0.28) |
| 1991                      | -0.58                            | (9.29) to 8.21  | -3.07                              | (4.41) to (1.87) | -0.47                            | (0.68) to (0.29) | -0.64                                | (0.93) to (0.39) | -0.09                                 | (0.13) to (0.06) | -0.59                              | (0.85) to (0.36) |
| 1992                      | -0.58                            | (10.09) to 7.53 | -2.97                              | (4.27) to (1.80) | -0.45                            | (0.64) to (0.27) | -0.68                                | (0.98) to (0.41) | -0.09                                 | (0.14) to (0.06) | -0.61                              | (0.87) to (0.37) |
| 1993                      | -1.42                            | (11.08) to 7.10 | -3.30                              | (4.74) to (1.99) | -0.47                            | (0.67) to (0.28) | -0.83                                | (1.19) to (0.50) | -0.11                                 | (0.16) to (0.07) | -0.64                              | (0.92) to (0.39) |
| 1994                      | -1.70                            | (11.69) to 6.71 | -3.41                              | (4.90) to (2.07) | -0.45                            | (0.65) to (0.27) | -0.95                                | (1.37) to (0.58) | -0.12                                 | (0.17) to (0.07) | -0.65                              | (0.94) to (0.40) |
| 1995                      | -1.70                            | (11.87) to 6.57 | -3.25                              | (4.68) to (1.97) | -0.47                            | (0.68) to (0.29) | -0.98                                | (1.40) to (0.59) | -0.12                                 | (0.17) to (0.07) | -0.62                              | (0.90) to (0.38) |
| 1996                      | -1.04                            | (10.48) to 6.62 | -3.05                              | (4.39) to (1.85) | -0.45                            | (0.65) to (0.27) | -0.94                                | (1.36) to (0.57) | -0.12                                 | (0.18) to (0.07) | -0.61                              | (0.88) to (0.37) |
| 1997                      | -0.94                            | (10.25) to 6.86 | -2.90                              | (4.14) to (1.79) | -0.43                            | (0.61) to (0.26) | -0.99                                | (1.40) to (0.61) | -0.13                                 | (0.18) to (0.08) | -0.61                              | (0.87) to (0.38) |
| 1998                      | -1.57                            | (12.85) to 6.73 | -3.21                              | (4.66) to (1.95) | -1.09                            | (1.98) to (0.26) | -1.12                                | (1.61) to (0.69) | -0.15                                 | (0.21) to (0.09) | -0.64                              | (0.91) to (0.40) |
| 1999                      | -1.35                            | (12.60) to 6.91 | -2.89                              | (4.18) to (1.76) | -1.09                            | (1.97) to (0.26) | -1.12                                | (1.60) to (0.69) | -0.15                                 | (0.22) to (0.09) | -0.63                              | (0.90) to (0.39) |
| 2000                      | -1.54                            | (11.92) to 6.16 | -3.08                              | (4.46) to (1.88) | -1.07                            | (1.95) to (0.25) | -1.20                                | (1.72) to (0.74) | -0.16                                 | (0.23) to (0.10) | -0.65                              | (0.93) to (0.40) |
| 2001                      | -1.52                            | (12.0) to 6.27  | -2.92                              | (4.23) to (1.76) | -1.05                            | (1.92) to (0.23) | -1.23                                | (1.77) to (0.76) | -0.16                                 | (0.24) to (0.10) | -0.63                              | (0.90) to (0.39) |
| 2002                      | -2.52                            | (13.87) to 6.82 | -2.66                              | (3.89) to (1.59) | -1.03                            | (1.85) to (0.27) | -1.16                                | (1.68) to (0.71) | -0.16                                 | (0.24) to (0.10) | -0.59                              | (0.84) to (0.36) |
| 2003                      | -2.56                            | (13.99) to 7.48 | -2.43                              | (3.57) to (1.44) | -0.96                            | (1.77) to (0.20) | -1.09                                | (1.58) to (0.65) | -0.15                                 | (0.22) to (0.09) | -0.56                              | (0.81) to (0.34) |
| 2004                      | -1.25                            | (10.87) to 8.56 | -2.69                              | (3.96) to (1.59) | -0.97                            | (1.78) to (0.21) | -1.13                                | (1.65) to (0.68) | -0.15                                 | (0.23) to (0.09) | -0.59                              | (0.85) to (0.36) |
| 2005                      | -1.33                            | (10.99) to 8.25 | -2.46                              | (3.63) to (1.45) | -0.99                            | (1.97) to (0.07) | -1.09                                | (1.61) to (0.65) | -0.15                                 | (0.23) to (0.09) | -0.55                              | (0.81) to (0.33) |
| 2006                      | -1.45                            | (11.33) to 8.26 | -2.05                              | (3.10) to (1.15) | -0.95                            | (1.93) to (0.04) | -0.96                                | (1.45) to (0.54) | -0.14                                 | (0.21) to (0.08) | -0.49                              | (0.73) to (0.28) |
| 2007                      | -1.53                            | (11.40) to 7.89 | -1.74                              | (2.66) to (0.95) | -0.88                            | (1.86) to 0.03   | -0.90                                | (1.37) to (0.50) | -0.13                                 | (0.20) to (0.07) | -0.45                              | (0.69) to (0.26) |
| 2008                      | -1.54                            | (11.53) to 8.03 | -1.58                              | (2.43) to (0.87) | -0.86                            | (1.83) to 0.05   | -0.86                                | (1.31) to (0.47) | -0.13                                 | (0.19) to (0.07) | -0.35                              | (0.54) to (0.20) |
| 2009                      | -1.30                            | (10.95) to 7.01 | -1.52                              | (2.34) to (0.83) | -0.85                            | (1.82) to 0.06   | -0.85                                | (1.30) to (0.47) | -0.12                                 | (0.19) to (0.07) | -0.29                              | (0.43) to (0.16) |
| 2010                      | -1.36                            | (10.92) to 7.37 | -1.45                              | (2.23) to (0.79) | -0.84                            | (1.81) to 0.07   | -0.84                                | (1.29) to (0.46) | -0.13                                 | (0.19) to (0.07) | -0.24                              | (0.37) to (0.14) |
| 2011                      | -0.98                            | (10.17) to 7.89 | -1.44                              | (2.22) to (0.78) | -0.83                            | (1.80) to 0.08   | -0.84                                | (1.29) to (0.46) | -0.13                                 | (0.19) to (0.07) | -0.24                              | (0.37) to (0.14) |
| 2012                      | -0.51                            | (9.33) to 8.63  | -1.43                              | (2.20) to (0.77) | -0.81                            | (1.78) to 0.09   | -0.83                                | (1.28) to (0.45) | -0.12                                 | (0.19) to (0.07) | -0.24                              | (0.36) to (0.13) |
| 2013                      | -0.26                            | (8.75) to 9.07  | -1.42                              | (2.20) to (0.76) | -0.81                            | (1.78) to 0.10   | -0.83                                | (1.27) to (0.45) | -0.12                                 | (0.19) to (0.07) | -0.24                              | (0.36) to (0.13) |
| 2014                      | -0.26                            | (8.81) to 9.26  | -1.42                              | (2.19) to (0.76) | -0.81                            | (1.78) to 0.10   | -0.83                                | (1.27) to (0.44) | -0.12                                 | (0.19) to (0.07) | -0.24                              | (0.36) to (0.13) |
| <b>Organic Soils</b>      |                                  |                 |                                    |                  |                                  |                  |                                      |                  |                                       |                  |                                    |                  |
| 1990                      | 3.46                             | 9.59 to 0.54    | 0.23                               | .98 to 0.01      | 0.00                             | 0.03 to 0.03     | 0.01                                 | 0.06 to 0.0      | 0.00                                  | 0.0 to 0.08      | 0.01                               | 0.20 to 0.0      |
| 1991                      | 3.39                             | 9.47 to 0.53    | 0.23                               | .97 to 0.01      | 0.00                             | 0.03 to 0.03     | 0.01                                 | 0.06 to 0.0      | 0.00                                  | 0.0 to 0.08      | 0.01                               | 0.20 to 0.0      |
| 1992                      | 3.31                             | 9.25 to 0.53    | 0.23                               | .97 to 0.01      | 0.00                             | 0.03 to 0.02     | 0.00                                 | 0.05 to 0.0      | 0.00                                  | 0.0 to 0.09      | 0.01                               | 0.22 to 0.0      |

|      |      |              |      |              |      |              |      |              |      |              |      |             |
|------|------|--------------|------|--------------|------|--------------|------|--------------|------|--------------|------|-------------|
| 1993 | 3.29 | 9.19 to 0.64 | 0.29 | 1.15 to 0.02 | 0.01 | 0.04 to 0.02 | 0.00 | 0.06 to 0.01 | 0.00 | 0.02 to 0.15 | 0.05 | 0.30 to 0.0 |
| 1994 | 3.25 | 9.07 to 0.73 | 0.32 | 1.32 to 0.02 | 0.01 | 0.04 to 0.02 | 0.00 | 0.05 to 0.01 | 0.00 | 0.02 to 0.18 | 0.07 | 0.32 to 0.0 |
| 1995 | 3.15 | 8.84 to 0.74 | 0.33 | 1.34 to 0.02 | 0.01 | 0.04 to 0.02 | 0.00 | 0.05 to 0.01 | 0.00 | 0.02 to 0.18 | 0.07 | 0.33 to 0.0 |
| 1996 | 3.14 | 8.74 to 0.72 | 0.32 | 1.29 to 0.02 | 0.01 | 0.03 to 0.02 | 0.00 | 0.05 to 0.01 | 0.00 | 0.02 to 0.18 | 0.08 | 0.33 to 0.0 |
| 1997 | 3.06 | 8.53 to 0.70 | 0.31 | 1.27 to 0.02 | 0.01 | 0.03 to 0.01 | 0.00 | 0.04 to 0.01 | 0.00 | 0.02 to 0.18 | 0.08 | 0.34 to 0.0 |
| 1998 | 2.82 | 8.61 to 0.82 | 0.38 | 1.46 to 0.02 | 0.00 | 0.06 to 0.01 | 0.00 | 0.04 to 0.01 | 0.00 | 0.03 to 0.19 | 0.10 | 0.31 to 0.0 |
| 1999 | 2.74 | 8.44 to 0.76 | 0.35 | 1.35 to 0.02 | 0.00 | 0.05 to 0.01 | 0.00 | 0.04 to 0.01 | 0.00 | 0.03 to 0.19 | 0.10 | 0.31 to 0.0 |
| 2000 | 2.72 | 8.43 to 0.82 | 0.39 | 1.44 to 0.02 | 0.00 | 0.06 to 0.01 | 0.00 | 0.04 to 0.01 | 0.00 | 0.03 to 0.19 | 0.10 | 0.31 to 0.0 |
| 2001 | 2.54 | 7.63 to 0.89 | 0.43 | 1.57 to 0.02 | 0.00 | 0.06 to 0.01 | 0.00 | 0.04 to 0.01 | 0.00 | 0.03 to 0.19 | 0.10 | 0.31 to 0.0 |
| 2002 | 2.49 | 7.50 to 0.98 | 0.46 | 1.74 to 0.02 | 0.00 | 0.06 to 0.01 | 0.00 | 0.05 to 0.01 | 0.00 | 0.03 to 0.17 | 0.08 | 0.29 to 0.0 |
| 2003 | 2.43 | 7.35 to 0.91 | 0.42 | 1.62 to 0.02 | 0.00 | 0.06 to 0.01 | 0.00 | 0.04 to 0.01 | 0.00 | 0.03 to 0.15 | 0.08 | 0.24 to 0.0 |
| 2004 | 2.31 | 7.38 to 1.01 | 0.41 | 1.89 to 0.02 | 0.00 | 0.06 to 0.0  | 0.00 | 0.0 to 0.01  | 0.00 | 0.03 to 0.16 | 0.09 | 0.27 to 0.0 |
| 2005 | 2.30 | 7.38 to 1.02 | 0.41 | 1.91 to 0.02 | 0.00 | 0.06 to 0.0  | 0.00 | 0.0 to 0.01  | 0.00 | 0.03 to 0.19 | 0.10 | 0.31 to 0.0 |
| 2006 | 2.28 | 7.37 to 0.99 | 0.39 | 1.85 to 0.02 | 0.00 | 0.06 to 0.0  | 0.00 | 0.0 to 0.01  | 0.00 | 0.03 to 0.22 | 0.11 | 0.38 to 0.0 |
| 2007 | 2.26 | 7.34 to 0.97 | 0.38 | 1.82 to 0.01 | 0.00 | 0.05 to 0.0  | 0.00 | 0.0 to 0.01  | 0.00 | 0.03 to 0.25 | 0.13 | 0.43 to 0.0 |
| 2008 | 2.27 | 7.35 to 0.95 | 0.37 | 1.81 to 0.02 | 0.00 | 0.05 to 0.01 | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.26 | 0.13 | 0.44 to 0.0 |
| 2009 | 2.28 | 7.35 to 1.18 | 0.43 | 2.33 to 0.0  | 0.00 | 0.01 to 0.01 | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.30 | 0.15 | 0.52 to 0.0 |
| 2010 | 2.25 | 7.25 to 1.14 | 0.42 | 2.25 to 0.0  | 0.00 | 0.01 to 0.01 | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.30 | 0.15 | 0.52 to 0.0 |
| 2011 | 2.23 | 7.20 to 1.16 | 0.43 | 2.29 to 0.0  | 0.00 | 0.01 to 0.01 | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.30 | 0.15 | 0.52 to 0.0 |
| 2012 | 2.23 | 7.19 to 1.15 | 0.43 | 2.24 to 0.0  | 0.00 | 0.01 to 0.01 | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.30 | 0.15 | 0.52 to 0.0 |
| 2013 | 2.23 | 7.16 to 1.16 | 0.44 | 2.27 to 0.0  | 0.00 | 0.01 to 0.01 | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.30 | 0.15 | 0.52 to 0.0 |
| 2014 | 2.23 | 7.18 to 1.16 | 0.42 | 2.27 to 0.0  | 0.00 | .01 to 0.01  | 0.00 | 0.03 to 0.01 | 0.00 | 0.03 to 0.30 | 0.15 | 0.52 to 0.0 |

Note: Estimates after 2010 are based on NRI data from 2010 and therefore do not fully reflect changes occurring in the latter part of the time series.

### **Step 3: Estimate Soil Organic C Stock Changes and Direct N<sub>2</sub>O Emissions from Organic Soils**

In this step, soil organic C losses and N<sub>2</sub>O emissions are estimated for organic soils that are drained for agricultural production.

#### ***Step 3a: Direct N<sub>2</sub>O Emissions Due to Drainage of Organic Soils in Cropland and Grassland***

To estimate annual N<sub>2</sub>O emissions from drainage of organic soils in cropland and grassland, the area of drained organic soils in croplands and grasslands for temperate regions is multiplied by the IPCC (2006) default emission factor for temperate soils and the corresponding area in sub-tropical regions is multiplied by the average (12 kg N<sub>2</sub>O-N/ha cultivated) of IPCC (2006) default emission factors for temperate (8 kg N<sub>2</sub>O-N/ha cultivated) and tropical (16 kg N<sub>2</sub>O-N/ha cultivated) organic soils. The uncertainty is determined based on simple error propagation methods (IPCC 2006), including uncertainty in the default emission factor ranging from 2–24 kg N<sub>2</sub>O-N/ha (IPCC 2006).

#### ***Step 3b: Soil Organic C Stock Changes Due to Drainage of Organic Soils in Cropland and Grassland***

Change in soil organic C stocks due to drainage of cropland and grassland soils are estimated annually from 1990 through 2010, based on the land-use and management activity data in conjunction with appropriate loss rate emission factors. The activity data are based on annual data from 1990 through 2010 from the NRI. The results for 2010 are applied to the years 2011 through 2014. Organic Soil emission factors representative of U.S. conditions have been estimated from published studies (Ogle et al. 2003), based on subsidence studies in the United States and Canada (Table A-246). PDFs are constructed as normal densities based on the mean C loss rates and associated variances. Input values are randomly selected from PDFs in a Monte Carlo analysis to estimate SOC change for 50,000 times and produce a 95 percent confidence interval for the inventory results. Losses of soil organic C from drainage of cropland and grassland soils are provided in Table A-239 and Table A-242.

### **Step 4: Estimate Indirect N<sub>2</sub>O Emissions for Croplands and Grasslands**

In this step, N<sub>2</sub>O emissions are estimated for the two indirect emission pathways (N<sub>2</sub>O emissions due to volatilization, and N<sub>2</sub>O emissions due to leaching and runoff of N), which are summed to yield total indirect N<sub>2</sub>O emissions from croplands and grasslands.

#### ***Step 4a: Indirect Soil N<sub>2</sub>O Emissions Due to Volatilization***

Indirect emissions from volatilization of N inputs from synthetic and commercial organic fertilizers, and PRP manure, are calculated according to the amount of mineral N that is transported in gaseous forms from the soil profile and later emitted as soil N<sub>2</sub>O following atmospheric deposition. See Step 1e for additional information about the methods used to compute N losses due to volatilization. The estimated N volatilized is multiplied by the IPCC default emission factor of 0.01 kg N<sub>2</sub>O-N/kg N (IPCC 2006) to estimate total N<sub>2</sub>O emissions from volatilization. The uncertainty is estimated using simple error propagation methods (IPCC 2006), by combining uncertainties in the amount of N volatilized, with uncertainty in the default emission factor ranging from 0.002–0.05 kg N<sub>2</sub>O-N/kg N (IPCC 2006). The estimates and uncertainties are provided in Table A-247.

#### ***Step 4b: Indirect Soil N<sub>2</sub>O Emissions Due to Leaching and Runoff***

The amount of mineral N from synthetic fertilizers, commercial organic fertilizers, PRP manure, crop residue, N mineralization, symbiotic fixation that is transported from the soil profile in aqueous form is used to calculate indirect emissions from leaching of mineral N from soils and losses in runoff of water associated with overland flow. See Step 1e for additional information about the methods used to compute N losses from soils due to leaching and runoff in overland water flows. The total amount of N transported from soil profiles through leaching and surface runoff is multiplied by the IPCC default emission factor of 0.0075 kg N<sub>2</sub>O-N/kg N (IPCC 2006) to estimate emissions for this source. The emission estimates are provided in Table A-248. The uncertainty is estimated based on simple error propagation methods (IPCC 2006), including uncertainty in the default emission factor ranging from 0.0005 to 0.025 kg N<sub>2</sub>O-N/kg N (IPCC 2006).

### **Step 5: Estimate Total Soil Organic C Stock Changes and N<sub>2</sub>O Emissions for U.S. Soils**

#### ***Step 5a: Estimate Total Soil N<sub>2</sub>O Emissions***


Total N<sub>2</sub>O emissions are estimated by adding total direct emissions (from mineral cropland soils, drainage and cultivation of organic soils, and grassland management) to indirect emissions. Uncertainties in the final estimate are

combined using simple error propagation methods (IPCC 2006), and expressed as a 95 percent confidence interval. Estimates are provided in Table A-249.


Direct and indirect simulated emissions of soil N<sub>2</sub>O vary regionally in croplands as a function of N input amount and timing of fertilization, tillage intensity, crop rotation sequence, weather, and soil type. Note that there are other management practices, such as fertilizer formulation (Halvorson et al. 2013), that influence emissions but are not represented in the model simulations. The highest total N<sub>2</sub>O emissions occur in Iowa, Illinois, Kansas, Minnesota, Nebraska and Texas (Table A-251). On a per area unit basis, direct N<sub>2</sub>O emissions are high in the Midwest, Northeast and many of the Mississippi River Basin states where there are high N inputs to hay, corn and soybean crops, and in some western states where irrigated crops are grown that require high N inputs (Figure A-16). Note that although the total crop area in the northeast is relatively low, emissions are high on a per unit area basis because a large portion of the cropland area in these states is used for hay production that receives large N inputs from both fertilizer and symbiotic fixation. Indirect emissions also tend to be high on a per unit of area basis in the Midwest, some Northeastern states and Florida (Figure A-17). In Florida, the high emission rates are driven by relatively high rainfall and coarse textured soils that facilitate N losses from leaching and runoff. Some Great Plains states also have indirect emissions where irrigation can contribute to leaching and runoff.

Direct and indirect emissions from non-federal grasslands are typically lower than the emissions from croplands (Table A-251, Figure A-18, and Figure A-19) because N inputs tend to be lower, particularly from synthetic fertilizer. Texas, Oklahoma, Kansas, Nebraska, Missouri, South Dakota and Montana are the highest emitters for this category due to large land areas used for pastures and rangeland (Table A-251). On a per unit of area basis, direct N<sub>2</sub>O emissions are higher in the Northeastern United States, Lower Mississippi River Basin, some of the Great Lakes and Midwestern states because these grasslands are more intensively managed (legume seeding, fertilization) while western rangelands receive few, if any, N inputs (Figure A-18). Also, rainfall is limited in most of the western United States, and grasslands are not typically irrigated so minimal leaching and runoff of N occurs in these grasslands, and therefore there are lower indirect N<sub>2</sub>O emissions (Figure A-19).


**Figure A-16: Tier 3 Cropland, 2014 Annual Direct N<sub>2</sub>O Emissions, Estimated Using the DAYCENT Model, (kg N/ha/year)**




**Figure A-17: Tier 3 Crops, 2014 Annual N Losses Leading to Indirect N<sub>2</sub>O Emissions, Estimated Using the DAYCENT Model, (kg N/ha/year)**



**Figure A-18: Non-federal Grasslands, 2014 Annual Direct N<sub>2</sub>O Emissions, Estimated Using the DAYCENT Model, (kg N/ha/year)**




**Figure A-19: Non-federal Grasslands, 2014 Annual N Losses Leading to Indirect N<sub>2</sub>O Emissions, Estimated Using the DAYCENT Model, (kg N/ha/year)**




**Step 5b: Estimate Total Soil Organic Stock Change**

The sum of total CO<sub>2</sub> emissions and removals from the Tier 3 DAYCENT Model Approach, Tier 2 IPCC Methods and additional land-use and management considerations are provided in Table A-252. The total change in soil organic C stocks (as seen in the *Land Use, Land-Use Change, and Forestry* chapter) as well as per hectare rate of change varies among the states (Figure A-20 and Figure A-21). The states with highest total amounts of C sequestration are California, Illinois, Iowa, Kentucky, Missouri, North Dakota and Tennessee (Table A-253). On a per hectare basis, the highest rates of C accumulation occur in states found in the Southeast, Northeast, Midwest and Pacific Coast. For organic soils, emission rates are highest in the regions that contain the majority of drained organic soils, including California, Florida, Indiana, Michigan, Minnesota, North Carolina and Wisconsin. On a per unit of area basis, the emission rate patterns are very similar to the total emissions in each state, with the highest rates in coastal states of the Southeast, states surrounding the Great Lakes, and California.

**Figure A-20: Net C Stock Change, per Hectare, for Mineral Soils Under Agricultural Management, 2014**



**Figure A-21: Net C Stock Change, per Hectare, for Organic Soils Under Agricultural Management, 2014**



**Step 5c: Estimate Total CH<sub>4</sub> Emissions from Rice Cultivation**

The sum of total CH<sub>4</sub> emissions from the Tier 3 DAYCENT Model Approach and Tier 1 IPCC Methods are provided in Table A-250. The states with highest total emissions are Arkansas, California, Louisiana and Texas (Table A-254). These states also have the largest areas of rice cultivation, and Louisiana and Texas have a relatively large proportion of fields with a second ratoon crop each year. Ratoon crops extend the period of time under flooded conditions, which leads to more CH<sub>4</sub> emissions.

**Table A-245: Assumptions and Calculations to Estimate the Contribution to Soil Organic Carbon Stocks from Application of Sewage Sludge to Mineral Soils**

|   | 1990          | 1991          | 1992          | 1993          | 1994          | 1995          | 1996          | 1997          | 1998          | 1999          | 2000          | 2001          |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Sewage Sludge N Applied to Agricultural Land (Mg N) <sup>a</sup>    | 51,848        | 55,107        | 58,480        | 61,971        | 64,721        | 67,505        | 72,081        | 75,195        | 78,353        | 80,932        | 83,523        | 86,124        |
| Assimilative Capacity (Mg N/ha) <sup>b</sup>                        | 0.120         | 0.120         | 0.120         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         |
| Area covered by Available Sewage Sludge N (ha) <sup>c</sup>         | 432,067       | 459,226       | 487,336       | 507,957       | 530,503       | 553,322       | 590,828       | 616,357       | 642,240       | 663,381       | 684,612       | 705,932       |
| Average Annual Rate of C storage (Mg C/ha-yr) <sup>d</sup>          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          |
| <b>Contribution to Soil C (MMT CO<sub>2</sub>/yr)<sup>e,f</sup></b> | <b>(0.60)</b> | <b>(0.64)</b> | <b>(0.68)</b> | <b>(0.71)</b> | <b>(0.74)</b> | <b>(0.77)</b> | <b>(0.82)</b> | <b>(0.86)</b> | <b>(0.89)</b> | <b>(0.92)</b> | <b>(0.95)</b> | <b>(0.98)</b> |

|   | 2002          | 2003          | 2004          | 2005          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| Sewage Sludge N Applied to Agricultural Land (Mg N) <sup>a</sup>    | 88,736        | 91,358        | 93,991        | 98,400        | 101,314       | 104,222       | 107,123       | 110,018       | 112,909       | 115,797       | 118,681       | 121,563       | 121,563       |
| Assimilative Capacity (Mg N/ha) <sup>b</sup>                        | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         | 0.122         |
| Area covered by Available Sewage Sludge N (ha) <sup>c</sup>         | 727,341       | 748,836       | 770,418       | 806,559       | 830,447       | 854,276       | 878,055       | 901,790       | 925,487       | 949,154       | 972,796       | 996,417       | 996,417       |
| Average Annual Rate of C storage (Mg C/ha-yr) <sup>d</sup>          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          | 0.38          |
| <b>Contribution to Soil C (MMT CO<sub>2</sub>/yr)<sup>e,f</sup></b> | <b>(1.01)</b> | <b>(1.04)</b> | <b>(1.07)</b> | <b>(1.12)</b> | <b>(1.16)</b> | <b>(1.19)</b> | <b>(1.22)</b> | <b>(1.26)</b> | <b>(1.29)</b> | <b>(1.32)</b> | <b>(1.36)</b> | <b>(1.39)</b> | <b>(1.39)</b> |

<sup>a</sup> N applied to soils described in Step 1d.

<sup>b</sup> Assimilative Capacity is the national average amount of manure-derived N that can be applied on cropland without buildup of nutrients in the soil (Kellogg et al., 2000).

<sup>c</sup> Area covered by sewage sludge N available for application to soils is the available N applied at the assimilative capacity rate. The 1992 assimilative capacity rate was applied to 1990 – 1992 and the 1997 rate was applied to 1993-2014.

<sup>d</sup> Annual rate of C storage based on national average increase in C storage for grazing lands that is attributed to organic matter amendments (0.38 Mg/ha-yr)

<sup>e</sup> Contribution to Soil C is estimated as the product of the area covered by the available sewage sludge N and the average annual C storage attributed to an organic matter amendment.

<sup>f</sup> Some small, undetermined fraction of this applied N is probably not applied to agricultural soils, but instead is applied to forests, home gardens, and other lands.

Note: Values in parentheses indicate net C storage.

**Table A-246: Carbon Loss Rates for Organic Soils Under Agricultural Management in the United States, and IPCC Default Rates (Metric Ton C/ha-yr)**

| Region                                      | Cropland |              | Grassland |                      |
|---|----------|--------------|-----------|----------------------|
|   | IPCC     | U.S. Revised | IPCC      | U.S. Revised         |
| Cold Temperate, Dry & Cold Temperate, Moist | 1        | 11.2±2.5     | 0.25      | 2.8±0.5 <sup>a</sup> |
| Warm Temperate, Dry & Warm Temperate, Moist | 10       | 14.0±2.5     | 2.5       | 3.5±0.8 <sup>a</sup> |
| Sub-Tropical, Dry & Sub-Tropical, Moist     | 1        | 11.2±2.5     | 0.25      | 2.8±0.5 <sup>a</sup> |

<sup>a</sup> There are not enough data available to estimate a U.S. value for C losses from grassland. Consequently, estimates are 25 percent of the values for cropland, which is an assumption that is used for the IPCC default organic soil C losses on grassland.

**Table A-247: Indirect N<sub>2</sub>O Emissions from Volatilization (MMT CO<sub>2</sub> Eq.)**

| Activity     | 1990        | 1991        | 1992        | 1993        | 1994        | 1995        | 1996        | 1997        | 1998        | 1999        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Croplands    | 13.0        | 13.2        | 12.8        | 12.8        | 13.8        | 14.3        | 14.0        | 14.1        | 14.5        | 13.6        | 13.4        | 13.5        | 13.7        | 14.1        | 13.8        | 13.8        | 14.4        | 14.0        | 14.1        | 13.4        | 13.9        | 14.3        | 14.5        | 14.2        | 14.2        |
| Grasslands   | 4.4         | 4.4         | 4.4         | 4.4         | 4.5         | 4.7         | 4.7         | 4.7         | 4.8         | 4.4         | 4.2         | 4.3         | 4.4         | 4.6         | 4.9         | 4.7         | 4.7         | 4.7         | 4.7         | 4.7         | 4.8         | 4.7         | 4.6         | 4.5         | 4.5         |
| <b>Total</b> | <b>17.3</b> | <b>17.5</b> | <b>17.2</b> | <b>17.3</b> | <b>18.3</b> | <b>19.0</b> | <b>18.7</b> | <b>18.7</b> | <b>19.3</b> | <b>18.0</b> | <b>17.7</b> | <b>17.9</b> | <b>18.0</b> | <b>18.7</b> | <b>18.7</b> | <b>18.5</b> | <b>19.1</b> | <b>18.7</b> | <b>18.7</b> | <b>18.1</b> | <b>18.6</b> | <b>18.9</b> | <b>19.1</b> | <b>18.8</b> | <b>18.7</b> |

**Table A-248: Indirect N<sub>2</sub>O Emissions from Leaching and Runoff (MMT CO<sub>2</sub> Eq.)**

| Activity     | 1990        | 1991        | 1992        | 1993        | 1994        | 1995        | 1996        | 1997        | 1998        | 1999        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Croplands    | 23.2        | 18.7        | 21.7        | 32.0        | 14.7        | 24.5        | 22.2        | 19.2        | 26.0        | 20.8        | 16.4        | 20.8        | 19.4        | 18.6        | 24.5        | 20.2        | 22.1        | 25.1        | 22.6        | 22.4        | 25.8        | 26.4        | 26.6        | 26.0        | 26.0        |
| Grasslands   | 17.7        | 13.5        | 15.5        | 16.2        | 13.0        | 14.6        | 18.7        | 12.6        | 15.9        | 12.3        | 13.1        | 16.9        | 16.3        | 14.6        | 19.2        | 10.2        | 16.3        | 13.7        | 15.5        | 20.2        | 12.4        | 13.2        | 12.9        | 12.6        | 12.6        |
| <b>Total</b> | <b>40.9</b> | <b>32.1</b> | <b>37.1</b> | <b>48.2</b> | <b>27.8</b> | <b>39.1</b> | <b>40.9</b> | <b>31.7</b> | <b>41.9</b> | <b>33.0</b> | <b>29.6</b> | <b>37.6</b> | <b>35.7</b> | <b>33.3</b> | <b>43.7</b> | <b>30.4</b> | <b>38.3</b> | <b>38.7</b> | <b>38.1</b> | <b>42.6</b> | <b>38.2</b> | <b>39.6</b> | <b>39.4</b> | <b>38.7</b> | <b>38.6</b> |

**Table A-249: Total N<sub>2</sub>O Emissions from Agricultural Soil Management (MMT CO<sub>2</sub> Eq.)**

| Activity   | 1990         | 1991         | 1992         | 1993         | 1994         | 1995         | 1996         | 1997         | 1998         | 1999         | 2000         | 2001         |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Total Direct</b>  | <b>245.0</b> | <b>241.9</b> | <b>243.5</b> | <b>245.1</b> | <b>243.0</b> | <b>243.7</b> | <b>248.8</b> | <b>246.8</b> | <b>261.5</b> | <b>241.8</b> | <b>245.4</b> | <b>242.2</b> |
| <b>Direct Emissions from Mineral Cropland Soils</b>          | <b>168.6</b> | <b>165.5</b> | <b>166.2</b> | <b>167.2</b> | <b>168.0</b> | <b>166.2</b> | <b>168.4</b> | <b>167.2</b> | <b>177.7</b> | <b>169.0</b> | <b>171.1</b> | <b>167.7</b> |
| Synthetic Fertilizer   | 59.2         | 58.8         | 60.8         | 58.3         | 63.3         | 57.3         | 62.4         | 60.7         | 61.4         | 59.5         | 61.7         | 58.6         |
| Organic Amendment <sup>a</sup>                               | 11.9         | 12.0         | 12.2         | 11.9         | 12.4         | 12.3         | 12.4         | 12.5         | 12.7         | 12.7         | 13.0         | 12.8         |
| Residue N <sup>b</sup>                                       | 25.9         | 28.1         | 25.9         | 26.3         | 25.5         | 27.5         | 26.7         | 26.4         | 25.6         | 28.9         | 27.6         | 26.3         |
| Mineralization and Asymbiotic Fixation                       | 71.6         | 66.6         | 67.3         | 70.7         | 66.8         | 69.1         | 67.0         | 67.5         | 78.0         | 67.9         | 68.7         | 70.1         |
| <b>Direct Emissions from Drained Organic Cropland Soils</b>  | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.1</b>   | <b>3.1</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.3</b>   |
| <b>Direct Emissions from Mineral Grassland Soils</b>         | <b>70.3</b>  | <b>70.4</b>  | <b>71.4</b>  | <b>71.9</b>  | <b>69.0</b>  | <b>71.5</b>  | <b>74.4</b>  | <b>73.7</b>  | <b>77.8</b>  | <b>66.8</b>  | <b>68.3</b>  | <b>68.2</b>  |
| Synthetic Mineral Fertilizer                                 | 1.1          | 1.1          | 1.1          | 1.2          | 1.1          | 1.1          | 1.1          | 1.1          | 1.4          | 1.1          | 1.2          | 1.2          |
| PRP Manure   | 13.4         | 12.8         | 13.3         | 13.9         | 14.7         | 15.3         | 16.0         | 14.6         | 14.7         | 12.3         | 12.5         | 12.0         |
| Managed Manure   | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            |
| Sewage Sludge  | 0.2          | 0.3          | 0.3          | 0.3          | 0.3          | 0.3          | 0.3          | 0.4          | 0.4          | 0.4          | 0.4          | 0.4          |
| Residue <sup>b</sup>   | 19.7         | 20.1         | 20.3         | 20.5         | 18.3         | 20.3         | 19.8         | 20.4         | 20.2         | 20.3         | 19.3         | 19.8         |
| Mineralization and Asymbiotic Fixation                       | 35.8         | 36.1         | 36.3         | 35.8         | 34.4         | 34.4         | 36.9         | 37.1         | 41.1         | 32.5         | 34.7         | 34.6         |
| <b>Direct Emissions from Drained Organic Grassland Soils</b> | <b>2.9</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>2.9</b>   | <b>2.9</b>   | <b>2.9</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>3.0</b>   |
| <b>Total Indirect</b>  | <b>58.2</b>  | <b>49.7</b>  | <b>54.4</b>  | <b>65.4</b>  | <b>46.1</b>  | <b>58.1</b>  | <b>59.6</b>  | <b>50.5</b>  | <b>61.2</b>  | <b>51.1</b>  | <b>47.2</b>  | <b>55.5</b>  |
| Volatilization   | 17.3         | 17.5         | 17.2         | 17.3         | 18.3         | 19.0         | 18.7         | 18.7         | 19.3         | 18.0         | 17.7         | 17.9         |
| Leaching/Runoff  | 40.9         | 32.1         | 37.1         | 48.2         | 27.8         | 39.1         | 40.9         | 31.7         | 41.9         | 33.0         | 29.6         | 37.6         |
| <b>Total Emissions</b>                                       | <b>303.3</b> | <b>291.5</b> | <b>297.9</b> | <b>310.5</b> | <b>289.1</b> | <b>301.8</b> | <b>308.4</b> | <b>297.3</b> | <b>322.8</b> | <b>292.9</b> | <b>292.6</b> | <b>297.6</b> |

**Table A-249 continued: Total N<sub>2</sub>O Emissions from Agricultural Soil Management (MMT CO<sub>2</sub> Eq.)**

| Activity   | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Total Direct</b>  | <b>245.4</b> | <b>249.7</b> | <b>268.7</b> | <b>248.3</b> | <b>252.2</b> | <b>260.4</b> | <b>251.2</b> | <b>253.5</b> | <b>263.8</b> | <b>264.5</b> | <b>264.5</b> | <b>261.2</b> | <b>261.0</b> |
| <b>Direct Emissions from Mineral Cropland Soils</b>          | <b>168.4</b> | <b>172.4</b> | <b>181.3</b> | <b>171.2</b> | <b>172.2</b> | <b>178.8</b> | <b>172.1</b> | <b>171.8</b> | <b>182.6</b> | <b>183.9</b> | <b>184.9</b> | <b>182.2</b> | <b>182.0</b> |
| Synthetic Fertilizer   | 60.3         | 61.2         | 63.8         | 61.4         | 62.3         | 65.6         | 61.7         | 59.9         | 59.3         | 61.0         | 61.8         | 59.5         | 59.3         |
| Organic Amendment <sup>a</sup>                               | 13.0         | 13.1         | 12.9         | 12.9         | 13.4         | 13.5         | 13.3         | 13.2         | 13.4         | 13.5         | 13.6         | 13.5         | 13.5         |
| Residue N <sup>b</sup>                                       | 26.7         | 27.6         | 25.5         | 26.6         | 26.1         | 26.3         | 25.6         | 25.2         | 27.8         | 27.6         | 27.5         | 27.5         | 27.6         |
| Mineralization and Asymbiotic Fixation                       | 68.3         | 70.5         | 79.1         | 70.3         | 70.3         | 73.3         | 71.5         | 73.4         | 82.2         | 81.8         | 82.0         | 81.6         | 81.6         |
| <b>Direct Emissions from Drained Organic Cropland Soils</b>  | <b>3.3</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.2</b>   | <b>3.1</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   | <b>3.0</b>   |
| <b>Direct Emissions from Mineral Grassland Soils</b>         | <b>70.8</b>  | <b>71.2</b>  | <b>81.3</b>  | <b>71.0</b>  | <b>74.0</b>  | <b>75.6</b>  | <b>73.1</b>  | <b>75.9</b>  | <b>75.5</b>  | <b>74.9</b>  | <b>74.0</b>  | <b>73.3</b>  | <b>73.3</b>  |
| Synthetic Mineral Fertilizer                                 | 1.3          | 1.3          | 1.5          | 1.3          | 1.4          | 1.3          | 1.3          | 1.4          | 1.3          | 1.2          | 1.2          | 1.2          | 1.4          |
| PRP Manure   | 12.2         | 12.5         | 12.9         | 12.3         | 13.6         | 12.8         | 12.8         | 13.1         | 12.5         | 11.9         | 11.0         | 10.3         | 10.3         |
| Managed Manure   | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            | +            |
| Sewage Sludge  | 0.4          | 0.4          | 0.4          | 0.5          | 0.5          | 0.5          | 0.5          | 0.5          | 0.5          | 0.5          | 0.6          | 0.6          | 0.6          |
| Residue <sup>b</sup>   | 20.1         | 20.2         | 20.9         | 21.0         | 20.3         | 21.5         | 20.7         | 20.3         | 21.8         | 21.7         | 21.7         | 21.7         | 21.6         |
| Mineralization and Asymbiotic Fixation                       | 36.7         | 36.6         | 45.3         | 35.8         | 38.0         | 39.3         | 37.7         | 40.5         | 39.1         | 39.4         | 39.4         | 39.4         | 39.2         |
| <b>Direct Emissions from Drained Organic Grassland Soils</b> | <b>3.0</b>   | <b>2.9</b>   | <b>2.9</b>   | <b>2.9</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>2.8</b>   | <b>2.7</b>   | <b>2.7</b>   | <b>2.7</b>   | <b>2.7</b>   | <b>2.7</b>   | <b>2.7</b>   |
| <b>Total Indirect</b>  | <b>53.7</b>  | <b>51.9</b>  | <b>62.4</b>  | <b>48.9</b>  | <b>57.4</b>  | <b>57.5</b>  | <b>56.8</b>  | <b>60.8</b>  | <b>56.9</b>  | <b>58.6</b>  | <b>58.5</b>  | <b>57.4</b>  | <b>57.3</b>  |
| Volatilization   | 18.0         | 18.7         | 18.7         | 18.5         | 19.1         | 18.7         | 18.7         | 18.1         | 18.6         | 18.9         | 19.1         | 18.8         | 18.7         |
| Leaching/Runoff  | 35.7         | 33.3         | 43.7         | 30.4         | 38.3         | 38.7         | 38.1         | 42.6         | 38.2         | 39.6         | 39.4         | 38.7         | 38.6         |
| <b>Total Emissions</b>                                       | <b>299.1</b> | <b>301.7</b> | <b>331.1</b> | <b>297.2</b> | <b>309.6</b> | <b>317.8</b> | <b>308.0</b> | <b>314.2</b> | <b>320.7</b> | <b>323.1</b> | <b>323.1</b> | <b>318.6</b> | <b>318.4</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> Organic amendment inputs include managed manure amendments, daily spread manure and other commercial organic fertilizer (i.e., dried blood, tankage, compost, and other).

<sup>b</sup> Residue N inputs include unharvested fixed N from legumes as well as crop residue N.

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

**Table A-250: Total CH<sub>4</sub> Emissions from Cultivation of Rice Estimated with Tier 1 and 3 Inventory Approaches (MMT CO<sub>2</sub> Eq.)**

| Approach     | Rice Methane (MMT CO <sub>2</sub> Eq) |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |              |             |             |              |              |              |              |              |              |
|--------------|---------------------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|-------------|-------------|--------------|--------------|--------------|--------------|--------------|--------------|
|              | 1990                                  | 1991         | 1992         | 1993         | 1994         | 1995         | 1996         | 1997         | 1998         | 1999         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007        | 2008        | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
| Tier 1       | 1.05                                  | 1.04         | 1.10         | 0.98         | 1.07         | 0.71         | 0.95         | 0.81         | 1.09         | 1.21         | 1.56         | 1.08         | 1.18         | 1.13         | 1.33         | 1.08         | 1.02         | 0.99        | 1.10        | 1.08         | 1.14         | 1.13         | 1.18         | 1.20         | 1.18         |
| Tier 3       | 12.08                                 | 11.93        | 11.91        | 12.65        | 9.32         | 12.41        | 11.45        | 10.47        | 12.39        | 11.37        | 12.83        | 10.97        | 14.01        | 9.87         | 9.70         | 11.93        | 9.25         | 8.95        | 7.84        | 9.88         | 10.71        | 10.71        | 10.72        | 10.71        | 10.72        |
| <b>Total</b> | <b>13.13</b>                          | <b>12.97</b> | <b>13.01</b> | <b>13.64</b> | <b>10.39</b> | <b>13.12</b> | <b>12.40</b> | <b>11.28</b> | <b>13.48</b> | <b>12.59</b> | <b>14.39</b> | <b>12.05</b> | <b>15.18</b> | <b>11.01</b> | <b>11.03</b> | <b>13.01</b> | <b>10.27</b> | <b>9.95</b> | <b>8.94</b> | <b>10.96</b> | <b>11.85</b> | <b>11.84</b> | <b>11.90</b> | <b>11.91</b> | <b>11.89</b> |

**Table A-251: Total 2014 N<sub>2</sub>O Emissions (Direct and Indirect) from Agricultural Soil Management by State (MMT CO<sub>2</sub> Eq.)**

| State           | Croplands <sup>a</sup> | Grasslands <sup>b</sup> | Total | Lower Bound | Upper Bound |
|-----------------|------------------------|-------------------------|-------|-------------|-------------|
| AL              | 1.45                   | 1.46                    | 2.94  | 2.34        | 4.16        |
| AR              | 5.53                   | 2.88                    | 8.48  | 6.80        | 11.62       |
| AZ              | 1.00                   | 1.00                    | 2.07  | 1.60        | 3.44        |
| CA              | 7.12                   | 0.84                    | 8.41  | 5.88        | 17.07       |
| CO              | 4.37                   | 2.96                    | 7.37  | 6.09        | 9.86        |
| CT              | 0.09                   | 0.04                    | 0.13  | 0.09        | 0.21        |
| DE              | 0.26                   | 0.02                    | 0.29  | 0.21        | 0.43        |
| FL              | 2.45                   | 2.98                    | 5.52  | 4.29        | 9.63        |
| GA              | 1.98                   | 1.00                    | 3.01  | 2.42        | 4.43        |
| HI <sup>c</sup> | 0.02                   | 0.15                    | 0.17  | 0.06        | 0.33        |
| IA              | 22.77                  | 3.05                    | 25.93 | 19.94       | 37.32       |
| ID              | 2.90                   | 0.71                    | 3.67  | 2.91        | 5.63        |
| IL              | 14.98                  | 1.17                    | 16.17 | 12.84       | 21.28       |
| IN              | 8.08                   | 0.77                    | 8.93  | 7.27        | 11.78       |
| KS              | 13.39                  | 4.15                    | 17.61 | 14.42       | 23.72       |
| KY              | 2.63                   | 2.97                    | 5.64  | 4.49        | 7.93        |
| LA              | 3.24                   | 1.46                    | 4.77  | 3.91        | 6.16        |
| MA              | 0.12                   | 1.24                    | 0.19  | 0.15        | 0.28        |
| MD              | 0.83                   | 0.19                    | 1.04  | 0.80        | 1.53        |
| ME              | 0.20                   | 0.26                    | 0.32  | 0.23        | 0.54        |
| MI              | 4.91                   | 0.95                    | 6.01  | 4.89        | 8.58        |
| MN              | 14.61                  | 1.88                    | 17.10 | 13.83       | 23.42       |
| MO              | 9.21                   | 6.43                    | 15.69 | 12.52       | 22.12       |
| MS              | 3.42                   | 1.14                    | 4.53  | 3.72        | 6.05        |
| MT              | 4.54                   | 4.98                    | 9.55  | 7.94        | 12.10       |
| NC              | 2.05                   | 0.64                    | 2.79  | 1.95        | 5.02        |
| ND              | 9.14                   | 1.53                    | 10.62 | 8.61        | 13.49       |
| NE              | 16.40                  | 3.63                    | 20.11 | 15.93       | 28.48       |
| NH              | 0.06                   | 0.04                    | 0.11  | 0.08        | 0.25        |
| NJ              | 0.24                   | 0.07                    | 0.32  | 0.25        | 0.45        |
| NM              | 1.08                   | 2.61                    | 3.61  | 2.86        | 5.21        |
| NV              | 0.37                   | 0.95                    | 0.76  | 0.56        | 1.25        |
| NY              | 3.20                   | 1.39                    | 4.65  | 3.73        | 6.75        |
| OH              | 7.75                   | 0.93                    | 8.89  | 7.09        | 12.93       |
| OK              | 3.99                   | 5.54                    | 9.52  | 7.65        | 13.22       |
| OR              | 1.37                   | 1.13                    | 2.51  | 2.08        | 3.35        |
| PA              | 2.92                   | 0.93                    | 3.89  | 3.09        | 5.81        |
| RI              | 0.02                   | 0.01                    | 0.02  | 0.02        | 0.05        |
| SC              | 0.91                   | 0.32                    | 1.25  | 1.01        | 1.83        |
| SD              | 9.59                   | 3.93                    | 13.51 | 11.17       | 17.30       |
| TN              | 2.15                   | 2.50                    | 4.69  | 3.69        | 6.77        |
| TX              | 13.55                  | 14.02                   | 27.68 | 23.39       | 35.40       |
| UT              | 0.78                   | 0.73                    | 1.53  | 1.22        | 2.22        |
| VA              | 1.06                   | 1.55                    | 2.61  | 2.10        | 3.61        |
| VT              | 0.54                   | 0.34                    | 0.89  | 0.70        | 1.37        |
| WA              | 2.44                   | 0.66                    | 3.19  | 2.62        | 4.56        |
| WI              | 9.50                   | 1.91                    | 11.81 | 9.10        | 18.40       |
| WV              | 0.28                   | 0.70                    | 0.98  | 0.71        | 1.56        |
| WY              | 1.49                   | 2.60                    | 4.13  | 3.37        | 5.47        |

<sup>a</sup> Emissions from non-manure organic N inputs for crops not simulated by DAYCENT were not estimated (NE) at the state level.

<sup>b</sup> Emissions from sewage sludge applied to grasslands and were not estimated (NE) at the state level

<sup>c</sup> N<sub>2</sub>O emissions are not reported for Hawaii except from cropland organic soils.

**Table A-252: Annual Soil C Stock Change in Cropland Remaining Cropland (CRC), Land Converted to Cropland (LCC), Grassland Remaining Grassland (GRG), and Land Converted to Grassland (LCG), in U.S. Agricultural Soils (MMT CO<sub>2</sub> Eq.)**

|   | 1990   | 1991   | 1992   | 1993   | 1994   | 1995   | 1996   | 1997   | 1998   | 1999   | 2000   | 2001   | 2002   | 2003   | 2004   | 2005   | 2006   | 2007   | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   |        |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| <b>Net emissions based on Tier 3 Century-based analysis (Step 2)</b>  |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| CRC   | (56.2) | (57.0) | (58.3) | (37.1) | (48.5) | (43.0) | (50.6) | (47.1) | (33.5) | (46.5) | (51.1) | (50.9) | (42.2) | (33.0) | (38.1) | (37.2) | (37.7) | (37.8) | (29.7) | (22.3) | (22.0) | (36.7) | (36.7) | (36.7) | (36.7) | (36.7) |
| GCC   | 12.9   | 12.8   | 13.6   | 11.6   | 6.5    | 13.6   | 11.2   | 13.1   | 7.6    | 8.5    | 8.4    | 8.2    | 8.6    | 7.7    | 6.9    | 7.9    | 7.9    | 5.8    | 6.8    | 5.8    | 6.6    | 4.5    | 4.5    | 4.5    | 4.5    | 4.5    |
| GRG   | (17.9) | (8.4)  | (9.3)  | (8.7)  | (24.2) | (7.4)  | (16.5) | (9.9)  | (13.1) | (3.8)  | (19.2) | (8.5)  | (6.8)  | (5.1)  | (7.4)  | (5.3)  | (14.4) | 1.3    | (8.4)  | (10.8) | (9.0)  | 1.1    | 1.1    | 1.1    | 1.1    | 1.1    |
| CCG   | (3.8)  | (4.7)  | (4.7)  | (5.1)  | (5.4)  | (5.6)  | (6.0)  | (6.6)  | (6.8)  | (6.9)  | (7.1)  | (7.6)  | (8.2)  | (7.3)  | (8.0)  | (7.2)  | (8.0)  | (7.9)  | (8.1)  | (8.0)  | (7.7)  | (7.2)  | (7.2)  | (7.2)  | (7.2)  | (7.2)  |
| <b>Net emissions based on the IPCC Tier 2 analysis (Step 3)</b>   |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| <b>Mineral Soils</b>  |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| CRC   | (6.2)  | (6.7)  | (7.4)  | (7.1)  | (7.0)  | (6.5)  | (6.1)  | (7.7)  | (7.3)  | (6.6)  | (7.1)  | (6.9)  | (6.8)  | (6.1)  | (5.5)  | (5.6)  | (4.7)  | (4.6)  | (4.0)  | (3.9)  | (4.0)  | (3.7)  | (3.3)  | (3.1)  | (3.1)  | (3.2)  |
| GCC   | 1.3    | 1.2    | 1.3    | 1.4    | 1.4    | 1.5    | 1.8    | 1.5    | 1.6    | 1.6    | 1.5    | 1.4    | 1.2    | 1.1    | 1.2    | 1.2    | 1.4    | 1.4    | 1.4    | 1.3    | 1.3    | 1.3    | 1.3    | 1.3    | 1.3    | 1.3    |
| FCC   | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    |
| OCC   | 0.2    | 0.1    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.1    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    |
| SCC   | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.0    | 0.0    | 0.0    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    |
| WCC   | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.2    | 0.2    | 0.1    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.1    | 0.1    | 0.1    | 0.2    | 0.2    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    |
| GRG   | (0.5)  | (0.6)  | (0.9)  | (1.4)  | (1.7)  | (1.7)  | (1.0)  | (0.9)  | (1.6)  | (1.3)  | (1.5)  | (1.5)  | (2.5)  | (2.6)  | (1.2)  | (1.3)  | (1.5)  | (1.5)  | (1.5)  | (1.3)  | (1.4)  | (1.0)  | (0.5)  | (0.3)  | (0.3)  | (0.3)  |
| CCG   | (3.1)  | (3.1)  | (3.0)  | (3.3)  | (3.4)  | (3.3)  | (3.1)  | (2.9)  | (3.2)  | (2.9)  | (3.1)  | (2.9)  | (2.7)  | (2.4)  | (2.7)  | (2.4)  | (2.5)  | (2.1)  | (1.7)  | (1.6)  | (1.5)  | (1.4)  | (1.4)  | (1.4)  | (1.4)  | (1.4)  |
| FCG   | (0.5)  | (0.5)  | (0.4)  | (0.5)  | (0.4)  | (0.5)  | (0.4)  | (0.4)  | (1.1)  | (1.1)  | (1.1)  | (1.0)  | (1.0)  | (1.0)  | (1.0)  | (1.0)  | (1.0)  | (0.9)  | (0.9)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  |
| OCG   | (0.6)  | (0.6)  | (0.7)  | (0.8)  | (1.0)  | (1.0)  | (0.9)  | (1.0)  | (1.1)  | (1.1)  | (1.2)  | (1.2)  | (1.2)  | (1.1)  | (1.1)  | (1.1)  | (1.0)  | (0.9)  | (0.9)  | (0.9)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  | (0.8)  |
| SCG   | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.2)  | (0.2)  | (0.2)  | (0.2)  | (0.1)  | (0.2)  | (0.2)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  | (0.1)  |
| WCG   | (0.5)  | (0.6)  | (0.6)  | (0.6)  | (0.7)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.6)  | (0.5)  | (0.5)  | (0.4)  | (0.3)  | (0.2)  | (0.2)  | (0.2)  | (0.2)  | (0.2)  | (0.2)  |
| <b>Organic Soils</b>  |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| CRC   | 28.0   | 27.5   | 27.5   | 27.2   | 27.2   | 27.1   | 27.0   | 26.9   | 27.1   | 26.4   | 26.3   | 27.1   | 27.4   | 27.6   | 28.8   | 28.7   | 28.5   | 27.6   | 27.2   | 27.7   | 27.8   | 27.8   | 27.8   | 27.8   | 27.8   | 27.8   |
| GCC   | 3.2    | 3.2    | 3.2    | 3.1    | 3.4    | 3.7    | 3.7    | 3.8    | 4.3    | 4.4    | 4.1    | 5.4    | 5.3    | 5.1    | 4.2    | 4.2    | 4.2    | 4.0    | 3.9    | 3.7    | 3.8    | 3.8    | 3.8    | 3.8    | 3.8    | 3.8    |
| FCC   | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    |
| OCC   | 0.1    | 0.1    | 0.0    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    |
| SCC   | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.0    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    |
| WCC   | 0.6    | 0.6    | 0.6    | 0.7    | 0.9    | 0.9    | 0.9    | 0.9    | 1.0    | 0.9    | 0.8    | 0.8    | 0.7    | 0.6    | 0.5    | 0.5    | 0.5    | 0.5    | 0.5    | 0.4    | 0.4    | 0.4    | 0.4    | 0.4    | 0.4    | 0.4    |
| GRG   | 6.1    | 6.0    | 5.9    | 5.8    | 5.7    | 5.6    | 5.5    | 5.4    | 5.3    | 5.2    | 5.2    | 4.7    | 4.6    | 4.5    | 4.5    | 4.4    | 4.4    | 4.4    | 4.4    | 4.4    | 4.4    | 4.4    | 4.3    | 4.3    | 4.3    | 4.3    |
| CCG   | 0.5    | 0.5    | 0.5    | 0.6    | 0.7    | 0.7    | 0.7    | 0.7    | 0.8    | 0.8    | 0.8    | 0.9    | 1.0    | 0.9    | 1.0    | 1.0    | 1.0    | 1.0    | 1.0    | 1.2    | 1.1    | 1.2    | 1.2    | 1.2    | 1.2    | 1.2    |
| FCG   | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    |
| OCG   | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    |
| SCG   | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    | 0.0    |
| WCG   | 0.1    | 0.1    | 0.1    | 0.1    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.1    | 0.2    | 0.2    | 0.2    | 0.2    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    |
| <b>Additional changes in net emissions from mineral soils based on application of sewage sludge to agricultural land (Step 4)</b> |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| GRG   | (0.6)  | (0.6)  | (0.7)  | (0.7)  | (0.7)  | (0.8)  | (0.8)  | (0.9)  | (0.9)  | (0.9)  | (1.0)  | (1.0)  | (1.0)  | (1.0)  | (1.1)  | (1.1)  | (1.2)  | (1.2)  | (1.2)  | (1.3)  | (1.3)  | (1.3)  | (1.4)  | (1.4)  | (1.4)  | (1.4)  |
| <b>Additional changes in net emissions from mineral soils based on additional enrollment of CRP land (Step 4)</b>                 |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| CRC   | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | -      | 0.1    | 1.1    | 2.8    | 3.7    | 3.7    |
| <b>Total Stock Changes by Land Use/Land-Use Change Category (Step 5)</b>  |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |        |
| CRC   | (34.3) | (36.2) | (38.2) | (17.0) | (28.3) | (22.3) | (29.7) | (27.9) | (13.7) | (26.8) | (31.8) | (30.6) | (21.6) | (11.5) | (14.8) | (14.1) | (13.8) | (14.7) | (6.5)  | 1.5    | 1.8    | (12.5) | (11.2) | (9.3)  | (8.4)  | (8.4)  |
| GCC   | 17.4   | 17.3   | 18.1   | 16.2   | 11.3   | 18.9   | 16.7   | 18.4   | 13.5   | 14.5   | 13.9   | 15.0   | 15.1   | 13.9   | 12.3   | 13.3   | 13.5   | 11.3   | 12.1   | 10.7   | 11.8   | 9.6    | 9.6    | 9.7    | 9.7    | 9.7    |
| FCC   | 0.3    | 0.3    | 0.3    | 0.3    | 0.2    | 0.3    | 0.3    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    |
| OCC   | 0.3    | 0.3    | 0.2    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.3    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    | 0.2    |
| SCC   | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    | 0.1    |
| WCC   | 0.7    | 0.7    | 0.7    | 0.9    | 1.0    | 1.1    | 1.1    | 1.1    | 1.2    | 1.1    | 1.0    | 1.0    | 0.9    | 0.8    | 0.7    | 0.6    | 0.7    | 0.7    | 0.6    | 0.6    | 0.6    | 0.6    | 0.6    | 0.6    | 0.6    | 0.6    |
| GRG   | (12.9) | (3.7)  | (5.1)  | (5.0)  | (20.9) | (4.2)  | (12.8) | (6.3)  | (10.3) | (0.9)  | (16.6) | (6.2)  | (5.7)  | (4.2)  | (5.2)  | (3.3)  | (12.5) | 3.0    | (6.8)  | (8.9)  | (7.3)  | 3.1    | 3.6    | 3.8    | 3.8    | 3.8    |

|                |               |               |               |               |               |               |               |               |               |               |               |               |               |               |               |               |               |               |               |              |              |              |              |              |              |
|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|
| CCG            | (6.4)         | (7.3)         | (7.1)         | (7.8)         | (8.1)         | (8.1)         | (8.3)         | (8.8)         | (9.2)         | (9.0)         | (9.4)         | (9.7)         | (9.8)         | (8.8)         | (9.7)         | (8.7)         | (9.0)         | (8.6)         | (8.8)         | (8.3)        | (8.0)        | (7.4)        | (7.4)        | (7.4)        | (7.4)        |
| FCG            | (0.5)         | (0.5)         | (0.4)         | (0.4)         | (0.4)         | (0.5)         | (0.4)         | (0.4)         | (1.1)         | (1.1)         | (1.1)         | (1.0)         | (1.0)         | (0.9)         | (0.9)         | (1.0)         | (0.9)         | (0.9)         | (0.8)         | (0.8)        | (0.8)        | (0.8)        | (0.8)        | (0.8)        | (0.8)        |
| OCC            | (0.6)         | (0.6)         | (0.7)         | (0.8)         | (0.9)         | (1.0)         | (0.9)         | (1.0)         | (1.1)         | (1.1)         | (1.2)         | (1.2)         | (1.1)         | (1.1)         | (1.1)         | (1.1)         | (1.0)         | (0.9)         | (0.8)         | (0.8)        | (0.8)        | (0.8)        | (0.8)        | (0.8)        | (0.8)        |
| SCG            | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.2)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)         | (0.1)        | (0.1)        | (0.1)        | (0.1)        | (0.1)        | (0.1)        |
| WCG            | (0.4)         | (0.5)         | (0.5)         | (0.5)         | (0.5)         | (0.4)         | (0.4)         | (0.4)         | (0.4)         | (0.4)         | (0.5)         | (0.4)         | (0.4)         | (0.4)         | (0.4)         | (0.4)         | (0.3)         | (0.2)         | (0.1)         | 0.0          | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          |
| <b>Total *</b> | <b>(36.4)</b> | <b>(30.2)</b> | <b>(32.7)</b> | <b>(13.9)</b> | <b>(46.3)</b> | <b>(16.0)</b> | <b>(34.2)</b> | <b>(25.0)</b> | <b>(20.8)</b> | <b>(23.2)</b> | <b>(45.1)</b> | <b>(32.9)</b> | <b>(23.5)</b> | <b>(12.1)</b> | <b>(19.0)</b> | <b>(14.3)</b> | <b>(23.1)</b> | <b>(10.1)</b> | <b>(10.8)</b> | <b>(5.9)</b> | <b>(2.5)</b> | <b>(7.9)</b> | <b>(6.2)</b> | <b>(4.0)</b> | <b>(3.1)</b> |

Note: Totals may not sum due to independent rounding.

**Table A-253: Soil C Stock Change for Mineral and Organic Soils in 2014 within individual states (MMT CO<sub>2</sub> Eq.)**

| State | Mineral Soil | Organic Soil | Total  |
|-------|--------------|--------------|--------|
| AL    | (1.29)       | 0.01         | (1.27) |
| AR    | (0.78)       | -            | (0.78) |
| AZ    | 1.14         | -            | 1.14   |
| CA    | (3.88)       | 1.58         | (2.29) |
| CO    | (0.84)       | 0.00         | (0.83) |
| CT    | 0.00         | 0.01         | 0.01   |
| DE    | (0.10)       | -            | (0.10) |
| FL    | 0.21         | 12.21        | 12.42  |
| GA    | (0.75)       | -            | (0.75) |
| HI    | (0.10)       | 0.77         | 0.67   |
| IA    | (2.33)       | 0.73         | (1.60) |
| ID    | (0.76)       | 0.03         | (0.73) |
| IL    | (3.15)       | 0.52         | (2.63) |
| IN    | (1.68)       | 2.36         | 0.68   |
| KS    | (1.41)       | -            | (1.41) |
| KY    | (2.74)       | -            | (2.74) |
| LA    | (1.44)       | 0.51         | (0.93) |
| MA    | (0.07)       | 0.28         | 0.22   |
| MD    | (0.42)       | 0.01         | (0.42) |
| ME    | (0.11)       | 0.01         | (0.10) |
| MI    | (0.62)       | 3.40         | 2.78   |
| MN    | (1.79)       | 7.65         | 5.86   |
| MO    | (3.77)       | -            | (3.77) |
| MS    | (0.55)       | 0.01         | (0.53) |
| MT    | 0.46         | 0.15         | 0.61   |
| NC    | (1.24)       | 1.89         | 0.65   |
| ND    | (3.45)       | 0.01         | (3.44) |
| NE    | (1.44)       | 0.00         | (1.43) |
| NH    | (0.03)       | 0.02         | (0.01) |
| NJ    | (0.07)       | 0.12         | 0.05   |
| NM    | 0.24         | -            | 0.24   |
| NV    | (0.08)       | 0.00         | (0.08) |
| NY    | (1.14)       | 0.53         | (0.61) |
| OH    | (1.45)       | 0.48         | (0.97) |
| OK    | 0.48         | -            | 0.48   |
| OR    | (0.73)       | 0.30         | (0.43) |
| PA    | (0.46)       | 0.05         | (0.41) |
| RI    | 0.01         | 0.02         | 0.03   |
| SC    | (0.90)       | 0.02         | (0.88) |
| SD    | (0.84)       | -            | (0.84) |
| TN    | (2.23)       | -            | (2.23) |
| TX    | (0.07)       | -            | (0.07) |
| UT    | (0.02)       | 0.08         | 0.06   |
| VA    | (1.46)       | 0.00         | (1.46) |
| VT    | (0.12)       | 0.06         | (0.05) |
| WA    | (1.02)       | 0.38         | (0.64) |
| WI    | 0.55         | 2.90         | 3.45   |
| WV    | (0.55)       | -            | (0.55) |
| WY    | 0.30         | -            | 0.30   |

Note: Parentheses indicate net C accumulation. Estimates do not include soil C stock change associated with federal croplands and grasslands, CRP enrollment after 2010, or sewage sludge application to soils, which were only estimated at the national scale. The sum of state results will not match the national results because state results are generated in a separate programming package, the sewage sludge and CRP enrollment after 2010 are not included, and differences arise due to rounding of values in this table.

**Table A-254: Total 2014 CH<sub>4</sub> Emissions from Rice Cultivation by State (MMT CO<sub>2</sub> Eq.)**

| State | Total |
|-------|-------|
| AL    | -     |
| AR    | 4.55  |
| AZ    | -     |
| CA    | 2.36  |
| CO    | -     |

|    |      |
|----|------|
| CT | -    |
| DE | -    |
| FL | -    |
| GA | -    |
| HI | -    |
| IA | -    |
| ID | -    |
| IL | -    |
| IN | -    |
| KS | -    |
| KY | -    |
| LA | 2.66 |
| MA | -    |
| MD | -    |
| ME | -    |
| MI | -    |
| MN | 0.01 |
| MO | 0.72 |
| MS | 0.25 |
| MT | -    |
| NC | -    |
| ND | -    |
| NE | -    |
| NH | -    |
| NJ | -    |
| NM | -    |
| NV | -    |
| NY | -    |
| OH | -    |
| OK | -    |
| OR | -    |
| PA | -    |
| RI | -    |
| SC | -    |
| SD | -    |
| TN | -    |
| TX | 1.34 |
| UT | -    |
| VA | -    |
| VT | -    |
| WA | -    |
| WI | -    |
| WV | -    |
| WY | -    |

---

## References

- AAPFCO (1995 through 2000b, 2002 through 2009) Commercial Fertilizers. Association of American Plant Food Control Officials. University of Kentucky, Lexington, KY.
- AAPFCO (2000a) 1999-2000 Commercial Fertilizers Data, ASCII files. Available from David Terry, Secretary, Association of American Plant Food Control Officials.
- Abdalla, M., Jones, J. Yeluripati, P. Smith, J. Burke and D M. Williams (2010) Testing DayCent and DNDC model simulations of N<sub>2</sub>O fluxes and assessing the impacts of climate change on the gas flux and biomass production from a humid pasture. *Atmos. Environ.* 44: 2961–2970.
- BLM (2014) Rangeland Inventory, Monitoring, and Evaluation Reports. Bureau of Land Management. U.S. Department of the Interior. Available online at: [http://www.blm.gov/wo/st/en/prog/more/rangeland\\_management/rangeland\\_inventory.html](http://www.blm.gov/wo/st/en/prog/more/rangeland_management/rangeland_inventory.html).
- BOEM (2014) Year 2011 Gulfwide Emissions Inventory Study (BOEM 2014-666) Bureau of Ocean Energy Management, U.S. Department of the Interior (November 2014) <<http://www.data.boem.gov/PI/PDFImages/ESPIS/5/5440.pdf>>.
- Cantens, G. (2004 through 2005) Personal Communication. Janet Lewis, Assistant to Gaston Cantens, Vice President of Corporate Relations, Florida Crystals Company and ICF International.
- Cheng, K., S.M. Ogle, W.J. Parton, G. Pan (2014) Simulating greenhouse gas mitigation potentials for Chinese croplands using the DAYCENT ecosystem model. *Global Change Biology* 20:948-962.
- Cheng, K., S.M. Ogle, W.J. Parton and G. Pan (2013) Predicting methanogenesis from rice paddies using the DAYCENT ecosystem model. *Ecological Modelling* 261-262:19-31.
- Cibrowski, P. (1996) Personal Communication. Peter Cibrowski, Minnesota Pollution Control Agency and Heike Mainhardt, ICF Incorporated. July 29, 1996.
- Coulston, J.W., Woodall, C.W., Domke, G.M., and Walters, B.F. (in preparation). Refined Delineation between Woodlands and Forests with Implications for U.S. National Greenhouse Gas Inventory of Forests. *Climatic Change*.
- CTIC (2004) 2004 Crop Residue Management Survey. Conservation Technology Information Center. Available online at <<http://www.ctic.purdue.edu/CRM/>>.
- Daly, C., G.H. Taylor, W.P. Gibson, T. Parzybok, G.L. Johnson, and P.A. Pasteris (1998) “Development of high-quality spatial datasets for the United States.” Proc., 1<sup>st</sup> International Conference on Geospatial Information in Agriculture and Forestry, Lake Buena Vista, FL, I-512-I-519. June 1-3, 1998.
- Daly, C., R.P. Neilson, and D.L. Phillips (1994) “A statistical-topographic model for mapping climatological precipitation over mountainous terrain.” *Journal of Applied Meteorology*, 33:140-158.
- Dean, W. E., and E. Gorham (1998) Magnitude and significance of carbon burial in lakes, reservoirs, and peatlands. *Geology* 26:535-538.
- Del Grosso, S.J., S.M. Ogle, W.J. Parton. (2011) Soil Organic Matter Cycling and Greenhouse Gas Accounting Methodologies, Chapter 1, pp 3-13 DOI: 10.1021/bk-2011-1072.ch001. In: L. Guo, A. Gunasekara, L. McConnell (Eds.) *Understanding Greenhouse Gas Emissions from Agricultural Management*, American Chemical Society, Washington, D.C.
- Del Grosso, S.J., W.J. Parton, C.A. Keough, and M. Reyes-Fox. (2011) Special features of the DayCent modeling package and additional procedures for parameterization, calibration, validation, and applications, in *Methods of Introducing System Models into Agricultural Research*, L.R. Ahuja and Liwang Ma, editors, p. 155-176, American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Madison, WI. USA.
- Del Grosso, S.J., S.M. Ogle, W.J. Parton, and F.J. Breidt (2010) “Estimating Uncertainty in N<sub>2</sub>O Emissions from U.S. Cropland Soils.” *Global Biogeochemical Cycles*, 24, GB1009, doi:10.1029/2009GB003544.
- Del Grosso, S.J., T. Wirth, S.M. Ogle, W.J. Parton (2008) Estimating agricultural nitrous oxide emissions. *EOS* 89, 529-530.
- Del Grosso, S.J., A.R. Mosier, W.J. Parton, and D.S. Ojima (2005) “DAYCENT Model Analysis of Past and Contemporary Soil N<sub>2</sub>O and Net Greenhouse Gas Flux for Major Crops in the USA.” *Soil Tillage and Research*, 83: 9-24. doi: 10.1016/j.still.2005.02.007.

- Del Grosso, S.J., W.J. Parton, A.R. Mosier, M.D. Hartman, J. Brenner, D.S. Ojima, and D.S. Schimel (2001) "Simulated Interaction of Carbon Dynamics and Nitrogen Trace Gas Fluxes Using the DAYCENT Model." In Schaffer, M., L. Ma, S. Hansen, (eds.); Modeling Carbon and Nitrogen Dynamics for Soil Management. CRC Press. Boca Raton, Florida. 303-332.
- Del Grosso, S.J., W.J. Parton, A.R. Mosier, D.S. Ojima, A.E. Kulmala and S. Phongpan (2000) General model for N<sub>2</sub>O and N<sub>2</sub> gas emissions from soils due to denitrification. *Global Biogeochem. Cycles*, 14:1045-1060.
- Delgado, J.A., S.J. Del Grosso, and S.M. Ogle (2009) "15N isotopic crop residue cycling studies and modeling suggest that IPCC methodologies to assess residue contributions to N<sub>2</sub>O-N emissions should be reevaluated." *Nutrient Cycling in Agroecosystems*, DOI 10.1007/s10705-009-9300-9.
- Deren, C. (2002) Personal Communication and Dr. Chris Deren, Everglades Research and Education Centre at the University of Florida and Caren Mintz, ICF International. August 15, 2002.
- Domke, G.M., Woodall, C.W., Smith, J.E., Westfall, J.A., McRoberts, R.E. (2012) Consequences of alternative tree-level biomass estimation procedures on U.S. forest carbon stock estimates. *Forest Ecology and Management*. 270: 108-116.
- Domke, G.M., Smith, J.E., and Woodall, C.W. (2011) Accounting for density reduction and structural loss in standing dead trees: Implications for forest biomass and carbon stock estimates in the United States. *Carbon Balance and Management*. 6:14.
- Domke, G.M., Woodall, C.W., Walters, B.F., McRoberts, R.E., Hatfield, M.A. (In Review) Strategies to compensate for the effects of nonresponse on forest carbon baseline estimates from the national forest inventory of the United States. *Forest Ecology and Management*.
- Domke, G.M., Woodall, C.W., Walters, B.F., Smith, J.E. (2013) From models to measurements: comparing down dead wood carbon stock estimates in the U.S. forest inventory. *PLoS ONE* 8(3): e59949.
- Domke, G.M., Perry, C.H., Walters, B.F., Woodall, C.W., and Smith, J.E. (in preparation). Estimation of forest floor carbon using the national forest inventory of the United States. Intended outlet: *Geoderma*.
- Easter, M., S. Williams, and S. Ogle. (2008) Gap-filling NRI data for the Soil C Inventory. Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO. Report provided to the U.S. Environmental Protection Agency, Tom Wirth.
- Edmonds, L., N. Gollehon, R.L. Kellogg, B. Kintzer, L. Knight, C. Lander, J. Lemunyon, D. Meyer, D.C. Moffitt, and J. Schaeffer (2003) "Costs Associated with Development and Implementation of Comprehensive Nutrient Management Plans." Part 1. Nutrient Management, Land Treatment, Manure and Wastewater Handling and Storage, and Recordkeeping. Natural Resource Conservation Service, U.S. Department of Agriculture.
- EIA (2007) Voluntary Greenhouse Gas Reports for EIA Form 1605B (Reporting Year 2006). Available online at <ftp://ftp.eia.doe.gov/pub/oiaf/1605/cdrom/>.
- Euliss, N., and R. Gleason (2002) Personal communication regarding wetland restoration factor estimates and restoration activity data. Ned Euliss and Robert Gleason of the U.S. Geological Survey, Jamestown, ND, to Stephen Ogle of the National Resource Ecology Laboratory, Fort Collins, CO. August 2002.
- Fleskes, J.P., Perry, W.M., Petrik, K.L., Spell, R., and Reid, F. (2005) Change in area of winter-flood and dry rice in the northern Central Valley of California determined by satellite imagery. *California Fish and Game*, 91: 207-215.
- Fry, J., Xian, G., Jin, S., Dewitz, J., Homer, C., Yang, L., Barnes, C., Herold, N., and Wickham, J. (2011) Completion of the 2006 National Land Cover Database for the Conterminous United States, PE&RS, Vol. 77(9):858-864.
- Gonzalez, R. (2007 through 2014) Email correspondence. Rene Gonzalez, Plant Manager, Sem-Chi Rice Company and ICF International.
- Gurung, R.B., F.J. Breidt, A. Dutin, and S.M. Ogle (2009) Predicting Enhanced Vegetation Index (EVI) for ecosystem modeling applications. *Remote Sensing of Environment* 113:2186-2193.
- Halvorson, A.D., C.S. Snyder, A.D. Blaylock, and S.J. Del Grosso (2013) Enhanced Efficiency Nitrogen Fertilizers: Potential Role in Nitrous Oxide Emission Mitigation. *Agronomy Journal*, doi:10.2134/agronj2013.0081

- Hardke, J.T. (2015) Trends in Arkansas rice production, 2014. B.R. Wells Arkansas Rice Research Studies 2014. Norman, R.J., and Moldenhauer, K.A.K., (Eds.). Research Series 626, Arkansas Agricultural Experiment Station, University of Arkansas.
- Hardke, J.T., and Wilson, C.E. Jr. (2013) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2012. Norman, R.J., and Moldenhauer, K.A.K., (Eds.). Research Series 609, Arkansas Agricultural Experiment Station, University of Arkansas.
- Hardke, J.T., and Wilson, C.E. Jr. (2014) Trends in Arkansas rice production, 2013. B.R. Wells Arkansas Rice Research Studies 2013. Norman, R.J., and Moldenhauer, K.A.K., (Eds.). Research Series 617, Arkansas Agricultural Experiment Station, University of Arkansas.
- Harmon, M.E., C.W. Woodall, B. Fasth, J. Sexton, M. Yatkov. (2011) Differences between standing and downed dead tree wood density reduction factors: A comparison across decay classes and tree species. Res. Paper. NRS-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 40 p
- Hollier, C. A. (ed) (1999) Louisiana rice production handbook. Louisiana State University Agricultural Center. LCES Publication Number 2321. 116 pp.
- Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones and A. Jarvis (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- IPCC (2003) *Good Practice Guidance for Land Use, Land-Use Change, and Forestry*. The Intergovernmental Panel on Climate Change, National Greenhouse Gas Inventories Programme, J. Penman, et al., eds. August 13, 2004. Available online at <<http://www.ipcc-nggip.iges.or.jp/public/gpplulucf/gpplulucf.htm>>.
- Johnson, D.M., and R. Mueller (2010) The 2009 Cropland Data Layer. Photogrammetric engineering and remote sensing 76:1201-1205.
- Kirstein, A. (2003 through 2004, 2006) Personal Communication. Arthur Kirstein, Coordinator, Agricultural Economic Development Program, Palm Beach County Cooperative Extension Service, FL and ICF International.
- Kraft, D.L. and H.C. Orender (1993) "Considerations for Using Sludge as a Fuel." *Tappi Journal*, 76(3): 175-183.
- Li, Y., D. Chen, Y. Zhang, R. Edis and H. Ding (2005) Comparison of three modeling approaches for simulating denitrification and nitrous oxide emissions from loam-textured arable soils. *Global Biogeochemical Cycles*, 19, GB3002.
- LSU (2015) Louisiana ratoon crop and conservation: Ratoon & Conservation Tillage Estimates. Louisiana State University, College of Agriculture AgCenter. Online at: [www.lsuagcenter.com](http://www.lsuagcenter.com)
- McGill, W.B., and C.V. Cole (1981) Comparative aspects of cycling of organic C, N, S and P through soil organic matter. *Geoderma* 26:267-286.
- Metherell, A.K., L.A. Harding, C.V. Cole, and W.J. Parton (1993) "CENTURY Soil Organic Matter Model Environment." Agroecosystem version 4.0. Technical documentation, GPSR Tech. Report No. 4, USDA/ARS, Ft. Collins, CO.
- Miller, M.R., Garr, J.D., and Coates, P.S. (2010) Changes in the status of harvested rice fields in the Sacramento Valley, California: Implications for wintering waterfowl. *Wetlands*, 30: 939-947.
- Miner, C. (1998) *Harvesting the High Plains: John Kriss and the business of wheat farming, 1920-1950*. University Press of Kansas, Lawrence, KS.
- Miner, R. (2008) "Calculations documenting the greenhouse gas emissions from the pulp and paper industry." Memorandum from Reid Miner, National Council for Air and Stream Improvement, Inc. (NCASI) to Becky Nicholson, RTI International, May 21, 2008.
- Mosier, A.R., Duxbury, J.M., Freney, J.R., Heinemeyer, O., and Minami, K. (1998) Assessing and mitigating N<sub>2</sub>O emissions from agricultural soils. *Climatic Change* 40:7-38.

- Nair, P.K.R. and V.D. Nair. (2003) Carbon storage in North American Agroforestry systems. In Kimble J., Heath L.S., Birdsey R.A., Lal R., editors. The potential of U.S. forest soils to sequester carbon and mitigate the greenhouse effect. CRC Press. Boca Raton, FL, 333–346.
- NASS (2004) Agricultural Chemical Usage: 2003 Field Crops Summary. Report AgCh1(04)a, National Agricultural Statistics Service, U.S. Department of Agriculture. Available online at <<http://usda.mannlib.cornell.edu/reports/nassr/other/pcu-bb/agcs0504.pdf>>.
- NASS (1999) Agricultural Chemical Usage: 1998 Field Crops Summary. Report AgCh1(99). National Agricultural Statistics Service, U.S. Department of Agriculture. Available online at <<http://usda.mannlib.cornell.edu/reports/nassr/other/pcu-bb/agch0599.pdf>>.
- NASS (1992) Agricultural Chemical Usage: 1991 Field Crops Summary. Report AgCh1(92). National Agricultural Statistics Service, U.S. Department of Agriculture. Available online at <<http://usda.mannlib.cornell.edu/reports/nassr/other/pcu-bb/agch0392.txt>>.
- Nevison, C.D., (2000) Review of the IPCC methodology for estimating nitrous oxide emissions associated with agricultural leaching and runoff, *Chemosphere – Global Change Science* 2, 493-500.
- NRAES (1992) On-Farm Composting Handbook (NRAES-54). Natural Resource, Agriculture, and Engineering Service. Available online at <[http://compost.css.cornell.edu/OnFarmHandbook/onfarm\\_TOC.html](http://compost.css.cornell.edu/OnFarmHandbook/onfarm_TOC.html)>.
- NRCS (1997) “National Soil Survey Laboratory Characterization Data,” Digital Data, Natural Resources Conservation Service, U.S. Department of Agriculture. Lincoln, NE.
- NRCS (1981) Land Resource Regions and Major Land Resource Areas of the United States, USDA Agriculture Handbook 296, United States Department of Agriculture, Natural Resources Conservation Service, National Soil Survey Center, Lincoln, NE, pp. 156.
- NRIAI (2003) Regional Budget and Cost Information. U.S. Department of Agriculture, Natural Resources Conservation Service, Natural Resources Inventory and Analysis Institute. Available online at <http://www.economics.nrcs.usda.gov/care/budgets/index.html>
- Nusser, S.M., F.J. Breidt, and W.A. Fuller (1998) “Design and Estimation for Investigating the Dynamics of Natural Resources, *Ecological Applications*, 8:234-245.
- Nusser, S.M., J.J. Goebel (1997) The national resources inventory: a long term monitoring programme. *Environmental and Ecological Statistics*, 4, 181-204.
- Ogle, S.M., Woodall, C.W., Swan, A., Smith, J., and Wirth, T. (in preparation). Determining the Managed Land Base for Delineating Carbon Sources and Sinks in the United States. *Environmental Science and Policy*.
- Ogle, S.M., F.J. Breidt, M. Easter, S. Williams, K. Killian, and K. Paustian (2010) “Scale and uncertainty in modeled soil organic carbon stock changes for U.S. croplands using a process-based model.” *Global Change Biology* 16:810-822.
- Ogle, S.M., F.J. Breidt, M. Easter, S. Williams and K. Paustian. (2007) “Empirically-Based Uncertainty Associated with Modeling Carbon Sequestration Rates in Soils.” *Ecological Modeling* 205:453-463.
- Ogle, S.M., F.J. Breidt, and K. Paustian. (2006) “Bias and variance in model results due to spatial scaling of measurements for parameterization in regional assessments.” *Global Change Biology* 12:516-523.
- Ogle, S.M., M.D. Eve, F.J. Breidt, and K. Paustian (2003) “Uncertainty in estimating land use and management impacts on soil organic carbon storage for U.S. agroecosystems between 1982 and 1997.” *Global Change Biology* 9:1521-1542.
- Parton, W.J., D.S. Schimel, C.V. Cole, D.S. Ojima (1987) “Analysis of factors controlling soil organic matter levels in Great Plains grasslands.” *Soil Science Society of America Journal* 51:1173-1179.
- Parton, W. J., J. M. O. Scurlock, D. S. Ojima, T. G. Gilmanov, R. J. Scholes, D. S. Schimel, T. Kirchner, J.-C. Menaut, T. Seastedt, E. G. Moya, A. Kamnalrut, and J. I. Kinyamario (1993) Observations and modeling of biomass and soil organic matter dynamics for grassland biomes worldwide. *Global Biogeochemical Cycles* 7:785-809.
- Parton, W.J., D.S. Ojima, C.V. Cole, and D.S. Schimel (1994) “A General Model for Soil Organic Matter Dynamics: Sensitivity to litter chemistry, texture and management,” in *Quantitative Modeling of Soil Forming Processes*. Special Publication 39, Soil Science Society of America, Madison, WI, 147-167.

- Parton, W.J., M.D. Hartman, D.S. Ojima, and D.S. Schimel (1998) "DAYCENT: Its Land Surface Submodel: Description and Testing". *Glob. Planet. Chang.* 19: 35-48.
- Parton, W.J., E.A. Holland, S.J. Del Grosso, M.D. Hartman, R.E. Martin, A.R. Mosier, D.S. Ojima, and D.S. Schimel (2001) Generalized model for NO<sub>x</sub> and N<sub>2</sub>O emissions from soils. *Journal of Geophysical Research.* 106 (D15):17403-17420.
- Peer, R., S. Thorneloe, and D. Epperson (1993) "A Comparison of Methods for Estimating Global Methane Emissions from Landfills." *Chemosphere*, 26(1-4):387-400.
- Potter, C. S., J.T. Randerson, C.B. Fields, P.A. Matson, P.M. Vitousek, H.A. Mooney, and S.A. Klooster. (1993) "Terrestrial ecosystem production: a process model based on global satellite and surface data." *Global Biogeochemical Cycles* 7:811-841.
- Potter, C., S. Klooster, A. Huete, and V. Genovese (2007) Terrestrial carbon sinks for the United States predicted from MODIS satellite data and ecosystem modeling. *Earth Interactions* 11, Article No. 13, DOI 10.1175/EI228.1.
- PRISM Climate Group, Oregon State University, <<http://prism.oregonstate.edu>>, created 24 July 2015.
- Quam, V.C., J. Gardner, J.R. Brandle, and T.K. Boes (1992) Windbreaks in Sustainable Agricultural Systems. EC-91-1772. University of Nebraska Extension. Lincoln, NE.
- Ruddy B.C., D.L. Lorenz, and D.K. Mueller (2006) County-level estimates of nutrient inputs to the land surface of the conterminous United States, 1982-2001. Scientific Investigations Report 2006-5012. U.S. Department of the Interior.
- Saghafi, Abouna (2013) Estimation of fugitive emissions from open cut coal mining and measurable gas content, 13th Coal Operators' Conference, University of Wollongong, The Australian Institute of Mining and Metallurgy & Mine Managers Association of Australia, 2013, 306-313.
- Sass, R.L., F.M. Fisher, S.T. Lewis, M.F. Jund, and F.T. Turner (1994) "Methane emissions from rice fields: effect of soil texture." *Global Biogeochemical Cycles* 8:135-140.
- Savitzky, A., and M. J. E. Golay (1964) Smoothing and Differentiation of Data by Simplified Least Squares Procedures. *Analytical Chemistry* 36:1627-1639.
- Saxton, K.E., W.J. Rawls, J.S. Romberger, and R.I. Papendick (1986) "Estimating Generalized Soil-Water Characteristics From Texture." *Soil Sci. Soc. Am. J.* 50:1031-1036.
- Schueneman, T. (1997, 1999 through 2001) Personal Communication. Tom Schueneman, Agricultural Extension Agent, Palm Beach County, FL and ICF International.
- Smith, J. (2008) E-mail correspondence between Jean Kim, ICF, and Jim Smith, U.S. Forest Service, December 3, 2008.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. (2015) State Soil Geographic (STATSGO) Database for State. Available online at <<http://www.ncgc.nrcs.usda.gov/products/datasets/statsgo/index.html>>.
- Spencer, S., S.M. Ogle, F.J. Breidt, J. Goebel, and K. Paustian (2011) Designing a national soil carbon monitoring network to support climate change policy: a case example for U.S. agricultural lands. *Greenhouse Gas Management & Measurement* 1:167-178.
- Stehfest, E., and C. Müller (2004), Simulation of N<sub>2</sub>O emissions from a urine-affected pasture in New Zealand with the ecosystem model DayCent, *J. Geophys. Res.*, 109, D03109, doi:10.1029/2003JD004261.
- Strehler, A., and W. Stützel (1987) "Biomass Residues." In Hall, D.O. and Overend, R.P. (eds.). *Biomass*. John Wiley and Sons, Ltd. Chichester, UK.
- TAMU (2015) Texas Rice Crop Survey. Texas A&M AgriLIFE Research Center at Beaumont. Online at: <<https://beaumont.tamu.edu/>>.
- Towery, D. (2001) Personal Communication. Dan Towery regarding adjustments to the CTIC (1998) tillage data to reflect long-term trends, Conservation Technology Information Center, West Lafayette, IN, and Marlen Eve, National Resource Ecology Laboratory, Fort Collins, CO. February 2001.
- TVA (1992b) Fertilizer Summary Data 1992. Tennessee Valley Authority, Muscle Shoals, AL.
- TVA (1991 through 1992a, 1993 through 1994) Commercial Fertilizers. Tennessee Valley Authority, Muscle Shoals, AL.

- UCCE, 2015. Rice Production Manual. Revised 2015. University of California Cooperative Extension, Davis, in collaboration with the California Rice Research Board.
- USDA (2010a) Crop Production 2009 Summary, National Agricultural Statistics Service, Agricultural Statistics Board, U.S. Department of Agriculture, Washington, DC. Available online at <http://usda.mannlib.cornell.edu>.
- USDA (2015) Quick Stats: U.S. & All States Data - Crops. National Agricultural Statistics Service, U.S. Department of Agriculture. Washington, DC. U.S. Department of Agriculture, National Agricultural Statistics Service. Washington, D.C., Available online at <http://quickstats.nass.usda.gov/>
- USDA (2003, 2005 through 2006, 2008 through 2009) Crop Production Summary, National Agricultural Statistics Service, Agricultural Statistics Board, U.S. Department of Agriculture, Washington, DC. Available online at <http://usda.mannlib.cornell.edu>,
- USDA (1998) Field Crops Final Estimates 1992-1997. Statistical Bulletin Number 947a. National Agricultural Statistics Service, U.S. Department of Agriculture. Washington, DC. Available online at <http://usda.mannlib.cornell.edu>. Accessed July 2001.
- USDA (1996) Agricultural Waste Management Field Handbook, National Engineering Handbook (NEH), Part 651. Natural Resources Conservation Service, U.S. Department of Agriculture. July 1996.
- USDA (1994) Field Crops: Final Estimates, 1987-1992. Statistical Bulletin Number 896, National Agriculture Statistics Service, U.S. Department of Agriculture. Washington, DC. Available online at <http://usda.mannlib.cornell.edu/datasets/crops/94896/sb896.txt>.
- USDA (1991) *State Soil Geographic (STATSGO) Data Base Data use information*. Miscellaneous Publication Number 1492, National Soil Survey Center, Natural Resources Conservation Service, U.S. Department of Agriculture, Fort Worth, TX.
- USDA-ERS (2015) Agricultural Resource Management Survey (ARMS) Farm Financial and Crop Production Practices: Tailored Reports. Online at: <http://www.ers.usda.gov/data-products/arms-farm-financial-and-crop-production-practices/arms-data.aspx>.
- USDA-ERS (1997) Cropping Practices Survey Data—1995. Economic Research Service, United States Department of Agriculture. Available online at <http://www.ers.usda.gov/data/archive/93018/>.
- USDA-FSA (2014) Conservation Reserve Program Monthly Summary – September 2014. U.S. Department of Agriculture, Farm Service Agency, Washington, DC, Available online at <https://www.fsa.usda.gov/Assets/USDA-FSA-Public/usdfiles/Conservation/PDF/summarysept2014.pdf>.
- USDA-NASS (2015) Quick Stats: U.S. & All States Data; Crops; Production and Area Harvested; 1990 - 2014. National Agricultural Statistics Service, U.S. Department of Agriculture. Washington, D.C. U.S. Department of Agriculture, National Agricultural Statistics Service. Washington, D.C., Available online at: <http://quickstats.nass.usda.gov/>.
- USDA-NRCS (2015) Summary Report: 2012 National Resources Inventory, Natural Resources Conservation Service, Washington, D.C., and Center for Survey Statistics and Methodology, Iowa State University, Ames, Iowa. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcseprd396218.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcseprd396218.pdf).
- USDA-NRCS (2013) Summary Report: 2010 National Resources Inventory, Natural Resources Conservation Service, Washington, D.C., and Center for Survey Statistics and Methodology, Iowa State University, Ames, Iowa. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/stelprdb1167354.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb1167354.pdf).
- USFWS (2010) Strategic Plan: The Partners for Fish and Wildlife Program, Stewardship of Fish and Wildlife Through Voluntary Conservation. U.S. Fish and Wildlife Service, Washington, DC, USA. <http://www.fws.gov/partners/docs/783.pdf>
- Vogelman, J.E., S.M. Howard, L. Yang, C. R. Larson, B. K. Wylie, and J. N. Van Driel (2001) “Completion of the 1990’s National Land Cover Data Set for the conterminous United States.” Photogrammetric Engineering and Remote Sensing, 67:650-662.
- Way, M.O., McCauley, G.M., Zhou, X.G., Wilson, L.T., and Morace, B. (Eds.). (2014) 2014 Texas Rice Production Guidelines. Texas A&M AgriLIFE Research Center at Beaumont.
- Williams, S.A. (2006) Data compiled for the Consortium for Agricultural Soils Mitigation of Greenhouse Gases (CASMGs) from an unpublished manuscript. Natural Resource Ecology Laboratory, Colorado State University.

- Williams, S. and K. Paustian (2005) Developing Regional Cropping Histories for Century Model U.S.-level Simulations. Colorado State University, Natural Resources Ecology Laboratory, Fort Collins, CO.
- Wilson, C.E. Jr., and Branson, J.W. (2006) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2005. Norman, R.J., Meullenet, J.-F., and Moldenhauer, K.A.K., (Eds.). Research Series 540, Arkansas Agricultural Experiment Station, University of Arkansas.
- Wilson, C.E. Jr., and Branson, J.W. (2005) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2004. Norman, R.J., Meullenet, J.-F., and Moldenhauer, K.A.K., (Eds.). Research Series 529, Arkansas Agricultural Experiment Station, University of Arkansas.
- Wilson, C.E. Jr., and Runsick, S.K. (2008) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2007. Norman, R.J., Meullenet, J.-F., and Moldenhauer, K.A.K., (Eds.). Research Series 560, Arkansas Agricultural Experiment Station, University of Arkansas.
- Wilson, C.E. Jr., and Runsick, S.K. (2007) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2006. Norman, R.J., Meullenet, J.-F., and Moldenhauer, K.A.K., (Eds.). Research Series 550, Arkansas Agricultural Experiment Station, University of Arkansas.
- Wilson, C.E. Jr., Runsick, S.K., Mazzanti, R. (2009) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2008. Norman, R.J., Meullenet, J.-F., and Moldenhauer, K.A.K., (Eds.). Research Series 571, Arkansas Agricultural Experiment Station, University of Arkansas.
- Wilson, C.E. Jr., Runsick, S.K., and Mazzanti, R. (2010) Trends in Arkansas rice production. B.R. Wells Arkansas Rice Research Studies 2009. Norman, R.J., and Moldenhauer, K.A.K., (Eds.). Research Series 581, Arkansas Agricultural Experiment Station, University of Arkansas.
- Zomer RJ, Trabucco A, Bossio DA, van Straaten O, Verchot LV (2008) Climate Change Mitigation: A Spatial Analysis of Global Land Suitability for Clean Development Mechanism Afforestation and Reforestation. *Agric. Ecosystems and Envir.* 126: 67-80.
- Zomer RJ, Bossio DA, Trabucco A, Yuanjie L, Gupta DC & Singh VP (2007) Trees and Water: Smallholder Agroforestry on Irrigated Lands in Northern India. Colombo, Sri Lanka: International Water Management Institute. pp 45. (IWMI Research Report 122).

### **3.13. Methodology for Estimating Net Carbon Stock Changes in *Forest Land Remaining Forest Land* and *Land Converted to Forest Land***


This sub-annex expands on the methodology used to estimate net changes in carbon (C) stocks in forest ecosystems and harvested wood products for *Forest Land Remaining Forest Land* and *Land Converted to Forest Land* as well as non-CO<sub>2</sub> emissions from forest fires. Full details of the C conversion factors and procedures may be found in the cited references. For details on the methods used to estimate changes in soil C stocks in the *Land Converted to Forest Land* section please refer to Annex 3.12.

#### **Carbon stocks and net stock change in forest ecosystems**

The inventory-based methodologies for estimating forest C stocks are based on Woodall et al. (2015a) and are consistent with IPCC (2003, 2006) stock-difference methods. Estimates of ecosystem C are based on data from the a network of annual inventory plots established and measured by the Forest Inventory and Analysis program within the USDA Forest Service; either direct measurements or attributes of forest inventories are the basis for estimating metric tons of C per hectare in IPCC pools (i.e., above- and belowground biomass, dead wood, litter, and soil organic carbon). Plot-level estimates are used to inform land area (by use) and stand age transition matrices across time which can be summed annually for an estimate of forest C stock change for *Forest Land Remaining Forest Land* and *Land Converted to Forest Land*. Recent publications (Coulston et al. 2015; Woodall et al. 2015a) detail the land use and stand age transition matrices that are informed by the annual forest inventory of the U.S. and were used in the accounting framework used in this Inventory. The annual forest inventories in the eastern U.S. have been remeasured which allows for empirical estimation of forest C stock net change within the accounting framework. In contrast, as numerous western states have not yet been remeasured, theoretical age transition matrices have been developed (Figure A-22).

A new accounting approach (Woodall et al. 2015a) was used to in this Inventory to be compliant with IPCC (2006) and UNFCCC reporting requirements. The United States' prior approach of comparing spatially and temporally inconsistent forest inventories of the 1980's and 1990's to the nationally consistent inventories of the 2000's not only resulted in highly uncertain estimates of individual forest C pools (see standing dead trees, Woodall et al. 2012), but also had an inability to disaggregate C stock changes by *Forest Land Remaining Forest Land* and *Land Converted to Forest Land* across the entire baseline (i.e., 1990 to the present) (Woodall et al. 2015a). The following subsections of this annex will describe the accounting framework used this year (Figure A-22) including the methods for estimating individual pools of forest C in addition to the eastern versus western approach to informing land use and stand age transitions.

**Figure A-22: Flowchart of the inputs necessary in the accounting framework, including the methods for estimating individual pools of forest C in the eastern and western conterminous U.S. states and coastal Alaska**



Note: An empirical age class transition matrix was used in the Eastern U.S. while a theoretical age class transition matrix was used in the Western U.S.

## Forest Land Definition

The definition of forest land within the U.S. and used for this Inventory is defined in Oswalt et al. (2014) as “Land at least 120 feet (37 meters) wide and at least 1 acre (0.4 hectare) in size with at least 10 percent cover (or equivalent stocking) by live trees including land that formerly had such tree cover and that will be naturally or artificially regenerated. Trees are woody plants having a more or less erect perennial stem(s) capable of achieving at least 3 inches (7.6 cm) in diameter at breast height, or 5 inches (12.7 cm) diameter at root collar, and a height of 16.4 feet (5 meters) at maturity in situ. The definition here includes all areas recently having such conditions and currently regenerating or capable of attaining such condition in the near future. Forest land also includes transition zones, such as areas between forest and non-forest lands that have at least 10 percent cover (or equivalent stocking) with live trees and forest areas adjacent to urban and built-up lands. Unimproved roads and trails, streams, and clearings in forest areas are classified as forest if they are less than 120 feet (36.6 meters) wide or an acre (0.4 hectare) in size. Forest land does not include land that is predominantly under agricultural or urban land use.” Timberland is productive forest land, which is on unreserved land and is producing or capable of producing crops of industrial wood. This is an important subclass of forest land because timberland is the primary source of C incorporated into harvested wood products. Productivity for timberland is at a minimum rate of 20 cubic feet per acre (1.4 cubic meters per hectare) per year of industrial wood (Woudenberg and Farrenkopf 1995). There are about 205 million hectares of timberland in the conterminous U.S., which represents 80 percent of all forest lands over the same area (Oswalt et al. 2014).


## Forest Inventory Data

The estimates of forest C stocks are based on data from forest inventory surveys. Forest inventory data were obtained from the USDA Forest Service, Forest Inventory and Analysis (FIA) Program (Frayer and Furnival 1999; USDA Forest Service 2015a; USDA Forest Service 2015b). Forest Inventory and Analysis data include remote sensing information and a collection of measurements in the field at sample locations called plots. Tree measurements include diameter at breast height, height, and species. On a subset of plots, additional measurements or samples are taken of downed dead wood, litter, and soil attributes. The technical advances needed to estimate C stocks from these data are ongoing (Woodall et al. 2015a) with the latest research incorporated on an annual basis (see Domke et al. 2016). The field protocols are thoroughly documented and available for download from the USDA Forest Service (2015c). Bechtold and Patterson (2005) provide the estimation procedures for standard forest inventory results. The data are freely available for download at USDA Forest Service (2011b) as the FIA Database (FIADB) Version 6.0 (USDA Forest Service 2015b; USDA Forest Service 2015c); these data are the primary sources of forest inventory data used to estimate forest C stocks. In addition to the field sampling component, fine-scale remotely sensed imagery (National Agriculture Imagery Program; NAIP 2015; Woodall et al. 2015b) is used to assign the land use at each sample location which has a nominal spatial resolution (raster cell size) of 1 m<sup>2</sup>. Prior to field measurement of each year’s collection of annual plots due for measurement (i.e., panel), each sample location in the panel (i.e., systematic distribution of plots within each state each year) is photo-interpreted manually by a forester to determine land use. As annual forest inventories have only just begun in the U.S. territories and in Hawaii, there is an assumption that these areas account for a net C change of zero. Survey data are available for the temperate oceanic ecoregion of Alaska (southeast and south central). These inventory data are publicly available for 6 million hectares of forest land, and these inventoried lands, representing an estimated 12 percent of the total forest land in Alaska, contribute to the forest C stocks presented here. Agroforestry systems are also not currently accounted for in the U.S. Inventory, since they are not explicitly inventoried by either of the two primary national natural resource inventory programs: the FIA program of the USDA Forest Service and the National Resources Inventory (NRI) of the USDA Natural Resources Conservation Service (Perry et al. 2005). The majority of these tree-based practices do not meet the size and definitions for forests within each of these resource inventories.

A national plot design and annualized sampling (USDA Forest Service 2015a) were introduced by FIA with most new annual inventories beginning after 1998. These are the only forest inventories used in the current accounting framework and subsequently in this submission. These surveys involve the sampling of all forest land including reserved and lower productivity lands. Almost all states have annualized inventory data available with substantial remeasurement in the eastern US (Figure A-23). Annualized sampling means that a portion of plots throughout the state is sampled each year, with the goal of measuring all plots once every 5 to 10 years, depending on the region of the U.S. The full unique set of data with all measured plots, such that each plot has been measured one time, is called a cycle. Sampling is designed such that partial inventory cycles provide usable, unbiased samples of forest inventory within the state, but with higher sampling errors than the full cycle. After all plots have been measured once, the sequence continues with remeasurement of the first year’s plots, starting the next new cycle. Most eastern states have completed one or two cycles of the annualized inventories, and some western states have begun remeasuring with a second annual cycle. Annually updated estimates of forest C stocks are affected by the redundancy in the data used to generate the annual updates of C stock. For example, a typical annual inventory update for an eastern state will include new data from remeasurement on 20 percent of plots; data from the remaining 80 percent of plots is identical to that included in the previous year’s annual update. The interpretation and use

of the annual inventory data can affect trend estimates of C stocks and stock changes (e.g., estimates based on 60 percent of an inventory cycle will be different than estimates with a complete (100 percent) cycle). In general, the C stock and stock change calculations use annual inventory summaries (updates) with unique sets of plot-level data (that is, without redundant sets); the most-recent annual update (i.e., 2015) is the exception because it is included in stock change calculations in order to include the most recent available data for each state. The specific inventories used in this report are listed in Table A-255 and this list can be compared with the full set of summaries available for download (USDA Forest Service 2015b).

**Figure A-23: Annual FIA plots (remeasured and not remeasured) across the U.S. including coastal Alaska through the 2014 field season**



Note: Due to the vast number of plots (where land use is measured even if no forest is present) they appear as spatially contiguous when displayed at the scale and resolution presented in this figure.

It should be noted that as the FIA program explores expansion of its vegetation inventory beyond the forest land use to other land uses (e.g., woodlands and urban areas) subsequent inventory observations will need to be delineated between forest and other land uses as opposed to a strict forest land use inventory. The forest C estimates provided here represent C stocks and stock change on managed forest lands (IPCC 2006, see Section 6.1 Representation of the U.S. Land Base), which is how all forest lands are classified on the 48 conterminous states. However, Alaska is considered to have significant areas of both managed and unmanaged forest lands. A new model delineating managed versus unmanaged lands for the U.S. (Ogle et al. in preparation), and used in this Inventory, is consistent with the assumption of managed forest lands on the 48 states. However, the model of Ogle et al. (in preparation) identifies some of the forest land in south central and southeastern coastal Alaska as unmanaged; this is in contrast to past assumptions of “managed” for these forest lands included in the FIADB. Therefore, the estimates for coastal Alaska as included here reflect that adjustment, which effectively reduces the forest area included here by about 5 percent. A second modification to the use of the FIADB-defined forest land introduced this year is to identify plots that do not meet the height component of the definition of forestland (Coulston et al. in preparation). These plots were identified as “other wooded lands” (i.e., not forest land use) and were removed from forest estimates and classified as grassland.<sup>105</sup> Note that minor differences in identifying and classifying

<sup>105</sup> Efforts are underway to account for the C stock changes in these “other wooded lands” that have been reclassified as grassland. The estimates should be included in the 2017 submission.

woodland as “forest” versus “other wooded” exist between the current Resources Planning Act Assessment (RPA) data (Oswalt et al. 2014) and the FIADB (USDA Forest Service 2015b) due to a refined modelling approach developed specifically for this submission (Coulston et al. in preparation).

**Table A-255: Specific annual forest inventories by state used in development of forest C stock and stock change estimates**

| Remeasured Annual Plots |                          |                                | Split Annual Cycle Plots |                   |                   |
|-------------------------|--------------------------|--------------------------------|--------------------------|-------------------|-------------------|
| State                   | Time 1 Year Range        | Time 2 Year Range              | State                    | Time 1 Year Range | Time 2 Year Range |
| Alabama                 | 2001 - 2009              | 2006 - 2014                    | Alaska (Coastal)         | 2004 - 2008       | 2009 - 2013       |
| Arkansas                | 2000, 2002, 2004 - 2010  | 2009 - 2014                    | Arizona                  | 2004 - 2008       | 2009 - 2013       |
| Connecticut             | 2004 - 2008              | 2009 - 2013                    | California               | 2001 - 2005       | 2006 - 2010       |
| Delaware                | 2004 - 2008              | 2009 - 2013                    | Colorado                 | 2004 - 2008       | 2009 - 2013       |
| Florida                 | 2002 - 2004, 2006 - 2007 | 2009 - 2013                    | Idaho                    | 2004 - 2008       | 2009 - 2013       |
| Georgia                 | 1998 - 2009              | 2005 - 2007, 2009 - 2013       | Montana                  | 2004 - 2008       | 2009 - 2013       |
| Illinois                | 2004 - 2009              | 2009 - 2014                    | Nevada                   | 2004 - 2008       | 2009 - 2013       |
| Indiana                 | 2004 - 2008              | 2009 - 2013                    | New Mexico               | 1999              | 2005 - 2013       |
| Iowa                    | 2004 - 2009              | 2009 - 2014                    | Oklahoma (West)          | 2009 - 2010       | 2011 - 2013       |
| Kansas                  | 2004 - 2009              | 2009 - 2014                    | Oregon                   | 2001 - 2005       | 2006 - 2010       |
| Kentucky                | 2000 - 2009              | 2005 - 2006, 2008 - 2012       | Texas (West)             | 2004 - 2007       | 2008 - 2012       |
| Louisiana               | 2001 - 2005, 2008        | 2009 - 2013                    | Utah                     | 2004 - 2008       | 2009 - 2013       |
| Maine                   | 2004 - 2008              | 2009 - 2013                    | Washington               | 2002 - 2006       | 2007 - 2011       |
| Maryland                | 2004 - 2008              | 2009 - 2013                    | Wyoming                  | 2000              | 2011 - 2013       |
| Massachusetts           | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| Michigan                | 2004 - 2009              | 2009 - 2014                    |                          |                   |                   |
| Minnesota               | 2005 - 2009              | 2010 - 2014                    |                          |                   |                   |
| Mississippi             | 2006                     | 2009 - 2014                    |                          |                   |                   |
| Missouri                | 2004 - 2009              | 2009 - 2014                    |                          |                   |                   |
| Nebraska                | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| New Hampshire           | 2003 - 2008              | 2009 - 2013                    |                          |                   |                   |
| New Jersey              | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| New York                | 2003 - 2008              | 2009 - 2013                    |                          |                   |                   |
| North Carolina          | 2002 - 2007              | 2003, 2005 - 2007, 2009 - 2014 |                          |                   |                   |
| North Dakota            | 2004 - 2009              | 2009 - 2014                    |                          |                   |                   |
| Ohio                    | 2003 - 2008              | 2009 - 2013                    |                          |                   |                   |
| Oklahoma (East)         | 2008                     | 2010 - 2013                    |                          |                   |                   |
| Pennsylvania            | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| Rhode Island            | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| South Carolina          | 2002 - 2010              | 2007 - 2013                    |                          |                   |                   |
| South Dakota            | 2004 - 2009              | 2009 - 2014                    |                          |                   |                   |
| Tennessee               | 2000 - 2009              | 2005 - 2012                    |                          |                   |                   |
| Texas (East)            | 2002 - 2008              | 2005, 2007 - 2012              |                          |                   |                   |
| Vermont                 | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| Virginia                | 2002 - 2003, 2005 - 2010 | 2008 - 2013                    |                          |                   |                   |
| West Virginia           | 2004 - 2008              | 2009 - 2013                    |                          |                   |                   |
| Wisconsin               | 2004 - 2009              | 2009 - 2014                    |                          |                   |                   |

Note: Remeasured annual plots represent a complete inventory cycle between measurements of the same plots while split annual cycle plots represent a single inventory cycle of plots that are split where remeasurements have yet to occur.

## Estimating Forest Inventory Plot-Level C-Density

For each inventory plot in each state, field data from the FIA program are used alone or in combination with auxiliary information (e.g., climate, surficial geology, elevation) to predict C density for each IPCC pool (i.e., aboveground and belowground biomass, dead wood, litter, SOC). In the past, most of the conversion factors and models used for inventory-based forest C estimates (Smith et al. 2010; Heath et al. 2011) were initially developed as an offshoot of the forest C simulation model FORCARB (Heath et al. 2010). The conversion factors and model coefficients were usually categorized by region and forest type. Thus, region and type are specifically defined for each set of estimates. More recently, the coarse approaches of the past have been updated with empirical information regarding C attributes of individual forest C pools such as dead wood and litter (e.g., Domke et al. 2013 and Domke et al. 2016). Factors are applied to the forest inventory data at the scale of FIA inventory plots which are a systematic sample of all forests attributes and land uses within each state. The results are estimates of C density (T per hectare) for the various forest pools. Carbon density for live trees, standing dead trees, understory vegetation, downed dead wood, litter, and soil organic matter are estimated. All non-soil C pools except litter can be separated into aboveground and belowground components. The live tree and understory C pools are combined into the biomass pool in this inventory. Similarly, standing dead trees and downed dead wood are pooled as dead wood in this inventory. C stocks and fluxes for *Forest Land Remaining Forest Land* are reported in pools following IPCC (2006).

### Live tree C pools

Live tree C pools include aboveground and belowground (coarse root) biomass of live trees with diameter at diameter breast height (d.b.h.) of at least 2.54 cm at 1.37 m above the forest floor. Separate estimates are made for above- and below-ground biomass components. If inventory plots include data on individual trees, tree C is based on Woodall et al. (2011), which is also known as the component ratio method (CRM), and is a function of volume, species, diameter, and, in some regions, tree height and site quality. The estimated sound volume (i.e., after rotten/missing deductions) provided in the tree table of the FIADB is the principal input to the CRM biomass calculation for each tree (Woodall et al. 2011). The estimated volumes of wood and bark are converted to biomass based on the density of each. Additional components of the trees such as tops, branches, and coarse roots, are estimated according to adjusted component estimates from Jenkins et al. (2003). Live trees with d.b.h. of less than 12.7 cm do not have estimates of sound volume in the FIADB, and CRM biomass estimates follow a separate process (see Woodall et al. 2011 for details). An additional component of foliage, which was not explicitly included in Woodall et al. (2011), was added to each tree following the same CRM method. Carbon is estimated by multiplying the estimated oven-dry biomass by a C constant of 0.5 because biomass is 50 percent of dry weight (IPCC 2006). Further discussion and example calculations are provided in Woodall et al. 2011 and Domke et al. 2012.

### Understory vegetation

Understory vegetation is a minor component of total forest ecosystem biomass. Understory vegetation is defined as all biomass of undergrowth plants in a forest, including woody shrubs and trees less than one-inch d.b.h. In this inventory, it is assumed that 10 percent of understory C mass is belowground. This general root-to-shoot ratio (0.11) is near the lower range of temperate forest values provided in IPCC (2006) and was selected based on two general assumptions: ratios are likely to be lower for light-limited understory vegetation as compared with larger trees, and a greater proportion of all root mass will be less than 2 mm diameter.

Estimates of C density are based on information in Birdsey (1996), which was applied to FIA permanent plots. These were fit to the model:

$$\text{Ratio} = e^{(A - B \times \ln(\text{live tree C density}))} \quad (1)$$

In this model, the ratio is the ratio of understory C density (T C/ha) to live tree C density (above- and below-ground) according to Jenkins et al. (2003) and expressed in T C/ha. An additional coefficient is provided as a maximum ratio; that is, any estimate predicted from the model that is greater than the maximum ratio is set equal to the maximum ratio. A full set of coefficients are in Table A-256. Regions and forest types are the same classifications described in Smith et al. (2003). As an example, the basic calculation for understory C in aspen-birch forests in the Northeast is:

$$\text{Understory (T C/ha)} = (\text{live tree C density}) \times e^{(0.855 - 1.03 \times \ln(\text{tree C density}))} \quad (2)$$

This calculation is followed by three possible modifications. First, the maximum value for the ratio is set to 2.02 (see value in column “maximum ratio”); this also applies to stands with zero tree C, which is undefined in the above model. Second, the minimum ratio is set to 0.005 (Birdsey 1996). Third, nonstocked (i.e., currently lacking tree cover but still in the forest land use) and pinyon/juniper forest types (see Table A-256) are set to coefficient A, which is a C density (T C/ha) for these types only.

**Table A-256: Coefficients for Estimating the Ratio of C Density of Understory Vegetation (above- and belowground, T C/ha)<sup>a</sup> by Region and Forest Type**

| Region <sup>b</sup> | Forest Type <sup>b</sup> | A     | B     | Maximum ratio <sup>c</sup> |
|---------------------|--------------------------|-------|-------|----------------------------|
| NE                  | Aspen-Birch              | 0.855 | 1.032 | 2.023                      |
|                     | MBB/Other Hardwood       | 0.892 | 1.079 | 2.076                      |
|                     | Oak-Hickory              | 0.842 | 1.053 | 2.057                      |
|                     | Oak-Pine                 | 1.960 | 1.235 | 4.203                      |
|                     | Other Pine               | 2.149 | 1.268 | 4.191                      |
|                     | Spruce-Fir               | 0.825 | 1.121 | 2.140                      |
|                     | White-Red-Jack Pine      | 1.000 | 1.116 | 2.098                      |
|                     | Nonstocked               | 2.020 | 2.020 | 2.060                      |
| NLS                 | Aspen-Birch              | 0.777 | 1.018 | 2.023                      |
|                     | Lowland Hardwood         | 0.650 | 0.997 | 2.037                      |
|                     | Maple-Beech-Birch        | 0.863 | 1.120 | 2.129                      |
|                     | Oak-Hickory              | 0.965 | 1.091 | 2.072                      |
|                     | Pine                     | 0.740 | 1.014 | 2.046                      |
|                     | Spruce-Fir               | 1.656 | 1.318 | 2.136                      |
|                     | Nonstocked               | 1.928 | 1.928 | 2.117                      |
| NPS                 | Conifer                  | 1.189 | 1.190 | 2.114                      |
|                     | Lowland Hardwood         | 1.370 | 1.177 | 2.055                      |
|                     | Maple-Beech-Birch        | 1.126 | 1.201 | 2.130                      |
|                     | Oak-Hickory              | 1.139 | 1.138 | 2.072                      |
|                     | Oak-Pine                 | 2.014 | 1.215 | 4.185                      |
|                     | Nonstocked               | 2.052 | 2.052 | 2.072                      |
| PSW                 | Douglas-fir              | 2.084 | 1.201 | 4.626                      |
|                     | Fir-Spruce               | 1.983 | 1.268 | 4.806                      |
|                     | Hardwoods                | 1.571 | 1.038 | 4.745                      |
|                     | Other Conifer            | 4.032 | 1.785 | 4.768                      |
|                     | Pinyon-Juniper           | 4.430 | 4.430 | 4.820                      |
|                     | Redwood                  | 2.513 | 1.312 | 4.698                      |
|                     | Nonstocked               | 4.431 | 4.431 | 4.626                      |
| PWE                 | Douglas-fir              | 1.544 | 1.064 | 4.626                      |
|                     | Fir-Spruce               | 1.583 | 1.156 | 4.806                      |
|                     | Hardwoods                | 1.900 | 1.133 | 4.745                      |
|                     | Lodgepole Pine           | 1.790 | 1.257 | 4.823                      |
|                     | Pinyon-Juniper           | 2.708 | 2.708 | 4.820                      |
|                     | Ponderosa Pine           | 1.768 | 1.213 | 4.768                      |
|                     | Nonstocked               | 4.315 | 4.315 | 4.626                      |
| PWW                 | Douglas-fir              | 1.727 | 1.108 | 4.609                      |
|                     | Fir-Spruce               | 1.770 | 1.164 | 4.807                      |
|                     | Other Conifer            | 2.874 | 1.534 | 4.768                      |
|                     | Other Hardwoods          | 2.157 | 1.220 | 4.745                      |
|                     | Red Alder                | 2.094 | 1.230 | 4.745                      |
|                     | Western Hemlock          | 2.081 | 1.218 | 4.693                      |
|                     | Nonstocked               | 4.401 | 4.401 | 4.589                      |
| RMN                 | Douglas-fir              | 2.342 | 1.360 | 4.731                      |
|                     | Fir-Spruce               | 2.129 | 1.315 | 4.749                      |
|                     | Hardwoods                | 1.860 | 1.110 | 4.745                      |
|                     | Lodgepole Pine           | 2.571 | 1.500 | 4.773                      |
|                     | Other Conifer            | 2.614 | 1.518 | 4.821                      |
|                     | Pinyon-Juniper           | 2.708 | 2.708 | 4.820                      |
|                     | Ponderosa Pine           | 2.099 | 1.344 | 4.776                      |
|                     | Nonstocked               | 4.430 | 4.430 | 4.773                      |
| RMS                 | Douglas-fir              | 5.145 | 2.232 | 4.829                      |
|                     | Fir-Spruce               | 2.861 | 1.568 | 4.822                      |
|                     | Hardwoods                | 1.858 | 1.110 | 4.745                      |
|                     | Lodgepole Pine           | 3.305 | 1.737 | 4.797                      |
|                     | Other Conifer            | 2.134 | 1.382 | 4.821                      |
|                     | Pinyon-Juniper           | 2.757 | 2.757 | 4.820                      |
|                     | Ponderosa Pine           | 3.214 | 1.732 | 4.820                      |
|                     | Nonstocked               | 4.243 | 4.243 | 4.797                      |
| SC                  | Bottomland Hardwood      | 0.917 | 1.109 | 1.842                      |
|                     | Misc. Conifer            | 1.601 | 1.129 | 4.191                      |
|                     | Natural Pine             | 2.166 | 1.260 | 4.161                      |

|    |                     |       |       |       |
|----|---------------------|-------|-------|-------|
|    | Oak-Pine            | 1.903 | 1.190 | 4.173 |
|    | Planted Pine        | 1.489 | 1.037 | 4.124 |
|    | Upland Hardwood     | 2.089 | 1.235 | 4.170 |
|    | Nonstocked          | 4.044 | 4.044 | 4.170 |
|    | Bottomland Hardwood | 0.834 | 1.089 | 1.842 |
|    | Misc. Conifer       | 1.601 | 1.129 | 4.191 |
|    | Natural Pine        | 1.752 | 1.155 | 4.178 |
| SE | Oak-Pine            | 1.642 | 1.117 | 4.195 |
|    | Planted Pine        | 1.470 | 1.036 | 4.141 |
|    | Upland Hardwood     | 1.903 | 1.191 | 4.182 |
|    | Nonstocked          | 4.033 | 4.033 | 4.182 |

<sup>a</sup> Prediction of ratio of understory C to live tree C is based on the model:  $\text{Ratio} = \exp(A - B \times \ln(\text{tree\_carbon\_tph}))$ , where "ratio" is the ratio of understory C density to live tree (above-and below-ground) C density, and "tree\_carbon\_density" is live tree (above-and below-ground) C density in T C/ha. Note that this ratio is multiplied by tree C density on each plot to produce understory vegetation.

<sup>b</sup> Regions and types as defined in Smith et al. (2003).

<sup>c</sup> Maximum ratio: any estimate predicted from the model that is greater than the maximum ratio is set equal to the maximum ratio.

## Dead Wood

The standing dead tree estimates are primarily based on plot-level measurements (Domke et al. 2011; Woodall et al. 2011). This C pool includes aboveground and belowground (coarse root) mass and includes trees of at least 12.7 cm d.b.h. Calculations follow the basic CRM method applied to live trees (Woodall et al. 2011) with additional modifications to account for decay and structural loss. In addition to the lack of foliage, two characteristics of standing dead trees that can significantly affect C mass are decay, which affects density and thus specific C content (Domke et al. 2011; Harmon et al. 2011), and structural loss such as branches and bark (Domke et al. 2011). Dry weight to C mass conversion is by multiplying by 0.5.

Downed dead wood, inclusive of logging residue, are sampled on a subset of FIA plots. Despite a reduced sample intensity, a single down woody material population estimate (Woodall et al. 2010; Domke et al. 2013; Woodall et al. 2013) per state is now incorporated into these empirical downed dead wood estimates. Downed dead wood is defined as pieces of dead wood greater than 7.5 cm diameter, at transect intersection, that are not attached to live or standing dead trees. It also includes stumps and roots of harvested trees. Ratio estimates of downed dead wood to live tree biomass were developed using FORCARB2 simulations and applied at the plot level (Smith et al. 2004). Estimates for downed dead wood correspond to the region and forest type classifications described in Smith et al. (2003). A full set of ratios is provided in Table A-257. An additional component of downed dead wood is a regional average estimate of logging residue based on Smith et al. (2006) applied at the plot level. These are based on a regional average C density at age zero and first order decay; initial densities and decay coefficients are provided in Table A-258. These amounts are added to explicitly account for downed dead wood following harvest. The sum of these two components are then adjusted by the ratio of population totals; that is, the ratio of plot-based to modeled estimates (Domke et al. 2013). An example of this 3-part calculation for downed dead wood in a 25-year-old naturally regenerated loblolly pine forest with 82.99 T C/ha in live trees (Jenkins et al. 2003) in Louisiana is as follows:

First, an initial estimate from live tree C density and Table A-257 (SC, Natural Pine)

$$\text{C density} = 82.99 \times 0.068 = 5.67 \text{ (T C/ha)}$$

Second, an average logging residue from age and Table A-257 (SC, softwood)

$$\text{C density} = 5.5 \times e^{(-25/17.9)} = 1.37 \text{ (T C/ha)}$$

Third, adjust the sum by the downed dead wood ratio plot-to-model for Louisiana, which was  $27.6/31.1 = 0.886$

$$\text{C density} = (5.67 + 1.37) \times 0.886 = 6.24 \text{ (T C/ha)}$$

**Table A-257: Ratio for Estimating Downed Dead Wood by Region and Forest Type**

| Region <sup>a</sup> | Forest type <sup>a</sup> | Ratio <sup>b</sup> | Region (cont'd) | Forest type (cont'd) | Ratio <sup>b</sup> (cont'd) |
|---------------------|--------------------------|--------------------|-----------------|----------------------|-----------------------------|
| NE                  | Aspen-Birch              | 0.078              | PWW             | Douglas-fir          | 0.100                       |
|                     | MBB/Other Hardwood       | 0.071              |                 | Fir-Spruce           | 0.090                       |
|                     | Oak-Hickory              | 0.068              |                 | Other Conifer        | 0.073                       |
|                     | Oak-Pine                 | 0.061              |                 | Other Hardwoods      | 0.062                       |
|                     | Other Pine               | 0.065              |                 | Red Alder            | 0.095                       |
|                     | Spruce-Fir               | 0.092              |                 | Western Hemlock      | 0.099                       |
|                     | White-Red-Jack Pine      | 0.055              |                 | Nonstocked           | 0.020                       |

|     |                   |       |                 |                     |       |
|-----|-------------------|-------|-----------------|---------------------|-------|
| NLS | Nonstocked        | 0.019 | RMN             | Douglas-fir         | 0.062 |
|     | Aspen-Birch       | 0.081 |                 | Fir-Spruce          | 0.100 |
|     | Lowland Hardwood  | 0.061 |                 | Hardwoods           | 0.112 |
|     | Maple-Beech-Birch | 0.076 |                 | Lodgepole Pine      | 0.058 |
|     | Oak-Hickory       | 0.077 |                 | Other Conifer       | 0.060 |
|     | Pine              | 0.072 |                 | Pinyon-Juniper      | 0.030 |
|     | Spruce-Fir        | 0.087 |                 | Ponderosa Pine      | 0.087 |
|     | Nonstocked        | 0.027 |                 | Nonstocked          | 0.018 |
| NPS | Conifer           | 0.073 | RMS             | Douglas-fir         | 0.077 |
|     | Lowland Hardwood  | 0.069 |                 | Fir-Spruce          | 0.079 |
|     | Maple-Beech-Birch | 0.063 |                 | Hardwoods           | 0.064 |
|     | Oak-Hickory       | 0.068 |                 | Lodgepole Pine      | 0.098 |
|     | Oak-Pine          | 0.069 |                 | Other Conifer       | 0.060 |
|     | Nonstocked        | 0.026 |                 | Pinyon-Juniper      | 0.030 |
| PSW | Douglas-fir       | 0.091 | SC              | Ponderosa Pine      | 0.082 |
|     | Fir-Spruce        | 0.109 |                 | Nonstocked          | 0.020 |
|     | Hardwoods         | 0.042 |                 | Bottomland Hardwood | 0.063 |
|     | Other Conifer     | 0.100 |                 | Misc. Conifer       | 0.068 |
|     | Pinyon-Juniper    | 0.031 |                 | Natural Pine        | 0.068 |
|     | Redwood           | 0.108 |                 | Oak-Pine            | 0.072 |
|     | Nonstocked        | 0.022 |                 | Planted Pine        | 0.077 |
| PWE | Douglas-fir       | 0.103 | SE              | Upland Hardwood     | 0.067 |
|     | Fir-Spruce        | 0.106 |                 | Nonstocked          | 0.013 |
|     | Hardwoods         | 0.027 |                 | Bottomland Hardwood | 0.064 |
|     | Lodgepole Pine    | 0.093 |                 | Misc. Conifer       | 0.081 |
|     | Pinyon-Juniper    | 0.032 |                 | Natural Pine        | 0.081 |
|     | Ponderosa Pine    | 0.103 |                 | Oak-Pine            | 0.063 |
|     | Nonstocked        | 0.024 |                 | Planted Pine        | 0.075 |
|     |                   |       | Upland Hardwood | 0.059               |       |
|     |                   |       | Nonstocked      | 0.012               |       |

<sup>a</sup> Regions and types as defined in Smith et al. (2003).

<sup>b</sup> The ratio is multiplied by the live tree C density on a plot to produce downed dead wood C density (T C/ha).

**Table A-258: Coefficients for Estimating Logging Residue Component of Downed Dead Wood**

| Region <sup>a</sup> | Forest Type Group <sup>b</sup> |                             | Decay Coefficient |
|---------------------|--------------------------------|-----------------------------|-------------------|
|                     | (softwood/<br>hardwood)        | Initial C Density<br>(T/ha) |                   |
| Alaska              | hardwood                       | 6.9                         | 12.1              |
| Alaska              | softwood                       | 8.6                         | 32.3              |
| NE                  | hardwood                       | 13.9                        | 12.1              |
| NE                  | softwood                       | 12.1                        | 17.9              |
| NLS                 | hardwood                       | 9.1                         | 12.1              |
| NLS                 | softwood                       | 7.2                         | 17.9              |
| NPS                 | hardwood                       | 9.6                         | 12.1              |
| NPS                 | softwood                       | 6.4                         | 17.9              |
| PSW                 | hardwood                       | 9.8                         | 12.1              |
| PSW                 | softwood                       | 17.5                        | 32.3              |
| PWE                 | hardwood                       | 3.3                         | 12.1              |
| PWE                 | softwood                       | 9.5                         | 32.3              |
| PWW                 | hardwood                       | 18.1                        | 12.1              |
| PWW                 | softwood                       | 23.6                        | 32.3              |
| RMN                 | hardwood                       | 7.2                         | 43.5              |
| RMN                 | softwood                       | 9.0                         | 18.1              |
| RMS                 | hardwood                       | 5.1                         | 43.5              |
| RMS                 | softwood                       | 3.7                         | 18.1              |
| SC                  | hardwood                       | 4.2                         | 8.9               |
| SC                  | softwood                       | 5.5                         | 17.9              |
| SE                  | hardwood                       | 6.4                         | 8.9               |
| SE                  | softwood                       | 7.3                         | 17.9              |

<sup>a</sup> Regions are defined in Smith et al. (2003) with the addition of coastal Alaska.

<sup>b</sup> Forest types are according to majority hardwood or softwood species.

## Litter carbon

Carbon in the litter layer is currently sampled on a subset of the FIA plots. Litter C is the pool of organic C (including material known as duff, humus, and fine woody debris) above the mineral soil and includes woody fragments with diameters of up to 7.5 cm. Because litter attributes are only collected on a subset of FIA plots, a model was developed to predict C density based on plot/site attributes for plots that lacked litter information (Domke et al. 2016).

As the litter, or forest floor, estimates are an entirely new model this year, a more detailed overview of the methods is provided here. The first step in model development was to evaluate all relevant variables—those that may influence the formation, accumulation, and decay of forest floor organic matter—from annual inventories collected on FIADB plots (P2) using all available estimates of forest floor C ( $n = 4,530$ ) from the P3 plots (hereafter referred to as the research dataset) compiled from 2000 through 2014 (Domke et al. 2016).

Random forest, a machine learning tool (Domke et al. 2016), was used to evaluate the importance of all relevant forest floor C predictors available from P2 plots in the research dataset. Given many of the variables were not available due to regional differences in sampling protocols during periodic inventories, the objective was to reduce the random forest regression model to the minimum number of relevant predictors without substantial loss in explanatory power. The form of the full random forest model was:

$$P(FFC_{Full}) = f(lat, lon, elev, fortypgrp, above, ppt, tmax, gmi) + u \quad (3)$$

where: *lat* = latitude, *lon* = longitude, *elev* = elevation, *fortypgrp* = forest type group, *above* = aboveground live tree C (trees  $\geq 2.54$  cm dbh), *ppt* = mean annual precipitation, *tmax* = average maximum temperature, *gmi* = the ratio of precipitation to potential evapotranspiration, *u* = the uncertainty in the prediction resulting from the sample-based estimates of the model parameters and observed residual variability around this prediction.

For each replacement, *u* was independently and randomly generated from a  $N(0, \sigma)$  distribution with  $\sigma$  incorporating the variability from both sources. This process of randomly selecting and incorporating *u* may be considered an imputation. Each model prediction was replaced independently *m* times and *m* separate estimates were combined where  $m = 1,000$  in this analysis.

Due to data limitation in certain regions and inventory periods a series of reduced random forest regression models were used rather than replacing missing variables with imputation techniques in random forest. Database records used to compile estimates for this report were grouped by variable availability and the approaches described herein were applied to replace forest floor model predictions from Smith and Heath (2002). Forest floor C predictions are expressed in  $T \cdot ha^{-1}$ .

## Soil organic carbon

Soil organic carbon (SOC) is the largest terrestrial C sink, and management of this pool is a critical component of efforts to mitigate atmospheric C concentrations. In the U.S., SOC in forests is monitored by the national forest inventory conducted by the FIA program (O'Neill et al. 2005). In previous C inventory submissions, SOC predictions were based, in part, on a model using the State Soil Geographic (STATSGO) database compiled by the Natural Resources Conservation Service (NRCS) (Amichev and Glabraith 2004), hereafter referred to as the country-specific ( $CS_{soc}$ ) model. Estimates of forest SOC found in the STATSGO database may be based on expert opinion and/or lack systematic field observations, but these country-specific model predictions have been used in past C inventory submissions. The FIA program has been consistently measuring soil attributes as part of the inventory since 2001 and has amassed an extensive inventory of SOC in forest land in the conterminous U.S. and coastal Alaska (O'Neill et al. 2005). More than 5,000 profile observations of SOC on forest land from FIA and the International Soil Carbon Monitoring Network (ISCN 2015) were used to develop and implement a modeling framework used for UNFCCC reporting that includes site-, stand-, and climate-specific variables that yield predictions of SOC stocks and stock changes specific to forest land in the U.S. This section provides a summary of the methodology used to predict SOC for this report. A complete description of the approach is being written by Domke et al. (In prep.).

The data used to develop the new modeling framework to predict SOC on forest land came from the FIA program and the ISCN. Since 2001, the FIA program has collected soil samples on every 16<sup>th</sup> base intensity plot distributed approximately every 38,848 ha, where at least one forested condition exists (Woodall et al. 2010). On fully forested plots, mineral and organic soils were sampled adjacent to subplots 2, 3, and 4 by taking a single core at each location from two layers: 0 to 10.16 cm and 10.16 to 20.32 cm. The texture of each soil layer was estimated in the field, and physical and chemical properties were determined in the laboratory (U.S. Forest Service 2011). For this analysis, estimates of SOC from the FIA program were calculated following O'Neill et al. (2005):

$$\sum SOC_{FIA\_TOTAL} = C_i \cdot BD_i \cdot t_i \cdot ucf \quad (4)$$

Where  $\sum SOC_{FIA\_TOTAL}$  = total mass (Mg C ha<sup>-1</sup>) of the mineral and organic soil C over all *i*th layers,  $C_i$  = percent organic C in the *i*th layer,  $BD_i$  = bulk density calculated as weight per unit volume of soil (g·cm<sup>-3</sup>) at the *i*th soil layer,  $t_i$  = thickness (cm) of the *i*th soil layer (either 0 to 10.16 cm or 10.16 to 20.32 cm), and  $ucf$  = unit conversion factor (100).

The  $SOC_{FIA\_TOTAL}$  estimates from each plot were assigned by forest condition on each plot, resulting in 3,667 profiles with SOC layer observations at 0 to 10.16 and 10.16 to 20.32 cm depths. Since the U.S. has historically reported SOC estimates to a depth of 100 cm (Heath et al. 2011, USEPA 2015), ISCN data from forests in the U.S. were harmonized with the FIA soil layer observations to develop model functions of SOC by soil order to a depth of 100 cm. All observations used from the ISCN were contributed by the Natural Resources Conservation Service. A total of 16,504 soil layers from 2,037 profiles were used from ISCN land uses defined as deciduous, evergreen, or mixed forest. The FIA-ISCN harmonized dataset used for model selection and prediction included a total of 5,704 profiles with 23,838 layer observations at depths ranging from 0 to 1,148 cm.

The modeling framework developed to predict SOC for this report was built around strategic-level forest and soil inventory information and auxiliary variables available for all FIA plots in the U.S. The first phase of the new estimation approach involved fitting models using the midpoint of each soil layer from the harmonized dataset and SOC estimates at those midpoints. Several linear and nonlinear models were evaluated, and a log-log model provided the optimal fit to the harmonized data:

$$\log_{10} SOC_i = I + \log_{10} Depth \quad (5)$$

Where  $\log_{10} SOC_i$  = SOC density (Mg C ha<sup>-1</sup> cm depth<sup>-1</sup>) at the midpoint depth,  $I$  = intercept,  $\log_{10} Depth$  = profile midpoint depth (cm).

The model was validated by partitioning the complete harmonized dataset multiple times into training and testing groups and then repeating this step for each soil order to evaluate model performance by soil order. Extra sum of squares F tests were used to evaluate whether there were statistically significant differences between the model coefficients from the model fit to the complete harmonized dataset and models fit to subsets of the data by soil order. Model coefficients for each soil order were used to predict SOC for the 20.32 to 100 cm layer for all FIA plots with soil profile observations. Next, the SOC layer observations from the FIA and predictions over the 100 cm profile for each FIA plot were summed:

$$SOC_{100} = SOC_{FIA\_TOTAL} + SOC_{20-100} \quad (6)$$

Where  $SOC_{100}$  = total estimated SOC density from 0-100 cm for each forest condition with a soil sample in the FIA database,  $SOC_{FIA\_TOTAL}$  as previously defined in model (1),  $SOC_{20-100}$  = predicted SOC from 20.32 to 100 cm from model (2). Note that bias correction factors will be incorporated into the  $SOC_{20-100}$  predictions and evaluated in Domke et al. (In prep.) but were not included in the  $SOC_{100}$  estimates used in this analysis.

In the second phase of the modeling framework,  $SOC_{100}$  estimates for FIA plots were used to predict SOC for plots lacking  $SOC_{100}$  estimates using Random forests, a machine learning tool that uses bootstrap aggregating (i.e., bagging) to develop models to improve prediction (Breimen 2001). Random forests also relies on random variable selection to develop a forest of uncorrelated regression trees. These trees recognize the relationship between a dependent variable, in this case  $SOC_{100}$ , and a set of predictor variables. All relevant predictor variables—those that may influence the formation, accumulation, and loss of SOC—from annual inventories collected on all base intensity plots and auxiliary climate, soil, and topographic variables obtained from the PRISM climate group (Northwest Alliance 2015), Natural Resources Conservation Service (NRCS 2015), and U.S. Geological Survey (Danielson and Gesch 2011), respectively, were included in the RF analysis. Due to regional differences in sampling protocols, many of the predictor variables included in the RF variable selection process were not available for all base intensity plots. To avoid problems with data limitations, pruning was used

to reduce the RF models to the minimum number of relevant predictors (including both continuous and categorical variables) without substantial loss in explanatory power or increase in root mean squared error (RMSE). The general form of the full RF models were:

$$P(SOC) = f(lat, lon, elev, fortypgrp, ppt, tmax, gmi, order, surfgeo) + u \quad (7)$$

where *lat* = latitude, *lon* = longitude, *elev* = elevation, *fortypgrp* = forest type group, *ppt* = mean annual precipitation, *tmax* = average maximum temperature, *gmi* = the ratio of precipitation to potential evapotranspiration, *order* = soil order, *surfgeo* = surficial geological description, and *u* = the uncertainty in the prediction resulting from the sample-based estimates of the model parameters and observed residual variability around this prediction. For each replacement, *u* was independently and randomly generated from a  $N(0, \sigma)$  distribution, with  $\sigma$  incorporating the variability from both sources. This process of randomly selecting and incorporating *u* may be considered an imputation. Each model prediction was replaced independently *m* times, and *m* separate predictions were combined following Rubin (1987):

$$\hat{C} = \frac{1}{m} \sum_{k=1}^m \hat{C}^k \quad (8)$$

where  $\hat{C}$  is the estimate for the *k*th completion of the data set. In this analysis, *m* = 1000, which is markedly larger than the *m* recommended by Rubin (1987), but given the extremely high level of replacement in this study, it was deemed necessary (Bodner 2008).

## Moving the Annual Forest Inventory Backwards and Forwards in Time: Transition Matrices

The accounting framework used this year is fundamentally driven by the annual forest inventory system conducted by the FIA program of the U.S. Forest Service (2015a-d). Unfortunately, the annual inventory system does not extend into the 1990's and the periodic data are not consistent (e.g., different plot design) with the annual inventory necessitating the adoption of a system to "backcast" the annual C estimates. Likewise, forecasting the annual inventory can enable the monitoring of U.S. greenhouse gas emission reduction targets, however, that is an activity beyond the scope of this document. To facilitate the backcasting of the U.S. annual forest inventory C estimates, the accounting framework is comprised of a forest dynamics module (age transition matrices) and a land use dynamics module (land area transition matrices). The forest dynamics module assesses forest sequestration, forest aging, and disturbance effects (i.e., disturbances such as wind, fire, and floods identified by foresters on inventory plots). The land use dynamics module assesses C stock transfers associated with afforestation and deforestation (e.g., Woodall et al. 2015b). Both modules are developed from land use area statistics and C stock change or C stock transfer by age class. The required inputs are estimated from more than 625,000 forest and nonforest observations in the FIA national database (U.S. Forest Service 2015a-c). Model predictions for before or after the annual inventory period are constructed from the accounting framework using only the annual observations. This modeling framework includes opportunities for user-defined scenarios to evaluate the impacts of land use change and disturbance rates on future C stocks and stock changes. As annual forest inventories in the eastern U.S. have largely completed at least one cycle and been remeasured, age and area transition matrices can be empirically informed. In contrast, as annual inventories in western states are still undergoing their first complete cycle they are still in the process of being remeasured, and as a result theoretical transition matrices need to be developed.

Wear and Coulston (2015) and Coulston et al. (2015) provide the framework for the projection model. The overall objective is to estimate unmeasured historical changes and future changes in forest C consistent with annual forest inventory measurements. For most regions, forest conditions are observed at time  $t_0$  and at a subsequent time  $t_1 = t_0 + s$ , where *s* is the time step (time measured in years) and is indexed by discrete (5 year) forest age classes. The inventory from  $t_0$  is then backcasted to the year 1990 (on average about 16 years) and projected from  $t_1$  to 2016 (about 5 years for the next inventory report). This backcasting/projection approach requires simulating changes in the age-class distribution resulting from forest aging and disturbance events and then applying C density estimates for each age class. For the North, South (except for west Texas and west Oklahoma), and Rocky Mountains regions of the country, age class transition matrices are estimated from observed changes in age classes between  $t_0$  and  $t_1$ . In the remainder of the regions (Pacific Coast including Alaska, west Texas, and west Oklahoma), only one inventory was available ( $t_0$ ) so transition matrices were derived from theory but informed by the condition of the observed inventory to backcast from  $t_0$  to 1990 and project from  $t_0$  to 2016.

### Theoretical Age Transition Matrices

Without any mortality-inducing disturbance, a projection of forest conditions would proceed by increasing all forest ages by the length of the time step until all forest resided in a terminal age class where the forest is retained indefinitely (this is by assumption, where forest C per unit area reaches a stable maximum). For the most basic case, disturbances (e.g., wildfire or timber harvesting) can reset some of the forest to the first age class. Disturbance can also alter the age class in more subtle ways. If a portion of trees in a multiple-age forest dies, the trees comprising the average age calculation change, thereby shifting the average age higher or lower (generally by one age class).

With  $n$  age classes, the age transition matrix ( $\mathbf{T}$ ) is an  $n \times n$  matrix, and each element ( $\mathbf{T}_{qr}$ ) defines the proportion of forest area in class  $q$  transitioning to class  $r$  during the time step ( $s$ ). The values of the elements of  $\mathbf{T}$  depend on a number of factors, including forest disturbances such as harvests, fire, storms, and the value of  $s$ , especially relative to the span of the age classes. For example, holding area fixed, allowing for no mortality, defining the time step  $s$  equivalent to the span of age classes, and defining five age classes results in:

$$\mathbf{T} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{pmatrix} \quad (9)$$

where all forest area progresses to the next age class and forests within the terminal age class are retained forever. With this version of  $\mathbf{T}$ , after five time steps all forests would be in the terminal age class. Relaxing these assumptions changes the structure of  $\mathbf{T}$ . If all disturbances, including harvesting and fire, that result in stand regeneration are accounted for and stochastic elements in forest aging are allowed,  $\mathbf{T}$  defines a traditional Lefkovich matrix population model (e.g., Caswell 2001) and becomes:

$$\mathbf{T} = \begin{pmatrix} 1 - t_1 - d_1 & d_2 & d_3 & d_4 & d_5 \\ t_1 & 1 - t_2 - d_2 & 0 & 0 & 0 \\ 0 & t_2 & 1 - t_3 - d_3 & 0 & 0 \\ 0 & 0 & t_3 & 1 - t_4 - d_4 & 0 \\ 0 & 0 & 0 & t_4 & 1 - d_5 \end{pmatrix} \quad (10)$$

Where  $t_q$  is the proportion of forest of age class  $q$  transitioning to age class  $q+1$ ,  $d_q$  is the proportion of age class  $q$  that experiences a stand-replacing disturbance, and  $(1 - t_q - d_q)$  is the proportion retained within age class  $q$  ( $\mathbf{T}_{qr}$ ).

### Projections and Backcast for Pacific Coast, Rocky Mountains, West Texas, and West Oklahoma

Projections of forest C in the Pacific (including Alaska), Rocky Mountains, west Texas and west Oklahoma are based on a life stage model:

$$\Delta C_t = C_{t+m} - C_t = (\mathbf{F}_t \mathbf{T} - \mathbf{F}_t) \cdot \mathbf{Den} + \mathbf{L}_t \cdot \mathbf{Den} \quad (11)$$

In this framework  $\mathbf{T}$  is an age transition matrix that shifts the age distribution of the forest  $\mathbf{F}$ . The difference in forest area by age class between time  $t$  and  $t+s$  is  $\mathbf{F}_t \mathbf{T} - \mathbf{F}_t$ . This quantity is multiplied by C density by age class ( $\mathbf{Den}$ ) to estimate C stock change of forest remaining forest between  $t$  and  $t+s$ . Land use change is accounted for by the addition of  $\mathbf{L}_t \cdot \mathbf{Den}$ , where  $\mathbf{L}_t$  identifies the age distribution of net land shifts into or out of forests. A query of the forest inventory databases provides estimates of  $\mathbf{F}$  and  $\mathbf{Den}$ , while inventory observations and modeling assumptions are used to estimate  $\mathbf{T}$ . By expanding  $\mathbf{Den}$  to a matrix of C contained in all the constituent pools of forest carbon, projections for all pools are generated.

Land use change is incorporated as a  $1 \times n$  vector  $\mathbf{L}$ , with positive entries indicating increased forest area and negative entries indicating loss of forest area, which provides insights of net change only. Implementing a forest area change

requires some information and assumptions about the distribution of the change across age classes (the n dimension of  $\mathbf{L}$ ). In the eastern states, projections are based on the projection of observed gross area changes by age class. In western states, total forest area changes are applied using rules. When net gains are positive, the area is added to the youngest forest age class; when negative, area is subtracted from all age classes in proportion to the area in each age class category.

Backcasting forest C inventories generally involve the same concepts as forecasting. An initial age class distribution is shifted at regular time steps backwards through time, using a transition matrix ( $\mathbf{B}$ ):

$$\mathbf{F}_{t-s} = \mathbf{F}_t \cdot \mathbf{B} \quad (12)$$

$\mathbf{B}$  is constructed based on similar logic used for creating  $\mathbf{T}$ . The matrix cannot simply be derived as the inverse of  $\mathbf{T}$  ( $\mathbf{F}_{t-s} = \mathbf{F}_t \mathbf{T}^{-1}$ ) because of the accumulating final age class (i.e.,  $\mathbf{T}$  does not contain enough information to determine the proportion of the final age class derived from the n-1 age class and the proportion that is retained in age class n from the previous time step).<sup>106</sup> However,  $\mathbf{B}$  can be constructed using observed changes from the inventory and assumptions about transition/accumulation including nonstationary elements of the transition model:

$$\mathbf{B} = \begin{pmatrix} 1 - \sum_q d_q & b_2 & 0 & 0 & 0 \\ d_1 & 1 - b_2 & b_3 & 0 & 0 \\ d_2 & 0 & 1 - b_3 & b_4 & 0 \\ d_3 & 0 & 0 & 1 - b_4 & b_r \\ d_4 & 0 & 0 & 0 & 1 - b_r \end{pmatrix} \quad (13)$$

Forest area changes need to be accounted for in the backcasts as well:

$$\mathbf{F}_{t-s} = \mathbf{F}_t \mathbf{B} - \mathbf{L}_t \quad (14)$$

Where  $\mathbf{L}_t$  is the forest area change between  $t_1$  and  $t_0$  as previously defined.

In the Rocky Mountains, age class transition matrices were empirically derived from observed changes in age classes between  $t_0$  and  $t_1$ . The frequency of transitions was constructed between age classes observed at  $t_0$  and  $t_1$  to define  $\mathbf{T}$  and between age classes  $t_1$  and  $t_0$  to define  $\mathbf{B}$ . In the Pacific Coast region, including Alaska, west Texas, and west Oklahoma, the theoretical life-stage models described by matrices (9) and (10) were applied. The disturbance factors ( $d$ ) in both  $\mathbf{T}$  and  $\mathbf{B}$  are derived from the current inventory by assuming that the area of forest in age class 1 resulted from disturbance in the previous period, the area in age class 2 resulted from disturbance in the period before that, and so on. The source of disturbed forest was assumed to be proportional to the area of forest in each age class. For projections ( $\mathbf{T}$ ), the average of implied disturbance for the previous two periods was applied. For the backcast ( $\mathbf{B}$ ), we move the disturbance frequencies implied by the age class distribution for each time step. For areas with empirical transition matrices, change in forest area ( $\mathbf{L}_t$ ) was backcasted/projected using the change in forest area observed for the period  $t_0$  to  $t_1$ . In the Pacific, including Alaska, west Texas, and west Oklahoma, it was assumed that total forest land area remained constant for the time period examined.

### *Projections and Backcast for North, South, east Texas, and east Oklahoma*

For the eastern U.S. a full set of remeasured plots were available. When remeasured data are available, the previously described approach is extended to estimate change more directly; in this case  $\Delta C_t = F_t \cdot \delta C$  where  $\Delta C$  is net stock change by pool within the analysis area,  $F$  is as previously defined, and  $\delta C$  is an  $n \times cp$  matrix of per unit area forest C stock change per year by pool (cp) arrayed by forest age class. Inter-period forest C dynamics are previously described, and the age transition matrix ( $\mathbf{T}$ ) is estimated from the observed data directly. Forest C change at the end of the next period is defined as:  $\Delta C_{t+s} = F_t \cdot \mathbf{T} \cdot \delta C$ . Land use change and disturbances such as cutting, fire, weather, insects, and diseases were incorporated by generalizing to account for the change vectors and undisturbed forest remaining as undisturbed forest:

<sup>106</sup> Simulation experiments show that a population that evolves as a function of  $\mathbf{T}$  can be precisely backcast using  $\mathbf{T}^{-1}$ . However, applying the inverse to a population that is not consistent with the long-run outcomes of the transition model can result in projections of negative areas within some stage age classes.

$$\Delta C_{t+s} = \sum_{d \in L} (A_{td} \cdot T_d \cdot \delta C_d) \quad (15)$$

Where  $A_{td}$  = area by age class of each mutually exclusive land category in L which includes d disturbances at time t.

$L = (FF, NFF, FNF, F_{cut}, F_{fire}, F_{weather}, F_{id})$  where FF=undisturbed forest remaining as undisturbed forest, NFF=nonforest to forest conversion, FNF=forest to nonforest conversion,  $F_{cut}$ =cut forest remaining as forest,  $F_{fire}$ =forest remaining as forest disturbed by fire,  $F_{weather}$ =forest remaining as forest disturbed by weather, and  $F_{id}$ =forest remaining as forest disturbed by insects and diseases. In the case of land transfers (FNF and NFF),  $T_d$  is an  $n \times n$  identity matrix and  $\delta C_d$  is a C stock transfer rate by age. Paired measurements for all plots in the inventory provide direct estimates of all elements of  $\delta C$ ,  $T_d$ , and  $A_{td}$  matrices.

Projections are developed by specifying either  $F_{t+s}$  or  $A_{t+sd}$  for either a future or a past state. To move the system forward, T is specified so that the age transition probabilities are set up as the probability between a time 0 and a time 1 transition. To move the system backward, T is replaced by B so that the age transition probabilities are for transitions from time 1 to time 0. Forecasts were developed by assuming the observed land use transitions and disturbance rates would continue for the next 5 years. Backcasts were developed using a Markov Chain process for land use transitions, observed disturbance rates for fire, weather, and insects. Historical forest cutting was incorporated by using the relationship between the area of forest cutting estimated from the inventory plots and the volume of roundwood production from the Timber Products Output program (U.S. Forest Service 2015d). This relationship allowed for the modification of  $F_{cut}$  such that it followed trends described by Oswald et al. (2014).

## Carbon in Harvested Wood Products

Estimates of the Harvested Wood Product (HWP) contribution to forest C sinks and emissions (hereafter called “HWP Contribution”) are based on methods described in Skog (2008) using the WOODCARB II model and the U.S. forest products module (Ince et al. 2011). These methods are based on IPCC (2006) guidance for estimating HWP C. The 2006 IPCC Guidelines provide methods that allow Parties to report HWP Contribution using one of several different accounting approaches: production, stock change, and atmospheric flow, as well as a default method. The various approaches are described below. The approaches differ in how HWP Contribution is allocated based on production or consumption as well as what processes (atmospheric fluxes or stock changes) are emphasized.

- **Production approach:** Accounts for the net changes in C stocks in forests and in the wood products pool, but attributes both to the producing country.
- **Stock-change approach:** Accounts for changes in the product pool within the boundaries of the consuming country.
- **Atmospheric-flow approach:** Accounts for net emissions or removals of C to and from the atmosphere within national boundaries. Carbon removal due to forest growth is accounted for in the producing country while C emissions to the atmosphere from oxidation of wood products are accounted for in the consuming country.
- **Default approach:** Assumes no change in C stocks in HWP. IPCC (2006) requests that such an assumption be justified if this is how a Party is choosing to report.

The U.S. uses the production accounting approach (as in previous years) to report HWP Contribution (Table A-259). Annual estimates of change are calculated by tracking the additions to and removals from the pool of products held in end uses (i.e., products in use such as housing or publications) and the pool of products held in solid waste disposal sites (SWDS).

The annual HWP variables that were used to estimate HWP contribution using the production approach are:

- (1) annual change of C in wood and paper products in use in the U.S. and other countries where the wood came from trees harvested in the U.S., and
- (2) annual change of C in wood and paper products in SWDS in the U.S. and other countries where the wood came from trees harvested in the U.S.

The sum of these variables yield estimates for HWP contribution under the production accounting approach.

**Table A-259: Harvested Wood Products from Wood Harvested in the U.S.—Annual Additions of C to Stocks and Total Stocks Under the Production Approach (Parentheses Indicate Net C Sequestration (i.e., a Net Removal of C from the Atmosphere))**

| Year | Net C additions per year (MMT C per year) |                 |                  | Total C stocks (MMT C) |                 |                  |
|------|---|-----------------|------------------|------------------------|-----------------|------------------|
|      | Total                                     | Products in use | Products in SWDS | Total                  | Products in use | Products in SWDS |
|      |   | Total           | Total            |                        |                 |                  |
| 1990 | -34.0                                     | -18.8           | -15.2            | 1,897                  | 1,250           | 647              |
| 1991 | -33.8                                     | -17.4           | -16.4            | 1,931                  | 1,265           | 665              |
| 1992 | -33.0                                     | -18.0           | -15.1            | 1,964                  | 1,282           | 683              |
| 1993 | -33.5                                     | -17.5           | -16.0            | 1,997                  | 1,297           | 701              |
| 1994 | -32.6                                     | -17.3           | -15.4            | 2,031                  | 1,313           | 718              |
| 1995 | -30.9                                     | -16.6           | -14.3            | 2,064                  | 1,328           | 736              |
| 1996 | -32.2                                     | -17.3           | -14.9            | 2,094                  | 1,342           | 752              |
| 1997 | -31.2                                     | -17.8           | -13.5            | 2,127                  | 1,357           | 769              |
| 1998 | -32.6                                     | -18.4           | -14.2            | 2,158                  | 1,371           | 787              |
| 1999 | -30.9                                     | -18.0           | -12.9            | 2,190                  | 1,385           | 806              |
| 2000 | -25.7                                     | -16.8           | -8.9             | 2,221                  | 1,398           | 824              |
| 2001 | -26.8                                     | -17.2           | -9.5             | 2,247                  | 1,407           | 840              |
| 2002 | -25.6                                     | -16.2           | -9.4             | 2,274                  | 1,416           | 858              |
| 2003 | -28.3                                     | -16.3           | -12.0            | 2,299                  | 1,426           | 874              |
| 2004 | -28.0                                     | -16.3           | -11.7            | 2,328                  | 1,438           | 890              |
| 2005 | -29.3                                     | -17.3           | -12.1            | 2,356                  | 1,449           | 906              |
| 2006 | -28.0                                     | -17.4           | -10.6            | 2,385                  | 1,461           | 924              |
| 2007 | -20.9                                     | -17.1           | -3.8             | 2,413                  | 1,472           | 941              |
| 2008 | -14.8                                     | -16.7           | 1.9              | 2,434                  | 1,476           | 958              |
| 2009 | -25.2                                     | -17.2           | -8.0             | 2,449                  | 1,474           | 975              |
| 2010 | -25.9                                     | -17.6           | -8.3             | 2,474                  | 1,482           | 992              |
| 2011 | -27.0                                     | -17.9           | -9.0             | 2,500                  | 1,490           | 1,010            |
| 2012 | -28.2                                     | -18.3           | -9.9             | 2,527                  | 1,499           | 1,028            |
| 2013 | -29.4                                     | -18.6           | -10.8            | 2,555                  | 1,509           | 1,046            |
| 2014 | -30.6                                     | -19.0           | -11.7            | 2,584                  | 1,520           | 1,065            |
| 2015 | -   | -               | -                | 2,615                  | 1,531           | 1,084            |

- Not reported or zero

**Table A-260: Harvested Wood Products Sectoral Background Data for LULUCF—U.S. (Production Approach)**

| Inventory year | 1A  | 1B   | 2   |
|----------------|---|--|---|
|                | Annual Change in stock of HWP in use produced from domestic harvest | Annual Change in stock of HWP in SWDS produced from domestic harvest | HWP Contribution to AFOLU CO <sub>2</sub> emissions/ removals |
|                | $\Delta C_{HWP\ IU\ DH}$  | $\Delta C_{HWP\ SWDS\ DH}$   | kt CO <sub>2</sub> /yr  |
| 1990           | -15,162   | -18,845  | -124,692  |
| 1991           | -16,393   | -17,446  | -124,076  |
| 1992           | -15,059   | -17,971  | -121,110  |
| 1993           | -16,001   | -17,500  | -122,836  |
| 1994           | -15,361   | -17,269  | -119,644  |
| 1995           | -14,339   | -16,552  | -113,267  |
| 1996           | -14,864   | -17,294  | -117,911  |
| 1997           | -13,456   | -17,765  | -114,476  |
| 1998           | -14,176   | -18,384  | -119,389  |
| 1999           | -12,946   | -18,001  | -113,469  |
| 2000           | -8,877  | -16,817  | -94,213   |
| 2001           | -9,527  | -17,236  | -98,131   |
| 2002           | -9,414  | -16,198  | -93,910   |
| 2003           | -12,042   | -16,297  | -103,910  |
| 2004           | -11,674   | -16,315  | -102,626  |
| 2005           | -12,059   | -17,289  | -107,609  |
| 2006           | -10,604   | -17,408  | -102,711  |
| 2007           | -3,830  | -17,110  | -76,779   |
| 2008           | 1,888   | -16,663  | -54,175   |
| 2009           | -7,968  | -17,222  | -92,364   |
| 2010           | -8,286  | -17,578  | -94,835   |
| 2011           | -9,034  | -17,937  | -98,894   |
| 2012           | -9,919  | -18,289  | -103,431  |
| 2013           | -10,790   | -18,630  | -107,873  |
| 2014           | -11,650   | -18,965  | -112,254  |

Annual estimates of variables 1A and 1B were calculated by tracking the additions to and removals from the pool of products held in end uses (e.g., products in uses such as housing or publications) and the pool of products held in SWDS. The pools include products exported and held in other countries and the pools in the U.S. exclude products made from wood harvested in other countries. Solidwood products added to pools include lumber and panels. End-use categories for solidwood include single and multifamily housing, alteration and repair of housing, and other end uses. There is one product category and one end-use category for paper. Additions to and removals from pools are tracked beginning in 1900, with the exception that additions of softwood lumber to housing begins in 1800. Solidwood and paper product production and trade data are from USDA Forest Service and other sources (Hair and Ulrich 1963; Hair 1958; USDC Bureau of Census 1976; Ulrich, 1985, 1989; Steer 1948; AF&PA 2006a, 2006b; Howard 2003).

The rate of removals from products in use and the rate of decay of products in SWDS are specified by first order (exponential) decay curves with given half-lives (time at which half of amount placed in use will have been discarded from use). Half-lives for products in use, determined after calibration of the model to meet two criteria, are shown in Table A-261. The first criterion is that the WOODCARB II model estimate of C in houses standing in 2001 needed to match an independent estimate of C in housing based on U.S. Census and USDA Forest Service survey data. The second criterion is that the WOODCARB II model estimate of wood and paper being discarded to SWDS needed to match EPA estimates of discards over the period 1990 to 2000. This calibration influences the estimate of variable 2A. The calibration also determines the amounts going to SWDS. In addition, WOODCARB II landfill decay rates have been validated by making sure that estimates of methane emissions from landfills based on EPA data are reasonable in comparison to methane estimates based on WOODCARB II landfill decay rates. Decay parameters for products in SWDS are shown in Table A-262.

A key assumption for estimating these variables is that products exported from the U.S. and held in pools in other countries have the same half-lives for products in use, the same percentage of discarded products going to SWDS, and the same decay rates in SWDS. Summaries of net fluxes and stocks for harvested wood in products and SWDS are in Land Converted to Forest Land – Soil C Methods.

**Table A-261: Half-life of Solidwood and Paper Products in End-Uses**

| Parameter  | Value | Units |
|--|-------|-------|
| Half-life of wood in single family housing 1920 and before           | 78.0  | Years |
| Half-life of wood in single family housing 1920–1939                 | 78.0  | Years |
| Half-life of wood in single family housing 1940–1959                 | 80.0  | Years |
| Half-life of wood in single family housing 1960–1979                 | 81.9  | Years |
| Half-life of wood in single family housing 1980 +                    | 83.9  | Years |
| Ratio of multifamily half live to single family half life            | 0.61  |       |
| Ratio of repair and alterations half-life to single family half life | 0.30  |       |
| Half-life for other solidwood product in end uses                    | 38.0  | Years |
| Half-life of paper in end uses                                       | 2.54  | Years |

Source: Skog, K.E. (2008) "Sequestration of C in harvested wood products for the U.S." *Forest Products Journal* 58:56–72.

**Table A-262: Parameters Determining Decay of Wood and Paper in SWDS**

| Parameter   | Value | Units   |
|---|-------|---------|
| Percentage of wood and paper in dumps that is subject to decay    | 100   | Percent |
| Percentage of wood in landfills that is subject to decay          | 23    | Percent |
| Percentage of paper in landfills that is subject to decay         | 56    | Percent |
| Half-life of wood in landfills / dumps (portion subject to decay) | 29    | Years   |
| Half-life of paper in landfills/ dumps (portion subject to decay) | 14.5  | Years   |

Source: Skog, K.E. (2008) "Sequestration of C in harvested wood products for the U.S." *Forest Products Journal* 58:56–72

**Table A-263: Net CO<sub>2</sub> Flux from Forest Pools in *Forest Land Remaining Forest Land* and *Harvested Wood Pools* (MMT CO<sub>2</sub> Eq.)**

| Carbon Pool           | 1990           | 1991           | 1992           | 1993           | 1994           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|-----------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Forest</b>         | <b>(598.8)</b> | <b>(605.4)</b> | <b>(605.0)</b> | <b>(605.1)</b> | <b>(598.8)</b> | <b>(594.1)</b> | <b>(563.6)</b> | <b>(560.8)</b> | <b>(564.4)</b> | <b>(568.6)</b> | <b>(570.7)</b> | <b>(584.3)</b> | <b>(596.0)</b> | <b>(605.5)</b> | <b>(613.2)</b> | <b>(619.9)</b> | <b>(647.2)</b> | <b>(637.8)</b> | <b>(632.4)</b> | <b>(631.2)</b> | <b>(630.1)</b> |
| Aboveground Biomass   | (312.4)        | (312.3)        | (312.1)        | (312.0)        | (307.5)        | (311.3)        | (259.1)        | (264.5)        | (267.1)        | (269.5)        | (272.1)        | (310.3)        | (310.4)        | (316.4)        | (321.0)        | (324.5)        | (331.2)        | (329.4)        | (324.6)        | (323.5)        | (322.5)        |
| Belowground Biomass   | (66.6)         | (66.6)         | (66.5)         | (66.4)         | (65.5)         | (66.5)         | (54.3)         | (55.6)         | (56.0)         | (56.5)         | (57.0)         | (65.7)         | (65.4)         | (66.7)         | (67.6)         | (68.2)         | (69.6)         | (69.3)         | (68.2)         | (67.9)         | (67.6)         |
| Dead Wood             | (34.8)         | (36.3)         | (36.7)         | (37.0)         | (37.2)         | (39.9)         | (47.9)         | (37.7)         | (38.0)         | (38.6)         | (37.6)         | (44.0)         | (46.2)         | (46.7)         | (47.4)         | (48.2)         | (50.2)         | (52.9)         | (53.7)         | (53.9)         | (54.2)         |
| Litter                | (35.9)         | (35.3)         | (35.2)         | (35.1)         | (34.8)         | (34.4)         | (24.5)         | (26.2)         | (26.5)         | (26.8)         | (27.1)         | (28.5)         | (28.0)         | (28.6)         | (29.2)         | (29.0)         | (34.5)         | (33.9)         | (33.1)         | (32.9)         | (32.7)         |
| Soil Organic Carbon   | (149.2)        | (154.8)        | (154.5)        | (154.6)        | (153.8)        | (142.1)        | (177.8)        | (176.8)        | (176.8)        | (177.3)        | (176.8)        | (135.8)        | (145.9)        | (147.1)        | (148.1)        | (150.0)        | (161.7)        | (152.4)        | (152.8)        | (152.9)        | (153.1)        |
| <b>Harvested Wood</b> | <b>(124.7)</b> | <b>(124.1)</b> | <b>(121.1)</b> | <b>(122.8)</b> | <b>(119.6)</b> | <b>(113.3)</b> | <b>(94.2)</b>  | <b>(98.1)</b>  | <b>(93.9)</b>  | <b>(103.9)</b> | <b>(102.6)</b> | <b>(107.6)</b> | <b>(102.7)</b> | <b>(76.8)</b>  | <b>(54.2)</b>  | <b>(92.4)</b>  | <b>(94.8)</b>  | <b>(98.9)</b>  | <b>(103.4)</b> | <b>(107.9)</b> | <b>(112.3)</b> |
| Products in Use       | (55.6)         | (60.1)         | (55.2)         | (58.7)         | (56.3)         | (52.6)         | (32.6)         | (34.9)         | (34.5)         | (44.2)         | (42.8)         | (44.2)         | (38.9)         | (14.0)         | 6.9            | (29.2)         | (30.4)         | (33.1)         | (36.4)         | (39.6)         | (42.7)         |
| SWDS                  | (69.1)         | (64.0)         | (65.9)         | (64.2)         | (63.3)         | (60.7)         | (61.7)         | (63.2)         | (59.4)         | (59.8)         | (59.8)         | (63.4)         | (63.8)         | (62.7)         | (61.1)         | (63.1)         | (64.5)         | (65.8)         | (67.1)         | (68.3)         | (69.5)         |
| <b>Total Net Flux</b> | <b>(723.5)</b> | <b>(729.4)</b> | <b>(726.1)</b> | <b>(728.0)</b> | <b>(718.5)</b> | <b>(707.4)</b> | <b>(657.8)</b> | <b>(658.9)</b> | <b>(658.3)</b> | <b>(672.5)</b> | <b>(673.3)</b> | <b>(691.9)</b> | <b>(698.7)</b> | <b>(682.2)</b> | <b>(667.4)</b> | <b>(712.3)</b> | <b>(742.0)</b> | <b>(736.7)</b> | <b>(735.8)</b> | <b>(739.1)</b> | <b>(742.3)</b> |

**Table A-264: Net C Flux from Forest Pools in *Forest Land Remaining Forest Land* and *Harvested Wood Pools* (MMT C)**

| Carbon Pool           | 1990           | 1991           | 1992           | 1993           | 1994           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|-----------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Forest</b>         | <b>(163.3)</b> | <b>(165.1)</b> | <b>(165.0)</b> | <b>(165.0)</b> | <b>(163.3)</b> | <b>(162.0)</b> | <b>(153.7)</b> | <b>(152.9)</b> | <b>(153.9)</b> | <b>(155.1)</b> | <b>(155.6)</b> | <b>(159.3)</b> | <b>(162.5)</b> | <b>(165.1)</b> | <b>(167.2)</b> | <b>(169.1)</b> | <b>(176.5)</b> | <b>(173.9)</b> | <b>(172.5)</b> | <b>(172.2)</b> | <b>(171.8)</b> |
| Aboveground Biomass   | (85.2)         | (85.2)         | (85.1)         | (85.1)         | (83.9)         | (84.9)         | (70.7)         | (72.1)         | (72.8)         | (73.5)         | (74.2)         | (84.6)         | (84.7)         | (86.3)         | (87.5)         | (88.5)         | (90.3)         | (89.8)         | (88.5)         | (88.2)         | (88.0)         |
| Belowground Biomass   | (18.2)         | (18.2)         | (18.1)         | (18.1)         | (17.9)         | (18.1)         | (14.8)         | (15.2)         | (15.3)         | (15.4)         | (15.6)         | (17.9)         | (17.8)         | (18.2)         | (18.4)         | (18.6)         | (19.0)         | (18.9)         | (18.6)         | (18.5)         | (18.4)         |
| Dead Wood             | (9.5)          | (9.9)          | (10.0)         | (10.1)         | (10.2)         | (10.9)         | (13.1)         | (10.3)         | (10.4)         | (10.5)         | (10.2)         | (12.0)         | (12.6)         | (12.7)         | (12.9)         | (13.1)         | (13.7)         | (14.4)         | (14.6)         | (14.7)         | (14.8)         |
| Litter                | (9.8)          | (9.6)          | (9.6)          | (9.6)          | (9.5)          | (9.4)          | (6.7)          | (7.2)          | (7.2)          | (7.3)          | (7.4)          | (7.8)          | (7.6)          | (7.8)          | (8.0)          | (7.9)          | (9.4)          | (9.3)          | (9.0)          | (9.0)          | (8.9)          |
| Soil Organic Carbon   | (40.7)         | (42.2)         | (42.1)         | (42.2)         | (41.9)         | (38.8)         | (48.5)         | (48.2)         | (48.2)         | (48.3)         | (48.2)         | (37.0)         | (39.8)         | (40.1)         | (40.4)         | (40.9)         | (44.1)         | (41.6)         | (41.7)         | (41.7)         | (41.7)         |
| <b>Harvested Wood</b> | <b>(34.0)</b>  | <b>(33.8)</b>  | <b>(33.0)</b>  | <b>(33.5)</b>  | <b>(32.6)</b>  | <b>(30.9)</b>  | <b>(25.7)</b>  | <b>(26.8)</b>  | <b>(25.6)</b>  | <b>(28.3)</b>  | <b>(28.0)</b>  | <b>(29.3)</b>  | <b>(28.0)</b>  | <b>(20.9)</b>  | <b>(14.8)</b>  | <b>(25.2)</b>  | <b>(25.9)</b>  | <b>(27.0)</b>  | <b>(28.2)</b>  | <b>(29.4)</b>  | <b>(30.6)</b>  |
| Products in Use       | (15.2)         | (16.4)         | (15.1)         | (16.0)         | (15.4)         | (14.3)         | (8.9)          | (9.5)          | (9.4)          | (12.0)         | (11.7)         | (12.1)         | (10.6)         | (3.8)          | 1.9            | (8.0)          | (8.3)          | (9.0)          | (9.9)          | (10.8)         | (11.7)         |
| SWDS                  | (18.8)         | (17.4)         | (18.0)         | (17.5)         | (17.3)         | (16.6)         | (16.8)         | (17.2)         | (16.2)         | (16.3)         | (16.3)         | (17.3)         | (17.4)         | (17.1)         | (16.7)         | (17.2)         | (17.6)         | (17.9)         | (18.3)         | (18.6)         | (19.0)         |
| <b>Total Net Flux</b> | <b>(197.3)</b> | <b>(198.9)</b> | <b>(198.0)</b> | <b>(198.5)</b> | <b>(196.0)</b> | <b>(192.9)</b> | <b>(179.4)</b> | <b>(179.7)</b> | <b>(179.5)</b> | <b>(183.4)</b> | <b>(183.6)</b> | <b>(188.7)</b> | <b>(190.5)</b> | <b>(186.1)</b> | <b>(182.0)</b> | <b>(194.3)</b> | <b>(202.4)</b> | <b>(200.9)</b> | <b>(200.7)</b> | <b>(201.6)</b> | <b>(202.5)</b> |

**Table A-265: Forest area (1,000 ha) and C Stocks in *Forest Land Remaining Forest Land* and Harvested Wood Pools (MMT C)**

|                                  | 1990           | 1995           | 2000           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           | 2015           |
|----------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>Forest Area<br/>(1000 ha)</b> | <b>261,796</b> | <b>263,780</b> | <b>265,918</b> | <b>266,336</b> | <b>266,754</b> | <b>267,179</b> | <b>267,604</b> | <b>268,029</b> | <b>268,454</b> | <b>268,874</b> | <b>269,271</b> | <b>269,668</b> | <b>270,065</b> | <b>270,462</b> | <b>270,871</b> | <b>271,280</b> | <b>271,719</b> | <b>272,158</b> |
| <b>Carbon Pools<br/>(MMT C)</b>  |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |                |
| <b>Forest</b>                    | <b>84,891</b>  | <b>85,713</b>  | <b>86,500</b>  | <b>86,653</b>  | <b>86,806</b>  | <b>86,960</b>  | <b>87,115</b>  | <b>87,271</b>  | <b>87,430</b>  | <b>87,593</b>  | <b>87,758</b>  | <b>87,925</b>  | <b>88,094</b>  | <b>88,271</b>  | <b>88,445</b>  | <b>88,617</b>  | <b>88,789</b>  | <b>88,961</b>  |
| Aboveground<br>Biomass           | 11,896         | 12,321         | 12,713         | 12,784         | 12,856         | 12,929         | 13,002         | 13,076         | 13,161         | 13,246         | 13,332         | 13,419         | 13,508         | 13,598         | 13,688         | 13,777         | 13,865         | 13,953         |
| Belowground<br>Biomass           | 2,442          | 2,532          | 2,615          | 2,630          | 2,645          | 2,660          | 2,676          | 2,691          | 2,709          | 2,727          | 2,745          | 2,764          | 2,782          | 2,801          | 2,820          | 2,839          | 2,857          | 2,876          |
| Dead Wood                        | 2,404          | 2,454          | 2,519          | 2,532          | 2,543          | 2,553          | 2,564          | 2,574          | 2,586          | 2,599          | 2,611          | 2,624          | 2,637          | 2,651          | 2,665          | 2,680          | 2,695          | 2,710          |
| Litter                           | 5,833          | 5,881          | 5,922          | 5,929          | 5,936          | 5,943          | 5,950          | 5,958          | 5,966          | 5,973          | 5,981          | 5,989          | 5,997          | 6,006          | 6,016          | 6,025          | 6,034          | 6,042          |
| Soil Organic<br>Carbon           | 62,316         | 62,525         | 62,730         | 62,779         | 62,827         | 62,875         | 62,923         | 62,972         | 63,009         | 63,048         | 63,089         | 63,129         | 63,170         | 63,214         | 63,255         | 63,297         | 63,339         | 63,381         |
| <b>Harvested Wood</b>            | <b>1,897</b>   | <b>2,064</b>   | <b>2,221</b>   | <b>2,247</b>   | <b>2,274</b>   | <b>2,299</b>   | <b>2,328</b>   | <b>2,356</b>   | <b>2,385</b>   | <b>2,413</b>   | <b>2,434</b>   | <b>2,449</b>   | <b>2,474</b>   | <b>2,500</b>   | <b>2,527</b>   | <b>2,555</b>   | <b>2,584</b>   | <b>2,615</b>   |
| Products in Use                  | 1,250          | 1,328          | 1,398          | 1,407          | 1,416          | 1,426          | 1,438          | 1,449          | 1,461          | 1,472          | 1,476          | 1,474          | 1,482          | 1,490          | 1,499          | 1,509          | 1,520          | 1,531          |
| SWDS                             | 647            | 736            | 824            | 840            | 858            | 874            | 890            | 906            | 924            | 941            | 958            | 975            | 992            | 1,010          | 1,028          | 1,046          | 1,065          | 1,084          |
| <b>Total Stock</b>               | <b>86,788</b>  | <b>87,777</b>  | <b>88,721</b>  | <b>88,900</b>  | <b>89,080</b>  | <b>89,260</b>  | <b>89,443</b>  | <b>89,627</b>  | <b>89,815</b>  | <b>90,006</b>  | <b>90,192</b>  | <b>90,374</b>  | <b>90,568</b>  | <b>90,771</b>  | <b>90,972</b>  | <b>91,172</b>  | <b>91,374</b>  | <b>91,576</b>  |

## ***Land Converted to Forest Land – Soil C Methods***

A Tier 2 method is applied to estimate soil C stock changes for *Land Converted to Forest Land* (Ogle et al. 2003, 2006; IPCC 2006). For this method, land is stratified by climate, soil types, land-use, and land management activity, and then assigned reference C levels and factors for the forest land and the previous land use. The difference between the stocks is reported as the stock change under the assumption that the change occurs over 20 years. Reference C stocks have been estimated from data in the National Soil Survey Characterization Database (USDA-NRCS 1997), and U.S.-specific stock change factors have been derived from published literature (Ogle et al. 2003; Ogle et al. 2006). Land use and land use change patterns are determined from a combination of the Forest Inventory and Analysis Dataset (FIA), the 2010 National Resources Inventory (NRI) (USDA-NRCS 2013), and National Land Cover Dataset (NLCD) (Homer et al. 2007). See Annex 3.12 for more information about this method (Methodology for Estimating N<sub>2</sub>O Emissions, CH<sub>4</sub> Emissions and Soil Organic C Stock Changes from Agricultural Soil Management).

Table A-266 summarizes the annual change in mineral soil C stocks from U.S. soils that were estimated using a Tier 2 method (Tg C/yr). The range is a 95 percent confidence interval from 50,000 simulations (Ogle et al. 2003, 2006). Table A-267 summarizes the total land areas by land use/land use change subcategory for mineral soils between 1990 and 2014 estimated with a Tier 2 approach and based on analysis of USDA National Resources Inventory data (USDA-NRCS 2013).

**Table A-266: Annual change in Mineral Soil C stocks from U.S. agricultural soils that were estimated using a Tier 2 method (Tg C/yr)**

| Category                                     | 1990                   | 1991                   | 1992                   | 1993                   | 1994                   | 1995                   | 1996                   | 1997                   | 1998                   | 1999                   |
|--|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| <b>Cropland Converted to Forest Land</b>     | 0.04<br>(0.02 to 0.07) | 0.05<br>(0.02 to 0.08) | 0.04<br>(0.01 to 0.08) | 0.05<br>(0.02 to 0.09) | 0.05<br>(0.02 to 0.09) | 0.06<br>(0.02 to 0.1)  | 0.06<br>(0.02 to 0.1)  | 0.06<br>(0.02 to 0.1)  | 0.06<br>(0.02 to 0.11) | 0.07<br>(0.02 to 0.12) |
| <b>Grassland Converted to Forest Land</b>    | 0.12<br>(0.06 to 0.19) | 0.13<br>(0.05 to 0.22) | 0.12<br>(0.03 to 0.21) | 0.14<br>(0.05 to 0.25) | 0.15<br>(0.05 to 0.26) | 0.17<br>(0.07 to 0.28) | 0.16<br>(0.06 to 0.27) | 0.17<br>(0.07 to 0.29) | 0.17<br>(0.05 to 0.29) | 0.18<br>(0.06 to 0.32) |
| <b>Other Lands Converted to Forest Land</b>  | 0.02<br>(0.01 to 0.03) | 0.02<br>(0.01 to 0.03) | 0.02<br>(0.01 to 0.03) | 0.02<br>(0.01 to 0.04) | 0.02<br>(0.01 to 0.04) | 0.03<br>(0.01 to 0.04) | 0.03<br>(0.01 to 0.04) | 0.03<br>(0.01 to 0.04) | 0.03<br>(0.01 to 0.05) | 0.03<br>(0.01 to 0.05) |
| <b>Settlements Converted to Forest Land</b>  | 0.00<br>(0 to 0)       | 0.00<br>(0 to 0.01)    |
| <b>Wetlands Converted to Forest Land</b>     | 0.00<br>(0 to 0)       | 0.00<br>(0 to 0.01)    | 0.01<br>(0 to 0.01)    | 0.01<br>(0 to 0.01)    | 0.01<br>(0 to 0.01)    | 0.01<br>(0 to 0.01)    |
| <b>Total Lands Converted to Forest Lands</b> | <b>0.18</b>            | <b>0.20</b>            | <b>0.19</b>            | <b>0.22</b>            | <b>0.23</b>            | <b>0.26</b>            | <b>0.25</b>            | <b>0.27</b>            | <b>0.27</b>            | <b>0.29</b>            |

Note: The range is a 95 percent confidence interval from 50,000 simulations (Ogle et al. 2003, 2006).

| Category                                    | 2000                   | 2001                   | 2002                   | 2003                   | 2004                   | 2005                   | 2006                   | 2007                   | 2008                   | 2009                   |
|---|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|
| <b>Cropland Converted to Forest Land</b>    | 0.06<br>(0.02 to 0.11) | 0.06<br>(0.02 to 0.11) | 0.05<br>(0.02 to 0.09) | 0.05<br>(0.01 to 0.1)  | 0.05<br>(0.01 to 0.1)  | 0.05<br>(0.01 to 0.09) | 0.02<br>(0 to 0.05)    | 0.02<br>(0.01 to 0.05) | 0.02<br>(0 to 0.04)    | 0.02<br>(0 to 0.04)    |
| <b>Grassland Converted to Forest Land</b>   | 0.17<br>(0.05 to 0.31) | 0.17<br>(0.05 to 0.31) | 0.15<br>(0.04 to 0.27) | 0.15<br>(0.04 to 0.28) | 0.14<br>(0.03 to 0.26) | 0.14<br>(0.03 to 0.26) | 0.07<br>(0.01 to 0.13) | 0.07<br>(0.02 to 0.14) | 0.07<br>(0.01 to 0.13) | 0.07<br>(0.01 to 0.13) |
| <b>Other Lands Converted to Forest Land</b> | 0.03<br>(0.01 to 0.05) | 0.03<br>(0.01 to 0.05) | 0.03<br>(0.01 to 0.05) | 0.03<br>(0.01 to 0.05) | 0.02<br>(0.01 to 0.05) | 0.03<br>(0.01 to 0.05) | 0.01<br>(0 to 0.02)    | 0.01<br>(0 to 0.02)    | 0.01<br>(0 to 0.02)    | 0.01<br>(0 to 0.02)    |
| <b>Settlements Converted to Forest Land</b> | 0.00                   | 0.00                   | 0.00                   | 0.00                   | 0.00                   | 0.00                   | 0.00                   | 0.00                   | 0.00                   | 0.00                   |

|  |             |             |             |             |             |             |             |             |             |             |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
|  | (0 to 0.01) | (0 to 0)    | (0 to 0)    | (0 to 0)    | (0 to 0)    |
| <b>Wetlands Converted to Forest Land</b>     | 0.01        | 0.01        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        | 0.00        |
|  | (0 to 0.01) | (0 to 0)    | (0 to 0)    | (0 to 0)    | (0 to 0)    |
| <b>Total Lands Converted to Forest Lands</b> | <b>0.27</b> | <b>0.27</b> | <b>0.24</b> | <b>0.24</b> | <b>0.23</b> | <b>0.22</b> | <b>0.11</b> | <b>0.11</b> | <b>0.10</b> | <b>0.10</b> |

Note: The range is a 95 percent confidence interval from 50,000 simulations (Ogle et al. 2003, 2006).

| <b>Category</b>                              | <b>2010</b>    | <b>2011</b>    | <b>2012</b> | <b>2013</b> | <b>2014</b> |
|--|----------------|----------------|-------------|-------------|-------------|
| <b>Cropland Converted to Forest Land</b>     | 0.02           | 0.02           | 0.02        | 0.02        | 0.02        |
|  | (0 to 0.04)    | (0 to 0.04)    | (0 to 0.04) | (0 to 0.04) | (0 to 0.04) |
| <b>Grassland Converted to Forest Land</b>    | 0.07           | 0.07           | 0.06        | 0.06        | 0.06        |
|  | (0.01 to 0.13) | (0.01 to 0.13) | (0 to 0.13) | (0 to 0.13) | (0 to 0.13) |
| <b>Other Lands Converted to Forest Land</b>  | 0.01           | 0.01           | 0.01        | 0.01        | 0.01        |
|  | (0 to 0.01)    | (0 to 0.01)    | (0 to 0.01) | (0 to 0.01) | (0 to 0.01) |
| <b>Settlements Converted to Forest Land</b>  | 0.00           | 0.00           | 0.00        | 0.00        | 0.00        |
|  | (0 to 0)       | (0 to 0)       | (0 to 0)    | (0 to 0)    | (0 to 0)    |
| <b>Wetlands Converted to Forest Land</b>     | 0.00           | 0.00           | 0.00        | 0.00        | 0.00        |
|  | (0 to 0)       | (0 to 0)       | (0 to 0)    | (0 to 0)    | (0 to 0)    |
| <b>Total Lands Converted to Forest Lands</b> | <b>0.10</b>    | <b>0.10</b>    | <b>0.10</b> | <b>0.10</b> | <b>0.10</b> |

Note: The range is a 95 percent confidence interval from 50,000 simulations (Ogle et al. 2003, 2006).

**Table A-267: Total land areas (hectares) by land use/land use change subcategory for mineral soils between 1990 and 2014**

| Conversion Land Areas (Hectares x 10 <sup>6</sup> ) | 1990        | 1991        | 1992        | 1993        | 1994        | 1995        | 2000        | 2001        | 2002        | 2003        | 2004        | 2005        | 2006        | 2007        | 2008        | 2009        | 2010        | 2011        | 2012        | 2013        | 2014        |
|---|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Cropland Converted to Forest Land                   | 0.22        | 0.21        | 0.21        | 0.22        | 0.21        | 0.25        | 0.23        | 0.23        | 0.23        | 0.23        | 0.24        | 0.23        | 0.22        | 0.21        | 0.21        | 0.21        | 0.21        | 0.21        | 0.20        | 0.20        | 0.20        |
| Grassland Converted to Forest Land                  | 0.58        | 0.59        | 0.59        | 0.62        | 0.64        | 0.70        | 0.65        | 0.66        | 0.66        | 0.64        | 0.64        | 0.64        | 0.65        | 0.63        | 0.64        | 0.65        | 0.65        | 0.65        | 0.66        | 0.66        | 0.66        |
| Other Lands Converted to Forest Land                | 0.09        | 0.09        | 0.09        | 0.10        | 0.10        | 0.11        | 0.11        | 0.11        | 0.11        | 0.11        | 0.11        | 0.12        | 0.12        | 0.08        | 0.08        | 0.07        | 0.07        | 0.07        | 0.07        | 0.07        | 0.07        |
| Settlements Converted to Forest Land                | 0.01        | 0.02        | 0.02        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.01        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        |
| Wetlands Converted to Forest Land                   | 0.01        | 0.01        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        | 0.02        |
| <b>Total Lands Converted to Forest Lands</b>        | <b>0.91</b> | <b>0.91</b> | <b>0.92</b> | <b>0.97</b> | <b>0.98</b> | <b>1.09</b> | <b>1.03</b> | <b>1.03</b> | <b>1.04</b> | <b>1.02</b> | <b>1.02</b> | <b>1.02</b> | <b>1.03</b> | <b>0.96</b> | <b>0.97</b> |

Note: Estimated with a Tier 2 approach and based on analysis of USDA National Resources Inventory data (USDA-NRCS 2013).

## Uncertainty Analysis

The uncertainty analyses for total net flux of forest C (see Table 6-14 in the FLRFL section) are consistent with the IPCC-recommended Tier 1 methodology (IPCC 2006). Specifically, they are considered approach 1 (propagation of error [Section 3.2.3.1]) (IPCC 2006). To better understand the effects of covariance, the contributions of sampling error and modeling error were parsed out. In addition, separate analyses were produced for forest ecosystem and HWP flux.

Estimates of forest C stocks in the U.S. are based on C estimates assigned to each of several thousand inventory plots from a regular grid. Uncertainty in these estimates and uncertainty associated with change estimates arise from many sources including sampling error and modeling error. Here we focus on these two types of error but acknowledge several other sources of error are present in the overall stock and stock change estimates. In terms of sampling based uncertainty, Design based estimators described by Bechtold and Patterson (2005) were used to quantify the variance of C stock estimates. In this section we denote the estimate of C stock at time  $t$  as  $C_t$  and the variances of the estimate of C stock for time  $t$  as  $\text{Var}(C_t)$ . These calculations follow Bechtold and Patterson (2005). The variance of stock change is then:

$$\text{Var}(C_{t2}-C_{t1})=\text{Var}(C_{t2})+\text{Var}(C_{t1})-2\cdot\text{Cov}(C_{t2},C_{t1}) \quad (16)$$

The uncertainty of a stock estimate associated with sampling error is  $U(C_t)_s=\text{Var}(C_t)^{0.5}$ . The uncertainty of a stock changes estimate associated with sampling error is  $U(\Delta C)_s=\text{Var}(C_{t2}-C_{t1})^{0.5}$ .

Model-based uncertainty is important because the pool-level C models have error. The total modeling mean-squared error ( $\text{MSE}_m$ ) is approximately  $1,622 \text{ (Mg/ha)}^2$ . The percent modeling error at time  $t$  is

$$\%U(C_t)_m = 100 \cdot \text{MSE}_m / d_t \quad (17)$$

Where  $d_t$  is the total C stock density at time  $t$  calculated as  $C_t/A_t$  where  $A_t$  is the forest area at time  $t$ .

The uncertainty of  $C_t$  from modeling error is

$$U(C_t)_m = C_t \cdot \%U(C_t)_m / 100 \quad (18)$$

The model-based uncertainty with respect to stock change is then

$$U(\Delta C)_m = (U(C_{t1})_m + U(C_{t2})_m - 2 \cdot \text{Cov}(U(C_{t1}), U(C_{t2})))^{0.5} \quad (19)$$

The sampling and model based uncertainty are combined for an estimate of total uncertainty. We considered these sources of uncertainty independent and combined as follow for stock change for stock change ( $\Delta C$ ):

$$U(\Delta C) = (U(\Delta C)_m^2 + U(\Delta C)_s^2)^{0.5} \text{ and the 95 percent confidence bounds was } \pm 2 \cdot U(\Delta C) \quad (20)$$

The mean square error (MSE) of pool models was ( $\text{MSE}, [\text{Mg C/ha}]^2$ ): soil C (1,143.0), litter (78.0), live tree (259.6), dead trees (101.5), understory (0.9), down dead wood (38.9), total MSE (1,621.9).

Numerous assumptions were adopted for creation of the forest ecosystem uncertainty estimates. Potential pool error correlations were ignored. Given the magnitude of the MSE for soil, including correlation among pool error would not appreciably change the modeling error contribution. Modeling error correlation between time 1 and time 2 was assumed to be 1. Because the MSE was fixed over time we assumed a linear relationship dependent on either the measurements at two points in time or an interpolation of measurements to arrive at annual flux estimates. Error associated with interpolation to arrive at annual flux is not included.

Uncertainty about net C flux in HWP is based on Skog et al. (2004) and Skog (2008). Latin hypercube sampling is the basis for the HWP Monte Carlo simulation. Estimates of the HWP variables and HWP Contribution under the production approach are subject to many sources of uncertainty. An estimate of uncertainty is provided that evaluated the effect of uncertainty in 13 sources, including production and trade data and parameters used to make the estimate. Uncertain data and parameters include data on production and trade and factors to convert them to C, the census-based estimate of C in housing in 2001, the EPA estimate of wood and paper discarded to SWDS for 1990 to 2000, the limits on decay of wood and paper in SWDS, the decay rate (half-life) of wood and paper in SWDS, the proportion of products produced in the United States made with wood harvested in the United States, and the rate of storage of wood and paper C in other countries that came from U.S. harvest, compared to storage in the United States.

The uncertainty about HWP and forest ecosystem net C flux were combined and assumed to be additive. Typically when propagating error from two estimates the variances of the estimates are additive. However, the uncertainty around the HWP flux was approximated using a Monte Carlo approach which resulted in the lack of a variance estimate for HWP C flux. Therefore, we considered the uncertainty additive between the HWP sequestration and the *Forest Land Remaining Forest Land* sequestration. Further, we assumed there was no covariance between the two estimates which is plausible as the observations used to construct each estimate are independent.

## Emissions from Forest Fires

### CO<sub>2</sub> Emissions from Forest Fires

As stated in other sections, the forest inventory approach implicitly accounts for emissions due to disturbances. Net C stock change is estimated from successive C stock estimates. A disturbance, such as a forest fire, removes C from the forest. The inventory data, on which net C stock estimates are based, already reflects the C loss from such disturbances because only C remaining in the forest is estimated. Estimating the CO<sub>2</sub> emissions from a disturbance such as fire and adding those emissions to the net CO<sub>2</sub> change in forests would result in double-counting the loss from fire because the inventory data already reflect the loss. There is interest, however, in the size of the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions from disturbances such as fire. These estimated emissions from forest fires are based on IPCC (2006) methodology, which includes a combination of U.S. specific data on area burned and potential fuel available for combustion along with IPCC default combustion and emission factors.

Emissions were calculated following IPCC (2006) methodology, according to equation 2.27 of IPCC (2006, Volume 4, Chapter 2), which in general terms is:

$$\text{Emissions} = \text{Area burned} \times \text{Fuel available} \times \text{Combustion factor} \times \text{Emission factor} \times 10^{-3} \quad (21)$$

Where the estimate for emissions is in units of metric tons (MT), which is generally summarized as million metric tons (MMT) per year. Area burned is the annual total area of forest fire in hectares. Fuel available is the mass of fuel available for combustion in metric tons dry weight per hectare. Combustion factor is the proportion of fuel consumed by fire and is unitless. The emission factor is gram of emission (in this case CO<sub>2</sub>) per kilogram dry matter burnt, and the “10<sup>-3</sup>” balances units. The first two factors are based on datasets specific to U.S. forests, whereas the last two factors employ IPCC (2006) default values.

Area burned is based on annual area of forest fires according to Monitoring Trends in Burn Severity (MTBS) MTBS Data Summaries 2015; Eidenshink et al. 2007) dataset summaries,<sup>107</sup> which include fire data for all 49 states that are a part of these estimates. That is, the MTBS data used here include the 48 conterminous states as well as Alaska, including interior Alaska; but note that the fire data used are also reduced to only include managed land. Summary information includes fire identity, year, location, area burned, fire intensity, and other fire characteristics. In addition to forest fires, the MTBS data include all wildland and prescribed fires on other ecosystems such as grasslands and rangelands; the “forest fire” distinction is not included as a part of identifying information for each fire. An additional spatial dataset – National MTBS Burned Area Boundaries – provides information to locate fires.<sup>108</sup> These individual-fire boundary data were used to partition the area burned in each fire to forest versus non-forest.

The MTBS fire data records include land cover information from the National Land Cover (NLCD) dataset (Homer et al. 2015), which can be used to distinguish forest fires from other wildland fires within the MTBS data. However, the forest land cover of the NLCD data, including the 2011 land cover (Homer et al. 2015) provides an estimate of forest land that is approximately 20 percent lower than forest area identified by the forest inventory of the USDA Forest Service (USDA Forest Service 2015b, e.g., data as of 2 June 2015) for the conterminous U.S. This suggests that annual area of forest fires identified with the NLCD cover data may underestimate area of forest burned, but the difference between USDA Forest Service (2015) and Homer et al. (2015) for each individual fire, if any, is dependent on specific areas where the fires actually occur. As an alternative data source, forest area for conterminous U.S. and Alaska are defined by Ruefenacht et al. (2008). The forest area for the conterminous states representative of approximately 2002 is within 2 percent of the forest areas estimated for 1990 through 2014 in US EPA (2015). These data were used to partition the perimeter data to forest for each fire (that is, area of forest relative to entire area of the fire for each MTBS fire). We assume that while changes in forests have occurred both before and since the data for Ruefenacht et al. (2008) were compiled, changes in forest versus non-forest status on lands subject to wildfires are likely minimal enough to make this dataset appropriate for this use. In addition, the Alaska forest area was allocated to managed and unmanaged areas according to Ogle et al. (in preparation), as discussed in more detail above.

The burned area perimeter dataset also was used to identify Alaska fires that were co-located with the area of permanent inventory plots of the USDA Forest Service’s (2015b) forest inventory along the southern coastal portion of the state. The only MTBS-identified burned forest areas in Alaska that coincide with the Forest Service’s permanent plot

<sup>107</sup> See <<http://www.mtbs.gov/dataaccess.html>>.

<sup>108</sup> See <<http://www.mtbs.gov/dataaccess.html>>.

inventoried area were on the northern (or Cook Inlet) side of the Kenai Peninsula, which is generally identified as boreal forest. From this, all MTBS fires of interest identified in Alaska are considered boreal forests.

Estimates of fuel availability are based on plot level forest inventory data, which are summarized by state and applied to all fires within the respective states. Plot level C stocks are defined by C conversion factors applied to current USDA Forest Service inventory data (USDA Forest Service 2015b; US EPA 2015; Smith et al. 2010) and summarized by state. We assume that while changes in forests have occurred over the years since the 1990 start of the reporting interval, the current general range of plot level C densities as determined by forest types and stand structures can be used as a representation of the potential fuel availability over the forest lands of a given state. We use the current forest inventory data<sup>109</sup> and the distribution of metric tons dry matter per hectare as the inputs for fuel availability. Fuel estimated for wildfires included all aboveground biomass (live trees and understory) as well as standing dead trees, down dead wood, and forest floor litter; whereas, fuel estimated for prescribed fires was based on the non-living components only.

The combustion factor used here for temperate forests is 0.45 (see Table 2.6 Volume 4, Chapter 2 of IPCC 2006). Similarly, the emission factor is an IPCC (2006) default, which for CO<sub>2</sub> is 1,569 g CO<sub>2</sub> per kg dry matter of fuel (see Table 2.5 Volume 4, Chapter 2 of IPCC 2006). With the application of equation 2.27 of IPCC (2006, in Volume 4, Chapter 2) defaults were used for mass of fuel available for the Alaska estimates because of the very limited coverage of boreal forests in the available U.S. forest inventories (see Table 2.4 Volume 4, Chapter 2 of IPCC 2006). Note that the values used for Alaska (Table 2.4 of IPCC 2006) represent the product of fuel available and the combustion factor.

Table A-268 provides summary values of annual area burned, area identified as forest fire, and emissions calculated according to equation 2.27 of IPCC (2006, in Volume 4, Chapter 2). The emission factor for CO<sub>2</sub> from Table 2.5 Volume 4, Chapter 2 of IPCC (2006) is provided in Table A-269. Separate calculations were made for each wild and prescribed fire in each state for each year. The results as MT CO<sub>2</sub> were summed to the MMT CO<sub>2</sub> per year values represented in Table A-268, and C emitted per year (Table A-268 and Table A-271) was based on multiplying by the conversion factor 12/44 (IPCC 2006).

**Table A-268: Areas (Hectares) from Wildfire Statistics and Corresponding Estimates of C and CO<sub>2</sub> (MMT/yr) Emissions for Wildfires and Prescribed Fires<sup>a</sup>**

| Year | Conterminous 48 States - Wildfires |                              |                    |                                  | Alaska - Wildfires             |                              |                    |                                  | Prescribed Fires (all 49 states) |                              |                    |                                  |
|------|------------------------------------|------------------------------|--------------------|----------------------------------|--------------------------------|------------------------------|--------------------|----------------------------------|----------------------------------|------------------------------|--------------------|----------------------------------|
|      | Reported area burned (1000 ha)     | Forest area burned (1000 ha) | C emitted (MMT/yr) | CO <sub>2</sub> emitted (MMT/yr) | Reported area burned (1000 ha) | Forest area burned (1000 ha) | C emitted (MMT/yr) | CO <sub>2</sub> emitted (MMT/yr) | Reported area burned (1000 ha)   | Forest area burned (1000 ha) | C emitted (MMT/yr) | CO <sub>2</sub> emitted (MMT/yr) |
| 1990 | 464                                | 185                          | 5.8                | 21.3                             | 572                            | 304                          | 5.3                | 19.5                             | 10                               | 6                            | 0.0                | 0.2                              |
| 1991 | 549                                | 309                          | 9.3                | 34.2                             | 308                            | 186                          | 3.3                | 12.0                             | 12                               | 8                            | 0.1                | 0.2                              |
| 1992 | 598                                | 180                          | 5.5                | 20.3                             | 12                             | 8                            | 0.1                | 0.5                              | 5                                | 4                            | 0.0                | 0.1                              |
| 1993 | 422                                | 80                           | 1.2                | 4.4                              | 161                            | 114                          | 2.0                | 7.3                              | 4                                | 2                            | 0.0                | 0.0                              |
| 1994 | 1,305                              | 629                          | 19.1               | 70.2                             | 97                             | 68                           | 1.2                | 4.4                              | 16                               | 10                           | 0.1                | 0.2                              |
| 1995 | 544                                | 131                          | 2.5                | 9.0                              | 13                             | 10                           | 0.2                | 0.6                              | 16                               | 11                           | 0.1                | 0.2                              |
| 1996 | 2,047                              | 509                          | 13.0               | 47.7                             | 0                              | 0                            | 0.0                | 0.0                              | 25                               | 18                           | 0.1                | 0.5                              |
| 1997 | 233                                | 74                           | 1.6                | 6.0                              | 0                              | 0                            | 0.0                | 0.0                              | 40                               | 10                           | 0.1                | 0.3                              |
| 1998 | 648                                | 271                          | 6.4                | 23.5                             | 71                             | 51                           | 0.9                | 3.3                              | 52                               | 21                           | 0.1                | 0.5                              |
| 1999 | 1,743                              | 463                          | 13.6               | 49.9                             | 349                            | 164                          | 2.9                | 10.5                             | 103                              | 49                           | 0.3                | 1.1                              |
| 2000 | 2,266                              | 1,020                        | 26.1               | 95.8                             | 312                            | 161                          | 2.8                | 10.4                             | 83                               | 23                           | 0.2                | 0.6                              |
| 2001 | 1,113                              | 469                          | 13.8               | 50.6                             | 44                             | 40                           | 0.7                | 2.6                              | 56                               | 30                           | 0.2                | 0.8                              |
| 2002 | 1,594                              | 1,114                        | 29.2               | 107.0                            | 857                            | 636                          | 11.2               | 40.9                             | 30                               | 16                           | 0.1                | 0.5                              |
| 2003 | 1,453                              | 792                          | 20.4               | 74.7                             | 234                            | 157                          | 2.8                | 10.1                             | 19                               | 13                           | 0.1                | 0.3                              |
| 2004 | 491                                | 321                          | 6.5                | 23.8                             | 2,656                          | 1,963                        | 34.4               | 126.3                            | 62                               | 35                           | 0.2                | 0.9                              |
| 2005 | 1,730                              | 601                          | 11.7               | 42.9                             | 1,964                          | 1,254                        | 22.0               | 80.7                             | 107                              | 62                           | 0.3                | 1.3                              |
| 2006 | 3,609                              | 947                          | 26.5               | 97.1                             | 106                            | 81                           | 1.4                | 5.2                              | 111                              | 80                           | 0.6                | 2.1                              |
| 2007 | 3,208                              | 1,488                        | 40.3               | 147.9                            | 239                            | 81                           | 1.4                | 5.2                              | 158                              | 97                           | 0.6                | 2.2                              |
| 2008 | 1,614                              | 724                          | 23.9               | 87.7                             | 38                             | 17                           | 0.3                | 1.1                              | 322                              | 252                          | 1.6                | 6.0                              |
| 2009 | 1,488                              | 493                          | 9.6                | 35.0                             | 1,143                          | 683                          | 12.0               | 44.0                             | 412                              | 318                          | 2.1                | 7.8                              |
| 2010 | 582                                | 141                          | 3.3                | 12.2                             | 311                            | 174                          | 3.1                | 11.2                             | 771                              | 658                          | 5.0                | 18.4                             |
| 2011 | 3,166                              | 1,226                        | 20.2               | 73.9                             | 96                             | 55                           | 1.0                | 3.5                              | 998                              | 245                          | 1.6                | 5.9                              |
| 2012 | 3,429                              | 1,463                        | 36.5               | 133.7                            | 102                            | 42                           | 0.7                | 2.7                              | 151                              | 110                          | 0.8                | 2.9                              |

<sup>109</sup> Retrieved from <<http://apps.fs.fed.us/fiadb-downloads/datamart.html>> on June 2, 2015.

|                   |       |     |      |      |     |     |     |      |     |     |     |     |
|-------------------|-------|-----|------|------|-----|-----|-----|------|-----|-----|-----|-----|
| 2013              | 1,091 | 637 | 17.6 | 64.7 | 498 | 347 | 6.1 | 22.3 | 277 | 267 | 1.5 | 5.3 |
| 2014 <sup>b</sup> | 1,091 | 637 | 17.6 | 64.7 | 498 | 347 | 6.1 | 22.3 | 277 | 267 | 1.5 | 5.3 |

<sup>a</sup> These emissions have already been accounted for in the estimates of net annual changes in C stocks, which accounts for the amount sequestered minus any emissions, including the assumption that combusted wood may continue to decay through time.

<sup>b</sup> The data for 2014 were incomplete when these estimates were summarized; therefore 2013, the most recent available estimate, is applied to 2014.

**Table A-269: Emission Factors for Extra Tropical Forest Burning and 100-year GWP (AR4), or equivalence ratios, of CH<sub>4</sub> and N<sub>2</sub>O to CO<sub>2</sub>**

| Emission Factor (g per kg dry matter burned) <sup>a</sup> |       | Equivalence Ratios <sup>b</sup>     |     |
|---|-------|-------------------------------------|-----|
| CH <sub>4</sub>   | 4.70  | CH <sub>4</sub> to CO <sub>2</sub>  | 25  |
| N <sub>2</sub> O  | 0.26  | N <sub>2</sub> O to CO <sub>2</sub> | 298 |
| CO <sub>2</sub>   | 1,569 | CO <sub>2</sub> to CO <sub>2</sub>  | 1   |

<sup>a</sup> IPCC (2006)

<sup>b</sup> IPCC (2007)

The set of fire emissions estimates using MODIS imagery and post-fire observations developed for Alaska by Veraverbeke et al. (2015a) is used here to provide a comparison with the estimates developed here (i.e., Table A-16). The spatial Alaskan Fire Emissions Database (AKFED, Veraverbeke et al. 2015b) was partitioned to forest land based on both Ruefenacht et al. (2008) and Homer et al. (2015) as well as managed/unmanaged (Ogle et al. in preparation). The estimates of annual C emitted from fire are in Table A-270, which also includes the estimates for managed forest land (both wildland and prescribed) that underlie the values provided in Table A-268. Note that the values in the six rightmost columns effectively partition the C emissions estimates provided in Veraverbeke et al. (2015a, see Table 2). That is, Table A-270, column 2 provides the estimates developed for this Inventory while each of columns 3-5 and 6-8 sum to the emissions estimates of Veraverbeke et al. (2015a); the differences between the two sets are how they are partitioned according to forest land.

**Table A-270: Estimated C emissions (MMT/yr) for fire based on the AKFED, and partitioned to managed forest land in Alaska**

| Year <sup>a</sup>    | Managed forest land (Table A-14) <sup>b</sup> | Forest land based on Ruefenacht et al. (2008) |                       |                 | Forest land based on Homer et al. (2015) |                       |                 |
|----------------------|---|---|-----------------------|-----------------|--|-----------------------|-----------------|
|                      |   | Managed forest land                           | Unmanaged forest land | Non-forest land | Managed forest land                      | Unmanaged forest land | Non-forest land |
| C emitted (MMT/year) |   |   |                       |                 |  |                       |                 |
| 2001                 | 0.7   | 0.8   | 0.3                   | 0.0             | 0.1                                      | 0.0                   | 1.1             |
| 2002                 | 11.2  | 12.7  | 3.3                   | 0.8             | 1.5                                      | 0.4                   | 14.8            |
| 2003                 | 2.8   | 4.0   | 1.4                   | 0.0             | 0.6                                      | 0.2                   | 4.7             |
| 2004                 | 34.4  | 51.8  | 16.6                  | 1.0             | 7.0                                      | 2.5                   | 59.9            |
| 2005                 | 22.0  | 29.8  | 14.1                  | 1.7             | 4.1                                      | 1.9                   | 39.6            |
| 2006                 | 1.4   | 0.7   | 0.1                   | 0.0             | 0.1                                      | 0.0                   | 0.7             |
| 2007                 | 1.5   | 1.4   | 1.0                   | 2.9             | 0.3                                      | 0.1                   | 4.9             |
| 2008                 | 0.3   | 0.4   | 0.4                   | 0.1             | 0.1                                      | 0.0                   | 0.8             |
| 2009                 | 12.0  | 16.3  | 9.8                   | 0.2             | 1.5                                      | 0.7                   | 24.1            |
| 2010                 | 4.7   | 4.6   | 1.1                   | 0.3             | 0.7                                      | 0.1                   | 5.1             |
| 2011                 | 1.0   | 1.5   | 0.3                   | 0.1             | 0.8                                      | 0.2                   | 0.9             |
| 2012                 | 0.8   | 0.8   | 0.2                   | 0.2             | 0.4                                      | 0.2                   | 0.6             |
| 2013                 | 6.1   | 7.4   | 2.5                   | 0.3             | 4.7                                      | 1.7                   | 3.7             |

<sup>a</sup> The AKFED data include the years 2001-2013 (Veraverbeke et al. 2015b).

<sup>b</sup> Values include both wildland and prescribed fires in Alaska.

## Non-CO<sub>2</sub> Emissions from Forest Fires

Emissions of non-CO<sub>2</sub> gasses – specifically, methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) – from forest fires are estimated using the same methodology described above (i.e., equation 2.27 of IPCC 2006, Volume 4, Chapter 2). The only difference in calculations is the gas-specific emission factors, which are listed in Table A-269. The summed annual estimates are provided in Table A-271. Conversion of the CH<sub>4</sub> and N<sub>2</sub>O estimates to CO<sub>2</sub> equivalents (as provided in Chapter 6-2) is based on global warming potentials (GWPs) provided in the IPCC Fourth Assessment Report (AR4) (IPCC 2007), which are the equivalence ratios listed in Table A-269. An example application of these ratios for the current year's estimate of CH<sub>4</sub> emissions is: 7.34 MMT CO<sub>2</sub> equiv. = 293,836 MT CH<sub>4</sub> × (25 kg CO<sub>2</sub> / 1 kg CH<sub>4</sub>) × 10<sup>-6</sup>.

Uncertainty about the non-CO<sub>2</sub> estimates is based on assigning a probability distribution to represent the estimated precision of each factor in equation 2.27 (of IPCC 2006). These probability distributions are randomly sampled with each calculation, and this is repeated a large number of times to produce a histogram, or frequency distribution of values for the calculated emissions. That is, a simple Monte Carlo (“Approach 2”) method was employed to propagate uncertainty in the equation (IPCC 2006). In general, probability densities are normal and also considered marginal distributions.

Estimates of burned forest area from the MTBS data (MTBS Data Summaries 2015; Ruefenacht et al. 2008; Ogle et al. in preparation) are assigned a normal distribution with relatively low uncertainty with a standard deviation of 4 percent, and these were sampled independently by year (Homer et al. 2015; Hao and Larkin 2014; Eidenshink et al. 2007). Fuel available is based on the distribution of plot level C densities (as metric tons dry matter per hectare) as defined within the current USDA Forest Service inventory data (USDA Forest Service 2015; US EPA 2015). We assume that current data adequately represent the general range of plot level C densities within a state’s forest land, given the limitations of the older inventory data as discussed elsewhere in this report. The plot-level C densities are summarized as dry weight densities (metric tons per hectare) for each plot with all aboveground dry weight summed as potential fuel for wildfires and all non-living components of aboveground dry weight assigned as potential fuel for prescribed fires. Frequency distributions of the plot data indicate that densities are distributed approximately lognormally. Each state’s data are fit to a lognormal distribution, and these were sampled independently by state and year. Note that each state has separate lognormal distributions for wild versus prescribed fire fuels, yet the same sampling sequence was used (i.e., jointly distributed within each state by year). Estimates for the Alaska fuel-by-combustion value as well as the combustion factor and emission factors are normal distributions with mean and standard deviations as defined in the tables (IPCC 2006 Tables 2.4, 2.5, and 2.6). These were sampled independently by year, and truncated to positive values where necessary. The equivalence ratios (Table A-269) to represent estimates as CO<sub>2</sub> equivalent were not considered uncertain values for these results.

**Table A-271: Estimated C Released and Estimates of Non-CO<sub>2</sub> Emissions (MMT/yr) for U.S. forests**

| Year | C emitted<br>(MMT/yr) | CH <sub>4</sub> emitted<br>(MMT/yr) | N <sub>2</sub> O<br>(MMT/yr) |
|------|-----------------------|-------------------------------------|------------------------------|
| 1990 | 11.2                  | 0.13                                | 0.007                        |
| 1991 | 12.6                  | 0.15                                | 0.008                        |
| 1992 | 5.7                   | 0.07                                | 0.004                        |
| 1993 | 3.2                   | 0.04                                | 0.002                        |
| 1994 | 20.4                  | 0.23                                | 0.013                        |
| 1995 | 2.7                   | 0.03                                | 0.002                        |
| 1996 | 13.2                  | 0.15                                | 0.008                        |
| 1997 | 1.7                   | 0.02                                | 0.001                        |
| 1998 | 7.4                   | 0.09                                | 0.005                        |
| 1999 | 16.8                  | 0.20                                | 0.011                        |
| 2000 | 29.1                  | 0.33                                | 0.018                        |
| 2001 | 14.7                  | 0.17                                | 0.010                        |
| 2002 | 40.5                  | 0.48                                | 0.026                        |
| 2003 | 23.2                  | 0.26                                | 0.015                        |
| 2004 | 41.2                  | 0.48                                | 0.027                        |
| 2005 | 34.1                  | 0.40                                | 0.022                        |
| 2006 | 28.5                  | 0.33                                | 0.018                        |
| 2007 | 42.4                  | 0.49                                | 0.027                        |
| 2008 | 25.8                  | 0.32                                | 0.018                        |
| 2009 | 23.7                  | 0.28                                | 0.016                        |
| 2010 | 11.4                  | 0.13                                | 0.007                        |
| 2011 | 22.7                  | 0.27                                | 0.015                        |
| 2012 | 38.0                  | 0.44                                | 0.024                        |
| 2013 | 25.2                  | 0.29                                | 0.016                        |

|                   |      |      |       |
|-------------------|------|------|-------|
| 2014 <sup>a</sup> | 25.2 | 0.29 | 0.016 |
|-------------------|------|------|-------|

<sup>a</sup> The data for 2014 were incomplete when these estimates were summarized; therefore 2013, the most recent available estimate, is applied to 2014.

## References

- AF&PA. (2006a and earlier). Statistical roundup. (Monthly). Washington, DC: American Forest & Paper Association.
- AF&PA. (2006b and earlier). Statistics of paper, paperboard and wood pulp. Washington, DC: American Forest & Paper Association.
- Amichev, B. Y. and J. M. Galbraith (2004) "A Revised Methodology for Estimation of Forest Soil Carbon from Spatial Soils and Forest Inventory Data Sets." *Environmental Management* 33(Suppl. 1):S74-S86.
- Bechtold, W.A.; Patterson, P.L. (2005) The enhanced forest inventory and analysis program—national sampling design and estimation procedures. Gen. Tech. Rep. SRS-80. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 85 p.
- Birdsey, R. (1996) "Carbon Storage for Major Forest Types and Regions in the Conterminous United States." In R.N. Sampson and D. Hair, (eds); *Forest and Global Change, Volume 2: Forest Management Opportunities for Mitigating Carbon Emissions*. American Forests. Washington, DC, 1-26 and 261-379 (appendices 262 and 263).
- Bodner, T.E. (2008) What improves with increased missing data imputations? *Structural Equation Modeling*. 15: 651-675.
- Breiman L. (2001) Random forests. *Machine Learning*. 45(1):5-32.
- Caswell, H. (2001) *Matrix population models*. Sunderland, MA: Sinauer Associates, Inc. 722 p.
- Coulston, J.W., Wear, D.N., and Vose, J.M. (2015) Complex forest dynamics indicate potential for slowing carbon accumulation in the southeastern United States. *Scientific Reports*. 5: 8002.
- Coulston, J.W. (In preparation). Tier 1 approaches to approximate the carbon implications of disturbances. On file with J.W. Coulston (jcoulston@fs.fed.us).
- Coulston, J.W., Woodall, C.W., Domke, G.M., and Walters, B.F. (in preparation). Refined Delineation between Woodlands and Forests with Implications for U.S. National Greenhouse Gas Inventory of Forests.
- Danielson J.J.; Gesch D.B. (2011) Global multi-resolution terrain elevation data 2010 (GMTED2010). Open-file report 2011-1073. Reston, VA: U.S. Department of the Interior, Geological Survey. 26 p.
- De Vos, B.; Cools, N.; Ilvesniemi, H.; Vesterdal, L.; Vanguelova, E.; Carnicelli, S. (2015) Benchmark values for forest soil carbon stocks in Europe: results from a large scale forest soil survey. *Geoderma*. 251: 33-46.
- Domke, G.M., Woodall, C.W., Smith, J.E., Westfall, J.A., McRoberts, R.E. (2012) Consequences of alternative tree-level biomass estimation procedures on U.S. forest carbon stock estimates. *Forest Ecology and Management*. 270: 108-116.
- Domke, G.M., Smith, J.E., and Woodall, C.W. (2011) Accounting for density reduction and structural loss in standing dead trees: Implications for forest biomass and carbon stock estimates in the United States. *Carbon Balance and Management*. 6:14.
- Domke, G.M., Woodall, C.W., Walters, B.F., Smith, J.E. (2013) From models to measurements: comparing down dead wood carbon stock estimates in the U.S. forest inventory. *PLoS ONE* 8(3): e59949.
- Domke, G.M., Perry, C.H., Walters, B.F., Woodall, C.W., and Smith, J.E. (2016). A framework for estimating litter carbon stocks in forests of the United States. *Science of the Total Environment*. 557-558, 469-478.
- Domke, G.M., Perry, C.H., Walters, B.F., Nave, L.E., Woodall, C.W., Swanston, C.W. (In Preparation) Estimating soil organic carbon in forest land of the United States.
- Eidenshink, J., B. Schwind, K. Brewer, Z. Zhu, B. Quayle, and S. Howard. (2007) A project for monitoring trends in burn severity. *Fire Ecology* 3(1): 3-21.
- Frayser, W.E., and G.M. Furnival (1999) "Forest Survey Sampling Designs: A History." *Journal of Forestry* 97(12): 4-10.
- Freed, R. (2004) Open-dump and Landfill timeline spreadsheet (unpublished). ICF International. Washington, D.C.
- Hair, D. and A.H. Ulrich (1963) The Demand and price situation for forest products – 1963. U.S. Department of Agriculture Forest Service, Misc Publication No. 953. Washington, DC.
- Hair, D. (1958) "Historical forestry statistics of the United States." *Statistical Bull.* 228. U.S. Department of Agriculture Forest Service, Washington, DC.

- Hao, W.M. and N.K. Larkin. (2014) Wildland fire emissions, carbon, and climate: Wildland fire detection and burned area in the United States. *Forest Ecology and Management* 317: 20–25.
- Harmon, M.E., C.W. Woodall, B. Fasth, J. Sexton, M. Yatkov. (2011) Differences between standing and downed dead tree wood density reduction factors: A comparison across decay classes and tree species. Res. Paper. NRS-15. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 40 p.
- Heath, L.S., M.C. Nichols, J.E. Smith, and J.R. Mills. (2010) FORCARB2: An updated version of the U.S. Forest Carbon Budget Model. Gen. Tech. Rep. NRS-67. USDA Forest Service, Northern Research Station, Newtown Square, PA. 52 p. [CD-ROM].
- Heath, L.S., J.E. Smith, K.E. Skog, D.J. Nowak, and C.W. Woodall. (2011) Managed forest carbon estimates for the U.S. Greenhouse Gas Inventory, 1990-2008. *Journal of Forestry* 109(3):167-173.
- Homer, C.G., J.A. Dewitz, L. Yang, S. Jin, P. Danielson, G. Xian, J. Coulston, N.D. Herold, J.D. Wickham, and K. Megown. (2015) Completion of the 2011 National Land Cover Database for the conterminous United States- Representing a decade of land cover change information. *Photogrammetric Engineering and Remote Sensing* 81(5): 345-354.
- Homer, C., Dewitz, J., Fry, J., Coan, M., Hossain, N., Larson, C., Herold, N., McKerrow, A., VanDriel, J.N., and Wickham, J. (2007) Completion of the 2001 National Land Cover Database for the Conterminous United States. *Photogrammetric Engineering and Remote Sensing*, Vol. 73, No. 4, pp 337-341.
- Howard, James L. (2003) *U.S. timber production, trade, consumption, and price statistics 1965 to 2002*. Res. Pap. FPL-RP-615. Madison, WI: USDA, Forest Service, Forest Products Laboratory. Available online at <<http://www.fpl.fs.fed.us/documnts/fplrp/fplrp615/fplrp615.pdf>>.
- Ince, P.J., Kramp, A.D., Skog, K.E., Spelter, H.N. and Wear, D.N. (2011) U.S. Forest Products Module: a technical document supporting the forest service 2010 RPA assessment. Research Paper-Forest Products Laboratory, USDA Forest Service, (FPL-RP-662).
- IPCC (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change, H.S. Eggleston, L. Buendia, K. Miwa, T Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- IPCC (2003) *Good Practice Guidance for Land Use, Land-Use Change, and Forestry*. The Intergovernmental Panel on Climate Change, National Greenhouse Gas Inventories Programme, J. Penman, et al., eds. August 13, 2004. Available online at <<http://www.ipcc-nggip.iges.or.jp/public/gpplulucf/gpplulucf.htm>>.
- ISCN. (2015) International Soil Carbon Monitoring Network (<http://iscn.fluxdata.org/>) database.
- Jandl, R., Rodeghiero, M., Martinez, C., Cotrufo, M. F., Bampa, F., van Wesemael, B., Harrison, R.B., Guerrini, I.A., deB Richter Jr., D., Rustad, L., Lorenz, K., Chabbi, A., Miglietta, F. 2014. Current status, uncertainty and future needs in soil organic carbon monitoring. *Science of the Total Environment*, 468, 376-383.
- Jenkins, J.C., D.C. Chojnacky, L.S. Heath, and R.A. Birdsey (2003) “National-scale biomass estimators for United States tree species.” *Forest Science* 49(1):12-35.
- Jobbagy, E.G.; Jackson, R.B. (2000) The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications*. 10: 423-436.
- Lal, R. (2005) Forest soils and carbon sequestration. *Forest Ecology and Management*. 220(1): 242-258.
- MTBS Data Summaries. (2015) MTBS Project, data last revised April 2015. (USDA Forest Service/U.S. Geological Survey). Available online at <<http://mtbs.gov/data/search.html>>. Accessed 25 August 2015.
- NAIP. (2015) National Agriculture Imagery Program. U.S. Department of Agriculture, Washington, DC. <http://www.fsa.usda.gov/programs-and-services/aerial-photography/imagery-programs/naip-imagery/> <last accessed November 23, 2015>

- Natural Resources Conservation Service [NRCS]. (2015) Soil geography: Description of STATSGO2 database. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geo/?cid=nrcs142p2\\_053629](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geo/?cid=nrcs142p2_053629) (accessed October 6, 2015).
- Northwest Alliance for Computational Science and Engineering. (2015) PRISM Climate Data. Available at <http://prism.oregonstate.edu> (accessed October 6, 2015).
- Ogle, S.M., M.D. Eve, F.J. Breidt, and K. Paustian (2003) "Uncertainty in estimating land use and management impacts on soil organic carbon storage for U.S. agroecosystems between 1982 and 1997." *Global Change Biology* 9:1521-1542.
- Ogle, S.M., F.J. Breidt, and K. Paustian. (2006) "Bias and variance in model results due to spatial scaling of measurements for parameterization in regional assessments." *Global Change Biology* 12:516-523.
- Ogle, S.M., Woodall, C.W., Swan, A., Smith, J.E., Wirth, T. (In preparation) Determining the Managed Land Base for Delineating Carbon Sources and Sinks in the United States. *Environmental Science and Policy*.
- O'Neill, K.P., Amacher, M.C., Perry, C.H. (2005) Soils as an indicator of forest health: a guide to the collection, analysis, and interpretation of soil indicator data in the Forest Inventory and Analysis program. Gen. Tech. Rep. NC-258. St. Paul, MN: US Department of Agriculture, Forest Service, North Central Research Station. 53 p.
- Oswalt, S.N.; Smith, W.B; Miles, P.D.; Pugh, S.A. (2014) Forest Resources of the United States, 2012: a technical document supporting the Forest Service 2015 update of the RPA Assessment. Gen. Tech. Rep. WO-91. Washington, DC: U.S. Department of Agriculture, Forest Service, Washington Office. 218 p.
- Perry, C.H., C.W. Woodall, and M. Schoeneberger (2005) Inventorying trees in agricultural landscapes: towards an accounting of "working trees". In: "Moving Agroforestry into the Mainstream." *Proc. 9th N. Am. Agroforestry Conf.*, Brooks, K.N. and Ffolliott, P.F. (eds). 12-15 June 2005, Rochester, MN [CD-ROM]. Dept. of Forest Resources, Univ. Minnesota, St. Paul, MN, 12 p. Available online at <<http://cinram.umn.edu/afta2005/>> (verified 23 Sept 2006).
- Ruefenacht, B., M.V. Finco, M.D. Nelson, R. Czaplewski, E.H. Helmer, J.A. Blackard, G.R. Holden, A.J. Lister, D. Salajanu, D. Weyermann, K. Winterberger. 2008. Conterminous U.S. and Alaska Forest Type Mapping Using Forest Inventory and Analysis. USDA Forest Service - Forest Inventory and Analysis Program & Remote Sensing Applications Center. Available online at < [http://data.fs.usda.gov/geodata/rastergateway/forest\\_type/](http://data.fs.usda.gov/geodata/rastergateway/forest_type/)>. Accessed 8 September 2015.
- Skog, K.E., K. Pingoud, and J.E. Smith (2004) "A method countries can use to estimate changes in carbon stored in harvested wood products and the uncertainty of such estimates." *Environmental Management* 33(Suppl. 1):S65-S73.
- Skog, K.E. (2008) "Sequestration of Carbon in harvested wood products for the United States." *Forest Products Journal*, 58(6): 56-72.
- Smith, J.E., L.S. Heath, and M.C. Nichols (2010). U.S. Forest Carbon Calculation Tool User's Guide: Forestland Carbon Stocks and Net Annual Stock Change. General Technical Report NRS-13 revised, U.S. Department of Agriculture Forest Service, Northern Research Station.
- Smith, J.E., L.S. Heath, K.E. Skog, R.A. Birdsey. (2006) Methods for calculating forest ecosystem and harvested carbon with standard estimates for forest types of the United States. Gen. Tech. Rep. NE-343. U.S. Department of Agriculture, Forest Service, Northeastern Research Station. Newtown Square, PA.
- Smith, J.E., L. S. Heath, and P. B. Woodbury (2004) "How to estimate forest carbon for large areas from inventory data." *Journal of Forestry* 102:25-31.
- Smith, J.E., L. S. Heath, and J. C. Jenkins (2003) *Forest Volume-to-Biomass Models and Estimates of Mass for Live and Standing Dead Trees of U.S. Forests*. General Technical Report NE-298, USDA Forest Service, Northeastern Research Station, Newtown Square, PA.
- Smith, J.E., and L.S. Heath (2002) "A model of forest floor carbon mass for United States forest types." Res. Paper NE-722. USDA Forest Service, Northeastern Research Station, Newtown Square, PA.
- Steer, Henry B. (1948) *Lumber production in the United States*. Misc. Pub. 669, U.S. Department of Agriculture Forest Service. Washington, DC.
- Sun, O.J.; Campbell, J.; Law, B.E.; Wolf, V. (2004) Dynamics of carbon stocks in soils and detritus across chronosequences of different forest types in the Pacific Northwest, USA. *Global Change Biology*. 10(9): 1470-1481.

- Tan, Z.X.; Lal, R.; Smeck, N.E.; Calhoun, F.G. (2004) Relationships between surface soil organic carbon pool and site variables. *Geoderma*. 121(3): 187-195.
- Thompson, J.A.; Kolka, R.K. (2005) Soil carbon storage estimation in a forested watershed using quantitative soil-landscape modeling. *Soil Science Society of America Journal*. 69(4): 1086-1093.
- Ulrich, A.H. (1989) *U.S. Timber Production, Trade, Consumption, and Price Statistics, 1950-1987*. USDA Miscellaneous Publication No. 1471, U.S. Department of Agriculture Forest Service. Washington, DC, 77.
- Ulrich, A.H. (1985) *U.S. Timber Production, Trade, Consumption, and Price Statistics 1950-1985*. Misc. Pub. 1453, U.S. Department of Agriculture Forest Service. Washington, DC.
- United Nations Framework Convention on Climate Change. (2013) Report on the individual review of the inventory submission of the United States of America submitted in 2012. FCCC/ARR/2012/USA. 42 p.
- USDC Bureau of Census (1976) *Historical Statistics of the United States, Colonial Times to 1970, Vol. 1*. Washington, DC.
- USDA Forest Service (2015a) Forest Inventory and Analysis National Program: Program Features. U.S. Department of Agriculture Forest Service. Washington, D.C. Available online at <<http://fia.fs.fed.us/program-features/>>. Accessed 17 September 2015.
- USDA Forest Service. (2015b) Forest Inventory and Analysis National Program: FIA Data Mart. U.S. Department of Agriculture Forest Service. Washington, D.C. Available online at <<http://apps.fs.fed.us/fiadb-downloads/datamart.html>>. Accessed 17 September 2015.
- USDA Forest Service. (2015c) Forest Inventory and Analysis National Program, FIA library: Field Guides, Methods and Procedures. U.S. Department of Agriculture Forest Service. Washington, D.C. Available online at <<http://www.fia.fs.fed.us/library/field-guides-methods-proc/>>. Accessed 17 September 2015.
- USDA Forest Service (2015d) Forest Inventory and Analysis National Program, FIA library: Database Documentation. U.S. Department of Agriculture, Forest Service, Washington Office. Available online at <<http://fia.fs.fed.us/library/database-documentation/>>. Accessed 17 September 2015.
- USDA-NRCS (1997) "National Soil Survey Laboratory Characterization Data," Digital Data, Natural Resources Conservation Service, U.S. Department of Agriculture. Lincoln, NE.
- USDA-NRCS (2013) Summary Report: 2010 National Resources Inventory, Natural Resources Conservation Service, Washington, D.C., and Center for Survey Statistics and Methodology, Iowa State University, Ames, Iowa. <[http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/stelprdb1167354.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb1167354.pdf)>.
- U.S. EPA. (2015) Annex 3.13 Methodology for estimating net carbon stock changes in forest lands remaining forest lands. in Inventory of US Greenhouse Gas Emissions and Sinks: 1990-2013. U.S. Environmental Protection Agency. EPA 430-R-15-004.
- Veraverbeke, S., B.M. Rogers, and J.T. Randerson. (2015a) Daily burned area and carbon emissions from boreal fires in Alaska. *Biogeosciences*, 12:3579–3601.
- Veraverbeke, S., B.M. Rogers, and J.T. Randerson. (2015b) CARVE: Alaskan Fire Emissions Database (AKFED), 2001-2013. ORNL DAAC, Oak Ridge, Tennessee, USA. <http://dx.doi.org/10.3334/ORNLLDAAC/1282>. Accessed 19 October 2015.
- Wear, D.N., Coulston, J.W. (2015) From sink to source: Regional variation in U.S. forest carbon futures. *Scientific Reports*. 5: 16518.
- Wellek, S. (2003) Testing statistical hypotheses of equivalence. London, England: Chapman & Hall.
- Woldeslassie, M.; Van Miegroet, H.; Gruselle, M.C.; Hambly, N. (2012) Storage and stability of soil organic carbon in aspen and conifer forest soils of northern Utah. *Soil Science Society of America Journal*. 76(6): 2230-2240.
- Woodall, C.W., L.S. Heath, G.M. Domke, and M.C. Nichols. (2011) Methods and equations for estimating aboveground volume, biomass, and carbon for trees in the U.S. forest inventory, 2010. Gen. Tech. Rep. NRS-88. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 30 p.
- Woodall, C.W., B.L. Conkling, M.C. Amacher, J.W. Coulston, S. Jovan, C.H. Perry, B. Schulz, G.C. Smith, S. Will Wolf. (2010). The Forest Inventory and Analysis Database Version 4.0: Database Description and Users Manual for Phase

3. Gen. Tech. Rep. NRS-61. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 180 p.
- Woodall, C.W., Coulston, J.W., Domke, G.M., Walters, B.F., Wear, D.N., Smith, J.E., Anderson, H.-E., Clough, B.J., Cohen, W.B., Griffith, D.M., Hagan, S.C., Hanou, I.S.; Nichols, M.C., Perry, C.H., Russell, M.B., Westfall, J.A., Wilson, B.T. (2015a) The U.S. Forest Carbon Accounting Framework: Stocks and Stock change 1990-2016. Gen. Tech. Rep. NRS-154. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 49 pp.
- Woodall, C.W., Walters, B.F., Coulston, J.W., D'Amato, A.W., Domke, G.M., Russell, M.B., Sowers, P.A. (2015b) Monitoring network confirms land use change is a substantial component of the forest carbon sink in the eastern United States. *Scientific Reports*. 5: 17028.
- Woodall, C.W., Domke, G.M., MacFarlane, D.W., Oswald, C.M. (2012) Comparing Field- and Model-Based Standing Dead Tree Carbon Stock Estimates Across Forests of the United States. *Forestry* 85(1): 125-133.
- Woodall, C.W., Walters, B.F., Oswald, S.N., Domke, G.M., Toney, C., Gray, A.N. (2013) Biomass and carbon attributes of downed woody materials in forests of the United States. *Forest Ecology and Management* 305: 48-59.
- Woodall, C.W., Domke, G.M., MacFarlane, D.W., Oswald, C.M. (2012) Comparing field- and model-based standing dead tree carbon stock estimates across forests of the United States. *Forestry*. 85: 125-133.
- Woudenberg, S.W. and T.O. Farrenkopf (1995) The Westwide forest inventory data base: user's manual. General Technical Report INT-GTR-317. U.S. Department of Agriculture Forest Service, Intermountain Research Station.

### 3.14. Methodology for Estimating CH<sub>4</sub> Emissions from Landfills

Landfill gas is a mixture of substances generated when bacteria decompose the organic materials contained in solid waste. By volume, landfill gas is about half CH<sub>4</sub> and half CO<sub>2</sub>.<sup>110</sup> The amount and rate of CH<sub>4</sub> generation depends upon the quantity and composition of the landfilled material, as well as the surrounding landfill environment.

Not all CH<sub>4</sub> generated within a landfill is emitted to the atmosphere. The CH<sub>4</sub> can be extracted and either flared or utilized for energy, thus oxidizing the CH<sub>4</sub> to CO<sub>2</sub> during combustion. Of the remaining CH<sub>4</sub>, a portion oxidizes to CO<sub>2</sub> as it travels through the top layer of the landfill cover. In general, landfill-related CO<sub>2</sub> emissions are of biogenic origin and primarily result from the decomposition, either aerobic or anaerobic, of organic matter such as food or yard wastes.<sup>111</sup> To estimate the amount of CH<sub>4</sub> produced in a landfill in a given year, information is needed on the type and quantity of waste in the landfill, as well as the landfill characteristics (e.g., size, aridity, waste density). This information is not available for the majority of landfills in the United States. Consequently, to estimate CH<sub>4</sub> generation, a methodology was developed (i.e., the first order decay waste model) based on the quantity of waste placed in landfills nationwide each year and model parameters from the analysis of measured CH<sub>4</sub> generation rates for U.S. landfills with gas recovery systems.

From various studies, surveys, and regulatory reporting programs (i.e., the Greenhouse Gas Reporting Program [GHGRP]) of the generation and disposal of solid waste, estimates of the amount of waste placed in MSW and industrial waste landfills were developed for previous Inventory years. The current Inventory relied heavily on the GHGRP data for municipal solid waste (MSW) landfills for current and historical (since the year the landfill started accepting waste) reported waste disposal quantities. A database of measured CH<sub>4</sub> generation rates at MSW landfills with gas recovery systems was compiled and analyzed. The results of this analysis and other studies were used to develop an estimate of the CH<sub>4</sub> generation potential for use in the first order decay model. In addition, the analysis and other studies provided estimates of the CH<sub>4</sub> generation rate constant as a function of precipitation. The first order decay (FOD) model was applied to annual waste disposal estimates for each year and for three ranges of precipitation to estimate CH<sub>4</sub> generation rates nationwide for the years of interest. Based on the organic content of industrial wastes and the estimates of the fraction of these wastes sent to industrial waste landfills, CH<sub>4</sub> emissions from industrial waste landfills were also estimated using the first order decay model. Total CH<sub>4</sub> emissions were estimated by adding the CH<sub>4</sub> generation from MSW and industrial landfills and subtracting the amounts of CH<sub>4</sub> recovered for energy or flaring at MSW landfills<sup>112</sup> and the amount oxidized in the soil at MSW and industrial landfills. The steps taken to estimate CH<sub>4</sub> emissions from U.S. landfills for the years 1990 through the current inventory year are discussed in greater detail below.

Figure A-24 presents the CH<sub>4</sub> emissions process—from waste generation to emissions—in graphical format.

#### Step 1: Estimate Annual Quantities of Solid Waste Placed in Landfills

For 1989 to 2014, estimates of the annual quantity of waste placed in MSW landfills were developed from a survey of State agencies as reported in the State of Garbage (SOG) in America surveys (Shin 2014; BioCycle 2010), adjusted to include U.S. territories.<sup>113</sup> The SOG survey is the only continually updated nationwide survey of waste disposed in landfills in the United States. Table A-272 shows estimates of waste quantities contributing to CH<sub>4</sub> emissions. The table shows SOG estimates of total waste generated and total waste landfilled (adjusted for U.S. territories) for various years over the 1990 to 2013 timeframe.

State-specific landfill waste generation data and a national average disposal factor for 1989 through 2008 were obtained from the SOG survey for every two years (i.e., 2002, 2004, 2006, and 2008 as published in BioCycle 2006; 2008, and 2010). The most recent SOG survey was published in 2014 (Shin 2014) for the 2011 year. A linear interpolation was used for the amount of waste generated in 2001, 2003, 2005, 2007, 2009, 2010, 2012, 2013, and 2014 because no SOG surveys were published for those years. Upon publication of the next SOG survey, the waste landfilled for 2012 to 2014 will be updated. Estimates of the quantity of waste landfilled from 1989 to the current inventory year are determined by applying a waste disposal factor to the total amount of waste generated (i.e., the SOG data). A waste disposal factor is determined for each year a SOG survey is published and is the ratio of the total amount of waste landfilled to the total amount of waste

<sup>110</sup> Typically, landfill gas also contains small amounts of nitrogen, oxygen, and hydrogen, less than 1 percent nonmethane volatile organic compounds (NMVOCs), and trace amounts of inorganic compounds.

<sup>111</sup> See Box 7-1 “Biogenic Emissions and Sinks of Carbon” in the Waste chapter for additional background on how biogenic emissions of landfill CO<sub>2</sub> are addressed in the U.S. Inventory.


<sup>112</sup> Landfill gas recovery is only estimated for MSW landfills due to a lack of national data on industrial waste landfills. Approximately 1 percent of the industrial waste landfills reporting under the GHGRP have active landfill gas collection systems.

<sup>113</sup> Since the SOG survey does not include U.S. territories, waste landfilled in U.S. territories was estimated using population data for the U.S. territories (U.S. Census Bureau 2013) and the per capita rate for waste landfilled from BioCycle (2010).

generated. The waste disposal factor is interpolated for the years in-between the SOG surveys and extrapolated for years after the last SOG survey. Methodological changes have occurred over the time that the SOG survey has been published, and this has affected the fluctuating trends observed in the data (RTI 2013).

Historical waste data, preferably since 1940, are required for the FOD model to estimate CH<sub>4</sub> generation for the Inventory time series. Estimates of waste placed in landfills in the 1940s and 1950s were developed based on U.S. population for each year and the per capital disposal rates from the 1960s. Estimates of the annual quantity of waste placed in landfills from 1960 through 1983 were developed from EPA's 1993 Report to Congress (EPA 1993) and a 1986 survey of MSW landfills (EPA 1988).

**Figure A-24: Methane Emissions Resulting from Landfilling Municipal and Industrial Waste**



<sup>a</sup> MSW waste generation is not calculated because annual quantities of waste disposal are available through EPA 2015c; annual production data used for industrial waste (Lockwood Post's Directory and the USDA).

<sup>b</sup> 1940 through 1988 based on EPA 1988 and EPA 1993; 1989 through 2008 based on *BioCycle* 2010; 2009 through 2014 based on Shin 2014.

<sup>c</sup> 2006 *IPCC Guidelines* – First Order Decay Model.

<sup>d</sup> EIA 2007, flare vendor database, EPA (GHGRP) 2015b.

<sup>e</sup> EIA 2007, EPA (LMOP) 2015a, and EPA (GHGRP) 2015b.

<sup>f</sup> 2006 *IPCC Guidelines*; Mancinelli and McKay 1985; Czepiel et al 1996.

**Table A-272: Solid Waste in MSW Landfills Contributing to CH<sub>4</sub> Emissions (MMT unless otherwise noted)**

|                                   | 1990  | 2005   | 2006   | 2007   | 2008   | 2009   | 2010   | 2011   | 2012   | 2013   | 2014   |
|-----------------------------------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Total MSW Generated <sup>a</sup>  | 270   | 458    | 454    | 429    | 403    | 405    | 406    | 408    | 411    | 414    | 418    |
| Percent of MSW                    |       |        |        |        |        |        |        |        |        |        |        |
| Landfilled <sup>a</sup>           | 77%   | 64%    | 65%    | 67%    | 69%    | 66%    | 66%    | 63%    | 63%    | 63%    | 63%    |
| Total MSW Landfilled              | 205   | 290    | 289    | 283    | 275    | 265    | 266    | 256    | 258    | 259    | 262    |
| MSW last 30 years <sup>b</sup>    | 4,876 | 6,281  | 6,415  | 6,539  | 6,653  | 6,753  | 6,851  | 6,935  | 7,018  | 7,099  | 7,180  |
| MSW since 1940 <sup>c</sup>       | 6,808 | 10,214 | 10,503 | 10,786 | 11,061 | 11,326 | 11,591 | 11,847 | 12,104 | 12,364 | 12,625 |
| Total Industrial Waste Landfilled | 10    | 11     | 11     | 11     | 11     | 10     | 11     | 11     | 11     | 11     | 11     |

<sup>a</sup> This estimate represents the waste that has been in place for 30 years or less, which contributes about 90 percent of the CH<sub>4</sub> generation. Values are based on EPA (1993) for years 1940 to years 1988 (not presented in table), *BioCycle* 2010 and Shin 2014 for years 1989 to 2014 (years prior to 1990 and 1991 to 2004 are not presented in table).

<sup>b</sup> Source: Data for years 1990 through 2014 are from the GHGRP for MSW landfills EPA (GHGRP) 2015, EPA (LMOP) 2015, and state agencies (EPA 2015c).

<sup>c</sup> This estimate represents the cumulative amount of waste that has been placed in landfills since 1940 to the year indicated and is the sum of the annual disposal rates used in the first order decay model. Values are based on EPA 1993, *BioCycle* 2010 and Shin 2014.

## Step 2: Estimate CH<sub>4</sub> Generation at Municipal Solid Waste Landfills

The CH<sub>4</sub> generation was estimated from the integrated form of the FOD model using the procedures and spreadsheets from IPCC (2006) for estimating CH<sub>4</sub> emissions from solid waste disposal. The form of the FOD model that was applied incorporates a time delay of 6 months after waste disposal before the generation of CH<sub>4</sub> begins.

The input parameters needed for the FOD model equations are the mass of waste disposed each year, which was discussed in the previous section, degradable organic carbon (DOC), and the decay rate constant (k). The DOC is determined from the CH<sub>4</sub> generation potential (L<sub>0</sub> in m<sup>3</sup> CH<sub>4</sub>/Mg waste), which is discussed in more detail in subsequent paragraphs, and the following equation:

$$\text{DOC} = [L_0 \times 6.74 \times 10^{-4}] \div [F \times 16/12 \times \text{DOC}_f \times \text{MCF}]$$

where,

|                         |   |   |
|-------------------------|---|---|
| DOC                     | = | degradable organic carbon (fraction, kt C/kt waste),  |
| L <sub>0</sub>          | = | CH <sub>4</sub> generation potential (m <sup>3</sup> CH <sub>4</sub> /Mg waste),                                    |
| 6.74 × 10 <sup>-4</sup> | = | CH <sub>4</sub> density (Mg/m <sup>3</sup> ),   |
| F                       | = | fraction of CH <sub>4</sub> by volume in generated landfill gas (equal to 0.5)                                      |
| 16/12                   | = | molecular weight ratio CH <sub>4</sub> /C,  |
| DOC <sub>f</sub>        | = | fraction of DOC that can decompose in the anaerobic conditions in the landfill (fraction equal to 0.5 for MSW), and |
| MCF                     | = | methane correction factor for year of disposal (fraction equal to 1 for anaerobic managed sites).                   |

The DOC value used in the CH<sub>4</sub> generation estimates from MSW landfills is 0.203 based on the CH<sub>4</sub> generation potential of 100 m<sup>3</sup> CH<sub>4</sub>/Mg waste as described below. Data from a set of 52 representative landfills across the U.S. in different precipitation ranges were chosen to evaluate L<sub>0</sub>, and ultimately the country-specific DOC value. The 2004 Chartwell Municipal Solid Waste Facility Directory confirmed that each of the 52 landfills chosen accepted or accepts both MSW and construction and demolition (C&D) waste (Chartwell 2004; RTI 2009).

The methane generation potential (L<sub>0</sub>) varies with the amount of organic content of the waste material. A higher L<sub>0</sub> occurs with a higher content of organic waste. Waste composition data are not collected for all landfills nationwide; thus a default value must be used. Values for L<sub>0</sub> were evaluated from landfill gas recovery data for this set of 52 landfills, which resulted in a best fit value for L<sub>0</sub> of 99 m<sup>3</sup>/Mg of waste (RTI 2004). This value compares favorably with a range of 50 to 162 (midrange of 106) m<sup>3</sup>/Mg presented by Peer, Thorneloe, and Epperson (1993); a range of 87 to 91 m<sup>3</sup>/Mg from a detailed analysis of 18 landfills sponsored by the Solid Waste Association of North America (SWANA 1998); and a value of 100 m<sup>3</sup>/Mg recommended in EPA's compilation of emission factors (EPA 1998; EPA 2008) based on data from 21 landfills. Based on the results from these studies, a value of 100 m<sup>3</sup>/Mg appears to be a reasonable best estimate to use in the FOD model for the national inventory.

The FOD model was applied to the gas recovery data for the 52 landfills to calculate the decay rate constant (k) directly for L<sub>0</sub> = 100 m<sup>3</sup>/Mg. The rate constant was found to increase with annual average precipitation; consequently, average values of k were developed for three ranges of precipitation, shown in Table A-273 and recommended in EPA's compilation of emission factors (EPA 2008).

**Table A-273: Average Values for Rate Constant (k) by Precipitation Range (yr<sup>-1</sup>)**

| Precipitation range (inches/year) | k (yr <sup>-1</sup> ) |
|-----------------------------------|-----------------------|
| <20                               | 0.020                 |
| 20-40                             | 0.038                 |
| >40                               | 0.057                 |

These values for k show reasonable agreement with the results of other studies. For example, EPA's compilation of emission factors (EPA 1998; EPA 2008) recommends a value of 0.02 yr<sup>-1</sup> for arid areas (less than 20 inches/year of precipitation) and 0.04 yr<sup>-1</sup> for non-arid areas. The SWANA (1998) study of 18 landfills reported a range in values of k from 0.03 to 0.06 yr<sup>-1</sup> based on CH<sub>4</sub> recovery data collected generally in the time frame of 1986 to 1995.

Using data collected primarily for the year 2000, the distribution of waste in place versus precipitation was developed from over 400 landfills (RTI 2004). A distribution was also developed for population vs. precipitation for comparison. The two distributions were very similar and indicated that population in areas or regions with a given precipitation range was a reasonable proxy for waste landfilled in regions with the same range of precipitation. Using U.S. Census data and rainfall data, the distributions of population versus rainfall were developed for each Census decade from

1950 through 2000. The distributions showed that the U.S. population has shifted to more arid areas over the past several decades. Consequently, the population distribution was used to apportion the waste landfilled in each decade according to the precipitation ranges developed for k, as shown in Table A-274.

**Table A-274: Percent of U.S. Population within Precipitation Ranges (%)**

| Precipitation Range (inches/year) | 1950 | 1960 | 1970 | 1980 | 1990 | 2000 | 2010 |
|-----------------------------------|------|------|------|------|------|------|------|
| <20                               | 10   | 13   | 14   | 16   | 19   | 19   | 18   |
| 20-40                             | 40   | 39   | 37   | 36   | 34   | 33   | 44   |
| >40                               | 50   | 48   | 48   | 48   | 48   | 48   | 38   |

Source: Years 1950 through 2000 are from RTI (2004) using population data from the U.S. Census Bureau and precipitation data from the National Climatic Data Center's National Oceanic and Atmospheric Administration. Year 2010 is based on the methodology from RTI (2004) and the U.S. Bureau of Census and precipitation data from the National Climatic Data Center's National Oceanic and Atmospheric Administration where available.

In developing the Inventory, the proportion of waste disposed of in managed landfills versus open dumps prior to 1980 was re-evaluated. Based on the historical data presented by Mintz et al. (2003), a timeline was developed for the transition from the use of open dumps for solid waste disposed to the use of managed landfills. Based on this timeline, it was estimated that 6 percent of the waste that was land disposed in 1940 was disposed of in managed landfills and 94 percent was managed in open dumps. Between 1940 and 1980, the fraction of waste that was land disposed transitioned towards managed landfills until 100 percent of the waste was disposed of in managed landfills in 1980. For wastes disposed of in dumps, a methane correction factor (MCF) of 0.6 was used based on the recommended IPCC default value for uncharacterized land disposal (IPCC 2006); this MCF is equivalent to assuming 50 percent of the open dumps are deep and 50 percent are shallow. The recommended IPCC default value for the MCF for managed landfills of 1 was used for the managed landfills (IPCC 2006).

### Step 3: Estimate CH<sub>4</sub> Generation at Industrial Landfills

Industrial waste landfills receive waste from factories, processing plants, and other manufacturing activities. In national inventories prior to the 1990 through 2005 inventory, CH<sub>4</sub> generation at industrial landfills was estimated as seven percent of the total CH<sub>4</sub> generation from MSW landfills, based on a study conducted by EPA (1993). For the 1990 through current inventories, the methodology was updated and improved by using activity factors (industrial production levels) to estimate the amount of industrial waste landfilled each year and by applying the FOD model to estimate CH<sub>4</sub> generation. A nationwide survey of industrial waste landfills found that over 99 percent of the organic waste placed in industrial landfills originated from two sectors: food processing (meat, vegetables, fruits) and pulp and paper (EPA 1993). Data for annual nationwide production for the food processing and pulp and paper sectors were taken from industry and government sources for recent years; estimates were developed for production for the earlier years for which data were not available. For the pulp and paper sector, production data published by the Lockwood-Post's Directory were used for years 1990 to 2001 and production data published by the U.S. Department of Agriculture were used for years 2002 through 2013. An extrapolation based on U.S. real gross domestic product was used for years 1940 through 1964. For the food processing sector, production levels were obtained or developed from the U.S. Department of Agriculture for the years 1990 through 2013 (ERG 2014). An extrapolation based on U.S. population was used for the years 1940 through 1989, and for the year 2014.

In addition to production data for the pulp and paper and food processing sectors, the following inputs are needed to use the FOD model for estimating CH<sub>4</sub> generation from industrial landfills: 1) quantity of waste that is disposed in industrial waste landfills (as a function of production), 2) CH<sub>4</sub> generation potential ( $L_0$ ) or DOC, and 3) the decay rate constant (k). Research into waste generation and disposal in landfills for the pulp and paper sector indicated that the quantity of waste landfilled was about 0.050 MT/MT of product compared to 0.046 MT/MT product for the food processing sector (RTI 2006). These factors were applied to estimates of annual production to estimate annual waste disposal in industrial waste landfills. Estimates for DOC were derived from available data (EPA, 2015b; Heath et al., 2010; NCASI, 2005; Kraft and Orender, 1993; NCASI 2008; Flores et al. 1999 as documented in RTI 2015b). The DOC value for industrial pulp and paper waste is estimated at 0.15 ( $L_0$  of 49 m<sup>3</sup>/MT); the DOC value for industrial food waste is estimated as 0.26 ( $L_0$  of 128 m<sup>3</sup>/MT) (RTI 2015b; RTI 2014). Estimates for k were taken from the default values in the 2006 IPCC Guidelines; the value of k given for food waste with disposal in a wet temperate climate is 0.19 yr<sup>-1</sup>, and the value given for paper waste is 0.06 yr<sup>-1</sup>.

A literature review was conducted for the 1990 to 2010 and 1990 to 2014 inventory years with the intent of updating values for  $L_0$  (specifically DOC) and k in the pulp and paper sector. Where pulp and paper mill wastewater treatment residuals or sludge are the primary constituents of pulp and paper waste landfilled, values for k available in the literature

range from 0.01/yr to 0.1/yr, while values for  $L_o$  range from 50 m<sup>3</sup>/Mt to 200 m<sup>3</sup>/Mt.<sup>114</sup> Values for these factors are highly variable and are dependent on the soil moisture content, which is generally related to rainfall amounts. At this time, sufficient data were available through the GHGRP to warrant a change to the  $L_o$  (DOC) from 99 to 49 m<sup>3</sup>/MT, but sufficient data were not obtained to warrant a change to  $k$  for the current inventory year. EPA will consider an update to the  $k$  values for the pulp and paper sector as new data arises and will work with stakeholders to gather data and other feedback on potential changes to these values.

As with MSW landfills, a similar trend in disposal practices from open dumps to managed landfills was expected for industrial waste landfills; therefore, the same time line that was developed for MSW landfills was applied to the industrial landfills to estimate the average MCF. That is, between 1940 and 1980, the fraction of waste that was land disposed transitioned from 6 percent managed landfills in 1940 and 94 percent open dumps to 100 percent managed landfills in 1980 and on. For wastes disposed of in dumps, an MCF of 0.6 was used and for wastes disposed of in managed landfills, an MCF of 1 was used, based on the recommended IPCC default values (IPCC 2006).

The parameters discussed above were used in the integrated form of the FOD model to estimate CH<sub>4</sub> generation from industrial waste landfills.

#### **Step 4: Estimate CH<sub>4</sub> Emissions Avoided**

The estimated landfill gas recovered per year (R) at MSW landfills was based on a combination of four databases and including recovery from flares and/or landfill gas-to-energy projects:

- EPA's GHGRP dataset for MSW landfills (EPA 2015b)
- a database developed by the Energy Information Administration (EIA) for the voluntary reporting of greenhouse gases (EIA 2007)
- a database of LFGTE projects that is primarily based on information compiled by EPA LMOP (EPA 2015a), and
- the flare vendor database (contains updated sales data collected from vendors of flaring equipment).

EPA's GHGRP MSW landfills database was first introduced as a data source for the 1990 to 2013 Inventory. The GHGRP MSW landfills database contains facility-reported data that undergoes rigorous verification and is considered to contain the least uncertain data of the four databases. However, this database is unique in that it only contains a portion of the landfills in the U.S. (although, presumably the highest emitters since only those landfills that meet the methane generation threshold must report) and only contains data from 2010 and later.

The destruction efficiencies reported through EPA's GHGRP were applied to the landfills in the GHGRP MSW landfills database. The median value of the reported destruction efficiencies was 99 percent for all reporting years (2010 through 2014, EPA 2015b). A destruction efficiency of 99 percent was applied to CH<sub>4</sub> recovered to estimate CH<sub>4</sub> emissions avoided for the other three databases. This value for destruction efficiency was selected based on the range of efficiencies (86 to 99+ percent) recommended for flares in EPA's *AP-42 Compilation of Air Pollutant Emission Factors*, Draft Chapter 2.4, Table 2.4-3 (EPA 2008). A typical value of 97.7 percent was presented for the non-methane components (i.e., volatile organic compounds and non-methane organic compounds) in test results (EPA 2008). An arithmetic average of 98.3 percent and a median value of 99 percent are derived from the test results presented in EPA 2008. Thus, a value of 99 percent for the destruction efficiency of flares has been used in Inventory methodology. Other data sources supporting a 99 percent destruction efficiency include those used to establish New Source Performance Standards (NSPS) for landfills and in recommendations for closed flares used in the EPA's LMOP.

#### **Step 4a: Estimate CH<sub>4</sub> Emissions Avoided Through Landfill Gas-to-Energy (LFGTE) and Flaring Projects**

The quantity of CH<sub>4</sub> avoided due to LFGTE systems was estimated based on information from three sources: (1) a database developed by the EIA for the voluntary reporting of greenhouse gases (EIA 2007); (2) a database compiled by LMOP and referred to as the LFGTE database for the purposes of this inventory (EPA 2015a); and (3) the GHGRP MSW landfills dataset (EPA 2015b). The EIA database included location information for landfills with LFGTE projects, estimates of CH<sub>4</sub> reductions, descriptions of the projects, and information on the methodology used to determine the CH<sub>4</sub> reductions. In general, the CH<sub>4</sub> reductions for each reporting year were based on the measured amount of landfill gas collected and the percent CH<sub>4</sub> in the gas. For the LFGTE database, data on landfill gas flow and energy generation (i.e., MW capacity) were used to estimate the total direct CH<sub>4</sub> emissions avoided due to the LFGTE project. The GHGRP MSW landfills database contains the most detailed data on landfills that reported under the GHGRP for years 2010 through 2014, however the amount of CH<sub>4</sub> recovered is not specifically allocated to a flare versus a LFGTE project. The allocation into flares or LFGTE was

<sup>114</sup> Sources reviewed included Heath et al. 2010; Miner 2008; Skog 2008; Upton et al. 2008; Barlaz 2006; Sonne 2006; NCASI 2005; Barlaz 1998; and Skog and Nicholson 2000.

performed by matching landfills to the EIA and LMOP databases for LFGTE projects and to the flare database for flares. Detailed information on the landfill name, owner or operator, city, and state are available for both the EIA and LFGTE databases; consequently, it was straightforward to identify landfills that were in both databases against those in EPA's GHGRP MSW landfills database. EPA's GHGRP MSW landfills database was given priority because CH<sub>4</sub> recovery and other supporting information were reported for each year and were based on direct measurements. The EIA database was given second priority (for landfills not in the GHGRP MSW landfills database) for years 1990 through 2005 because CH<sub>4</sub> recovery based on direct measurements was reported by landfill during this period. Landfills in the LMOP database that were also in EPA's GHGRP MSW landfills database and/or EIA database were not included in the total recovery values to avoid double or triple counting across the databases.

If a landfill in EPA's GHGRP MSW landfills database was also in the EIA, LFGTE, and/or flare vendor database, the avoided emissions were only based on EPA's GHGRP MSW landfills database to avoid double or triple counting the recovery amounts. In other words, the recovery from the same landfill was not included in the total recovery from the EIA, LFGTE, or flare vendor databases.

If a landfill in the EIA database was also in the LFGTE and/or the flare vendor database, the CH<sub>4</sub> recovery was based on the EIA data because landfill owners or operators directly reported the amount of CH<sub>4</sub> recovered using gas flow concentration and measurements, and because the reporting accounted for changes over time. However, as the EIA database only includes facility-reported data through 2006, the amount of CH<sub>4</sub> recovered for years 2007 and later were assumed to be the same as in 2006 for landfills that are in the EIA database, but not in the GHGRP or LFGTE databases. This quantity likely underestimates flaring because the EIA database does not have information on all flares in operation for the years after 2006. However, nearly all (93 percent) of landfills in the EIA database also report to the GHGRP, which means that only seven percent of landfills in the EIA database are counted in the total recovery.

If both the flare data and LFGTE recovery data were available for any of the remaining landfills (i.e., not in the EIA or GHGRP databases), then the avoided emissions were based on the LFGTE data, which provides reported landfill-specific data on gas flow for direct use projects and project capacity (i.e., megawatts) for electricity projects. The LFGTE database is based on the most EPA LMOP database (published annually). The remaining portion of avoided emissions is calculated by the flare vendor database, which estimates CH<sub>4</sub> combusted by flares using the midpoint of a flare's reported capacity. New flare vendor sales data were unable to be obtained for the current Inventory year. Given that each LFGTE project is likely to also have a flare, double counting reductions from flares and LFGTE projects in the LFGTE database was avoided by subtracting emission reductions associated with LFGTE projects for which a flare had not been identified from the emission reductions associated with flares (referred to as the flare correction factor).

#### ***Step 4b: Estimate CH<sub>4</sub> Emissions Avoided Through Flaring for the Flare Database***

To avoid double counting, flares associated with landfills in EPA's GHGRP, EIA and LFGTE databases were not included in the total quantity of CH<sub>4</sub> recovery from the flare vendor database. As with the LFGTE projects, reductions from flaring landfill gas in the EIA database were based on measuring the volume of gas collected and the percent of CH<sub>4</sub> in the gas. The information provided by the flare vendors included information on the number of flares, flare design flow rates or flare dimensions, year of installation, and generally the city and state location of the landfill. When a range of design flare flow rates was provided by the flare vendor, the median landfill gas flow rate was used to estimate CH<sub>4</sub> recovered from each remaining flare (i.e., for each flare not associated with a landfill in the EIA, GHGRP, or LFGTE databases). Several vendors provided information on the size of the flare rather than the flare design gas flow rate. To estimate a median flare gas flow rate for flares associated with these vendors, the size of the flare was matched with the size and corresponding flow rates provided by other vendors. Some flare vendors reported the maximum capacity of the flare. An analysis of flare capacity versus measured CH<sub>4</sub> flow rates from the EIA database showed that the flares operated at 51 percent of capacity when averaged over the time series and at 72 percent of capacity for the highest flow rate for a given year. For those cases when the flare vendor supplied maximum capacity, the actual flow was estimated as 50 percent of capacity. Total CH<sub>4</sub> avoided through flaring from the flare vendor database was estimated by summing the estimates of CH<sub>4</sub> recovered by each flare for each year. Flare sales data were not provided to the EPA for the previous and current Inventory year.

#### ***Step 4c: Reduce CH<sub>4</sub> Emissions Avoided Through Flaring***

As mentioned in Step 4b, flares in the flare vendor database associated with landfills in the GHGRP MSW landfills, EIA, and LMOP databases were not accounted for in the flare reduction estimates in the flare vendor database. If comprehensive data on flares were available, each LFGTE project in EPA's GHGRP, EIA, and LFGTE databases would have an identified flare because it is assumed that most LFGTE projects have flares. However, given that the flare vendor database only covers approximately 50 to 75 percent of the flare population, an associated flare was not identified for all LFGTE projects. These LFGTE projects likely have flares, yet flares were unable to be identified for one of two reasons: 1)

inadequate identifier information in the flare vendor data, or 2) a lack of the flare in the flare vendor database. For those projects for which a flare was not identified due to inadequate information, CH<sub>4</sub> avoided would be overestimated, as both the CH<sub>4</sub> avoided from flaring and the LFGTE project would be counted. To avoid overestimating emissions avoided from flaring, the CH<sub>4</sub> avoided from LFGTE projects with no identified flares was determined and the flaring estimate from the flare vendor database was reduced by this quantity (referred to as a flare correction factor) on a state-by-state basis. This step likely underestimates CH<sub>4</sub> avoided due to flaring, but was applied to be conservative in the estimates of CH<sub>4</sub> emissions avoided.

Additional effort was undertaken to improve the methodology behind the flare correction factor for the 1990 to 2009 and 1990 to 2014 inventory years to reduce the total number of flares in the flare vendor database that were not matched to landfills and/or LFGTE projects in the EIA and LFGTE databases. Each flare in the flare vendor database not associated with a LFGTE project in the EIA, LFGTE, or GHGRP databases was investigated to determine if it could be matched. For some unmatched flares, the location information was missing or incorrectly transferred to the flare vendor database and was corrected during the review. In other instances, the landfill names were slightly different between what the flare vendor provided and the actual landfill name as listed in the EIA, LFGTE and GHGRP databases. The remaining flares did not have adequate information through the name, location, or owner to identify it to a landfill in any of the recovery databases or through an Internet search; it is these flares that are included in the flare correction factor for the current inventory year.

A large majority of the unmatched flares are associated with landfills in the LFGTE database that are currently flaring, but are also considering LFGTE. These landfills projects considering a LFGTE project are labeled as candidate, potential, or construction in the LFGTE database. The flare vendor database was improved in the 1990 to 2009 inventory year to match flares with operational, shutdown as well as candidate, potential, and construction LFGTE projects, thereby reducing the total number of unidentified flares in the flare vendor database, all of which are used in the flare correction factor. The results of this effort significantly decreased the number of flares used in the flare correction factor, and consequently, increased recovered flare emissions, and decreased net emissions from landfills for the 1990 through 2009 Inventory. The revised state-by-state flare correction factors were applied to the entire Inventory time series.

#### **Step 5: Estimate CH<sub>4</sub> Oxidation**

A portion of the CH<sub>4</sub> escaping from a landfill oxidizes to CO<sub>2</sub> in the top layer of the soil. The amount of oxidation depends upon the characteristics of the soil and the environment. For purposes of this analysis, it was assumed that of the CH<sub>4</sub> generated, minus the amount of gas recovered for flaring or LFGTE projects, 10 percent was oxidized in the soil (Jensen and Pipatti 2002; Mancinelli and McKay 1985; Czepiel et al 1996). The factor of 10 percent is consistent with the value recommended in the 2006 IPCC revised guidelines for managed and covered landfills, and was therefore applied to the estimates of CH<sub>4</sub> generation minus recovery for both MSW and industrial landfills

A literature review was conducted in 2011 (RTI 2011) to provide recommendations for the most appropriate oxidation rate assumptions. It was found that oxidation values are highly variable and range from zero to over 100 percent (i.e., the landfill is considered to be an atmospheric sink by virtue of the landfill gas extraction system pulling atmospheric methane down through the cover). There is considerable uncertainty and variability surrounding estimates of the rate of oxidation because oxidation is difficult to measure and varies considerably with the presence of a gas collection system, thickness and type of the cover material, size and area of the landfill, climate, and the presence of cracks and/or fissures in the cover material through which methane can escape. IPCC (2006) notes that test results from field and laboratory studies may lead to over-estimations of oxidation in landfill cover soils because they largely determine oxidation using uniform and homogeneous soil layers. In addition, a number of studies note that gas escapes more readily through the side slopes of a landfill as compared to moving through the cover thus complicating the correlation between oxidation and cover type or gas recovery.

Sites with landfill gas collection systems are generally designed and managed better to improve gas recovery. More recent research (2006 to 2012) on landfill cover methane oxidation has relied on stable isotope techniques that may provide a more reliable measure of oxidation. Results from this recent research consistently point to higher cover soil methane oxidation rates than the IPCC (2006) default of 10 percent. A continued effort will be made to review the peer-reviewed literature to better understand how climate, cover type, and gas recovery influence the rate of oxidation at active and closed landfills. At this time, the IPCC recommended oxidation factor of 10 percent will continue to be used for all landfills.

#### **Step 6: Estimate Total CH<sub>4</sub> Emissions**

Total CH<sub>4</sub> emissions were calculated by adding emissions from MSW and industrial landfills, and subtracting CH<sub>4</sub> recovered and oxidized, as shown in Table A-275.

**Table A-275: CH<sub>4</sub> Emissions from Landfills (kt)**

|  | 1990         | 1995           | 2001           | 2002           | 2003           | 2004           | 2005           | 2006           | 2007           | 2008           | 2009           | 2010           | 2011           | 2012           | 2013           | 2014           |
|--|--------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| MSW CH <sub>4</sub> Generation           | 8,214        | 9,140          | 10,084         | 10,383         | 10,769         | 11,135         | 11,482         | 11,809         | 12,118         | 12,395         | 12,638         | 12,839         | 13,008         | 13,144         | 13,280         | 13,418         |
| Industrial CH <sub>4</sub> Generation    | 484          | 537            | 612            | 618            | 625            | 629            | 636            | 639            | 643            | 648            | 653            | 656            | 657            | 659            | 661            | 665            |
| <b>Potential Emissions</b>               | <b>8,698</b> | <b>9,676</b>   | <b>10,696</b>  | <b>11,001</b>  | <b>11,394</b>  | <b>11,765</b>  | <b>12,117</b>  | <b>12,448</b>  | <b>12,760</b>  | <b>13,043</b>  | <b>13,291</b>  | <b>13,495</b>  | <b>13,665</b>  | <b>13,803</b>  | <b>13,942</b>  | <b>14,083</b>  |
| <b>Emissions Avoided</b>                 | <b>(718)</b> | <b>(1,935)</b> | <b>(4,616)</b> | <b>(4,894)</b> | <b>(4,995)</b> | <b>(5,304)</b> | <b>(5,272)</b> | <b>(5,707)</b> | <b>(5,954)</b> | <b>(6,176)</b> | <b>(6,424)</b> | <b>(7,178)</b> | <b>(7,249)</b> | <b>(7,480)</b> | <b>(7,529)</b> | <b>(7,507)</b> |
| <b>Oxidation at MSW Landfills</b>        | (750)        | (720)          | (547)          | (549)          | (577)          | (583)          | (621)          | (610)          | (616)          | (622)          | (621)          | (566)          | (576)          | (566)          | (575)          | (591)          |
| <b>Oxidation at Industrial Landfills</b> | (48)         | (54)           | (61)           | (62)           | (63)           | (63)           | (64)           | (64)           | (64)           | (65)           | (65)           | (66)           | (66)           | (66)           | (66)           | (67)           |
| <b>Net Emissions</b>                     | <b>7,182</b> | <b>6,967</b>   | <b>5,472</b>   | <b>5,496</b>   | <b>5,759</b>   | <b>5,815</b>   | <b>6,161</b>   | <b>6,066</b>   | <b>6,126</b>   | <b>6,180</b>   | <b>6,180</b>   | <b>5,685</b>   | <b>5,774</b>   | <b>5,691</b>   | <b>5,772</b>   | <b>5,919</b>   |

Notes: MSW generation in Table A-248 represents emissions before oxidation. In other tables throughout the text, MSW generation estimates account for oxidation. Totals may not sum exactly to the last significant figure due to rounding. Parentheses denote negative values.

## References

- Barlaz, M.A. (2006) "Forest Products Decomposition in Municipal Solid Waste Landfills." *Waste Management*, 26(4): 321-333.
- Barlaz, M.A. (1998) "Carbon Storage During Biodegradation of Municipal Solid Waste Components in Laboratory-scale Landfills." *Global Biogeochemical Cycles*, 12(2): 373-380, June 1998.
- BioCycle (2010) "The State of Garbage in America" By L. Arsova, R. Van Haaren, N. Goldstein, S. Kaufman, and N. Themelis. *BioCycle*. December 2010. Available online at <[http://www.jgpress.com/archives/\\_free/002191.html](http://www.jgpress.com/archives/_free/002191.html)>.
- Chartwell (2004) Municipal Solid Waste Directory. The Envirobiz Group.
- Czepiel, P., B. Mosher, P. Crill, and R. Harriss (1996) "Quantifying the Effect of Oxidation on Landfill Methane Emissions." *Journal of Geophysical Research*, 101(D11):16721-16730.
- EIA (2007) Voluntary Greenhouse Gas Reports for EIA Form 1605B (Reporting Year 2006). Available online at <<ftp://ftp.eia.doe.gov/pub/oiaf/1605/cdrom/>>.
- EPA (2015a) *Landfill Gas-to-Energy Project Database*. Landfill Methane and Outreach Program. August 2015.
- EPA (2015b) Greenhouse Gas Reporting Program (GHGRP). 2015 Envirofacts. Subpart HH: Municipal Solid Waste Landfills. Available online at <<http://www.epa.gov/enviro/facts/ghg/search.html>>.
- EPA (2008) *Compilation of Air Pollution Emission Factors, Publication AP-42*, Draft Section 2.4 Municipal Solid Waste Landfills. October 2008.
- EPA (1998) *Compilation of Air Pollution Emission Factors, Publication AP-42*, Section 2.4 Municipal Solid Waste Landfills. November 1998.
- EPA (1993) *Anthropogenic Methane Emissions in the United States, Estimates for 1990: Report to Congress*, U.S. Environmental Protection Agency, Office of Air and Radiation. Washington, D.C. EPA/430-R-93-003. April 1993.
- EPA (1988) *National Survey of Solid Waste (Municipal) Landfill Facilities*, U.S. Environmental Protection Agency. Washington, D.C. EPA/530-SW-88-011. September 1988.
- ERG (2014) Draft Production Data Supplied by ERG for 1990-2013 for Pulp and Paper, Fruits and Vegetables, and Meat. August.
- Flores, R.A., C.W. Shanklin, M. Loza-Garay, S.H. Wie (1999) "Quantification and Characterization of Food Processing Wastes/Residues." *Compost Science & Utilization*, 7(1): 63-71.
- Heath, L.S. et al. 2010. Greenhouse Gas and Carbon Profile of the U.S. Forest Products Industry Value Chain. *Environmental Science and Technology* 44(2010) 3999-4005.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Hayama, Kanagawa, Japan.
- Jensen, J.E.F., and R. Pipatti (2002) "CH<sub>4</sub> Emissions from Solid Waste Disposal." Background paper for the Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories.
- Kraft, D.L. and H.C. Orender (1993) "Considerations for Using Sludge as a Fuel." *Tappi Journal*, 76(3): 175-183.
- Mancinelli, R. and C. McKay (1985) "Methane-Oxidizing Bacteria in Sanitary Landfills." *Proc. First Symposium on Biotechnical Advances in Processing Municipal Wastes for Fuels and Chemicals*, Minneapolis, MN, 437-450. August.
- Miner, R. (2008). "Calculations documenting the greenhouse gas emissions from the pulp and paper industry." Memorandum from Reid Minor, National Council for Air and Stream Improvement, Inc. (NCASI) to Becky Nicholson, RTI International, May 21, 2008.
- Mintz C., R. Freed, and M. Walsh (2003) "Timeline of Anaerobic Land Disposal of Solid Waste." Memorandum to T. Wirth (EPA) and K. Skog (USDA), December 31, 2003.
- National Council for Air and Stream Improvement, Inc. (NCASI) (2008) "Calculations Documenting the Greenhouse Gas Emissions from the Pulp and Paper Industry." Memorandum to R. Nicholson (RTI).
- National Council for Air and Stream Improvement, Inc. (NCASI) (2005) "Calculation Tools for Estimating Greenhouse Gas Emissions from Pulp and Paper Mills, Version 1.1." July 8, 2005.
- Peer, R., S. Thorneloe, and D. Epperson (1993) "A Comparison of Methods for Estimating Global Methane Emissions from Landfills." *Chemosphere*, 26(1-4):387-400.

- RTI (2014) Analysis of DOC Values for Industrial Solid Waste for the Pulp and Paper Industry and the Food Industry. Memorandum prepared by J. Coburn for R. Schmeltz (EPA), October 28, 2014.
- RTI (2011) Updated Research on Methane Oxidation in Landfills. Memorandum prepared by K. Weitz (RTI) for R. Schmeltz (EPA), January 14, 2011.
- RTI (2009) GHG Inventory Improvement – Construction & Demolition Waste DOC and L<sub>0</sub> Value. Memorandum prepared by J. Coburn and K. Bronstein (RTI) for R. Schmeltz, April 15, 2010.
- RTI (2006) Methane Emissions for Industrial Landfills. Memorandum prepared by K. Weitz and M. Bahner for M. Weitz (EPA), September 5, 2006.
- RTI (2004) Documentation for Changes to the Methodology for the Inventory of Methane Emissions from Landfills. Memorandum prepared by M. Branscome and J. Coburn (RTI) to E. Scheehle (EPA), August 26, 2004.
- Shin, D. (2014). Generation and Disposition of Municipal Solid Waste (MSW) in the United States – A National Survey. Master of Science thesis submitted to the Department of Earth and Environmental Engineering Fu Foundation School of Engineering and Applied Science, Columbia University. January 3, 2014. Available online at <[http://www.seas.columbia.edu/earth/wtert/sofos/Dolly\\_Shin\\_Thesis.pdf](http://www.seas.columbia.edu/earth/wtert/sofos/Dolly_Shin_Thesis.pdf)>.
- Skog, K.E. (2008) “Sequestration of Carbon in harvested wood products for the United States.” *Forest Products Journal*, 58(6): 56-72.
- Skog, K. and G.A. Nicholson (2000) “Carbon Sequestration in Wood and Paper Products.” USDA Forest Service Gen. Tech. Rep. RMRS-GTR-59.
- Solid Waste Association of North America (SWANA) (1998) *Comparison of Models for Predicting Landfill Methane Recovery*. Publication No. GR-LG 0075. March 1998.
- Sonne, E. (2006) “Greenhouse Gas Emissions from Forestry Operations: A Life Cycle Assessment.” *J. Environ. Qual.* 35:1439-1450.
- Upton, B., R. Miner, M. Spinney, L.S. Heath (2008) “The Greenhouse Gas and Energy Impacts of Using Wood Instead of Alternatives in Residential Construction in the United States.” *Biomass and Bioenergy*, 32: 1-10.

# ANNEX 4 IPCC Reference Approach for Estimating CO<sub>2</sub> Emissions from Fossil Fuel Combustion

It is possible to estimate carbon dioxide (CO<sub>2</sub>) emissions from fossil fuel consumption using alternative methodologies and different data sources than those described in the Estimating Emissions from Fossil Fuel Combustion Annex. For example, the United Nations Framework Convention on Climate Change (UNFCCC) reporting guidelines request that countries, in addition to their “bottom-up” sectoral methodology, complete a “top-down” Reference Approach for estimating CO<sub>2</sub> emissions from fossil fuel combustion. Volume 2: Energy, Chapter 6: Reference Approach of the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006) states, “comparability between the sectoral and reference approaches continues to allow a country to produce a second independent estimate of CO<sub>2</sub> emissions from fuel combustion with limited additional effort and data requirements.” This reference method estimates fossil fuel consumption by adjusting national aggregate fuel production data for imports, exports, and stock changes rather than relying on end-user consumption surveys. The basic principle is that once carbon (C)-based fuels are brought into a national economy, they are either saved in some way (e.g., stored in products, kept in fuel stocks, or left unoxidized in ash) or combusted, and therefore the C in them is oxidized and released into the atmosphere. Accounting for actual consumption of fuels at the sectoral or sub-national level is not required. The following discussion provides the detailed calculations for estimating CO<sub>2</sub> emissions from fossil fuel combustion from the United States using the IPCC-recommended Reference Approach.

## Step 1: Collect and Assemble Data in Proper Format

To ensure the comparability of national inventories, the Intergovernmental Panel on Climate Change (IPCC) has recommended that countries report energy data using the International Energy Agency (IEA) reporting convention. National energy statistics were collected in physical units from several Energy Information Administration (EIA) documents in order to obtain the necessary data on production, imports, exports, and stock changes.

It was necessary to make a number of modifications to these data to generate more accurate apparent consumption estimates of these fuels. The first modification adjusts for consumption of fossil fuel feedstocks accounted for in the Industrial Processes and Product Use chapter, which include the following: unspecified coal for coal coke used in iron and steel production; natural gas, distillate fuel, and coal used in iron and steel production; natural gas used for ammonia production; petroleum coke used in the production of aluminum, ferroalloys, titanium dioxide, ammonia, and silicon carbide; and other oil and residual fuel oil used in the manufacture of C black. The second modification adjusts for the fact that EIA energy statistics include synthetic natural gas in coal and natural gas data. The third modification adjusts for the inclusion of ethanol in motor gasoline statistics. Ethanol is a biofuel, and net carbon fluxes from changes in biogenic carbon reservoirs in croplands are accounted for in the estimates for Land Use, Land-Use Change, and Forestry (see Chapter 6). The fourth modification adjusts for consumption of bunker fuels, which refer to quantities of fuels used for international transportation estimated separately from U.S. totals. The fifth modification consists of the addition of U.S. Territories data that are typically excluded from the national aggregate energy statistics. The territories include Puerto Rico, U.S. Virgin Islands, Guam, American Samoa, Wake Island, and U.S. Pacific Islands. These data, as well as the production, import, export, and stock change statistics, are presented in Table A-276.

The C content of fuel varies with the fuel's heat content. Therefore, for an accurate estimation of CO<sub>2</sub> emissions, fuel statistics were provided on an energy content basis (e.g., Btu or joules). Because detailed fuel production statistics are typically provided in physical units (as in Table A-276 for 2014), they were converted to units of energy before CO<sub>2</sub> emissions were calculated. Fuel statistics were converted to their energy equivalents by using conversion factors provided by EIA. These factors and their data sources are displayed in Table A-277. The resulting fuel type-specific energy data for 2014 are provided in Table A-278.

## Step 2: Estimate Apparent Fuel Consumption

The next step of the IPCC Reference Approach is to estimate “apparent consumption” of fuels within the country. This requires a balance of primary fuels produced, plus imports, minus exports, and adjusting for stock changes. In this way, C enters an economy through energy production and imports (and decreases in fuel stocks) and is transferred out of the country through exports (and increases in fuel stocks). Thus, apparent consumption of primary fuels (including crude oil, natural gas liquids, anthracite, bituminous, subbituminous and lignite coal, and natural gas) can be calculated as follows:

$$\text{Apparent Consumption} = \text{Production} + \text{Imports} - \text{Exports} - \text{Stock Change}$$

Flows of secondary fuels (e.g., gasoline, residual fuel, coke) should be added to primary apparent consumption. The production of secondary fuels, however, should be ignored in the calculations of apparent consumption since the C contained in these fuels is already accounted for in the supply of primary fuels from which they were derived (e.g., the estimate for apparent consumption of crude oil already contains the C from which gasoline would be refined). Flows of secondary fuels should therefore be calculated as follows:

$$\text{Secondary Consumption} = \text{Imports} - \text{Exports} - \text{Stock Change}$$

Note that this calculation can result in negative numbers for apparent consumption of secondary fuels. This result is perfectly acceptable since it merely indicates a net export or stock increase in the country of that fuel when domestic production is not considered.

Next, the apparent consumption and secondary consumption need to be adjusted for feedstock uses of fuels accounted for in the Industrial Processes and Product Use chapter, international bunker fuels, and U.S. territory fuel consumption. Bunker fuels and feedstocks accounted for in the Industrial Processes and Product Use chapter are subtracted from these estimates, while fuel consumption in U.S. Territories is added.

The IPCC Reference Approach calls for estimating apparent fuel consumption before converting to a common energy unit. However, certain primary fuels in the United States (e.g., natural gas and steam coal) have separate conversion factors for production, imports, exports, and stock changes. In these cases, it is not appropriate to multiply apparent consumption by a single conversion factor since each of its components has different heat contents. Therefore, United States fuel statistics were converted to their heat equivalents before estimating apparent consumption. Results are provided in Table A-277.

### Step 3: Estimate Carbon Emissions

Once apparent consumption is estimated, the remaining calculations are similar to those for the “bottom-up” Sectoral Approach (see Estimating Emissions from Fossil Fuel Combustion Annex). Potential CO<sub>2</sub> emissions were estimated using fuel-specific C coefficients (see Table A-278).<sup>115</sup> The C in products from non-energy uses of fossil fuels (e.g., plastics or asphalt) was then estimated and subtracted (see Table A-280). This step differs from the Sectoral Approach in that emissions from both fuel combustion and non-energy uses are accounted for in this approach. Finally, to obtain actual CO<sub>2</sub> emissions, net emissions were adjusted for any C that remained unoxidized as a result of incomplete combustion (e.g., C contained in ash or soot). The fraction oxidized was assumed to be 100 percent for petroleum, coal, and natural gas based on guidance in IPCC (2006) (see Estimating Emissions from Fossil Fuel Combustion Annex).

### Step 4: Convert to CO<sub>2</sub> Emissions

Because the 2006 IPCC Guidelines recommend that countries report greenhouse gas emissions on a full molecular weight basis, the final step in estimating CO<sub>2</sub> emissions from fossil fuel consumption was converting from units of C to units of CO<sub>2</sub>. Actual C emissions were multiplied by the molecular-to-atomic weight ratio of CO<sub>2</sub> to C (44/12) to obtain total CO<sub>2</sub> emitted from fossil fuel combustion in million metric tons (MMT). The results are contained in Table A-279.

## Comparison Between Sectoral and Reference Approaches

These two alternative approaches can both produce reliable estimates that are comparable within a few percent. Note that the reference approach *includes* emissions from non-energy uses. Therefore, these totals should be compared to the aggregation of fuel use and emission totals from Emissions of CO<sub>2</sub> from Fossil Fuel Combustion and Carbon Emitted from Non-Energy Uses of Fossil Fuels Annexes. These two sections together are henceforth referred to as the Sectoral Approach. Other than this distinction, the major difference between methodologies employed by each approach lies in the energy data used to derive C emissions (i.e., the actual surveyed consumption for the Sectoral Approach versus apparent consumption derived for the Reference Approach). In theory, both approaches should yield identical results. In practice, however, slight discrepancies occur. An examination of past Common Reporting Format (CRF) table submissions during UNFCCC reviews has highlighted the need to further investigate these discrepancies. The investigation found that the most recent (two to three) inventory years tend to have larger differences in consumption and emissions estimates occurring earlier in the time series. This is a result of annual energy consumption data revisions in the EIA energy statistics, and the revisions have the greatest impact on the most recent few years of inventory estimates. As a result, the differences between the Sectoral and Reference Approach decrease and are resolved over time. For the United States, these differences are discussed below.

---

<sup>115</sup> Carbon coefficients from EIA were used wherever possible. Because EIA did not provide coefficients for coal, the IPCC-recommended emission factors were used in the top-down calculations for these fuels. See notes in Table A-279 for more specific source information.

## Differences in Total Amount of Energy Consumed

Table A-282 summarizes the differences between the Reference and Sectoral Approaches in estimating total energy consumption in the United States. Although theoretically the two methods should arrive at the same estimate for U.S. energy consumption, the Reference Approach provides an energy consumption total that is 2.5 percent lower than the Sectoral Approach for 2014. The greatest differences lie in lower estimates for coal and petroleum consumption for the Reference Approach (2.5 percent and 4.7 percent, respectively) and higher estimates for natural gas consumption for the Reference Approach (0.4 percent).

There are several potential sources for the discrepancies in consumption estimates:

- *Product Definitions.* The fuel categories in the Reference Approach are different from those used in the Sectoral Approach, particularly for petroleum. For example, the Reference Approach estimates apparent consumption for crude oil. Crude oil is not typically consumed directly, but refined into other products. As a result, the United States does not focus on estimating the energy content of the various grades of crude oil, but rather estimating the energy content of the various products resulting from crude oil refining. The United States does not believe that estimating apparent consumption for crude oil, and the resulting energy content of the crude oil, is the most reliable method for the United States to estimate its energy consumption. Other differences in product definitions include using sector-specific coal statistics in the Sectoral Approach (i.e., residential, commercial, industrial coking, industrial other, and transportation coal), while the Reference Approach characterizes coal by rank (i.e., anthracite, bituminous, etc.). Also, the liquefied petroleum gas (LPG) statistics used in the bottom-up calculations are actually a composite category composed of natural gas liquids (NGL) and LPG.
- *Heat Equivalents.* It can be difficult to obtain heat equivalents for certain fuel types, particularly for categories such as "crude oil" where the key statistics are derived from thousands of producers in the United States and abroad.
- *Possible inconsistencies in U.S. Energy Data.* The United States has not focused its energy data collection efforts on obtaining the type of aggregated information used in the Reference Approach. Rather, the United States believes that its emphasis on collection of detailed energy consumption data is a more accurate methodology for the United States to obtain reliable energy data. Therefore, top-down statistics used in the Reference Approach may not be as accurately collected as bottom-up statistics applied to the Sectoral Approach.
- *Balancing Item.* The Reference Approach uses *apparent* consumption estimates while the Sectoral Approach uses *reported* consumption estimates. While these numbers should be equal, there always seems to be a slight difference that is often accounted for in energy statistics as a "balancing item."

## Differences in Estimated CO<sub>2</sub> Emissions

Given these differences in energy consumption data, the next step for each methodology involved estimating emissions of CO<sub>2</sub>. Table A-283 summarizes the differences between the two methods in estimated C emissions.

As mentioned above, for 2014, the Reference Approach resulted in a 2.5 percent lower estimate of energy consumption in the United States than the Sectoral Approach. The resulting emissions estimate for the Reference Approach was 2.3 percent lower. Estimates of natural gas emissions from the Reference Approach are higher (0.4 percent), and coal and petroleum emission estimates are lower (2.6 percent and 3.9 percent, respectively) than the Sectoral Approach. Potential reasons for these differences may include:

- *Product Definitions.* Coal data is aggregated differently in each methodology, as noted above. The format used for the Sectoral Approach likely results in more accurate estimates than in the Reference Approach. Also, the Reference Approach relies on a "crude oil" category for determining petroleum-related emissions. Given the many sources of crude oil in the United States, it is not an easy matter to track potential differences in C content between many different sources of crude; particularly since information on the C content of crude oil is not regularly collected.
- *Carbon Coefficients.* The Reference Approach relies on several default C coefficients by rank provided by IPCC (2006), while the Sectoral Approach uses annually updated category-specific coefficients by sector that are likely to be more accurate. Also, as noted above, the C coefficient for crude oil is more uncertain than that for specific secondary petroleum products, given the many sources and grades of crude oil consumed in the United States.

Although the two approaches produce similar results, the United States believes that the "bottom-up" Sectoral Approach provides a more accurate assessment of CO<sub>2</sub> emissions at the fuel level. This improvement in accuracy is largely

a result of the data collection techniques used in the United States, where there has been more emphasis on obtaining the detailed products-based information used in the Sectoral Approach than obtaining the aggregated energy flow data used in the Reference Approach. The United States believes that it is valuable to understand both methods.

## References

- EIA (2016). Supplemental Tables on Petroleum Product detail. *Monthly Energy Review, February 2016*, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2016/02).
- EIA (1995-2015). *Petroleum Supply Annual*, Energy Information Administration, U.S. Department of Energy, Washington, DC, Volume I. DOE/EIA-0340.
- EIA (1992). Coal and lignite production. *EIA State Energy Data Report 1992*, Energy Information Administration, U.S. Department of Energy, Washington, DC.
- EPA (2010). Carbon Content Coefficients Developed for EPA's Mandatory Reporting Rule. Office of Air and Radiation, Office of Atmospheric Programs, U.S. Environmental Protection Agency, Washington, D.C.
- IPCC (2006). *2006 IPCC Guidelines for National Greenhouse Gas Inventories*, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T., and Tanabe K. (eds.). Published: IGES, Japan.
- SAIC (2004). Analysis prepared by Science Applications International Corporation for EPA, Office of Air and Radiation, Market Policies Branch.
- USGS (1998). *CoalQual Database Version 2.0*, U.S. Geological Survey.

**Table A-276: 2014 U.S. Energy Statistics (Physical Units)**

| Fuel Category (Units)                    | Fuel Type                                    | Production | Imports   | Exports   | Stock Change | Adjustment | Bunkers | U.S. Territories |
|--|--|------------|-----------|-----------|--------------|------------|---------|------------------|
| <b>Solid Fuels (Thousand Short Tons)</b> | Anthracite Coal                              | 1,638      | [a]       | [a]       | [a]          |            |         |                  |
|  | Bituminous Coal                              | 451,467    | [a]       | [a]       | [a]          |            |         |                  |
|  | Sub-bituminous Coal                          | 474,797    | [a]       | [a]       | [a]          | 367        |         |                  |
|  | Lignite                                      | 72,147     | [a]       | [a]       | [a]          | 4,440      |         |                  |
|  | Coke   |            | 76        | 946       |              | (9)        |         |                  |
|  | Unspecified Coal                             |            | 11,350    | 97,257    |              | 3,052      | 2,317   | 1,653            |
| <b>Gas Fuels (Million Cubic Feet)</b>    | Natural Gas                                  | 25,767,267 | 2,695,378 | 1,514,242 | 253,464      | 283,639    |         | 58,713           |
| <b>Liquid Fuels (Thousand Barrels)</b>   | Crude Oil                                    | 3,178,555  | 2,680,626 | 128,233   | 31,268       |            |         |                  |
|  | Nat Gas Liquids and Liquefied Refinery Gases | 1,100,298  | 52,031    | 256,587   | 46,939       |            |         | 3,503            |
|  | Other Liquids                                | 0          | 409,161   | 147,584   | 22,160       |            |         |                  |
|  | Motor Gasoline                               | (102,366)  | 17,987    | 161,225   | (7,740)      |            |         | 34,763           |
|  | Aviation Gasoline                            |            | 28        | 0         | 220          |            |         |                  |
|  | Kerosene                                     |            | 7         | 3,537     | 238          |            |         | 1,262            |
|  | Jet Fuel                                     |            | 34,479    | 59,464    | 1,091        |            | 169,671 | 9,731            |
|  | Distillate Fuel                              |            | 71,050    | 401,875   | 8,743        | 397        | 6,240   | 12,856           |
|  | Residual Fuel                                |            | 63,294    | 132,798   | (4,482)      | 14,000     | 58,721  | 23,657           |
|  | Naphtha for petrochemical feedstocks         |            | 10,136    | 0         | 351          |            |         |                  |
|  | Petroleum Coke                               |            | 2,088     | 197,491   | (1,620)      | 8,331      |         |                  |
|  | Other Oil for petrochemical feedstocks       |            | 2,370     | 0         | (624)        | 1,240      |         |                  |
|  | Special Naphthas                             |            | 5,048     | 0         | 181          |            |         |                  |
|  | Lubricants                                   |            | 12,110    | 24,495    | 2,123        |            |         | 172              |
|  | Waxes  |            | 1,915     | 1,967     | (134)        |            |         |                  |
|  | Asphalt/Road Oil                             |            | 10,527    | 8,496     | (581)        |            |         |                  |
|  | Still Gas                                    |            | 0         | 0         | 0            |            |         |                  |
| Misc. Products                           |  |            | 47        | 417       | 8            |            |         | 9,061            |

[a] Included in Unspecified Coal

Note: Parentheses indicate negative values.

Sources: Solid and Gas Fuels: EIA (2016); Liquid Fuels: EIA (1995-2015).

**Table A-277: Conversion Factors to Energy Units (Heat Equivalents)**

| Fuel Category (Units)                      | Fuel Type                                    | Production | Imports | Exports | Stock Change | Adjustment | Bunkers | U.S. Territories |
|--|--|------------|---------|---------|--------------|------------|---------|------------------|
| <b>Solid Fuels (Million Btu/Short Ton)</b> | Anthracite Coal                              | 22.57      |         |         |              |            |         |                  |
|  | Bituminous Coal                              | 23.89      |         |         |              |            |         |                  |
|  | Sub-bituminous Coal                          | 17.14      |         |         |              | 28.16      |         |                  |
|  | Lignite                                      | 12.87      |         |         |              | 12.87      |         |                  |
|  | Coke   |            | 21.86   | 25.41   | 21.86        |            |         |                  |
|  | Unspecified                                  |            | 25.00   | 25.97   | 20.86        | 266.92     |         | 25.14            |
| <b>Natural Gas (BTU/Cubic Foot)</b>        |  | 1,032      | 1,025   | 1,009   | 1,032        | 1,031      |         | 1,032            |
| <b>Liquid Fuels (Million Btu/Barrel)</b>   | Crude Oil                                    | 5.80       | 6.04    | 5.80    | 5.80         |            | 5.80    | 5.80             |
|  | Nat Gas Liquids and Liquefied Refinery Gases | 3.72       | 3.72    | 3.72    | 3.72         |            | 3.72    | 3.72             |
|  | Other Liquids                                | 5.83       | 5.83    | 5.83    | 5.83         |            | 5.83    | 5.83             |
|  | Motor Gasoline                               | 5.06       | 5.06    | 5.06    | 5.06         |            | 5.06    | 5.06             |
|  | Aviation Gasoline                            |            | 5.05    | 5.05    | 5.05         |            | 5.05    | 5.05             |
|  | Kerosene                                     |            | 5.67    | 5.67    | 5.67         |            | 5.67    | 5.67             |
|  | Jet Fuel                                     |            | 5.67    | 5.67    | 5.67         |            | 5.81    | 5.67             |
|  | Distillate Fuel                              |            | 5.83    | 5.83    | 5.83         | 5.83       | 5.83    | 5.83             |
|  | Residual Oil                                 |            | 6.29    | 6.29    | 6.29         | 6.29       | 6.29    | 6.29             |
|  | Naphtha for petrochemical feedstocks         |            | 5.25    | 5.25    | 5.25         |            | 5.25    | 5.25             |
|  | Petroleum Coke                               |            | 6.02    | 6.02    | 6.02         | 6.02       | 6.02    | 6.02             |
|  | Other Oil for petrochemical feedstocks       |            | 5.83    | 5.83    | 5.83         | 5.83       | 5.83    | 5.83             |
|  | Special Naphthas                             |            | 5.25    | 5.25    | 5.25         |            | 5.25    | 5.25             |
|  | Lubricants                                   |            | 6.07    | 6.07    | 6.07         |            | 6.07    | 6.07             |
|  | Waxes  |            | 5.54    | 5.54    | 5.54         |            | 5.54    | 5.54             |
|  | Asphalt/Road Oil                             |            | 6.64    | 6.64    | 6.64         |            | 6.64    | 6.64             |
|  | Still Gas                                    |            | 6.00    | 6.00    | 6.00         |            | 6.00    | 6.00             |
| Misc. Products                             |  | 5.80       | 5.80    | 5.80    |              | 5.80       | 5.80    |                  |

Sources: Coal and lignite production: EIA (1992); Unspecified Solid Fuels, Coke, Natural Gas and Petroleum Products: EIA (1995-2015).

**Table A-278: 2014 Apparent Consumption of Fossil Fuels (Tbtu)**

| Fuel Category       | Fuel Type                                    | Production      | Imports         | Exports         | Stock Change | Adjustment     | Bunkers        | U.S. Territories | Apparent Consumption |
|---------------------|--|-----------------|-----------------|-----------------|--------------|----------------|----------------|------------------|----------------------|
| <b>Solid Fuels</b>  | Anthracite Coal                              | 37.0            |                 |                 |              |                |                |                  | 37.0                 |
|                     | Bituminous Coal                              | 10,785.5        |                 |                 |              |                |                |                  | 10,785.5             |
|                     | Sub-bituminous Coal                          | 8,138.0         |                 |                 |              | 10.3           |                |                  | 8,127.7              |
|                     | Lignite                                      | 928.2           |                 |                 |              | 57.1           |                |                  | 871.1                |
|                     | Coke   |                 | 1.7             | 24.0            | (0.2)        |                |                |                  | (22.2)               |
|                     | Unspecified                                  |                 |                 | 283.7           | 2,526.0      | 63.7           | 618.6          | 41.6             | (2,882.9)            |
| <b>Gas Fuels</b>    | Natural Gas                                  | 26,591.8        | 2,762.8         | 1,527.9         | 261.6        | 292.3          |                | 60.6             | 27,333.4             |
| <b>Liquid Fuels</b> | Crude Oil                                    | 18,435.6        | 16,177.6        | 743.8           | 181.4        |                |                |                  | 33,688.1             |
|                     | Nat Gas Liquids and Liquefied Refinery Gases | 4,096.4         | 193.7           | 955.3           | 174.8        |                |                | 13.0             | 3,173.1              |
|                     | Other Liquids                                |                 | 2,383.4         | 859.7           | 129.1        |                |                |                  | 1,394.6              |
|                     | Motor Gasoline                               | (518.0)         | 91.0            | 815.8           | (39.2)       |                |                | 175.9            | (1,027.7)            |
|                     | Aviation Gasoline                            |                 | 0.1             | 1.1             | 1.1          |                |                |                  | (2.1)                |
|                     | Kerosene                                     |                 | 0.0             | 20.1            | 1.3          |                |                | 7.2              | (14.2)               |
|                     | Jet Fuel                                     |                 | 195.5           | 337.2           | 6.2          |                |                | 985.1            | (1,077.7)            |
|                     | Distillate Fuel                              |                 | 413.9           | 2,340.9         | 50.9         |                | 2.3            | 36.3             | (1,941.8)            |
|                     | Residual Oil                                 |                 | 397.9           | 834.9           | (28.2)       |                | 88.0           | 369.2            | (717.3)              |
|                     | Naphtha for petrochemical feedstocks         |                 | 53.2            |                 | 1.8          |                |                |                  | 51.4                 |
|                     | Petroleum Coke                               |                 | 12.6            | 1,189.7         | (9.8)        |                | 50.2           |                  | (1,217.5)            |
|                     | Other Oil for petrochemical feedstocks       |                 | 13.8            |                 | (3.6)        |                | 7.2            |                  | 10.2                 |
|                     | Special Naphthas                             |                 | 26.5            |                 | 0.9          |                |                |                  | 25.5                 |
|                     | Lubricants                                   |                 | 73.4            | 148.6           | 12.9         |                |                | 1.0              | (86.9)               |
|                     | Waxes  |                 | 10.6            | 10.9            | (0.7)        |                |                |                  | 0.5                  |
|                     | Asphalt/Road Oil                             |                 | 69.9            | 56.4            | (3.9)        |                |                |                  | 17.3                 |
| Still Gas           |  |                 |                 |                 |              |                |                |                  |                      |
| Misc. Products      |  |                 | 0.3             | 2.4             | 0.0          |                |                | 52.5             | 50.3                 |
| <b>Total</b>        |  | <b>68,494.7</b> | <b>23,161.6</b> | <b>12,394.4</b> | <b>800.2</b> | <b>1,126.0</b> | <b>1,390.6</b> | <b>630.6</b>     | <b>76,575.5</b>      |

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values.

**Table A-279: 2014 Potential CO<sub>2</sub> Emissions**

| Fuel Category       | Fuel Type                              | Apparent Consumption (QBtu) | Carbon Coefficients<br>(MMT Carbon/QBtu) | Potential Emissions<br>(MMT CO <sub>2</sub> Eq.) |
|---------------------|--|-----------------------------|--|--|
| <b>Solid Fuels</b>  | Anthracite Coal                        | 0.04                        | 28.28                                    | 3.8  |
|                     | Bituminous Coal                        | 10.79                       | 25.44                                    | 1,006.1  |
|                     | Sub-bituminous Coal                    | 8.13                        | 26.50                                    | 789.8  |
|                     | Lignite                                | 0.87                        | 26.65                                    | 85.1   |
|                     | Coke                                   | (0.02)                      | 31.00                                    | (2.5)  |
|                     | Unspecified                            | (2.88)                      | 25.34                                    | (267.8)  |
| <b>Gas Fuels</b>    | Natural Gas                            | 27.33                       | 14.46                                    | 1,448.7  |
| <b>Liquid Fuels</b> | Crude Oil                              | 33.69                       | 20.31                                    | 2,508.2  |
|                     | Nat Gas Liquids and LRGs               | 3.17                        | 16.89                                    | 196.5  |
|                     | Other Liquids                          | 1.39                        | 20.31                                    | 103.8  |
|                     | Motor Gasoline                         | (1.03)                      | 19.46                                    | (73.3)   |
|                     | Aviation Gasoline                      | (0.00)                      | 18.86                                    | (0.1)  |
|                     | Kerosene                               | (0.01)                      | 19.96                                    | (1.0)  |
|                     | Jet Fuel                               | (1.08)                      | 19.70                                    | (77.8)   |
|                     | Distillate Fuel                        | (1.94)                      | 20.17                                    | (143.6)  |
|                     | Residual Oil                           | (0.72)                      | 20.48                                    | (53.9)   |
|                     | Naphtha for petrochemical feedstocks   | 0.05                        | 18.55                                    | 3.5  |
|                     | Petroleum Coke                         | (1.22)                      | 27.85                                    | (124.3)  |
|                     | Other Oil for petrochemical feedstocks | 0.01                        | 20.17                                    | 0.8  |
|                     | Special Naphthas                       | 0.03                        | 19.74                                    | 1.8  |
|                     | Lubricants                             | (0.09)                      | 20.20                                    | (6.4)  |
|                     | Waxes                                  | 0.00                        | 19.80                                    | 0.0  |
|                     | Asphalt/Road Oil                       | 0.02                        | 20.55                                    | 1.3  |
|                     | Still Gas                              | -                           | 18.20                                    | -  |
| Misc. Products      | 0.05                                   | 20.31                       | 3.7                                      |  |
| <b>Total</b>        |  |                             |  | <b>5,402.4</b>                                   |

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values.

Sources: C content coefficients by coal rank from USGS (1998) and SAIC (2004); Unspecified Solid Fuels, EIA (1995-2015), Natural Gas and Liquid Fuels: EPA (1995-2015).

**Table A-280: 2014 Non-Energy Carbon Stored in Products**

| Fuel Type                        | Consumption<br>for Non-Energy<br>Use (Tbtu) | Carbon                               |        | Carbon<br>Content<br>(MMT Carbon) | Fraction<br>Sequestered | Carbon Stored<br>(MMT CO <sub>2</sub> Eq.) |
|----------------------------------|---|--------------------------------------|--------|-----------------------------------|-------------------------|--|
|                                  |   | Coefficients<br>(MMT<br>Carbon/Qbtu) | Carbon |                                   |                         |  |
| Coal                             | 23.0  | 31.00                                | 0.71   | 0.71                              | 0.10                    | 0.9  |
| Natural Gas                      | 305.1                                       | 14.46                                | 4.41   | 4.41                              | 0.65                    | 10.6                                       |
| Asphalt & Road Oil               | 792.6                                       | 20.55                                | 16.29  | 16.29                             | 1.00                    | 59.5                                       |
| LPG                              | 2,109.4                                     | 17.06                                | 35.99  | 35.99                             | 0.65                    | 86.4                                       |
| Lubricants                       | 281.1                                       | 20.20                                | 5.68   | 5.68                              | 0.09                    | 1.9  |
| Pentanes Plus                    | 43.5  | 19.10                                | 0.83   | 0.83                              | 0.65                    | 2.0  |
| Petrochemical Feedstocks         | [a]   | [a]                                  | [a]    | [a]                               | [a]                     | 37.7                                       |
| Petroleum Coke                   | 0.0   | 27.85                                | 0.00   | 0.00                              | 0.30                    | 0.0  |
| Special Naphtha                  | 104.4                                       | 19.74                                | 2.06   | 2.06                              | 0.65                    | 4.9  |
| Waxes/Misc.                      | [a]   | [a]                                  | [a]    | [a]                               | [a]                     | 0.8  |
| Misc. U.S. Territories Petroleum | [a]   | [a]                                  | [a]    | [a]                               | [a]                     | 0.4  |
| <b>Total</b>                     |   |                                      |        |                                   |                         | <b>205.2</b>                               |

[a] Values for Misc. U.S. Territories Petroleum, Petrochemical Feedstocks and Waxes/Misc. are not shown because these categories are aggregates of numerous smaller components.

Note: Totals may not sum due to independent rounding.

**Table A-281: 2014 Reference Approach CO<sub>2</sub> Emissions from Fossil Fuel Consumption (MMT CO<sub>2</sub> Eq. unless otherwise noted)**

| Fuel Category | Potential<br>Emissions | Carbon<br>Sequestered | Net<br>Emissions | Fraction<br>Oxidized | Total<br>Emissions |
|---------------|------------------------|-----------------------|------------------|----------------------|--------------------|
| Coal          | 1,614.5                | 0.9                   | 1,613.6          | 100.0%               | 1,613.6            |
| Petroleum     | 2,339.1                | 193.7                 | 2,145.5          | 100.0%               | 2,145.5            |
| Natural Gas   | 1,448.7                | 10.6                  | 1,438.1          | 100.0%               | 1,438.1            |
| <b>Total</b>  | <b>5,402.4</b>         | <b>205.2</b>          | <b>5,197.2</b>   |                      | <b>5,197.2</b>     |

Note: Totals may not sum due to independent rounding.

**Table A-282: Fuel Consumption in the United States by Estimating Approach (Tbtu)<sup>a</sup>**

| Approach                    | 1990          | 1995          | 2000          | 2001          | 2002          | 2003          | 2004          | 2005          | 2006          | 2007          | 2008          | 2009          | 2010          | 2011          | 2012          | 2013          | 2014          |
|-----------------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <b>Sectoral</b>             | <b>69,724</b> | <b>74,941</b> | <b>82,559</b> | <b>81,135</b> | <b>81,933</b> | <b>82,332</b> | <b>83,957</b> | <b>83,919</b> | <b>82,734</b> | <b>83,937</b> | <b>81,237</b> | <b>76,443</b> | <b>78,920</b> | <b>77,497</b> | <b>75,693</b> | <b>77,762</b> | <b>78,517</b> |
| Coal                        | 18,072        | 19,187        | 21,748        | 21,121        | 21,192        | 21,625        | 21,893        | 22,187        | 21,834        | 22,067        | 21,753        | 19,231        | 20,267        | 19,071        | 16,827        | 17,459        | 17,355        |
| Natural Gas                 | 19,184        | 22,170        | 23,392        | 22,466        | 23,163        | 22,561        | 22,623        | 22,282        | 21,960        | 23,371        | 23,594        | 23,193        | 24,312        | 24,679        | 25,832        | 26,560        | 27,232        |
| Petroleum                   | 32,468        | 33,585        | 37,418        | 37,548        | 37,578        | 38,145        | 39,441        | 39,449        | 38,940        | 38,498        | 35,891        | 34,019        | 34,342        | 33,747        | 33,034        | 33,743        | 33,931        |
| <b>Reference (Apparent)</b> | <b>68,730</b> | <b>74,018</b> | <b>81,524</b> | <b>80,676</b> | <b>81,431</b> | <b>81,724</b> | <b>83,600</b> | <b>83,495</b> | <b>82,061</b> | <b>83,899</b> | <b>80,398</b> | <b>76,464</b> | <b>77,829</b> | <b>76,499</b> | <b>75,299</b> | <b>76,005</b> | <b>76,576</b> |
| Coal                        | 17,573        | 18,567        | 20,957        | 20,710        | 20,797        | 21,081        | 21,735        | 21,986        | 21,534        | 21,577        | 21,391        | 19,243        | 19,620        | 18,756        | 16,483        | 16,948        | 16,916        |
| Natural Gas                 | 19,276        | 22,274        | 23,484        | 22,535        | 23,238        | 22,630        | 22,690        | 22,349        | 22,029        | 23,441        | 23,666        | 23,277        | 24,409        | 24,778        | 25,924        | 26,637        | 27,333        |
| Petroleum                   | 31,882        | 33,177        | 37,083        | 37,431        | 37,395        | 38,013        | 39,175        | 39,160        | 38,498        | 38,881        | 35,341        | 33,944        | 33,800        | 32,965        | 32,891        | 32,420        | 32,326        |
| <b>Difference</b>           | <b>-1.4%</b>  | <b>-1.2%</b>  | <b>-1.3%</b>  | <b>-0.6%</b>  | <b>-0.6%</b>  | <b>-0.7%</b>  | <b>-0.4%</b>  | <b>-0.5%</b>  | <b>-0.8%</b>  | <b>0.0%</b>   | <b>-1.0%</b>  | <b>0.0%</b>   | <b>-1.4%</b>  | <b>-1.3%</b>  | <b>-0.5%</b>  | <b>-2.3%</b>  | <b>-2.5%</b>  |
| Coal                        | -2.8%         | -3.2%         | -3.6%         | -1.9%         | -1.9%         | -2.5%         | -0.7%         | -0.9%         | -1.4%         | -2.2%         | -1.7%         | 0.1%          | -3.2%         | -1.7%         | -2.0%         | -2.9%         | -2.5%         |
| Natural Gas                 | 0.5%          | 0.5%          | 0.4%          | 0.3%          | 0.3%          | 0.3%          | 0.3%          | 0.3%          | 0.3%          | 0.3%          | 0.3%          | 0.4%          | 0.4%          | 0.4%          | 0.4%          | 0.3%          | 0.4%          |
| Petroleum                   | -1.8%         | -1.2%         | -0.9%         | -0.3%         | -0.5%         | -0.3%         | -0.7%         | -0.7%         | -1.1%         | 1.0%          | -1.5%         | -0.2%         | -1.6%         | -2.3%         | -0.4%         | -3.9%         | -4.7%         |

<sup>a</sup> Includes U.S. Territories. Does not include international bunker fuels.

Note: Totals may not sum due to independent rounding.

**Table A-283: CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Estimating Approach (MMT CO<sub>2</sub> Eq.)<sup>a</sup>**

| Approach                    | 1990         | 1995         | 2000         | 2001         | 2002         | 2003         | 2004         | 2005         | 2006         | 2007         | 2008         | 2009         | 2010         | 2011         | 2012         | 2013         | 2014         |
|-----------------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| <b>Sectoral</b>             | <b>4,858</b> | <b>5,169</b> | <b>5,734</b> | <b>5,657</b> | <b>5,697</b> | <b>5,750</b> | <b>5,857</b> | <b>5,886</b> | <b>5,800</b> | <b>5,876</b> | <b>5,692</b> | <b>5,299</b> | <b>5,472</b> | <b>5,336</b> | <b>5,130</b> | <b>5,279</b> | <b>5,322</b> |
| Coal                        | 1,719        | 1,823        | 2,071        | 2,011        | 2,022        | 2,066        | 2,093        | 2,121        | 2,083        | 2,106        | 2,076        | 1,835        | 1,935        | 1,820        | 1,607        | 1,667        | 1,656        |
| Natural Gas                 | 1,007        | 1,164        | 1,227        | 1,178        | 1,215        | 1,183        | 1,189        | 1,172        | 1,156        | 1,230        | 1,242        | 1,221        | 1,277        | 1,297        | 1,357        | 1,397        | 1,432        |
| Petroleum                   | 2,133        | 2,182        | 2,435        | 2,468        | 2,460        | 2,501        | 2,575        | 2,593        | 2,560        | 2,540        | 2,373        | 2,243        | 2,260        | 2,219        | 2,166        | 2,215        | 2,234        |
| <b>Reference (Apparent)</b> | <b>4,793</b> | <b>5,132</b> | <b>5,683</b> | <b>5,654</b> | <b>5,697</b> | <b>5,738</b> | <b>5,883</b> | <b>5,891</b> | <b>5,782</b> | <b>5,888</b> | <b>5,653</b> | <b>5,334</b> | <b>5,407</b> | <b>5,285</b> | <b>5,127</b> | <b>5,168</b> | <b>5,197</b> |
| Coal                        | 1,654        | 1,756        | 1,988        | 1,967        | 1,976        | 2,002        | 2,065        | 2,087        | 2,049        | 2,053        | 2,036        | 1,832        | 1,868        | 1,789        | 1,573        | 1,615        | 1,614        |
| Natural Gas                 | 1,013        | 1,170        | 1,233        | 1,182        | 1,220        | 1,188        | 1,194        | 1,176        | 1,160        | 1,235        | 1,247        | 1,226        | 1,283        | 1,303        | 1,363        | 1,401        | 1,438        |
| Petroleum                   | 2,126        | 2,206        | 2,462        | 2,505        | 2,501        | 2,548        | 2,625        | 2,627        | 2,573        | 2,600        | 2,370        | 2,276        | 2,256        | 2,193        | 2,190        | 2,152        | 2,145        |
| <b>Difference</b>           | <b>-1.4%</b> | <b>-0.7%</b> | <b>-0.9%</b> | <b>-0.1%</b> | <b>0.0%</b>  | <b>-0.2%</b> | <b>0.5%</b>  | <b>0.1%</b>  | <b>-0.3%</b> | <b>0.2%</b>  | <b>-0.7%</b> | <b>0.6%</b>  | <b>-1.2%</b> | <b>-1.0%</b> | <b>-0.1%</b> | <b>-2.1%</b> | <b>-2.3%</b> |
| Coal                        | -3.8%        | -3.7%        | -4.0%        | -2.2%        | -2.3%        | -3.1%        | -1.3%        | -1.6%        | -1.7%        | -2.5%        | -1.9%        | -0.2%        | -3.4%        | -1.8%        | -2.1%        | -3.1%        | -2.6%        |
| Natural Gas                 | 0.6%         | 0.6%         | 0.5%         | 0.3%         | 0.4%         | 0.4%         | 0.4%         | 0.3%         | 0.3%         | 0.3%         | 0.3%         | 0.4%         | 0.5%         | 0.5%         | 0.4%         | 0.3%         | 0.4%         |
| Petroleum                   | -0.3%        | 1.1%         | 1.1%         | 1.5%         | 1.7%         | 1.9%         | 1.9%         | 1.3%         | 0.5%         | 2.4%         | -0.1%        | 1.5%         | -0.2%        | -1.1%        | 1.1%         | -2.9%        | -3.9%        |

<sup>a</sup> Includes U.S. Territories. Does not include international bunker fuels.

Note: Totals may not sum due to independent rounding.

# ANNEX 5 Assessment of the Sources and Sinks of Greenhouse Gas Emissions Not Included

Although this report is intended to be a comprehensive assessment of anthropogenic<sup>116</sup> sources and sinks of greenhouse gas emissions for the United States, certain sources have been identified but not included in the estimates presented for various reasons. Before discussing these sources, however, it is important to note that processes or activities that are not *anthropogenic in origin* or do not result in a *net source or sink* of greenhouse gas emissions are intentionally excluded from a national inventory of anthropogenic greenhouse gas emissions, in line with guidance from the IPCC in their guidelines for national inventories.

Given a source category that is both anthropogenic and results in net greenhouse gas emissions, reasons for not including a source related to an anthropogenic activity include one or more of the following:

- Though an estimating method has been developed, data were not adequately available to calculate emissions.
- Emissions were implicitly accounted for within another source category (e.g., CO<sub>2</sub> from Fossil Fuel Combustion).

It is also important to note that the United States believes that the sources discussed below are very low in comparison with the overall estimate of total U.S. greenhouse gas emissions, and not including them introduces a very minor bias. In general, the emission sources described in this annex are for source categories with methodologies introduced in the *2006 IPCC Guidelines* for which data collection has not been sufficient to pursue an initial estimation of greenhouse gases. Reporting of inventories to the UNFCCC under Decision 24/CP.19 requests “Where methodological or data gaps in inventories exist, information on these gaps should be presented in a transparent manner.” Furthermore, these revised reporting guidelines allow a country to indicate that a disproportionate amount of effort would be required to collect data for a gas from a specific category that would be insignificant in terms of the overall level and trend in national emissions.<sup>117</sup> Specifically, where the notation key “NE” is used in the Common Reporting Format tables that accompany this Inventory report submission to the UNFCCC, countries are required to describe why such emissions or removals have not been estimated (UNFCCC 2013).

With this guidance, the United States will consider the next steps in providing transparent information on these categories in future inventories.

## Source Categories Not Estimated

The following section is arranged by sector and source where additional explanations are available to describe the reason the source was not estimated.

### Energy

#### IPCC Category 1.A.5.a: CO<sub>2</sub> from Non-Hazardous Industrial Waste Incineration and Medical Waste Incineration

Waste incineration of the municipal waste stream and hazardous waste incineration of fossil fuel-derived materials are reported two sections of the Energy chapter of the Inventory: in the section on CO<sub>2</sub> emissions from waste incineration, and in the calculation of emissions and storage from non-energy uses of fossil fuels.

Two additional categories of waste incineration that are not included in our calculus are industrial non-hazardous waste and medical waste incineration. Data are not readily available for these sources.

An analysis was conducted based on a study of hospital/ medical/ infectious waste incinerator (HMIWI) facilities in the United States showing that medical waste incineration emissions could be considered insignificant. Based on that

---

<sup>116</sup> The term “anthropogenic,” in this context, refers to greenhouse gas emissions and removals that are a direct result of human activities or are the result of natural processes that have been affected by human activities (*2006 IPCC Guidelines for National Greenhouse Gas Inventories*).

<sup>117</sup> Paragraph 37(b) of Decision 24/CP.19 “Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention.” See < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>>.

study's information of waste throughput and an analyses of fossil-based composition of the waste, it was determined that annual greenhouse gas emissions for medical waste incineration would be below 500 kt CO<sub>2</sub> Eq./year and considered insignificant for the purposes of inventory reporting under the UNFCCC.<sup>118</sup>

#### **IPCC Category 1.B.1.a.iii: CO<sub>2</sub> from Abandoned Underground Coal Mines**

Emissions from this source are not estimated due to difficulties in obtaining data. Inclusion of emissions from this source will be investigated for future Inventories.

#### **IPCC Category 1.C CO<sub>2</sub> Transport and Storage**

Carbon dioxide is produced, captured, transported, and used for Enhanced Oil Recovery (EOR) as well as commercial and non-EOR industrial applications. This CO<sub>2</sub> is produced from both naturally-occurring CO<sub>2</sub> reservoirs and from industrial sources such as natural gas processing plants and ammonia plants. In the Inventory, emissions from naturally-produced CO<sub>2</sub> are estimated based on the specific application. In the Inventory, CO<sub>2</sub> that is used in non-EOR industrial and commercial applications (e.g., food processing, chemical production) is assumed to be emitted to the atmosphere during its industrial use. These emissions are discussed in the Carbon Dioxide Consumption section. The naturally-occurring CO<sub>2</sub> used in EOR operations is assumed to be fully sequestered. Additionally, all anthropogenic CO<sub>2</sub> emitted from natural gas processing and ammonia plants is assumed to be emitted to the atmosphere, regardless of whether the CO<sub>2</sub> is captured or not. These emissions are currently included in the Natural Gas Systems and the Ammonia Production sections of the Inventory report, respectively.

In the United States, facilities that produce CO<sub>2</sub> for various end-use applications (including capture facilities such as acid gas removal plants and ammonia plants), importers of CO<sub>2</sub>, exporters of CO<sub>2</sub>, facilities that conduct geologic sequestration of CO<sub>2</sub>, and facilities that inject CO<sub>2</sub> underground (including facilities conducting EOR), are required to report greenhouse gas data annually to EPA through its GHGRP. EPA will continue to evaluate the availability of additional GHGRP data and other opportunities for improving the emission estimates.

### **Industrial Processes and Product Use**

#### **IPCC Category 2.B.4: N<sub>2</sub>O from Caprolactam Production**

Caprolactam is a widely used chemical intermediate, primarily to produce nylon-6. All processes for producing caprolactam involve the catalytic oxidation of ammonia, with N<sub>2</sub>O being produced as a byproduct. More research is required to determine this source's significance because there is currently insufficient information available on caprolactam production to estimate emissions in the United States.

#### **IPCC Category 2.B.5.b: CO<sub>2</sub> and CH<sub>4</sub> from Calcium Carbide Production**

CO<sub>2</sub> is formed by the oxidation of petroleum coke in the production of calcium carbide. These CO<sub>2</sub> emissions are implicitly accounted for in the storage factor calculation for the non-energy use of petroleum coke in the Energy chapter. CH<sub>4</sub> may also be emitted from the production of calcium carbide because the petroleum coke used in the process contains volatile organic compounds, which form CH<sub>4</sub> during thermal decomposition. EPA will continue research to determine if calcium carbide production and consumption data are available for the United States. If these data are available, calcium carbide emission estimates will be included in this source category.

#### **IPCC Category 2.E.3: SF<sub>6</sub> Emissions from Photovoltaics**

Along with more emissions information for semiconductor manufacturing, EPA's GHGRP requires the reporting of emissions from other types of electronics manufacturing, including micro-electro-mechanical systems, flat panel displays, and photovoltaic cells. There currently are no flat panel display and photovoltaic cell manufacturing facilities that are reporting to EPA's GHGRP, and five reporting MEMs manufacturers. The MEMs manufacturers also report emissions from semiconductor manufacturing and do not distinguish between these two types of manufacturing in their report; thus, emissions from MEMs manufacturers are included in emissions from semiconductor manufacturing. Emissions from manufacturing of flat panel displays and photovoltaic cells may be included in future Inventory reports; however, estimation methodologies would need to be developed.

#### **IPCC Category 2.E.4: SF<sub>6</sub> from Heat Transfer Fluid**

Fluorinated heat transfer fluids (HTFs), of which some are liquid perfluorinated compounds, are used for temperature control, device testing, cleaning substrate surfaces and other parts, and soldering in certain types of

---

<sup>118</sup> Paragraph 37(b) of Decision 24/CP.19 "Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention." See < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>>.

semiconductor manufacturing production processes. Evaporation of these fluids is a source of fluorinated emissions (EPA 2006). Further research is needed to develop new emission factors for F-GHGs from HTFs.

## **Agriculture**

### **IPCC Category 3.A.4: CH<sub>4</sub> from Camels and Llamas Enteric Fermentation**

Enteric fermentation emissions from camels and llamas are not estimated because there is no significant population of camels and llamas in the United States. Additional analyses will be conducted to quantitatively justify emissions reporting as “not estimated” and considered insignificant.<sup>119</sup>

### **IPCC Category 3.A.4: CH<sub>4</sub> and N<sub>2</sub>O from Camels and Llamas Manure Management**

Manure management emissions from camels and llamas are not estimated because there is no significant population of camels and llamas in the United States. Additional analyses will be conducted to quantitatively justify emissions reporting as “not estimated” and considered insignificant.<sup>120</sup>

### **IPCC Category 3.F.1.2: CH<sub>4</sub> and N<sub>2</sub>O from Field Burning of Agricultural Residues—Barley, Oats, Rye, Potatoes**

There is no significant burning of barley, oats, rye, and potatoes in the United States, and therefore emissions from field burning of agricultural residues from these crops are not currently estimated. Additional analyses will be conducted to quantitatively justify emissions reporting as “not estimated” and considered insignificant.

### **IPCC Category 3.F.1.2 N<sub>2</sub>O from Prescribed Burning of Savannas—Grassland**

This Inventory does not currently include the non-CO<sub>2</sub> greenhouse gas emissions that occur with biomass burning. Grassland burning is not as common in the United States as in other regions of the world, but fires do occur through both natural ignition sources and prescribed burning.

---

<sup>119</sup> Paragraph 37(b) of Decision 24/CP.19 “Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention.” See < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>>.

<sup>120</sup> Paragraph 37(b) of Decision 24/CP.19 “Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention.” See < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>>.

# ANNEX 6 Additional Information

## 6.1. Global Warming Potential Values

Global Warming Potential (GWP) is intended as a quantified measure of the globally averaged relative radiative forcing impacts of a particular greenhouse gas. It is defined as the cumulative radiative forcing – both direct and indirect effects—integrated over a specific period of time from the emission of a unit mass of gas relative to some reference gas (IPCC 2007). Carbon dioxide (CO<sub>2</sub>) was chosen as this reference gas. Direct effects occur when the gas itself is a greenhouse gas. Indirect radiative forcing occurs when chemical transformations involving the original gas produce a gas or gases that are greenhouse gases, or when a gas influences other radiatively important processes such as the atmospheric lifetimes of other gases. The relationship between kilotons (kt) of a gas and million metric tons of CO<sub>2</sub> equivalents (MMT CO<sub>2</sub> Eq.) can be expressed as follows:

$$\text{MMT CO}_2 \text{ Eq.} = (\text{kt of gas}) \times (\text{GWP}) \times \left( \frac{\text{MMT}}{1,000 \text{ kt}} \right)$$

where,

|                         |   |   |
|-------------------------|---|---|
| MMT CO <sub>2</sub> Eq. | = | Million metric tons of Carbon Dioxide Equivalents |
| kt                      | = | Kilotons (equivalent to a thousand metric tons)   |
| GWP                     | = | Global Warming Potential                          |
| MMT                     | = | Million metric tons                               |

GWP values allow policy makers to compare the impacts of emissions and reductions of different gases. According to the IPCC, GWP values typically have an uncertainty of ±35 percent, though some GWP values have larger uncertainty than others, especially those in which lifetimes have not yet been ascertained. In the following decision, the parties to the UNFCCC have agreed to use consistent GWP values from the *IPCC Fourth Assessment Report (AR4)*, based upon a 100 year time horizon, although other time horizon values are available (see Table A-284). While noting the specific reporting requirements of the UNFCCC this Inventory uses agreed upon GWP values, it is also noted that unweighted gas emissions and sinks in kilotons (kt) are provided in the Trends chapter of this report (Table 2-2).

*...the global warming potential values used by Parties included in Annex I to the Convention (Annex I Parties) to calculate the carbon dioxide equivalence of anthropogenic emissions by sources and removals by sinks of greenhouse gases shall be those listed in the column entitled "Global warming potential for given time horizon" in table 2.14 of the errata to the contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, based on the effects of greenhouse gases over a 100-year time horizon...*<sup>121</sup>

Greenhouse gases with relatively long atmospheric lifetimes (e.g., CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs, SF<sub>6</sub>, and NF<sub>3</sub>) tend to be evenly distributed throughout the atmosphere, and consequently global average concentrations can be determined. However, the short-lived gases such as water vapor, carbon monoxide, tropospheric ozone, other indirect greenhouse gases (e.g., NO<sub>x</sub> and NMVOCs), and tropospheric aerosols (e.g., SO<sub>2</sub> products and black carbon) vary spatially, and consequently it is difficult to quantify their global radiative forcing impacts. GWP values are generally not attributed to these gases that are short-lived and spatially inhomogeneous in the atmosphere.

**Table A-284: IPCC AR4 Global Warming Potentials (GWP) and Atmospheric Lifetimes (Years) of Gases Used in this Report**

| Gas                                     | Atmospheric Lifetime      | 100-year GWP <sup>a</sup> | 20-year GWP | 500-year GWP |
|---|---------------------------|---------------------------|-------------|--------------|
| Carbon dioxide (CO <sub>2</sub> )       | See footnote <sup>b</sup> | 1                         | 1           | 1            |
| Methane (CH <sub>4</sub> ) <sup>c</sup> | 12.4 <sup>d</sup>         | 25                        | 72          | 7.6          |
| Nitrous oxide (N <sub>2</sub> O)        | 121 <sup>d</sup>          | 298                       | 289         | 153          |
| HFC-23                                  | 222                       | 14,800                    | 12,000      | 12,200       |
| HFC-32                                  | 5.2                       | 675                       | 2,330       | 205          |

<sup>121</sup> United Nations Framework Convention on Climate Change; < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf> >; 31 January 2014; Report of the Conference of the Parties at its nineteenth session; held in Warsaw from 11 to 23 November 2013; Addendum; Part two: Action taken by the Conference of the Parties at its nineteenth session; Decision 24/CP.19; Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention; p. 2. (UNFCCC 2014)

|                                 |                     |        |        |        |
|---------------------------------|---------------------|--------|--------|--------|
| HFC-125                         | 28.2                | 3,500  | 6,350  | 1,100  |
| HFC-134a                        | 13.4                | 1,430  | 3,830  | 435    |
| HFC-143a                        | 47.1                | 4,470  | 5,890  | 1,590  |
| HFC-152a                        | 1.5                 | 124    | 437    | 38     |
| HFC-227ea                       | 38.9                | 3,220  | 5,310  | 1,040  |
| HFC-236fa                       | 242                 | 9,810  | 8,100  | 7,660  |
| HFC-43-10mee                    | 16.1                | 1,640  | 4,140  | 500    |
| CF <sub>4</sub>                 | 50,000 <sup>d</sup> | 7,390  | 5,210  | 11,200 |
| C <sub>2</sub> F <sub>6</sub>   | 10,000              | 12,200 | 8,630  | 18,200 |
| C <sub>3</sub> F <sub>8</sub>   | 2,600               | 8,830  | 6,310  | 12,500 |
| C <sub>4</sub> F <sub>10</sub>  | 2,600               | 8,860  | 6,330  | 12,500 |
| c-C <sub>4</sub> F <sub>8</sub> | 3,200               | 10,300 | 7,310  | 14,700 |
| C <sub>5</sub> F <sub>12</sub>  | 4,100               | 9,160  | 6,510  | 13,300 |
| C <sub>6</sub> F <sub>14</sub>  | 3,100               | 9,300  | 6,600  | 13,300 |
| SF <sub>6</sub>                 | 3,200               | 22,800 | 16,300 | 32,600 |
| NF <sub>3</sub>                 | 500                 | 17,200 | 12,300 | 20,700 |

Source: IPCC (2007)

<sup>a</sup> GWP values used in this report are calculated over 100 year time horizon.

<sup>b</sup> For a given amount of carbon dioxide emitted, some fraction of the atmospheric increase in concentration is quickly absorbed by the oceans and terrestrial vegetation, some fraction of the atmospheric increase will only slowly decrease over a number of years, and a small portion of the increase will remain for many centuries or more.

<sup>c</sup> The methane GWP includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO<sub>2</sub> is not included.

Table A-285 presents direct GWP values for ozone-depleting substances (ODSs). Ozone-depleting substances directly absorb infrared radiation and contribute to positive radiative forcing; however, their effect as ozone-depleters also leads to a negative radiative forcing because ozone itself is a potent greenhouse gas. There is considerable uncertainty regarding this indirect effect; therefore, a range of net GWP values is provided for ozone depleting substances. The IPCC Guidelines and the UNFCCC do not include reporting instructions for estimating emissions of ODSs because their use is being phased-out under the Montreal Protocol (see note below Table A-285). The effects of these compounds on radiative forcing are not addressed in this report.

**Table A-285: 100-year Direct Global Warming Potentials for Select Ozone Depleting Substances**

| Gas                              | Direct GWP |
|----------------------------------|------------|
| CFC-11                           | 4,750      |
| CFC-12                           | 10,900     |
| CFC-113                          | 6,130      |
| HCFC-22                          | 1,810      |
| HCFC-123                         | 77         |
| HCFC-124                         | 609        |
| HCFC-141b                        | 725        |
| HCFC-142b                        | 2,310      |
| CH <sub>3</sub> CCl <sub>3</sub> | 146        |
| CCl <sub>4</sub>                 | 1,400      |
| CH <sub>3</sub> Br               | 5          |
| Halon-1211                       | 1,890      |
| Halon-1301                       | 7,140      |

Source: IPCC (2007)

Note: Because these compounds have been shown to deplete stratospheric ozone, they are typically referred to as ozone depleting substances (ODSs). However, they are also potent greenhouse gases. Recognizing the harmful effects of these compounds on the ozone layer, in 1987 many governments signed the *Montreal Protocol on Substances that Deplete the Ozone Layer* to limit the production and importation of a number of CFCs and other halogenated compounds. The United States furthered its commitment to phase-out ODSs by signing and ratifying the Copenhagen Amendments to the Montreal Protocol in 1992. Under these amendments, the United States committed to ending the production and importation of halons by 1994, and CFCs by 1996.

The IPCC published its *Fifth Assessment Report (AR5)* in 2013, providing the most current and comprehensive scientific assessment of climate change (IPCC 2013). Within this report, the GWP values were revised relative to the IPCC's *Second Assessment Report (SAR)* (IPCC 1996), IPCC's *Third Assessment Report (TAR)* (IPCC 2001), and the IPCC's *Fourth Assessment Report (AR4)* (IPCC 2007). Although the AR4 GWP values are used throughout this report in line with UNFCCC inventory reporting guidelines, it is informative to review the changes to the GWP values and the impact they have on the total GWP-weighted emissions of the United States. All GWP values use CO<sub>2</sub> as a reference gas; a change in

the radiative efficiency of CO<sub>2</sub> thus impacts the GWP of all other greenhouse gases. Since the SAR and TAR, the IPCC has applied an improved calculation of CO<sub>2</sub> radiative forcing and an improved CO<sub>2</sub> response function. The GWP values are drawn from IPCC (2007), with updates for those cases where new laboratory or radiative transfer results have been published. Additionally, the atmospheric lifetimes of some gases have been recalculated, and updated background concentrations were used. Table A-286 shows how the GWP values of the other gases relative to CO<sub>2</sub> tend to be larger in AR4 and AR5 because the revised radiative forcing of CO<sub>2</sub> is lower than in earlier assessments, taking into account revisions in lifetimes. Comparisons of GWP values are based on the 100-year time horizon required for UNFCCC inventory reporting. However, there were some instances in which other variables, such as the radiative efficiency or the chemical lifetime, were altered that resulted in further increases or decreases in particular GWP values. In addition, the values for radiative forcing and lifetimes have been calculated for a variety of halocarbons. Updates in some well-mixed HFC compounds (including HFC-23, HFC-32, HFC-134a, and HFC-227ea) for AR4 result from investigation into radiative efficiencies in these compounds, with some GWP values changing by up to 40 percent; with this change, the uncertainties associated with these well-mixed HFCs are thought to be approximately 12 percent.

It should be noted that the use of IPCC AR4 GWP values for the current Inventory applies across the entire time series of the Inventory (i.e., from 1990 to 2014). As such, GWP comparisons throughout this chapter are presented relative to AR4 GWPs.

**Table A-286: Comparison of GWP values and Lifetimes Used in the SAR, AR4, and AR5**

| Gas                                     | Lifetime (years) |                      |        | GWP (100 year) |        |                  |                                 | Difference in GWP (Relative to AR4) |         |                  |         |                                 |                                     |
|---|------------------|----------------------|--------|----------------|--------|------------------|---------------------------------|-------------------------------------|---------|------------------|---------|---------------------------------|-------------------------------------|
|   | SAR              | AR4                  | AR5    | SAR            | AR4    | AR5 <sup>a</sup> | AR5 with feedbacks <sup>b</sup> | SAR                                 | SAR (%) | AR5 <sup>a</sup> | AR5 (%) | AR5 with feedbacks <sup>b</sup> | AR5 with feedbacks <sup>b</sup> (%) |
| Carbon dioxide (CO <sub>2</sub> )       | c                | d                    | d      | 1              | 1      | 1                | 1                               | NC                                  | NC      | NC               | NC      | NC                              | NC                                  |
| Methane (CH <sub>4</sub> ) <sup>d</sup> | 12±3             | 8.7/12 <sup>e</sup>  | 12.4   | 21             | 25     | 28               | 34                              | (4)                                 | (16%)   | 3                | 12%     | 9                               | 36%                                 |
| Nitrous oxide (N <sub>2</sub> O)        | 120              | 120/114 <sup>e</sup> | 121    | 310            | 298    | 265              | 298                             | 12                                  | 4%      | (33)             | (11%)   | 0                               | 0%                                  |
| <b>Hydrofluorocarbons</b>               |                  |                      |        |                |        |                  |                                 |                                     |         |                  |         |                                 |                                     |
| HFC-23                                  | 264              | 270                  | 222    | 11,700         | 14,800 | 12,400           | 13,856                          | (3,100)                             | (21%)   | (2,400)          | (16%)   | (944)                           | (6)%                                |
| HFC-32                                  | 5.6              | 4.9                  | 5.2    | 650            | 675    | 677              | 817                             | (25)                                | (4%)    | 2                | +       | 142                             | 21%                                 |
| HFC-125                                 | 32.6             | 29                   | 28.2   | 2,800          | 3,500  | 3,170            | 3,691                           | (700)                               | (20%)   | (330)            | (9%)    | 191                             | 5%                                  |
| HFC-134a                                | 14.6             | 14                   | 13.4   | 1,300          | 1,430  | 1,300            | 1,549                           | (130)                               | (9%)    | (130)            | (9%)    | 119                             | 8%                                  |
| HFC-143a                                | 48.3             | 52                   | 47.1   | 3,800          | 4,470  | 4,800            | 5,508                           | (670)                               | (15%)   | 330              | 7%      | 1,038                           | 23%                                 |
| HFC-152a                                | 1.5              | 1.4                  | 1.5    | 140            | 124    | 138              | 167                             | 16                                  | 13%     | 14               | 11%     | 43                              | 35%                                 |
| HFC-227ea                               | 36.5             | 34.2                 | 38.9   | 2,900          | 3,220  | 3,350            | 3,860                           | (320)                               | (10%)   | 130              | 4%      | 640                             | 20%                                 |
| HFC-236fa                               | 209              | 240                  | 242    | 6,300          | 9,810  | 8,060            | 8,998                           | (3,510)                             | (36%)   | (1,750)          | (18%)   | (812)                           | (8)%                                |
| HFC-245fa                               | NA               | 7.6                  | 7.7    | NA             | 1,030  | 858              | 1032                            | NA                                  | NA      | (172)            | (17%)   | 2                               | +                                   |
| HFC-365mfc                              | NA               | 6.6                  | 8.7    | NA             | 794    | 804              | 966                             | NA                                  | NA      | 10               | 1%      | 172                             | 22%                                 |
| HFC-43-10mee                            | 17.1             | 15.9                 | 16.1   | 1,300          | 1,640  | 1,650            | 1,952                           | (340)                               | (21%)   | 10               | 1%      | 312                             | 19%                                 |
| <b>Fully Fluorinated Species</b>        |                  |                      |        |                |        |                  |                                 |                                     |         |                  |         |                                 |                                     |
| SF <sub>6</sub>                         | 3,200            | 3,200                | 3,200  | 23,900         | 22,800 | 23,500           | 26,087                          | 1,100                               | 5%      | 700              | 3%      | 3,287                           | 14%                                 |
| CF <sub>4</sub>                         | 50,000           | 50,000               | 50,000 | 6,500          | 7,390  | 6,630            | 7,349                           | (890)                               | (12%)   | (760)            | (10%)   | (41)                            | (1)%                                |
| C <sub>2</sub> F <sub>6</sub>           | 10,000           | 10,000               | 10,000 | 9,200          | 12,200 | 11,100           | 12,340                          | (3,000)                             | (25%)   | (1,100)          | (9%)    | 140                             | 1%                                  |
| C <sub>3</sub> F <sub>8</sub>           | 2,600            | 2,600                | 2,600  | 7,000          | 8,830  | 8,900            | 9,878                           | (1,830)                             | (21%)   | 70               | 1%      | 1,048                           | 12%                                 |
| C <sub>4</sub> F <sub>10</sub>          | 2,600            | 2,600                | 2,600  | 7,000          | 8,860  | 9,200            | 10,213                          | (1,860)                             | (21%)   | 340              | 4%      | 1,353                           | 15%                                 |
| c-C <sub>4</sub> F <sub>8</sub>         | 3,200            | 3,200                | 3,200  | 8,700          | 10,300 | 9,540            | 10,592                          | (1,600)                             | (16%)   | (760)            | (7%)    | 292                             | 3%                                  |
| C <sub>5</sub> F <sub>12</sub>          | 4,100            | 4,100                | 4,100  | 7,500          | 9,160  | 8,550            | 9,484                           | (1,660)                             | (18%)   | (610)            | (7%)    | 324                             | 4%                                  |
| C <sub>6</sub> F <sub>14</sub>          | 3,200            | 3,200                | 3,100  | 7,400          | 9,300  | 7,910            | 8,780                           | (1,900)                             | (20%)   | (1,390)          | (15%)   | (520)                           | (6)%                                |
| NF <sub>3</sub>                         | NA               | 740                  | 500    | NA             | 17,200 | 16,100           | 17,885                          | NA                                  | NA      | (1,100)          | (6%)    | 685                             | 4%                                  |

+ Does not exceed 0.05 or 0.05 percent.

NC (No Change)

NA (Not Applicable)

<sup>a</sup> The GWP values presented here are the ones most consistent with the methodology used in the AR4 report.

<sup>b</sup> The GWP values presented here from the AR5 report include climate-carbon feedbacks for the non-CO<sub>2</sub> gases in order to be consistent with the approach used in calculating the CO<sub>2</sub> lifetime. Additionally, the AR5 reported separate values for fossil versus biogenic methane in order to account for the CO<sub>2</sub> oxidation product.

<sup>c</sup> For a given amount of carbon dioxide emitted, some fraction of the atmospheric increase in concentration is quickly absorbed by the oceans and terrestrial vegetation, some fraction of the atmospheric increase will only slowly decrease over a number of years, and a small portion of the increase will remain for many centuries or more.

<sup>d</sup> No single lifetime can be determined for CO<sub>2</sub>. (See IPCC 2001)

<sup>e</sup> The methane GWP includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO<sub>2</sub> is only included in the value from AR5 that includes climate-carbon feedbacks.

<sup>f</sup> Methane and nitrous oxide have chemical feedback systems that can alter the length of the atmospheric response, in these cases, global mean atmospheric lifetime (LT) is given first, followed by perturbation time (PT).

Note: Parentheses indicate negative values. Source: IPCC (2013), IPCC (2007), IPCC (2001), IPCC (1996).

The choice of GWP values between the SAR, AR4, and AR5 with or without climate-carbon feedbacks has an impact on both the overall emissions estimated by the Inventory, as well as the trend in emissions over time. To summarize, Table A-287 shows the overall trend in U.S. greenhouse gas emissions, by gas, from 1990 through 2014 using the four GWP sets. The table also presents the impact of SAR and AR5 GWP values with or without feedbacks on the total emissions for 1990 and for 2014.

**Table A-287: Effects on U.S. Greenhouse Gas Emissions Using SAR, AR4, and AR5 GWP values (MMT CO<sub>2</sub>Eq.)**

| Gas  | Trend from 1990 to 2014 |              |                  |                  | Revisions to Annual Emission Estimates (Relative to AR4) |             |              |                  |             |              |
|--|-------------------------|--------------|------------------|------------------|--|-------------|--------------|------------------|-------------|--------------|
|  |                         |              |                  |                  | SAR  |             |              | AR5 <sup>a</sup> |             |              |
|  | SAR                     | AR4          | AR5 <sup>a</sup> | AR5 <sup>b</sup> | 1990   |             |              | 2014             |             |              |
| CO <sub>2</sub>                                    | 440.9                   | 440.9        | 440.9            | 440.9            | NC   | NC          | NC           | NC               | NC          | NC           |
| CH <sub>4</sub>                                    | (36.1)                  | (43.0)       | (48.2)           | (58.5)           | (123.8)  | 92.9        | 278.6        | (116.9)          | 87.7        | 263.1        |
| N <sub>2</sub> O                                   | (2.8)                   | (2.7)        | (2.4)            | (2.7)            | 16.4   | (45.0)      | 0.0          | 16.2             | (44.7)      | 0.0          |
| HFCs, PFCs, SF <sub>6</sub><br>and NF <sub>3</sub> | 60.8                    | 78.1         | 71.3             | 88.3             | (11.8)   | (8.9)       | 1.6          | (29.2)           | (15.7)      | 11.7         |
| <b>Total</b>                                       | <b>462.7</b>            | <b>473.3</b> | <b>461.7</b>     | <b>468.0</b>     | <b>(119.3)</b>   | <b>38.9</b> | <b>280.2</b> | <b>(129.9)</b>   | <b>27.3</b> | <b>274.8</b> |
| <b>Percent Change</b>                              | <b>7.4%</b>             | <b>7.4%</b>  | <b>7.2%</b>      | <b>7.0%</b>      | <b>(1.9)%</b>  | <b>0.6%</b> | <b>4.4%</b>  | <b>(1.9)%</b>    | <b>0.4%</b> | <b>4.0%</b>  |

NC (No Change)

<sup>a</sup> The GWP values presented here are the ones most consistent with the methodology used in the AR4 report, and exclude climate-carbon feedbacks.

<sup>b</sup> The GWP values presented here from the AR5 report include climate-carbon feedbacks for the non-CO<sub>2</sub> gases in order to be consistent with the approach used in calculating the CO<sub>2</sub> lifetime. Additionally, the AR5 reported separate values for fossil versus biogenic methane in order to account for the CO<sub>2</sub> oxidation product.

Note: Totals may not sum due to independent rounding. Excludes sinks. Parentheses indicate negative values.

When the GWP values from the SAR are applied to the emission estimates presented in this report, total emissions for the year 2014 are 6,740.6 MMT CO<sub>2</sub> Eq., as compared to the official emission estimate of 6,870.5 MMT CO<sub>2</sub> Eq. using AR4 GWP values (i.e., the use of SAR GWPs results in a 1.9 percent decrease relative to emissions estimated using AR4 GWPs). Table A-288 provides a detailed summary of U.S. greenhouse gas emissions and sinks for 1990 through 2014, using the GWP values from the SAR. The percent change in emissions is equal to the percent change in the GWP; however, in cases where multiple gases are emitted in varying amounts the percent change is variable over the years, such as with substitutes for ozone depleting substances. Table A-289 summarizes the resulting change in emissions from using SAR GWP values relative to emissions using AR4 values for 1990 through 2014, including the percent change for 2014.

**Table A-288: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks using the SAR GWP values (MMT CO<sub>2</sub> Eq.)**

| Gas/Source                                | 1990           | 2005           | 2010           | 2011           | 2012           | 2013           | 2014           |
|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| <b>CO<sub>2</sub></b>                     | <b>5,115.1</b> | <b>6,122.7</b> | <b>5,688.8</b> | <b>5,559.5</b> | <b>5,349.2</b> | <b>5,502.6</b> | <b>5,556.0</b> |
| Fossil Fuel Combustion                    | 4,740.7        | 5,747.1        | 5,358.3        | 5,227.7        | 5,024.7        | 5,157.6        | 5,208.2        |
| <i>Electricity Generation</i>             | 1,820.8        | 2,400.9        | 2,258.4        | 2,157.7        | 2,022.2        | 2,038.1        | 2,039.3        |
| <i>Transportation</i>                     | 1,493.8        | 1,887.0        | 1,728.3        | 1,707.6        | 1,696.8        | 1,713.0        | 1,737.6        |
| <i>Industrial</i>                         | 842.5          | 828.0          | 775.5          | 773.3          | 782.9          | 812.2          | 813.3          |
| <i>Residential</i>                        | 338.3          | 357.8          | 334.6          | 326.8          | 282.5          | 329.7          | 345.1          |
| <i>Commercial</i>                         | 217.4          | 223.5          | 220.1          | 220.7          | 196.7          | 221.0          | 231.9          |
| <i>U.S. Territories</i>                   | 27.9           | 49.9           | 41.4           | 41.5           | 43.6           | 43.5           | 41.0           |
| Non-Energy Use of Fuels                   | 118.1          | 138.9          | 114.1          | 108.5          | 105.6          | 121.7          | 114.3          |
| Iron and Steel Production & Metallurgical |                |                |                |                |                |                |                |
| Coke Production                           | 99.7           | 66.5           | 55.7           | 59.9           | 54.2           | 52.2           | 55.4           |
| Natural Gas Systems                       | 37.7           | 30.1           | 32.4           | 35.7           | 35.2           | 38.5           | 42.4           |
| Cement Production                         | 33.3           | 45.9           | 31.3           | 32.0           | 35.1           | 36.1           | 38.8           |
| Petrochemical Production                  | 21.6           | 27.4           | 27.2           | 26.3           | 26.5           | 26.4           | 26.5           |
| Lime Production                           | 11.7           | 14.6           | 13.4           | 14.0           | 13.7           | 14.0           | 14.1           |
| Other Process Uses of Carbonates          | 4.9            | 6.3            | 9.6            | 9.3            | 8.0            | 10.4           | 12.1           |
| Ammonia Production                        | 13.0           | 9.2            | 9.2            | 9.3            | 9.4            | 10.0           | 9.4            |
| Incineration of Waste                     | 8.0            | 12.5           | 11.0           | 10.5           | 10.4           | 9.4            | 9.4            |
| Carbon Dioxide Consumption                | 1.5            | 1.4            | 4.4            | 4.1            | 4.0            | 4.2            | 4.5            |

|   |              |              |              |              |              |              |              |
|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Urea Consumption for Non-Agricultural Purposes            | 3.8          | 3.7          | 4.7          | 4.0          | 4.4          | 4.2          | 4.0          |
| Petroleum Systems   | 3.6          | 3.9          | 4.2          | 4.2          | 3.9          | 3.7          | 3.6          |
| Aluminum Production                                       | 6.8          | 4.1          | 2.7          | 3.3          | 3.4          | 3.3          | 2.8          |
| Soda Ash Production and Consumption                       | 2.8          | 3.0          | 2.7          | 2.7          | 2.8          | 2.8          | 2.8          |
| Ferroalloy Production                                     | 2.2          | 1.4          | 1.7          | 1.7          | 1.9          | 1.8          | 1.9          |
| Titanium Dioxide Production                               | 1.2          | 1.8          | 1.8          | 1.7          | 1.5          | 1.7          | 1.8          |
| Glass Production  | 1.5          | 1.9          | 1.5          | 1.3          | 1.2          | 1.3          | 1.3          |
| Phosphoric Acid Production                                | 1.5          | 1.3          | 1.1          | 1.2          | 1.1          | 1.1          | 1.1          |
| Zinc Production   | 0.6          | 1.0          | 1.2          | 1.3          | 1.5          | 1.4          | 1.0          |
| Lead Production   | 0.5          | 0.6          | 0.5          | 0.5          | 0.5          | 0.5          | 0.5          |
| Silicon Carbide Production and Consumption                | 0.4          | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          |
| Magnesium Production and Processing                       | +            | +            | +            | +            | +            | +            | +            |
| Wood Biomass and Ethanol Consumption <sup>a</sup>         | 219.4        | 229.8        | 265.1        | 268.1        | 267.7        | 286.3        | 293.7        |
| International Bunker Fuels <sup>b</sup>                   | 103.5        | 113.1        | 117.0        | 111.7        | 105.8        | 99.8         | 103.2        |
| <b>CH<sub>4</sub></b>                                     | <b>650.0</b> | <b>602.6</b> | <b>606.8</b> | <b>602.6</b> | <b>600.1</b> | <b>606.0</b> | <b>613.9</b> |
| Natural Gas Systems                                       | 173.7        | 149.0        | 139.6        | 142.9        | 145.0        | 147.5        | 147.9        |
| Enteric Fermentation                                      | 137.9        | 141.8        | 143.9        | 141.9        | 140.1        | 139.0        | 138.0        |
| Landfills   | 150.8        | 129.4        | 119.4        | 121.3        | 119.5        | 121.2        | 124.3        |
| Petroleum Systems   | 32.5         | 41.0         | 45.4         | 47.3         | 49.0         | 54.4         | 57.2         |
| Coal Mining   | 81.1         | 53.9         | 69.2         | 59.8         | 55.8         | 54.3         | 56.8         |
| Manure Management   | 31.2         | 47.3         | 51.2         | 51.7         | 53.5         | 51.5         | 51.4         |
| Wastewater Treatment                                      | 13.1         | 13.4         | 13.0         | 12.8         | 12.6         | 12.4         | 12.3         |
| Rice Cultivation  | 11.0         | 10.9         | 10.0         | 9.9          | 10.0         | 10.0         | 10.0         |
| Stationary Combustion                                     | 7.1          | 6.2          | 5.9          | 5.9          | 5.6          | 6.7          | 6.8          |
| Abandoned Underground Coal Mines                          | 6.0          | 5.5          | 5.5          | 5.4          | 5.2          | 5.2          | 5.3          |
| Composting  | 0.3          | 1.6          | 1.5          | 1.6          | 1.6          | 1.7          | 1.7          |
| Mobile Combustion   | 4.7          | 2.3          | 1.9          | 1.9          | 1.8          | 1.8          | 1.7          |
| Field Burning of Agricultural Residues                    | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          |
| Petrochemical Production                                  | 0.2          | 0.1          | +            | +            | 0.1          | 0.1          | 0.1          |
| Ferroalloy Production                                     | +            | +            | +            | +            | +            | +            | +            |
| Silicon Carbide Production and Consumption                | +            | +            | +            | +            | +            | +            | +            |
| Iron and Steel Production & Metallurgical Coke Production | +            | +            | +            | +            | +            | +            | +            |
| Incineration of Waste                                     | +            | +            | +            | +            | +            | +            | +            |
| International Bunker Fuels <sup>b</sup>                   | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          |
| <b>N<sub>2</sub>O</b>                                     | <b>422.6</b> | <b>413.6</b> | <b>426.8</b> | <b>433.3</b> | <b>425.8</b> | <b>419.6</b> | <b>419.8</b> |
| Agricultural Soil Management                              | 315.5        | 309.1        | 333.6        | 336.1        | 336.1        | 331.5        | 331.2        |
| Stationary Combustion                                     | 12.4         | 21.0         | 23.1         | 22.2         | 22.2         | 23.8         | 24.4         |
| Manure Management   | 14.6         | 17.2         | 17.9         | 18.1         | 18.2         | 18.2         | 18.2         |
| Mobile Combustion   | 42.9         | 35.8         | 24.6         | 23.3         | 20.8         | 18.9         | 17.0         |
| Nitric Acid Production                                    | 12.6         | 11.8         | 12.0         | 11.3         | 10.9         | 11.1         | 11.4         |
| Adipic Acid Production                                    | 15.8         | 7.4          | 4.4          | 10.6         | 5.8          | 4.1          | 5.7          |
| Wastewater Treatment                                      | 3.5          | 4.5          | 4.7          | 4.9          | 4.9          | 5.0          | 5.0          |
| N <sub>2</sub> O from Product Uses                        | 4.4          | 4.4          | 4.4          | 4.4          | 4.4          | 4.4          | 4.4          |
| Composting  | 0.4          | 1.7          | 1.7          | 1.7          | 1.8          | 1.9          | 1.9          |
| Incineration of Waste                                     | 0.5          | 0.4          | 0.3          | 0.3          | 0.3          | 0.3          | 0.3          |
| Semiconductor Manufacture                                 | +            | 0.1          | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          |
| Field Burning of Agricultural Residues                    | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          | 0.1          |
| International Bunker Fuels <sup>b</sup>                   | 0.9          | 1.0          | 1.0          | 1.0          | 1.0          | 0.9          | 0.9          |
| <b>HFCs</b>   | <b>36.9</b>  | <b>102.2</b> | <b>126.2</b> | <b>129.5</b> | <b>130.3</b> | <b>132.3</b> | <b>138.2</b> |
| Substitution of Ozone Depleting Substances <sup>c</sup>   | 0.3          | 86.2         | 119.7        | 122.4        | 125.8        | 128.8        | 133.9        |
| HFC-22 Production   | 36.4         | 15.8         | 6.4          | 6.9          | 4.3          | 3.2          | 4.0          |
| Semiconductor Manufacture                                 | 0.2          | 0.2          | 0.2          | 0.2          | 0.2          | 0.1          | 0.2          |
| Magnesium Production and Processing                       | +            | +            | +            | +            | +            | 0.1          | 0.1          |
| <b>PFCs</b>   | <b>20.6</b>  | <b>5.5</b>   | <b>3.7</b>   | <b>5.8</b>   | <b>5.3</b>   | <b>4.8</b>   | <b>5.0</b>   |
| Semiconductor Manufacture                                 | 18.4         | 3.0          | 1.6          | 2.9          | 2.8          | 2.5          | 2.5          |
| Aluminum Production                                       | 2.2          | 2.6          | 2.1          | 2.9          | 2.5          | 2.3          | 2.5          |
| <b>SF<sub>6</sub></b>                                     | <b>32.6</b>  | <b>14.7</b>  | <b>9.9</b>   | <b>10.5</b>  | <b>7.9</b>   | <b>7.6</b>   | <b>7.7</b>   |

|  |                |                |                |                |                |                |                |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Electrical Transmission and Distribution | 26.6           | 11.1           | 7.4            | 7.1            | 5.8            | 5.7            | 5.9            |
| Magnesium Production and Processing      | 5.4            | 2.9            | 2.2            | 2.9            | 1.7            | 1.5            | 1.1            |
| Semiconductor Manufacture                | 0.5            | 0.7            | 0.4            | 0.4            | 0.4            | 0.4            | 0.8            |
| <b>NF<sub>3</sub></b>                    | <b>NA</b>      |
| Semiconductor Manufacture                | NA             |
| <b>Total Emissions</b>                   | <b>6,277.9</b> | <b>7,261.4</b> | <b>6,862.3</b> | <b>6,741.2</b> | <b>6,518.7</b> | <b>6,672.9</b> | <b>6,740.6</b> |
| <b>LULUCF Emissions<sup>d</sup></b>      | <b>14.7</b>    | <b>27.0</b>    | <b>17.5</b>    | <b>22.1</b>    | <b>31.0</b>    | <b>23.2</b>    | <b>23.7</b>    |
| <b>LULUCF Total Net Flux<sup>e</sup></b> | <b>(753.0)</b> | <b>(726.7)</b> | <b>(784.3)</b> | <b>(784.9)</b> | <b>(782.0)</b> | <b>(783.7)</b> | <b>(787.0)</b> |
| <b>LULUCF Sector Total<sup>f</sup></b>   | <b>(738.3)</b> | <b>(699.7)</b> | <b>(766.7)</b> | <b>(762.8)</b> | <b>(751.0)</b> | <b>(760.4)</b> | <b>(763.3)</b> |
| <b>Net Emissions (Sources and Sinks)</b> | <b>5,539.5</b> | <b>6,561.7</b> | <b>6,095.5</b> | <b>5,978.5</b> | <b>5,767.7</b> | <b>5,912.4</b> | <b>5,977.2</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Note: Total emissions presented without LULUCF. Net emissions presented with LULUCF.

<sup>a</sup> Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry.

<sup>b</sup> Emissions from International Bunker Fuels are not included in totals.

<sup>c</sup> Small amounts of PFC emissions also result from this source.

<sup>d</sup> LULUCF emissions include the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions reported for Non-CO<sub>2</sub> Emissions from Forest Fires, N<sub>2</sub>O Fluxes from Forest Soils, CO<sub>2</sub> Emissions from Agricultural Liming, CO<sub>2</sub> Emissions from Urea Fertilization, Peatlands Remaining Peatlands, and N<sub>2</sub>O Fluxes from Settlement Soils.

<sup>e</sup> Net CO<sub>2</sub> flux is the net C stock change from the following categories: *Forest Land Remaining Forest Land, Land Converted to Forest Land, Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, Land Converted to Grassland, Settlements Remaining Settlements, and Other.*

Notes: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

**Table A-289: Change in U.S. Greenhouse Gas Emissions Using SAR GWP values relative to AR4 GWP values (MMT CO<sub>2</sub> Eq.)**

| Gas/Source  | 2010-2014      |                |                |                |                |                |                | Percent        |
|---|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
|   | 1990           | 2005           | 2010           | 2011           | 2012           | 2013           | 2014           | Change in 2014 |
| <b>CO<sub>2</sub></b>                                     | <b>NC</b>      |
| <b>CH<sub>4</sub></b>                                     | <b>(123.8)</b> | <b>(114.8)</b> | <b>(115.6)</b> | <b>(114.8)</b> | <b>(114.3)</b> | <b>(115.4)</b> | <b>(116.9)</b> | <b>(8.0%)</b>  |
| Natural Gas Systems                                       | (33.1)         | (28.4)         | (26.6)         | (27.2)         | (27.6)         | (28.1)         | (28.2)         | (8.0%)         |
| Enteric Fermentation                                      | (26.3)         | (27.0)         | (27.4)         | (27.0)         | (26.7)         | (26.5)         | (26.3)         | (8.0%)         |
| Landfills   | (28.7)         | (24.6)         | (22.7)         | (23.1)         | (22.8)         | (23.1)         | (23.7)         | (8.0%)         |
| Petroleum Systems   | (6.2)          | (7.8)          | (8.7)          | (9.0)          | (9.3)          | (10.4)         | (10.9)         | (8.0%)         |
| Coal Mining   | (15.4)         | (10.3)         | (13.2)         | (11.4)         | (10.6)         | (10.3)         | (10.8)         | (8.0%)         |
| Manure Management   | (5.9)          | (9.0)          | (9.7)          | (9.8)          | (10.2)         | (9.8)          | (9.8)          | (8.0%)         |
| Wastewater Treatment                                      | (2.5)          | (2.5)          | (2.5)          | (2.4)          | (2.4)          | (2.4)          | (2.4)          | (8.0%)         |
| Rice Cultivation  | (2.1)          | (2.1)          | (1.9)          | (1.9)          | (1.9)          | (1.9)          | (1.9)          | (8.0%)         |
| Stationary Combustion                                     | (1.4)          | (1.2)          | (1.1)          | (1.1)          | (1.1)          | (1.3)          | (1.3)          | (8.0%)         |
| Abandoned Underground Coal Mines                          | (1.2)          | (1.1)          | (1.1)          | (1.0)          | (1.0)          | (1.0)          | (1.0)          | (8.0%)         |
| Composting  | (0.1)          | (0.3)          | (0.3)          | (0.3)          | (0.3)          | (0.3)          | (0.3)          | (8.0%)         |
| Mobile Combustion   | (0.9)          | (0.4)          | (0.4)          | (0.4)          | (0.3)          | (0.3)          | (0.3)          | (8.0%)         |
| Field Burning of Agricultural Residues                    | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| Petrochemical Production                                  | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| Ferroalloy Production                                     | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| Silicon Carbide Production and Consumption                | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| Iron and Steel Production & Metallurgical Coke Production | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| Incineration of Waste                                     | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| <i>International Bunker Fuels<sup>a</sup></i>             | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (+)            | (8.0%)         |
| <b>N<sub>2</sub>O</b>                                     | <b>16.4</b>    | <b>16.0</b>    | <b>16.5</b>    | <b>16.8</b>    | <b>16.5</b>    | <b>16.2</b>    | <b>16.2</b>    | <b>(0.7%)</b>  |
| Agricultural Soil Management                              | 12.2           | 12.0           | 12.9           | 13.0           | 13.0           | 12.8           | 12.8           | (0.7%)         |
| Stationary Combustion                                     | 0.5            | 0.8            | 0.9            | 0.9            | 0.9            | 0.9            | 0.9            | (0.7%)         |
| Manure Management   | 0.6            | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            | 0.7            | (0.7%)         |
| Mobile Combustion   | 1.7            | 1.4            | 1.0            | 0.9            | 0.8            | 0.7            | 0.7            | (0.7%)         |
| Nitric Acid Production                                    | 0.5            | 0.5            | 0.5            | 0.4            | 0.4            | 0.4            | 0.4            | (0.7%)         |
| Adipic Acid Production                                    | 0.6            | 0.3            | 0.2            | 0.4            | 0.2            | 0.2            | 0.2            | (0.7%)         |

|   |               |               |               |               |               |               |               |                |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|----------------|
| Wastewater Treatment                                    | 0.1           | 0.2           | 0.2           | 0.2           | 0.2           | 0.2           | 0.2           | (0.7%)         |
| N <sub>2</sub> O from Product Uses                      | 0.2           | 0.2           | 0.2           | 0.2           | 0.2           | 0.2           | 0.2           | (0.7%)         |
| Composting  | +             | 0.1           | 0.1           | 0.1           | 0.1           | 0.1           | 0.1           | (0.7%)         |
| Incineration of Waste                                   | +             | +             | +             | +             | +             | +             | +             | (0.7%)         |
| Semiconductor Manufacture                               | +             | +             | +             | +             | +             | +             | +             | (0.7%)         |
| Field Burning of Agricultural Residues                  | +             | +             | +             | +             | +             | +             | +             | (0.7%)         |
| International Bunker Fuels <sup>a</sup>                 | +             | +             | +             | +             | +             | +             | +             | (0.7%)         |
| <b>HFCs</b>   | <b>(9.7)</b>  | <b>(17.7)</b> | <b>(23.2)</b> | <b>(24.8)</b> | <b>(25.5)</b> | <b>(26.7)</b> | <b>(28.5)</b> | <b>(10.1%)</b> |
| Substitution of Ozone Depleting Substances <sup>b</sup> | 0.0           | (13.5)        | (21.5)        | (22.9)        | (24.3)        | (25.8)        | (27.3)        | (9.8%)         |
| HCFC-22 Production                                      | (9.7)         | (4.2)         | (1.7)         | (1.8)         | (1.1)         | (0.9)         | (1.1)         | (18.9%)        |
| Semiconductor Manufacture                               | (+)           | (+)           | (+)           | (+)           | (+)           | (+)           | (0.1)         | (18.9%)        |
| Magnesium Production and Processing                     | +             | +             | (+)           | (+)           | (+)           | (+)           | (+)           | (23.1%)        |
| <b>PFCs</b>   | <b>(3.6)</b>  | <b>(1.1)</b>  | <b>(0.8)</b>  | <b>(1.2)</b>  | <b>(0.7)</b>  | <b>(1.0)</b>  | <b>(0.6)</b>  | <b>(9.5%)</b>  |
| Semiconductor Manufacture                               | 15.6          | (0.3)         | (1.1)         | (0.6)         | (0.2)         | (0.4)         | (0.5)         | (12.4%)        |
| Aluminum Production                                     | (19.2)        | (0.9)         | 0.3           | (0.6)         | (0.4)         | (0.6)         | +             | (6.1%)         |
| <b>SF<sub>6</sub></b>                                   | <b>1.5</b>    | <b>0.7</b>    | <b>0.5</b>    | <b>0.5</b>    | <b>0.4</b>    | <b>0.3</b>    | <b>0.4</b>    | <b>(2.6%)</b>  |
| Electrical Transmission and Distribution                | 1.2           | 0.5           | 0.3           | 0.3           | 0.3           | 0.3           | 0.3           | (2.6%)         |
| Magnesium Production and Processing                     | 0.3           | 0.1           | 0.1           | 0.1           | 0.1           | 0.1           | +             | (2.6%)         |
| Semiconductor Manufacture                               | +             | +             | +             | +             | +             | +             | +             | (2.6%)         |
| <b>NF<sub>3</sub></b>                                   | <b>NA</b>     | <b>(37.2%)</b> |
| Semiconductor Manufacture                               | NA            | (37.2%)        |
| <b>Total Emissions</b>                                  | <b>(78.7)</b> | <b>(76.3)</b> | <b>(77.4)</b> | <b>(77.9)</b> | <b>(76.8)</b> | <b>(77.7)</b> | <b>(78.8)</b> | <b>(1.8%)</b>  |

NC (No Change)

NA (Not Applicable)

+ Absolute value does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup>Emissions from International Bunker Fuels are not included in totals.

<sup>b</sup>Small amounts of PFC emissions also result from this source.

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values.

Table A-290 below shows a comparison of total emissions estimates by sector using both the IPCC SAR and AR4 GWP values. For most sectors, the change in emissions that result from using SAR relative to AR4 GWP values was minimal. The effect on emissions from waste was by far the greatest (15.2 percent decrease in 2014 using SAR GWP values, relative to emissions using AR4 GWP values), due the predominance of CH<sub>4</sub> emissions in this sector. Emissions from all other sectors were comprised of mainly CO<sub>2</sub> or a mix of gases, which moderated the effect of the changes.

**Table A-290: Comparison of Emissions by Sector using IPCC AR4 and SAR GWP Values (MMT CO<sub>2</sub>Eq.)**

| Sector                                      | 1990    | 2005    | 2010    | 2011    | 2012    | 2013    | 2014    |
|---|---------|---------|---------|---------|---------|---------|---------|
| <b>Energy</b>                               |         |         |         |         |         |         |         |
| AR4 GWP, Used In Inventory                  | 5,324.9 | 6,294.5 | 5,884.6 | 5,744.0 | 5,533.9 | 5,693.5 | 5,746.2 |
| SAR GWP                                     | 5,269.0 | 6,247.6 | 5,835.5 | 5,695.6 | 5,485.6 | 5,643.8 | 5,695.3 |
| Difference (%)                              | (1.1%)  | (0.7%)  | (0.8%)  | (0.8%)  | (0.9%)  | (0.9%)  | (0.9%)  |
| <b>Industrial Processes and Product Use</b> |         |         |         |         |         |         |         |
| AR4 GWP, Used In Inventory                  | 340.9   | 354.3   | 353.0   | 370.5   | 360.1   | 363.5   | 379.2   |
| SAR GWP                                     | 330.3   | 336.5   | 329.6   | 345.3   | 334.4   | 336.3   | 350.8   |
| Difference (%)                              | (3.1%)  | (5.0%)  | (6.6%)  | (6.8%)  | (7.1%)  | (7.5%)  | (7.5%)  |
| <b>Agriculture</b>                          |         |         |         |         |         |         |         |
| AR4 GWP, Used In Inventory                  | 532.0   | 552.2   | 582.3   | 583.1   | 583.3   | 575.3   | 573.6   |
| SAR GWP                                     | 510.5   | 526.7   | 556.8   | 558.0   | 558.2   | 550.5   | 549.1   |
| Difference (%)                              | (4.1%)  | (4.6%)  | (4.4%)  | (4.3%)  | (4.3%)  | (4.3%)  | (4.3%)  |
| <b>LULUCF</b>                               |         |         |         |         |         |         |         |
| AR4 GWP, Used In Inventory                  | (738.0) | (698.5) | (766.4) | (762.0) | (749.7) | (759.6) | (762.5) |
| SAR GWP                                     | (738.3) | (699.7) | (766.7) | (762.8) | (751.0) | (760.4) | (763.3) |
| Difference (%)                              | 0.1%    | 0.2%    | +       | 0.1%    | 0.2%    | 0.1%    | 0.1%    |
| <b>Waste</b>                                |         |         |         |         |         |         |         |
| AR4 GWP, Used In Inventory                  | 199.3   | 177.8   | 165.5   | 167.8   | 165.7   | 167.8   | 171.4   |
| SAR GWP                                     | 168.1   | 150.6   | 140.3   | 142.3   | 140.5   | 142.3   | 145.3   |
| Difference (%)                              | (15.6%) | (15.3%) | (15.3%) | (15.2%) | (15.2%) | (15.2%) | (15.2%) |
| <b>Net Emissions</b>                        |         |         |         |         |         |         |         |
| AR4 GWP, Used In Inventory                  | 5,659.2 | 6,680.3 | 6,219.0 | 6,103.4 | 5,893.3 | 6,040.4 | 6,108.0 |
| SAR GWP                                     | 5,539.5 | 6,561.7 | 6,095.5 | 5,978.5 | 5,767.7 | 5,912.4 | 5,977.2 |

**Difference (%)** (2.1%) (1.8%) (2.0%) (2.0%) (2.1%) (2.1%) (2.1%)

+ Does not exceed 0.05 percent.

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values.

Further, Table A-291 and Table A-292 show the comparison of emission estimates using AR5 GWP values relative to AR4 GWP values without climate-carbon feedbacks, on an emissions and percent change basis. Table A-293 and Table A-294 show the comparison of emission estimates using AR5 GWP values with climate-carbon feedbacks. The use of AR5 GWP values without climate-carbon feedbacks<sup>122</sup> results in an increase in emissions of CH<sub>4</sub> and SF<sub>6</sub> relative to AR4 GWP values, but a decrease in emissions of other gases. The use of AR5 GWP values with climate-carbon feedbacks does not impact N<sub>2</sub>O emissions; however, it results in an increase in emissions of CH<sub>4</sub> and SF<sub>6</sub> relative to AR4 GWP values, and a decrease in emissions of other gases. Overall, these comparisons of AR4 and AR5 GWP values do not have a significant effect on U.S. emission trends, resulting in an increase in emissions of less than 1 percent using AR5 GWP values, or 4 percent when using AR5 GWP values with climate-carbon feedbacks. As with the comparison of SAR and AR4 GWP values presented above, the percent change in emissions is equal to the percent change in the GWP for each gas; however, in cases where multiple gases are emitted in varying amounts the percent change is variable over the years, such as with substitutes for ozone depleting substances.

**Table A-291: Change in U.S. Greenhouse Gas Emissions Using AR5<sup>a</sup> without Climate-Carbon Feedbacks Relative to AR4 GWP Values (MMT CO<sub>2</sub> Eq.)**

| Gas              | 1990        | 2005        | 2010        | 2011        | 2012        | 2013        | 2014        |
|------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| CO <sub>2</sub>  | NC          |
| CH <sub>4</sub>  | 92.9        | 86.1        | 86.7        | 86.1        | 85.7        | 86.6        | 87.7        |
| N <sub>2</sub> O | (45.0)      | (44.0)      | (45.4)      | (46.1)      | (45.3)      | (44.7)      | (44.7)      |
| HFCs             | (7.5)       | (13.4)      | (14.7)      | (15.0)      | (14.8)      | (14.9)      | (15.8)      |
| PFCs             | (2.4)       | (0.6)       | (0.4)       | (0.7)       | (0.2)       | (0.6)       | (0.2)       |
| SF <sub>6</sub>  | 1.0         | 0.4         | 0.3         | 0.3         | 0.2         | 0.2         | 0.2         |
| NF <sub>3</sub>  | (+)         | (+)         | (+)         | (+)         | (+)         | (+)         | (+)         |
| <b>Total</b>     | <b>38.9</b> | <b>28.4</b> | <b>26.4</b> | <b>24.5</b> | <b>25.5</b> | <b>26.6</b> | <b>27.3</b> |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

NC (No change)

Note: Total emissions presented without LULUCF.

<sup>a</sup> The GWP values presented here are the ones most consistent with the methodology used in the AR4 report. The AR5 report has also calculated GWP values (shown in Table A-293) where climate-carbon feedbacks have been included for the non-CO<sub>2</sub> gases in order to be consistent with the approach used in calculating the CO<sub>2</sub> lifetime. Additionally, the AR5 reported separate values for fossil versus biogenic methane in order to account for the CO<sub>2</sub> oxidation product.

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values.

**Table A-292: Change in U.S. Greenhouse Gas Emissions Using AR5<sup>a</sup> without Climate-Carbon Feedbacks Relative to AR4 GWP Values (Percent)**

| Gas/Source                                       | 1990    | 2005    | 2010    | 2011    | 2012    | 2013    | 2014    |
|--|---------|---------|---------|---------|---------|---------|---------|
| CO <sub>2</sub>                                  | NC      |
| CH <sub>4</sub>                                  | 12.0%   | 12.0%   | 12.0%   | 12.0%   | 12.0%   | 12.0%   | 12.0%   |
| N <sub>2</sub> O                                 | (11.1%) | (11.1%) | (11.1%) | (11.1%) | (11.1%) | (11.1%) | (11.1%) |
| HFCs   | (16.0%) | (11.2%) | (9.8%)  | (9.7%)  | (9.5%)  | (9.4%)  | (9.4%)  |
| Substitution of Ozone Depleting Substances       | 11.3%   | (10.2%) | (9.4%)  | (9.3%)  | (9.3%)  | (9.2%)  | (9.2%)  |
| HCFC-22 Production <sup>a</sup>                  | (16.2%) | (16.2%) | (16.2%) | (16.2%) | (16.2%) | (16.2%) | (16.2%) |
| Semiconductor Manufacture <sup>b</sup>           | (16.2%) | (16.2%) | (16.2%) | (16.2%) | (16.2%) | (16.2%) | (16.2%) |
| Magnesium Production and Processing <sup>c</sup> | 0.0%    | 0.0%    | (9.1%)  | (9.1%)  | (9.1%)  | (9.1%)  | (9.1%)  |
| PFCs   | (10.0%) | (9.6%)  | (9.4%)  | (9.4%)  | (3.7%)  | (10.2%) | (3.3%)  |
| Semiconductor Manufacture <sup>b</sup>           | (9.4%)  | (9.1%)  | (9.1%)  | (8.9%)  | (9.1%)  | (9.1%)  | (9.2%)  |

<sup>122</sup> The IPCC AR5 report provides additional information on emission metrics. See <[https://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5\\_Chapter08\\_FINAL.pdf](https://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_Chapter08_FINAL.pdf)>.

|                                  |             |             |             |             |             |             |             |
|----------------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Aluminum Production <sup>d</sup> | (10.1%)     | (10.1%)     | (10.0%)     | (10.0%)     | 2.0%        | (11.2%)     | 3.8%        |
| SF <sub>6</sub>                  | 3.1%        | 3.1%        | 3.1%        | 3.1%        | 3.1%        | 3.1%        | 3.1%        |
| NF <sub>3</sub>                  | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      |
| <b>Total</b>                     | <b>0.6%</b> | <b>0.4%</b> | <b>0.4%</b> | <b>0.4%</b> | <b>0.4%</b> | <b>0.4%</b> | <b>0.4%</b> |

Note: Total emissions presented without LULUCF.

NC (No change)

<sup>a</sup> The GWP values presented here are the ones most consistent with the methodology used in the AR4 report. The AR5 report has also calculated GWP values (shown in Table A-294) where climate-carbon feedbacks have been included for the non-CO<sub>2</sub> gases in order to be consistent with the approach used in calculating the CO<sub>2</sub> lifetime. Additionally, the AR5 reported separate values for fossil versus biogenic methane in order to account for the CO<sub>2</sub> oxidation product.

<sup>a</sup> HFC-23 emitted

<sup>b</sup> Emissions from HFC-23, CF<sub>4</sub>, C<sub>2</sub>F<sub>6</sub>, C<sub>3</sub>F<sub>8</sub>, SF<sub>6</sub>, and NF<sub>3</sub>.

<sup>c</sup> Zero change in beginning of time series since emissions were zero.

<sup>d</sup> PFC emissions from CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub>

Note: Parentheses indicate negative values.

**Table A-293: Change in U.S. Greenhouse Gas Emissions Using AR5 with Climate-Carbon Feedbacks<sup>a</sup> Relative to AR4 GWP Values (MMT CO<sub>2</sub> Eq.)**

| Gas              | 1990         | 2005         | 2010         | 2011         | 2012         | 2013         | 2014         |
|------------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| CO <sub>2</sub>  | NC           |
| CH <sub>4</sub>  | 278.6        | 258.2        | 260.1        | 258.3        | 257.2        | 259.7        | 263.1        |
| N <sub>2</sub> O | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          | 0.0          |
| HFCs             | (2.9)        | 5.0          | 8.9          | 9.1          | 9.6          | 10.0         | 10.2         |
| PFCs             | (+)          | +            | +            | +            | 0.4          | (+)          | 0.4          |
| SF <sub>6</sub>  | 4.5          | 2.0          | 1.4          | 1.4          | 1.1          | 1.0          | 1.1          |
| NF <sub>3</sub>  | +            | +            | +            | +            | +            | +            | +            |
| <b>Total</b>     | <b>280.2</b> | <b>265.3</b> | <b>270.4</b> | <b>268.9</b> | <b>268.4</b> | <b>270.8</b> | <b>274.8</b> |

Note: Total emissions presented without LULUCF.

NC (No Change)

+ Absolute value does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> The GWP values presented here from the AR5 report include climate-carbon feedbacks for the non-CO<sub>2</sub> gases in order to be consistent with the approach used in calculating the CO<sub>2</sub> lifetime. Additionally, the AR5 reported separate values for fossil versus biogenic methane in order to account for the CO<sub>2</sub> oxidation product.

Notes: Totals may not sum due to independent rounding. Parentheses indicate negative values.

**Table A-294: Change in U.S. Greenhouse Gas Emissions Using AR5 with Climate-Carbon Feedbacks<sup>a</sup> Relative to AR4 GWP Values (Percent)**

| Gas/Source                                       | 1990        | 2005        | 2010        | 2011        | 2012        | 2013        | 2014        |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| CO <sub>2</sub>                                  | NC          |
| CH <sub>4</sub>                                  | 36.0%       | 36.0%       | 36.0%       | 36.0%       | 36.0%       | 36.0%       | 36.0%       |
| N <sub>2</sub> O                                 | 0.0%        | 0.0%        | 0.0%        | 0.0%        | 0.0%        | 0.0%        | 0.0%        |
| HFCs   | (6.1%)      | 4.2%        | 6.0%        | 5.9%        | 6.2%        | 6.3%        | 6.1%        |
| Substitution of Ozone Depleting Substances       | 34.7%       | 6.3%        | 6.7%        | 6.7%        | 6.7%        | 6.6%        | 6.6%        |
| HFC-22 Production <sup>a</sup>                   | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      |
| Semiconductor Manufacture <sup>b</sup>           | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      | (6.4%)      |
| Magnesium Production and Processing <sup>c</sup> | 0.0%        | 0.0%        | 8.3%        | 8.3%        | 8.3%        | 8.3%        | 8.3%        |
| PFCs   | (0.2%)      | 0.3%        | 0.5%        | 0.5%        | 6.9%        | (0.3%)      | 7.3%        |
| Semiconductor Manufacture <sup>b</sup>           | 0.6%        | 0.9%        | 0.9%        | 1.1%        | 0.9%        | 0.9%        | 0.8%        |
| Aluminum Production <sup>d</sup>                 | (0.3%)      | (0.3%)      | (0.1%)      | (0.2%)      | 13.2%       | (1.5%)      | 15.1%       |
| SF <sub>6</sub>                                  | 14.4%       | 14.4%       | 14.4%       | 14.4%       | 14.4%       | 14.4%       | 14.4%       |
| NF <sub>3</sub>                                  | 4.0%        | 4.0%        | 4.0%        | 4.0%        | 4.0%        | 4.0%        | 4.0%        |
| <b>Total</b>                                     | <b>4.4%</b> | <b>3.6%</b> | <b>3.9%</b> | <b>3.9%</b> | <b>4.0%</b> | <b>4.0%</b> | <b>4.0%</b> |

Note: Total emissions presented without LULUCF.

NC (No Change)

<sup>a</sup> The GWP values presented here from the AR5 report include climate-carbon feedbacks for the non-CO<sub>2</sub> gases in order to be consistent with the approach used in calculating the CO<sub>2</sub> lifetime. Additionally, the AR5 reported separate values for fossil versus biogenic methane in order to account for the CO<sub>2</sub> oxidation product.

<sup>a</sup> HFC-23 emitted

<sup>b</sup> Emissions from HFC-23, CF<sub>4</sub>, C<sub>2</sub>F<sub>6</sub>, C<sub>3</sub>F<sub>8</sub>, SF<sub>6</sub>, and NF<sub>3</sub>.

<sup>c</sup> Zero change in beginning of time series since emissions were zero.

<sup>d</sup> PFC emissions from CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub>

Notes: Excludes Sinks. Parentheses indicate negative values.

## 6.2. Ozone Depleting Substance Emissions

Ozone is present in both the stratosphere,<sup>123</sup> where it shields the earth from harmful levels of ultraviolet radiation, and at lower concentrations in the troposphere,<sup>124</sup> where it is the main component of anthropogenic photochemical “smog.” Chlorofluorocarbons (CFCs), halons, carbon tetrachloride, methyl chloroform, and hydrochlorofluorocarbons (HCFCs), along with certain other chlorine and bromine containing compounds, have been found to deplete the ozone levels in the stratosphere. These compounds are commonly referred to as ozone depleting substances (ODSs). If left unchecked, stratospheric ozone depletion could result in a dangerous increase of ultraviolet radiation reaching the earth’s surface. In 1987, nations around the world signed the *Montreal Protocol on Substances that Deplete the Ozone Layer*. This landmark agreement created an international framework for limiting, and ultimately eliminating, the production of most ozone depleting substances. ODSs have historically been used in a variety of industrial applications, including refrigeration and air conditioning, foam blowing, fire extinguishing, sterilization, solvent cleaning, and as an aerosol propellant.

In the United States, the Clean Air Act Amendments of 1990 provide the legal instrument for implementation of the *Montreal Protocol* controls. The Clean Air Act classifies ozone depleting substances as either Class I or Class II, depending upon the ozone depletion potential (ODP) of the compound.<sup>125</sup> The production of CFCs, halons, carbon tetrachloride, and methyl chloroform – all Class I substances – has already ended in the United States. However, large amounts of these chemicals remain in existing equipment,<sup>126</sup> and stockpiles of the ODSs, as well as material recovered from equipment being decommissioned, are used for maintaining the existing equipment. As a result, emissions of Class I compounds will continue, albeit in ever decreasing amounts, for many more years. Class II designated substances, all of which are HCFCs, have been, or are being, phased out at later dates than Class I compounds because they have lower ozone depletion potentials. These compounds served, and in some cases continue to serve, as interim replacements for Class I compounds in many industrial applications. The use and emissions of HCFCs in the United States is anticipated to continue for several decades as equipment that use Class II substances are retired from use. Under current controls, however, the production for domestic use of all HCFCs in the United States will end by the year 2030.

In addition to contributing to ozone depletion, CFCs, halons, carbon tetrachloride, methyl chloroform, and HCFCs are also potent greenhouse gases. However, the depletion of the ozone layer has a cooling effect on the climate that counteracts the direct warming from tropospheric emissions of ODSs. Stratospheric ozone influences the earth’s radiative balance by absorption and emission of longwave radiation from the troposphere as well as absorption of shortwave radiation from the sun; overall, stratospheric ozone has a warming effect.

The IPCC has prepared both direct GWP values and net (combined direct warming and indirect cooling) GWP ranges for some of the most common ozone depleting substances (IPCC 2007). See Annex 6.1, Global Warming Potential Values, for a listing of the net GWP values for ODS.

Although the IPCC emission inventory guidelines do not require the reporting of emissions of ozone depleting substances, the United States believes that no inventory is complete without the inclusion of these compounds. Emission estimates for several ozone depleting substances are provided in Table A-295.

**Table A-295: Emissions of Ozone Depleting Substances (kt)**

| Compound             | 1990 | 2005 | 2010 | 2011 | 2012 | 2013 | 2014 |
|----------------------|------|------|------|------|------|------|------|
| <b>Class I</b>       |      |      |      |      |      |      |      |
| CFC-11               | 29   | 12   | 8    | 8    | 8    | 7    | 7    |
| CFC-12               | 124  | 23   | 3    | 3    | 2    | 2    | 1    |
| CFC-113              | 59   | +    | +    | +    | +    | +    | +    |
| CFC-114              | 5    | +    | +    | +    | +    | +    | +    |
| CFC-115              | 8    | 2    | +    | +    | +    | +    | +    |
| Carbon Tetrachloride | 4    | +    | +    | +    | +    | +    | +    |

<sup>123</sup> The stratosphere is the layer from the top of the troposphere up to about 50 kilometers. Approximately 90 percent of atmospheric ozone is within the stratosphere. The greatest concentration of ozone occurs in the middle of the stratosphere, in a region commonly called the ozone layer.

<sup>124</sup> The troposphere is the layer from the ground up to about 11 kilometers near the poles and 16 kilometers in equatorial regions (i.e., the lowest layer of the atmosphere, where humans live). It contains roughly 80 percent of the mass of all gases in the atmosphere and is the site for weather processes including most of the water vapor and clouds.

<sup>125</sup> Substances with an ozone depletion potential of 0.2 or greater are designated as Class I. All other designated substances that deplete stratospheric ozone but which have an ODP of less than 0.2 are Class II.

<sup>126</sup> Older refrigeration and air-conditioning equipment, fire extinguishing systems, meter-dose inhalers, and foam products blown with CFCs/HCFCs may still contain ODS.

|                   |     |    |    |    |    |    |    |
|-------------------|-----|----|----|----|----|----|----|
| Methyl Chloroform | 223 | +  | +  | +  | +  | +  | +  |
| Halon-1211        | 2   | 1  | 1  | 1  | 1  | 1  | 1  |
| Halon-1301        | 2   | +  | +  | +  | +  | +  | +  |
| <b>Class II</b>   |     |    |    |    |    |    |    |
| HCFC-22           | 49  | 82 | 82 | 80 | 76 | 73 | 69 |
| HCFC-123          | +   | 1  | 1  | 1  | 1  | 1  | 1  |
| HCFC-124          | +   | 2  | 1  | 1  | 1  | 1  | 1  |
| HCFC-141b         | 1   | 4  | 9  | 9  | 9  | 10 | 10 |
| HCFC-142b         | 2   | 3  | 1  | 1  | 1  | 1  | 1  |
| HCFC-225ca/cb     | +   | +  | +  | +  | +  | +  | +  |

+ Does not exceed 0.5 kt.

## Methodology and Data Sources

Emissions of ozone depleting substances were estimated using the EPA's Vintaging Model. The model, named for its method of tracking the emissions of annual "vintages" of new equipment that enter into service, is a "bottom-up" model. It models the consumption of chemicals based on estimates of the quantity of equipment or products sold, serviced, and retired each year, and the amount of the chemical required to manufacture and/or maintain the equipment. The Vintaging Model makes use of this market information to build an inventory of the in-use stocks of the equipment in each of the end-uses. Emissions are estimated by applying annual leak rates, service emission rates, and disposal emission rates to each population of equipment. By aggregating the emission and consumption output from the different end-uses, the model produces estimates of total annual use and emissions of each chemical. Please see Annex 3.9, Methodology for Estimating HFC and PFC Emissions from Substitution of Ozone Depleting Substances, of this Inventory for a more detailed discussion of the Vintaging Model.

## Uncertainties

Uncertainties exist with regard to the levels of chemical production, equipment sales, equipment characteristics, and end-use emissions profiles that are used by these models. Please see the ODS Substitutes section of this report for a more detailed description of the uncertainties that exist in the Vintaging Model.

### 6.3. Sulfur Dioxide Emissions

Sulfur dioxide (SO<sub>2</sub>), emitted into the atmosphere through natural and anthropogenic processes, affects the Earth's radiative budget through photochemical transformation into sulfate aerosols that can (1) scatter sunlight back to space, thereby reducing the radiation reaching the Earth's surface; (2) affect cloud formation; and (3) affect atmospheric chemical composition (e.g., stratospheric ozone, by providing surfaces for heterogeneous chemical reactions). The overall effect of SO<sub>2</sub>-derived aerosols on radiative forcing is believed to be negative (IPCC 2007). However, because SO<sub>2</sub> is short-lived and unevenly distributed through the atmosphere, its radiative forcing impacts are highly uncertain. Sulfur dioxide emissions have been provided below in Table A-296.

The major source of SO<sub>2</sub> emissions in the United States is the burning of sulfur containing fuels, mainly coal. Metal smelting and other industrial processes also release significant quantities of SO<sub>2</sub>. The largest contributor to U.S. emissions of SO<sub>2</sub> is electricity generation, accounting for 64.0 percent of total SO<sub>2</sub> emissions in 2014 (see Table A-297); coal combustion accounted for approximately 92.0 percent of that total. The second largest source was industrial fuel combustion, which produced 13.3 percent of 2014 SO<sub>2</sub> emissions. Overall, SO<sub>2</sub> emissions in the United States decreased by 78.4 percent from 1990 to 2014. The majority of this decline came from reductions from electricity generation, primarily due to increased consumption of low sulfur coal from surface mines in western states.

Sulfur dioxide is important for reasons other than its effect on radiative forcing. It is a major contributor to the formation of urban smog and acid rain. As a contributor to urban smog, high concentrations of SO<sub>2</sub> can cause significant increases in acute and chronic respiratory diseases. In addition, once SO<sub>2</sub> is emitted, it is chemically transformed in the atmosphere and returns to earth as the primary contributor to acid deposition, or acid rain. Acid rain has been found to accelerate the decay of building materials and paints, cause the acidification of lakes and streams, and damage trees. As a result of these harmful effects, the United States has regulated the emissions of SO<sub>2</sub> under the Clean Air Act. The EPA has also developed a strategy to control these emissions via four programs: (1) the National Ambient Air Quality Standards program,<sup>127</sup> (2) New Source Performance Standards,<sup>128</sup> (3) the New Source Review/Prevention of Significant Deterioration Program,<sup>129</sup> and (4) the Sulfur Dioxide Allowance Program.<sup>130</sup>

**Table A-296: SO<sub>2</sub> Emissions (kt)**

| Sector/Source                               | 1990          | 2005          | 2010         | 2011         | 2012         | 2013         | 2014         |
|---|---------------|---------------|--------------|--------------|--------------|--------------|--------------|
| <b>Energy</b>                               | <b>19,628</b> | <b>12,364</b> | <b>6,396</b> | <b>5,273</b> | <b>4,106</b> | <b>4,020</b> | <b>3,923</b> |
| Stationary Sources                          | 18,407        | 11,541        | 6,120        | 5,008        | 3,859        | 3,790        | 3,710        |
| Oil and Gas Activities                      | 390           | 180           | 117          | 108          | 108          | 108          | 108          |
| Mobile Sources                              | 793           | 619           | 144          | 142          | 125          | 108          | 90           |
| Waste Combustion                            | 38            | 25            | 16           | 15           | 15           | 15           | 15           |
| <b>Industrial Processes and Product Use</b> | <b>1,307</b>  | <b>831</b>    | <b>617</b>   | <b>604</b>   | <b>604</b>   | <b>604</b>   | <b>604</b>   |
| Miscellaneous                               | 11            | 114           | 146          | 179          | 179          | 179          | 179          |
| Other Industrial Processes                  | 362           | 327           | 190          | 171          | 171          | 171          | 171          |
| Metals Processing                           | 659           | 158           | 141          | 131          | 131          | 131          | 131          |
| Chemical and Allied Product                 |               |               |              |              |              |              |              |
| Manufacturing                               | 269           | 228           | 132          | 115          | 115          | 115          | 115          |
| Storage and Transport                       | 6             | 2             | 7            | 8            | 8            | 8            | 8            |
| <b>Solvent Use</b>                          | <b>0</b>      | <b>+</b>      | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     |
| Degreasing                                  | 0             | 0             | 0            | 0            | 0            | 0            | 0            |
| Graphic Arts                                | 0             | 0             | 0            | 0            | 0            | 0            | 0            |
| Dry Cleaning                                | NA            | 0             | 0            | 0            | 0            | 0            | 0            |
| Surface Coating                             | 0             | 0             | 0            | 0            | 0            | 0            | 0            |
| Other Industrial                            | 0             | +             | +            | +            | +            | +            | +            |
| Nonindustrial                               | NA            | NA            | NA           | NA           | NA           | NA           | NA           |
| <b>Agriculture</b>                          | <b>NA</b>     | <b>NA</b>     | <b>NA</b>    | <b>NA</b>    | <b>NA</b>    | <b>NA</b>    | <b>NA</b>    |
| Agricultural Burning                        | NA            | NA            | NA           | NA           | NA           | NA           | NA           |
| <b>Waste</b>                                | <b>+</b>      | <b>1</b>      | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     | <b>+</b>     |
| Landfills                                   | +             | 1             | +            | +            | +            | +            | +            |
| Wastewater Treatment                        | +             | 0             | 0            | 0            | 0            | 0            | 0            |

<sup>127</sup> [42 U.S.C § 7409, CAA § 109]

<sup>128</sup> [42 U.S.C § 7411, CAA § 111]

<sup>129</sup> [42 U.S.C § 7473, CAA § 163]

<sup>130</sup> [42 U.S.C § 7651, CAA § 401]

|               |   |               |               |              |              |              |              |              |
|---------------|---|---------------|---------------|--------------|--------------|--------------|--------------|--------------|
| Miscellaneous | + | 0             | 0             | 0            | 0            | 0            | 0            |              |
| <b>Total</b>  |   | <b>20,935</b> | <b>13,196</b> | <b>7,014</b> | <b>5,877</b> | <b>4,711</b> | <b>4,625</b> | <b>4,528</b> |

+ Does not exceed 0.5 kt\* Miscellaneous includes other combustion and fugitive dust categories.

Note: Totals may not sum due to independent rounding.

Source: Data taken from EPA (2015) and disaggregated based on EPA (2003).

**Table A-297: SO<sub>2</sub> Emissions from Electricity Generation (kt)**

| Fuel Type                 | 1990          | 2005         | 2010         | 2011         | 2012         | 2013         | 2014         |
|---------------------------|---------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Coal                      | 13,808        | 8,680        | 4,752        | 3,859        | 2,802        | 2,739        | 2,665        |
| Oil                       | 580           | 458          | 251          | 204          | 148          | 144          | 141          |
| Gas                       | 1             | 174          | 95           | 77           | 56           | 55           | 53           |
| Misc. Internal Combustion | 45            | 57           | 31           | 25           | 18           | 18           | 18           |
| Other                     | NA            | 71           | 39           | 31           | 23           | 22           | 22           |
| <b>Total</b>              | <b>14,433</b> | <b>9,439</b> | <b>5,168</b> | <b>4,196</b> | <b>3,047</b> | <b>2,978</b> | <b>2,898</b> |

Note: Totals may not sum due to independent rounding.

Source: Data taken from EPA (2015) and disaggregated based on EPA (2003).

## 6.4. Complete List of Source Categories

| Chapter/Source  | Gas(es)   |
|---|---|
| <b>Energy</b>   |   |
| Fossil Fuel Combustion                                    | CO <sub>2</sub>   |
| Non-Energy Use of Fossil Fuels                            | CO <sub>2</sub>   |
| Stationary Combustion (excluding CO <sub>2</sub> )        | CH <sub>4</sub> , N <sub>2</sub> O, CO, NO <sub>x</sub> , NMVOC                   |
| Mobile Combustion (excluding CO <sub>2</sub> )            | CH <sub>4</sub> , N <sub>2</sub> O, CO, NO <sub>x</sub> , NMVOC                   |
| Coal Mining   | CH <sub>4</sub>   |
| Abandoned Underground Coal Mines                          | CH <sub>4</sub>   |
| Petroleum Systems   | CH <sub>4</sub>   |
| Natural Gas Systems                                       | CH <sub>4</sub>   |
| Incineration of Waste                                     | CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O, NO <sub>x</sub> , CO, NMVOC |
| <b>Industrial Processes and Product Use</b>               |   |
| Titanium Dioxide Production                               | CO <sub>2</sub>   |
| Aluminum Production                                       | CO <sub>2</sub> , CF <sub>4</sub> , C <sub>2</sub> F <sub>6</sub>                 |
| Iron and Steel Production & Metallurgical Coke Production | CO <sub>2</sub> , CH <sub>4</sub>   |
| Ferroalloy Production                                     | CO <sub>2</sub> , CH <sub>4</sub>   |
| Ammonia Production  | CO <sub>2</sub>   |
| Urea Consumption for Non-Agricultural Purposes            | CO <sub>2</sub>   |
| Cement Production   | CO <sub>2</sub>   |
| Lime Production   | CO <sub>2</sub>   |
| Other Process Uses of Carbonates                          | CO <sub>2</sub>   |
| Soda Ash Production and Consumption                       | CO <sub>2</sub>   |
| Glass Production  | CO <sub>2</sub>   |
| Carbon Dioxide Consumption                                | CO <sub>2</sub>   |
| Phosphoric Acid Production                                | CO <sub>2</sub>   |
| Petrochemical Production                                  | CH <sub>4</sub> , CO <sub>2</sub>   |
| Silicon Carbide Production and Consumption                | CH <sub>4</sub> , CO <sub>2</sub>   |
| Lead Production   | CO <sub>2</sub>   |
| Zinc Production   | CO <sub>2</sub>   |
| Adipic Acid Production                                    | N <sub>2</sub> O  |
| Nitric Acid Production                                    | N <sub>2</sub> O  |
| N <sub>2</sub> O from Product Uses                        | N <sub>2</sub> O  |
| Substitution of Ozone Depleting Substances                | HFCs, PFCs <sup>a</sup>   |
| HFC-22 Production   | HFC-23  |
| Semiconductor Manufacture                                 | N <sub>2</sub> O, HFCs, PFCs <sup>b</sup> , SF <sub>6</sub> , NF <sub>3</sub>     |
| Electrical Transmission and Distributing                  | SF <sub>6</sub>   |
| Magnesium Production and Processing                       | CO <sub>2</sub> , HFCs, SF <sub>6</sub>   |
| <b>Agriculture</b>  |   |
| Enteric Fermentation                                      | CH <sub>4</sub>   |
| Manure Management   | CH <sub>4</sub> , N <sub>2</sub> O  |
| Rice Cultivation  | CH <sub>4</sub>   |
| Field Burning of Agricultural Residues                    | CH <sub>4</sub> , N <sub>2</sub> O, NO <sub>x</sub> , CO                          |
| Agricultural Soil Management                              | N <sub>2</sub> O  |
| <b>Land Use, Land-Use Change, and Forestry</b>            |   |
| Net CO <sub>2</sub> Flux                                  | CO <sub>2</sub> (sink)  |
| Cropland Remaining Cropland                               | CO <sub>2</sub>   |
| Land Converted to Cropland                                | CO <sub>2</sub>   |
| Grassland Remaining Grassland                             | CO <sub>2</sub>   |
| Land Converted to Grassland                               | CO <sub>2</sub>   |
| Settlements Remaining Settlements                         | CO <sub>2</sub> , N <sub>2</sub> O  |
| Forest Land Remaining Forest Land                         | CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O, NO <sub>x</sub> , CO        |
| Land Converted to Forest Land                             | CO <sub>2</sub>   |
| Wetlands Remaining Wetlands                               | CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O                              |
| Other (Landfilled Yard Trimmings and Food Scraps)         | CO <sub>2</sub>   |
| <b>Waste</b>  |   |
| Landfills   | CH <sub>4</sub> , NO <sub>x</sub> , CO, NMVOC                                     |
| Wastewater Treatment                                      | CH <sub>4</sub> , N <sub>2</sub> O, NO <sub>x</sub> , CO, NMVOC                   |
| Composting  | CH <sub>4</sub> , N <sub>2</sub> O  |

<sup>a</sup> Includes HFC-23, HFC-32, HFC-125, HFC-134a, HFC-143a, HFC-236fa, CF<sub>4</sub>, HFC-152a, HFC-227ea, HFC-245fa, HFC-4310mee, and PFC/PPFes.

<sup>b</sup> Includes such gases as HFC-23, CF<sub>4</sub>, C<sub>2</sub>F<sub>6</sub>.

## 6.5. Constants, Units, and Conversions

### Metric Prefixes

Although most activity data for the United States is gathered in customary U.S. units, these units are converted into metric units per international reporting guidelines. Table A-298 provides a guide for determining the magnitude of metric units.

**Table A-298: Guide to Metric Unit Prefixes**

| Prefix/Symbol   | Factor     |
|-----------------|------------|
| atto (a)        | $10^{-18}$ |
| femto (f)       | $10^{-15}$ |
| pico (p)        | $10^{-12}$ |
| nano (n)        | $10^{-9}$  |
| micro ( $\mu$ ) | $10^{-6}$  |
| milli (m)       | $10^{-3}$  |
| centi (c)       | $10^{-2}$  |
| deci (d)        | $10^{-1}$  |
| deca (da)       | 10         |
| hecto (h)       | $10^2$     |
| kilo (k)        | $10^3$     |
| mega (M)        | $10^6$     |
| giga (G)        | $10^9$     |
| tera (T)        | $10^{12}$  |
| peta (P)        | $10^{15}$  |
| exa (E)         | $10^{18}$  |

### Unit Conversions

|              |   |                 |   |                    |
|--------------|---|-----------------|---|--------------------|
| 1 kilogram   | = | 2.205 pounds    |   |                    |
| 1 pound      | = | 0.454 kilograms |   |                    |
| 1 short ton  | = | 2,000 pounds    | = | 0.9072 metric tons |
| 1 metric ton | = | 1,000 kilograms | = | 1.1023 short tons  |

|                |   |                      |
|----------------|---|----------------------|
| 1 cubic meter  | = | 35.315 cubic feet    |
| 1 cubic foot   | = | 0.02832 cubic meters |
| 1 U.S. gallon  | = | 3.785412 liters      |
| 1 barrel (bbl) | = | 0.159 cubic meters   |
| 1 barrel (bbl) | = | 42 U.S. gallons      |
| 1 liter        | = | 0.001 cubic meters   |

|             |   |                  |
|-------------|---|------------------|
| 1 foot      | = | 0.3048 meters    |
| 1 meter     | = | 3.28 feet        |
| 1 mile      | = | 1.609 kilometers |
| 1 kilometer | = | 0.622 miles      |

|               |   |                            |   |                 |   |                     |
|---------------|---|----------------------------|---|-----------------|---|---------------------|
| 1 acre        | = | 43,560 square feet         | = | 0.4047 hectares | = | 4,047 square meters |
| 1 square mile | = | 2.589988 square kilometers |   |                 |   |                     |

To convert degrees Fahrenheit to degrees Celsius, subtract 32 and multiply by 5/9

To convert degrees Celsius to Kelvin, add 273.15 to the number of Celsius degrees

## Density Conversions<sup>131</sup>

|                          |               |   |                                |
|--------------------------|---------------|---|--------------------------------|
| Methane                  | 1 cubic meter | = | 0.67606 kilograms              |
| Carbon dioxide           | 1 cubic meter | = | 1.85387 kilograms              |
| Natural gas liquids      | 1 metric ton  | = | 11.6 barrels = 1,844.2 liters  |
| Unfinished oils          | 1 metric ton  | = | 7.46 barrels = 1,186.04 liters |
| Alcohol                  | 1 metric ton  | = | 7.94 barrels = 1,262.36 liters |
| Liquefied petroleum gas  | 1 metric ton  | = | 11.6 barrels = 1,844.2 liters  |
| Aviation gasoline        | 1 metric ton  | = | 8.9 barrels = 1,415.0 liters   |
| Naphtha jet fuel         | 1 metric ton  | = | 8.27 barrels = 1,314.82 liters |
| Kerosene jet fuel        | 1 metric ton  | = | 7.93 barrels = 1,260.72 liters |
| Motor gasoline           | 1 metric ton  | = | 8.53 barrels = 1,356.16 liters |
| Kerosene                 | 1 metric ton  | = | 7.73 barrels = 1,228.97 liters |
| Naphtha                  | 1 metric ton  | = | 8.22 barrels = 1,306.87 liters |
| Distillate               | 1 metric ton  | = | 7.46 barrels = 1,186.04 liters |
| Residual oil             | 1 metric ton  | = | 6.66 barrels = 1,058.85 liters |
| Lubricants               | 1 metric ton  | = | 7.06 barrels = 1,122.45 liters |
| Bitumen                  | 1 metric ton  | = | 6.06 barrels = 963.46 liters   |
| Waxes                    | 1 metric ton  | = | 7.87 barrels = 1,251.23 liters |
| Petroleum coke           | 1 metric ton  | = | 5.51 barrels = 876.02 liters   |
| Petrochemical feedstocks | 1 metric ton  | = | 7.46 barrels = 1,186.04 liters |
| Special naphtha          | 1 metric ton  | = | 8.53 barrels = 1,356.16 liters |
| Miscellaneous products   | 1 metric ton  | = | 8.00 barrels = 1,271.90 liters |

## Energy Conversions

### Converting Various Energy Units to Joules

The common energy unit used in international reports of greenhouse gas emissions is the joule. A joule is the energy required to push with a force of one Newton for one meter. A terajoule (TJ) is one trillion ( $10^{12}$ ) joules. A British thermal unit (Btu, the customary U.S. energy unit) is the quantity of heat required to raise the temperature of one pound of water one degree Fahrenheit at or near 39.2 degrees Fahrenheit.

|        |   |
|--------|---|
| 1 TJ = | 2.388×10 <sup>11</sup> calories           |
|        | 23.88 metric tons of crude oil equivalent |
|        | 947.8 million Btus                        |
|        | 277,800 kilowatt-hours                    |

### Converting Various Physical Units to Energy Units

Data on the production and consumption of fuels are first gathered in physical units. These units must be converted to their energy equivalents. The conversion factors in Table A-299 can be used as default factors, if local data are not available. See Appendix A of EIA's *Monthly Energy Review February 2016* (EIA 2016) for more detailed information on the energy content of various fuels.

---

<sup>131</sup> Reference: EIA (2007)

**Table A-299: Conversion Factors to Energy Units (Heat Equivalents)**

| <b>Fuel Type (Units)</b>                   | <b>Factor</b> |
|--|---------------|
| <b>Solid Fuels (Million Btu/Short ton)</b> |               |
| Anthracite coal                            | 22.573        |
| Bituminous coal                            | 23.89         |
| Sub-bituminous coal                        | 17.14         |
| Lignite                                    | 12.866        |
| Coke                                       | 23.367        |
| <b>Natural Gas (Btu/Cubic foot)</b>        | <b>1,032</b>  |
| <b>Liquid Fuels (Million Btu/Barrel)</b>   |               |
| Motor gasoline                             | 5.060         |
| Aviation gasoline                          | 5.048         |
| Kerosene                                   | 5.670         |
| Jet fuel, kerosene-type                    | 5.670         |
| Distillate fuel                            | 5.825         |
| Residual oil                               | 6.287         |
| Naphtha for petrochemicals                 | 5.248         |
| Petroleum coke                             | 6.024         |
| Other oil for petrochemicals               | 5.825         |
| Special naphthas                           | 5.248         |
| Lubricants                                 | 6.065         |
| Waxes                                      | 5.537         |
| Asphalt                                    | 6.636         |
| Still gas                                  | 6.000         |
| Misc. products                             | 5.796         |

Note: For petroleum and natural gas, *Monthly Energy Review February 2016* (EIA 2016). For coal ranks, *State Energy Data Report 1992* (EIA 1993). All values are given in higher heating values (gross calorific values).

## 6.6. Abbreviations

|                 |  |
|-----------------|--|
| AAPFCO          | American Association of Plant Food Control Officials                   |
| ABS             | Acrylonitrile butadiene styrene  |
| AC              | Air conditioner  |
| ACC             | American Chemistry Council   |
| AEDT            | FAA Aviation Environmental Design Tool                                 |
| AEO             | Annual Energy Outlook  |
| AFEAS           | Alternative Fluorocarbon Environmental Acceptability Study             |
| AFV             | Alternative fuel vehicle   |
| AGA             | American Gas Association   |
| AHEF            | Atmospheric and Health Effect Framework                                |
| AISI            | American Iron and Steel Institute                                      |
| ANGA            | American Natural Gas Alliance  |
| ANL             | Argonne National Laboratory  |
| APC             | American Plastics Council  |
| API             | American Petroleum Institute   |
| APTA            | American Public Transportation Association                             |
| AR4             | IPCC Fourth Assessment Report  |
| AR5             | IPCC Fifth Assessment Report   |
| ARI             | Advanced Resources International                                       |
| ASAE            | American Society of Agricultural Engineers                             |
| ASTM            | American Society for Testing and Materials                             |
| BCEF            | Biomass conversion and expansion factors                               |
| BEA             | Bureau of Economic Analysis, U.S. Department of Commerce               |
| BLM             | Bureau of Land Management  |
| BoC             | Bureau of Census   |
| BOD             | Biological oxygen demand   |
| BOD5            | Biochemical oxygen demand over a 5-day period                          |
| BOEM            | Bureau of Ocean Energy Management                                      |
| BOEMRE          | Bureau of Ocean Energy Management, Regulation and Enforcement          |
| BOF             | Basic oxygen furnace   |
| BRS             | Biennial Reporting System  |
| BTS             | Bureau of Transportation Statistics, U.S. Department of Transportation |
| Btu             | British thermal unit   |
| C               | Carbon   |
| C&EN            | Chemical and Engineering News  |
| CAAA            | Clean Air Act Amendments of 1990                                       |
| CAPP            | Canadian Association of Petroleum Producers                            |
| CARB            | California Air Resources Board   |
| CBI             | Confidential business information                                      |
| C-CAP           | Coastal Change Analysis Program  |
| CDAP            | Chemical Data Access Tool  |
| CEFM            | Cattle Enteric Fermentation Model                                      |
| CEMS            | Continuous emission monitoring system                                  |
| CFC             | Chlorofluorocarbon   |
| CFR             | Code of Federal Regulations  |
| CGA             | Compressed Gas Association   |
| CH <sub>4</sub> | Methane  |
| CHP             | Combined heat and power  |
| CIGRE           | International Council on Large Electric Systems                        |
| CKD             | Cement kiln dust   |
| CLE             | Crown Light Exposure   |
| CMA             | Chemical Manufacturer's Association                                    |
| CMM             | Coal mine methane  |
| CMOP            | Coalbed Methane Outreach Program                                       |
| CMR             | Chemical Market Reporter   |
| CNG             | Compressed natural gas   |
| CO              | Carbon monoxide  |
| CO <sub>2</sub> | Carbon dioxide   |
| COD             | Chemical oxygen demand   |
| COGCC           | Colorado Oil and Gas Conservation Commission                           |

|         |  |
|---------|--|
| CRF     | Common Reporting Format                                      |
| CRM     | Component ratio method                                       |
| CRP     | Conservation Reserve Program                                 |
| CTIC    | Conservation Technology Information Center                   |
| CVD     | Chemical vapor deposition                                    |
| CWNS    | Clean Watershed Needs Survey                                 |
| d.b.h   | Diameter breast height                                       |
| DE      | Digestible energy  |
| DESC    | Defense Energy Support Center-DoD's defense logistics agency |
| DFAMS   | Defense Fuels Automated Management System                    |
| DHS     | Department of Homeland Security                              |
| DM      | Dry matter   |
| DOC     | Degradable organic carbon                                    |
| DOC     | U.S. Department of Commerce                                  |
| DoD     | U.S. Department of Defense                                   |
| DOE     | U.S. Department of Energy                                    |
| DOI     | U.S. Department of the Interior                              |
| DOT     | U.S. Department of Transportation                            |
| DRI     | Direct Reduced Iron  |
| EAF     | Electric arc furnace   |
| EDB     | Aircraft Engine Emissions Databank                           |
| EDF     | Environmental Defense Fund                                   |
| EER     | Energy economy ratio   |
| EF      | Emission factor  |
| EFMA    | European Fertilizer Manufacturers Association                |
| EJ      | Exajoule   |
| EGR     | Exhaust gas recirculation                                    |
| EGU     | Electric generating unit                                     |
| EIA     | Energy Information Administration, U.S. Department of Energy |
| EIIP    | Emissions Inventory Improvement Program                      |
| EOR     | Enhanced oil recovery  |
| EPA     | U.S. Environmental Protection Agency                         |
| ERS     | Economic Research Service                                    |
| ETMS    | Enhanced Traffic Management System                           |
| EV      | Electric vehicle   |
| EVI     | Enhanced Vegetation Index                                    |
| FAA     | Federal Aviation Administration                              |
| FAO     | Food and Agricultural Organization                           |
| FAOSTAT | Food and Agricultural Organization database                  |
| FCCC    | Framework Convention on Climate Change                       |
| FEB     | Fiber Economics Bureau                                       |
| FERC    | Federal Energy Regulatory Commission                         |
| FGD     | Flue gas desulfurization                                     |
| FHWA    | Federal Highway Administration                               |
| FIA     | Forest Inventory and Analysis                                |
| FIADB   | Forest Inventory and Analysis Database                       |
| FIPR    | Florida Institute of Phosphate Research                      |
| FQSV    | First-quarter of silicon volume                              |
| FSA     | Farm Service Agency  |
| FTP     | Federal Test Procedure                                       |
| g       | Gram   |
| GCV     | Gross calorific value  |
| GDP     | Gross domestic product                                       |
| GHG     | Greenhouse gas   |
| GHGRP   | Greenhouse Gas Reporting Program                             |
| GJ      | Gigajoule  |
| GOADS   | Gulf Offshore Activity Data System                           |

|        |   |
|--------|---|
| GPG    | Good Practice Guidance  |
| GRI    | Gas Research Institute  |
| GSAM   | Gas Systems Analysis Model  |
| GTI    | Gas Technology Institute  |
| GWP    | Global warming potential  |
| ha     | Hectare   |
| HBFC   | Hydrobromofluorocarbon  |
| HC     | Hydrocarbon   |
| HCFC   | Hydrochlorofluorocarbon   |
| HDDV   | Heavy duty diesel vehicle   |
| HDGV   | Heavy duty gas vehicle  |
| HDPE   | High density polyethylene   |
| HFC    | Hydrofluorocarbon   |
| HFE    | Hydrofluoroethers   |
| HHV    | Higher Heating Value  |
| HMA    | Hot Mix Asphalt   |
| HMIWI  | Hospital/medical/infectious waste incinerator                       |
| HTF    | Heat Transfer Fluid   |
| HTS    | Harmonized Tariff Schedule  |
| HWP    | Harvested wood product  |
| IBF    | International bunker fuels  |
| IC     | Integrated Circuit  |
| ICAO   | International Civil Aviation Organization                           |
| ICE    | Internal combustion engine  |
| IDB    | Integrated Database   |
| IEA    | International Energy Agency   |
| IFO    | Intermediate Fuel Oil   |
| IISRP  | International Institute of Synthetic Rubber Products                |
| ILENR  | Illinois Department of Energy and Natural Resources                 |
| IMO    | International Maritime Organization                                 |
| IPAA   | Independent Petroleum Association of America                        |
| IPCC   | Intergovernmental Panel on Climate Change                           |
| IPPU   | Industrial Processes and Product Use                                |
| ITC    | U.S. International Trade Commission                                 |
| ITRS   | International Technology Roadmap for Semiconductors                 |
| JWR    | Jim Walters Resources   |
| KCA    | Key category analysis   |
| kg     | Kilogram  |
| kt     | Kiloton   |
| kWh    | Kilowatt hour   |
| LDDT   | Light duty diesel truck   |
| LDDV   | Light duty diesel vehicle   |
| LDGT   | Light duty gas truck  |
| LDGV   | Light duty gas vehicle  |
| LDPE   | Low density polyethylene  |
| LDT    | Light-duty truck  |
| LDV    | Light-duty vehicle  |
| LEV    | Low emission vehicles   |
| LFG    | Landfill gas  |
| LFGTE  | Landfill gas-to-energy  |
| LHV    | Lower Heating Value   |
| LKD    | Lime kiln dust  |
| LLDPE  | Linear low density polyethylene                                     |
| LMOP   | EPA's Landfill Methane Outreach Program                             |
| LNG    | Liquefied natural gas   |
| LPG    | Liquefied petroleum gas(es)   |
| LTO    | Landing and take-off  |
| LULUCF | Land use, land-use change, and forestry                             |
| MARPOL | International Convention for the Prevention of Pollution from Ships |
| MC     | Motorcycle  |
| MCF    | Methane conversion factor   |
| MCL    | Maximum Contaminant Levels  |

|                         |  |
|-------------------------|--|
| MCFD                    | Thousand cubic feet per day                              |
| MDI                     | Metered dose inhalers                                    |
| MECS                    | EIA Manufacturer's Energy Consumption Survey             |
| MEM                     | Micro-electromechanical systems                          |
| MER                     | Monthly Energy Review                                    |
| MGO                     | Marine gas oil   |
| MJ                      | Megajoule  |
| MLRA                    | Major Land Resource Area                                 |
| mm                      | Millimeter   |
| MMBtu                   | Million British thermal units                            |
| MMCF                    | Million cubic feet                                       |
| MMCFD                   | Million cubic feet per day                               |
| MMS                     | Minerals Management Service                              |
| MMT                     | Million Metric Tons                                      |
| MMTCE                   | Million metric tons carbon equivalent                    |
| MMT CO <sub>2</sub> Eq. | Million metric tons carbon dioxide equivalent            |
| MODIS                   | Moderate Resolution Imaging Spectroradiometer            |
| MoU                     | Memorandum of Understanding                              |
| MOVES                   | U.S. EPA's Motor Vehicle Emission Simulator model        |
| MPG                     | Miles per gallon   |
| MRLC                    | Multi-Resolution Land Characteristics Consortium         |
| MRV                     | Monitoring, reporting, and verification                  |
| MSHA                    | Mine Safety and Health Administration                    |
| MSW                     | Municipal solid waste                                    |
| MT                      | Metric ton   |
| MTBE                    | Methyl Tertiary Butyl Ether                              |
| MTBS                    | Monitoring Trends in Burn Severity                       |
| MVAC                    | Motor vehicle air conditioning                           |
| MY                      | Model year   |
| N <sub>2</sub> O        | Nitrous oxide  |
| NA                      | Not available  |
| NACWA                   | National Association of Clean Water Agencies             |
| NAHMS                   | National Animal Health Monitoring System                 |
| NAICS                   | North American Industry Classification System            |
| NAPAP                   | National Acid Precipitation and Assessment Program       |
| NARR                    | North American Regional Reanalysis Product               |
| NASA                    | National Aeronautics and Space Administration            |
| NASF                    | National Association of State Foresters                  |
| NASS                    | USDA's National Agriculture Statistics Service           |
| NC                      | No change  |
| NCASI                   | National Council of Air and Stream Improvement           |
| NCV                     | Net calorific value                                      |
| NE                      | Not estimated  |
| NEI                     | National Emissions Inventory                             |
| NEMA                    | National Electrical Manufacturers Association            |
| NEMS                    | National Energy Modeling System                          |
| NESHAP                  | National Emission Standards for Hazardous Air Pollutants |
| NEU                     | Non-Energy Use   |
| NEV                     | Neighborhood Electric Vehicle                            |
| NF <sub>3</sub>         | Nitrogen trifluoride                                     |
| NGHGI                   | National Greenhouse Gas Inventory                        |
| NGL                     | Natural gas liquids                                      |
| NIR                     | National Inventory Report                                |
| NLA                     | National Lime Association                                |
| NLCD                    | National Land Cover Dataset                              |
| NMOC                    | Non-methane organic compounds                            |
| NM VOC                  | Non-methane volatile organic compound                    |

|                 |  |
|-----------------|--|
| NO              | Nitric oxide   |
| NO              | Not occurring  |
| NO <sub>2</sub> | Nitrogen Dioxide                                       |
| NO <sub>x</sub> | Nitrogen oxides  |
| NOAA            | National Oceanic and Atmospheric Administration        |
| NPRA            | National Petroleum and Refiners Association            |
| NRC             | National Research Council                              |
| NRCS            | Natural Resources Conservation Service                 |
| NRI             | National Resources Inventory                           |
| NSCEP           | National Service Center for Environmental Publications |
| NSCR            | Non-selective catalytic reduction                      |
| NSPS            | New source performance standards                       |
| NWS             | National Weather Service                               |
| OAG             | Official Airline Guide                                 |
| OAP             | EPA Office of Atmospheric Programs                     |
| OAQPS           | EPA Office of Air Quality Planning and Standards       |
| ODP             | Ozone depleting potential                              |
| ODS             | Ozone depleting substances                             |
| OECD            | Organization of Economic Co-operation and Development  |
| OEM             | Original equipment manufacturers                       |
| OGJ             | Oil & Gas Journal                                      |
| OH              | Hydroxyl radical                                       |
| OMS             | EPA Office of Mobile Sources                           |
| ORNL            | Oak Ridge National Laboratory                          |
| OSHA            | Occupational Safety and Health Administration          |
| OTA             | Office of Technology Assessment                        |
| OTAQ            | EPA Office of Transportation and Air Quality           |
| PAH             | Polycyclic aromatic hydrocarbons                       |
| PCC             | Precipitate calcium carbonate                          |
| PDF             | Probability Density Function                           |
| PECVD           | Plasma enhanced chemical vapor deposition              |
| PET             | Polyethylene terephthalate                             |
| PET             | Potential evapotranspiration                           |
| PEVM            | PFC Emissions Vintage Model                            |
| PFC             | Perfluorocarbon  |
| PFPE            | Perfluoropolyether                                     |
| PHMSA           | Pipeline and Hazardous Materials Safety Administration |
| PI              | Productivity index                                     |
| POTW            | Publicly Owned Treatment Works                         |
| ppbv            | Parts per billion (10 <sup>9</sup> ) by volume         |
| ppm             | Parts per million                                      |
| ppmv            | Parts per million (10 <sup>6</sup> ) by volume         |
| pptv            | Parts per trillion (10 <sup>12</sup> ) by volume       |
| PRP             | Pasture/Range/Paddock                                  |
| PS              | Polystyrene  |
| PSU             | Primary Sample Unit                                    |
| PU              | Polyurethane   |
| PVC             | Polyvinyl chloride                                     |
| PV              | Photovoltaic   |
| QA/QC           | Quality Assurance and Quality Control                  |
| QBtu            | Quadrillion Btu  |
| R&D             | Research and Development                               |
| RECs            | Reduced Emissions Completions                          |
| RCRA            | Resource Conservation and Recovery Act                 |
| RMA             | Rubber Manufacturers' Association                      |
| RPA             | Resources Planning Act                                 |
| RTO             | Regression-through-the-origin                          |
| SAE             | Society of Automotive Engineers                        |
| SAGE            | System for assessing Aviation's Global Emissions       |
| SAN             | Styrene Acrylonitrile                                  |
| SAR             | IPCC Second Assessment Report                          |
| SCR             | Selective catalytic reduction                          |

|                 |   |
|-----------------|---|
| SCSE            | South central and southeastern coastal                              |
| SEC             | Securities and Exchange Commission                                  |
| SEMI            | Semiconductor Equipment and Materials Industry                      |
| SF <sub>6</sub> | Sulfur hexafluoride   |
| SICAS           | Semiconductor International Capacity Statistics                     |
| SNAP            | Significant New Alternative Policy Program                          |
| SNG             | Synthetic natural gas   |
| SO <sub>2</sub> | Sulfur dioxide  |
| SOC             | Soil Organic Carbon   |
| SOG             | State of Garbage survey   |
| SOHIO           | Standard Oil Company of Ohio  |
| SSURGO          | Soil Survey Geographic Database                                     |
| STMC            | Scrap Tire Management Council                                       |
| SULEV           | Super Ultra Low Emissions Vehicle                                   |
| SWANA           | Solid Waste Association of North America                            |
| SWDS            | Solid waste disposal sites  |
| TA              | Treated anaerobically (wastewater)                                  |
| TAM             | Typical animal mass   |
| TAME            | Tertiary amyl methyl ether  |
| TAR             | IPCC Third Assessment Report  |
| TBtu            | Trillion Btu  |
| TDN             | Total digestible nutrients  |
| TEDB            | Transportation Energy Data Book                                     |
| TFI             | The Fertilizer Institute  |
| TIGER           | Topologically Integrated Geographic Encoding and Referencing survey |
| TJ              | Terajoule   |
| TLEV            | Traditional low emissions vehicle                                   |
| TMLA            | Total Manufactured Layer Area                                       |
| TRI             | Toxic Release Inventory   |
| TSDF            | Hazardous waste treatment, storage, and disposal facility           |
| TVA             | Tennessee Valley Authority  |
| UAN             | Urea ammonium nitrate   |
| UDI             | Utility Data Institute  |
| UFORE           | U.S. Forest Service's Urban Forest Effects model                    |
| UG              | Underground (coal mining)   |
| U.S.            | United States   |
| U.S. ITC        | United States International Trade Commission                        |
| UEP             | United Egg Producers  |
| ULEV            | Ultra low emission vehicle  |
| UNEP            | United Nations Environmental Programme                              |
| UNFCCC          | United Nations Framework Convention on Climate Change               |
| USAA            | U.S. Aluminum Association   |
| USAF            | United States Air Force   |
| USDA            | United States Department of Agriculture                             |
| USFS            | United States Forest Service  |
| USGS            | United States Geological Survey                                     |
| VAIP            | EPA's Voluntary Aluminum Industrial Partnership                     |
| VAM             | Ventilation air methane   |
| VKT             | Vehicle kilometers traveled   |
| VMT             | Vehicle miles traveled  |
| VOCs            | Volatile organic compounds  |
| VS              | Volatile solids   |
| WERF            | Water Environment Research Federation                               |
| WFF             | World Fab Forecast (previously WFW, World Fab Watch)                |
| WGC             | World Gas Conference  |
| WIP             | Waste in place  |
| WMO             | World Meteorological Organization                                   |

|      |                            |
|------|----------------------------|
| WMS  | Waste management systems   |
| WTE  | Waste-to-energy            |
| WW   | Wastewater                 |
| WWTP | Wastewater treatment plant |
| ZEVs | Zero emissions vehicles    |

## 6.7. Chemical Formulas

**Table A-300: Guide to Chemical Formulas**

| Symbol  | Name                                 |
|---|--------------------------------------|
| Al  | Aluminum                             |
| Al <sub>2</sub> O <sub>3</sub>                    | Aluminum Oxide                       |
| Br  | Bromine                              |
| C   | Carbon                               |
| CH <sub>4</sub>                                   | Methane                              |
| C <sub>2</sub> H <sub>6</sub>                     | Ethane                               |
| C <sub>3</sub> H <sub>8</sub>                     | Propane                              |
| CF <sub>4</sub>                                   | Perfluoromethane                     |
| C <sub>2</sub> F <sub>6</sub>                     | Perfluoroethane, hexafluoroethane    |
| c-C <sub>3</sub> F <sub>6</sub>                   | Perfluorocyclopropane                |
| C <sub>3</sub> F <sub>8</sub>                     | Perfluoropropane                     |
| c-C <sub>4</sub> F <sub>8</sub>                   | Perfluorocyclobutane                 |
| C <sub>4</sub> F <sub>10</sub>                    | Perfluorobutane                      |
| C <sub>5</sub> F <sub>12</sub>                    | Perfluoropentane                     |
| C <sub>6</sub> F <sub>14</sub>                    | Perfluorohexane                      |
| CF <sub>3</sub> I                                 | Trifluoroiodomethane                 |
| CFCl <sub>3</sub>                                 | Trichlorofluoromethane (CFC-11)      |
| CF <sub>2</sub> Cl <sub>2</sub>                   | Dichlorodifluoromethane (CFC-12)     |
| CF <sub>3</sub> Cl                                | Chlorotrifluoromethane (CFC-13)      |
| C <sub>2</sub> F <sub>3</sub> Cl <sub>3</sub>     | Trichlorotrifluoroethane (CFC-113)*  |
| CCl <sub>3</sub> CF <sub>3</sub>                  | CFC-113a*                            |
| C <sub>2</sub> F <sub>4</sub> Cl <sub>2</sub>     | Dichlorotetrafluoroethane (CFC-114)  |
| C <sub>2</sub> F <sub>5</sub> Cl                  | Chloropentafluoroethane (CFC-115)    |
| CHCl <sub>2</sub> F                               | HCFC-21                              |
| CHF <sub>2</sub> Cl                               | Chlorodifluoromethane (HCFC-22)      |
| C <sub>2</sub> F <sub>3</sub> HCl <sub>2</sub>    | HCFC-123                             |
| C <sub>2</sub> F <sub>4</sub> HCl                 | HCFC-124                             |
| C <sub>2</sub> FH <sub>3</sub> Cl <sub>2</sub>    | HCFC-141b                            |
| C <sub>2</sub> H <sub>3</sub> F <sub>2</sub> Cl   | HCFC-142b                            |
| CF <sub>3</sub> CF <sub>2</sub> CHCl <sub>2</sub> | HCFC-225ca                           |
| CClF <sub>2</sub> CF <sub>2</sub> CHClF           | HCFC-225cb                           |
| CCl <sub>4</sub>                                  | Carbon tetrachloride                 |
| CHClCCl <sub>2</sub>                              | Trichloroethylene                    |
| CCl <sub>2</sub> CCl <sub>2</sub>                 | Perchloroethylene, tetrachloroethene |
| CH <sub>3</sub> Cl                                | Methylchloride                       |
| CH <sub>3</sub> CCl <sub>3</sub>                  | Methylchloroform                     |
| CH <sub>2</sub> Cl <sub>2</sub>                   | Methylenechloride                    |
| CHCl <sub>3</sub>                                 | Chloroform, trichloromethane         |
| CHF <sub>3</sub>                                  | HFC-23                               |
| CH <sub>2</sub> F <sub>2</sub>                    | HFC-32                               |
| CH <sub>3</sub> F                                 | HFC-41                               |
| C <sub>2</sub> HF <sub>5</sub>                    | HFC-125                              |
| C <sub>2</sub> H <sub>2</sub> F <sub>4</sub>      | HFC-134                              |
| CH <sub>2</sub> FCF <sub>3</sub>                  | HFC-134a                             |
| C <sub>2</sub> H <sub>3</sub> F <sub>3</sub>      | HFC-143*                             |
| C <sub>2</sub> H <sub>3</sub> F <sub>3</sub>      | HFC-143a*                            |
| CH <sub>2</sub> FCH <sub>2</sub> F                | HFC-152*                             |
| C <sub>2</sub> H <sub>4</sub> F <sub>2</sub>      | HFC-152a*                            |
| CH <sub>3</sub> CH <sub>2</sub> F                 | HFC-161                              |
| C <sub>3</sub> HF <sub>7</sub>                    | HFC-227ea                            |
| CF <sub>3</sub> CF <sub>2</sub> CH <sub>2</sub> F | HFC-236cb                            |
| CF <sub>3</sub> CHFCHF <sub>2</sub>               | HFC-236ea                            |

|  |                                     |
|--|-------------------------------------|
| C <sub>3</sub> H <sub>2</sub> F <sub>6</sub>                                       | HFC-236fa                           |
| C <sub>3</sub> H <sub>3</sub> F <sub>5</sub>                                       | HFC-245ca                           |
| CHF <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>                                   | HFC-245fa                           |
| CF <sub>3</sub> CH <sub>2</sub> CF <sub>2</sub> CH <sub>3</sub>                    | HFC-365mfc                          |
| C <sub>6</sub> H <sub>2</sub> F <sub>10</sub>                                      | HFC-43-10mee                        |
| CF <sub>3</sub> OCHF <sub>2</sub>  | HFE-125                             |
| CF <sub>2</sub> HOCHF <sub>2</sub> H   | HFE-134                             |
| CH <sub>3</sub> OCF <sub>3</sub>   | HFE-143a                            |
| CF <sub>3</sub> CHFOCF <sub>3</sub>  | HFE-227ea                           |
| CF <sub>3</sub> CHClOCHF <sub>2</sub>  | HCFE-235da2                         |
| CF <sub>3</sub> CHFOCHF <sub>2</sub>   | HFE-236ea2                          |
| CF <sub>3</sub> CH <sub>2</sub> OCF <sub>3</sub>                                   | HFE-236fa                           |
| CF <sub>3</sub> CF <sub>2</sub> OCH <sub>3</sub>                                   | HFE-245cb2                          |
| CHF <sub>2</sub> CH <sub>2</sub> OCF <sub>3</sub>                                  | HFE-245fa1                          |
| CF <sub>3</sub> CH <sub>2</sub> OCHF <sub>2</sub>                                  | HFE-245fa2                          |
| CHF <sub>2</sub> CF <sub>2</sub> OCH <sub>3</sub>                                  | HFE-254cb2                          |
| CF <sub>3</sub> CH <sub>2</sub> OCH <sub>3</sub>                                   | HFE-263fb2                          |
| CF <sub>3</sub> CF <sub>2</sub> OCF <sub>2</sub> CHF <sub>2</sub>                  | HFE-329mcc2                         |
| CF <sub>3</sub> CF <sub>2</sub> OCH <sub>2</sub> CF <sub>3</sub>                   | HFE-338mcf2                         |
| CF <sub>3</sub> CF <sub>2</sub> CF <sub>2</sub> OCH <sub>3</sub>                   | HFE-347mcc3                         |
| CF <sub>3</sub> CF <sub>2</sub> OCH <sub>2</sub> CHF <sub>2</sub>                  | HFE-347mcf2                         |
| CF <sub>3</sub> CHFCF <sub>2</sub> OCH <sub>3</sub>                                | HFE-356mec3                         |
| CHF <sub>2</sub> CF <sub>2</sub> CF <sub>2</sub> OCH <sub>3</sub>                  | HFE-356pcc3                         |
| CHF <sub>2</sub> CF <sub>2</sub> OCH <sub>2</sub> CHF <sub>2</sub>                 | HFE-356pcf2                         |
| CHF <sub>2</sub> CF <sub>2</sub> CH <sub>2</sub> OCHF <sub>2</sub>                 | HFE-356pcf3                         |
| CF <sub>3</sub> CF <sub>2</sub> CH <sub>2</sub> OCH <sub>3</sub>                   | HFE-365mcf3                         |
| CHF <sub>2</sub> CF <sub>2</sub> OCH <sub>2</sub> CH <sub>3</sub>                  | HFE-374pcf2                         |
| C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub>                                     | HFE-7100                            |
| C <sub>4</sub> F <sub>9</sub> OC <sub>2</sub> H <sub>5</sub>                       | HFE-7200                            |
| CHF <sub>2</sub> OCF <sub>2</sub> OC <sub>2</sub> F <sub>4</sub> OCHF <sub>2</sub> | H-Galden 1040x                      |
| CHF <sub>2</sub> OCF <sub>2</sub> OCHF <sub>2</sub>                                | HG-10                               |
| CHF <sub>2</sub> OCF <sub>2</sub> CF <sub>2</sub> OCHF <sub>2</sub>                | HG-01                               |
| CH <sub>3</sub> OCH <sub>3</sub>   | Dimethyl ether                      |
| CH <sub>2</sub> Br <sub>2</sub>  | Dibromomethane                      |
| CH <sub>2</sub> BrCl   | Dibromochloromethane                |
| CHBr <sub>3</sub>  | Tribromomethane                     |
| CHBrF <sub>2</sub>   | Bromodifluoromethane                |
| CH <sub>3</sub> Br   | Methylbromide                       |
| CF <sub>2</sub> BrCl   | Bromodichloromethane (Halon 1211)   |
| CF <sub>3</sub> Br(CBrF <sub>3</sub> )   | Bromotrifluoromethane (Halon 1301)  |
| CF <sub>3</sub> I  | FIC-1311                            |
| CO   | Carbon monoxide                     |
| CO <sub>2</sub>  | Carbon dioxide                      |
| CaCO <sub>3</sub>  | Calcium carbonate, Limestone        |
| CaMg(CO <sub>3</sub> ) <sub>2</sub>  | Dolomite                            |
| CaO  | Calcium oxide, Lime                 |
| Cl   | atomic Chlorine                     |
| F  | Fluorine                            |
| Fe   | Iron                                |
| Fe <sub>2</sub> O <sub>3</sub>   | Ferric oxide                        |
| FeSi   | Ferrosilicon                        |
| H, H <sub>2</sub>  | atomic Hydrogen, molecular Hydrogen |
| H <sub>2</sub> O   | Water                               |
| H <sub>2</sub> O <sub>2</sub>  | Hydrogen peroxide                   |
| OH   | Hydroxyl                            |
| N, N <sub>2</sub>  | atomic Nitrogen, molecular Nitrogen |
| NH <sub>3</sub>  | Ammonia                             |
| NH <sub>4</sub> <sup>+</sup>   | Ammonium ion                        |
| HNO <sub>3</sub>   | Nitric acid                         |
| MgO  | Magnesium oxide                     |
| NF <sub>3</sub>  | Nitrogen trifluoride                |
| N <sub>2</sub> O   | Nitrous oxide                       |

|                                  |                                      |
|----------------------------------|--------------------------------------|
| NO                               | Nitric oxide                         |
| NO <sub>2</sub>                  | Nitrogen dioxide                     |
| NO <sub>3</sub>                  | Nitrate radical                      |
| Na                               | Sodium                               |
| Na <sub>2</sub> CO <sub>3</sub>  | Sodium carbonate, soda ash           |
| Na <sub>3</sub> AlF <sub>6</sub> | Synthetic cryolite                   |
| O, O <sub>2</sub>                | atomic Oxygen, molecular Oxygen      |
| O <sub>3</sub>                   | Ozone                                |
| S                                | atomic Sulfur                        |
| H <sub>2</sub> SO <sub>4</sub>   | Sulfuric acid                        |
| SF <sub>6</sub>                  | Sulfur hexafluoride                  |
| SF <sub>5</sub> CF <sub>3</sub>  | Trifluoromethylsulphur pentafluoride |
| SO <sub>2</sub>                  | Sulfur dioxide                       |
| Si                               | Silicon                              |
| SiC                              | Silicon carbide                      |
| SiO <sub>2</sub>                 | Quartz                               |

\* Distinct isomers.

## References

- EIA (2016) *Monthly Energy Review, February 2016*. Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE/EIA-0035(2016/02). February 2016.
- EIA (2007) *Emissions of Greenhouse Gases in the United States 2006, Draft Report*. Office of Integrated Analysis and Forecasting, Energy Information Administration, U.S. Department of Energy, Washington, DC. DOE-EIA-0573(2006).
- EIA (1998) *Emissions of Greenhouse Gases in the United States*, DOE/EIA-0573(97), Energy Information Administration, U.S. Department of Energy. Washington, DC. October.
- EIA (1993) *State Energy Data Report 1992*, DOE/EIA-0214(93), Energy Information Administration, U.S. Department of Energy. Washington, DC. December.
- EPA (2015) "1970 - 2014 Average annual emissions, all criteria pollutants in MS Excel." National Emissions Inventory (NEI) Air Pollutant Emissions Trends Data. Office of Air Quality Planning and Standards, March 2015. Available online at <<http://www.epa.gov/ttn/chieftrends/index.html>>.
- EPA (2003) E-mail correspondence. Air pollutant data. Office of Air Pollution to the Office of Air Quality Planning and Standards, U.S. Environmental Protection Agency (EPA). December 22, 2003.
- IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- IPCC (2007) *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.). Cambridge University Press. Cambridge, United Kingdom 996 pp.
- IPCC (2001) *Climate Change 2001: The Scientific Basis. Intergovernmental Panel on Climate Change*, J.T. Houghton, Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, C.A. Johnson, and K. Maskell (eds.). Cambridge University Press. Cambridge, United Kingdom.
- IPCC (1996) *Climate Change 1995: The Science of Climate Change. Intergovernmental Panel on Climate Change*, J.T. Houghton, L.G. Meira Filho, B.A. Callander, N. Harris, A. Kattenberg, and K. Maskell. (eds.). Cambridge University Press. Cambridge, United Kingdom.

# ANNEX 7 Uncertainty

The annual U.S. Inventory presents the best effort to produce estimates for greenhouse gas source and sink categories in the United States. These estimates were generated according to the UNFCCC reporting guidelines, following the recommendations set forth in the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC 2006). This Annex provides an overview of the uncertainty analysis conducted to support the U.S. Inventory, describes the sources of uncertainty characterized throughout the Inventory associated with various source categories (including emissions and sinks), and describes the methods through which uncertainty information was collected, quantified, and presented.

## 7.1. Overview

The current inventory emission estimates for some source categories, such as for CO<sub>2</sub> Emissions from Fossil Fuel Combustion, have relatively low level of uncertainty associated with them. However, for some other source categories, the inventory emission estimates are considered less certain. The two major types of uncertainty associated with these emission estimates are (1) model uncertainty, which arises when the emission and/or removal estimation models used in developing the inventory estimates do not fully and accurately characterize the respective emission and/or removal processes (due to a lack of technical details or other resources), resulting in the use of incorrect or incomplete estimation methodologies and (2) parameter uncertainty, which arises due to a lack of precise input data such as emission factors and activity data.

The model uncertainty can be partially analyzed by comparing the model results with those of other models developed to characterize the same emission (or removal) process, after taking into account the differences in their conceptual framework, capabilities, data, and assumptions. However, it would be very difficult—if not impossible—to quantify the model uncertainty associated with the emission estimates (primarily because, in most cases, only a single model has been developed to estimate emissions from any one source). Therefore, model uncertainty was not quantified in this report. Nonetheless, it has been discussed qualitatively, where appropriate, along with the individual source category description and inventory estimation methodology.

Parameter uncertainty is, therefore, the principal type and source of uncertainty associated with the national inventory emission estimates and is the main focus of the quantitative uncertainty analyses in this report. Parameter uncertainty has been quantified for all of the emission sources and sinks in the U.S. Inventory, with the exception of one very small emission source category, CH<sub>4</sub> emissions from Incineration of Waste, which was included in the *1990 through 2008 National GHG Inventory* for the first time, and three other source categories (International Bunker Fuels Energy Sources of Indirect Greenhouse Gas Emissions, and Wood Biomass and Ethanol Consumption) whose emissions are not included in the Inventory totals.

The primary purpose of the uncertainty analysis conducted in support of the U.S. Inventory is (i) to determine the quantitative uncertainty associated with the emission (and removal) estimates presented in the main body of this report [based on the uncertainty associated with the input parameters used in the emission (and removal) estimation methodologies] and (ii) to evaluate the relative importance of the input parameters in contributing to uncertainty in the associated source category inventory estimate and in the overall inventory estimate. Thus, the U.S. Inventory uncertainty analysis provides a strong foundation for developing future improvements and revisions to the Inventory estimation process. For each source category, the analysis highlights opportunities for changes to data measurement, data collection, and calculation methodologies. These are presented in the “Planned Improvements” sections of each source category’s discussion in the main body of the report.

## 7.2. Methodology and Results

The United States has developed a quality assurance and quality control (QA/QC) and uncertainty management plan (EPA 2002). Like the QA/QC plan, the uncertainty management plan is part of a continually evolving process. The uncertainty management plan provides for a quantitative assessment of the inventory analysis itself, thereby contributing to continuing efforts to understand both what causes uncertainty and how to improve inventory quality. Although the plan provides both general and specific guidelines for implementing quantitative uncertainty analysis, its components are intended to evolve over time, consistent with the inventory estimation process. The U.S. plan includes procedures and guidelines, and forms and templates, for developing quantitative assessments of uncertainty in the national Inventory estimates (EPA 2002). For the 1990 through 2014 Inventory, EPA has used the uncertainty management plan as well as the methodology presented in the *2006 IPCC Guidelines*.

The *2006 IPCC Guidelines* recommends two methods—Approach 1 and Approach 2—for developing quantitative estimates of uncertainty in the inventory estimate of individual source categories and the overall inventory. Of these, the

Approach 2 method is both more flexible and reliable than Approach 1; both approaches are described in the next section. The United States is in the process of implementing a multi-year strategy to develop quantitative estimates of uncertainty for all source categories using the Approach 2. In following the UNFCCC requirement under Article 4.1, emissions from International Bunker Fuels, Energy Sources of Indirect Greenhouse Gas Emissions, and Wood Biomass and Ethanol Consumption are not included in the total emissions estimated for the U.S. Inventory; therefore, no quantitative uncertainty estimates have been developed for these source categories.<sup>132</sup> Emissions from Biomass and Ethanol Consumption are accounted for implicitly in the Land Use, Land-Use Change and Forestry (LULUCF) chapter through the calculation of changes in carbon stocks. The Energy sector does provide an estimate of CO<sub>2</sub> emissions from Biomass and Ethanol Consumption provided as a memo item for informational purposes consistent with the UNFCCC reporting requirements.

### **Approach 1 and Approach 2 Methods**

The Approach 1 method for estimating uncertainty is based on the error propagation equation. This equation combines the uncertainty associated with the activity data and the uncertainty associated with the emission (or the other) factors. The Approach 1 method is applicable where emissions (or removals) are usually estimated as the product of an activity value and an emission factor or as the sum of individual sub-source category values. Inherent in employing the Approach 1 method are the assumptions that, for each source category, (i) both the activity data and the emission factor values are approximately normally distributed, (ii) the coefficient of variation (i.e., the ratio of the standard deviation to the mean) associated with each input variable is less than 30 percent, and (iii) the input variables within and across (sub-) source categories are not correlated (i.e., value of each variable is independent of the values of other variables).

The Approach 2 method is preferred (i) if the uncertainty associated with the input variables is significantly large, (ii) if the distributions underlying the input variables are not normal, (iii) if the estimates of uncertainty associated with the input variables are correlated, and/or (iv) if a sophisticated estimation methodology and/or several input variables are used to characterize the emission (or removal) process correctly. In practice, the Approach 2 is the preferred method of uncertainty analysis for all source categories where sufficient and reliable data are available to characterize the uncertainty of the input variables.

The Approach 2 method employs the Monte Carlo Stochastic Simulation technique (also referred to as the Monte Carlo method). Under this method, estimates of emissions (or removals) for a particular source category are generated many times (equal to the number of simulations specified) using an uncertainty model, which is an emission (or removal) estimation equation that imitates or is the same as the inventory estimation model for a particular source category. These estimates are generated using the respective, randomly-selected values for the constituent input variables using commercially available simulation software such as @RISK.

### **Characterization of Uncertainty in Input Variables**

Both Approach 1 and Approach 2 uncertainty analyses require that all the input variables are well-characterized in terms of their Probability Density Functions (PDFs). In the absence of particularly convincing data measurements, sufficient data samples, or expert judgments that determined otherwise, the PDFs incorporated in the current source category uncertainty analyses were limited to normal, lognormal, uniform, triangular, and beta distributions. The choice among these five PDFs depended largely on the observed or measured data and expert judgment.

### **Source Category Inventory Uncertainty Estimates**

Discussion surrounding the input parameters and sources of uncertainty for each source category appears in the body of this report. Table A-301 summarizes results based on assessments of source category-level uncertainty. The table presents base year (1990 or 1995) and current year (2014) emissions for each source category. The combined uncertainty (at the 95 percent confidence interval) for each source category is expressed as the percentage deviation above and below the total 2014 emissions estimated for that source category. Source category trend uncertainty is described subsequently in this Appendix.

---

<sup>132</sup> However, because the input variables that determine the emissions from the Fossil Fuel Combustion and the International Bunker Fuels source categories are correlated, uncertainty associated with the activity variables in the International Bunker Fuels was taken into account in estimating the uncertainty associated with the Fossil Fuel Combustion.

**Table A-301: Summary Results of Source Category Uncertainty Analyses**

| Source Category   | Base Year Emissions <sup>h,a</sup> | 2014 Emissions <sup>a</sup> | 2014 Uncertainty <sup>b</sup> |            |
|---|------------------------------------|-----------------------------|-------------------------------|------------|
|   | MMT CO <sub>2</sub> Eq.            | MMT CO <sub>2</sub> Eq.     | Low                           | High       |
| <b>CO<sub>2</sub></b>                                     | <b>5,114.7</b>                     | <b>5,555.6</b>              | <b>-2%</b>                    | <b>5%</b>  |
| Fossil Fuel Combustion <sup>c</sup>                       | 4,740.3                            | 5,207.8                     | -2%                           | 5%         |
| Non-Energy Use of Fuels                                   | 118.1                              | 114.3                       | -25%                          | 42%        |
| Iron and Steel Production & Metallurgical Coke Production | 99.7                               | 55.4                        | -15%                          | 15%        |
| Natural Gas Systems                                       | 37.7                               | 42.4                        | -19%                          | 30%        |
| Cement Production   | 33.3                               | 38.8                        | -6%                           | 6%         |
| Petrochemical Production                                  | 21.6                               | 26.5                        | -5%                           | 5%         |
| Lime Production   | 11.7                               | 14.1                        | -3%                           | 3%         |
| Other Process Uses of Carbonates                          | 4.9                                | 12.1                        | -12%                          | 15%        |
| Ammonia Production  | 13.0                               | 9.4                         | -8%                           | 8%         |
| Incineration of Waste                                     | 8.0                                | 9.4                         | -10%                          | 14%        |
| Carbon Dioxide Consumption                                | 1.5                                | 4.5                         | -12%                          | 13%        |
| Urea Consumption for Non-Agricultural Purposes            | 3.8                                | 4.0                         | -12%                          | 12%        |
| Petroleum Systems   | 3.6                                | 3.6                         | -24%                          | 149%       |
| Aluminum Production                                       | 6.8                                | 2.8                         | -2%                           | 2%         |
| Soda Ash Production and Consumption                       | 2.8                                | 2.8                         | -7%                           | 6%         |
| Ferroalloy Production                                     | 2.2                                | 1.9                         | -12%                          | 12%        |
| Titanium Dioxide Production                               | 1.2                                | 1.8                         | -12%                          | 13%        |
| Glass Production  | 1.5                                | 1.3                         | -4%                           | 5%         |
| Phosphoric Acid Production                                | 1.5                                | 1.1                         | -19%                          | 20%        |
| Zinc Production   | 0.6                                | 1.0                         | -19%                          | 21%        |
| Lead Production   | 0.5                                | 0.5                         | -15%                          | 16%        |
| Silicon Carbide Production and Consumption                | 0.4                                | 0.2                         | -9%                           | 9%         |
| Magnesium Production and Processing                       | +                                  | +                           | -5%                           | 5%         |
| <i>Wood Biomass and Ethanol Consumption<sup>d</sup></i>   | 219.4                              | 293.7                       | NE                            | NE         |
| <i>International Bunker Fuels<sup>e</sup></i>             | 103.5                              | 103.2                       | NE                            | NE         |
| <b>CH<sub>4</sub></b>                                     | <b>773.9</b>                       | <b>730.8</b>                | <b>-8%</b>                    | <b>26%</b> |
| Natural Gas Systems                                       | 206.8                              | 176.1                       | -19%                          | 30%        |
| Enteric Fermentation                                      | 164.2                              | 164.3                       | 11%                           | 18%        |
| Landfills   | 179.6                              | 148.0                       | -30%                          | 61%        |
| Petroleum Systems   | 38.7                               | 68.1                        | -24%                          | 149%       |
| Coal Mining   | 96.5                               | 67.6                        | -12%                          | 15%        |
| Manure Management   | 37.2                               | 61.2                        | -18%                          | 20%        |
| Wastewater Treatment                                      | 15.7                               | 14.7                        | -39%                          | 2%         |
| Rice Cultivation  | 13.1                               | 11.9                        | -17%                          | 17%        |
| Stationary Combustion                                     | 8.5                                | 8.1                         | -41%                          | 155%       |
| Abandoned Underground Coal Mines                          | 7.2                                | 6.3                         | -18%                          | 24%        |
| Composting  | 0.4                                | 2.1                         | -50%                          | 50%        |
| Mobile Combustion   | 5.6                                | 2.0                         | -12%                          | 18%        |
| Field Burning of Agricultural Residues                    | 0.2                                | 0.3                         | -40%                          | 40%        |
| Petrochemical Production                                  | 0.2                                | 0.1                         | -55%                          | 45%        |
| Ferroalloy Production                                     | +                                  | +                           | -12%                          | 12%        |
| Silicon Carbide Production and Consumption                | +                                  | +                           | -9%                           | 10%        |
| Iron and Steel Production & Metallurgical Coke Production | +                                  | +                           | -19%                          | 19%        |
| Incineration of Waste                                     | +                                  | +                           | NE                            | NE         |
| <i>International Bunker Fuels<sup>e</sup></i>             | 0.2                                | 0.1                         | NE                            | NE         |
| <b>N<sub>2</sub>O</b>                                     | <b>406.2</b>                       | <b>403.5</b>                | <b>-20%</b>                   | <b>11%</b> |
| Agricultural Soil Management                              | 303.3                              | 318.4                       | -18%                          | 47%        |
| Stationary Combustion                                     | 11.9                               | 23.4                        | -24%                          | 46%        |
| Manure Management   | 14.0                               | 17.5                        | -16%                          | 24%        |
| Mobile Combustion   | 41.2                               | 16.3                        | -4%                           | 27%        |
| Nitric Acid Production                                    | 12.1                               | 10.9                        | -5%                           | 5%         |
| Adipic Acid Production                                    | 15.2                               | 5.4                         | -4%                           | 4%         |
| Wastewater Treatment                                      | 3.4                                | 4.8                         | -76%                          | 108%       |
| N <sub>2</sub> O from Product Uses                        | 4.2                                | 4.2                         | -24%                          | 24%        |

|   |                |                |             |            |
|---|----------------|----------------|-------------|------------|
| Composting  | 0.3            | 1.8            | -50%        | 50%        |
| Incineration of Waste                                   | 0.5            | 0.3            | -53%        | 163%       |
| Semiconductor Manufacture                               | +              | 0.2            | -12%        | 12%        |
| Field Burning of Agricultural Residues                  | 0.1            | 0.1            | -29%        | 29%        |
| <i>International Bunker Fuels<sup>e</sup></i>           | <i>1.1</i>     | <i>1.0</i>     | <i>NE</i>   | <i>NE</i>  |
| <b>HFCs, PFCs, SF<sub>6</sub>, and NF<sub>3</sub></b>   | <b>132.6</b>   | <b>175.3</b>   | <b>-4%</b>  | <b>6%</b>  |
| Substitution of Ozone Depleting Substances <sup>f</sup> | 31.0           | 156.4          | -1%         | 10%        |
| Electrical Transmission and Distribution                | 25.4           | 5.6            | -17%        | 23%        |
| HFC-22 Production                                       | 46.1           | 5.0            | -7%         | 10%        |
| Semiconductor Manufacture                               | 3.6            | 4.5            | -6%         | 6%         |
| Aluminum Production                                     | 21.5           | 2.5            | -6%         | 6%         |
| Magnesium Production                                    | 5.2            | 1.2            | -12%        | 15%        |
| <b>Total Emissions<sup>g</sup></b>                      | <b>6,427.4</b> | <b>6,865.2</b> | <b>-2%</b>  | <b>5%</b>  |
| <b>LULUCF Emissions<sup>i</sup></b>                     | <b>15.0</b>    | <b>24.6</b>    | <b>-48%</b> | <b>58%</b> |
| <b>LULUCF Total Net Flux<sup>j</sup></b>                | <b>(753.0)</b> | <b>(787.0)</b> | <b>-18%</b> | <b>34%</b> |
| <b>LULUCF Sector Total<sup>k</sup></b>                  | <b>(738.0)</b> | <b>(762.5)</b> | <b>-18%</b> | <b>35%</b> |
| <b>Net Emissions (Sources and Sinks)<sup>g</sup></b>    | <b>5,689.5</b> | <b>6,102.7</b> | <b>-4%</b>  | <b>6%</b>  |

Notes: Total emissions (excluding emissions for which uncertainty was not quantified) is presented without LULUCF. Net emissions is presented with LULUCF.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

NE - Not Estimated

<sup>a</sup> Emission estimates reported in this table correspond to emissions from only those source categories for which quantitative uncertainty was performed for the current inventory. Thus the totals reported for 2013 in this table exclude approximately 5.7 MMT CO<sub>2</sub> Eq. of emissions for which quantitative uncertainty was not assessed. Hence, these emission estimates do not match the final total U.S. greenhouse gas emission estimates presented in this inventory. All uncertainty estimates correspond only to the totals reported in this table.

<sup>b</sup> The uncertainty estimates correspond to a 95 percent confidence interval, with the lower bound corresponding to 2.5<sup>th</sup> percentile and the upper bound corresponding to 97.5<sup>th</sup> percentile.

<sup>c</sup> This source category's inventory estimates exclude CO<sub>2</sub> emissions from geothermal sources, as quantitative uncertainty analysis was not performed for that sub-source category. Hence, for this source category, the emissions reported in this table do not match the emission estimates presented in the Energy chapter of the inventory.

<sup>d</sup> Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals.

<sup>e</sup> Emissions from International Bunker Fuels are not included in the totals.

<sup>f</sup> This source category's estimate for 2013 excludes 5.3 MMT of CO<sub>2</sub> Eq. from several very small emission sources, as uncertainty associated with those sources was not assessed. Hence, for this source category, the emissions reported in this table do not match the emission estimates presented in the Industrial Processes and Product Use chapter of the inventory.

<sup>g</sup> Totals exclude emissions for which uncertainty was not quantified.

<sup>h</sup> Base Year is 1990 for all sources except Substitution of Ozone Depleting Substances, for which the United States has chosen 1995.

<sup>i</sup> LULUCF emissions include the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions reported for Non-CO<sub>2</sub> Emissions from Forest Fires, N<sub>2</sub>O Fluxes from Forest Soils, CO<sub>2</sub> Emissions from Agricultural Liming, CO<sub>2</sub> Emissions from Urea Fertilization, Peatlands Remaining Peatlands, and N<sub>2</sub>O Fluxes from Settlement Soils.

<sup>j</sup> Net CO<sub>2</sub> flux is the net C stock change from the following categories: *Forest Land Remaining Forest Land, Land Converted to Forest Land, Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, Land Converted to Grassland, Settlements Remaining Settlements, and Other.*

<sup>k</sup> The LULUCF Sector Total is the net sum of all emissions (i.e., sources) of greenhouse gases to the atmosphere plus removals of CO<sub>2</sub> (i.e., sinks or negative emissions) from the atmosphere.

Notes: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

## Overall (Aggregate) Inventory Level Uncertainty Estimates

The overall level uncertainty estimate for the U.S. Greenhouse Gas Emissions Inventory was developed using the IPCC Approach 2 uncertainty estimation methodology. The uncertainty models of all the emission source categories could not be directly integrated to develop the overall uncertainty estimates due to software constraints in integrating multiple, large uncertainty models. Therefore, an alternative approach was adopted to develop the overall uncertainty estimates. The Monte Carlo simulation output data for each emission source category uncertainty analysis were combined by type of gas and the probability distributions were fitted to the combined simulation output data, where such simulated output data were available. If such detailed output data were not available for particular emissions sources, individual probability distributions were assigned to those source category emission estimates based on the most detailed data available from the quantitative uncertainty analysis performed.

For Composting and parts of Agricultural Soil Management source categories, Approach 1 uncertainty results were used in the overall uncertainty analysis estimation. However, for all other emission sources (excluding international bunker fuels, CO<sub>2</sub> from biomass combustion, and CH<sub>4</sub> from incineration of waste), Approach 2 uncertainty results were used in the overall uncertainty estimation.

The overall uncertainty model results indicate that the 2014 U.S. greenhouse gas emissions are estimated to be within the range of approximately 6,765.4 to 7,223.9 MMT CO<sub>2</sub> Eq., reflecting a relative 95 percent confidence interval uncertainty range of -2 percent to 5 percent with respect to the total U.S. greenhouse gas emission estimate of approximately 6,865.2 MMT CO<sub>2</sub> Eq. The uncertainty interval associated with total CO<sub>2</sub> emissions, which constitute about 81 percent of the total U.S. greenhouse gas emissions in 2014, ranges from -2 percent to 5 percent of total CO<sub>2</sub> emissions estimated. The results indicate that the uncertainty associated with the inventory estimate of the total CH<sub>4</sub> emissions ranges from -8 percent to 26 percent, uncertainty associated with the total inventory N<sub>2</sub>O emission estimate ranges from -20 percent to 11 percent, and uncertainty associated with high GWP gas emissions ranges from -4 percent to 6 percent.

A summary of the overall quantitative uncertainty estimates is shown below.

**Table A-302: Quantitative Uncertainty Assessment of Overall National Inventory Emissions (MMT CO<sub>2</sub> Eq. and Percent)**

| Gas  | 2014 Emission                                      |                          | Uncertainty Range Relative to Emission Estimate <sup>b</sup> |  |             |  | Standard  |  |
|--|--|--------------------------|--|--|-------------|--|---|--|
|  | Estimate <sup>a</sup><br>(MMT CO <sub>2</sub> Eq.) | Range                    |  | Relative to Emission Estimate <sup>b</sup> |             | Mean <sup>c</sup><br>(MMT CO <sub>2</sub> Eq.) | Deviation <sup>c</sup><br>(MMT CO <sub>2</sub> Eq.) |  |
|  |  | Lower Bound <sup>d</sup> | Upper Bound <sup>d</sup>                                     | Lower Bound                                | Upper Bound |  |   |  |
| CO <sub>2</sub>  | 5,555.6  | 5,459.4                  | 5,830.0  | -2%  | 5%          | 5,643.8  | 94.9  |  |
| CH <sub>4</sub> <sup>e</sup>                                 | 730.8  | 674.3                    | 917.5  | -8%  | 26%         | 785.0  | 60.2  |  |
| N <sub>2</sub> O <sup>e</sup>                                | 403.5  | 322.5                    | 447.9  | -20%                                       | 11%         | 378.6  | 32.2  |  |
| PFC, HFC, SF <sub>6</sub> , and NF <sub>3</sub> <sup>e</sup> | 175.3  | 172.3                    | 190.9  | -4%  | 6%          | 181.6  | 4.7   |  |
| <b>Total Emissions</b>                                       | <b>6,865.2</b>                                     | <b>6,765.4</b>           | <b>7,223.9</b>   | <b>-2%</b>                                 | <b>5%</b>   | <b>6,989.0</b>                                 | <b>117.5</b>  |  |
| <b>LULUCF Emissions<sup>f</sup></b>                          | <b>24.6</b>  | <b>12.8</b>              | <b>38.9</b>  | <b>-48%</b>                                | <b>58%</b>  | <b>23.0</b>                                    | <b>6.8</b>  |  |
| <b>LULUCF Total Net Flux<sup>g</sup></b>                     | <b>(787.0)</b>                                     | <b>(1,051.4)</b>         | <b>(647.8)</b>   | <b>-18%</b>                                | <b>34%</b>  | <b>(847.2)</b>                                 | <b>102.9</b>  |  |
| <b>LULUCF Sector Total<sup>h</sup></b>                       | <b>(762.5)</b>                                     | <b>(1,029.8)</b>         | <b>(622.5)</b>   | <b>-18%</b>                                | <b>35%</b>  | <b>(824.2)</b>                                 | <b>103.0</b>  |  |
| <b>Net Emissions (Sources and Sinks)</b>                     | <b>6,102.7</b>                                     | <b>5,861.6</b>           | <b>6,477.6</b>   | <b>-4%</b>                                 | <b>6%</b>   | <b>6,164.8</b>                                 | <b>156.7</b>  |  |

Notes: Total emissions (excluding emissions for which uncertainty was not quantified) is presented without LULUCF. Net emissions is presented with LULUCF

<sup>a</sup> Emission estimates reported in this table correspond to emissions from only those source categories for which quantitative uncertainty was performed this year. Thus the totals reported in this table exclude approximately 5.3 MMT CO<sub>2</sub> Eq. of emissions for which quantitative uncertainty was not assessed. Hence, these emission estimates do not match the final total U.S. greenhouse gas emission estimates presented in this Inventory.

<sup>b</sup> The lower and upper bounds for emission estimates correspond to a 95 percent confidence interval, with the lower bound corresponding to 2.5<sup>th</sup> percentile and the upper bound corresponding to 97.5<sup>th</sup> percentile.

<sup>c</sup> Mean value indicates the arithmetic average of the simulated emission estimates; standard deviation indicates the extent of deviation of the simulated values from the mean.

<sup>d</sup> The lower and upper bound emission estimates for the sub-source categories do not sum to total emissions because the low and high estimates for total emissions were calculated separately through simulations.

<sup>e</sup> The overall uncertainty estimates did not take into account the uncertainty in the GWP values for CH<sub>4</sub>, N<sub>2</sub>O and high GWP gases used in the inventory emission calculations for 2014.

<sup>f</sup> LULUCF emissions include the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions reported for Non-CO<sub>2</sub> Emissions from Forest Fires, N<sub>2</sub>O Fluxes from Forest Soils, CO<sub>2</sub> Emissions from Agricultural Liming, CO<sub>2</sub> Emissions from Urea Fertilization, Peatlands Remaining Peatlands, and N<sub>2</sub>O Fluxes from Settlement Soils.

<sup>g</sup> Net CO<sub>2</sub> flux is the net C stock change from the following categories: *Forest Land Remaining Forest Land, Land Converted to Forest Land, Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, Land Converted to Grassland, Settlements Remaining Settlements, and Other.*

<sup>h</sup> The LULUCF Sector Total is the net sum of all emissions (i.e., sources) of greenhouse gases to the atmosphere plus removals of CO<sub>2</sub> (i.e., sinks or negative emissions) from the atmosphere.

Notes: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

### Trend Uncertainty

In addition to the estimates of uncertainty associated with the current year's emission estimates, this Annex also presents the estimates of trend uncertainty. The 2006 IPCC Guidelines defines trend as the difference in emissions between the base year (i.e., 1990) and the current year (i.e., 2014) inventory estimates. However, for purposes of understanding the concept of trend uncertainty, the emission trend is defined in this Inventory as the percentage change in the emissions (or removal) estimated for the current year, relative to the emission (or removal) estimated for the base year. The uncertainty associated with this emission trend is referred to as trend uncertainty.

Under the Approach 1 method, the trend uncertainty for a source category is estimated using the sensitivity of the calculated difference between the base year and the current year (i.e., 2014) emissions to an incremental (i.e., 1 percent)

increase in one or both of these values for that source category. The two sensitivities are expressed as percentages: Type A sensitivity highlights the effect on the difference between the base and the current year emissions caused by a 1 percent change in both, while Type B sensitivity highlights the effect caused by a change to only the current year's emissions. Both sensitivities are simplifications introduced in order to analyze the correlation between the base and the current year estimates. Once calculated, the two sensitivities are combined using the error propagation equation to estimate the overall trend uncertainty.

Under the Approach 2 method, the trend uncertainty is estimated using the Monte Carlo Stochastic Simulation technique. The trend uncertainty analysis takes into account the fact that the base and the current year estimates often share input variables. For purposes of the current Inventory, a simple approach has been adopted, under which the base year source category emissions (or removals) are assumed to exhibit the same uncertainty characteristics as the current year emissions (or removals). Source category-specific PDFs for base year estimates were developed using current year (i.e., 2014) uncertainty output data. These were adjusted to account for differences in magnitude between the two years' inventory estimates. Then, for each source category, a trend uncertainty estimate was developed using the Monte Carlo method. The overall inventory trend uncertainty estimate was developed by combining all source category-specific trend uncertainty estimates. These trend uncertainty estimates present the range of likely change from base year to 2014, and are shown in Table A-303.

**Table A-303: Quantitative Assessment of Trend Uncertainty (MMT CO<sub>2</sub> Eq. and Percent)**

| Gas/Source  | Base Year                 | 2014                   | Emissions          | Trend Range <sup>a,b</sup> |             |
|---|---------------------------|------------------------|--------------------|----------------------------|-------------|
|   | Emissions <sup>a</sup>    | Emissions <sup>a</sup> | Trend <sup>a</sup> | Trend Range <sup>a,b</sup> |             |
|   | (MMT CO <sub>2</sub> Eq.) |                        | (%)                | (%)                        |             |
|   |                           |                        |                    | Lower Bound                | Upper Bound |
| <b>CO<sub>2</sub></b>                                     | <b>5,114.7</b>            | <b>5,555.6</b>         | <b>9%</b>          | <b>4%</b>                  | <b>14%</b>  |
| Fossil Fuel Combustion <sup>c</sup>                       | 4,740.3                   | 5,207.8                | 10%                | 5%                         | 15%         |
| Non-Energy Use of Fuels                                   | 118.1                     | 114.3                  | -3%                | -41%                       | 59%         |
| Iron and Steel Production & Metallurgical Coke Production | 99.7                      | 55.4                   | -44%               | -55%                       | -32%        |
| Natural Gas Systems                                       | 37.7                      | 42.4                   | 12%                | -21%                       | 57%         |
| Cement Production   | 33.3                      | 38.8                   | 16%                | 7%                         | 27%         |
| Petrochemical Production                                  | 21.6                      | 26.5                   | 23%                | 15%                        | 31%         |
| Lime Production   | 11.7                      | 14.1                   | 21%                | 16%                        | 25%         |
| Other Process Uses of Carbonates                          | 4.9                       | 12.1                   | -146%              | 104%                       | 197%        |
| Ammonia Production  | 13.0                      | 9.4                    | -28%               | -36%                       | -19%        |
| Incineration of Waste                                     | 8.0                       | 9.4                    | 18%                | 0%                         | 40%         |
| Carbon Dioxide Consumption                                | 1.5                       | 4.5                    | 204%               | 154%                       | 264%        |
| Urea Consumption for Non-Agricultural Purposes            | 3.8                       | 4.0                    | 6%                 | -10%                       | 25%         |
| Petroleum Systems   | 3.6                       | 3.6                    | 0%                 | -55%                       | 134%        |
| Aluminum Production                                       | 6.8                       | 2.8                    | -59%               | -60%                       | -57%        |
| Soda Ash Production and Consumption                       | 2.8                       | 2.8                    | 0%                 | -9%                        | 10%         |
| Ferroalloy Production                                     | 2.2                       | 1.9                    | -11%               | -25%                       | 6%          |
| Titanium Dioxide Production                               | 1.2                       | 1.8                    | 47%                | 22%                        | 76%         |
| Glass Production  | 1.5                       | 1.3                    | -13%               | -18%                       | -7%         |
| Phosphoric Acid Production                                | 1.5                       | 1.1                    | -28%               | -46%                       | -5%         |
| Zinc Production   | 0.6                       | 1.0                    | 51%                | 15%                        | 101%        |
| Lead Production   | 0.5                       | 0.5                    | 0%                 | -19%                       | 25%         |
| Silicon Carbide Production and Consumption                | 0.4                       | 0.2                    | -54%               | -60%                       | -47%        |
| Magnesium Production and Processing                       | 0.0                       | +                      | NA                 | NA                         | NA          |
| Wood Biomass and Ethanol Consumption <sup>e</sup>         | 219.4                     | 293.7                  | 2%                 | NE                         | NE          |
| International Bunker Fuels <sup>f</sup>                   | 103.5                     | 103.2                  | 0%                 | NE                         | NE          |
| <b>CH<sub>4</sub></b>                                     | <b>773.9</b>              | <b>730.8</b>           | <b>-6%</b>         | <b>-23%</b>                | <b>17%</b>  |
| Natural Gas Systems                                       | 206.8                     | 176.1                  | -15%               | -40%                       | 20%         |
| Enteric Fermentation                                      | 164.2                     | 164.3                  | 0%                 | -18%                       | 23%         |
| Landfills   | 179.6                     | 148.0                  | -18%               | -66%                       | 105%        |
| Petroleum Systems   | 38.7                      | 68.1                   | 76%                | -23%                       | 311%        |
| Coal Mining   | 96.5                      | 67.6                   | -30%               | -46%                       | -17%        |
| Manure Management   | 37.2                      | 61.2                   | 65%                | 12%                        | 131%        |
| Wastewater Treatment                                      | 15.7                      | 14.7                   | -6%                | -69%                       | 4%          |

|   |                |                |             |             |             |
|---|----------------|----------------|-------------|-------------|-------------|
| Rice Cultivation  | 13.1           | 11.9           | -9%         | -66%        | 137%        |
| Stationary Combustion                                     | 8.5            | 8.1            | -5%         | -70%        | 200%        |
| Abandoned Underground Coal Mines                          | 7.2            | 6.3            | -12%        | -44%        | 26%         |
| Composting  | 0.4            | 2.1            | 439%        | 136%        | 1118%       |
| Mobile Combustion   | 5.6            | 2.0            | -64%        | -71%        | -55%        |
| Field Burning of Agricultural Residues                    | 0.2            | 0.3            | 18%         | -37%        | 123%        |
| Petrochemical Production                                  | 0.2            | 0.1            | -43%        | -76%        | 38%         |
| Ferroalloy Production                                     | +              | +              | -21%        | -33%        | -6%         |
| Silicon Carbide Production and Consumption                | +              | +              | -67%        | -71%        | -62%        |
| Iron and Steel Production & Metallurgical Coke Production | +              | +              | -44%        | -67%        | -42%        |
| Incineration of Waste                                     | +              | +              | NE          | NE          | NE          |
| <i>International Bunker Fuels<sup>f</sup></i>             | <i>0.2</i>     | <i>0.1</i>     | <i>-52%</i> | <i>NE</i>   | <i>NE</i>   |
| <b>N<sub>2</sub>O</b>                                     | <b>406.2</b>   | <b>403.5</b>   | <b>-1%</b>  | <b>-31%</b> | <b>44%</b>  |
| Agricultural Soil Management                              | 303.3          | 318.4          | 5%          | -35%        | 69%         |
| Stationary Combustion                                     | 11.9           | 23.4           | 96%         | 22%         | 213%        |
| Manure Management   | 14.0           | 17.5           | 25%         | -6%         | 64%         |
| Mobile Combustion   | 41.2           | 16.3           | -60%        | -68%        | -52%        |
| Nitric Acid Production                                    | 12.1           | 10.9           | -10%        | -16%        | -2%         |
| Adipic Acid Production                                    | 15.2           | 5.4            | -64%        | -66%        | -62%        |
| Wastewater Treatment                                      | 3.4            | 4.8            | 44%         | -70%        | 511%        |
| N <sub>2</sub> O from Product Uses                        | 4.2            | 4.2            | 0%          | -32%        | 46%         |
| Composting  | 0.3            | 1.8            | 439%        | 135%        | 1111%       |
| Incineration of Waste                                     | 0.5            | 0.3            | -32%        | -81%        | 143%        |
| Field Burning of Agricultural Residues                    | 0.1            | 0.1            | 23%         | -21%        | 94%         |
| <i>International Bunker Fuels<sup>f</sup></i>             | <i>1.1</i>     | <i>1.0</i>     | <i>5%</i>   | <i>NE</i>   | <i>NE</i>   |
| <b>HFCs, PFCs, SF<sub>6</sub>, and NF<sub>3</sub></b>     | <b>132.6</b>   | <b>175.3</b>   | <b>32%</b>  | <b>26%</b>  | <b>45%</b>  |
| Substitution of Ozone Depleting Substances <sup>g</sup>   | 31.0           | 156.4          | 405%        | 366%        | 447%        |
| Electrical Transmission and Distribution                  | 25.4           | 5.6            | -78%        | -83%        | -71%        |
| HCFC-22 Production  | 46.1           | 5.0            | -89%        | -90%        | -88%        |
| Semiconductor Manufacture                                 | 3.6            | 4.5            | 27%         | 17%         | 38%         |
| Aluminum Production                                       | 21.5           | 2.5            | -88%        | -89%        | -87%        |
| Magnesium Production                                      | 5.2            | 1.2            | -80%        | -83%        | -77%        |
| <b>Total Emissions<sup>h</sup></b>                        | <b>6,427.4</b> | <b>6,865.2</b> | <b>7%</b>   | <b>2%</b>   | <b>12%</b>  |
| <b>LULUCF Emissions<sup>i</sup></b>                       | <b>15.0</b>    | <b>24.6</b>    | <b>63%</b>  | <b>-18%</b> | <b>229%</b> |
| <b>LULUCF Total Net Flux<sup>j</sup></b>                  | <b>(753.0)</b> | <b>(787.0)</b> | <b>5%</b>   | <b>-25%</b> | <b>46%</b>  |
| <b>LULUCF Sector Total<sup>k</sup></b>                    | <b>(738.0)</b> | <b>(762.5)</b> | <b>3%</b>   | <b>-27%</b> | <b>45%</b>  |
| <b>Net Emissions (Sources and Sinks)</b>                  | <b>5,689.5</b> | <b>6,102.7</b> | <b>7%</b>   | <b>0%</b>   | <b>16%</b>  |

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

Notes: Total emissions (excluding emissions for which uncertainty was not quantified) is presented without LULUCF. Net emissions is presented with LULUCF  
NE - Not Estimated

<sup>a</sup> Emission estimates reported in this table correspond to emissions from only those source categories for which quantitative uncertainty was performed for the current inventory. Thus the totals reported for 2014 in this table exclude approximately 5.3 MMT CO<sub>2</sub> Eq. of emissions for which quantitative uncertainty was not assessed. Hence, these emission estimates do not match the final total U.S. greenhouse gas emission estimates presented in this Inventory. All uncertainty estimates correspond only to the totals reported in this table.

<sup>b</sup> The trend range represents a 95 percent confidence interval for the emission trend, with the lower bound corresponding to 2.5th percentile value and the upper bound corresponding to 97.5th percentile value.

<sup>c</sup> This source category's inventory estimates exclude CO<sub>2</sub> emissions from geothermal sources, as quantitative uncertainty analysis was not performed for that sub-source category. Hence, for this source category, the emissions reported in this table do not match the emission estimates presented in the Energy chapter of the Inventory.

<sup>d</sup> Sinks are only included in Net Emissions.

<sup>e</sup> Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals.

<sup>f</sup> Emissions from International Bunker Fuels are not included in the totals.

<sup>g</sup> This source category's estimate for 2014 excludes 5.3 MMT of CO<sub>2</sub> Eq. from several very small emission sources, as uncertainty associated with those sources was not assessed. Hence, for this source category, the emissions reported in this table do not match the emission estimates presented in the Industrial Processes and Product Use chapter of the Inventory.

<sup>h</sup> Totals exclude emissions for which uncertainty was not quantified.

<sup>i</sup> Base Year is 1990 for all sources except Substitution of Ozone Depleting Substances, for which the United States has chosen 1995.

<sup>j</sup> LULUCF emissions include the CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions reported for Non-CO<sub>2</sub> Emissions from Forest Fires, N<sub>2</sub>O Fluxes from Forest Soils, CO<sub>2</sub> Emissions from Agricultural Liming, CO<sub>2</sub> Emissions from Urea Fertilization, Peatlands Remaining Peatlands, and N<sub>2</sub>O Fluxes from Settlement Soils.

<sup>k</sup> Net CO<sub>2</sub> flux is the net C stock change from the following categories: *Forest Land Remaining Forest Land, Land Converted to Forest Land, Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, Land Converted to Grassland, Settlements Remaining Settlements, and Other.*

<sup>k</sup> The LULUCF Sector Total is the net sum of all emissions (i.e., sources) of greenhouse gases to the atmosphere plus removals of CO<sub>2</sub> (i.e., sinks or negative emissions) from the atmosphere.

Notes: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

### 7.3. Planned Improvements

Identifying the sources of uncertainty in the emission and sink estimates of the Inventory and quantifying the magnitude of the associated uncertainty is the crucial first step towards improving those estimates. Quantitative assessment of the parameter uncertainty may also provide information about the relative importance of input parameters (such as activity data and emission factors), based on their relative contribution to the uncertainty within the source category estimates. Such information can be used to prioritize resources with a goal of reducing uncertainty over time within or among inventory source categories and their input parameters. In the current Inventory, potential sources of model uncertainty have been identified for some emission source categories, and uncertainty estimates based on their parameters' uncertainty have been developed for all the emission source categories, with the exception of CH<sub>4</sub> from Incineration of Waste, and the International Bunker Fuels, Energy Sources of Indirect Greenhouse Gas Emissions, and Wood Biomass and Ethanol Consumption source categories, which are not included in the energy sector totals. Emissions from Wood Biomass and Ethanol Consumption however are accounted for implicitly in the Land Use, Land-Use Change and Forestry (LULUCF) chapter through the calculation of changes in carbon stocks. The Energy sector does provide an estimate of CO<sub>2</sub> emissions from Wood Biomass and Ethanol Consumption provided as a memo item for informational purposes.

Specific areas that require further research include:

- *Incorporating excluded emission sources.* Quantitative estimates for some of the sources and sinks of greenhouse gas emissions, such as from some land-use activities, industrial processes, and parts of mobile sources, could not be developed at this time either because data are incomplete or because methodologies do not exist for estimating emissions from these source categories. See Annex 5 of this report for a discussion of the sources of greenhouse gas emissions and sinks excluded from this report. In the future, efforts will focus on estimating emissions from excluded emission sources and developing uncertainty estimates for all source categories for which emissions are estimated.
- *Improving the accuracy of emission factors.* Further research is needed in some cases to improve the accuracy of emission factors used to calculate emissions from a variety of sources. For example, the accuracy of current emission factors applied to CH<sub>4</sub> and N<sub>2</sub>O emissions from stationary and mobile combustion are highly uncertain.
- *Collecting detailed activity data.* Although methodologies exist for estimating emissions for some sources, problems arise in obtaining activity data at a level of detail in which aggregate emission factors can be applied. For example, the ability to estimate emissions of SF<sub>6</sub> from electrical transmission and distribution is limited due to a lack of activity data regarding national SF<sub>6</sub> consumption or average equipment leak rates.

In improving the quality of uncertainty estimates the following include areas that deserve further attention:

- *Refine Source Category and Overall Uncertainty Estimates.* For many individual source categories, further research is needed to more accurately characterize PDFs that surround emissions modeling input variables. This might involve using measured or published statistics or implementing rigorous elicitation protocol to elicit expert judgments, if published or measured data are not available.
- *Include GWP uncertainty in the estimation of overall level and trend uncertainty.* The current year's Inventory does not include the uncertainty associated with the GWP values in the estimation of the overall uncertainty for the Inventory. Including this source would contribute to a better characterization of overall uncertainty and help assess the level of attention that this source of uncertainty warrants in the future.
- *Improve characterization of trend uncertainty associated with base year Inventory estimates.* The characterization of base year uncertainty estimates could be improved, by developing explicit uncertainty models for the base year. This would then improve the analysis of trend uncertainty. However, not all of the simplifying assumptions described in the "Trend Uncertainty" section above may be eliminated through this process due to a lack of availability of more appropriate data.

## 7.4. Additional Information on Uncertainty Analyses by Source

The quantitative uncertainty estimates associated with each emission and sink source category are reported in each chapter of this Inventory following the discussions of inventory estimates and their estimation methodology. This section provides additional descriptions of the uncertainty analyses performed for some of the sources, including the models and methods used to calculate the emission estimates and the potential sources of uncertainty surrounding them. These sources are organized below in the same order as the sources in each chapter of the main section of this Inventory. To avoid repetition, the following uncertainty analysis discussions of individual source categories do not include descriptions of these source categories. Hence, to better understand the details provided below, refer to the respective chapters and sections in the main section of this Inventory, as needed. All uncertainty estimates are reported relative to the 2014 Inventory estimates for the 95 percent confidence interval, unless otherwise specified.

### Energy

The uncertainty analysis descriptions in this section correspond to source categories included in the Energy chapter of the Inventory.

#### CO<sub>2</sub> from Fossil Fuel Combustion

For estimates of CO<sub>2</sub> from fossil fuel combustion, there are uncertainties in the consumption data, carbon content of fuels and products, and carbon oxidation efficiencies.

Although statistics of total fossil fuel and other energy consumption are relatively accurate, the allocation of this consumption to individual end-use sectors (i.e., residential, commercial, industrial, and transportation) is less certain. For this uncertainty estimation, the inventory estimation model for CO<sub>2</sub> from fossil fuel combustion was integrated with the relevant variables from the inventory estimation model for International Bunker Fuels, to realistically characterize the interaction (or endogenous correlation) between the variables of these two models.

In developing the uncertainty estimation model, uniform distributions were assumed for all activity-related input variables and emission factors, based on the SAIC/EIA (2001) report.<sup>133</sup> Triangular distributions were assigned for the oxidization factors (or combustion efficiencies). The uncertainty ranges were assigned to the input variables based on the data reported in SAIC/EIA (2001) and on conversations with various agency personnel.<sup>134</sup>

The uncertainty ranges for the activity-related input variables were typically asymmetric around their inventory estimates; the uncertainty ranges for the emissions factors were symmetric. Bias (or systematic uncertainties) associated with these variables accounted for much of the uncertainties associated with these variables (SAIC/EIA 2001).<sup>135</sup> For purposes of this uncertainty analysis, each input variable was simulated 10,000 times through Monte Carlo sampling.

#### CH<sub>4</sub> and N<sub>2</sub>O from Stationary Combustion

The uncertainty estimation model for this source category was developed by integrating the CH<sub>4</sub> and N<sub>2</sub>O stationary source inventory estimation models with the model for CO<sub>2</sub> from fossil fuel combustion to realistically characterize the interaction (or endogenous correlation) between the variables of these three models. About 55 input variables were simulated for the uncertainty analysis of this source category (about 20 from the CO<sub>2</sub> emissions from fossil fuel combustion inventory estimation model and about 35 from the stationary source inventory models).

In developing the uncertainty estimation model, uniform distribution was assumed for all activity-related input variables and N<sub>2</sub>O emission factors, based on the SAIC/EIA (2001) report.<sup>136</sup> For these variables, the uncertainty ranges

<sup>133</sup> SAIC/EIA (2001) characterizes the underlying probability density function for the input variables as a combination of uniform and normal distributions (the former to represent the bias component and the latter to represent the random component). However, for purposes of the current uncertainty analysis, it was determined that uniform distribution was more appropriate to characterize the probability density function underlying each of these variables.

<sup>134</sup> In the SAIC/EIA (2001) report, the quantitative uncertainty estimates were developed for each of the three major fossil fuels used within each end-use sector; the variations within the sub-fuel types within each end-use sector were not modeled. However, for purposes of assigning uncertainty estimates to the sub-fuel type categories within each end-use sector in the current uncertainty analysis, SAIC/EIA (2001)-reported uncertainty estimates were extrapolated.

<sup>135</sup> Although, in general, random uncertainties are the main focus of statistical uncertainty analysis, when the uncertainty estimates are elicited from experts, their estimates include both random and systematic uncertainties. Hence, both these types of uncertainties are represented in this uncertainty analysis.

<sup>136</sup> SAIC/EIA (2001) characterizes the underlying probability density function for the input variables as a combination of uniform and normal distributions (the former distribution to represent the bias component and the latter to represent the random

were assigned to the input variables based on the data reported in SAIC/EIA (2001).<sup>137</sup> However, the CH<sub>4</sub> emission factors differ from those used by EIA. Since these factors were obtained from IPCC/UNEP/OECD/IEA (1997), uncertainty ranges were assigned based on IPCC default uncertainty estimates (IPCC 2006).

### **CH<sub>4</sub> and N<sub>2</sub>O from Mobile Combustion**

The uncertainty analysis was performed on 2014 estimates of CH<sub>4</sub> and N<sub>2</sub>O emissions, incorporating probability distribution functions associated with the major input variables. For the purposes of this analysis, the uncertainty was modeled for the following four major sets of input variables: (1) VMT data, by on-road vehicle and fuel type, (2) emission factor data, by on-road vehicle, fuel, and control technology type, (3) fuel consumption, data, by non-road vehicle and equipment type, and (4) emission factor data, by non-road vehicle and equipment type.

### **Carbon Emitted from Non-Energy Uses of Fossil Fuels**

An uncertainty analysis was conducted to quantify the uncertainty surrounding the estimates of emissions and storage factors from non-energy uses.

The non-energy use analysis is based on U.S.-specific storage factors for (1) feedstock materials (natural gas, LPG, pentanes plus, naphthas, other oils, still gas, special naphthas, and other industrial coal), (2) asphalt, (3) lubricants, and (4) waxes. To characterize uncertainty, five separate analyses were conducted, corresponding to each of the five categories. In all cases, statistical analyses or expert judgments of uncertainty were not available directly from the information sources for all the activity variables; thus, uncertainty estimates were determined using assumptions based on source category knowledge.

### **Incineration of Waste**

The uncertainties in the waste incineration emission estimates arise from both the assumptions applied to the data and from the quality of the data. Key factors include MSW incineration rate; fraction oxidized; missing data on waste composition; average C content of waste components; assumptions on the synthetic/biogenic C ratio; and combustion conditions affecting N<sub>2</sub>O emissions. The highest levels of uncertainty surround the variables that are based on assumptions (e.g., percent of clothing and footwear composed of synthetic rubber); the lowest levels of uncertainty surround variables that were determined by quantitative measurements (e.g., combustion efficiency, C content of C black).

### **Coal Mining**

The uncertainty associated with emission estimates from underground ventilation systems can be attributed to the fact that the actual measurement data from MSHA or EPA's Greenhouse Gas Reporting Program (GHGRP) used were not continuous but rather an average of quarterly instantaneous readings. Additionally, the measurement equipment used can be expected to have resulted in an average of 10 percent overestimation of annual CH<sub>4</sub> emissions (Mutmansky & Wang 2000). GHGRP data was used for a number of the mines beginning in 2013, however, the equipment uncertainty is applied to both MSHA and GHGRP data.

Estimates of CH<sub>4</sub> recovered by degasification systems are relatively certain for utilized CH<sub>4</sub> because of the availability of gas sales information. Many of the recovery estimates use data on wells within 100 feet of a mined area. However, uncertainty exists concerning the radius of influence of each well. The number of wells counted, and thus the avoided emissions, may vary if the drainage area is found to be larger or smaller than estimated.

### **Abandoned Underground Coal Mines**

The parameters for which values must be estimated for each mine in order to predict its decline curve are: 1) the coal's adsorption isotherm; 2) CH<sub>4</sub> flow capacity as expressed by permeability; and 3) pressure at abandonment. Because these parameters are not available for each mine, a methodological approach to estimating emissions was used that generates

component). However, for purposes of the current uncertainty analysis, it was determined that uniform distribution was more appropriate to characterize the probability density function underlying each of these variables.

<sup>137</sup> In the SAIC/EIA (2001) report, the quantitative uncertainty estimates were developed for each of the three major fossil fuels used within each end-use sector; the variations within the sub-fuel types within each end-use sector were not modeled. However, for purposes of assigning uncertainty estimates to the sub-fuel type categories within each end-use sector in the current uncertainty analysis, SAIC/EIA (2001)-reported uncertainty estimates were extrapolated.

a probability distribution of potential outcomes based on the most likely value and the probable range of values for each parameter. The range of values is not meant to capture the extreme values, but rather values that represent the highest and lowest quartile of the cumulative probability density function of each parameter. Once the low, mid, and high values are selected, they are applied to a probability density function.

### **Petroleum Systems**

The uncertainty analysis conducted for the 1990 through 2009 Inventory has not yet been updated for the 1990 through 2014 Inventory years; instead, the uncertainty percentage ranges calculated previously were applied to 2014 emission estimates. The majority of sources in the current Inventory were calculated using the same emission factors and activity data for which probability density functions were developed in the 1990 through 2009 uncertainty analysis.

### **Natural Gas Systems**

The uncertainty analysis conducted for the 1990 through 2009 Inventory has not yet been updated for the 1990 through 2014 Inventory years; instead, the uncertainty percentage ranges calculated previously were applied to 2014 emissions estimates. The majority of sources in the current Inventory were calculated using the same emission factors and activity data for which probability density functions were developed in the 1990 through 2009 uncertainty analysis.

### **International Bunker Fuels**

Emission estimates related to the consumption of international bunker fuels are subject to the same uncertainties as those from domestic aviation and marine mobile combustion emissions; however, additional uncertainties result from the difficulty in collecting accurate fuel consumption activity data for international transport activities separate from domestic transport activities. Uncertainties exist with regard to the total fuel used by military aircraft and ships, and in the activity data on military operations and training that were used to estimate percentages of total fuel use reported as bunker fuel emissions. There are also uncertainties in fuel end-uses by fuel-type, emissions factors, fuel densities, diesel fuel sulfur content, aircraft and vessel engine characteristics and fuel efficiencies, and the methodology used to back-calculate the data set to 1990 using the original set from 1995.

### **Wood Biomass and Ethanol Consumption**

It is assumed that the combustion efficiency for woody biomass is 100 percent, which is believed to be an overestimate of the efficiency of wood combustion processes in the United States. Decreasing the combustion efficiency would decrease emission estimates. Additionally, the heat content applied to the consumption of woody biomass in the residential, commercial, and electric power sectors is unlikely to be a completely accurate representation of the heat content for all the different types of woody biomass consumed within these sectors. Emission estimates from ethanol production are more certain than estimates from woody biomass consumption due to better activity data collection methods and uniform combustion techniques.

## **Industrial Processes and Product Use**

The uncertainty analysis descriptions in this section correspond to source categories included in the Industrial Processes and Product Use chapter of the Inventory.

### **Cement Production**

The uncertainties contained in these estimates are primarily due to uncertainties in the lime content of clinker and in the percentage of CKD recycled inside the cement kiln. Uncertainty is also associated with the assumption that all calcium-containing raw materials are  $\text{CaCO}_3$ , when a small percentage likely consists of other carbonate and non-carbonate raw materials.

### **Lime Production**

The uncertainties contained in these estimates can be attributed to slight differences in the chemical composition of lime products and  $\text{CO}_2$  recovery rates for on-site process use over the time series. Although the methodology accounts for various formulations of lime, it does not account for the trace impurities found in lime, such as iron oxide, alumina, and silica. In addition, a portion of the  $\text{CO}_2$  emitted during lime production will actually be reabsorbed when the lime is consumed, especially at captive lime production facilities. Another uncertainty is the assumption that calcination emissions for LKD are around 2 percent. There is limited data publicly available on LKD generation rates and also quantities, types of other byproducts/wastes produced at lime facilities.

### **Glass Production**

The uncertainty levels presented in this section arise in part due to variations in the chemical composition of limestone used in glass production. The uncertainty estimates also account for uncertainty associated with activity data. Large fluctuations in reported consumption exist, reflecting year-to-year changes in the number of survey responders. The accuracy of distribution by end use is also uncertain because this value is reported by the manufacturer of the input carbonates

(limestone, dolomite & soda ash) and not the end user. Additionally, there is significant inherent uncertainty associated with estimating withheld data points for specific end uses of limestone and dolomite. Lastly, much of the limestone consumed in the United States is reported as “other unspecified uses;” therefore, it is difficult to accurately allocate this unspecified quantity to the correct end-uses.

### **Other Process Uses of Carbonates**

The uncertainty levels presented in this section account for uncertainty associated with activity data. Data on limestone and dolomite consumption are collected by USGS through voluntary national surveys. Large fluctuations in reported consumption exist, reflecting year-to-year changes in the number of survey responders. The accuracy of distribution by end-use is also uncertain because this value is reported by the producer/mines and not the end-user. Additionally, there is significant inherent uncertainty associated with estimating withheld data points for specific end-uses of limestone and dolomite. Lastly, much of the limestone consumed in the United States is reported as “other unspecified uses;” therefore, it is difficult to accurately allocate this unspecified quantity to the correct end-uses. Uncertainty in the estimates also arises in part due to variations in the chemical composition of limestone.

### **Ammonia Production**

The uncertainties presented in this section are primarily due to how accurately the emission factor used represents an average across all ammonia plants using natural gas feedstock. Uncertainties are also associated with ammonia production estimates and the assumption that all ammonia production and subsequent urea production was from the same process. Uncertainty is also associated with the representativeness of the emission factor used for the petroleum coke-based ammonia process. It is also assumed that ammonia and urea are produced at collocated plants from the same natural gas raw material.

### **Urea Consumption for Non-Agricultural Purposes**

The primary uncertainties associated with this source category are associated with the accuracy of the estimates of urea production, urea imports, urea exports, and the amount of urea used as fertilizer as well as the fact that each estimate is obtained from a different data source. Because urea production estimates are no longer available from the USGS, there is additional uncertainty associated with urea produced beginning in 2011. There is also uncertainty associated with the assumption that all of the carbon in urea is released into the environment as CO<sub>2</sub> during use.

### **Nitric Acid Production**

Uncertainty associated with the parameters used to estimate N<sub>2</sub>O emissions includes that of production data, the share of U.S. nitric acid production attributable to each emission abatement technology over the time series (especially prior to 2010), and the associated emission factors applied to each abatement technology type.

### **Adipic Acid Production**

Uncertainty associated with N<sub>2</sub>O emission estimates includes the methods used by companies to monitor and estimate emissions.

### **Silicon Carbide Production and Consumption**

There is uncertainty associated with the emission factors used because they are based on stoichiometry as opposed to monitoring of actual SiC production plants. For CH<sub>4</sub>, there is also uncertainty associated with the hydrogen-containing volatile compounds in the petroleum coke (IPCC 2006). There is also uncertainty associated with the use or destruction of methane generated from the process in addition to uncertainty associated with levels of production, net imports, consumption levels, and the percent of total consumption that is attributed to metallurgical and other non-abrasive uses.

### **Titanium Dioxide Production**

As of 2004, the last remaining sulfate-process plant in the United States closed. Since annual TiO<sub>2</sub> production was not reported by USGS by the type of production process used (chloride or sulfate) prior to 2004 and only the percentage of total production capacity by process was reported, the percent of total TiO<sub>2</sub> production capacity that was attributed to the chloride process was multiplied by total TiO<sub>2</sub> production to estimate the amount of TiO<sub>2</sub> produced using the chloride process. Finally, the emission factor was applied uniformly to all chloride-process production, and no data were available to account for differences in production efficiency among chloride-process plants.

## **Soda Ash Production and Consumption**

Emission estimates from soda ash production have relatively low associated uncertainty levels in that reliable and accurate data sources are available for the emission factor and activity data. Soda ash production data was collected by the USGS from voluntary surveys. One source of uncertainty is the purity of the trona ore used for manufacturing soda ash. The primary source of uncertainty, however, results from the fact that emissions from soda ash consumption are dependent upon the type of processing employed by each end-use.

## **Petrochemical Production**

Sources of uncertainty on the CH<sub>4</sub> and CO<sub>2</sub> emission factors used for acrylonitrile and methanol production are derived from the use of default or average factors from a limited number of studies. There is some uncertainty in the applicability of the average emission factors for each petrochemical type across all prior years. While petrochemical production processes in the United States have not changed significantly since 1990, some operational efficiencies have been implemented at facilities over the time series.

## **HCFC-22 Production**

The uncertainty analysis presented in this section was based on a plant-level Monte Carlo Stochastic Simulation for 2006. A normal probability density function was assumed for all measurements and biases except the equipment leak estimates for one plant; a log-normal probability density function was used for this plant's equipment leak estimates. The simulation for 2006 yielded a 95-percent confidence interval for U.S. emissions of 6.8 percent below to 9.6 percent above the reported total.

The relative errors yielded by the Monte Carlo Stochastic Simulation for 2006 were applied to the U.S. emission estimate for 2014. The resulting estimates of absolute uncertainty are likely to be reasonably accurate because (1) the methods used by the three plants to estimate their emissions are not believed to have changed significantly since 2006, and (2) although the distribution of emissions among the plants may have changed between 2006 and 2014, the two plants that contribute significantly to emissions were estimated to have similar relative uncertainties in their 2006 (as well as 2005) emission estimates.

## **Carbon Dioxide Production**

Uncertainty is associated with the data reported through EPA's GHGRP, specifically the amount of CO<sub>2</sub> consumed for food and beverage applications given a threshold for reporting under GHGRP applicable to those reporting under Subpart PP, in addition to the exclusion of the amount of CO<sub>2</sub> transferred to all other end-use categories. Uncertainty is also associated with the exclusion of imports/exports data for CO<sub>2</sub> suppliers.

## **Phosphoric Acid Production**

Regional production for 2014 was estimated based on regional production data from previous years and multiplied by regionally-specific emission factors. There is uncertainty associated with the degree to which the estimated 2014 regional production data represents actual production in those regions.

An additional source of uncertainty is the carbonate composition of phosphate rock; the composition of phosphate rock varies depending upon where the material is mined, and may also vary over time. A third source of uncertainty is the assumption that all domestically-produced phosphate rock is used in phosphoric acid production and used without first being calcined.

## **Iron and Steel Production and Metallurgical Coke Production**

Uncertainty is associated with the total U.S. coking coal consumption, total U.S. coke production, and materials consumed during this process. Therefore, for the purpose of this analysis, uncertainty parameters are applied to primary data inputs to the calculation (i.e., coking coal consumption and metallurgical coke production) only.

There is uncertainty associated with the assumption that direct reduced iron and sinter consumption are equal to production. There is uncertainty associated with the assumption that all coal used for purposes other than coking coal is for direct injection coal; some of this coal may be used for electricity generation. There is also uncertainty associated with the C contents for pellets, sinter, and natural ore. For electric arc furnace (EAF) steel production, there is uncertainty associated with the amount of EAF anode and charge C consumed due to inconsistent data throughout the time series. Also for EAF steel production, there is uncertainty associated with the assumption that 100 percent of the natural gas attributed to "steelmaking furnaces" by AISI is process-related and nothing is combusted for energy purposes. Uncertainty is also associated with the use of process gases such as blast furnace gas and coke oven gas.

## **Ferroalloy Production**

Uncertainty for this source is associated with the type and availability of annual ferroalloy production data, which have varied over the time series. Such production data may or may not include details such as ferroalloy content, production

practices (e.g., biomass used as primary or secondary carbon source), amount of reducing agent used, and furnace specifics (e.g., type, operation technique, control technology).

### **Aluminum Production**

Uncertainty was assigned to the CO<sub>2</sub>, CF<sub>4</sub>, and C<sub>2</sub>F<sub>6</sub> emission values reported by each individual facility to EPA's GHGRP. Uncertainty surrounding the reported CO<sub>2</sub>, CF<sub>4</sub>, and C<sub>2</sub>F<sub>6</sub> emission values were determined to have a normal distribution with uncertainty ranges of ±6, ±16, and ±20 percent, respectively.

### **Magnesium Production**

Uncertainty surrounding the total estimated emissions in 2014 is attributed to the uncertainties around SF<sub>6</sub>, HFC-134a and CO<sub>2</sub> emission estimates. To estimate the uncertainty surrounding the estimated 2014 SF<sub>6</sub> emissions from magnesium production and processing, the uncertainties associated with three variables were estimated: (1) emissions reported by magnesium producers and processors for 2014 through EPA's GHGRP, (2) emissions estimated for magnesium producers and processors that reported via the Partnership in prior years but did not report 2014 emissions through EPA's GHGRP, and (3) emissions estimated for magnesium producers and processors that did not participate in the Partnership or report through EPA's GHGRP. Additional uncertainties exist in these estimates that are not addressed in this methodology, such as the basic assumption that SF<sub>6</sub> neither reacts nor decomposes during use.

### **Lead Production**

Uncertainty associated with lead production relates to the applicability of emission factors and the accuracy of primary and secondary production data provided by the USGS.

### **Zinc Production**

There is uncertainty associated with the amount of EAF dust consumed in the United States to produce secondary zinc using emission-intensive Waelz kilns.

There are also uncertainties associated with the accuracy of the emission factors used to estimate CO<sub>2</sub> emissions from secondary zinc production processes.

### **Semiconductor Manufacture**

The equation used to estimate uncertainty is:

$$\text{Total Emissions (E}_T\text{)} = \text{GHGRP Reported F-GHG Emissions (E}_{R,\text{F-GHG}}\text{)} + \text{Non-Reporters' Estimated F-GHG Emissions (E}_{NR,\text{F-GHG}}\text{)} + \text{GHGRP Reported N}_2\text{O Emissions (E}_{R,\text{N}_2\text{O}}\text{)} + \text{Non-Reporters' Estimated N}_2\text{O Emissions (E}_{NR,\text{N}_2\text{O}}\text{)}$$

where E<sub>R</sub> and E<sub>NR</sub> denote totals for the indicated subcategories of emissions for F-GHG and N<sub>2</sub>O, respectively.

The uncertainty estimate of E<sub>R, F-GHG</sub>, or GHGRP reported F-GHG emissions, is developed based on gas-specific uncertainty estimates of emissions for two industry segments, one processing 200 mm wafers and one processing 300 mm wafers. These gas and wafer-specific uncertainty estimates are applied to the total emissions of the facilities that did not abate emissions as reported under EPA's GHGRP.

For those facilities reporting abatement of emissions under EPA's GHGRP, estimates of uncertainties for the no abatement industry segments are modified to reflect the use of full and partial abatement. For all facilities reporting gas abatement, a triangular distribution of destruction or removal efficiency is assumed for each gas. For facilities reporting partial abatement, the distribution of fraction of the gas fed through the abatement device, for each gas, is assumed to be triangularly distributed as well. Gas-specific emission uncertainties were estimated by convolving the distributions of unabated emissions with the appropriate distribution of abatement efficiency for fully and partially abated facilities using a Montel Carlo simulation.

The uncertainty in E<sub>R, F-GHG</sub> is obtained by allocating the estimates of uncertainties to the total GHGRP-reported emissions from each of the six industry segments. The uncertainty in E<sub>R, N<sub>2</sub>O</sub> is obtained by assuming that the uncertainty in the emissions reported by each of the GHGRP reporting facilities results from the uncertainty in quantity of N<sub>2</sub>O consumed and the N<sub>2</sub>O emission factor (or utilization). The quantity of N<sub>2</sub>O utilized (the complement of the emission factor) was assumed to have a triangular distribution with a minimum value of 0 percent, mode of 20 percent and maximum value of 84

percent. The uncertainty for the total reported N<sub>2</sub>O emissions was then estimated by combining the uncertainties of each of the facilities reported emissions using Monte Carlo simulation. The estimate of uncertainty in E<sub>NR,F-GHG</sub> and E<sub>NR,N<sub>2</sub>O</sub> entailed developing estimates of uncertainties for the proportion of total emissions attributed to non-reporters for 2011 and 2012, which are subsequently dependent on the emissions factors for each non-reporting sub-category and the corresponding estimates of Total Manufactured Layer Area (TMLA) for the years 2011 and 2012.

The uncertainty in TMLA depends on the uncertainty of two variables – an estimate of the uncertainty in the average annual capacity utilization for each level of production of fabs (e.g., full scale or R&D production) and a corresponding estimate of the uncertainty in the number of layers manufactured. For both variables, the distributions of capacity utilizations and number of manufactured layers are assumed triangular for all categories of non-reporting fabs. For production fabs and for facilities that manufacture discrete devices, the most probable utilization is assumed to be 79 percent for 2011 and 65 percent for 2012, with the highest and lowest utilization assumed to be 84 percent for 2011 and 70 percent for 2012, and 62 percent for 2011 and 51 percent for 2012, respectively. The most probable values for utilization for R&D facilities are assumed to be 79 percent for 2011 and 65 percent for 2012, with the highest utilization also at 79 percent for 2011 and 65 percent for 2012, and the lowest utilization at 40 percent for 2011 and 33 percent for 2012. For the triangular distributions that govern the number of possible layers manufactured, it is assumed the most probable value is one layer less than reported in the ITRS.

The uncertainty bounds for the average capacity utilization and the number of layers manufactured are used as inputs in a separate Monte Carlo simulation to estimate the uncertainty around the TMLA of both individual facilities as well as the total non-reporting TMLA of each sub-population for both 2011 and 2012. The uncertainty around the emission factors for each non-reporting category of facilities (for both 2011 and 2012) is dependent on the uncertainty of the total emissions (MMT CO<sub>2</sub> Eq. units) and the TMLA of each reporting facility in that category. For simplicity, the results of the Monte Carlo simulations on the bounds of the gas- and wafer size-specific emissions as well as the TMLA and emission factors are assumed to be normally distributed and the uncertainty bounds are assigned at 1.96 standard deviations around the estimated mean. The departures from normality were observed to be small. The final step in estimating the uncertainty in emissions of non-reporting facilities is convolving the distribution of emission factors with the distribution of TMLA individually for 2011 and 2012, along with the distributions of GHGRP-reported emissions for 2011 and 2012 to estimate the uncertainty around the proportion of total emissions attributed to non-reporters for 2011 and 2012 using Monte Carlo simulation.

### **Substitution of Ozone Depleting Substances**

Given that emissions of ODS substitutes occur from thousands of different kinds of equipment and from millions of point and mobile sources throughout the United States, significant uncertainties exist with regard to the levels of equipment sales, equipment characteristics, and end-use emissions profiles that were used to estimate annual emissions for the various compounds.

The Vintaging Model estimates emissions from 66 end-uses. The uncertainty analysis quantifies the level of uncertainty associated with the aggregate emissions resulting from the top 21 end-uses, comprising over 95 percent of the total emissions, and 6 other end-uses. These 27 end-uses comprise 97 percent of the total emissions, equivalent to 156.4 MMT CO<sub>2</sub> Eq.

In order to calculate uncertainty, functional forms were developed to simplify some of the complex “vintaging” aspects of some end-use sectors, especially with respect to refrigeration and air-conditioning, and to a lesser degree, fire extinguishing. The functional forms used variables that included. Uncertainty was estimated around each variable within the functional forms (e.g., growth rates, emission factors, transition from ODSs, change in charge size as a result of the transition, disposal quantities, disposal emission rates, and either stock for the current year or original ODS consumption) based on expert judgment. The most significant sources of uncertainty for this source category include the emission factors for residential unitary AC, as well as the percent of non-MDI aerosol propellant that is HFC-152a.

### **Electrical Transmission and Distribution**

To estimate the uncertainty associated with emissions of SF<sub>6</sub> from Electrical Transmission and Distribution, uncertainties associated with four quantities were estimated: (1) emissions from Partners, (2) emissions from GHGRP-Only Reporters, (3) emissions from Non-Reporters, and (4) emissions from manufacturers of electrical equipment. Uncertainties were also estimated regarding (1) the quantity of SF<sub>6</sub> supplied with equipment by equipment manufacturers, which is projected from Partner provided nameplate capacity data and industry SF<sub>6</sub> nameplate capacity estimates, and (2) the manufacturers’ SF<sub>6</sub> emissions rate.

### **Nitrous Oxide from Product Uses**

The overall uncertainty associated with the 2014 N<sub>2</sub>O emission estimate from N<sub>2</sub>O product usage was calculated using the 2006 IPCC Guidelines (IPCC 2006) Approach 2 methodology. Uncertainty associated with the parameters used

to estimate N<sub>2</sub>O emissions include production data, total market share of each end use, and the emission factors applied to each end use, respectively.

## **Agriculture**

The uncertainty analysis descriptions in this section correspond to some source categories included in the Agriculture chapter of the Inventory.

### **Enteric Fermentation**

Uncertainty estimates were developed for the 1990 through 2001 Inventory report (i.e., 2003 submission to the UNFCCC). There have been no significant changes to the methodology since that time; consequently, these uncertainty estimates were directly applied to the 2014 emission estimates in this Inventory report.

A total of 185 primary input variables were identified as key input variables for the uncertainty analysis. A normal distribution was assumed for almost all activity- and emission factor-related input variables. Triangular distributions were assigned to three input variables to ensure only positive values would be simulated.

### **Manure Management**

An analysis (ERG 2003a) was conducted for the manure management emission estimates presented in the 1990 through 2001 Inventory report (i.e., 2003 submission to the UNFCCC) to determine the uncertainty associated with estimating CH<sub>4</sub> and N<sub>2</sub>O emissions from livestock manure management. These uncertainty estimates were directly applied to the 2014 emission estimates as there have not been significant changes in the methodology since that time.

### **Rice Cultivation**

Sources of uncertainty in the Tier 3 method include management practices, uncertainties in model structure (i.e., algorithms and parameterization), and variance associated with the NRI sample. Sources of uncertainty in the IPCC (2006) Tier 1 method include the emission factors, management practices, and variance associated with the NRI sample. A Monte Carlo analysis was used to propagate uncertainties in the Tier 1 and 3 methods, and the uncertainties from each approach are combined to produce the final CH<sub>4</sub> emissions estimate using simple error propagation (IPCC 2006).

### **Agricultural Soil Management**

Uncertainty was estimated for each of the following five components of N<sub>2</sub>O emissions from agricultural soil management: (1) direct emissions simulated by DAYCENT; (2) the components of indirect emissions (N volatilized and leached or runoff) simulated by DAYCENT; (3) direct emissions approximated with the IPCC (2006) Approach 1 method; (4) the components of indirect emissions (N volatilized and leached or runoff) approximated with the IPCC (2006) Approach 1 method; and (5) indirect emissions estimated with the IPCC (2006) Approach 1 method.

### **Field Burning of Agricultural Residues**

Due to data limitations, uncertainty resulting from the fact that emissions from burning of Kentucky bluegrass and “other crop” residues are not included in the emissions estimates was not incorporated into the uncertainty analysis.

## **Land Use, Land-Use Change, and Forestry**

The uncertainty analysis descriptions in this section correspond to source categories included in the Land Use, Land-Use Change, and Forestry chapter of the Inventory.

### **Forest Land Remaining Forest Land**

The uncertainty analysis descriptions in this section correspond to source categories included in the *Forest Land Remaining Forest Land* sub-chapter of Land Use, Land-Use Change, and Forestry chapter of the Inventory.

### **Changes in Forest Carbon Stocks**

A quantitative uncertainty analysis placed bounds on current flux for forest ecosystems as well as C in harvested wood products through Monte Carlo Stochastic Simulation of the Methods and probabilistic sampling of C conversion factors and inventory data.

## **Non-CO<sub>2</sub> Emissions from Forest Fires**

In order to quantify the uncertainties for emissions from forest fires calculated as described above, a Monte Carlo (IPCC Approach 2) sampling approach was employed to propagate uncertainty in the equation as it was applied for U.S. forest land. See IPCC (2006) and Annex 3.13 for the quantities and assumptions employed to define and propagate uncertainty.

### **Direct N<sub>2</sub>O fluxes from Forest Soils**

The amount of N<sub>2</sub>O emitted from forests depends not only on N inputs and fertilized area, but also on a large number of variables, including organic C availability, oxygen gas partial pressure, soil moisture content, pH, temperature, and tree planting/harvesting cycles. The effect of the combined interaction of these variables on N<sub>2</sub>O flux is complex and highly uncertain.

Uncertainties exist in the fertilization rates, annual area of forest lands receiving fertilizer, and the emission factors. The uncertainty ranges around the 2005 activity data and emission factor input variables were directly applied to the 2014 emissions estimates. IPCC (2006) provided estimates for the uncertainty associated with direct and indirect N<sub>2</sub>O emission factor for synthetic N fertilizer application to soils.

### **Cropland Remaining Cropland**

The uncertainty analysis descriptions in this section correspond to source categories included in the *Cropland Remaining Cropland* sub-chapter of Land Use, Land-Use Change, and Forestry chapter of the Inventory.

#### **Agricultural Soil Carbon Stock Change**

Uncertainty associated with the *Cropland Remaining Cropland* land-use category was addressed for changes in agricultural soil C stocks (including both mineral and organic soils).

#### **CO<sub>2</sub> Emissions from Liming**

Uncertainty regarding the amount of limestone and dolomite applied to soils was estimated at ±15 percent with normal densities (Tepordei 2003, Willett 2013b). Analysis of the uncertainty associated with the emission factors included the fraction of lime dissolved by nitric acid versus the fraction that reacts with carbonic acid, and the portion of bicarbonate that leaches through the soil and is transported to the ocean. The probability distribution functions for the fraction of lime dissolved by nitric acid and the portion of bicarbonate that leaches through the soil were represented as smoothed triangular distributions between ranges of zero and 100 percent of the estimates.

#### **CO<sub>2</sub> Emissions from Urea Fertilization**

The largest source of uncertainty was the default emission factor, which assumes that 100 percent of the C in CO(NH<sub>2</sub>)<sub>2</sub> applied to soils is ultimately emitted into the environment as CO<sub>2</sub>. In addition, each urea consumption data point has an associated uncertainty.

#### **Land Converted to Cropland**

Uncertainty analysis for mineral soil C stock changes using the Tier 3 and Tier 2 methodologies are based on the same method described for *Cropland Remaining Cropland*.

Uncertainty was estimated for each subsource (i.e., biomass C stocks, mineral soil C stocks and organic soil C stocks) and method that was used in the Inventory analysis (i.e., Tier 2 and Tier 3).

#### **Grassland Remaining Grassland**

Uncertainty was estimated for each subsource (i.e., mineral soil C stocks and organic soil C stocks) and disaggregated to the level of the inventory methodology employed (i.e., Tier 2 and Tier 3).

#### **Land Converted to Grassland**

Uncertainty was estimated for each subsource (i.e., biomass C stocks, mineral soil C stocks and organic soil C stocks) and disaggregated to the level of the inventory methodology employed (i.e., Tier 2 and Tier 3).

#### **Wetlands Remaining Wetlands**

The uncertainty analysis descriptions in this section correspond to source categories included in the *Wetlands Remaining Wetlands* sub-chapter of Land Use, Land-Use Change, and Forestry chapter of the Inventory.

#### **Peatlands Remaining Peatlands**

The uncertainty associated with peat production data was estimated to be ± 25 percent (Apodaca 2008), assumed to be normally distributed, and is attributed to the USGS receives data from the smaller peat producers but estimates production from some larger peat distributors. The uncertainty associated with the reported production data for Alaska was

assumed to be the same as for the lower 48 states, or  $\pm 25$  percent with a normal distribution. The uncertainty associated with the average bulk density values was estimated to be  $\pm 25$  percent with a normal distribution (Apodaca 2008). The uncertainty associated with the emission factors was assumed to be triangularly distributed. The uncertainty values surrounding the C fractions were based on IPCC (2006) and the uncertainty was assumed to be uniformly distributed. The uncertainty values associated with the fraction of peatland covered by ditches was assumed to be  $\pm 100$  percent with a normal distribution based on the assumption that greater than 10 percent coverage, the upper uncertainty bound, is not typical of drained organic soils outside of The Netherlands (IPCC 2013).

### **Settlements Remaining Settlements**

The uncertainty analysis descriptions in this section correspond to source categories included in the *Settlements Remaining Settlements* sub-chapter of Land Use, Land-Use Change, and Forestry chapter of the Inventory.

### **Changes in Carbon Stocks in Urban Trees**

Uncertainty associated with changes in C stocks in urban trees includes the uncertainty associated with urban area, percent urban tree coverage, and estimates of gross and net C sequestration for each of the 50 states and the District of Columbia. Additional uncertainty is associated with the biomass equations, conversion factors, and decomposition assumptions used to calculate C sequestration and emission estimates (Nowak et al. 2002).

### **N<sub>2</sub>O Fluxes from Settlement Soils**

The amount of N<sub>2</sub>O emitted from settlements depends not only on N inputs and fertilized area, but also on a large number of variables, including organic C availability, oxygen gas partial pressure, soil moisture content, pH, temperature, and irrigation/watering practices. The effect of the combined interaction of these variables on N<sub>2</sub>O flux is complex and highly uncertain.

Uncertainties exist in both the fertilizer N and sewage sludge application rates in addition to the emission factors. Uncertainty in the amounts of sewage sludge applied to non-agricultural lands and used in surface disposal was derived from variability in several factors. The uncertainty ranges around 2005 activity data and emission factor input variables were directly applied to the 2014 emission estimates.

### **Other**

The uncertainty analysis descriptions in this section correspond to source categories included in the Other sub-chapter of Land Use, Land-Use Change, and Forestry chapter of the Inventory.

### **Changes in Yard Trimming and Food Scrap Carbon Stocks in Landfills**

The uncertainty analysis for landfilled yard trimmings and food scraps includes an evaluation of the effects of uncertainty for the following data and factors: disposal in landfills per year (tons of C), initial C content, moisture content, decay rate, and proportion of C stored. The C storage landfill estimates are also a function of the composition of the yard trimmings (i.e., the proportions of grass, leaves and branches in the yard trimmings mixture). There are respective uncertainties associated with each of these factors.

## **Waste**

The uncertainty analysis descriptions in this section correspond to source categories included in the Waste chapter of the Inventory.

### **Landfills**

The primary uncertainty associated with the estimates of CH<sub>4</sub> emissions from MSW and industrial waste landfills concerns the characterization of landfills. There is also a high degree of uncertainty and variability associated with the first order decay model, particularly when a homogeneous waste composition and hypothetical decomposition rates are applied to heterogeneous landfills (IPCC 2006).

The lack of landfill-specific information regarding the number and type of industrial waste landfills in the United States is a primary uncertainty with respect to the industrial waste generation and emissions estimates. Uncertainty also exists in the estimates of the landfill gas oxidized. Another significant source of uncertainty lies with the estimates of CH<sub>4</sub> recovered by flaring and gas-to-energy projects at MSW landfills. Industrial waste landfills are shown with a lower range of uncertainty due to the smaller number of data sources and associated uncertainty involved.

### **Wastewater Treatment**

Uncertainty associated with the parameters used to estimate CH<sub>4</sub> emissions from wastewater treatment include that of numerous input variables used to model emissions from domestic wastewater, and wastewater from pulp and paper manufacture, meat and poultry processing, fruits and vegetable processing, ethanol production, and petroleum refining. Uncertainty associated with the parameters used to estimate N<sub>2</sub>O emissions include that of sewage sludge disposal, total U.S. population, average protein consumed per person, fraction of N in protein, non-consumption nitrogen factor, emission factors per capita and per mass of sewage-N, and for the percentage of total population using centralized wastewater treatment plants.

### **Composting**

The estimated uncertainty from the *2006 IPCC Guidelines* is  $\pm 50$  percent for the Approach 1 methodology.

## **References**

- EPA (2002) *Quality Assurance/Quality Control and Uncertainty Management Plan for the U.S. Greenhouse Gas Inventory: Procedures Manual for Quality Assurance/Quality Control and Uncertainty Analysis*, U.S. Greenhouse Gas Inventory Program, U.S. Environmental Protection Agency, Office of Atmospheric Programs, EPA 430-R-02-007B, June 2002.
- IPCC (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*, Paris: Intergovernmental Panel on Climate Change, United Nations Environment Programme, Organization for Economic Co-Operation and Development, International Energy Agency.
- USGS (2011) *2010 Mineral Yearbook; Aluminum [Advanced Release]*. U.S. Geological Survey, Reston, VA.

# ANNEX 8 QA/QC Procedures

## 8.1. Background

The purpose of this annex is to describe the Quality Assurance/Quality Control (QA/QC) procedures and information quality considerations that are used throughout the process of creating and compiling the *Inventory of U.S. Greenhouse Gas Emissions and Sinks*. This includes the evaluation of the quality and relevance of data and models used as inputs into the Inventory; proper management, incorporation, and aggregation of data; and review of the numbers and estimates to ensure that they are as accurate and transparent as possible. Quality control—in the form of both good practices (such as documentation procedures) and checks on whether good practices and procedures are being followed—is applied at every stage of inventory development and document preparation. In addition, quality assurance occurs at two stages—an expert review and a public review. While both phases can significantly contribute to the quality of the Inventory, the public review phase is also essential for promoting the openness of the Inventory development process and the transparency of the inventory data and methods.

## 8.2. Purpose

The *Quality Assurance/Quality Control and Uncertainty Management Plan* for the Inventory (QA/QC Management Plan) guides the process of ensuring the quality of the Inventory. The QA/QC Management Plan describes data and methodology checks, develops processes governing peer review and public comments, and provides guidance on conducting an analysis of the uncertainty surrounding the emission estimates. The QA/QC Management Plan procedures also stress continual improvement, providing for corrective actions that are designed to improve the inventory estimates over time.


Key attributes of the QA/QC Management Plan are summarized in Figure A-25. These attributes include:

- *Procedures and Forms*: detailed and specific systems that serve to standardize the process of documenting and archiving information, as well as to guide the implementation of QA/QC and the analysis of uncertainty
- *Implementation of Procedures*: application of QA/QC procedures throughout the whole Inventory development process from initial data collection, through preparation of the emission estimates, to publication of the Inventory
- *Quality Assurance*: expert and public reviews for both the Inventory estimates and the report (which is the primary vehicle for disseminating the results of the Inventory development process)
- *Quality Control*: consideration of secondary data and source-specific checks (Tier 2 QC) in parallel, and coordination with the uncertainty assessment; the development of protocols and templates, which provide for more structured communication and integration with the suppliers of secondary information
- *Tier 1 (general) and Tier 2 (source-specific) Checks*: quality controls and checks, as recommended by the *IPCC Good Practice Guidance*
- *Record Keeping*: provisions to track which procedures have been followed, the results of the QA/QC process, uncertainty analysis, and feedback mechanisms for corrective action based on the results of the investigations, which provide for continual data quality improvement and guided research efforts
- *Multi-Year Implementation*: a schedule for coordinating the application of QA/QC procedures across multiple years
- *Interaction and Coordination*: promoting communication within the EPA, across Federal agencies and departments, state government programs, and research institutions and consulting firms involved in supplying data or preparing estimates for the Inventory. The QA/QC Management Plan itself is intended to be revised to

reflect new information that becomes available as the program develops, methods are improved, or additional supporting documents become necessary.

In addition, based on the national QA/QC Management Plan for the Inventory, source-specific QA/QC plans have been developed for a number of sources. These plans follow the procedures outlined in the national QA/QC plan, tailoring the procedures to the specific text and spreadsheets of the individual sources. For each greenhouse gas emissions source or sink included in this Inventory, a minimum of a Tier 1 QA/QC analysis has been undertaken. Where QA/QC activities for a particular source go beyond the minimum Tier 1 level, further explanation is provided within the respective source category text.

**Figure A-25: U.S. QA/QC Plan Summary**



### 8.3. Assessment Factors

The *Inventory of U.S. Greenhouse Gas Emissions and Sinks* development process follows guidance outlined in EPA’s *Guidelines for Ensuring and Maximizing the Quality, Objectivity, Utility, and Integrity, of Information Disseminated by the Environmental Protection Agency*<sup>138</sup> and *A Summary of General Assessment Factors for Evaluating the Quality of Scientific and Technical Information*.<sup>139</sup> This includes evaluating the data and models used as inputs into the Inventory against the five general assessment factors: soundness, applicability and utility, clarity and completeness,

<sup>138</sup> EPA report #260R-02-008, October 2002, Available online at <<http://www.epa.gov/quality/guidelines-ensuring-and-maximizing-quality-objectivity-utility-and-integrity-information>>.

<sup>139</sup> EPA report #100/B-03/001, June 2003, Available online at <<http://www.epa.gov/risk/guidance-evaluating-and-documenting-quality-existing-scientific-and-technical-information>>, and Addendum to: *A Summary of General Assessment Factors for Evaluating the Quality of Scientific and Technical Information*, December 2012, Available online at <<http://www.epa.gov/risk/summary-general-assessment-factors-evaluating-quality-scientific-and-technical-information>>.

uncertainty and variability, evaluation and review. Table A-304 defines each factor and explains how it was considered during the process of creating the current Inventory.

**Table A-304: Assessment Factors and Definitions**

| General Assessment Factor         | Definition  | How the Factor was Considered   |
|-----------------------------------|---|---|
| Soundness (AF1)                   | The extent to which the scientific and technical procedures, measures, methods or models employed to generate the information are reasonable for, and consistent with their intended application. | <p>The underlying data, methodologies, and models used to generate the <i>Inventory of U.S. Greenhouse Gas Emissions and Sinks</i> are reasonable for and consistent with their intended application, to provide information regarding all sources and sinks of greenhouse gases in the United States for the Inventory year, as required per UNFCCC Annex I country reporting requirements.</p> <p>The U.S. emissions calculations follow the <i>2006 IPCC Guidelines</i> developed specifically for UNFCCC inventory reporting. They are based on the best available, peer-reviewed scientific information, and have been used by the international community for over 20 years. When possible, Tier 2 and Tier 3 methodologies from the <i>2006 IPCC Guidelines</i> are applied to calculate U.S. emissions more accurately.</p> |
| Applicability and Utility (AF2)   | The extent to which the information is relevant for the Agency's intended use.  | The Inventory's underlying data, methodology, and models are relevant for their intended application because they generate the sector-specific greenhouse gas emissions trends necessary for assessing and understanding all sources and sinks of greenhouse gases in the United States for the Inventory year. They are relevant for communicating U.S. emissions information to domestic audiences, and they are consistent with the <i>2006 IPCC Guidelines</i> developed specifically for UNFCCC reporting purposes of international greenhouse gas inventories.  |
| Clarity and Completeness (AF3)    | The degree of clarity and completeness with which the data, assumptions, methods, quality assurance, sponsoring organizations and analyses employed to generate the information are documented.   | The methodological and calculation approaches applied to generate the <i>Inventory of U.S. Greenhouse Gas Emissions and Sinks</i> are extensively documented in the <i>2006 IPCC Guidelines</i> . The Inventory report describes its adherence to the <i>2006 IPCC Guidelines</i> , and the U.S. Government agencies provide data to implement the <i>2006 IPCC Guidelines</i> approaches. Any changes made to calculations, due to updated data and methods, are explained and documented in the report consistent with UNFCCC reporting guidelines.   |
| Uncertainty and Variability (AF4) | The extent to which the variability and uncertainty (quantitative and qualitative)  | The evaluation of uncertainties for underlying data is documented in the Uncertainty section of the Annex to the <i>Inventory of U.S. Greenhouse Gas Emissions and Sinks</i> . In   |

|                                    |   |  |
|------------------------------------|---|--|
|                                    | <p>in the information or in the procedures, measures, methods or models are evaluated and characterized.</p>                                    | <p>accordance with the <i>2006 IPCC Guidelines</i>, the uncertainty associated with the Inventory's underlying data, methodology, and models was evaluated by running a Monte-Carlo uncertainty analysis on source category emissions data to produce a 95 percent confidence interval for the annual greenhouse gas emissions for that source. To develop overall uncertainty estimates, the Monte Carlo simulation output data for each emission source category uncertainty analysis were combined by type of gas, and the probability distributions were fitted to the combined simulation output data where such simulated output data were available.</p>  |
| <p>Evaluation and Review (AF5)</p> | <p>The extent of independent verification, validation and peer review of the information or of the procedures, measures, methods or models.</p> | <p>The majority of the underlying methodology, calculations, and models used to generate the <i>Inventory of U.S. Greenhouse Gas Emissions and Sinks</i> have been independently verified and peer reviewed as part of their publication in the <i>2006 IPCC Guidelines</i>. In cases where the methodology differs slightly from the <i>2006 IPCC Guidelines</i>, these were independently verified and validated by technical experts during the annual expert review phase of the Inventory development process.</p> <p>For the data used in calculating greenhouse gas emissions for each source, multiple levels of evaluation and review occur. Data are compared to results from previous years, and calculations and equations are continually evaluated and updated as appropriate. Throughout the process, inventory data and methodological improvements are planned and incorporated.</p> <p>The Inventory undergoes annual cycles of expert and public review before publication. This process ensures that both experts and the general public can review each source of emissions and sinks, and have an extended opportunity to provide feedback on the methodologies used, calculations, data sources, and presentation of information.</p> |