


## 2. Trends in Greenhouse Gas Emissions

### 2.1 Recent Trends in U.S. Greenhouse Gas Emissions and Sinks


In 2013, total U.S. greenhouse gas emissions were 6,742.2 MMT or million metric tons CO<sub>2</sub> Eq. Total U.S. emissions have increased by 7.0 percent from 1990 to 2013, and emissions increased from 2012 to 2013 by 1.8 percent (118.8 MMT CO<sub>2</sub> Eq.). The increase from 2012 to 2013 was due to an increase in the carbon intensity of fuels consumed to generate electricity due to an increase in coal consumption, with decreased natural gas consumption. Additionally, cold winter conditions lead to an increase in fuels for the residential and commercial sectors for heating. Lastly, transportation emissions increased as a result of a small increase in vehicle miles traveled (VMT) and fuel use across on-road transportation modes. Since 1990, U.S. emissions have increased at an average annual rate of 0.3 percent. Figure 2-1 through Figure 2-3 illustrate the overall trend in total U.S. emissions by gas, annual changes, and absolute changes since 1990.

**Figure 2-1: U.S. Greenhouse Gas Emissions by Gas**


Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



1 **Figure 2-2: Annual Percent Change in U.S. Greenhouse Gas Emissions**



2  
3 **Figure 2-3: Cumulative Change in Annual U.S. Greenhouse Gas Emissions Relative to 1990**  
4 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



6  
7 Overall, from 1990 to 2013, total emissions of CO<sub>2</sub> increased by 429.2 MMT CO<sub>2</sub> Eq. (8.4 percent), while total  
8 emissions of CH<sub>4</sub> decreased by 86.4 MMT CO<sub>2</sub> Eq. (11.7 percent), and total emissions of N<sub>2</sub>O increased 25.0 MMT  
9 CO<sub>2</sub> Eq. (7.6 percent). During the same period, aggregate weighted emissions of HFCs, PFCs, SF<sub>6</sub>, and NF<sub>3</sub> rose by  
10 75.7 MMT CO<sub>2</sub> Eq. (74.2 percent). Despite being emitted in smaller quantities relative to the other principal  
11 greenhouse gases, emissions of HFCs, PFCs, SF<sub>6</sub>, and NF<sub>3</sub> are significant because many of them have extremely  
12 high GWPs and, in the cases of PFCs SF<sub>6</sub>, and NF<sub>3</sub>, long atmospheric lifetimes. Conversely, U.S. greenhouse gas  
13 emissions were partly offset by C sequestration in managed forests, trees in urban areas, agricultural soils, and  
14 landfilled yard trimmings. These were estimated to offset 13.1 percent of total emissions in 2013.

15 As the largest contributor to U.S. greenhouse gas emissions, carbon dioxide (CO<sub>2</sub>) from fossil fuel combustion has  
16 accounted for approximately 77 percent of global warming potential (GWP) weighted emissions for the entire time  
17 series since 1990, from 75 percent of total GWP-weighted emissions in 1990 to 77 percent in 2013. Emissions from  
18 this source category grew by 9.6 percent (454.8 MMT CO<sub>2</sub> Eq.) from 1990 to 2013 and were responsible for most of  
19 the increase in national emissions during this period. From 2012 to 2013, these emissions increased by 2.6 percent  
20 (133.2 MMT CO<sub>2</sub> Eq.). Historically, changes in emissions from fossil fuel combustion have been the dominant  
21 factor affecting U.S. emission trends.

22 Changes in CO<sub>2</sub> emissions from fossil fuel combustion are influenced by many long-term and short-term factors,  
23 including population and economic growth, energy price fluctuations, technological changes, and seasonal  
24 temperatures. On an annual basis, the overall consumption of fossil fuels in the United States fluctuates primarily in  
25 response to changes in general economic conditions, energy prices, weather, and the availability of non-fossil  
26 alternatives. For example, in a year with increased consumption of goods and services, low fuel prices, severe  
27 summer and winter weather conditions, nuclear plant closures, and lower precipitation feeding hydroelectric dams,  
28 there would likely be proportionally greater fossil fuel consumption than in a year with poor economic performance,  
29 high fuel prices, mild temperatures, and increased output from nuclear and hydroelectric plants.

1 In the longer-term, energy consumption patterns respond to changes that affect the scale of consumption (e.g.,  
2 population, number of cars, and size of houses), the efficiency with which energy is used in equipment (e.g., cars,  
3 power plants, steel mills, and light bulbs) and behavioral choices (e.g., walking, bicycling, or telecommuting to work  
4 instead of driving).

5 Energy-related CO<sub>2</sub> emissions also depend on the type of fuel or energy consumed and its carbon (C) intensity.  
6 Producing a unit of heat or electricity using natural gas instead of coal, for example, can reduce the CO<sub>2</sub> emissions  
7 because of the lower C content of natural gas.

8 A brief discussion of the year to year variability in fuel combustion emissions is provided below, beginning with  
9 2009.

10 From 2009 to 2010, CO<sub>2</sub> emissions from fossil fuel combustion increased by 3.3 percent, which represents one of  
11 the largest annual increases in CO<sub>2</sub> emissions from fossil fuel combustion for the twenty four-year period from 1990  
12 to 2013.<sup>48</sup> This increase is primarily due to an increase in economic output from 2009 to 2010, and increased  
13 industrial production and manufacturing output (FRB 2014). Carbon dioxide emissions from fossil fuel combustion  
14 in the industrial sector increased by 6.5 percent, including increased emissions from the combustion of fuel oil,  
15 natural gas and coal. Overall, coal consumption increased by 5.8 percent, the largest annual increase in coal  
16 consumption for the twenty four-year period between 1990 and 2013. In 2010, weather conditions remained fairly  
17 constant in the winter and were much hotter in the summer compared to 2009, as heating degree days decreased  
18 slightly by 0.4 percent and cooling degree days increased by 17.3 percent to their highest levels in the twenty one-  
19 year period from 1990 to 2010. As a result of the more energy-intensive summer weather conditions, electricity  
20 sales to the residential and commercial end-use sectors in 2010 increased approximately 6.0 percent and 1.8 percent,  
21 respectively.

22 From 2010 to 2011, CO<sub>2</sub> emissions from fossil fuel combustion decreased by 2.5 percent. This decrease is a result of  
23 multiple factors including: (1) a decrease in the carbon intensity of fuels consumed to generate electricity due to a  
24 decrease in coal consumption, with increased natural gas consumption and a significant increase in hydropower  
25 used; (2) a decrease in transportation-related energy consumption due to higher fuel costs, improvements in fuel  
26 efficiency, and a reduction in miles traveled; and (3) relatively mild winter conditions resulting in an overall  
27 decrease in energy demand in most sectors. Changing fuel prices played a role in the decreasing emissions. A  
28 significant increase in the price of motor gasoline in the transportation sector was a major factor leading to a  
29 decrease in energy consumption by 1.1 percent. In addition, an increase in the price of coal and a concurrent  
30 decrease in natural gas prices led to a 5.7 percent decrease and a 2.5 percent increase in fuel consumption of these  
31 fuels by electric generators. This change in fuel prices also reduced the carbon intensity of fuels used to produce  
32 electricity in 2011, further contributing to the decrease in fossil fuel combustion emissions.

33 From 2011 to 2012, CO<sub>2</sub> emissions from fossil fuel combustion decreased by 3.9 percent, with emissions from fossil  
34 fuel combustion at their lowest level since 1995. This decrease from 2011 to 2012 is primarily a result of the  
35 decrease in the carbon intensity of fuels used to generate electricity due to a slight increase in the price of coal, and a  
36 significant decrease in the price of natural gas. The consumption of coal used to generate electricity decreased by  
37 12.3 percent, while consumption of natural gas for electricity generation increased by 20.4 percent. Also, emissions  
38 declined in the transportation sector largely due to a small increase in fuel efficiency across different transportation  
39 modes and limited new demand for passenger transportation. In 2012, weather conditions remained fairly constant in  
40 the summer and were much warmer in the winter compared to 2011, as cooling degree days increased by 1.7 percent  
41 while heating degree days decreased 12.6 percent. This decrease in heating degree days resulted in a decreased  
42 demand for heating fuel in the residential and commercial sector, which had a decrease in natural gas consumption  
43 of 11.7 and 8.0 percent, respectively.  
44

45 From 2012 to 2013, CO<sub>2</sub> emissions from fossil fuel combustion increased by 2.6 percent, this increase is primarily a  
46 result of the increased energy consumption in the residential and commercial sectors, as heating degree days  
47 increased 18.6 percent in 2013 as compared to 2012. The cooler weather led to an increase of 16.9 and 12.6 percent  
48 direct use of fuels in the residential and commercial sectors, respectively. In addition, there was an increase of 1.2  
49 and 0.9 percent in electricity consumption in the residential and commercial sectors, respectively, due to regions that  
50 heat their homes with electricity. The consumption of natural gas used to generate electricity decreased by 10.2

---

<sup>48</sup> This increase also represents the largest absolute and percentage increase since 1988 (EIA 2011a).

1 percent due to an increase in the price of natural gas. Electric power plants shifted some consumption from natural  
 2 gas to coal, and as a result increased coal consumption to generate electricity by 4.2 percent. Lastly, industrial  
 3 production increased 2.9 percent from 2012 to 2013, resulting in an increase in the in CO<sub>2</sub> emissions from fossil fuel  
 4 combustion from the industrial sector by 4.3 percent.

5 Table 2-1 summarizes emissions and sinks from all U.S. anthropogenic sources in weighted units of MMT CO<sub>2</sub> Eq.,  
 6 while unweighted gas emissions and sinks in kilotons (kt) are provided in Table 2-2.

7 **Table 2-1: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (MMT CO<sub>2</sub> Eq.)**

Gas/Source	1990	2005	2009	2010	2011	2012	2013
<b>CO<sub>2</sub></b>	<b>5,126.8</b>	<b>6,156.4</b>	<b>5,553.0</b>	<b>5,754.2</b>	<b>5,618.7</b>	<b>5,418.7</b>	<b>5,556.0</b>
Fossil Fuel Combustion	4,740.7	5,753.5	5,226.1	5,401.2	5,266.6	5,062.3	5,195.5
Electricity Generation	1,820.8	2,402.1	2,146.4	2,259.2	2,158.5	2,022.7	2,040.5
Transportation	1,493.8	1,891.0	1,747.0	1,764.1	1,745.4	1,735.9	1,754.0
Industrial	842.5	828.8	728.3	775.6	773.7	783.9	817.3
Residential	338.3	357.9	336.7	335.1	327.5	283.4	329.9
Commercial	217.4	223.7	223.8	220.5	221.3	197.5	221.5
U.S. Territories	27.9	50.0	43.8	46.6	40.2	39.0	32.3
Non-Energy Use of Fuels	120.8	155.5	129.3	130.1	122.9	128.9	133.0
Iron and Steel Production & Metallurgical Coke Production	99.8	66.7	43.0	55.7	60.0	54.3	52.3
Natural Gas Systems	37.6	30.0	32.2	32.3	35.6	34.8	37.8
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1
Petrochemical Production	21.6	28.1	23.7	27.5	26.4	26.5	26.3
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2
Incineration of Waste	8.0	12.5	11.3	11.0	10.5	10.4	10.1
Cropland Remaining Cropland	7.1	7.9	7.2	8.6	8.0	10.0	9.9
Petroleum Systems	4.4	4.9	4.7	4.2	4.5	5.1	6.0
Urea Consumption for Non- Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7
Other Process Uses of Carbonates	4.9	6.3	7.6	9.6	9.3	8.0	4.4
Aluminum Production	6.8	4.1	3.0	2.7	3.3	3.4	3.3
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9
Wetlands Remaining Wetlands	1.0	1.1	1.0	1.0	0.9	0.8	0.8
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.1	0.2	0.2	0.2	0.2
Magnesium Production and Processing	+	+	+	+	+	+	+
<i>Land Use, Land-Use Change, and   Forestry<sup>a</sup> (Sink)<sup>a</sup></i>	<i>(774.1)</i>	<i>(912.6)</i>	<i>(871.3)</i>	<i>(872.0)</i>	<i>(881.3)</i>	<i>(880.7)</i>	<i>(882.0)</i>
<i>Wood Biomass and Ethanol   Consumption<sup>b</sup></i>	<i>219.4</i>	<i>229.8</i>	<i>250.5</i>	<i>265.1</i>	<i>268.1</i>	<i>267.7</i>	<i>283.3</i>
<i>International Bunker Fuels<sup>c</sup></i>	<i>103.5</i>	<i>113.1</i>	<i>106.4</i>	<i>117.0</i>	<i>111.7</i>	<i>105.8</i>	<i>99.8</i>
<b>CH<sub>4</sub></b>	<b>740.4</b>	<b>703.9</b>	<b>724.6</b>	<b>684.3</b>	<b>678.5</b>	<b>666.2</b>	<b>654.1</b>
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5
Natural Gas Systems	175.1	171.8	170.3	162.2	162.2	157.5	159.9
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6
Manure Management	37.2	56.3	59.7	60.9	61.4	63.7	61.4

Petroleum Systems	30.5	24.0	34.3	35.8	36.7	38.8	40.4
Wastewater Treatment	15.7	15.9	15.6	15.5	15.3	15.2	15.0
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3
Stationary Combustion	8.5	7.4	7.4	7.1	7.1	6.6	8.0
Abandoned Underground Coal							
Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2
Forest Land Remaining Forest Land	2.5	8.3	5.8	4.7	14.6	15.7	5.8
Mobile Combustion	5.6	3.0	2.3	2.3	2.3	2.2	2.1
Composting	0.4	1.9	1.9	1.8	1.9	1.9	2.0
Iron and Steel Production &							
Metallurgical Coke Production	1.1	0.9	0.4	0.6	0.7	0.7	0.7
Field Burning of Agricultural							
Residues	0.3	0.2	0.3	0.3	0.3	0.3	0.3
Petrochemical Production	0.2	0.1	0.1	0.1	+	0.1	0.1
Ferroalloy Production	+	+	+	+	+	+	+
Silicon Carbide Production and							
Consumption	+	+	+	+	+	+	+
Wetlands Remaining Wetlands	+	+	+	+	+	+	+
Incineration of Waste	+	+	+	+	+	+	+
<i>International Bunker Fuels<sup>c</sup></i>	0.2	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O</b>	<b>329.5</b>	<b>355.2</b>	<b>355.4</b>	<b>359.4</b>	<b>371.1</b>	<b>364.9</b>	<b>354.5</b>
Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7
Stationary Combustion	11.9	20.2	20.4	22.2	21.3	21.4	22.9
Mobile Combustion	41.2	38.1	24.6	23.7	22.5	20.2	18.4
Manure Management	13.8	16.4	17.0	17.1	17.3	17.3	17.3
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7
Wastewater Treatment	3.4	4.3	4.6	4.7	4.8	4.9	4.9
N <sub>2</sub> O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Forest Land Remaining Forest Land	1.7	5.8	4.2	3.5	10.0	10.7	4.2
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0
Settlements Remaining Settlements	1.0	1.8	1.7	1.8	1.9	1.9	1.8
Composting	0.3	1.7	1.7	1.6	1.7	1.7	1.8
Incineration of Waste	0.5	0.4	0.3	0.3	0.3	0.3	0.3
Semiconductor Manufacture	+	0.1	0.1	0.1	0.2	0.2	0.2
Field Burning of Agricultural							
Residues	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Wetlands Remaining Wetlands	+	+	+	+	+	+	+
<i>International Bunker Fuels<sup>c</sup></i>	0.9	1.0	0.9	1.0	1.0	0.9	0.9
<b>HFCs</b>	<b>46.6</b>	<b>131.4</b>	<b>142.9</b>	<b>152.6</b>	<b>157.4</b>	<b>159.2</b>	<b>164.3</b>
Substitution of Ozone Depleting							
Substances <sup>d</sup>	0.3	111.1	136.0	144.4	148.4	153.5	158.6
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	5.5
Semiconductor Manufacture	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Magnesium Production and							
Processing	0.0	0.0	+	+	+	+	0.1
<b>PFCs</b>	<b>24.3</b>	<b>6.6</b>	<b>3.9</b>	<b>4.4</b>	<b>6.9</b>	<b>6.0</b>	<b>5.8</b>
Aluminum Production	21.5	3.4	1.9	1.9	3.5	2.9	3.0
Semiconductor Manufacture	2.8	3.2	2.0	2.6	3.4	3.0	2.9
<b>SF<sub>6</sub></b>	<b>31.1</b>	<b>14.0</b>	<b>9.2</b>	<b>9.4</b>	<b>10.0</b>	<b>7.7</b>	<b>6.9</b>
Electrical Transmission and							
Distribution	25.4	10.6	7.3	6.9	6.8	5.7	5.1
Magnesium Production and							
Semiconductor Manufacture	5.2	2.7	1.6	2.1	2.8	1.6	1.4
<b>NF<sub>3</sub></b>	<b>+</b>	<b>0.5</b>	<b>0.4</b>	<b>0.5</b>	<b>0.7</b>	<b>0.6</b>	<b>0.6</b>
Semiconductor Manufacture	+	0.5	0.4	0.5	0.7	0.6	0.6
<b>Total</b>	<b>6,298.8</b>	<b>7,367.9</b>	<b>6,789.5</b>	<b>6,964.7</b>	<b>6,843.3</b>	<b>6,623.4</b>	<b>6,742.2</b>
<b>Net Emissions (Sources and Sinks)</b>	<b>5,524.7</b>	<b>6,455.4</b>	<b>5,918.2</b>	<b>6,092.7</b>	<b>5,962.0</b>	<b>5,742.7</b>	<b>5,860.2</b>

---

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> The net CO<sub>2</sub> flux total includes both emissions and sequestration, and constitutes a sink in the United States. Sinks are only included in net emissions total. Parentheses indicate negative values or sequestration.

<sup>b</sup> Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry.

<sup>c</sup> Emissions from International Bunker Fuels are not included in totals.

<sup>d</sup> Small amounts of PFC emissions also result from this source.

1 **Table 2-2: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks (kt)**

<b>Gas/Source</b>	<b>1990</b>	<b>2005</b>	<b>2009</b>	<b>2010</b>	<b>2011</b>	<b>2012</b>	<b>2013</b>
<b>CO<sub>2</sub></b>	<b>5,126,849</b>	<b>6,156,355</b>	<b>5,552,985</b>	<b>5,754,163</b>	<b>5,618,702</b>	<b>5,418,713</b>	<b>5,556,034</b>
Fossil Fuel Combustion	4,740,671	5,753,511	5,226,095	5,401,165	5,266,578	5,062,332	5,195,488
Electricity Generation	1,820,818	2,402,143	2,146,415	2,259,190	2,158,481	2,022,679	2,040,457
Transportation	1,493,758	1,891,048	1,747,047	1,764,146	1,745,384	1,735,860	1,753,991
Industrial	842,473	828,793	728,275	775,638	773,699	783,884	817,320
Residential	338,347	357,903	336,703	335,071	327,539	283,408	329,928
Commercial	217,393	223,664	223,837	220,533	221,319	197,519	221,533
U.S. Territories	27,882	49,960	43,818	46,586	40,157	38,982	32,259
Non-Energy Use of Fuels	120,842	155,455	129,338	130,067	122,868	128,949	133,040
Iron and Steel Production & Metallurgical Coke Production	99,781	66,666	43,029	55,746	60,008	54,327	52,288
Natural Gas Systems	37,622	29,982	32,201	32,334	35,551	34,764	37,806
Cement Production	33,278	45,910	29,432	31,256	32,010	35,051	36,146
Petrochemical Production	21,633	28,124	23,719	27,472	26,446	26,534	26,347
Lime Production	11,700	14,552	11,411	13,381	13,981	13,715	14,072
Ammonia Production	13,047	9,196	8,454	9,188	9,292	9,377	10,152
Incineration of Waste	7,972	12,454	11,295	11,026	10,550	10,363	10,137
Cropland Remaining Cropland	7,084	7,854	7,224	8,562	7,970	10,001	9,936
Petroleum Systems	4,442	4,902	4,675	4,171	4,486	5,079	6,020
Urea Consumption for Non-Agricultural Purposes	3,784	3,653	3,427	4,730	4,029	4,449	4,663
Other Process Uses of Carbonates	4,907	6,339	7,583	9,560	9,335	8,022	4,424
Aluminum Production	6,831	4,142	3,009	2,722	3,292	3,439	3,255
Soda Ash Production and Consumption	2,741	2,868	2,488	2,612	2,624	2,672	2,712
Ferroalloy Production	2,152	1,392	1,469	1,663	1,735	1,903	1,785
Titanium Dioxide Production	1,195	1,755	1,648	1,769	1,729	1,528	1,608
Zinc Production	632	1,030	943	1,182	1,286	1,486	1,429
Phosphoric Acid Production	1,586	1,395	1,016	1,130	1,198	1,138	1,173
Glass Production	1,535	1,928	1,045	1,481	1,299	1,248	1,160
Carbon Dioxide Consumption	1,472	1,375	1,795	1,206	802	841	903
Wetlands Remaining Wetlands	1,050	1,096	1,019	1,018	922	809	796
Lead Production	516	553	525	542	538	527	525
Silicon Carbide Production and Consumption	375	219	145	181	170	158	169
Magnesium Production and Processing	1	3	1	1	3	2	2
<i>Land Use, Land-Use Change, and Forestry (Sink)<sup>a</sup></i>	<i>(774,052)</i>	<i>(912,570)</i>	<i>(871,304)</i>	<i>(871,977)</i>	<i>(881,313)</i>	<i>(880,663)</i>	<i>(881,962)</i>
<i>Wood Biomass and Ethanol Consumption<sup>b</sup></i>	<i>219,413</i>	<i>229,844</i>	<i>250,491</i>	<i>265,110</i>	<i>268,064</i>	<i>267,730</i>	<i>283,337</i>
<i>International Bunker Fuels<sup>c</sup></i>	<i>103,463</i>	<i>113,139</i>	<i>106,410</i>	<i>116,992</i>	<i>111,660</i>	<i>105,805</i>	<i>99,763</i>
<b>CH<sub>4</sub></b>	<b>29,617</b>	<b>28,155</b>	<b>28,983</b>	<b>27,370</b>	<b>27,141</b>	<b>26,650</b>	<b>26,162</b>
Enteric Fermentation	6,566	6,755	6,908	6,844	6,750	6,653	6,581
Natural Gas Systems	7,003	6,873	6,814	6,488	6,487	6,299	6,396
Landfills	7,450	6,620	6,324	4,873	4,851	4,611	4,585
Coal Mining	3,860	2,565	3,194	3,293	2,849	2,658	2,584
Manure Management	1,486	2,254	2,388	2,437	2,457	2,548	2,456
Petroleum Systems	1,221	960	1,372	1,431	1,466	1,552	1,618
Wastewater Treatment	626	635	623	619	610	606	601
Rice Cultivation	366	358	378	444	339	372	332
Stationary Combustion	339	296	296	283	283	264	318
Abandoned Underground Coal Mines	288	264	254	263	257	249	249

Forest Land Remaining Forest Land	101	332	233	190	584	626	233
Mobile Combustion	225	121	93	92	91	88	86
Composting	15	75	75	73	75	77	79
Iron and Steel Production & Metallurgical Coke	46	34	17	25	28	29	28
Field Burning of Agricultural Residues	13	9	12	11	12	12	12
Petrochemical Production	9	6	2	2	2	3	3
Ferroalloy Production	1	+	+	+	+	1	+
Silicon Carbide Production and Consumption	1	+	+	+	+	+	+
Wetlands Remaining Wetlands	+	+	+	+	+	+	+
Incineration of Waste	+	+	+	+	+	+	+
<i>International Bunker Fuels<sup>c</sup></i>	7	5	5	6	5	4	3
<b>N<sub>2</sub>O</b>	<b>1,106</b>	<b>1,192</b>	<b>1,193</b>	<b>1,206</b>	<b>1,245</b>	<b>1,224</b>	<b>1,190</b>
Agricultural Soil Management	752	817	886	887	892	892	885
Stationary Combustion	40	68	69	74	71	72	77
Mobile Combustion	138	128	82	80	76	68	62
Manure Management	46	55	57	57	58	58	58
Nitric Acid Production	41	38	32	39	37	35	36
Wastewater Treatment	11	15	16	16	16	16	17
N <sub>2</sub> O from Product Uses	14	14	14	14	14	14	14
Forest Land Remaining Forest Land	6	20	14	12	33	36	14
Adipic Acid Production	51	24	9	14	34	19	13
Settlements Remaining	3	6	6	6	6	6	6
Composting	1	6	6	5	6	6	6
Incineration of Waste	2	1	1	1	1	1	1
Semiconductor Manufacture	+	+	+	+	1	1	1
Field Burning of Agricultural Residues	+	+	+	+	+	+	+
Wetlands Remaining Wetlands	+	+	+	+	+	+	+
<i>International Bunker Fuels<sup>c</sup></i>	3	3	3	3	3	3	3
<b>HFCs</b>	<b>M</b>						
Substitution of Ozone Depleting Substances <sup>d</sup>	M	M	M	M	M	M	M
HCFC-22 Production	3	1	+	1	1	+	+
Semiconductor Manufacture	+	+	+	+	+	+	+
Magnesium Production and Processing	0	0	+	+	+	+	+
<b>PFCs</b>	<b>M</b>						
Aluminum Production	M	M	M	M	M	M	M
Semiconductor Manufacture	M	M	M	M	M	M	M
<b>SF<sub>6</sub></b>	<b>1</b>	<b>+</b>	<b>+</b>	<b>+</b>	<b>+</b>	<b>+</b>	<b>+</b>
Electrical Transmission and Distribution	1	+	+	+	+	+	+
Magnesium Production and Processing	+	+	+	+	+	+	+
Semiconductor Manufacture	+	+	+	+	+	+	+
<b>NF<sub>3</sub></b>	<b>+</b>						
Semiconductor Manufacture	+	+	+	+	+	+	+

+ Does not exceed 0.5 kt.

M Mixture of multiple gases

<sup>a</sup> The net CO<sub>2</sub> flux total includes both emissions and sequestration, and constitutes a sink in the United States. Sinks are only included in net emissions total. Parentheses indicate negative values or sequestration.

<sup>b</sup> Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry

<sup>c</sup> Emissions from International Bunker Fuels are not included in totals.


<sup>d</sup> Small amounts of PFC emissions also result from this source.

Note: Totals may not sum due to independent rounding.

1 Emissions of all gases can be summed from each source category into a set of five sectors defined by the  
 2 Intergovernmental Panel on Climate Change (IPCC). Over the twenty four-year period of 1990 to 2013, total  
 3 emissions in the Energy, Industrial Processes and Product Use, and Agriculture sectors grew by 416.8 MMT CO<sub>2</sub>  
 4 Eq. (7.9 percent), 18.2 MMT CO<sub>2</sub> Eq. (5.3 percent), and 67.0 MMT CO<sub>2</sub> Eq. (14.9 percent), respectively. Emissions  
 5 from the Waste sector decreased by 67.7 MMT CO<sub>2</sub> Eq. (32.9 percent). Over the same period, estimates of net C  
 6 sequestration in the Land Use, Land-Use Change, and Forestry sector increased by 98.8 MMT CO<sub>2</sub> Eq. (13.0  
 7 percent).

8 **Figure 2-4: U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC Sector**

9 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



10

11 **Table 2-3: Recent Trends in U.S. Greenhouse Gas Emissions and Sinks by Chapter/IPCC**  
 12 **Sector (MMT CO<sub>2</sub> Eq.)**

Chapter/IPCC Sector	1990	2005	2009	2010	2011	2012	2013
<b>Energy</b>	<b>5,288.6</b>	<b>6,292.0</b>	<b>5,749.5</b>	<b>5,921.3</b>	<b>5,770.0</b>	<b>5,561.2</b>	<b>5,705.4</b>
Fossil Fuel Combustion	4,740.7	5,753.5	5,226.1	5,401.2	5,266.6	5,062.3	5,195.5
Natural Gas Systems	212.7	201.8	202.5	194.5	197.7	192.2	197.7
Non-Energy Use of Fuels	120.8	155.5	129.3	130.1	122.9	128.9	133.0
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6
Petroleum Systems	35.0	28.9	39.0	39.9	41.1	43.9	46.5
Stationary Combustion	20.4	27.6	27.8	29.3	28.4	28.0	30.8
Mobile Combustion	46.9	41.1	26.9	26.0	24.8	22.4	20.6
Incineration of Waste	8.4	12.8	11.6	11.4	10.9	10.7	10.4
Abandoned Underground Coal Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2
<b>Industrial Processes and Product Use</b>	<b>342.1</b>	<b>367.4</b>	<b>314.8</b>	<b>353.6</b>	<b>371.0</b>	<b>361.2</b>	<b>360.3</b>
Substitution of Ozone Depleting Substances	0.3	111.1	136.0	144.4	148.4	153.5	158.6
Iron and Steel Production & Metallurgical Coke Production	100.9	67.5	43.5	56.4	60.7	55.1	53.0
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1

Petrochemical Production	21.9	28.3	23.8	27.5	26.5	26.6	26.4
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2
Aluminum Production	28.3	7.6	4.9	4.6	6.8	6.4	6.2
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	5.5
Electrical Transmission and Distribution	25.4	10.6	7.3	6.9	6.8	5.7	5.1
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7
Other Process Uses of Carbonates	4.9	6.3	7.6	9.6	9.3	8.0	4.4
N <sub>2</sub> O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Semiconductor Manufacture	3.6	4.7	3.1	3.8	4.9	4.5	4.2
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.7	1.5
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2
<b>Agriculture</b>	<b>448.7</b>	<b>494.5</b>	<b>523.3</b>	<b>524.8</b>	<b>522.1</b>	<b>523.0</b>	<b>515.7</b>
Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5
Manure Management	51.0	72.8	76.7	78.0	78.7	81.0	78.7
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3
Field Burning of Agricultural Residues	0.4	0.3	0.4	0.4	0.4	0.4	0.4
<b>Land Use, Land-Use Change, and Forestry (Emissions)</b>	<b>13.4</b>	<b>24.8</b>	<b>20.0</b>	<b>19.6</b>	<b>35.3</b>	<b>39.0</b>	<b>22.6</b>
Forest Land Remaining Forest Land	4.3	14.1	10.0	8.2	24.6	26.3	10.0
Cropland Remaining Cropland	7.1	7.9	7.2	8.6	8.0	10.0	9.9
Settlements Remaining Settlements	1.0	1.8	1.7	1.8	1.9	1.9	1.8
Wetlands Remaining Wetlands	1.1	1.1	1.0	1.0	0.9	0.8	0.8
<b>Waste</b>	<b>206.0</b>	<b>189.2</b>	<b>181.8</b>	<b>145.5</b>	<b>144.9</b>	<b>138.9</b>	<b>138.3</b>
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6
Wastewater Treatment	19.0	20.2	20.2	20.2	20.1	20.0	20.0
Composting	0.7	3.5	3.6	3.5	3.5	3.7	3.7
<b>Total Emissions</b>	<b>6,298.8</b>	<b>7,367.9</b>	<b>6,789.5</b>	<b>6,964.7</b>	<b>6,843.3</b>	<b>6,623.4</b>	<b>6,742.2</b>
Net CO <sub>2</sub> Flux From Land Use, Land-Use Change and Forestry (Sinks) <sup>a</sup>	(774.1)	(912.6)	(871.3)	(872.0)	(881.3)	(880.7)	(882.0)
<b>Net Emission (Sources and Sinks)</b>	<b>5,524.7</b>	<b>6,455.4</b>	<b>5,918.2</b>	<b>6,092.7</b>	<b>5,962.0</b>	<b>5,742.7</b>	<b>5,860.2</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

<sup>a</sup> The net CO<sub>2</sub> flux total includes both emissions and sequestration, and constitutes a sink in the United States. Sinks are only included in net emissions total. Please refer to Table 2-8 for a breakout by source.


Note: Totals may not sum due to independent rounding. Parentheses indicate negative values or sequestration.

# 1 Energy

2 Energy-related activities, primarily fossil fuel combustion, accounted for the vast majority of U.S. CO<sub>2</sub> emissions for  
3 the period of 1990 through 2013. In 2013, approximately 82 percent of the energy consumed in the United States  
4 (on a Btu basis) was produced through the combustion of fossil fuels. The remaining 18 percent came from other  
5 energy sources such as hydropower, biomass, nuclear, wind, and solar energy (see Figure 2-5 and Figure 2-6). A  
6 discussion of specific trends related to CO<sub>2</sub> as well as other greenhouse gas emissions from energy consumption is  
7 presented in the Energy chapter. Energy-related activities are also responsible for CH<sub>4</sub> and N<sub>2</sub>O emissions (43  
8 percent and 12 percent of total U.S. emissions of each gas, respectively). Table 2-4 presents greenhouse gas  
9 emissions from the Energy chapter, by source and gas.


10 **Figure 2-5: 2013 Energy Chapter Greenhouse Gas Sources**

11 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



12

1 **Figure 2-6: 2013 U.S. Fossil Carbon Flows (MMT CO<sub>2</sub> Eq.)**  
 2 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.  
 3



4  
 5 **Table 2-4: Emissions from Energy (MMT CO<sub>2</sub> Eq.)**

Gas/Source	1990	2005	2009	2010	2011	2012	2013
<b>CO<sub>2</sub></b>	<b>4,911.5</b>	<b>5,956.3</b>	<b>5,403.6</b>	<b>5,578.8</b>	<b>5,440.0</b>	<b>5,241.5</b>	<b>5,382.5</b>
Fossil Fuel Combustion	4,740.7	5,753.5	5,226.1	5,401.2	5,266.6	5,062.3	5,195.5
Electricity Generation	1,820.8	2,402.1	2,146.4	2,259.2	2,158.5	2,022.7	2,040.5
Transportation	1,493.8	1,891.0	1,747.0	1,764.1	1,745.4	1,735.9	1,754.0
Industrial	842.5	828.8	728.3	775.6	773.7	783.9	817.3
Residential	338.3	357.9	336.7	335.1	327.5	283.4	329.9
Commercial	217.4	223.7	223.8	220.5	221.3	197.5	221.5
U.S. Territories	27.9	50.0	43.8	46.6	40.2	39.0	32.3
Non-Energy Use of Fuels	120.8	155.5	129.3	130.1	122.9	128.9	133.0
Natural Gas Systems	37.6	30.0	32.2	32.3	35.6	34.8	37.8
Incineration of Waste	8.0	12.5	11.3	11.0	10.5	10.4	10.1
Petroleum Systems	4.4	4.9	4.7	4.2	4.5	5.1	6.0
Biomass - Wood <sup>a</sup>	215.2	206.9	188.2	192.5	195.2	194.9	208.6
International Bunker Fuels <sup>b</sup>	103.5	113.1	106.4	117.0	111.7	105.8	99.8
Biomass - Ethanol <sup>a</sup>	4.2	22.9	62.3	72.6	72.9	72.8	74.7
<b>CH<sub>4</sub></b>	<b>323.4</b>	<b>277.0</b>	<b>300.6</b>	<b>296.3</b>	<b>285.8</b>	<b>277.8</b>	<b>281.3</b>
Natural Gas Systems	175.1	171.8	170.3	162.2	162.2	157.5	159.9
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6
Petroleum Systems	30.5	24.0	34.3	35.8	36.7	38.8	40.4
Stationary Combustion	8.5	7.4	7.4	7.1	7.1	6.6	8.0
Abandoned Underground Coal	7.2	6.6	6.4	6.6	6.4	6.2	6.2
Mobile Combustion	5.6	3.0	2.3	2.3	2.3	2.2	2.1
Incineration of Waste	+	+	+	+	+	+	+
International Bunker Fuels <sup>b</sup>	0.2	0.1	0.1	0.1	0.1	0.1	0.1
<b>N<sub>2</sub>O</b>	<b>53.6</b>	<b>58.7</b>	<b>45.4</b>	<b>46.2</b>	<b>44.1</b>	<b>41.9</b>	<b>41.6</b>
Stationary Combustion	11.9	20.2	20.4	22.2	21.3	21.4	22.9

Mobile Combustion	41.2	38.1	24.6	23.7	22.5	20.2	18.4
Incineration of Waste	0.5	0.4	0.3	0.3	0.3	0.3	0.3
<i>International Bunker Fuels<sup>b</sup></i>	<i>0.9</i>	<i>1.0</i>	<i>0.9</i>	<i>1.0</i>	<i>1.0</i>	<i>0.9</i>	<i>0.9</i>
<b>Total</b>	<b>5,288.6</b>	<b>6,292.0</b>	<b>5,749.5</b>	<b>5,921.3</b>	<b>5,770.0</b>	<b>5,561.2</b>	<b>5,705.4</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> Emissions from Wood Biomass and Ethanol Consumption are not included specifically in summing energy sector totals. Net carbon fluxes from changes in biogenic carbon reservoirs are accounted for in the estimates for Land Use, Land-Use Change, and Forestry.

<sup>b</sup> Emissions from International Bunker Fuels are not included in totals.

1  
2 Carbon dioxide emissions from fossil fuel combustion are presented in Table 2-5 based on the underlying U.S.  
3 energy consumer data collected by EIA. Estimates of CO<sub>2</sub> emissions from fossil fuel combustion are calculated from  
4 these EIA “end-use sectors” based on total consumption and appropriate fuel properties (any additional analysis and  
5 refinement of the EIA data is further explained in the Energy chapter of this report). EIA’s fuel consumption data for  
6 the electric power sector comprises electricity-only and combined-heat-and-power (CHP) plants within the NAICS  
7 22 category whose primary business is to sell electricity, or electricity and heat, to the public (nonutility power  
8 producers can be included in this sector as long as they meet they electric power sector definition). EIA statistics for  
9 the industrial sector include fossil fuel consumption that occurs in the fields of manufacturing, agriculture, mining,  
10 and construction. EIA’s fuel consumption data for the transportation sector consists of all vehicles whose primary  
11 purpose is transporting people and/or goods from one physical location to another. EIA’s fuel consumption data for  
12 the industrial sector consists of all facilities and equipment used for producing, processing, or assembling goods  
13 (EIA includes generators that produce electricity and/or useful thermal output primarily to support on-site industrial  
14 activities in this sector). EIA’s fuel consumption data for the residential sector consists of living quarters for private  
15 households. EIA’s fuel consumption data for the commercial sector consists of service-providing facilities and  
16 equipment from private and public organizations and businesses (EIA includes generators that produce electricity  
17 and/or useful thermal output primarily to support the activities at commercial establishments in this sector). Table  
18 2-5 and Figure 2-7 summarize CO<sub>2</sub> emissions from fossil fuel combustion by end-use sector. Figure 2-8 further  
19 describes the total emissions from fossil fuel combustion, separated by end-use sector, including CH<sub>4</sub> and N<sub>2</sub>O in  
20 addition to CO<sub>2</sub>.

21 **Table 2-5: CO<sub>2</sub> Emissions from Fossil Fuel Combustion by End-Use Sector (MMT CO<sub>2</sub> Eq.)**

End-Use Sector	1990	2005	2009	2010	2011	2012	2013
<b>Transportation</b>	<b>1,496.8</b>	<b>1,895.8</b>	<b>1,751.5</b>	<b>1,768.6</b>	<b>1,749.6</b>	<b>1,739.7</b>	<b>1,758.0</b>
Combustion	1,493.8	1,891.0	1,747.0	1,764.1	1,745.4	1,735.9	1,754.0
Electricity	3.0	4.7	4.5	4.5	4.3	3.9	4.0
<b>Industrial</b>	<b>1,529.2</b>	<b>1,565.7</b>	<b>1,330.3</b>	<b>1,416.7</b>	<b>1,398.6</b>	<b>1,376.8</b>	<b>1,400.0</b>
Combustion	842.5	828.8	728.3	775.6	773.7	783.9	817.3
Electricity	686.7	737.0	602.0	641.1	624.9	592.9	582.7
<b>Residential</b>	<b>931.4</b>	<b>1,214.7</b>	<b>1,123.2</b>	<b>1,175.5</b>	<b>1,118.5</b>	<b>1,008.9</b>	<b>1,070.8</b>
Combustion	338.3	357.9	336.7	335.1	327.5	283.4	329.9
Electricity	593.0	856.7	786.5	840.4	791.0	725.5	740.9
<b>Commercial</b>	<b>755.4</b>	<b>1,027.4</b>	<b>977.3</b>	<b>993.8</b>	<b>959.6</b>	<b>898.0</b>	<b>934.4</b>
Combustion	217.4	223.7	223.8	220.5	221.3	197.5	221.5
Electricity	538.0	803.7	753.5	773.3	738.3	700.4	712.8
<b>U.S. Territories<sup>a</sup></b>	<b>27.9</b>	<b>50.0</b>	<b>43.8</b>	<b>46.6</b>	<b>40.2</b>	<b>39.0</b>	<b>32.3</b>
<b>Total</b>	<b>4,740.7</b>	<b>5,753.5</b>	<b>5,226.1</b>	<b>5,401.2</b>	<b>5,266.6</b>	<b>5,062.3</b>	<b>5,195.5</b>
<b>Electricity Generation</b>	<b>1,820.8</b>	<b>2,402.1</b>	<b>2,146.4</b>	<b>2,259.2</b>	<b>2,158.5</b>	<b>2,022.7</b>	<b>2,040.5</b>


Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding. Combustion-related emissions from electricity generation are allocated based on aggregate national electricity consumption by each end-use sector.

<sup>a</sup> Fuel consumption by U.S. Territories (i.e., American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands) is included in this report.

1 **Figure 2-7: 2013 CO<sub>2</sub> Emissions from Fossil Fuel Combustion by Sector and Fuel Type**


2 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



3

4 **Figure 2-8: 2013 End-Use Sector Emissions of CO<sub>2</sub> from Fossil Fuel Combustion**

5 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



6

7 The main driver of emissions in the Energy sector is CO<sub>2</sub> from fossil fuel combustion. Electricity generation is the  
 8 largest emitter of CO<sub>2</sub>, and electricity generators consumed 34 percent of U.S. energy from fossil fuels and emitted  
 9 39 percent of the CO<sub>2</sub> from fossil fuel combustion in 2013. Electricity generation emissions can also be allocated to  
 10 the end-use sectors that are consuming that electricity, as presented in Table 2-5. The transportation end-use sector  
 11 accounted for 1,758.0 MMT CO<sub>2</sub> Eq. in 2013 or approximately 34 percent of total CO<sub>2</sub> emissions from fossil fuel  
 12 combustion. The industrial end-use sector accounted for 27 percent of CO<sub>2</sub> emissions from fossil fuel combustion.  
 13 The residential and commercial end-use sectors accounted for 21 and 18 percent, respectively, of CO<sub>2</sub> emissions  
 14 from fossil fuel combustion. Both of these end-use sectors were heavily reliant on electricity for meeting energy  
 15 needs, with electricity consumption for lighting, heating, air conditioning, and operating appliances contributing 69  
 16 and 76 percent of emissions from the residential and commercial end-use sectors, respectively. Significant trends in  
 17 emissions from energy source categories over the twenty four-year period from 1990 through 2013 included the  
 18 following:

- 19 • Total CO<sub>2</sub> emissions from fossil fuel combustion increased from 4,740.7 MMT CO<sub>2</sub> Eq. in 1990 to 5,195.5  
 20 MMT CO<sub>2</sub> Eq. in 2013—a 9.6 percent total increase over the twenty four-year period. From 2012 to 2013,  
 21 these emissions increased by 133.2 MMT CO<sub>2</sub> Eq. (2.6 percent).

- 1 • CH<sub>4</sub> emissions from natural gas systems were the second largest anthropogenic source of CH<sub>4</sub> emissions in  
2 the United States with 159.9 MMT CO<sub>2</sub> Eq. emitted into the atmosphere in 2013; emissions have decreased  
3 by 15.2 MMT CO<sub>2</sub> Eq. (8.7 percent) since 1990.
- 4 • CO<sub>2</sub> emissions from non-energy use of fossil fuels increased by 12.2 MMT CO<sub>2</sub> Eq. (10.1 percent) from  
5 1990 through 2013. Emissions from non-energy uses of fossil fuels were 133.0 MMT CO<sub>2</sub> Eq. in 2013,  
6 which constituted 2.4 percent of total national CO<sub>2</sub> emissions.
- 7 • N<sub>2</sub>O emissions from stationary combustion increased by 11.0 MMT CO<sub>2</sub> Eq. (92.0 percent) from 1990  
8 through 2013. N<sub>2</sub>O emissions from this source increased primarily as a result of an increase in the number  
9 of coal fluidized bed boilers in the electric power sector.
- 10 • CO<sub>2</sub> emissions from incineration of waste (10.1 MMT CO<sub>2</sub> Eq. in 2013) increased by 2.2 MMT CO<sub>2</sub> Eq.  
11 (27.2 percent) from 1990 through 2013, as the volume of plastics and other fossil carbon-containing  
12 materials in municipal solid waste grew.

13 The increase in CO<sub>2</sub> emissions from fossil fuel combustion in 2013 was a result of multiple factors including: (1) the  
14 increase in the price of natural gas led to an increase of coal-fired generation in the electric power sector; (2) much  
15 colder winter conditions resulted in an increased demand for heating fuel in the residential and commercial sectors;  
16 (3) an increase in industrial production across multiple sectors which resulted in increases in industrial sector  
17 emissions,<sup>49</sup> and (4) an increase in transportation emissions resulting from a small increase in vehicle miles traveled  
18 (VMT) and fuel use across on-road transportation modes.

## 19 Industrial Processes and Product Use

20 The Industrial Processes and Product Use (IPPU) section includes greenhouse gas emissions occurring from  
21 industrial processes and from the use of greenhouse gases in products. This section includes sources of emissions  
22 formerly represented in the ‘Industrial Processes’ and ‘Solvent and Other Product Use’ sectors in prior versions of  
23 this report.

24 Greenhouse gas emissions are produced as the by-products of many non-energy-related industrial activities. For  
25 example, industrial processes can chemically transform raw materials, which often release waste gases such as CO<sub>2</sub>,  
26 CH<sub>4</sub>, and N<sub>2</sub>O. These processes include iron and steel production and metallurgical coke production, cement  
27 production, ammonia production, urea consumption, lime production, other process uses of carbonates (e.g., flux  
28 stone, flue gas desulfurization, and glass manufacturing), soda ash production and consumption, titanium dioxide  
29 production, phosphoric acid production, ferroalloy production, CO<sub>2</sub> consumption, silicon carbide production and  
30 consumption, aluminum production, petrochemical production, nitric acid production, adipic acid production, lead  
31 production, zinc production, and N<sub>2</sub>O from product uses (see Figure 2-9). Industrial processes also release HFCs,  
32 PFCs, SF<sub>6</sub>, and NF<sub>3</sub>. In addition to their use as ODS substitutes, HFCs, PFCs, SF<sub>6</sub>, NF<sub>3</sub>, and other fluorinated  
33 compounds are employed and emitted by a number of other industrial sources in the United States. These industries  
34 include aluminum production, HCFC-22 production, semiconductor manufacture, electric power transmission and  
35 distribution, and magnesium metal production and processing. Table 2-6 presents greenhouse gas emissions from  
36 industrial processes by source category.


37

---

<sup>49</sup> Further details on industrial sector combustion emissions are provided by EPA’s GHGRP  
(<http://ghgdata.epa.gov/ghgp/main.do>).

1 **Figure 2-9: 2013 Industrial Processes and Product Use Chapter Greenhouse Gas Sources**

2 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



3

4 **Table 2-6: Emissions from Industrial Processes and Product Use (MMT CO<sub>2</sub> Eq.)**

Gas/Source	1990	2005	2009	2010	2011	2012	2013
<b>CO<sub>2</sub></b>	<b>207.2</b>	<b>191.1</b>	<b>141.1</b>	<b>165.8</b>	<b>169.8</b>	<b>166.4</b>	<b>162.8</b>
Iron and Steel Production & Metallurgical Coke Production	99.8	66.7	43.0	55.7	60.0	54.3	52.3
<i>Iron and Steel Production</i>	97.3	64.6	42.1	53.7	58.6	53.8	50.5
<i>Metallurgical Coke Production</i>	2.5	2.0	1.0	2.1	1.4	0.5	1.8
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1
Petrochemical Production	21.6	28.1	23.7	27.5	26.4	26.5	26.3
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7
Other Process Uses of Carbonates	4.9	6.3	7.6	9.6	9.3	8.0	4.4
Aluminum Production	6.8	4.1	3.0	2.7	3.3	3.4	3.3
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2

Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5
Silicon Carbide Production and Consumption	0.4	0.2	0.1	0.2	0.2	0.2	0.2
Magnesium Production and Processing	+	+	+	+	+	+	+
<b>CH<sub>4</sub></b>	<b>1.4</b>	<b>1.0</b>	<b>0.5</b>	<b>0.7</b>	<b>0.8</b>	<b>0.8</b>	<b>0.8</b>
Iron and Steel Production & Metallurgical Coke	1.1	0.9	0.4	0.6	0.7	0.7	0.7
<i>Iron and Steel Production</i>	1.1	0.9	0.4	0.6	0.7	0.7	0.7
<i>Metallurgical Coke Production</i>	+	+	+	+	+	+	+
Petrochemical Production	0.2	0.1	0.1	0.1	+	0.1	0.1
Ferroalloy Production	+	+	+	+	+	+	+
Silicon Carbide Production and Consumption	+	+	+	+	+	+	+
<b>N<sub>2</sub>O</b>	<b>31.6</b>	<b>22.8</b>	<b>16.7</b>	<b>20.1</b>	<b>25.5</b>	<b>20.4</b>	<b>19.1</b>
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7
N <sub>2</sub> O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0
Semiconductor Manufacturing	+	0.1	0.1	0.1	0.2	0.2	0.2
<b>HFCs</b>	<b>46.6</b>	<b>131.4</b>	<b>142.9</b>	<b>152.6</b>	<b>157.4</b>	<b>159.2</b>	<b>164.3</b>
Substitution of Ozone Depleting Substances <sup>a</sup>	0.3	111.1	136.0	144.4	148.4	153.5	158.6
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	5.5
Semiconductor Manufacturing	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Magnesium Production and Processing	+	+	+	+	+	+	0.1
<b>PFCs</b>	<b>24.3</b>	<b>6.6</b>	<b>3.9</b>	<b>4.4</b>	<b>6.9</b>	<b>6.0</b>	<b>5.8</b>
Aluminum Production	21.5	3.4	1.9	1.9	3.5	2.9	3.0
Semiconductor Manufacturing	2.8	3.2	2.0	2.6	3.4	3.0	2.9
<b>SF<sub>6</sub></b>	<b>31.1</b>	<b>14.0</b>	<b>9.2</b>	<b>9.4</b>	<b>10.0</b>	<b>7.7</b>	<b>6.9</b>
Electrical Transmission and Distribution	25.4	10.6	7.3	6.9	6.8	5.7	5.1
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.6	1.4
Semiconductor Manufacturing	0.5	0.7	0.3	0.4	0.4	0.4	0.4
<b>NF<sub>3</sub></b>	<b>+</b>	<b>0.5</b>	<b>0.4</b>	<b>0.5</b>	<b>0.7</b>	<b>0.6</b>	<b>0.6</b>
Semiconductor Manufacturing	+	0.5	0.4	0.5	0.7	0.6	0.6
<b>Total</b>	<b>342.1</b>	<b>367.4</b>	<b>314.8</b>	<b>353.6</b>	<b>371.0</b>	<b>361.2</b>	<b>360.3</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> Small amounts of PFC emissions also result from this source.

Note: Totals may not sum due to independent rounding.

- 1 Overall, emissions from the IPPU sector increased by 5.3 percent from 1990 to 2013. Significant trends in emissions  
2 from IPPU source categories over the twenty four-year period from 1990 through 2013 included the following:
- 3 • HFC emissions from ODS substitutes have been increasing from small amounts in 1990 to 158.6 MMT  
4 CO<sub>2</sub> Eq. in 2013. This increase was in large part the result of efforts to phase out CFCs and other ODSs in  
5 the United States. In the short term, this trend is expected to continue, and will likely continue over the  
6 next decade as HCFCs, which are interim substitutes in many applications, are themselves phased-out  
7 under the provisions of the Copenhagen Amendments to the *Montreal Protocol*.
  - 8 • Combined CO<sub>2</sub> and CH<sub>4</sub> emissions from iron and steel production and metallurgical coke production  
9 decreased by 3.8 percent to 53.0 MMT CO<sub>2</sub> Eq. from 2012 to 2013, and have declined overall by 47.9  
10 MMT CO<sub>2</sub> Eq. (47.5 percent) from 1990 through 2013, due to restructuring of the industry, technological  
11 improvements, and increased scrap steel utilization.
  - 12 • CO<sub>2</sub> emissions from ammonia production (10.2 MMT CO<sub>2</sub> Eq. in 2013) decreased by 2.9 MMT CO<sub>2</sub> Eq.  
13 (22.2 percent) since 1990. Ammonia production relies on natural gas as both a feedstock and a fuel, and as  
14 such, market fluctuations and volatility in natural gas prices affect the production of ammonia.
  - 15 • Urea consumption for non-agricultural purposes (4.7 MMT CO<sub>2</sub> Eq. in 2013) increased by 0.9 MMT CO<sub>2</sub>  
16 Eq. (23.2 percent) since 1990.

- 1 • In 2013, N<sub>2</sub>O emissions from product uses constituted 1.2 percent of U.S. N<sub>2</sub>O emissions. From 1990 to  
2 2013, emissions from this source category decreased by 0.4 percent, though slight increases occurred in  
3 intermediate years.
- 4 • N<sub>2</sub>O emissions from adipic acid production were 4.0 MMT CO<sub>2</sub> Eq. in 2013, and have decreased  
5 significantly since 1990 due to both the widespread installation of pollution control measures in the late  
6 1990s and plant idling in the late 2000s. Emissions from adipic acid production have decreased by 73.8  
7 percent since 1990 and by 76.4 percent since a peak in 1995.
- 8 • PFC emissions from aluminum production decreased by 86.2 percent (18.5 MMT CO<sub>2</sub> Eq.) from 1990 to  
9 2013, due to both industry emission reduction efforts and lower domestic aluminum production.


## 10 Agriculture

11 Agricultural activities contribute directly to emissions of greenhouse gases through a variety of processes, including  
12 the following source categories: enteric fermentation in domestic livestock, livestock manure management, rice  
13 cultivation, agricultural soil management, and field burning of agricultural residues.

14 In 2013, agricultural activities were responsible for emissions of 515.7 MMT CO<sub>2</sub> Eq., or 7.6 percent of total U.S.  
15 greenhouse gas emissions. CH<sub>4</sub> and N<sub>2</sub>O were the primary greenhouse gases emitted by agricultural activities. CH<sub>4</sub>  
16 emissions from enteric fermentation and manure management represented about 25.2 percent and 9.4 percent of total  
17 CH<sub>4</sub> emissions from anthropogenic activities, respectively, in 2013. Agricultural soil management activities, such as  
18 fertilizer use and other cropping practices, were the largest source of U.S. N<sub>2</sub>O emissions in 2013, accounting for  
19 74.4 percent.

20 **Figure 2-10: 2013 Agriculture Chapter Greenhouse Gas Sources**

21 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



22

23 **Table 2-7: Emissions from Agriculture (MMT CO<sub>2</sub> Eq.)**

Gas/Source	1990	2005	2009	2010	2011	2012	2013
<b>CH<sub>4</sub></b>	<b>210.8</b>	<b>234.4</b>	<b>242.1</b>	<b>243.4</b>	<b>238.9</b>	<b>239.6</b>	<b>234.5</b>
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5
Manure Management	37.2	56.3	59.7	60.9	61.4	63.7	61.4
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3
Field Burning of Agricultural Residues	0.3	0.2	0.3	0.3	0.3	0.3	0.3
<b>N<sub>2</sub>O</b>	<b>237.9</b>	<b>260.1</b>	<b>281.2</b>	<b>281.4</b>	<b>283.2</b>	<b>283.4</b>	<b>281.1</b>
Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7
Manure Management	13.8	16.4	17.0	17.1	17.3	17.3	17.3

Field Burning of Agricultural Residues	0.1	0.1	0.1	0.1	0.1	0.1	0.1
<b>Total</b>	<b>448.7</b>	<b>494.5</b>	<b>523.3</b>	<b>524.8</b>	<b>522.1</b>	<b>523.0</b>	<b>515.7</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

1 Some significant trends in U.S. emissions from Agriculture source categories include the following:

- 2 • Agricultural soils produced approximately 74.4 percent of N<sub>2</sub>O emissions in the United States in 2013. Estimated emissions from this source in 2013 were 263.7 MMT CO<sub>2</sub> Eq. Annual N<sub>2</sub>O emissions from agricultural soils fluctuated between 1990 and 2013, although overall emissions were 17.7 percent higher in 2013 than in 1990. Year-to-year fluctuations are largely a reflection of annual variation in weather patterns, synthetic fertilizer use, and crop production.
- 7 • Enteric fermentation is the largest anthropogenic source of CH<sub>4</sub> emissions in the United States. In 2013, enteric fermentation CH<sub>4</sub> emissions were 164.5 MMT CO<sub>2</sub> Eq. (25.2 percent of total CH<sub>4</sub> emissions), which represents an increase of 0.4 MMT CO<sub>2</sub> Eq. (0.2 percent) since 1990. This increase in emissions from 1990 to 2013 in enteric generally follows the increasing trends in cattle populations. From 1990 to 1995 emissions increased and then decreased from 1996 to 2004, mainly due to fluctuations in beef cattle populations and increased digestibility of feed for feedlot cattle. Emissions generally increased from 2005 to 2007, as both dairy and beef populations underwent increases and the literature for dairy cow diets indicated a trend toward a decrease in feed digestibility for those years. Emissions decreased again from 2008 to 2013 as beef cattle populations again decreased.
- 16 • Overall, emissions from manure management increased 54.4 percent between 1990 and 2013. This encompassed an increase of 65.2 percent for CH<sub>4</sub>, from 37.2 MMT CO<sub>2</sub> Eq. in 1990 to 61.4 MMT CO<sub>2</sub> Eq. in 2013; and an increase of 25.4 percent for N<sub>2</sub>O, from 13.8 MMT CO<sub>2</sub> Eq. in 1990 to 17.3 MMT CO<sub>2</sub> Eq. in 2013. The majority of the increase observed in CH<sub>4</sub> resulted from swine and dairy cow manure, where emissions increased 48 and 115 percent, respectively, from 1990 to 2013. From 2012 to 2013, there was a 3.6 percent decrease in total CH<sub>4</sub> emissions from manure management, mainly due to minor shifts in the animal populations and the resultant effects on manure management system allocations.

## 23 Land Use, Land-Use Change, and Forestry

24 When humans alter the terrestrial biosphere through land use, changes in land use, and land management practices, they also alter the background carbon fluxes between biomass, soils, and the atmosphere. Forest management practices, tree planting in urban areas, the management of agricultural soils, and the landfilling of yard trimmings and food scraps have resulted in an uptake (sequestration) of carbon in the United States, which offset about 13.1 percent of total U.S. greenhouse gas emissions in 2013. Forests (including vegetation, soils, and harvested wood) accounted for approximately 88 percent of total 2013 net CO<sub>2</sub> flux, urban trees accounted for 10 percent, mineral and organic soil carbon stock changes accounted for less than 0.5 percent, and landfilled yard trimmings and food scraps accounted for 1.5 percent of the total net flux in 2013. The net forest sequestration is a result of net forest growth, increasing forest area, and a net accumulation of carbon stocks in harvested wood pools. The net sequestration in urban forests is a result of net tree growth and increased urban forest size. In agricultural soils, mineral and organic soils sequester approximately 2.4 times as much C as is emitted from these soils through liming and urea fertilization. The mineral soil C sequestration is largely due to the conversion of cropland to hay production fields, the limited use of bare-summer fallow areas in semi-arid areas, and an increase in the adoption of conservation tillage practices. The landfilled yard trimmings and food scraps net sequestration is due to the long-term accumulation of yard trimming and food scraps carbon in landfills.

39 Land use, land-use change, and forestry activities in 2013 resulted in a net C sequestration of 882.0 MMT CO<sub>2</sub> Eq. (240.5 MMT C) (Table 2-8). This represents an offset of approximately 15.9 percent of total U.S. CO<sub>2</sub> emissions, or 13.1 percent of total greenhouse gas emissions in 2013. Between 1990 and 2013, total land use, land-use change, and forestry net C flux resulted in a 13.9 percent increase in CO<sub>2</sub> sequestration, primarily due to an increase in the rate of net C accumulation in forest C stocks, particularly in aboveground and belowground tree biomass, and harvested wood pools.

1 **Table 2-8: Net CO<sub>2</sub> Flux from Land Use, Land-Use Change, and Forestry (MMT CO<sub>2</sub> Eq.)**

Sink Category	1990	2005	2009	2010	2011	2012	2013
Forest Land Remaining Forest Land	(639.4)	(807.1)	(764.9)	(765.4)	(773.8)	(773.1)	(775.7)
Cropland Remaining Cropland	(65.2)	(28.0)	(27.5)	(25.9)	(25.8)	(25.0)	(23.4)
Land Converted to Cropland	24.5	19.8	16.2	16.2	16.2	16.1	16.1
Grassland Remaining Grassland	(1.9)	4.2	11.7	11.7	11.7	11.5	12.1
Land Converted to Grassland	(7.4)	(9.0)	(8.9)	(8.9)	(8.9)	(8.8)	(8.8)
Settlements Remaining Settlements	(60.4)	(80.5)	(85.0)	(86.1)	(87.3)	(88.4)	(89.5)
Other (Landfilled Yard Trimmings and Food Scraps)	(24.2)	(12.0)	(12.9)	(13.6)	(13.5)	(13.0)	(12.8)
<b>Total</b>	<b>(774.1)</b>	<b>(912.6)</b>	<b>(871.3)</b>	<b>(872.0)</b>	<b>(881.3)</b>	<b>(880.7)</b>	<b>(882.0)</b>

Note: Totals may not sum due to independent rounding. Parentheses indicate net sequestration.

2 Land use, land-use change, and forestry source categories also resulted in emissions of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O that are  
3 not included in the net CO<sub>2</sub> flux estimates presented in Table 2-8. Liming of agricultural soils and urea fertilization  
4 resulted in CO<sub>2</sub> emissions of 9.9 MMT CO<sub>2</sub> Eq. in 2013, an increase of about 40.3 percent relative to 1990. Lands  
5 undergoing peat extraction resulted in CO<sub>2</sub> emissions of 0.8 MMT CO<sub>2</sub> Eq. and CH<sub>4</sub> and N<sub>2</sub>O emissions of less than  
6 0.05 MMT CO<sub>2</sub> Eq. each. N<sub>2</sub>O emissions from the application of synthetic fertilizers to forest soils have increased  
7 from 0.1 MMT CO<sub>2</sub> Eq. in 1990 to 0.3 MMT CO<sub>2</sub> Eq. in 2013. Settlement soils in 2013 resulted in direct N<sub>2</sub>O  
8 emissions of 1.8 MMT CO<sub>2</sub> Eq., a 78.3 percent increase relative to 1990. Emissions from forest fires in 2013  
9 resulted in CH<sub>4</sub> emissions of 5.8 MMT CO<sub>2</sub> Eq., and in N<sub>2</sub>O emissions of 3.8 MMT CO<sub>2</sub> Eq. (Table 2-9).

10 **Table 2-9: Emissions from Land Use, Land-Use Change, and Forestry (MMT CO<sub>2</sub> Eq.)**

Source Category	1990	2005	2009	2010	2011	2012	2013
<b>CO<sub>2</sub></b>	<b>8.1</b>	<b>9.0</b>	<b>8.2</b>	<b>9.6</b>	<b>8.9</b>	<b>10.8</b>	<b>10.7</b>
Cropland Remaining Cropland: Liming of Agricultural Soils	4.7	4.3	3.7	4.8	3.9	5.8	5.9
Cropland Remaining Cropland: Urea Fertilization	2.4	3.5	3.6	3.8	4.1	4.2	4.0
Wetlands Remaining Wetlands: Peatlands Remaining Peatlands	1.0	1.1	1.0	1.0	0.9	0.8	0.8
<b>CH<sub>4</sub></b>	<b>2.5</b>	<b>8.3</b>	<b>5.8</b>	<b>4.8</b>	<b>14.6</b>	<b>15.7</b>	<b>5.8</b>
Forest Land Remaining Forest Land: Forest Fires	2.5	8.3	5.8	4.7	14.6	15.7	5.8
Wetlands Remaining Wetlands: Peatlands Remaining Peatlands	+	+	+	+	+	+	+
<b>N<sub>2</sub>O</b>	<b>2.7</b>	<b>7.6</b>	<b>5.9</b>	<b>5.3</b>	<b>11.8</b>	<b>12.6</b>	<b>6.0</b>
Forest Land Remaining Forest Land: Forest Fires	1.7	5.5	3.8	3.1	9.6	10.3	3.8
Settlements Remaining Settlements: Settlement Soils	0.1	0.3	0.3	0.3	0.3	0.3	0.3
Forest Land Remaining Forest Land: Forest Soils	1.0	1.8	1.7	1.8	1.9	1.9	1.8
Wetlands Remaining Wetlands: Peatlands Remaining Peatlands	+	+	+	+	+	+	+
<b>Total</b>	<b>13.4</b>	<b>24.8</b>	<b>20.0</b>	<b>19.6</b>	<b>35.3</b>	<b>39.0</b>	<b>22.6</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values. Totals may not sum due to independent rounding.

+ Less than 0.05 MMT CO<sub>2</sub> Eq.

11 Other significant trends from 1990 to 2013 in emissions from land use, land-use change, and forestry source  
12 categories include:

- 13 • Net C sequestration by forest land (i.e., carbon stock accumulation in the five carbon pools) has increased  
14 by approximately 21 percent. This is primarily due to increased forest management and the effects of  
15 previous reforestation. The increase in intensive forest management resulted in higher growth rates and  
16 higher biomass density. The tree planting and conservation efforts of the 1970s and 1980s continue to have

a significant impact on sequestration rates. Finally, the forested area in the United States increased over the past twenty four-years, although only at an average rate of 0.1 percent per year.


- Net sequestration of C by urban trees has increased by 48.1 percent over the period from 1990 to 2013. This is primarily due to an increase in urbanized land area in the United States.
- Annual C sequestration in landfilled yard trimmings and food scraps has decreased by 47.0 percent since 1990. Food scrap generation has grown by 53 percent since 1990, and though the proportion of food scraps discarded in landfills has decreased slightly from 82 percent in 1990 to 78 percent in 2013, the tonnage disposed in landfills has increased considerably (by 46 percent). Overall, the decrease in the landfill disposal rate of yard trimmings has more than compensated for the increase in food scrap disposal in landfills.

## Waste

Waste management and treatment activities are sources of greenhouse gas emissions (see Figure 2-11). In 2013, landfills were the third largest source of U.S. anthropogenic CH<sub>4</sub> emissions, accounting for 17.5 percent of total U.S. CH<sub>4</sub> emissions.<sup>50</sup> Additionally, wastewater treatment accounts for 14.4 percent of Waste emissions, 2.3 percent of U.S. CH<sub>4</sub> emissions, and 1.4 percent of N<sub>2</sub>O emissions. Emissions of CH<sub>4</sub> and N<sub>2</sub>O from composting grew from 1990 to 2013, and resulted in emissions of 3.7 MMT CO<sub>2</sub> Eq. in 2013. A summary of greenhouse gas emissions from the Waste chapter is presented in Table 2-10.

**Figure 2-11: 2013 Waste Chapter Greenhouse Gas Sources**

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



Overall, in 2013, waste activities generated emissions of 138.3 MMT CO<sub>2</sub> Eq., or 2.1 percent of total U.S. greenhouse gas emissions.

**Table 2-10: Emissions from Waste (MMT CO<sub>2</sub> Eq.)**

Gas/Source	1990	2005	2009	2010	2011	2012	2013
<b>CH<sub>4</sub></b>	<b>202.3</b>	<b>183.2</b>	<b>175.5</b>	<b>139.1</b>	<b>138.4</b>	<b>132.4</b>	<b>131.6</b>
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6
Wastewater Treatment	15.7	15.9	15.6	15.5	15.3	15.2	15.0

<sup>50</sup> Landfills also store carbon, due to incomplete degradation of organic materials such as wood products and yard trimmings, as described in the Land Use, Land-Use Change, and Forestry chapter.

Composting	0.4	1.9	1.9	1.8	1.9	1.9	2.0
<b>N<sub>2</sub>O</b>	<b>3.7</b>	<b>6.0</b>	<b>6.3</b>	<b>6.4</b>	<b>6.5</b>	<b>6.6</b>	<b>6.7</b>
Wastewater Treatment	3.4	4.3	4.6	4.7	4.8	4.9	4.9
Composting	0.3	1.7	1.7	1.6	1.7	1.7	1.8
<b>Total</b>	<b>206.0</b>	<b>189.2</b>	<b>181.8</b>	<b>145.5</b>	<b>144.9</b>	<b>138.9</b>	<b>138.3</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Totals may not sum due to independent rounding.

1 Some significant trends in U.S. emissions from waste source categories include the following:

- 2 • From 1990 to 2013, net CH<sub>4</sub> emissions from landfills decreased by 71.6 MMT CO<sub>2</sub> Eq. (38.4 percent), with  
3 small increases occurring in interim years. This downward trend in overall emissions is the result of  
4 increases in the amount of landfill gas collected and combusted as well as reduction in the amount of  
5 decomposable materials (i.e., paper and paperboard, food scraps, and yard trimmings) discarded in MSW  
6 landfills over the time series,<sup>51</sup> which has more than offset the additional CH<sub>4</sub> emissions resulting from an  
7 increase in the amount of municipal solid waste landfilled.
- 8 • Combined CH<sub>4</sub> and N<sub>2</sub>O emissions from composting have generally increased since 1990, from 0.7 MMT  
9 CO<sub>2</sub> Eq. to 3.7 MMT CO<sub>2</sub> Eq. in 2013, which represents slightly more than a five-fold increase over the  
10 time series. The growth in composting since the 1990s is attributable to primarily two factors: (1) steady  
11 growth in population and residential housing, and (2) the enactment of legislation by state and local  
12 governments that discouraged the disposal of yard trimmings in landfills.
- 13 • From 1990 to 2013, CH<sub>4</sub> and N<sub>2</sub>O emissions from wastewater treatment decreased by 0.6 MMT CO<sub>2</sub> Eq.  
14 (4.0 percent) and increased by 1.6 MMT CO<sub>2</sub> Eq. (46.5 percent), respectively. Methane emissions from  
15 domestic wastewater treatment have decreased since 1999 due to decreasing percentages of wastewater  
16 being treated in anaerobic systems, including reduced use of on-site septic systems and central anaerobic  
17 treatment systems. Nitrous oxide emissions from wastewater treatment processes gradually increased  
18 across the time series as a result of increasing U.S. population and protein consumption.

## 19 2.1 Emissions by Economic Sector

20 Throughout this report, emission estimates are grouped into six sectors (i.e., chapters) defined by the IPCC and  
21 detailed above: Energy; Industrial Processes; Solvent and Other Product Use; Agriculture; Land Use, Land-Use  
22 Change, and Forestry; and Waste. While it is important to use this characterization for consistency with UNFCCC  
23 reporting guidelines, it is also useful to allocate emissions into more commonly used sectoral categories. This  
24 section reports emissions by the following U.S. economic sectors: residential, commercial, industry, transportation,  
25 electricity generation, and agriculture, as well as U.S. territories.

26 Using this categorization, emissions from electricity generation accounted for the largest portion (31 percent) of  
27 U.S. greenhouse gas emissions in 2013. Transportation activities, in aggregate, accounted for the second largest  
28 portion (27 percent). Emissions from industry accounted for about 21 percent of U.S. greenhouse gas emissions in  
29 2013. In contrast to electricity generation and transportation, emissions from industry have in general declined over  
30 the past decade. The long-term decline in these emissions has been due to structural changes in the U.S. economy  
31 (i.e., shifts from a manufacturing-based to a service-based economy), fuel switching, and efficiency improvements.  
32 The remaining 21 percent of U.S. greenhouse gas emissions were contributed by the residential, agriculture, and  
33 commercial sectors, plus emissions from U.S. territories. The residential sector accounted for 6 percent, and  
34 primarily consisted of CO<sub>2</sub> emissions from fossil fuel combustion. Activities related to agriculture accounted for  
35 roughly 9 percent of U.S. emissions; unlike other economic sectors, agricultural sector emissions were dominated by  
36 N<sub>2</sub>O emissions from agricultural soil management and CH<sub>4</sub> emissions from enteric fermentation, rather than CO<sub>2</sub>  
37 from fossil fuel combustion. The commercial sector accounted for roughly 6 percent of emissions, while U.S.

<sup>51</sup> The CO<sub>2</sub> produced from combusted landfill CH<sub>4</sub> at landfills is not counted in national inventories as it is considered part of the natural C cycle of decomposition.


1 territories accounted for less than 1 percent. Carbon dioxide was also emitted and sequestered (in the form of C) by a  
 2 variety of activities related to forest management practices, tree planting in urban areas, the management of  
 3 agricultural soils, and landfilling of yard trimmings.

4 Table 2-11 presents a detailed breakdown of emissions from each of these economic sectors by source category, as  
 5 they are defined in this report. Figure 2-12 shows the trend in emissions by sector from 1990 to 2013.

6

7 **Figure 2-12: Emissions Allocated to Economic Sectors**

8 Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



9

10 **Table 2-11: U.S. Greenhouse Gas Emissions Allocated to Economic Sectors (MMT CO<sub>2</sub> Eq. and**  
 11 **Percent of Total in 2013)**

Sector/Source	1990	2005	2009	2010	2011	2012	2013	Percent <sup>a</sup>
<b>Electric Power Industry</b>	<b>1,864.8</b>	<b>2,445.2</b>	<b>2,186.4</b>	<b>2,301.2</b>	<b>2,198.9</b>	<b>2,061.3</b>	<b>2,077.7</b>	<b>30.8%</b>
CO <sub>2</sub> from Fossil Fuel Combustion	1,820.8	2,402.1	2,146.4	2,259.2	2,158.5	2,022.7	2,040.5	30.3%
Stationary Combustion	7.7	16.5	17.2	18.9	18.0	18.2	19.5	0.3%
Incineration of Waste	8.4	12.8	11.6	11.4	10.9	10.7	10.4	0.2%
Electrical Transmission and Distribution	25.4	10.6	7.3	6.9	6.8	5.7	5.1	0.1%
Other Process Uses of Carbonates	2.5	3.2	3.8	4.8	4.7	4.0	2.2	0.0%
<b>Transportation</b>	<b>1,551.3</b>	<b>2,021.0</b>	<b>1,862.0</b>	<b>1,875.7</b>	<b>1,849.2</b>	<b>1,831.0</b>	<b>1,841.8</b>	<b>27.3%</b>
CO <sub>2</sub> from Fossil Fuel Combustion	1,493.8	1,891.0	1,747.0	1,764.1	1,745.4	1,735.9	1,754.0	26.0%
Substitution of Ozone Depleting Substances	+	80.4	81.4	77.9	72.0	66.3	60.5	0.9%
Mobile Combustion	45.7	39.4	25.1	24.1	22.9	20.5	18.6	0.3%
Non-Energy Use of Fuels	11.8	10.2	8.5	9.5	9.0	8.3	8.8	0.1%
<b>Industry</b>	<b>1,585.8</b>	<b>1,476.4</b>	<b>1,311.3</b>	<b>1,385.8</b>	<b>1,384.5</b>	<b>1,381.1</b>	<b>1,424.1</b>	<b>21.1%</b>
CO <sub>2</sub> from Fossil Fuel Combustion	811.4	782.0	681.6	728.0	724.3	732.9	767.5	11.4%
Natural Gas Systems	212.7	201.8	202.5	194.5	197.7	192.2	197.7	2.9%
Non-Energy Use of Fuels	103.3	137.2	116.9	116.4	110.3	117.3	121.6	1.8%
Coal Mining	96.5	64.1	79.9	82.3	71.2	66.5	64.6	1.0%
Iron and Steel Production	100.9	67.5	43.5	56.4	60.7	55.1	53.0	0.8%
Petroleum Systems	35.0	28.9	39.0	39.9	41.1	43.9	46.5	0.7%
Cement Production	33.3	45.9	29.4	31.3	32.0	35.1	36.1	0.5%
Petrochemical Production	21.9	28.3	23.8	27.5	26.5	26.6	26.4	0.4%
Substitution of Ozone Depleting Substances	+	7.3	12.4	15.3	17.0	18.7	20.4	0.3%
Lime Production	11.7	14.6	11.4	13.4	14.0	13.7	14.1	0.2%
Nitric Acid Production	12.1	11.3	9.6	11.5	10.9	10.5	10.7	0.2%
Ammonia Production	13.0	9.2	8.5	9.2	9.3	9.4	10.2	0.2%

Abandoned Underground Coal Mines	7.2	6.6	6.4	6.6	6.4	6.2	6.2	0.1%
Aluminum Production	28.3	7.6	4.9	4.6	6.8	6.4	6.2	0.1%
HCFC-22 Production	46.1	20.0	6.8	8.0	8.8	5.5	5.5	0.1%
Urea Consumption for Non-Agricultural Purposes	3.8	3.7	3.4	4.7	4.0	4.4	4.7	0.1%
N <sub>2</sub> O from Product Uses	4.2	4.2	4.2	4.2	4.2	4.2	4.2	0.1%
Semiconductor Manufacture	3.6	4.7	3.1	3.8	4.9	4.5	4.2	0.1%
Adipic Acid Production	15.2	7.1	2.7	4.2	10.2	5.5	4.0	0.1%
Stationary Combustion	4.9	4.6	3.6	3.9	3.9	3.9	3.9	0.1%
Soda Ash Production and Consumption	2.7	2.9	2.5	2.6	2.6	2.7	2.7	0.0%
Other Process Uses of Carbonates	2.5	3.2	3.8	4.8	4.7	4.0	2.2	0.0%
Ferroalloy Production	2.2	1.4	1.5	1.7	1.7	1.9	1.8	0.0%
Titanium Dioxide Production	1.2	1.8	1.6	1.8	1.7	1.5	1.6	0.0%
Magnesium Production and Processing	5.2	2.7	1.6	2.1	2.8	1.7	1.5	0.0%
Mobile Combustion	0.9	1.3	1.3	1.4	1.4	1.4	1.5	0.0%
Zinc Production	0.6	1.0	0.9	1.2	1.3	1.5	1.4	0.0%
Phosphoric Acid Production	1.6	1.4	1.0	1.1	1.2	1.1	1.2	0.0%
Glass Production	1.5	1.9	1.0	1.5	1.3	1.2	1.2	0.0%
Carbon Dioxide Consumption	1.5	1.4	1.8	1.2	0.8	0.8	0.9	0.0%
Lead Production	0.5	0.6	0.5	0.5	0.5	0.5	0.5	0.0%
Silicon Carbide Production and Consumption	0.4	0.2	0.2	0.2	0.2	0.2	0.2	0.0%
<b>Agriculture</b>	<b>492.5</b>	<b>564.9</b>	<b>588.8</b>	<b>590.8</b>	<b>605.6</b>	<b>611.7</b>	<b>586.8</b>	<b>8.7%</b>
N <sub>2</sub> O from Agricultural Soil Management	224.0	243.6	264.1	264.3	265.8	266.0	263.7	3.9%
Enteric Fermentation	164.2	168.9	172.7	171.1	168.7	166.3	164.5	2.4%
Manure Management	51.0	72.8	76.7	78.0	78.7	81.0	78.7	1.2%
CO <sub>2</sub> from Fossil Fuel Combustion	31.0	46.8	46.7	47.6	49.4	51.0	49.8	0.7%
CH <sub>4</sub> and N <sub>2</sub> O from Forest Fires	4.2	13.8	9.7	7.9	24.2	26.0	9.7	0.1%
Rice Cultivation	9.2	8.9	9.4	11.1	8.5	9.3	8.3	0.1%
Liming of Agricultural Soils	4.7	4.3	3.7	4.8	3.9	5.8	5.9	0.1%
Urea Fertilization	2.4	3.5	3.6	3.8	4.1	4.2	4.0	0.1%
CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O from Managed Peatlands	1.1	1.1	1.0	1.0	0.9	0.8	0.8	0.0%
Mobile Combustion	0.3	0.5	0.5	0.5	0.5	0.56	0.5	0.0%
Field Burning of Agricultural Residues	0.4	0.3	0.4	0.4	0.4	0.4	0.4	0.0%
N <sub>2</sub> O from Forest Soils	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.0%
Stationary Combustion	+	+	+	+	+	+	+	0.0%
<b>Commercial</b>	<b>424.8</b>	<b>430.0</b>	<b>432.3</b>	<b>396.7</b>	<b>401.0</b>	<b>374.8</b>	<b>401.9</b>	<b>6.0%</b>
CO <sub>2</sub> from Fossil Fuel Combustion	217.4	223.7	223.8	220.5	221.3	197.5	221.5	3.3%
Landfills	186.2	165.5	158.1	121.8	121.3	115.3	114.6	1.7%
Substitution of Ozone Depleting Substances	+	15.7	25.2	29.3	33.4	37.1	40.8	0.6%
Wastewater Treatment	15.7	15.9	15.6	15.5	15.3	15.2	15.0	0.2%
Human Sewage	3.4	4.3	4.6	4.7	4.8	4.9	4.9	0.1%
Composting	0.7	3.5	3.6	3.5	3.5	3.7	3.7	0.1%
Stationary Combustion	1.4	1.4	1.4	1.4	1.4	1.2	1.3	0.0%
<b>Residential</b>	<b>345.9</b>	<b>372.3</b>	<b>360.7</b>	<b>363.5</b>	<b>360.2</b>	<b>321.2</b>	<b>374.7</b>	<b>5.6%</b>
CO <sub>2</sub> from Fossil Fuel Combustion	338.3	357.9	336.7	335.1	327.5	283.4	329.9	4.9%
Substitution of Ozone Depleting Substances	0.3	7.7	17.0	21.8	25.9	31.4	37.0	0.5%
Stationary Combustion	6.3	4.9	5.3	4.8	4.9	4.5	5.9	0.1%
Settlement Soil Fertilization	1.0	1.8	1.7	1.8	1.9	1.9	1.8	0.0%
<b>U.S. Territories</b>	<b>33.7</b>	<b>58.2</b>	<b>47.9</b>	<b>51.0</b>	<b>43.9</b>	<b>42.5</b>	<b>35.1</b>	<b>0.5%</b>
CO <sub>2</sub> from Fossil Fuel Combustion	27.9	50.0	43.8	46.6	40.2	39.0	32.3	0.5%
Non-Energy Use of Fuels	5.7	8.1	3.9	4.2	3.6	3.3	2.7	0.0%
Stationary Combustion	0.1	0.2	0.2	0.2	0.2	0.2	0.1	0.0%
<b>Total Emissions</b>	<b>6,298.8</b>	<b>7,367.9</b>	<b>6,789.5</b>	<b>6,964.7</b>	<b>6,843.3</b>	<b>6,623.4</b>	<b>6,742.2</b>	<b>100.0%</b>
<b>Sinks</b>	<b>(774.1)</b>	<b>(912.6)</b>	<b>(871.3)</b>	<b>(872.0)</b>	<b>(881.3)</b>	<b>(880.7)</b>	<b>(882.0)</b>	<b>-13.1%</b>
CO <sub>2</sub> Flux from Forests <sup>b</sup>	(639.4)	(807.1)	(764.9)	(765.4)	(773.8)	(773.1)	(775.7)	-11.5%

Urban Trees	(60.4)	(80.5)	(85.0)	(86.1)	(87.3)	(88.4)	(89.5)	-1.3%
Landfilled Yard Trimmings and Food Scraps	(24.2)	(12.0)	(12.9)	(13.6)	(13.5)	(13.0)	(12.8)	-0.2%
CO <sub>2</sub> Flux from Agricultural Soil Carbon Stocks	(50.0)	(13.0)	(8.5)	(6.9)	(6.7)	(6.1)	(4.0)	-0.1%
<b>Net Emissions</b>	<b>5,524.7</b>	<b>6,455.4</b>	<b>5,918.2</b>	<b>6,092.7</b>	<b>5,962.0</b>	<b>5,742.7</b>	<b>5,860.2</b>	<b>86.9%</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Includes all emissions of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs, SF<sub>6</sub>, and NF<sub>3</sub>. Parentheses indicate negative values or sequestration. Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq. or 0.05 percent.

<sup>a</sup> Percent of total emissions for year 2013.

<sup>b</sup> Includes the effects of net additions to stocks of carbon stored in harvested wood products.

## 1 Emissions with Electricity Distributed to Economic Sectors

2 It can also be useful to view greenhouse gas emissions from economic sectors with emissions related to electricity  
3 generation distributed into end-use categories (i.e., emissions from electricity generation are allocated to the  
4 economic sectors in which the electricity is consumed). The generation, transmission, and distribution of electricity,  
5 which is the largest economic sector in the United States, accounted for 31 percent of total U.S. greenhouse gas  
6 emissions in 2013. Emissions increased by 11 percent since 1990, as electricity demand grew and fossil fuels  
7 remained the dominant energy source for generation. Electricity generation-related emissions increased from 2012  
8 to 2013 by 0.8 percent, primarily due to increased CO<sub>2</sub> emissions from fossil fuel combustion. Electricity sales to  
9 the residential and commercial end-use sectors in 2013 increased approximately 1.2 percent and 0.9 percent,  
10 respectively. The trend in the residential and commercial sectors can largely be attributed to colder more energy-  
11 intensive winter conditions compared to 2012. Electricity sales to the industrial sector in 2013 decreased by  
12 approximately 3.1 percent. Overall, in 2013, the amount of electricity generated (in kWh) decreased by 0.1 percent  
13 from the previous year. Despite the decrease in generation, CO<sub>2</sub> emissions from the electric power sector increased  
14 by 0.8 percent as the consumption of CO<sub>2</sub> intensive coal and petroleum for electricity generation increased by 4.2  
15 percent and 19.3 percent, respectively, in 2013 and the consumption of natural gas for electricity generation,  
16 decreased by 10.2 percent. Table 2-12 provides a detailed summary of emissions from electricity generation-related  
17 activities.

18 **Table 2-12: Electricity Generation-Related Greenhouse Gas Emissions (MMT CO<sub>2</sub> Eq.)**

Gas/Fuel Type or Source	1990	2005	2009	2010	2011	2012	2013
<b>CO<sub>2</sub></b>	<b>1,831.2</b>	<b>2,417.8</b>	<b>2,161.5</b>	<b>2,275.0</b>	<b>2,173.7</b>	<b>2,037.1</b>	<b>2,052.8</b>
Fossil Fuel Combustion	1,820.8	2,402.1	2,146.4	2,259.2	2,158.5	2,022.7	2,040.5
<i>Coal</i>	<i>1,547.6</i>	<i>1,983.8</i>	<i>1,740.9</i>	<i>1,827.6</i>	<i>1,722.7</i>	<i>1,511.2</i>	<i>1,575.0</i>
<i>Natural Gas</i>	<i>175.3</i>	<i>318.8</i>	<i>372.2</i>	<i>399.0</i>	<i>408.8</i>	<i>492.2</i>	<i>441.9</i>
<i>Petroleum</i>	<i>97.5</i>	<i>99.2</i>	<i>33.0</i>	<i>32.2</i>	<i>26.6</i>	<i>18.8</i>	<i>23.1</i>
<i>Geothermal</i>	<i>0.4</i>						
Incineration of Waste	8.0	12.5	11.3	11.0	10.5	10.4	10.1
Other Process Uses of Carbonates	2.5	3.2	3.8	4.8	4.7	4.0	2.2
<b>CH<sub>4</sub></b>	<b>0.3</b>	<b>0.5</b>	<b>0.4</b>	<b>0.5</b>	<b>0.4</b>	<b>0.4</b>	<b>0.4</b>
Stationary Combustion <sup>a</sup>	0.3	0.5	0.4	0.5	0.4	0.4	0.4
Incineration of Waste	+	+	+	+	+	+	+
<b>N<sub>2</sub>O</b>	<b>7.8</b>	<b>16.4</b>	<b>17.1</b>	<b>18.8</b>	<b>17.9</b>	<b>18.1</b>	<b>19.4</b>
Stationary Combustion <sup>a</sup>	7.4	16.0	16.8	18.5	17.6	17.8	19.1
Incineration of Waste	0.5	0.4	0.3	0.3	0.3	0.3	0.3
<b>SF<sub>6</sub></b>	<b>25.4</b>	<b>10.6</b>	<b>7.3</b>	<b>6.9</b>	<b>6.8</b>	<b>5.7</b>	<b>5.1</b>
Electrical Transmission and Distribution	25.4	10.6	7.3	6.9	6.8	5.7	5.1
<b>Total</b>	<b>1,864.8</b>	<b>2,445.2</b>	<b>2,186.4</b>	<b>2,301.2</b>	<b>2,198.9</b>	<b>2,061.3</b>	<b>2,077.7</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding.

<sup>a</sup> Includes only stationary combustion emissions related to the generation of electricity.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.


To distribute electricity emissions among economic end-use sectors, emissions from the source categories assigned to the electricity generation sector were allocated to the residential, commercial, industry, transportation, and agriculture economic sectors according to each economic sector's share of retail sales of electricity consumption (EIA 2011, Duffield 2006). These source categories include CO<sub>2</sub> from Fossil Fuel Combustion, CH<sub>4</sub> and N<sub>2</sub>O from Stationary Combustion, Incineration of Waste, Other Process Uses of Carbonates, and SF<sub>6</sub> from Electrical Transmission and Distribution Systems. Note that only 50 percent of the Other Process Uses of Carbonates emissions were associated with electricity generation and distributed as described; the remainder of Other Process Uses of Carbonates emissions were attributed to the industrial processes economic end-use sector.<sup>52</sup>

When emissions from electricity are distributed among these sectors, industrial activities account for the largest share of total U.S. greenhouse gas emissions (29.0 percent), followed closely by emissions from transportation (27.4 percent). Emissions from the residential and commercial sectors also increase substantially when emissions from electricity are included. In all sectors except agriculture, CO<sub>2</sub> accounts for more than 80 percent of greenhouse gas emissions, primarily from the combustion of fossil fuels.

Table 2-13 presents a detailed breakdown of emissions from each of these economic sectors, with emissions from electricity generation distributed to them. Figure 2-13 shows the trend in these emissions by sector from 1990 to 2013.

**Figure 2-13: Emissions with Electricity Distributed to Economic Sectors**

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.



**Table 2-13: U.S. Greenhouse Gas Emissions by Economic Sector and Gas with Electricity-Related Emissions Distributed (MMT CO<sub>2</sub> Eq.) and Percent of Total in 2013**

Sector/Gas	1990	2005	2009	2010	2011	2012	2013	Percent <sup>a</sup>
<b>Industry</b>	<b>2,227.8</b>	<b>2,162.4</b>	<b>1,856.7</b>	<b>1,970.4</b>	<b>1,955.7</b>	<b>1,923.1</b>	<b>1,954.9</b>	<b>29.0%</b>
<b>Direct Emissions</b>	<b>1,585.8</b>	<b>1,476.4</b>	<b>1,311.3</b>	<b>1,385.8</b>	<b>1,384.5</b>	<b>1,381.1</b>	<b>1,424.1</b>	<b>21.1%</b>
CO <sub>2</sub>	1,161.5	1,142.0	972.7	1,041.9	1,039.7	1,052.5	1,093.5	16.2%
CH <sub>4</sub>	312.7	269.5	292.9	289.3	278.9	271.5	273.6	4.1%
N <sub>2</sub> O	35.3	26.6	19.9	23.6	28.9	23.8	22.5	0.3%
HFCs, PFCs, SF <sub>6</sub> , and NF <sub>3</sub>	76.3	38.3	25.8	31.1	37.0	33.3	34.5	0.5%
<b>Electricity-Related</b>	<b>642.0</b>	<b>686.1</b>	<b>545.4</b>	<b>584.6</b>	<b>571.2</b>	<b>542.1</b>	<b>530.7</b>	<b>7.9%</b>

<sup>52</sup> Emissions were not distributed to U.S. territories, since the electricity generation sector only includes emissions related to the generation of electricity in the 50 states and the District of Columbia.

CO <sub>2</sub>	630.4	678.4	539.2	577.9	564.6	535.7	524.4	7.8%
CH <sub>4</sub>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0%
N <sub>2</sub> O	2.7	4.6	4.3	4.8	4.7	4.8	5.0	0.1%
SF <sub>6</sub>	8.7	3.0	1.8	1.8	1.8	1.5	1.3	0.0%
<b>Transportation</b>	<b>1,554.4</b>	<b>2,025.8</b>	<b>1,866.6</b>	<b>1,880.3</b>	<b>1,853.6</b>	<b>1,834.9</b>	<b>1,845.9</b>	<b>27.4%</b>
<b>Direct Emissions</b>	<b>1,551.3</b>	<b>2,021.0</b>	<b>1,862.0</b>	<b>1,875.7</b>	<b>1,849.2</b>	<b>1,831.0</b>	<b>1,841.8</b>	<b>27.3%</b>
CO <sub>2</sub>	1,505.6	1,901.2	1,755.6	1,773.6	1,754.4	1,744.2	1,762.8	26.1%
CH <sub>4</sub>	5.4	2.7	2.0	1.9	1.9	1.8	1.7	0.0%
N <sub>2</sub> O	40.3	36.7	23.1	22.2	21.0	18.7	16.8	0.2%
HFCs <sup>b</sup>	+	80.4	81.4	77.9	72.0	66.3	60.5	0.9%
<b>Electricity-Related</b>	<b>3.1</b>	<b>4.8</b>	<b>4.6</b>	<b>4.6</b>	<b>4.3</b>	<b>3.9</b>	<b>4.1</b>	<b>0.1%</b>
CO <sub>2</sub>	3.1	4.8	4.5	4.5	4.3	3.9	4.0	0.1%
CH <sub>4</sub>	+	+	+	+	+	+	+	0.0%
N <sub>2</sub> O	+	+	+	+	+	+	+	0.0%
SF <sub>6</sub>	+	+	+	+	+	+	+	0.0%
<b>Commercial</b>	<b>975.8</b>	<b>1,248.1</b>	<b>1,199.8</b>	<b>1,184.4</b>	<b>1,153.1</b>	<b>1,088.6</b>	<b>1,127.8</b>	<b>16.7%</b>
<b>Direct Emissions</b>	<b>424.8</b>	<b>430.0</b>	<b>432.3</b>	<b>396.7</b>	<b>401.0</b>	<b>374.8</b>	<b>401.9</b>	<b>6.0%</b>
CO <sub>2</sub>	217.4	223.7	223.8	220.5	221.3	197.5	221.5	3.3%
CH <sub>4</sub>	203.3	184.3	176.6	140.2	139.4	133.3	132.7	2.0%
N <sub>2</sub> O	4.1	6.3	6.6	6.7	6.8	6.8	7.0	0.1%
HFCs	+	15.7	25.2	29.3	33.4	37.1	40.8	0.6%
<b>Electricity-Related</b>	<b>551.0</b>	<b>818.1</b>	<b>767.5</b>	<b>787.6</b>	<b>752.1</b>	<b>713.8</b>	<b>725.9</b>	<b>10.8%</b>
CO <sub>2</sub>	541.1	808.9	758.8	778.7	743.5	705.4	717.2	10.6%
CH <sub>4</sub>	0.1	0.2	0.2	0.2	0.1	0.1	0.1	0.0%
N <sub>2</sub> O	2.3	5.5	6.0	6.4	6.1	6.3	6.8	0.1%
SF <sub>6</sub>	7.5	3.5	2.6	2.4	2.3	2.0	1.8	0.0%
<b>Residential</b>	<b>953.3</b>	<b>1,244.4</b>	<b>1,161.8</b>	<b>1,219.5</b>	<b>1,166.0</b>	<b>1,060.5</b>	<b>1,129.1</b>	<b>16.7%</b>
<b>Direct Emissions</b>	<b>345.9</b>	<b>372.3</b>	<b>360.7</b>	<b>363.5</b>	<b>360.2</b>	<b>321.2</b>	<b>374.7</b>	<b>5.6%</b>
CO <sub>2</sub>	338.3	357.9	336.7	335.1	327.5	283.4	329.9	4.9%
CH <sub>4</sub>	5.2	4.1	4.4	4.0	4.0	3.7	5.0	0.1%
N <sub>2</sub> O	2.1	2.6	2.6	2.6	2.7	2.6	2.8	0.0%
HFCs	0.3	7.7	17.0	21.8	25.9	31.4	37.0	0.5%
<b>Electricity-Related</b>	<b>607.3</b>	<b>872.1</b>	<b>801.1</b>	<b>856.0</b>	<b>805.8</b>	<b>739.3</b>	<b>754.4</b>	<b>11.2%</b>
CO <sub>2</sub>	596.4	862.3	792.0	846.3	796.6	730.6	745.4	11.1%
CH <sub>4</sub>	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.0%
N <sub>2</sub> O	2.5	5.8	6.3	7.0	6.6	6.5	7.1	0.1%
SF <sub>6</sub>	8.3	3.8	2.7	2.6	2.5	2.0	1.8	0.0%
<b>Agriculture</b>	<b>553.8</b>	<b>629.0</b>	<b>656.7</b>	<b>659.2</b>	<b>671.0</b>	<b>673.9</b>	<b>649.5</b>	<b>9.6%</b>
<b>Direct Emissions</b>	<b>492.5</b>	<b>564.9</b>	<b>588.8</b>	<b>590.8</b>	<b>605.6</b>	<b>611.7</b>	<b>586.8</b>	<b>8.7%</b>
CO <sub>2</sub>	39.2	55.8	54.9	57.2	58.3	61.8	60.6	0.9%
CH <sub>4</sub>	213.4	242.9	248.2	248.4	253.7	255.5	240.6	3.6%
N <sub>2</sub> O	239.9	266.3	285.8	285.2	293.5	294.4	285.7	4.2%
<b>Electricity-Related</b>	<b>61.3</b>	<b>64.1</b>	<b>67.8</b>	<b>68.4</b>	<b>65.4</b>	<b>62.1</b>	<b>62.7</b>	<b>0.9%</b>
CO <sub>2</sub>	60.2	63.4	67.1	67.6	64.7	61.4	61.9	0.9%
CH <sub>4</sub>	+	+	+	+	+	+	+	0.0%
N <sub>2</sub> O	0.3	0.4	0.5	0.6	0.5	0.5	0.6	0.0%
SF <sub>6</sub>	0.8	0.3	0.2	0.2	0.2	0.2	0.2	0.0%
<b>U.S. Territories</b>	<b>33.7</b>	<b>58.2</b>	<b>47.9</b>	<b>51.0</b>	<b>43.9</b>	<b>42.5</b>	<b>35.1</b>	<b>0.5%</b>
<b>Total</b>	<b>6,298.8</b>	<b>7,367.9</b>	<b>6,789.5</b>	<b>6,964.7</b>	<b>6,843.3</b>	<b>6,623.4</b>	<b>6,742.2</b>	<b>100.0%</b>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Emissions from electricity generation are allocated based on aggregate electricity consumption in each end-use sector.

Totals may not sum due to independent rounding.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq. or 0.05 percent.

<sup>a</sup> Percent of total emissions for year 2013.

<sup>b</sup> Includes primarily HFC-134a.

## 1 Industry

2 The industrial end-use sector includes CO<sub>2</sub> emissions from fossil fuel combustion from all manufacturing facilities,  
3 in aggregate. This sector also includes emissions that are produced as a byproduct of the non-energy-related  
4 industrial process activities. The variety of activities producing these non-energy-related emissions includes  
5 methane emissions from petroleum and natural gas systems, fugitive CH<sub>4</sub> emissions from coal mining, by-product  
6 CO<sub>2</sub> emissions from cement manufacture, and HFC, PFC, SF<sub>6</sub>, and NF<sub>3</sub> byproduct emissions from semiconductor  
7 manufacture, to name a few. Since 1990, industrial sector emissions have declined. The decline has occurred both  
8 in direct emissions and indirect emissions associated with electricity use. In theory, emissions from the industrial  
9 end-use sector should be highly correlated with economic growth and industrial output, but heating of industrial  
10 buildings and agricultural energy consumption are also affected by weather conditions. In addition, structural  
11 changes within the U.S. economy that lead to shifts in industrial output away from energy-intensive manufacturing  
12 products to less energy-intensive products (e.g., from steel to computer equipment) also have a significant effect on  
13 industrial emissions.

## 14 Transportation

15 When electricity-related emissions are distributed to economic end-use sectors, transportation activities accounted  
16 for 27 percent of U.S. greenhouse gas emissions in 2013. The largest sources of transportation greenhouse gases in  
17 2013 were passenger cars (42.7 percent), freight trucks (22.1 percent), light duty trucks, which include sport utility  
18 vehicles, pickup trucks, and minivans (18.0 percent), commercial aircraft (6.2 percent), rail (2.6 percent), pipelines  
19 (2.6 percent), and ships and boats (2.2 percent). These figures include direct CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O emissions from  
20 fossil fuel combustion used in transportation and emissions from non-energy use (i.e., lubricants) used in  
21 transportation, as well as HFC emissions from mobile air conditioners and refrigerated transport allocated to these  
22 vehicle types.

23 In terms of the overall trend, from 1990 to 2013, total transportation emissions rose by 19 percent due, in large part,  
24 to increased demand for travel as fleetwide light-duty vehicle fuel economy was relatively stable (average new  
25 vehicle fuel economy declined slowly from 1990 through 2004 and then increased more rapidly from 2005 through  
26 2013). The number of vehicle miles traveled by light-duty motor vehicles (passenger cars and light-duty trucks)  
27 increased 36 percent from 1990 to 2013, as a result of a confluence of factors including population growth,  
28 economic growth, urban sprawl, and low fuel prices during the beginning of this period. The decline in new light-  
29 duty vehicle fuel economy between 1990 and 2004 reflected the increasing market share of light-duty trucks, which  
30 grew from about 30 percent of new vehicle sales in 1990 to 48 percent in 2004. Starting in 2005, the rate of VMT  
31 growth slowed considerably (and declined rapidly in 2008) while average new vehicle fuel economy began to  
32 increase. Average new vehicle fuel economy has improved almost every year since 2005, and the truck share has  
33 decreased to about 37 percent of new vehicles in MY 2013 (EPA, 2014d). Between 2012 and 2013, VMT increased  
34 by only 0.6 percent. Table 2-14 provides a detailed summary of greenhouse gas emissions from transportation-  
35 related activities with electricity-related emissions included in the totals.

36 From 2008 to 2009, CO<sub>2</sub> emissions from the transportation end-use sector declined 3.9 percent. The decrease in  
37 emissions could largely be attributed to decreased economic activity in 2009 and an associated decline in the  
38 demand for transportation. Modes such as medium- and heavy-duty trucks were significantly impacted by the  
39 decline in freight transport. From 2009 to 2013, CO<sub>2</sub> emissions from the transportation end-use sector stabilized  
40 even as economic activity rebounded slightly.

41 Almost all of the energy consumed for transportation was supplied by petroleum-based products, with more than  
42 half being related to gasoline consumption in automobiles and other highway vehicles. Other fuel uses, especially  
43 diesel fuel for freight trucks and jet fuel for aircraft, accounted for the remainder. The primary driver of  
44 transportation-related emissions was CO<sub>2</sub> from fossil fuel combustion, which increased by 17 percent from 1990 to  
45 2013. This rise in CO<sub>2</sub> emissions, combined with an increase in HFCs from close to zero emissions in 1990 to 60.5  
46 MMT CO<sub>2</sub> Eq. in 2013, led to an increase in overall emissions from transportation activities of 19 percent.

1 **Table 2-14: Transportation-Related Greenhouse Gas Emissions (MMT CO<sub>2</sub> Eq.)**

Gas/Vehicle	1990	2005	2009	2010	2011	2012	2013
<b>Passenger Cars</b>	<b>656.7</b>	<b>713.0</b>	<b>810.9</b>	<b>805.3</b>	<b>797.5</b>	<b>792.1</b>	<b>788.5</b>
CO <sub>2</sub>	629.3	661.9	766.0	763.7	760.1	759.7	760.8
CH <sub>4</sub>	3.2	1.4	1.2	1.2	1.2	1.1	1.1
N <sub>2</sub> O	24.1	18.0	13.8	12.9	12.3	10.7	9.4
HFCs	+	31.7	29.9	27.5	23.9	20.6	17.3
<b>Light-Duty Trucks</b>	<b>335.6</b>	<b>554.6</b>	<b>358.8</b>	<b>357.7</b>	<b>341.0</b>	<b>335.3</b>	<b>333.0</b>
CO <sub>2</sub>	321.1	505.6	317.4	317.6	303.8	301.2	301.9
CH <sub>4</sub>	1.7	0.9	0.4	0.4	0.4	0.4	0.3
N <sub>2</sub> O	12.8	14.8	5.8	5.5	5.1	4.4	3.9
HFCs	+	33.3	35.2	34.2	31.7	29.3	26.7
<b>Medium- and Heavy-Duty Trucks</b>	<b>231.1</b>	<b>409.9</b>	<b>390.3</b>	<b>403.8</b>	<b>402.0</b>	<b>401.9</b>	<b>407.1</b>
CO <sub>2</sub>	230.1	396.0	375.8	389.3	387.6	387.4	392.6
CH <sub>4</sub>	0.3	0.1	0.1	0.1	0.1	0.1	0.1
N <sub>2</sub> O	0.7	1.1	1.2	1.2	1.1	1.1	1.0
HFCs	+	12.7	13.2	13.2	13.3	13.3	13.3
<b>Buses</b>	<b>8.4</b>	<b>12.1</b>	<b>16.6</b>	<b>16.4</b>	<b>17.5</b>	<b>18.7</b>	<b>18.9</b>
CO <sub>2</sub>	8.4	11.8	16.0	15.9	16.9	18.1	18.3
CH <sub>4</sub>	+	+	+	+	+	+	+
N <sub>2</sub> O	+	+	0.1	0.1	0.1	0.1	0.1
HFCs	+	0.3	0.4	0.4	0.4	0.4	0.4
<b>Motorcycles</b>	<b>1.8</b>	<b>1.7</b>	<b>4.3</b>	<b>3.8</b>	<b>3.7</b>	<b>4.3</b>	<b>4.3</b>
CO <sub>2</sub>	1.7	1.6	4.2	3.8	3.7	4.2	4.2
CH <sub>4</sub>	+	+	+	+	+	+	+
N <sub>2</sub> O	+	+	+	+	+	+	+
<b>Commercial Aircraft<sup>a</sup></b>	<b>110.9</b>	<b>133.9</b>	<b>120.6</b>	<b>114.3</b>	<b>115.6</b>	<b>114.3</b>	<b>115.4</b>
CO <sub>2</sub>	109.9	132.7	119.5	113.3	114.6	113.3	114.3
CH <sub>4</sub>	+	+	+	+	+	+	+
N <sub>2</sub> O	1.0	1.2	1.1	1.0	1.1	1.0	1.1
<b>Other Aircraft<sup>b</sup></b>	<b>78.3</b>	<b>59.6</b>	<b>36.8</b>	<b>40.4</b>	<b>34.2</b>	<b>32.1</b>	<b>34.7</b>
CO <sub>2</sub>	77.5	59.1	36.4	40.1	33.9	31.8	34.4
CH <sub>4</sub>	0.1	0.1	+	+	+	+	+
N <sub>2</sub> O	0.7	0.5	0.3	0.4	0.3	0.3	0.3
<b>Ships and Boats<sup>c</sup></b>	<b>44.9</b>	<b>45.2</b>	<b>39.3</b>	<b>45.4</b>	<b>47.1</b>	<b>40.8</b>	<b>40.0</b>
CO <sub>2</sub>	44.3	44.5	38.7	44.6	46.2	40.0	39.3
CH <sub>4</sub>	+	+	+	+	+	+	+
N <sub>2</sub> O	0.6	0.6	0.5	0.8	0.8	0.7	0.7
HFCs	+	+	+	+	+	+	+
<b>Rail</b>	<b>39.0</b>	<b>53.3</b>	<b>43.7</b>	<b>46.5</b>	<b>48.1</b>	<b>46.8</b>	<b>47.5</b>
CO <sub>2</sub>	38.5	50.3	40.7	43.5	45.0	43.7	44.4
CH <sub>4</sub>	0.1	0.1	0.1	0.1	0.1	0.1	0.1
N <sub>2</sub> O	0.3	0.4	0.3	0.3	0.3	0.3	0.3
HFCs	+	2.5	2.6	2.6	2.6	2.6	2.6
Other Emissions from Electricity Generation <sup>d</sup>	0.1	0.1	+	+	+	+	+
<b>Pipelines<sup>e</sup></b>	<b>36.0</b>	<b>32.2</b>	<b>36.7</b>	<b>37.1</b>	<b>37.8</b>	<b>40.3</b>	<b>47.7</b>
CO <sub>2</sub>	36.0	32.2	36.7	37.1	37.8	40.3	47.7
<b>Lubricants</b>	<b>11.8</b>	<b>10.2</b>	<b>8.5</b>	<b>9.5</b>	<b>9.0</b>	<b>8.3</b>	<b>8.8</b>
CO <sub>2</sub>	11.8	10.2	8.5	9.5	9.0	8.3	8.8
<b>Total Transportation</b>	<b>1,554.4</b>	<b>2,025.8</b>	<b>1,866.6</b>	<b>1,880.3</b>	<b>1,853.6</b>	<b>1,834.9</b>	<b>1,845.9</b>
<i>International Bunker Fuels<sup>f</sup></i>	<i>104.5</i>	<i>114.2</i>	<i>107.5</i>	<i>118.1</i>	<i>112.8</i>	<i>106.8</i>	<i>100.7</i>

Note: Emissions values are presented in CO<sub>2</sub> equivalent mass units using IPCC AR4 GWP values.

Note: Totals may not sum due to independent rounding. Passenger cars and light-duty trucks include vehicles typically used for personal travel and less than 8,500 lbs; medium- and heavy-duty trucks include vehicles larger than

---

8,500 lbs. HFC emissions primarily reflect HFC-134a.

+ Does not exceed 0.05 MMT CO<sub>2</sub> Eq.

<sup>a</sup> Consists of emissions from jet fuel consumed by domestic operations of commercial aircraft (no bunkers).

<sup>b</sup> Consists of emissions from jet fuel and aviation gasoline consumption by general aviation and military aircraft.

<sup>c</sup> Fluctuations in emission estimates are associated with fluctuations in reported fuel consumption, and may reflect issues with data sources.

<sup>d</sup> Other emissions from electricity generation are a result of waste incineration (as the majority of municipal solid waste is combusted in “trash-to-steam” electricity generation plants), electrical transmission and distribution, and a portion of Other Process Uses of Carbonates (from pollution control equipment installed in electricity generation plants).

<sup>e</sup> CO<sub>2</sub> estimates reflect natural gas used to power pipelines, but not electricity. While the operation of pipelines produces CH<sub>4</sub> and N<sub>2</sub>O, these emissions are not directly attributed to pipelines in the US Inventory.

<sup>f</sup> Emissions from International Bunker Fuels include emissions from both civilian and military activities; these emissions are not included in the transportation totals.

## 1 Commercial

2 The commercial sector is heavily reliant on electricity for meeting energy needs, with electricity consumption for  
3 lighting, heating, air conditioning, and operating appliances. The remaining emissions were largely due to the direct  
4 consumption of natural gas and petroleum products, primarily for heating and cooking needs. Energy-related  
5 emissions from the residential and commercial sectors have generally been increasing since 1990, and are often  
6 correlated with short-term fluctuations in energy consumption caused by weather conditions, rather than prevailing  
7 economic conditions. Landfills and wastewater treatment are included in this sector, with landfill emissions  
8 decreasing since 1990 and wastewater treatment emissions decreasing slightly.

## 9 Residential

10 The residential sector is heavily reliant on electricity for meeting energy needs, with electricity consumption for  
11 lighting, heating, air conditioning, and operating appliances. The remaining emissions were largely due to the direct  
12 consumption of natural gas and petroleum products, primarily for heating and cooking needs. Emissions from the  
13 residential sectors have generally been increasing since 1990, and are often correlated with short-term fluctuations in  
14 energy consumption caused by weather conditions, rather than prevailing economic conditions. In the long-term,  
15 this sector is also affected by population growth, regional migration trends, and changes in housing and building  
16 attributes (e.g., size and insulation).

## 17 Agriculture

18 The agriculture sector includes a variety of processes, including enteric fermentation in domestic livestock, livestock  
19 manure management, and agricultural soil management. In 2013, agricultural soil management was the largest  
20 source of N<sub>2</sub>O emissions, and enteric fermentation was the largest source of CH<sub>4</sub> emissions in the United States.  
21 This sector also includes small amounts of CO<sub>2</sub> emissions from fossil fuel combustion by motorized farm equipment  
22 like tractors. The agriculture sector is less reliant on electricity than the other sectors.

23

### 24 **Box 2-1: Methodology for Aggregating Emissions by Economic Sector**

25 In presenting the Economic Sectors in the annual Inventory of U.S. Greenhouse Gas Emissions and Sinks, the  
26 Inventory expands upon the standard IPCC sectors common for UNFCCC reporting. Discussing greenhouse gas  
27 emissions relevant to U.S.-specific sectors improves communication of the report’s findings.

28 In the Electricity Generation economic sector, CO<sub>2</sub> emissions from the combustion of fossil fuels included in the  
29 EIA electric utility fuel consuming sector are apportioned to this economic sector. Stationary combustion emissions  
30 of CH<sub>4</sub> and N<sub>2</sub>O are also based on the EIA electric utility sector. Additional sources include CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O  
31 from waste incineration, as the majority of municipal solid waste is combusted in “trash-to-steam” electricity

1 generation plants. The Electricity Generation economic sector also includes SF<sub>6</sub> from Electrical Transmission and  
2 Distribution, and a portion of CO<sub>2</sub> from Other Process Uses of Carbonates (from pollution control equipment  
3 installed in electricity generation plants).

4 In the Transportation economic sector, the CO<sub>2</sub> emissions from the combustion of fossil fuels included in the EIA  
5 transportation fuel consuming sector are apportioned to this economic sector (additional analyses and refinement of  
6 the EIA data is further explained in the Energy chapter of this report). Additional emissions are apportioned from  
7 the CH<sub>4</sub> and N<sub>2</sub>O from Mobile Combustion, based on the EIA transportation sector. Substitutes of Ozone Depleting  
8 Substances are apportioned based on their specific end-uses within the source category, with emissions from  
9 transportation refrigeration/air-conditioning systems to this economic sector. Finally, CO<sub>2</sub> emissions from Non-  
10 Energy Uses of Fossil Fuels identified as lubricants for transportation vehicles are included in the Transportation  
11 economic sector.

12 For the Industry economic sector, the CO<sub>2</sub> emissions from the combustion of fossil fuels included in the EIA  
13 industrial fuel consuming sector, minus the agricultural use of fuel explained below, are apportioned to this  
14 economic sector. Stationary and mobile combustion emissions of CH<sub>4</sub> and N<sub>2</sub>O are also based on the EIA industrial  
15 sector, minus emissions apportioned to the Agriculture economic sector described below. Substitutes of Ozone  
16 Depleting Substances are apportioned based on their specific end-uses within the source category, with most  
17 emissions falling within the Industry economic sector (minus emissions from the other economic sectors).  
18 Additionally, all process-related emissions from sources with methods considered within the IPCC Industrial  
19 Process guidance have been apportioned to this economic sector. This includes the process-related emissions (i.e.,  
20 emissions from the actual process to make the material, not from fuels to power the plant) from such activities as  
21 Cement Production, Iron and Steel Production and Metallurgical Coke Production, and Ammonia Production.  
22 Additionally, fugitive emissions from energy production sources, such as Natural Gas Systems, Coal Mining, and  
23 Petroleum Systems are included in the Industry economic sector. A portion of CO<sub>2</sub> from Other Process Uses of  
24 Carbonates (from pollution control equipment installed in large industrial facilities) are also included in the Industry  
25 economic sector. Finally, all remaining CO<sub>2</sub> emissions from Non-Energy Uses of Fossil Fuels are assumed to be  
26 industrial in nature (besides the lubricants for transportation vehicles specified above), and are attributed to the  
27 Industry economic sector.

28 As agriculture equipment is included in EIA's industrial fuel consuming sector surveys, additional data is used to  
29 extract the fuel used by agricultural equipment, to allow for accurate reporting in the Agriculture economic sector  
30 from all sources of emissions, such as motorized farming equipment. Energy consumption estimates are obtained  
31 from Department of Agriculture survey data, in combination with separate EIA fuel sales reports. This  
32 supplementary data is used to apportion CO<sub>2</sub> emissions from fossil fuel combustion, and CH<sub>4</sub> and N<sub>2</sub>O emissions  
33 from stationary and mobile combustion (all data is removed from the Industrial economic sector, to avoid double-  
34 counting). The other emission sources included in this economic sector are intuitive for the agriculture sectors, such  
35 as N<sub>2</sub>O emissions from Agricultural Soils, CH<sub>4</sub> from Enteric Fermentation (i.e., exhalation from the digestive tracts  
36 of domesticated animals), CH<sub>4</sub> and N<sub>2</sub>O from Manure Management, CH<sub>4</sub> from Rice Cultivation, CO<sub>2</sub> emissions  
37 from Liming of Agricultural Soils and Urea Application, and CH<sub>4</sub> and N<sub>2</sub>O from Forest Fires. N<sub>2</sub>O emissions from  
38 the Application of Fertilizers to tree plantations (termed "forest land" by the IPCC) are also included in the  
39 Agriculture economic sector.

40 The Residential economic sector includes the CO<sub>2</sub> emissions from the combustion of fossil fuels reported for the  
41 EIA residential sector. Stationary combustion emissions of CH<sub>4</sub> and N<sub>2</sub>O are also based on the EIA residential fuel  
42 consuming sector. Substitutes of Ozone Depleting Substances are apportioned based on their specific end-uses  
43 within the source category, with emissions from residential air-conditioning systems to this economic sector. N<sub>2</sub>O  
44 emissions from the Application of Fertilizers to developed land (termed "settlements" by the IPCC) are also  
45 included in the Residential economic sector.

46 The Commercial economic sector includes the CO<sub>2</sub> emissions from the combustion of fossil fuels reported in the  
47 EIA commercial fuel consuming sector data. Stationary combustion emissions of CH<sub>4</sub> and N<sub>2</sub>O are also based on the  
48 EIA commercial sector. Substitutes of Ozone Depleting Substances are apportioned based on their specific end-uses  
49 within the source category, with emissions from commercial refrigeration/air-conditioning systems to this economic  
50 sector. Public works sources including direct CH<sub>4</sub> from Landfills and CH<sub>4</sub> and N<sub>2</sub>O from Wastewater Treatment and  
51 Composting are included in this economic sector.

52

**Box 2-2: Recent Trends in Various U.S. Greenhouse Gas Emissions-Related Data**

Total emissions can be compared to other economic and social indices to highlight changes over time. These comparisons include: (1) emissions per unit of aggregate energy consumption, because energy-related activities are the largest sources of emissions; (2) emissions per unit of fossil fuel consumption, because almost all energy-related emissions involve the combustion of fossil fuels; (3) emissions per unit of electricity consumption, because the electric power industry—utilities and non-utilities combined—was the largest source of U.S. greenhouse gas emissions in 2013; (4) emissions per unit of total gross domestic product as a measure of national economic activity; or (5) emissions per capita.

Table 2-15 provides data on various statistics related to U.S. greenhouse gas emissions normalized to 1990 as a baseline year. Greenhouse gas emissions in the United States have grown at an average annual rate of 0.3 percent since 1990. Since 1990, this rate is slightly slower than that for total energy and for fossil fuel consumption, and much slower than that for electricity consumption, overall gross domestic product and national population (see Table 2-15).

**Table 2-15: Recent Trends in Various U.S. Data (Index 1990 = 100)**

Chapter/IPCC Sector	1990	2005	2009	2010	2011	2012	2013	Growth <sup>a</sup>
Greenhouse Gas Emissions <sup>b</sup>	100	117	108	111	109	105	107	0.3%
Energy Consumption <sup>c</sup>	100	119	113	117	116	113	116	0.7%
Fossil Fuel Consumption <sup>c</sup>	100	119	109	113	111	108	111	0.5%
Electricity Consumption <sup>c</sup>	100	134	131	137	137	135	135	1.3%
GDP <sup>d</sup>	100	159	161	165	168	172	175	2.5%
Population <sup>e</sup>	100	118	123	124	125	125	126	1.0%

<sup>a</sup> Average annual growth rate


<sup>b</sup> GWP-weighted values

<sup>c</sup> Energy-content-weighted values (EIA 2014)

<sup>d</sup> Gross Domestic Product in chained 2009 dollars (BEA 2014)

<sup>e</sup> U.S. Census Bureau (2014)

**Figure 2-14: U.S. Greenhouse Gas Emissions Per Capita and Per Dollar of Gross Domestic Product**



Source: BEA (2014), U.S. Census Bureau (2014), and emission estimates in this report.

## 2.2 Indirect Greenhouse Gas Emissions (CO, NO<sub>x</sub>, NMVOCs, and SO<sub>2</sub>)

The reporting requirements of the UNFCCC<sup>53</sup> request that information be provided on indirect greenhouse gases, which include CO, NO<sub>x</sub>, NMVOCs, and SO<sub>2</sub>. These gases do not have a direct global warming effect, but indirectly affect terrestrial radiation absorption by influencing the formation and destruction of tropospheric and stratospheric ozone, or, in the case of SO<sub>2</sub>, by affecting the absorptive characteristics of the atmosphere. Additionally, some of these gases may react with other chemical compounds in the atmosphere to form compounds that are greenhouse gases. Carbon monoxide is produced when carbon-containing fuels are combusted incompletely. Nitrogen oxides (i.e., NO and NO<sub>2</sub>) are created by lightning, fires, fossil fuel combustion, and in the stratosphere from N<sub>2</sub>O. Non-CH<sub>4</sub> volatile organic compounds—which include hundreds of organic compounds that participate in atmospheric chemical reactions (i.e., propane, butane, xylene, toluene, ethane, and many others)—are emitted primarily from transportation, industrial processes, and non-industrial consumption of organic solvents. In the United States, SO<sub>2</sub> is primarily emitted from coal combustion for electric power generation and the metals industry. Sulfur-containing compounds emitted into the atmosphere tend to exert a negative radiative forcing (i.e., cooling) and therefore are discussed separately.

One important indirect climate change effect of NMVOCs and NO<sub>x</sub> is their role as precursors for tropospheric ozone formation. They can also alter the atmospheric lifetimes of other greenhouse gases. Another example of indirect greenhouse gas formation into greenhouse gases is CO's interaction with the hydroxyl radical—the major atmospheric sink for CH<sub>4</sub> emissions—to form CO<sub>2</sub>. Therefore, increased atmospheric concentrations of CO limit the number of hydroxyl molecules (OH) available to destroy CH<sub>4</sub>.

Since 1970, the United States has published estimates of emissions of CO, NO<sub>x</sub>, NMVOCs, and SO<sub>2</sub> (EPA 2014),<sup>54</sup> which are regulated under the Clean Air Act. Table 2-16 shows that fuel combustion accounts for the majority of emissions of these indirect greenhouse gases. Industrial processes—such as the manufacture of chemical and allied products, metals processing, and industrial uses of solvents—are also significant sources of CO, NO<sub>x</sub>, and NMVOCs.

**Table 2-16: Emissions of NO<sub>x</sub>, CO, NMVOCs, and SO<sub>2</sub> (kt)**

Gas/Activity	1990	2005	2009	2010	2011	2012	2013
<b>NO<sub>x</sub></b>	<b>21,771</b>	<b>17,394</b>	<b>13,533</b>	<b>12,709</b>	<b>12,663</b>	<b>11,958</b>	<b>11,229</b>
Mobile Fossil Fuel Combustion	10,862	10,295	7,855	7,348	7,250	6,768	6,286
Stationary Fossil Fuel Combustion	10,023	5,858	4,476	4,135	3,885	3,635	3,639
Oil and Gas Activities	139	321	469	548	626	626	626
Industrial Processes and Product Use	592	572	493	470	447	447	447
Forest Land Remaining Forest Land	64	212	149	121	373	400	149
Waste Combustion	82	128	82	78	73	73	73
Agricultural Burning	8	6	8	8	8	8	8
Waste	+	2	1	1	1	1	1
<b>CO</b>	<b>132,337</b>	<b>74,283</b>	<b>52,076</b>	<b>51,377</b>	<b>58,431</b>	<b>58,346</b>	<b>48,348</b>
Mobile Fossil Fuel Combustion	119,360	58,615	39,596	39,823	37,819	36,764	35,709
Forest Land Remaining Forest Land	2,300	7,550	5,313	4,323	13,291	14,262	5,310
Stationary Fossil Fuel Combustion	5,000	4,648	4,049	4,125	4,201	4,201	4,201
Industrial Processes and Product Use	4,129	1,557	1,332	1,280	1,233	1,233	1,233
Waste Combustion	978	1,403	1,165	1,085	1,006	1,006	1,006
Oil and Gas Activities	302	318	368	495	623	623	623
Agricultural Burning	268	184	247	241	255	253	262

<sup>53</sup> See < <http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2>>.

<sup>54</sup> NO<sub>x</sub> and CO emission estimates from Field Burning of Agricultural Residues were estimated separately, and therefore not taken from EPA (2014).

Waste	1	7	5	5	5	5	5
<b>NMVOCs</b>	<b>20,930</b>	<b>13,154</b>	<b>11,609</b>	<b>11,624</b>	<b>11,223</b>	<b>11,030</b>	<b>10,201</b>
Industrial Processes and Product Use	7,638	5,849	4,407	4,249	4,093	4,093	4,093
Mobile Fossil Fuel Combustion	10,932	5,724	4,688	4,629	4,148	3,955	3,126
Oil and Gas Activities	554	510	1,808	2,033	2,259	2,259	2,259
Stationary Fossil Fuel Combustion	912	716	554	578	603	603	603
Waste Combustion	222	241	103	92	81	81	81
Waste	673	114	49	44	38	38	38
Agricultural Burning	NA						
<b>SO<sub>2</sub></b>	<b>20,935</b>	<b>13,196</b>	<b>8,288</b>	<b>7,080</b>	<b>5,948</b>	<b>4,790</b>	<b>4,690</b>
Stationary Fossil Fuel Combustion	18,407	11,541	7,258	6,177	5,097	3,944	3,851
Industrial Processes and Product Use	1,307	831	657	623	621	621	621
Oil and Gas Activities	390	180	125	116	106	106	106
Mobile Fossil Fuel Combustion	793	619	229	148	110	104	97
Waste Combustion	38	25	17	16	15	15	15
Waste	+	1	1	+	+	+	+
Agricultural Burning	NA						

Source: (EPA 2014) except for estimates from Field Burning of Agricultural Residues.

NA (Not Available)

Note: Totals may not sum due to independent rounding.

+ Does not exceed 0.5 kt.

### Box 2-3: Sources and Effects of Sulfur Dioxide

Sulfur dioxide (SO<sub>2</sub>) emitted into the atmosphere through natural and anthropogenic processes affects the earth's radiative budget through its photochemical transformation into sulfate aerosols that can (1) scatter radiation from the sun back to space, thereby reducing the radiation reaching the earth's surface; (2) affect cloud formation; and (3) affect atmospheric chemical composition (e.g., by providing surfaces for heterogeneous chemical reactions). The indirect effect of sulfur-derived aerosols on radiative forcing can be considered in two parts. The first indirect effect is the aerosols' tendency to decrease water droplet size and increase water droplet concentration in the atmosphere. The second indirect effect is the tendency of the reduction in cloud droplet size to affect precipitation by increasing cloud lifetime and thickness. Although still highly uncertain, the radiative forcing estimates from both the first and the second indirect effect are believed to be negative, as is the combined radiative forcing of the two (IPCC 2001). However, because SO<sub>2</sub> is short-lived and unevenly distributed in the atmosphere, its radiative forcing impacts are highly uncertain.

Sulfur dioxide is also a major contributor to the formation of regional haze, which can cause significant increases in acute and chronic respiratory diseases. Once SO<sub>2</sub> is emitted, it is chemically transformed in the atmosphere and returns to the earth as the primary source of acid rain. Because of these harmful effects, the United States has regulated SO<sub>2</sub> emissions in the Clean Air Act.

Electricity generation is the largest anthropogenic source of SO<sub>2</sub> emissions in the United States, accounting for 63.0 percent in 2013. Coal combustion contributes nearly all of those emissions (approximately 92 percent). Sulfur dioxide emissions have decreased in recent years, primarily as a result of electric power generators switching from high-sulfur to low-sulfur coal and installing flue gas desulfurization equipment.