


Legend:

- | | | | |
|---------------|--|---|---------------------------|
| ⊕
% change | Location of 1994 sample. Symbol proportional to absolute mass LOSS relative to 1984. | + | NYSDEC 1984 Core Location |
| ⊙
% change | Location of 1994 sample. Symbol proportional to absolute mass GAIN relative to 1984. | X | NYSDEC 1984 Grab Location |
| ⊕ ⊙ | Be-7 detected in 1994 surface sample. | | |
| ⊕ ⊙ | Be-7 not detected in 1994 surface sample. | | |
- Marker Scale:**
- | | | |
|---------------------------------|------------------------------|------------------------------|
| NO CHANGE
0 g/m ² | LOSS
220 g/m ² | GAIN
220 g/m ² |
| ⊕ ⊙ | | |
- Sediment Classifications Based on Side Scan SONAR Images (R.Flood, 1993)**
- Rock or Rocky Sediment
 - Coarse or Coarser Sediment
 - Fine or Finer Sediment
 - Mounds (Dredge Spoils?)
 - TAMS Shoreline
 - NYSDEC Hot Spot Area



Notes:

1. Hot Spot locations were digitized from NYSDEC's "PCB Reclamation Project" Drawings (December 1985)
2. % Change is calculated as follows: $[(\text{Mass}/\text{Area}_{1994} - \text{Mass}/\text{Area}_{1984}) / \text{Mass}/\text{Area}_{1984}] * 100\%$