CASE GS0315	LINDANE	PM PM# 04/05/04
CHEM 009001	Lindane (gamm	a isomer of benzene hexac
BRANCH EEB D	DISC 40 TOPIC 05100542	
FORMULATION 00	- ACTIVE INGREDIENT	
FICHE/MASTER ID	00101191 CONTENT CA	T 01
birds of cer sings, J. S	ylor, A.; Wright, E.; et a rtain chlorinated insectic Sci, Fd Agric. 14(Aug):567 ission received on unknown ted by Shell Chemical Co.,	-577. (Also In unpub- date under unknown admin.
SUBST. CLASS =	S.	
OTHER SUBJECT D PRIM: EEB -35-0		
DIRECT RVW TIME	= L((MH) START-DATE	56/P5 END DATE 5/7/P5
REVIEWED BY: AT TITLE: AT ORG: LOC/TEL:		DATE: 5/7/05
APPROVED BY:	Section Chief EEB	
LOC/TEL:	Zenny 2. Croven	DATE: 6/6/85
SIGNATURE:	sterry 2. Crover	DATE: 6/6/63

DATA EVALUATION RECORD

- Chemical: Lindane, Dieldrin, Aldrin, Heptachlor 1.
- Test Material: Unknown percents ai 2.
- Avian Acute Oral Study Study/Action Type: 3. Pigeon (Columba livia)
- Turtle, E.; Taylor, A.; Wright E., et al., Study ID: 4. The Effects on Birds of Certain Chlorinated Insecticides used as Seed Dressings. Fd. Agric. 14 (Aug): 567-577. Submitted by Shell MRID: 00101191. Chemical Co., Washington, DC.
- Ann Stavola 5. Reviewed By:

Aquatic Biologist

HED/EEB

Signature: ONN Stevola

Date: June 5, 1965

Signature:

Date: Jerry 7. Cover

6/6/85

6. Approved By: Harry Craven

Supervisory Biologist

HED/EEB

Conclusions: 7.

> The study is not scientifically sound and does not meet our guideline requirements for an avian acute oral study.

Recommendations: 8.

N/A

Background: 9.

> This study was submitted for the data call-in for the Lindane Standard.

10. Materials and Methods:

- A. Test animals: Domestic pigeon (Columba livia); purchased from animal suppliers; of heterogeneous stock and unknown history.
- B. Dose: Lindane 290, 420, 600 mg/kg
 Dieldrin 20, 40, 80 mg/kg
 Aldrin 40, 46, 53, 61, 70, 80 mg/kg
 Heptachlor 40, 54, 73, 99, 133, 180, 243 mg/kg

All toxicants were administered as acute oral doses in gelatin capsules.

- C. Study Design: The birds were transferred to individual cages for the tests and were weighed 24 hrs prior to administration of the toxicants. The birds were deprived of food overnight before receiving the toxicants, and access to food was delayed up to 4 hrs after dosing. There were four birds/dose in the dieldrin test, and eight birds/dose for the other tests. Birds were analyzed for residues at the conclusion of the tests.
- D. Statistical Analysis: The method used to calculate the LD50 values was not given.

11. Reported Results:

Lindane	> 600
Dieldrin	67 (44-115)
Aldrin	55 (46-65)
Heptachlor	167 (133-245)

The residue analyses indicated that there was a wide variation in the distribution of residues through the flesh and organs. There was only a rough correlation between the occurrence of death and the size of a residue in the body.

The study was undertaken to ascertain if these insecticides were responsible for the deaths of a large number of wild birds in the spring in England.

12. Study Author's Conclusions/QA Measures:

The results of the residue analyses support the view that aldrin, dieldrin, and heptachlor were mainly responsible for the deaths.

The lab tests indicated that lindane has a low toxicity in comparison to the other three organochlorines.

No QA statements.

13. Reviewer's Discussion:

A. Test Procedures:

The test procedure does not follow those recommended by EPA for the following reasons: the report does not indicate the purity of the test substances; inappropriate test species; birds were not from the same stock and their histories and ages were not known; control birds were not used; small group sizes per treatment level.

B. Statistical Analysis:

Neither the raw data nor the method used % calculate the LD50 values were given. The results cannot be verified.

C. Discussion/Results:

Based upon problems with the test procedure and failure to include the raw data the results cannot be accepted.

D. Adequacy of the Study:

- 1. Conclusions: Invalid
- Rationale: Poor procedures, failure to include raw data.
- 3. Repairability: None