

**Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)**


PMRA Submission Number {.....}

EPA MRID Number 470528-14


**Data Requirement:** PMRA Data Code {.....}  
EPA DP Barcode D338050  
OECD Data Point {.....}  
EPA MRID 470528-14  
EPA Guideline OPPTS 850.1075 (72-1a)

**Test material:** Dimethyl Disulfide TC **Purity:** 998.89 g/kg (99.9%)  
**Common name:** DMDS  
**Chemical name:** IUPAC: Dimethyl Disulfide  
CAS name: Dimethyl Disulfide  
CAS No.: 624-92-0  
Synonyms: None Reported

**Primary Reviewer:** John Marton  
**Staff Scientist, Cambridge Environmental, Inc.**

**Signature:**   
**Date:** 05/21/07

**Secondary Reviewer:** Teri S. Myers  
**Senior Scientist, Cambridge Environmental, Inc.**

**Signature:**   
**Date:** 05/31/07

**Secondary Reviewer:** James Felkel *J. Felkel 10/30/08* **Date:** {.....}  
**{EPA/OECD/PMRA}**

**Secondary Reviewer(s):** {.....} **Date:** {.....}  
**{EPA/OECD/PMRA}**

**Reference/Submission No.:** {.....}

**Company Code** {.....} [For PMRA]  
**Active Code** {.....} [For PMRA]  
**Use Site Category:** {.....} [For PMRA]  
**EPA PC Code** 029088

**Date Evaluation Completed:** {dd-mm-yyyy}

**CITATION:** Scheerbaum, Dirk. 2007. Dimethyl Disulfide TC: Fish (Rainbow Trout), Acute Toxicity Test, Semi-Static, 96 H. Unpublished study performed by Dr. U Noack-Laboratorien, Kathe-Paulus-Str 1, D-31157 Sarstedt. Laboratory report number FAR106301. Study sponsored by Arkema/Thiochemistry Business Unit, Departement Securite Environnement Produit, France. Study completed January 8, 2007.

**DISCLAIMER:** This document provides guidance for EPA and PMRA reviewers on how to complete a data evaluation record after reviewing a scientific study concerning the acute toxicity of a pesticide to fish. It is not intended to prescribe conditions to any external party for conducting this study nor to establish absolute criteria regarding the assessment of whether the study is scientifically sound and whether the study satisfies any applicable data requirements. Reviewers are expected to review and to determine for each study, on a case-by-case basis, whether it is scientifically sound and provides sufficient information to satisfy applicable data requirements. Studies that fail to meet any of the conditions may be accepted, if appropriate; similarly, studies that meet all of the conditions may be rejected, if appropriate. In sum, the reviewer is to take into account the totality of factors related to the test methodology and results in determining the acceptability of the study.



2083569

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

This Data Evaluation Record may have been revised by the Environmental Fate and Effects Division subsequent to signing by Cambridge Environmental Inc. personnel.

## **EXECUTIVE SUMMARY:**

In a 96-h acute toxicity study, Rainbow Trout (*Oncorhynchus mykiss*) were exposed to Dimethyl Disulfide TC at initial measured concentrations of <0.010 (<LOQ; control), 0.303, 0.563, 1.36, 3.44 and 7.55 mg ai/L under static-renewal conditions; nominal concentrations were 0 (negative control), 0.625, 1.25, 2.5, 5 and 10 mg/L. The 96-h LC<sub>50</sub> was 0.960 mg ai/L. The NOAEC value, based on mortality and sub-lethal effects, was 0.563 mg ai/L. Sub-lethal effects (i.e., lethargy, staggering, slow escape reflex, missing escape reflex, lying on the bottom of the test vessel and hyperventilation) were observed in the groups exposed to initial measured concentrations of 1.36, 3.44 and 7.55 mg ai/L. Based on the results of this study, Dimethyl Disulfide TC would be classified as highly toxic to Rainbow Trout in accordance with the classification system of the U.S. EPA.

The study is considered scientifically sound and is classified ACCEPTABLE for an acute freshwater fish study. The NOAEC and LC<sub>50</sub> values were 0.563 and 0.960 mg ai/L, respectively.

## **Results Synopsis**

Test Organism Size/Age(mean weight or length): Mean Weight- 0.68 g; Mean Body Length- 4.60 cm  
Test Type (Flow-through, Static, Static Renewal): Static-Renewal

LC<sub>50</sub>: 0.960 mg ai/L      95% C.I.: 0.563-1.36 mg ai/L  
NOAEC: 0.563 mg ai/L      Probit Slope: N/A  
EC<sub>50</sub>: Not Determined  
Endpoint(s) Affected: Mortality and Sub-Lethal Effects

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

## I. MATERIALS AND METHODS:

**GUIDELINE FOLLOWED:** This study was reported to be conducted following guidelines outlined in EPA OPPTS Draft Guideline 850.1075 and that this guideline is "in accordance" with EC, OECD and Japanese (METI) guidelines.

There were no major deviations from OPPTS 850.1075 guidelines noted in the present review.

**COMPLIANCE:** Signed and dated No Data Confidentiality, GLP and Quality Assurance statements were provided; the study was declared to be conducted in compliance with the present OECD, EC and German principles of Good Laboratory Practice.

### **A. REPORTED MATERIALS:**

**1. Test material** Dimethyl Disulfide TC

**Description:** Light Yellow Liquid

**Lot No./Batch No. :** 17-08-04 (Batch Number)

**Purity:** 998.89 g/kg (99.9%)

**Stability of compound under test conditions:** The new solutions at 0 hours yielded recoveries of 45-76% of nominal. The aged solutions at 24 hours yielded recoveries of 64-84% of the initial measured 0 hour values. The new solutions at 72 hours yielded recoveries of 40-57% of nominal; only the three lowest treatment levels were analytically verified at 72 and 96 hours due to complete mortality at the two highest treatment levels. The aged solutions at 96 hours yielded recoveries of 70-104% of the new solutions at 72 hours.

*(OECD recommends water solubility, stability in water and light, pKa, Pow, and vapor pressure of test compound)*

**Storage conditions of test chemicals:** Stored at room temperature protected from moisture and light.

### **Physicochemical properties of Dimethyl Disulfide TC.**

Parameter	Values	Comments
Water solubility at 20EC	Insoluble @ 20°C	
Vapor pressure	20°C: 28 hPa (mbar) 25°C: 38 hPa (mbar)	
UV absorption	Not Reported	
PKa	Not Reported	

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

Parameter	Values	Comments
Kow	Not Reported	

## 2. Test organism:

**Species:** Rainbow Trout (*Oncorhynchus mykiss*) EPA recommends a cold water species (preferably rainbow trout *Oncorhynchus mykiss*) and a warm water species (preferably bluegill sunfish *Lepomis macrochirus*). OECD recommends choice of species at discretion of testing laboratory.

**Age at test initiation:** Juvenile

**Weight at study initiation:** Mean- 0.68 g, range not reported EPA recommends: mean 0.5 - 5 g.

**Length at study initiation:** Mean- 4.60 cm, range not reported; however, the study author stated that the longest fish was not more than twice the length of the shortest fish EPA recommends: Longest not > 2x shortest; OECD recommends 2.0 ± 1.0 cm for bluegill and 5.0 ± 1.0 cm for rainbow trout

**Source:** Forellenzucht Trostdadt GbR, Dofstr. 7, D-98646 Trostdadt/Thuringen  
EPA recommends that all organisms be from the same source

## B. REPORTED STUDY DESIGN:

### 1. Experimental Conditions

a. Range-finding study: A 72-hour static-renewal range-finding study was conducted using concentrations of 0 (negative control), 1, 10 and 100 mg/L. Complete mortality was observed after 24 hours at the 10 and 100 mg/L treatment levels, while no mortality occurred in the negative control or 1 mg/L treatment group.

### b. Definitive Study

Table 1: Experimental Parameters

Parameter	Details	Remarks
		Criteria
<u>Acclimation</u>		
Period:	At least 12 days	The recommended acclimation period is a minimum of 14 days; OECD guideline recommends a minimum of 12 days. Pretest mortality should be < 3% 48 h. prior to testing. OECD pretest mortality criteria: >10% = rejection of entire batch; ≥ 5 and ≤ 10% = continued acclimation for 7 days; <5% = acceptable.
Conditions: (same as test or not)	Same as test	
Feeding:	Fish were fed daily (4% of the fish body weight per feeding day) with SGP 493 (Krystal) Export. Fish were not fed within 48 hours of test initiation.	
Health: (any mortality observed)	Only healthy and normal fish were used. Mortality was <5% prior to test initiation.	
Duration of the test	96 Hours	The recommended test duration is 96 hours.

# **Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)**

PMRA Submission Number {.....}

EPA MRID Number 470528-14

Parameter	Details	Remarks
		Criteria
<u>Test condition</u>		
Static/flow-through	Static Renewal	
Type of dilution system - for flow-through method.	N/A	
Renewal rate for static renewal	Test solutions were renewed daily.	<i>A reproducible supply of toxicant is recommended. Consistent flow rate is usually 5-10 vol/24 hours; meter systems should be calibrated before and after study and checked twice daily during test period.</i>
Aeration, if any	No aeration was provided	Immediately following preparation of the test solutions, test media were treated with an ultraturrax for 1 minute at approximately 12000 rpm.
		<i>Aeration is not recommended; OECD guideline recommends aeration. If aeration is necessary, test solutions must be analyzed periodically to verify exposure.</i>

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

Parameter	Details	Remarks
		Criteria
<u>Test vessel</u>  Material: (glass/stainless steel)  Size:  Fill volume:	Glass  23 L  21 L	<p>Due to the high volatility of the test item, it was directly applied to the test vessels. The study author also reported that, with regard to the high volatility of the test item, a reduced gas phase above the water was set up.</p> <hr/> <p><i>Test vessel size is usually 19 L (5 gal) or 30 x 60 x 30 cm. Fill volume is usually 15-30 L of solution.</i></p>
Source of dilution water Quality:	Local tap water was filtered on activated charcoal and aerated for at least 24 hours to remove chlorine.	<hr/> <p><i>Recommended source of dilution water is soft, reconstituted water or water from a natural source. EPA does not recommend the use of dechlorinated tap water; however, its use may be supportable if the biological responses for the organisms and chemical analyses of residual chlorine meet conditions in the Agency's 850.1010 guidelines for dilution water (<a href="http://www.epa.gov/opptsfrs/OPPTS_Harmonized/850_Ecological_Effects_Test_Guidelines/Draft/850.1010.pdf">http://www.epa.gov/opptsfrs/OPPTS_Harmonized/850_Ecological_Effects_Test_Guidelines/Draft/850.1010.pdf</a>) Dilution water should be intensely aerated before the study. OECD permits dechlorinated tap water.</i></p>

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

Parameter	Details	Remarks
		Criteria
<u>Water parameters:</u> Hardness	60-79 mg/L as CaCO <sub>3</sub>	
pH	7.11-7.68 (New Solutions) 7.17-7.49 (Old Solutions)	<u>Hardness:</u> EPA recommends 40 - 48 mg/L as CaCO <sub>3</sub> (OECD recommends 10 - 250 mg/L)
Dissolved oxygen	82-99% DO Saturation	<u>pH:</u> EPA recommends 7.2 - 7.6; 8.0-8.3 for marine-stenohaline fishes, 7.7-8.0 for estuarine-euryhaline fishes, monthly range < 0.8; (OECD recommends pH 6.0 - 8.5)
Total Organic carbon	See Reviewer's Comments	<u>Dissolved Oxygen:</u> EPA recommends: Static: ≥ 60% during first 48 hrs and ≥ 40% during second 48 hrs; flow-through: ≥ 60%; (OECD guideline recommends at least 80% saturation value).
Particulate Matter	Not Reported	<u>Temperature:</u> EPA recommends 12 EC for coldwater species, 17 or 22 EC for warmwater species, and 22 ± 1 EC for estuarine/marine organisms. (OECD recommends 21 - 25°C for bluegill and 13 - 17°C for rainbow trout).
Metals	See Reviewer's Comments	<u>Salinity:</u> EPA recommends 30-34‰ (parts per thousand) for marine, 10-17‰ for estuarine fish, weekly range < 6‰
Pesticides	See Reviewer's Comments	Water quality should be measured at beginning of test and every 48 hours.
Chlorine	<0.01 mg/L (0 Hrs, Control)	
Temperature	11.8-12.7°C	
{Salinity for marine or estuarine species}	N/A	
Intervals of water quality measurement	Temperature, DO and pH were measured in all test vessels containing live fish before and after each renewal period. Temperature was also measured continuously in the control vessel.	
<u>Number of replicates/groups:</u> control:	1	Recommended number of replicates includes a control and five treatment levels. Each concentration should be 60% of the next highest concentration; concentrations should be in a geometric series.
solvent control:	N/A	
treated ones:	1	
<u>Number of organisms per replicate</u> <u>/groups:</u> control:	10	Number of organisms per replicate should be ≥ 10/concentration; OECD guideline recommends at least 7 fish/concentration.
solvent control:	N/A	
treated ones:	10	

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

Parameter	Details	Remarks
		Criteria
Biomass loading rate	<0.8 g/L	Recommended static conditions are # 0.8 g/L at # 17EC and # 0.5 g/L at > 17EC. Recommended flow-through conditions are # 1 g/L/day. OECD recommends a maximum of 1 g fish/L for static and semi-static, while higher rates are recommended for flow-through.
<u>Test concentrations:</u> nominal:  measured:	0 (negative control), 0.625, 1.25, 2.5, 5 and 10 mg/L  <0.010 (<LOQ; control), 0.303, 0.563, 1.36, 3.44 and 7.55 mg ai/L	Samples from the two highest treatment levels were not analyzed at 72 and 96 hours due to complete mortality. The measured concentrations from the old solutions at 24 hours yielded recoveries of 64-84% of nominal, while the measured concentrations from the aged solutions at 96 hours (from the three lowest treatment levels), yielded recoveries of 70-104% of the initial measured concentrations at 72 hours. As measured values were not determined before and after each renewal period for each treatment level, the reviewer was unable to calculate the time-weighted average or mean-measured concentrations. Further, the measured values in the aged solutions at 24 hours ranged <70% of the initial measured concentrations at 0 hours. Therefore, the reviewer reported the 0 hour measured concentrations as the "measured" values in the DER; additionally, the reviewer used these concentrations for all reporting and statistical analyses.
Solvent (type, percentage, if used)	N/A; a solvent was not used	The solvent should not exceed 0.5 ml/L for static tests or 0.1 ml/L for flow-through tests; OECD recommends that the solvent not exceed 100 mg/L.
Lighting	12L:12D with a 15-30 minute transition period	The recommended photo period is 16 hours of light and 8 hours of dark with a 15-30 minute transition period. OECD recommends a photo period of 12 -16 hours.
Feeding	Fish were not fed during the definitive test.	Fish should not feed during the study.



**Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)**

PMRA Submission Number {.....}

EPA MRID Number 470528-14

Parameter	Details	Remarks
		Criteria
<u>Recovery of chemical</u> Frequency of determination Level of quantization Level of detection	0, 24, 72 and 96 hours 0.010 mg ai/L Not Reported	
Positive control {if used, indicate the chemical and concentrations}	A positive control was not used	
Other parameters, if any	None Reported	

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

## 2. Observations:

Table 2: Observations

Parameter	Details	Remarks
		Criteria
Parameters measured including the sublethal effects/toxicity symptoms	Mortality and sub-lethal effects	
Observation intervals	6, 24, 48, 72 and 96 hours	Observation intervals should be a minimum of every 24 hours.
Were raw data included?	Yes	
Other observations, if any	None	

## II. RESULTS AND DISCUSSION:

### A. REPORTED MORTALITY:

By test termination (96 hours), mortality was 0% in the negative control and initial measured 0.303 and 0.563 mg ai/L treatment groups and 90, 100 and 100% in the initial measured 1.36, 3.44 and 7.55 mg ai/L treatment groups, respectively. Complete mortality was observed in the initial measured 3.44 and 7.55 mg ai/L treatment groups after 48 and 24 hours of exposure, respectively. The study author's analyses yielded NOAEC and LC<sub>50</sub> values of 0.541 and 0.97 (0.96-0.98) mg ai/L, respectively, based on the geometric mean measured test concentrations. The study author's NOAEC value corresponded to the initial measured concentration of 0.563 mg ai/L.

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

**Table 3: Effect of Dimethyl Disulfide on Mortality of *Oncorhynchus mykiss*.**

Initial Measured Concentrations (mg ai/L)	No. of Fish at Start of Study	Observation Period					
		Day 1		Day 3		Day 4	
		No Dead	% Mortality	No Dead	% Mortality	No Dead	% Mortality
<0.010 (Negative Control)	10	0	0	0	0	0	0
0.303	10	0	0	0	0	0	0
0.563	10	0	0	0	0	0	0
1.36	10	0	0	5	50	9	90
3.44	10	0	0	10	100	10	100
7.55	10	10	100	10	100	10	100
NOAEC*	0.541 mg ai/L						
LC <sub>50</sub> *	0.97 (0.96-0.98) mg ai/L						
Positive control, if used mortality: LC <sub>50</sub> :	N/A	N/A	N/A	N/A	N/A	N/A	N/A

\* The study author's toxicity values were derived using the geometric mean-measured concentrations. The reviewer's results are based on the initial measured concentrations.

N/A-Not Applicable as a positive control was not used.

## B. REPORTED NON-LETHAL TOXICITY ENDPOINTS:

Throughout the 96-hour exposure period, no sub-lethal effects were observed in the negative control or in the initial measured 0.303 and 0.563 mg ai/L treatment groups. The first effects were observed in the initial measured 7.55 mg ai/L treatment group after 6 hours, at which time one fish was lethargic while another was observed staggering. By 24 hours, all fish at this level were dead. After 24 hours of exposure, one fish in the initial measured 0.563 mg ai/L treatment group was lethargic and another was missing its escape reflex; complete mortality was observed by 48 hours. At 48 hours, one fish in the initial measured 1.36 mg ai/L treatment group was lethargic; at 72 hours, one fish was lethargic while the remaining live fish were either missing their escape reflex or lying on the bottom of the test vessel. The one surviving fish at this level was lethargic at test termination.

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

**Table 4: Sub-lethal Effect of Dimethyl Disulfide on *Oncorhynchus mykiss*.**

Initial Measured Concentrations (mg ai/L)	Observation Period		
	Endpoints at Day 1	Endpoints at Day 2	Endpoints at Day 4
	% Affected	% Affected	% Affected
<0.010 (Negative Control)	A.N.	A.N.	A.N.
0.303	A.N.	A.N.	A.N.
0.563	A.N.	A.N.	A.N.
1.36	A.N.	10%- Lethargic	100%- Lethargic
3.44	10%- Lethargic 10%- Missing escape reflex	--	--
7.55	--	--	--
NOAEC*	0.541 mg ai/L		
LOAEC*	1.26 mg ai/L		
EC <sub>50</sub>	Not Reported		
Positive control, if used % sublethal effect: EC <sub>50</sub> :	N/A	N/A	N/A

A.N.- All fish appear normal and healthy

\*- The study author's toxicity values were derived using the geometric mean-measured concentrations. The reviewer's results are based on the initial measured concentrations.

N/A- Not Applicable as a positive control was not used.

## C. REPORTED STATISTICS:

The LC<sub>50</sub> value was calculated by sigmoidal dose-response regression. Calculation of the confidence intervals for LC<sub>50</sub> was carried out using standard procedures according to Clopper and Pearson (1934). The concentrations leading to 0 and 100% mortality, as well as the NOAEC after 96 hours, were determined directly from the test results if applicable. All of the study author's toxicity values were determined using the geometric mean-measured concentrations.

## D. VERIFICATION OF STATISTICAL RESULTS:

Statistical Method(s): The 96-hour LC<sub>50</sub> (and 95% C.I.) was determined by analyzing the cumulative mortality using the binomial probability method via Toxanal Statistical software. The NOAEC value was visually determined based on the observed % mortality. As no quantitative sub-lethal measurements were taken, the reviewer was unable to determine an EC<sub>50</sub> value. All toxicity values were determined based on the initial measured concentrations.

LC<sub>50</sub>: 0.960 mg ai/L      95% C.I.: 0.563-1.36 mg ai/L  
 NOAEC: 0.563 mg ai/L  
 Probit Slope: N/A      95% C.I.: N/A

# Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)

PMRA Submission Number {.....}

EPA MRID Number 470528-14

## E. STUDY DEFICIENCIES:

There were no study deficiencies.

## F. REVIEWER'S COMMENTS:

The study author's 95% C.I. associated with the 96-hour  $LC_{50}$  (0.96-0.98 mg ai/L) was associated with a narrower 95% confidence interval than the reviewer's (0.563-1.36 mg ai/L); however, the study author conducted all analyses using the geometric mean-measured concentrations, while the reviewer used the initial measured concentrations due to low recoveries at 24 hours (<70% of 0-hour measurements in the nominal 0.625 and 2.5 mg/L treatment groups). Therefore, the reviewer's results are reported in the Executive Summary and Conclusions sections of this DER.

Samples from the two highest treatment levels were not analyzed at 72 and 96 hours due to complete mortality. The measured concentrations from the old solutions at 24 hours yielded recoveries of 64-84% of nominal, while the measured concentrations from the aged solutions at 96 hours (from the three lowest treatment levels), yielded recoveries of 70-104% of the initial measured concentrations at 72 hours. As measured values were not determined before and after each renewal period for each treatment level, the reviewer was unable to calculate the time-weighted average or mean-measured concentrations. Furthermore, the measured concentrations of the aged solutions at 24 hours were <70% of the initial measured concentrations at 0 hours. Therefore, the reviewer reported the 0 hour measured concentrations as the "measured" values in the DER; additionally, the reviewer used these concentrations for all reporting and statistical analyses.

An analysis of the tap water was provided in the study report; however, it is reported in German. Therefore, the reviewer is unsure about which analytes were detected and at which concentrations.

QC samples (fortified levels of 10, 100 and 100,000  $\mu$ g/L) were analyzed to determine the accuracy, precision and specificity of the analytical methodology. The mean recovery rates were 99, 102 and 89% of nominal for the 10, 100 and 100,000  $\mu$ g/L samples, respectively.

The beginning of the experimental portion of the definitive toxicity test began on September 9, 2006 and ended on September 18, 2006.

## G. CONCLUSIONS:

The study is considered scientifically sound and is classified ACCEPTABLE for an acute freshwater fish study. The NOAEC and  $LC_{50}$  values were 0.563 and 0.960 mg ai/L, respectively.

$LC_{50}$ : 0.960 mg ai/L      95% C.I.: 0.563-1.36 mg ai/L  
NOAEC: 0.563 mg ai/L      Probit Slope: N/A  
 $EC_{50}$ : Not Determined  
Endpoint(s) Affected: Mortality and Sub-Lethal Effects

## III. REFERENCES:

EPA OPPTS Draft Guideline 850.1075 (1996).

Clopper and Pearson (1934): Biometrika 26: 404-413 cited in GraphPad Prism Statistics Guide 4.0

SANCO/3029/99 rev. 4, Residues: Guidance for generating and reporting methods of analysis in support of pre-registration data requirements for Annex II (Part A, Section 4) and Annex III (part A, Section 5) of Directive 91/414 (11/07/00).

**Data Evaluation Report on the Acute Toxicity of Dimethyl Disulfide to Fish (*Oncorhynchus mykiss*)**

PMRA Submission Number {.....}

EPA MRID Number 470528-14

**APPENDIX I. OUTPUT OF REVIEWER'S STATISTICAL VERIFICATION:**

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
7.55	10	10	100	9.765625E-02
3.44	10	10	100	9.765625E-02
1.36	10	9	90	1.074219
.563	10	0	0	9.765625E-02
.303	10	0	0	9.765625E-02

THE BINOMIAL TEST SHOWS THAT .563 AND 1.36 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS, BECAUSE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS .9603242

WHEN THERE ARE LESS THAN TWO CONCENTRATIONS AT WHICH THE PERCENT DEAD IS BETWEEN 0 AND 100, NEITHER THE MOVING AVERAGE NOR THE PROBIT METHOD CAN GIVE ANY STATISTICALLY SOUND RESULTS.