

July 7/18/88

MRID 263579

DATA EVALUATION RECORD

1. **CHEMICAL:** Ortho Dibrom LVC 10.
2. **TEST MATERIAL:** Formulation: Ortho Dibrom LVC 10 (SX 1599, PN 5196); 15% as Naled technical (1,2-dibromo-2,2-dichloroethyl dimethyl phosphate); an amber-colored liquid.
3. **STUDY TYPE:** Daphnia magna 48-Hour Flow-Through Test.
Species Tested: Daphnia magna.
4. **CITATION:** Suprenant, D.C. 1986. Acute Toxicity of Ortho Dibrom LVC 10 to Daphnia magna Under Flow-Through Conditions. Prepared by Springborn Bionomics, Inc., Wareham, MA. Submitted by Chevron Environmental Health Center, Inc., Richmond, CA. Bionomics Report #EW-86-3-1952. MRID 263579. *MRID*
5. **REVIEWED BY:**

Prapimpan Kosalwat, Ph.D.
Staff Toxicologist
KEN Engineering and
Applied Sciences, Inc.

Signature: *Prapimpan Kosalwat*
Date: *5/13/88*
6. **APPROVED BY:**

Isabel C. Johnson, M.S.
Principal Scientist
KEN Engineering and
Applied Sciences, Inc.

Signature: *Isabel C. Johnson*
Date: *May 13, 1988*

for Henry T. Craven
Supervisor, EEB/HED
USEPA

Signature: *John Noles*
Date: *7/18/88*
7. **CONCLUSIONS:** This study is scientifically sound but does not meet the guideline requirements for freshwater invertebrate test. The 48-hour LC50 value of 2.9 ug/L nominal concentration classifies Ortho Dibrom LVC 10 as very highly toxic to Daphnia magna. The NOEL was 0.62 ug/L nominal concentration of Ortho Dibrom LVC 10.
8. **RECOMMENDATIONS:** N/A.

9. BACKGROUND:
10. DISCUSSION OF INDIVIDUAL TESTS: N/A.
11. MATERIALS AND METHODS:

A. Test Animals: The daphnids used in this toxicity test were obtained from laboratory cultures maintained at Springborn Bionomics, Inc., Wareham, MA. The culture water was prepared by fortifying well water based on the formula for hard water (EPA, 1975) and filtering it through a carbon filter and an Amberlite XAD-7 resin column to remove any potential organic contaminants. This water had total hardness and alkalinity ranges as CaCO_3 of 160-180 mg/L and 110-130 mg/L, respectively. Other parameters monitored were a pH range of 7.9-8.3, a temperature of $20 \pm 1^\circ\text{C}$, a dissolved oxygen concentration of greater than 60% of saturation and a specific conductance range of 400-600 micromhos per centimeter (umhos/cm).

The daphnid culture area received a regulated photoperiod of 16 hours of light and 8 hours darkness. Light at an intensity of 5-10 hectolux at the culture solution surfaces was provided by a combination of Sylvania Growlux and Cool White fluorescent bulbs. Daphnids were fed a solution of green algae (Ankistrodesmus sp. or Selenastrum sp.) and yeast suspension once daily. The ambient air temperature in the culture area was controlled in order to maintain the culture solution temperatures at $20 \pm 1^\circ\text{C}$.

B. Test System: The exposure system used in this study was a modified, proportional diluter, similar to that described by Mount and Brungs (1967) with a 0.50 dilution factor. The dilution water was from the same source as the water used in the culture vessels and was characterized as having total hardness of 160-170 mg/L as CaCO_3 , alkalinity of 120 mg/L as CaCO_3 , pH of 7.9-8.1, and a specific conductance of 500-600 umhos/cm during the study period.

The diluter delivered 5 nominal concentrations of Ortho Dibrom LVC 10, a solvent (acetone) control and a dilution water control to the test aquaria. Test vessels were glass battery jars having a volume capacity of 1.8 liters. Test solutions drained from aquaria through a 3 x 8 cm notch cut on the upper edge of the jars which maintained a solution depth of 15 cm. Drains were covered with a Nitex 40-mesh screen to prevent loss of the daphnids. Test solutions were delivered to the aquaria at a rate of 5 aquarium volumes per 24 hours. The ambient air temperature in the laboratory was controlled in order to maintain test solution temperatures at $20 \pm 1^\circ\text{C}$. Test solutions were not aerated. The test area was illuminated with Cool White and Grow Lux fluorescent lights at an intensity of 3-6 hectolux during a photoperiod of 16 hours light and 8 hours darkness.

A primary stock solution of 1130 ug/milliliter (mL) was prepared by diluting the 0.113 g of Ortho Dibrom LVC 10 with acetone to a volume of 100 mL. A secondary stock solution of 113 ug/mL was formulated by diluting 10 mL of the primary stock to 100 mL with acetone. A 50-mL gas-tight syringe with a stainless steel needle was activated during each diluter cycle by a mechanical injector to deliver 0.0345 mL of the 113 ug/mL Ortho Dibrom LVC 10 stock solution into the chemical mixing chamber of the diluter. The solution in the mixing chamber served as the highest treatment level which was subsequently diluted (0.5 dilution factor) to provide the exposure concentration range.

C. Dosage: 48-hour acute flow-through test.

D. Design: Procedures used in this acute toxicity test followed those described in the protocol entitled "Acute Toxicity of Ortho Dibrom LVC 10 to Rainbow Trout, Bluegill Sunfish and Daphnia magna in Flow-Through Test Systems," CEHC Protocol #S-2595; 1 February 1985. This protocol closely follows "Methods for acute toxicity tests with fish, macroinvertebrates, and amphibians (EPA, 1975).

Eighty daphnids, ≤ 24 hours old, were impartially distributed to each concentration (20 daphnids per replicate) at the initiation of the study. The test concentrations were 0.62, 1.2, 2.5, 5.0, and 10 ug/L as nominal concentrations of Ortho Dibrom LVC 10. All treatments and the controls were conducted in quadruplicate. The test daphnids were not fed. Biological observations and observations of the physical characteristics of each replicate test solution were also made and recorded at 0, 24, and 48 hours of exposure. The pH, dissolved oxygen concentration and temperature were measured at 0, 24, and 48 hours in one replicate vessel of all treatment levels and the controls.

Prior to initiating the definitive test, water samples (500 mL) were removed from each replicate solution of the controls, low, middle and high treatment levels two days prior to test initiation. These pretest samples were analyzed for Naled technical (active ingredient) to confirm that the proper concentration of test material was being delivered and maintained in the exposure aquaria. During the definitive test composite water samples (total volume of 500 mL) were removed from two replicates of each treatment level and the controls at 0, 24, and 48 hours. Alternate replicate solutions were sampled at each interval to provide a minimum of one analysis of each replicate.

The concentrations tested and the corresponding mortality data derived from the toxicity test were used to estimate the median lethal concentrations (LC50) and 95% confidence intervals at each 24-hour interval of the exposure period. In addition, the no-discernible-effect concentration (NDEC), defined as the highest

concentration tested at and below which there were no toxicant-related mortalities or observed behavioral and physical abnormalities (e.g., lethargy, flared carapace), was determined.

E. Statistics: A computer program by Stephan (1982) was used to calculate LC50 values. Three statistical methods, in the following order of preference, were available in the computer program: moving average angle analysis, probit analysis, and binomial probability.

12. **REPORTED RESULTS:** The method validation-recovery study conducted at Springborn Bionomics, Inc. for Naled technical in freshwater demonstrated that the minimum concentration of Naled technical which could accurately be measured in the exposure solutions was approximately 10 ug/L. During the definitive test exposing D. magna, the nominal concentration range was 0.62 to 10 ug/L of Ortho Dibrom LVC 10 (0.093 to 1.5 ug/L as Naled technical). Since the concentration of Naled technical in all exposure solutions was considerably lower than the established minimum detection limit, the amount of test material could not be analytically verified.

The diluter system which prepared and delivered the test solutions to the exposure aquaria functioned properly throughout the 48-hour study period. Daily observations of the test aquaria indicated that the test material was in solution at all treatment levels tested. The water quality parameters (pH = 7.6-7.7, dissolved oxygen = 87-93% of saturation, and temperature = 20-22°C) were unaffected by the concentrations of Ortho Dibrom LVC 10 tested.

The following table summarizes the nominal test concentrations with corresponding cumulative mortalities and observations made during the toxicity test.

Nominal Conc. of Ortho Dibrom LVC 10 (ug/L)	Cumulative Mortality (%)	
	24-hour	48-hour
10.0	100	100
5.0	60 ^a	91 ^a
2.5	0 ^a	11
1.2	0	6
0.62	0	0
Solvent Control	0	0
Control	0	0 ^b

a = Several daphnids were lethargic and on the bottom of the test vessels.

b = One daphnid was on the surface of the test solution.

The 24- and 48-hour LC50 values with their 95% confidence intervals estimated were 4.6 (2.5-10) and 2.9 (2.5-3.4) ug/L, respectively. The no-discernible-effect concentration was determined to be 0.62 ug/L of Ortho Dibrom LVC 10.

14. REVIEWER'S DISCUSSION AND INTERPRETATION OF STUDY RESULTS:

A. Test Procedure: The test procedures were in general accordance with the protocols recommended by the SEP's guidelines, except for the following deviations:

o The hardness of test water was between 160 and 170 mg/L as CaCO₃ which is much higher than the recommended water hardness of 40-48 mg/L as CaCO₃.

o There was no 15- to 30-minute transition period between light and dark as recommended by the guidelines.

o The test temperature was measured every 24 hours, instead of measuring continuously (hourly) as recommended by the guidelines.

o Each treatment concentration was only 50% of the next highest concentration. The SEP's guidelines recommend each designated

treatment group to be exposed to a concentration of toxicant that is at least 60% of the next highest concentration.

B. Statistical Analysis: Statistical analysis used by the author is appropriate and verified by the reviewer. However, only percent mortality was given in the report. The reviewer converted the percent mortality reported into number of dead animals before the statistical analysis. The resulting LC50 (3.1 ug/L with 95% C.L. of 2.8-3.4) was slightly different from that reported by the author.

C. Discussion/Results: The 48-hour LC50 of 2.9 ug/L as nominal concentration of Ortho Dibrom LVC 10 (95% confidence limits = 2.5-3.4 ug/L) is considered very highly toxic to Daphnia magna. The no-observed-effect level (NOEL) was 0.62 ug/L as nominal concentration of Ortho Dibrom LVC 10. The test concentrations could not be verified with the analytical method used.

D. Adequacy of the Study:

(1) Classification: Supplemental.

(2) Rationale: The hardness of the test water was much higher than the recommended hardness in the guidelines. The test concentrations could not be verified with the analytical method used.

(3) Repairability: No.

15. COMPLETION OF ONE-LINER: Yes, May 12, 1988.

KOSALWAT ORTHO DIBROM LVC 10 DAPHNIA MAGNA 5-6-88

CONC.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
10	80	80	100	0
5	80	73	91.25	0
2.5	80	9	11.25	0
1.2	80	5	6.25	0
.62	80	0	0	0

BECAUSE THE NUMBER OF ORGANISMS USED WAS SO LARGE, THE 95 PERCENT CONFIDENCE INTERVALS CALCULATED FROM THE BINOMIAL PROBABILITY ARE UNRELIABLE. USE THE INTERVALS CALCULATED BY THE OTHER TESTS.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 3.481689

RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	G	LC50	95 PERCENT CONFIDENCE LIMITS
4	1.120225E-02	3.109589	2.818416 3.448722

RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
6	1.105838	11.45898	0

A PROBABILITY OF 0 MEANS THAT IT IS LESS THAN 0.001.

SINCE THE PROBABILITY IS LESS THAN 0.05, RESULTS CALCULATED USING THE PROBIT METHOD PROBABLY SHOULD NOT BE USED.

SLOPE = 5.363157
95 PERCENT CONFIDENCE LIMITS = -.2766762 AND 11.00299

LC50 = 3.229266
95 PERCENT CONFIDENCE LIMITS = 0 AND +INFINITY

LC10 = 1.871988
95 PERCENT CONFIDENCE LIMITS = 0 AND 3.075873
