NOTICE OF RESEARCH PROJECT

X-TIPS

Toxicology Information Program 8600 Rockville Pike Bethesda, Maryland 20209 Tel: (301) 496-1131

SUPPORTING ORGANIZATION NUMBER(S):

SUPPORTING ORGANIZATION:

The Pennsylvania State University Agricultural Experiment Station

Contract No:

Project PA 2211 and/or

Control No:

Histopathologic and Histochemical Indices of Sublethal Pesticide Toxication in

Fish INVESTIGATOR(S) DEPARTMENT/SPECIALTY, SCHOOL OR DIVISION

Adam Anthony w. H. Neff

######!!

ii.

BAHARA

PERFORMING ORGANIZATION:

PERIOD FOR THIS NRP.

The Pennsylvania State University

Start Date: September 1976

208 Mueller Bldg.

End Date: June 1981

Name and Address. Including Zip Code. University Park, PA 16802

Annual Funding: \$15,500

Laboratory tests have been conducted to investigate histopathologic and histochemical changes in tissues of brook trout (Salvelinus fontinalis) exposed to sublethal dosages of the pesticides atrazine, carbaryl, 2,4-dichlorophenoxyacetic acid, and parathion for one week. Toxicant concentrations in ppb which kill 0% of the fish within seven days were used. Histochemical methods were selected to provide measures of cellular constitutents involved in the control or regulation of metabolism as well as end products of synthesis. Preliminary cytophotometric analyses were made of DNA and RNA changes in respiratory, hepatic, renal, and endocrine tissue. Quantification of nuclear DNA and cytoplasmic RNA content was made on an individual cell basis using scanning-integrating microdensitometry. The histopathologic analyses will be structured into a pathological index to measure tissue-specific and whole-fish pathology. The histochemical tests employed will be screened for potential short-term bioassays of toxicant effects.

CAS Registry No.

Atrazine (2-Chloro-4-ethylamino-6-isopropylamino-s-triazine) 1912-24-9 Carbaryl (1-Naphthyl N-methylcarbamate) 63-25-2 2,4-Dichlorophenoxyacetic acid 94-75-7

Parathion (Diethyl p-nitrophenyl phosphorothioate) 56-38-2

TOX-TIPS OCTOBER .1980 53-5

The bibliographic references (author, title, source) on this page have been obtained from searches of the following National Library of Medicine's on-line retrieval files: TOXLINE, CHEMLINE, MEDLINE, SDILINE, and CANCERLINE. The searches, made on the substances that are the subject of the test described in the Notice of Research Project, usually have retrieved the most recent citations from these files. Because these searches have not been exhaustive, the investigator may wish to obtain more complete searches on the test substances by searching the NLM on-line files or by contacting the Toxicology Information Response Center (TIRC) at the Oak Ridge National Laboratory, P.O. Box X, Building 20240ak Ridge, TN 37830 — Phone: 615–576–1743.

Anthony A ; Neff NH HISTOPATHOLOGIC AND HISTOCHEMICAL INDICES OF SUBLETHAL PESTICIDE TOXICATION IN FISH Toxicology Research Projects Directory, Vol.05 iss.08 1980

Bathe R; Ullmann L; Sachsse K; Hess R Relation between toxicity to fish and to mammals: a comparative study under defined laboratory conditions Proc. Eur. Soc. Toxicol.; VOL 17, ISS Predict. Chronic Toxic. Short Term Stud., Proc. Heet., 1975, 1976,351-5

Gunkel G ; Kausch H Acute toxicity of atrazine (s-triazine) on Coregonus fera Jurine under starvation conditions Arch. Hydrobiol., Suppl.; VOL 48, ISS 2, 1976,207-34

Weis P; Weis JS
Congenital abnormalities in estuarine fishes produced by environmental contaminants
Anim. Monit. Environ. Pollut., [Symp. Pathobiol. Environ. Pollut.: Anim. Models Wildl. Monit.]; 1979,94-107

建

into.

Tilak KS; Rao DM Ra; Devi AP; Murty AS Toxicity of carbaryl and 1-naphthol to the freshwater fish Labeo rohita Indian J. Exp. Biol.; VOL 18, ISS 1, 1980,75-6

Bansal SK; Verma SR; Gupta AK; Rani S; Dalela RC Pesticide-induced alterations in the oxygen uptake rate of a freshwater major carp Labeo rohita. Ecotoxicol. Environ. Saf. 3(4): 374-382 1979 (27 References)

Singh SP; Yadav NK Toxicity of some herbicides to major carp fingerlings. Indian J. Ecol. 5(2): 141-147 1978 (10 References)

Sigmon CF
Physiological effects of 2,4-D and 2,4,5T on selected aquatic organisms
Report; ISS WRRI-75, W79-05830, OWRT-A-038-SC(1); Order No. PB-294539,, 1979,30 pp.

Biro P
Acute effects of the sodium salt of 2,4-D on the early developmental stages of bleak, Alburnus alburnus.
J. Fish. Biol. 14(1): 101-109 1979 (25 References)

Kumari SDR; Nair NB Toxicity of some insecticides to Lepidocephalus thermalis (Cuv. & Val.). Proc. Indian Natl. Sci. Acad. Part B 44(3): 122-132 1978 (31 References)

Rand GM
The effect of exposure to a subacute concentration of parathion on the general locomotor behavior of the goldfish.
Bull. Environ. Contam. Toxicol. 18(2): 259-266 1977 (20
References)