

12/6/88

Accession No. 408409-04

DATA EVALUATION RECORD

1. **CHEMICAL:** Chlorpyrifos Technical
Shaughnessey No. 059101
2. **TEST MATERIAL:** Chlorpyrifos Technical; Lot No. 389318
95.9% Active Ingredient
3. **STUDY TYPE:** Flow-through Acute Toxicity Test for Freshwater
Fish. Species Tested: Lepomis macrochirus
4. **CITATION:** Bowman, Jane H. (1988); Acute Flow-through Toxicity
of Chlorpyrifos Technical to Bluegill Sunfish
(Lepomis macrochirus), Report No. 37189; prepared
by Analytical Bio-Chemistry Laboratories, Inc.
Columbia, Missouri; submitted by Makhteshim-Agan
(America) Inc. New York, New York; Accession No.
408409-04.

5. **REVIEWED BY:**

Kimberly D. Rhodes
Aquatic Toxicologist
Hunter/ESE

Signature: *Kimberly D. Rhodes*
Date:

6. **APPROVED BY:**

Prapimpan Kosalwat, Ph.D.
Staff Toxicologist
KBN Engineering and
Applied Sciences, Inc.

Signature: *P. Kosalwat*
Date: *11/28/88*

for Henry T. Craven, M.S.
Supervisor, EEB/HED
USEPA

Signature: *John Noles*
Date: *12/6/88*

7. **CONCLUSIONS:** This study appears scientifically sound and
fulfills the Guideline requirements for a 96-hour acute flow-
through toxicity test of freshwater fish. The 96-hour LC50
based upon mean measured concentrations of Chlorpyrifos
Technical to bluegill sunfish (Lepomis macrochirus) was
~~0.0062 mg/L~~, which classifies it as very highly toxic to
bluegill sunfish. The NOEC was determined to be 0.0011 mg/L
after 96 hours.

Best CLS
5.8 (4.7 - 7.5) ppb

8. **RECOMMENDATIONS:** N/A

B. Montague

9. BACKGROUND:

10. DISCUSSION OF INDIVIDUAL TESTS: N/A

11. MATERIALS AND METHODS:

A. Test Animals: Bluegill sunfish (Lepomis macrochirus) were obtained from a commercial supplier in Missouri. The fish were reared and maintained at ABC Laboratories in ABC well water and were fed newly hatched brine shrimp or a commercially available fish food daily. Forty-eight hours before the initiation of the test, bluegill sunfish were removed from the culture and placed in the temperature acclimation unit. During this time, the fish were held without food. The bluegill used as the control group during this study had a mean weight of 2.1 (± 0.87) grams and a mean length of 41 (± 5.5) millimeters at test termination. The biomass loading rate was 1.4 g/L. The laboratory environment was maintained on a 16-hour daylight photoperiod.

B. Test System: A proportional diluter system described by Mount and Brungs, utilizing a Hamilton Micro Lab 420 syringe dispenser, was used for the intermittent introduction of Chlorpyrifos Technical test solutions and diluent water into each test chamber. The proportional diluter system used for the project was set to provide test levels approximately 50 percent dilutions of each other. The diluter delivered one liter of test solution or control water to the test vessels at an average rate of 9.5 times per hour over the course of the study. This flow rate was sufficient to replace the 30-liter volume within the test chambers 7.6 times per day. Five concentrations of the test material with dilution water control and solvent controls were tested. The test chambers were immersed in a temperature controlled water bath held at $22 \pm 1^\circ\text{C}$. The lighting was maintained on a 16-hour daylight photoperiod.

Dilution water for the bluegill sunfish test was a blend of reverse osmosis water and ABC well water characterized as having a pH of 7.1 - 7.9, total hardness of 40 - 48 mg/L as CaCO_3 , total alkalinity of 44 - 56 mg/L as CaCO_3 and specific conductance of 100 - 160 umhos/cm.

C. Dosage: 96-hour flow-through acute test.

D. Design: Two static range-finding tests were conducted with Chlorpyrifos. Based on the results of the preliminary testing five concentrations were selected for

definitive testing. Twenty bluegill sunfish were tested per concentration. A control, solvent control, and nominal Chlorpyrifos Technical concentrations of 0.020, 0.010, 0.0050, 0.0025, and 0.0013 mg/L were tested. The nominal concentrations were not corrected as active ingredient. The concentration of acetone in the solvent control (0.03 mL of acetone per liter of water) was approximately equivalent to that received by the highest test concentration. The mean measured concentrations of Chlorpyrifos Technical were 0.0011, 0.0020, 0.0038, 0.0087 and 0.018 mg/L as active ingredient. All concentrations were observed once every 24 hours for mortality and abnormal effects. The water quality parameters (temperature, dissolved oxygen and pH) were measured in the control, low, middle, and high test levels at 0, 48, and 96 hours of testing. Analytical samples were collected from each test level and the diluter stock at 0 and 96 hours.

- E. Statistics: The concentration of toxicant lethal to 50% of the population (LC50's) and 95% confidence intervals was determined at 24-, 48-, 72-, and 96-hour exposure periods by the computer program developed by Stephan et al., 1978.
12. REPORTED RESULTS: The 24-, 48-, 72-, and 96-hour LC50 values for Chlorpyrifos were >0.018 (although >0.0087 was reported), 0.0081, 0.0073 and 0.0058 mg/L, respectively, based upon mean measured concentrations. The slope of the 96-hour dose-response line was 3.7 as calculated by probit computer analysis. Behavioral/sublethal effects, observed in all concentrations except the lowest (0.0011 mg/L), included surfacing, labored respiration, fish on bottom of test chamber, loss of equilibrium, quiescence, forward curved pectoral fins and light discoloration. Due to the lack of mortality or behavioral/sublethal effects in the lowest concentration a no-effect concentration of Chlorpyrifos Technical to bluegill sunfish was determined to be 0.0011 mg/L. Table 5 presents mortality and behavioral observations made during the test (attached). There was no mortality in the control or solvent control during the study. The dissolved oxygen concentrations ranged between 7.4 and 8.4 mg/L (88 to 100% saturation at 22°C) during the test.
13. STUDY AUTHOR'S CONCLUSIONS/QUALITY ASSURANCE MEASURES: The 96-hour LC50 value for Chlorpyrifos Technical was calculated by moving average method to be 0.0058 mg/L with 95 percent confidence limits of 0.0047 and 0.0075 based upon mean measured concentrations. The NOEC (No-Observed-Effect Concentration) was 0.0011 mg/L after 96 hours.

"The study was conducted following the intent of the Good Laboratory Practice Regulations and the final report was reviewed by Analytical Bio-Chemistry Laboratories' Quality Assurance Unit." A Quality Assurance Statement was included in the report.

14. REVIEWER'S DISCUSSION AND INTERPRETATION OF STUDY RESULTS:

- A. Test Procedure: The test procedures were generally in accordance with protocols recommended by the Guidelines, but deviated from the SEP as follows:
- o The SEP recommends that fish be acclimated to study conditions for at least two weeks prior to testing. The bluegill sunfish were removed from the culture tank and placed in the temperature acclimation unit forty-eight hours before the acclimation of the test.
 - o Six-hour temperature measurements were not recorded as required by the SEP for tests conducted in a water bath.
 - o The SEP states that each designated treatment group should be exposed to a concentration of toxicant that is at least 60% of the next highest concentration. Each designated treatment group for the test was only 50% of the next highest concentration.
 - o The SEP recommends a 16-hour light and an 8-hour dark photoperiod with a 15- to 30-minute transition period between light and dark. The report did not state whether a 15- to 30-minute transition period between light and dark was maintained.
- B. Statistical Analysis: The reviewer used the Toxanal computer program to calculate the LC50 values. These calculations are attached. The probit method provides a 96-hour LC50 value of 0.0062 mg/L with a 95 percent confidence interval of 0.0049 to 0.0079 mg/L which is similar to that reported by the author. The slope of the toxicity curve was estimated to be 3.7.
- C. Discussion/Results: The study results appear to be scientifically valid. The 96-hour LC50 value based upon mean measured concentrations was estimated to be 0.0062 mg/L. Therefore, Chlorpyrifos Technical is classified as very highly toxic to bluegill sunfish (Lepomis macrochirus).

D. Adequacy of the Study:

- (1) Classification: Core
- (2) Rationale: N/A
- (3) Repairability: N/A

15 COMPLETION OF ONE-LINER FOR STUDY: Yes, 11-17-88.

KIMBERLY SWOODED ONLOPPYRATES TECHNICAL LECTURE MICROCOMPUTER 11 27 80

COND.	NUMBER EXPOSED	NUMBER DEAD	PERCENT DEAD	BINOMIAL PROB. (PERCENT)
.018	20	20	100	9.536742E-05
.0087	20	12	60.00001	25.17223
.0038	20	5	25	2.049473
.002	20	1	5	2.002716E-03
.0011	20	0	0	9.536742E-05

THE BINOMIAL TEST SHOWS THAT .0038 AND .018 CAN BE
USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT
CONFIDENCE LIMITS. BECAUSE THE ACTUAL CONFIDENCE LEVEL
ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 6.907752E-03

RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

SPAN	S	LC50	95 PERCENT CONFIDENCE LIMITS
4	4.730058E-02	5.860047E-03	4.696116E-03 7.609292E-03

RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	S	H	GOODNESS OF FIT PROBABILITY
4	.1033621	1	.5128763

SLOPE = 3.661161
95 PERCENT CONFIDENCE LIMITS = 2.484099 AND 4.838223

LC50 = 6.172224E-03
95 PERCENT CONFIDENCE LIMITS = 4.90955E-03 AND 7.911971E-03

LC10 = 2.785877E-03
95 PERCENT CONFIDENCE LIMITS = .0017897 AND 3.654275E-03

Page 7 is not included in this copy.

Pages _____ through _____ are not included.

The material not included contains the following type of information:

- ☐ Identity of product inert ingredients.
 - ☐ Identity of product impurities.
 - ☐ Description of the product manufacturing process.
 - ☐ Description of quality control procedures.
 - ☐ Identity of the source of product ingredients.
 - ☐ Sales or other commercial/financial information.
 - ☐ A draft product label.
 - ☐ The product confidential statement of formula.
 - ☐ Information about a pending registration action.
 - ☒ FIFRA registration data.
 - ☐ The document is a duplicate of page(s) _____.
 - ☐ The document is not responsive to the request.
-

The information not included is generally considered confidential by product registrants. If you have any questions, please contact the individual who prepared the response to your request.
