#### DATA EVALUATION RECORD

CHEMICAL: Triphenyltin Hydroxide (TPTH) 1.

2. Triphenyltin (TPTH) Technical, Lot#RFRAM909K

Purity: 97.23%

STUDY TYPE: §72-3 (c) Acute toxicity test for estuarine and 3...

marine organisms: Mysid Shrimp (Mysidopsis bahia)

CITATION: 4.

Author:

Machado, M. W.

Title:

Triphenyltin (TPTH) - Acute Toxicity to Mysid Shrimp

(Mysidopsis bahia) under Flow-

Through Conditions

Date:

March 10, 1994

Laboratory Report #:

94-1-5124

Any Other Study #:

11117.0593.6103.515

Sponsor:

Griffin Corp. Valdosta, GA

Sponsor #:

N/A

Laboratory:

Springborn Laboratories, Inc.

MRID No.:

43212701

5. REVIEWED BY:

Dennis J. McLane, Wildlife Biologist

Signature:

Ecological Effects Branch

Environmental Fate and Effects Division (7507 C)

APPROVED BY: 6.

Les Touart, Chief, Section 1

Signature:

Ecological Effects Branch

Environmental Fate and Effects Division (7507C)

Date: 9-12-14

CONCLUSION This study fulfills the guideline requirements. The reported moving average angle was 4.3 (3.7-5.0)  $\mu g/L$ which will place TPTH in the very highly toxic category (<100  $\mu g/L$ ).

- 8. RECOMMENDATIONS N/A
- 9. BACKGROUND Submitted in response to the list A process.
- 10. MATERIALS AND METHODS
  - A. Test Organisms: Mysid Shrimp

Guideline Criteria	Reported Information		
Species (Scientific Name)	Mysidopsis bahia		
Mean Weight (> 0.5 grams)	0.001 g		
Supplier	Springborn Laboratories		
All shrimp from same source (yes or no)	yes		
All shrimp from the same year class (yes or no)	yes ≤24 hours old		
Other Comments			

# B. <u>Source/Acclimation</u>

Guideline Criteria	Reported Information		
Acclimation Period (minimum 10 days)	Cultured at this laboratory		
Wild caught 7 day quarantine (yes or no)	no		
Check for signs of disease or injury (yes or no, if yes describe)	Not reported		
If diseased it can be treated in 48-hr pretest no sign of the disease remains (Report hours prior to test in which no sign of disease or N/A)	N/A		
No feeding during the study (When last fed)	twice daily		
<3% mortality 48 hours prior to testing (% mortality, if any)	Not reported		

## C. <u>Test System</u>:

Guideline Criteria	Reported Information
Describe source of dilution water	Cape Cod Canal, Bourne, Massachusetts

A			
Does water support test animals without observable signs of stress?	Yes		
What was the salinity of the water used? (30-34% ppt for marine (stenohaline) shrimp and 10- 17% ppt for estuarine (euryhaline) shrimp.	3 2‰		
Water Temperature (22°C)	25°C		
pH 8.0-8.3 marine (stenohaline) shrimp 7.7-8.0 estuarine (euryhaline) shrimp	7.7-8.0		
Dissolved Oxygen (Static 1 <sup>st</sup> 48 hrs 40%; 2 <sup>nd</sup> 48 hrs 60%; Flow-through 60%) (% of lowest conc. & hour)	90-103%		
Total Organic Carbon	1.1-1.5 mg/l		
Test Aquaria 1. Material (glass or stainless steel) 2. a. Static volume (18.9 L (5 gal or 19000 cc) with 15 L solution) b. Static or flow-through volume (300x600x300 = 54000 cc.)	<ul> <li>1. glass and silicone sealant</li> <li>a. N/A</li> <li>b. 39 x 20 x 25 cm</li> <li>depth limited by a 14.5 cm</li> <li>standpipe for a volume of 11 L</li> </ul>		
Type of Dilution System (Reproducible supply of toxicant)	modified serial diluter system		
Flow rate Consistent flow rate-meter systems calibrated before study and checked 2*24 hours - 5 to 10 vol/24 hours	6.5 volume replacements per aquarium every 24 hours		

Biomass Loading Rate (Static no > 0.8 g/L ≤ 17°C; >17°C 0.5 g/L; Flow-through 1 g/L/24	0.00014 g
Photoperiod (16 L & 8 D)	16 L & 8 D
Solvents 1. (Do not exceed 0.5 ml/L for static tests) 2. (Do not exceed 0.1 ml/L for	1. N/A 2. 0.1 ml/L
flow-through) Other Comments	

## D. <u>Test Design</u>:

Guideline Criteria	Reported Information		
Range Finding Test (LC <sub>50</sub> >100 mg/L with 30 shrimp, no definitive test required.)	Mortality at 11 $\mu$ g/L		
<u>Definitive Test</u>			
Nominal Concentrations (control+5 treatment levels; dosage should be 60% of the next highest concentration; concentrations should be geometric series)	control, solvent control, 1.6, 2.6, 4,3, 7.2, and 12 $\mu g/L$		
Controls (Minimum control mortality; static 10%; flow-through 5%	0% mortality for both control and solvent control		
Number of Test Organisms; (Minimum 20/level can be divided among containers)	20/level		
All organisms must be randomly assigned to test vessels. (yes or no, describe if no)	"impartially selected and distributed"		
Biological Observations (yes or no)	Yes		

Water Parameter Measurements 1. Temperature - record every 6 hrs;>1°C.  2. D.O. beginning,48 hrs,end for control high, medium, and low dose. 3. pH beginning,48 hrs; end for control, high, medium, and low dose.	1. Continuous in replicate A range 24-25°C, others were daily and 25°C 2.daily in each aquarium 3.daily in each aquarium
Chemical Analysis (needed if aeration, volatile, insoluble, precipitate, not steel or glass, known to adsorb, and flow-through) (yes)	yes
Other Comments	

#### 11. REPORTED RESULTS:

Guideline Criteria	Reported Information		
Mean Measured Concentrations (report conc.)	1.4, 2.2, 3.8, 6.3, 10		
Recovery of Chemical (% recovery)	90, 83, 89, 88, 87		
Mortality & Observations (Describe observations & attach mortality tables)	See attached Table 3 for both mortality and observations		
Author's Comments			

12. STUDY AUTHOR'S CONCLUSIONS / QUALITY ASSURANCE MEASURES:
Author provided no conclusions. Quality assurance unit statement
was provided and signed by the Patricia D. Royal the Regulatory
Affairs and Quality Assurance Manager. Also a good laboratory
practice compliance statement was include with signature from the
Study director, Sponsor study monitor, and Applicant/Submitter.

#### 13. REVIEWER'S DISCUSSION AND INTERPRETATION

#### A. Test Procedure:

The following items did not meet the guideline

#### criteria:

- 1. Rather than randomly assigned to test vessels the shrimp were "impartially selected and distributed to test vessels.
  - 2. Disease in the shrimp was not discussed.
- 3. The percent mortality 48 prior to the test was not reported.

B. Statistical Analysis

Guideline Criteria	Reported Information		
Binomial (yes, no, or not reported)	No		
Moving Average Angle (yes, no, or not reported)	Yes, 4.3 (3.7-5.0) μg/L		
Probit (yes, no, or not reported)	No		
Other Comments	EEB used the Toxanal program which gives $LC_{50}$ values from binomial, moving average, and probit methods. They were as follows: binomial 4.7 (3.8-6.3) $\mu$ g/L, moving average 4.57 (3.96-5.336) $\mu$ g/L, and probit 4.74 (4.26-5.31) $\mu$ g/L. (see attached printout).		

C. <u>Discussion/Results</u>: The items under 13. <u>REVIEWER'S</u>
<u>DISCUSSION AND INTERPRETATION</u> are not expected in this case to effect the results of the study.

#### D. Adequacy of the Study:

- 1. Classification: Core
- 2. Rational: Fulfills guideline requirements.
- 3. Reparability: N/A

#### 14. COMPLETION DATE OF ONE-LINER FOR STUDY:

MCLANE TPTH MYSID 96-HOUR FLOW-THROOUGH ACUTE

\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*\*

CONC.	NUMBER	NUMBER	PERCENT	BINOMIAL
	EXPOSED	DEAD	DEAD	PROB. (PERCENT)
10	20	20	100	9.536742E-05
6.3	20	19	95	2.002716E-03
3.8	20	2	10	2.012253E-02
2.2	2.0	0	0	9.536742E-05
1.4	2.0	0	0	9.536742E-05

THE BINOMIAL TEST SHOWS THAT 3.8 AND 6.3 CAN BE USED AS STATISTICALLY SOUND CONSERVATIVE 95 PERCENT CONFIDENCE LIMITS, BECAUSE THE ACTUAL CONFIDENCE LEVEL ASSOCIATED WITH THESE LIMITS IS GREATER THAN 95 PERCENT.

AN APPROXIMATE LC50 FOR THIS SET OF DATA IS 4.785323

RESULTS CALCULATED USING THE MOVING AVERAGE METHOD

95 PERCENT CONFIDENCE LIMITS G LC50 SPAN 4 4.557268E-02

G

4.573244 3.968218

5.360389

RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS

GOODNESS OF FIT PROBABILITY

.1649973

.9999995

13.33154 SLOPE

95 PERCENT CONFIDENCE LIMITS = 7.91629 18.7468 AND

LC50 =4.74165

95 PERCENT CONFIDENCE LIMITS = 4.263238 AND 5.318471

3.807736

95 PERCENT CONFIDENCE LIMITS = 3.163379 AND 4.238397

\*