

DATA EVALUATION RECORD

1. CHEMICAL: Cuprous oxide plus triphenyltin hydroxide
2. FORMULATION: Rabamarine, which is a mixture of 2 formulated products and contains 36.5% cuprous oxide and 8.7% triphenyltin hydroxide.
3. CITATION: Sousa, J. (1981) Acute Toxicity of Kansai Rabamarine to the Rainbow Trout; received 7/21/81 under 46197-1; unpublished report prepared by E G & G Bionomics for M & T Chemicals, Inc, Rahway, New Jersey (in Acc # 245649).
4. REVIEWED BY: Stephen M Hopkins
Plant Physiologist
EEB/HED
5. DATE REVIEWED: 9/25/81
6. TEST TYPE: Fish acute LC₅₀ -- Rainbow trout
7. REPORTED RESULTS: The testing laboratory demonstrated that the 96hr LC₅₀ of Rabamarine to the rainbow trout is 120 ppb product, with 95% confidence limits of 99-160 ppb. This 120 ppb of product contains approximately 44 ppb of cuprous oxide and 10 ppb of triphenyltin hydroxide.
8. REVIEWER'S CONCLUSIONS: This study is scientifically sound, and meets EPA requirements for a fish acute LC₅₀ study using the formulated product.

Testing Laboratory Report

A. Test Procedure

The procedure generally followed the EPA proposed guidelines of July 10, 1978. Some specifics of note include:

Weight of fish - 0.98 g average
Number of fish - 10 per treatment level
Test vessel size - 20 liter glass jars containing 15 liters each
Temperature - $12 \pm 1^{\circ}\text{C}$
Loading - 0.66g/liter
Dilution water - Reconstituted deionized water
Treatment levels - 36, 60, 100, 170, and 280 ppb plus untreated and N,N-dimethylformamide controls
Test initiation - February 10, 1981
Test material - The test material was a mixture containing:
16 parts of solution A (57% cuprous oxide), and
9 parts of solution B (23.8% triphenyltin hydroxide).
The final mixture contained 36.5% cuprous oxide and 8.7% triphenyltin hydroxide.

B. Statistical Analysis

Mortality was analyzed by the moving average angle method.

C. Results

<u>Concentration</u>	<u>Mortality at 96hrs</u>
280 ppb product	100%
170	80
100	30
60	0
36	0
controls	0

The author calculated that the 96hr LC₅₀ of Rabamarine to the rainbow trout is 120 ppb product, with 95% confidence limits of 99-160 ppb. 60 ppb was a no-effect level.

Reviewer's Evaluation

A. Test Procedure

The procedure generally complied with the 1978 EPA guidelines.

B. Statistical Analysis

Mortality was analyzed by the probit method, the results of which agreed with the findings of the testing laboratory.

C. Results/Discussion

The testing laboratory demonstrated that the 96hr LC₅₀ of Rabamarine to the rainbow trout is 120 ppb product, with 95% confidence limits of 99-160 ppb. This 120 ppb of product contains approximately 44 ppb of cuprous oxide and 10 ppb of triphenyltin hydroxide.

D. Conclusion

1. Category: Core for formulated product
2. Rationale: NA
3. Repairability: NA