

Shaughnessy No.: 090501

Date out EAB: 04 APR 1984

To: Taylor
Product Manager 25
Registration Division TS-767

From: Samuel M. Creeger, Chief
Environmental Chemistry Review Section 1
Exposure Assessment Branch
Hazard Evaluation Division TS-769

Attach, please find the Eab Review of:

Reg./File No.: 324-316

Chemical: Alachlor

Type Product: H

Product Name: Lasso

Company Name: Monsanto

Submission Purpose: Exposure Assessment

ZBB Code: other

Action Code: 401

Date In: 3/1/84

EAB No.: 4199

Date Completed: 4/4/84

TAIS (Level II) Days

Deferrals To:

62 7

Ecological Effects Branch

Residue Chemistry Branch


X Toxicology Branch

I. INTRODUCTION

Monsanto has resubmitted data previously generated as a result of a field study conducted using alachlor formulations. This submission contains recalculated exposure numbers taking into account recovery data and a 70 kg body weight (rather than 60 kg). The EPA Accession No. of the original submission is 070591 (EAB review dated 5/25/82) and of the current submission is 252498.

II. Alachlor: Lasso

2-chloro-2',6'-diethyl-N-(methoxymethyl)-acetanilide



III. DISCUSSION

Because of the controversy surrounding any previously submitted exposure assessments, EAB has returned to the original study and recalculated potential exposures to personnel involved in the use of alachlor.

A major item of controversy was the fact that the registrant failed to correct his raw data for recovery. Consequently EAB in its current review of the data did take into account these values. From the raw data submitted in the original study by the registrant, average recovery values from all field or lab studies conducted are as follows:

	<u>Field</u>	<u>Lab</u>
<u>Dermal</u>		
gauze pads	86%	92%
cotton gloves	74	74
rubber gloves		87
<u>Respiratory</u>		
foam plugs	96	95
silica gel	89	96
charcoal	77	76

For this current exposure assessment, EAB chose to use for gauze pads a recovery value of 89% and for cotton gloves, 74%. Exposure to hands as determined only from the residue values found on cotton gloves (worn under rubber gloves) was considered. Respiratory exposure values were used as submitted by the registrant in this submission since foam and silica gel were the trapping media and recovery values are excellent for both.

The assumptions and steps used to determine the exposure values are:

- o The average of the residue values from two reps is used.
- o All residue values are corrected for recovery as explained above.
- o Dermal and respiratory absorption is 100%.
- o Three different exposure scenarios are considered:
 - No Protection: This assumes the person is naked with the exception of wearing rubber gloves. Residue values from pads located in the following body regions were used: hands (cotton gloves), forehead, chest, back, thigh, forearm (outer), bicep (outer), and ankle (outer).
 - With Protection (80%): This assumes the person is wearing protective clothing such as coveralls. (Normal work clothes are not considered to be protective clothing.) Residue values from the same regions as with No Protection are used but now it is assumed that the protective clothing reduces exposure by 80%. It is still assumed that the worker is wearing rubber gloves so that his hand exposure (cotton gloves) is not further reduced by 80%.

It is believed this scenario best represents exposure to alachlor under normal working conditions.
 - With Protection (100%): This assumes the person is wearing the same protective clothing as above but that now the clothing reduces exposure by 100%. Only the face, back of the neck, front of neck and "V" of the chest, and the hands (cotton gloves) are exposed to the active ingredient.
- o From information presented by the registrant at various meetings and supported generally by BUD, the following assumptions were used in appropriate calculations:
 - A private farmer may treat a 100 acre plot/day and may have up to 600 acres to treat. Consequently he may be exposed to alachlor from 1 - 6 days/year.
 - A custom applicator may treat a 100 acre plot/day but may be exposed to alachlor from 5 - 30 days/year.
 - An aerial applicator, used only in emergency situations, may treat up to 1000 acres/day and may be exposed to alachlor from 5 - 10 days/year.
 - For Lifetime exposure calculations, a 40 year working lifetime is assumed during a 70 year lifespan.

All appropriate tables, abstracted from the registrant's study, and calculations used by EAB to determine the potential exposure to alachlor are attached to this review.

ALACHLOR (LASSO EC) EXPOSURE

<u>Operation</u>	<u>DERMAL EXPOSURE</u>		<u>RESPIRATORY EXPOSURE</u>
	<u>No Protection</u>	<u>With Protection (80%)</u>	<u>With Protection (100%)</u>
Ground Open Tank Fill	0.39 mg/kg	0.091 mg/kg	0.023 mg/kg
	110. mg/hr	26. mg/hr	6.9 mg/hr
	1.6 mg/kg/hr	0.38 mg/kg/hr	0.099 mg/kg/hr
	0.00096 mg/kg/lb	0.0002 mg/kg/lb	0.000059 mg/kg/lb
	0.38 mg/kg/day	0.09 mg/kg/day	0.024 mg/kg/day
	0.38 - 2.28 mg/kg/yr (Private)	0.09 - 0.54 mg/kg/yr (Private)	0.024 - 0.14 mg/kg/yr (Private)
	0.00061-0.0036 mg/kg/day (Lifetime-Private)	0.00014 - 0.00086 mg/kg/day (Lifetime-Private)	0.000038 - 0.00022 mg/kg/day (Lifetime-Private)
	1.9 - 11.4 mg/kg/yr (Custom)	0.45 - 2.7 mg/kg/yr (Custom)	0.12 - 0.72 mg/kg/yr (Custom)
	0.003 - 0.018 mg/kg/day (Lifetime-Custom)	0.00072 - 0.0043 mg/kg/day (Lifetime-Custom)	0.00019 - 0.0012 mg/kg/day (Lifetime-Custom)
			0.000065 - 0.0004 mg/kg/yr

ALACHLOR (LASSO EC) EXPOSURE

<u>Operation</u>	<u>DERMAL EXPOSURE</u>		<u>RESPIRATORY EXPOSURE</u>
	<u>No Protection</u>	<u>With Protection (80%)</u>	<u>With Protection (100%)</u>
Ground Applicator	0.019 mg/kg	0.0074 mg/kg	0.0048 mg/kg
	1.2 mg/hr	0.47 mg/hr	0.3 mg/hr
	0.017 mg/kg/hr	0.0067 mg/kg/hr	0.0043 mg/kg/hr
	0.000046 mg/kg/lb	0.000018 mg/kg/lb	0.000012 mg/kg/lb
	0.018 mg/kg/day	0.0073 mg/kg/day	0.0047 mg/kg/day
	0.018 - 0.11 mg/kg/yr (Private)	0.0073 - 0.044 mg/kg/yr (Private)	0.0047 - 0.028 mg/kg/yr (Private)
	(2.9-18.)x10 ⁻⁵ mg/kg/day (Lifetime-Private)	(1.2 - 7.) x 10 ⁻⁵ mg/kg/day (Lifetime-Private)	(0.75 - 4.5) x 10 ⁻⁵ mg/kg/day (Lifetime-Private)
	0.09 - 0.54 mg/kg/yr (Custom)	0.036 - 0.22 mg/kg/yr (Custom)	0.024 - 0.14 mg/kg/yr (Custom)
	(14.-86.)x10 ⁻⁵ mg/kg/day (Lifetime-Custom)	(5.8 - 35.) x 10 ⁻⁵ mg/kg/day (Lifetime-Custom)	(3.8 - 22.) x 10 ⁻⁵ mg/kg/day (Lifetime-Custom)
			0.00003 - 0.00018 mg/kg/yr

ALACHLOR (LASSO EC) EXPOSURE

<u>Operation</u>	<u>DERMAL EXPOSURE</u>		<u>RESPIRATORY EXPOSURE</u>
	<u>No Protection</u>	<u>With Protection (80%)</u>	<u>With Protection (100%)</u>
Ground Probe Transfer 5 gal	0.02 mg/kg	0.0042 mg/kg	0.0056 mg/kg
	5.5 mg/hr	1.1 mg/hr	1.5 mg/hr
	0.079 mg/kg/hr	0.016 mg/kg/hr	0.021 mg/kg/hr
	0.000051 mg/kg/lb	0.00001 mg/kg/lb	0.000014 mg/kg/lb
	0.02 mg/kg/day	0.0042 mg/kg/day	0.0055 mg/kg/day
	0.02 - 0.12 mg/kg/yr (Private)	0.0042 - 0.025 mg/kg/yr (Private)	0.0055 - 0.033 mg/kg/yr (Private)
	(3.2-19.)x10 ⁻⁵ mg/kg/day (Lifetime-Private)	(0.67 - 4.) x 10 ⁻⁵ mg/kg/day (Lifetime-Private)	(0.88 - 5.3) x 10 ⁻⁵ mg/kg/day (Lifetime-Private)
	0.1 - 0.6 mg/kg/yr (Custom)	0.021 - 0.13 mg/kg/yr (Custom)	0.028 - 0.16 mg/kg/yr (Custom)
	(16.-96.)x10 ⁻⁵ mg/kg/day (Lifetime-Custom)	(3.4 - 21.) x 10 ⁻⁵ mg/kg/day (Lifetime-Custom)	(4.5 - 26.) x 10 ⁻⁵ mg/kg/day (Lifetime-Custom)

ALACHLOR (LASSO EC) EXPOSURE

<u>Operation</u>	<u>DERMAL EXPOSURE</u>		<u>RESPIRATORY EXPOSURE</u>
	<u>No Protection</u>	<u>With Protection (80%)</u>	<u>With Protection (100%)</u>
Ground Probe Transfer 55 gal	0.18 mg/kg	0.038 mg/kg	0.017 mg/kg
	65 mg/hr	13 mg/hr	5.9 mg/hr
	0.92 mg/kg/hr	0.19 mg/kg/hr	0.084 mg/kg/hr
	0.00046 mg/kg/lb	0.000095 mg/kg/lb	0.000042 mg/kg/lb
	0.18 mg/kg/day	0.038 mg/kg/day	0.017 mg/kg/day
	0.18 - 1.1 mg/kg/yr (Private)	0.038 - 0.23 mg/kg/yr (Private)	0.017 - 0.1 mg/kg/yr (Private)
	(29.-176.)x10 ⁻⁵ mg/kg/day (Lifetime-Private)	(6.1 - 37.) x 10 ⁻⁵ mg/kg/day (Lifetime-Private)	(2.7 - 16.) x 10 ⁻⁵ mg/kg/day (Lifetime-Private)
	0.9 - 5.4 mg/kg/yr (Custom)	0.19 - 1.1 mg/kg/yr (Custom)	0.085 - 0.51 mg/kg/yr (Custom)
	(144.-864.)x10 ⁻⁵ mg/kg/day (Lifetime-Custom)	(30. - 176.) x 10 ⁻⁵ mg/kg/day (Lifetime-Custom)	(14. - 82.) x 10 ⁻⁵ mg/kg/day (Lifetime-Custom)

ALACHLOR (LASSO EC) EXPOSURE

<u>Operation</u>	<u>DERMAL EXPOSURE</u>		<u>RESPIRATORY EXPOSURE</u>
	<u>No Protection</u>	<u>With Protection (80%)</u>	<u>With Protection (100%)</u>
Aerial Tank-fill	1.8 mg/kg	0.37 mg/kg	0.024 mg/kg
	640. mg/hr	130. mg/hr	8.2 mg/hr
	9.1 mg/kg/hr	1.9 mg/kg/hr	0.12 mg/kg/hr
	0.0061 mg/kg/lb	0.00126 mg/kg/lb	0.00008 mg/kg/lb
	18.3 mg/kg/day	3.8 mg/kg/day	0.24 mg/kg/day
	92. - 183. mg/kg/yr	19. - 38. mg/kg/yr	1.2 - 2.4 mg/kg/yr
	0.14 - 0.29 mg/kg/day (Lifetime)	0.03 - 0.061 mg/kg/day (Lifetime)	0.0019 - 0.0038 mg/kg/day (Lifetime)
Aerial Pilot	0.082 mg/kg	0.019 mg/kg	0.0053 mg/kg
	8.2 mg/hr	1.9 mg/hr	0.53 mg/hr
	0.12 mg/kg/hr	0.027 mg/kg/hr	0.0076 mg/kg/hr
	0.00028 mg/kg/lb	0.000063 mg/kg/lb	0.000018 mg/kg/lb
	0.84 mg/kg/day	0.19 mg/kg/day	0.053 mg/kg/day
	4.2 - 8.4 mg/kg/yr	0.95 - 1.9 mg/kg/yr	0.26 - 0.53 mg/kg/yr
	0.0067 - 0.013 mg/kg/day (Lifetime)	0.0015 - 0.003 mg/kg/day (Lifetime)	0.00042 - 0.00085 mg/kg/day (Lifetime)

ALACHLOR (LASSO EC) EXPOSURE

<u>Operation</u>	<u>DERMAL EXPOSURE</u>		<u>RESPIRATORY EXPOSURE</u>
	<u>No Protection</u>	<u>With Protection (80%)</u>	<u>With Protection (100%)</u>
Aerial Flagman	2.5 mg/kg	0.56 mg/kg	0.22 mg/kg
	440. mg/hr	97. mg/hr	38 mg/hr
	6.3 mg/kg/hr	1.4 mg/kg/hr	0.55 mg/kg/hr
	0.0084 mg/kg/lb	0.00186 mg/kg/lb	0.00073 mg/kg/lb
	25.2 mg/kg/day	5.6 mg/kg/day	2.2 mg/kg/day
	126. - 252. mg/kg/yr	28. - 56. mg/kg/yr	11. - 22. mg/kg/yr
	0.2 - 0.4 mg/kg/day (Lifetime)	0.045 - 0.09 mg/kg/day (Lifetime)	0.018 - 0.035 mg/kg/day (Lifetime)

IV. CONCLUSIONS and RECOMMENDATION

With the recalculation of potential exposure to alachlor under various scenarios, EAB makes the following conclusions:

- o Respiratory exposure is insignificant when compared to dermal exposure.
- o In terms of mg/kg/lb, the following comparison can be made for potential dermal exposure, ground open tank fill and ground application:

Mixer/Loader + Applicator:


$$2 \times 10^{-4} \text{ mg/kg/lb} + 0.2 \times 10^{-4} \text{ mg/kg/lb} = 2.2 \times 10^{-4} \text{ mg/kg/lb (EPA-80\% Protection)}$$

$$3 \times 10^{-4} \text{ mg/kg/lb} + 0.6 \times 10^{-4} \text{ mg/kg/lb} = 3.6 \times 10^{-4} \text{ mg/kg/lb (Monsanto)}$$

$$4 \times 10^{-4} \text{ mg/kg/lb} + 1.2 \times 10^{-4} \text{ mg/kg/lb} = 5.2 \times 10^{-4} \text{ mg/kg/lb (Canada)}$$

- o It is believed that the exposure scenarios developed represent reasonable exposure ranges to a vast majority of alachlor users.
- o The exposure studies conducted by Monsanto were well planned and executed and continue to provide valuable data on exposure to applicators.

EAB defers to Toxicology Branch for the conclusion of the risk assessment. If further number generation or explanation is needed, please call me.



Richard V. Moraski, Ph.D.

Chemist

Environmental Chemistry Review Section No. 1

CALCULATIONS

Ground Application
Open Tank Fill

A. No Protection

Residue values average of two reps:

Hands (cotton gloves)	0.99 ug/cm ²	x	820 cm ²	=	811.8 ug
Forehead	0.02	x	650	=	13.0
Chest	3.2	x	3700	=	11840.
Back	0.04	x	3610	=	144.
Thigh	3.94	x	2250	=	8865.
Forearm (outer)	0.06	x	1210	=	72.6
Bicep (outer)	1.23	x	1320	=	1623.6
Ankle (outer)	0.33	x	2380	=	785.4

Total Hands + Body = 24155.4 ug
Total Body only = 23343.6 ug

Corrected for recovery:

Hands 811.8 ug/0.74 = 1097. ug
Body 23343.6 ug/0.89 = 26228.7 ug

Total Hands + Body = 27325.7 ug

Assume 70 kg body weight:

27325.7 ug/70 kg = 390 ug/kg = 0.39 mg/kg

Assume 0.24 hr (from study):

27325.7 ug/0.24 hr = 113857 ug/hr = 110 mg/hr

mg/kg/hr:

27325.7 ug/70 kg/0.24 hr = 1625 ug/kg/hr = 1.6 mg/kg/hr

mg/kg/lb:

assumptions: 0.24 hr/100 acre; 400 lb ai/100 acre; 4 lb ai/acre (from study)

1.6 mg/kg/hr x 0.24 hr/100 acre x 100 acre/400 lb = 96×10^{-5} mg/kg/lb

Daily exposure: mg/kg/day

assumptions: 0.24 hr/100 acre; 400 lb ai/100 acre; 100 acre/day

1.6 mg/kg/hr x 0.24 hr/day = 0.38 mg/kg/day

96×10^{-5} mg/kg/lb x 400 lb/day = 0.38 mg/kg/day

Annual exposure: mg/kg/yr:

-Private: assumptions: 1 - 6 day/yr; 100 - 600 acre/yr; 400 - 2400 lb/yr; 4 lb/acre

$$0.38 \text{ mg/kg/day} \times 1 \text{ day/yr} = 0.38 \text{ mg/kg/yr}$$

$$0.38 \text{ mg/kg/day} \times 6 \text{ day/yr} = 2.28 \text{ mg/kg/yr}$$

-Custom: assumptions: 5 - 30 day/yr; 500 - 3000 acre/yr; 2000 - 12000 lb/yr; 4 lb/acre

$$0.38 \text{ mg/kg/day} \times 5 \text{ day/yr} = 1.9 \text{ mg/kg/yr}$$

$$0.38 \text{ mg/kg/day} \times 30 \text{ day/yr} = 11.4 \text{ mg/kg/yr}$$

Lifetime daily exposure: mg/kg/day:

-Private: assumptions: 40 yr working lifetime; 70 yr lifespan [40/70x365 = 0.0016]

$$0.38 \text{ mg/kg/yr} \times \frac{40 \text{ yr}}{70 \text{ yr} \times 365 \text{ day/yr}} = 0.00061 \text{ mg/kg/day}$$

$$2.28 \text{ mg/kg/yr} \times 0.0016 = 0.0036 \text{ mg/kg/day}$$

-Custom: assumptions: 40 yr working lifetime; 70 yr lifespan

$$1.9 \text{ mg/kg/yr} \times 0.0016 = 0.003 \text{ mg/kg/day}$$

$$11.4 \text{ mg/kg/yr} \times 0.0016 = 0.018 \text{ mg/kg/day}$$

B. With Protection (80%)

Assume 80% reduction in exposure to body only. (20% of exposure with no protection.)

$$26228.7 \text{ ug} \times 20\% = 5245.7 \text{ ug}$$

$$\text{Total: Hand + Body} = 1097 \text{ ug} + 5245.7 \text{ ug} = 6342.7 \text{ ug}$$

$$\text{mg/kg: } 6342.7 \text{ ug}/70 \text{ kg} = 0.091 \text{ mg/kg}$$

$$\text{mg/hr: } 6342.7 \text{ ug}/0.24 \text{ hr} = 26 \text{ mg/hr}$$

$$\text{mg/kg/hr: } 0.091 \text{ mg/kg}/0.24 \text{ hr} = 0.38 \text{ mg/kg/hr}$$

$$\text{mg/kg/lb: } 0.38 \text{ mg/kg/hr} \times 0.24 \text{ hr}/100 \text{ acre} \times 100 \text{ acre}/400 \text{ lb} = 2 \times 10^{-4} \text{ mg/kg/lb}$$

$$\text{Daily exposure: mg/kg/day: } 0.38 \text{ mg/kg/hr} \times 0.24 \text{ hr/day} = 0.09 \text{ mg/kg/day}$$

Annual exposure: mg/kg/yr:

Private

$$\begin{aligned} 0.09 \text{ mg/kg/day} &\times 1 \text{ day/yr} = 0.09 \text{ mg/kg/yr} \\ 0.09 \text{ mg/kg/day} &\times 6 \text{ day/yr} = 0.54 \text{ mg/kg/yr} \end{aligned}$$

Custom

$$\begin{aligned} 0.09 \text{ mg/kg/day} &\times 5 \text{ day/yr} = 0.45 \text{ mg/kg/yr} \\ 0.09 \text{ mg/kg/day} &\times 30 \text{ day/yr} = 2.7 \text{ mg/kg/yr} \end{aligned}$$

Lifetime daily exposure: mg/kg/day

Private

$$\begin{aligned} 0.09 \text{ mg/kg/yr} &\times 0.0016 = 0.00014 \text{ mg/kg/day} \\ 0.54 \text{ mg/kg/yr} &\times 0.0016 = 0.00086 \text{ mg/kg/day} \end{aligned}$$

Custom

$$\begin{aligned} 0.45 \text{ mg/kg/yr} &\times 0.0016 = 0.00072 \text{ mg/kg/day} \\ 2.7 \text{ mg/kg/yr} &\times 0.0016 = 0.0043 \text{ mg/kg/day} \end{aligned}$$

C. With Protection (100%)

Assume 100% protection to covered body parts. (Hand exposure determined by cotton gloves.) The following body areas are the only exposed parts to alachlor: face, front of neck ("V" of chest), back of neck and hands as determined by residues in cotton gloves.

Residue values average of two reps:

Hands (cotton gloves)	0.99 ug/cm ²	x	820 cm ²	=	812. ug
Chest	3.2	x	150	=	480.
Back	0.04	x	110	=	4.4
Forehead	0.02	x	650	=	13.

$$\begin{aligned} \text{Total Hands + Body} &= 1309.4 \text{ ug} \\ \text{Total Body only} &= 497. \text{ ug} \end{aligned}$$

Corrected for recovery:

$$\begin{aligned} \text{Hands: } 812 \text{ ug}/0.74 &= 1097.3 \text{ ug} \\ \text{Body: } 497.4 \text{ ug}/0.89 &= 558.8 \text{ ug} \end{aligned}$$

$$\text{Total Hands + Body} = 1656.2 \text{ ug}$$

$$\text{mg/kg: } 1656.2 \text{ ug}/70 \text{ kg} = 0.023 \text{ mg/kg}$$

$$\text{mg/hr: } 1656.2 \text{ ug}/0.24 \text{ hr} = 6.9 \text{ mg/hr}$$

$$\text{mg/kg/hr: } 1656.2 \text{ ug}/70 \text{ kg}/0.24 \text{ hr} = 0.099 \text{ mg/kg/hr}$$

$$\text{mg/kg/lb: } 0.099 \text{ mg/kg/hr} \times 0.24 \text{ hr}/100 \text{ acre} \times 100 \text{ acre}/400 \text{ lb} = 0.000059 \text{ mg/kg/lb}$$

Daily exposure: mg/kg/day:

$$0.099 \text{ mg/kg/hr} \times 0.24 \text{ hr/day} = 0.024 \text{ mg/kg/day}$$

mg/kg/yr:

Private

$$0.024 \text{ mg/kg/day} \times 1 \text{ day/yr} = 0.024 \text{ mg/kg/yr}$$

$$0.024 \text{ mg/kg/day} \times 6 \text{ day/yr} = 0.14 \text{ mg/kg/yr}$$

Custom

$$0.024 \text{ mg/kg/day} \times 5 \text{ day/yr} = 0.12 \text{ mg/kg/yr}$$

$$0.024 \text{ mg/kg/day} \times 30 \text{ day/yr} = 0.72 \text{ mg/kg/yr}$$

Lifetime daily exposure: mg/kg/day

Private

$$0.024 \text{ mg/kg/yr} \times 0.0016 = 0.000038 \text{ mg/kg/day}$$

$$0.14 \text{ mg/kg/yr} \times 0.0016 = 0.00022 \text{ mg/kg/day}$$

Custom

$$0.12 \text{ mg/kg/yr} \times 0.0016 = 0.00019 \text{ mg/kg/day}$$

$$0.72 \text{ mg/kg/yr} \times 0.0016 = 0.0012 \text{ mg/kg/day}$$

D. Respiratory Exposure

Respiratory exposure was abstracted from the current submission without change and multiplied by appropriate factors to obtain specified units. The highest residue value was chosen to determine the exposure (12.7 ug/m^3).

$$\text{mg/hr: } 12.7 \text{ ug/m}^3 \times 1.5 \text{ m}^3/\text{hr} = 19 \text{ ug/hr}$$

$$\text{mg/kg/hr: } 19 \text{ ug/hr}/70 \text{ kg} = 0.27 \text{ ug/kg/hr}$$

Daily exposure: mg/kg/day:

$$12.7 \text{ ug/m}^3 \times 1.5 \text{ m}^3/\text{hr} \times 0.24 \text{ hr/day} = 4.6 \text{ ug/day}/70 \text{ kg} = 0.065 \text{ ug/kg/day}$$

$$\text{mg/kg/lb: } 0.0000004 \text{ mg/kg/lb (from current submission)}$$

Ground Application
Applicator

A. No Protection

Residue values average of two reps:

Hands (cotton gloves)	0.27 ug/cm ²	x	820 cm ²	=	221.4 ug
Forehead	0.011	x	650	=	7.15
Chest	0.146	x	3700	=	540.2
Back	0.007	x	3610	=	25.3
Thigh	0.128	x	2250	=	288.0
Forearm (outer)	0.11	x	1210	=	13.3
Bicep (outer)	0.32	x	1320	=	42.2
Ankle (outer)	0.027	x	2380	=	64.3

Total Hands + Body	1201.85 ug
Total Body only	980.45 ug

Corrected for recovery:

Hands	221.4 ug/0.74	=	299.2 ug
Body	980.45 ug/0.89	=	1101.6 ug

Total Hands + Body 1330.8 ug

Assume 70 kg body weight:

1330.8 ug/70 kg = 19.0 ug/kg

Assume 1.1 hr (from study):

1330.8 ug/1.1 hr = 1209.8 ug/hr

mg/kg/hr:

1330.8 ug/70 kg/1.1 hr = 17.2 ug/kg/hr

mg/kg/lb:

assumptions: 1.1 hr/100 acre; 400 lb ai/100 acre; 4 lb ai/acre (from study)

0.017 mg/kg/hr x 1.1 hr/100 acre x 100 acre/400 lb = 0.046 ug/kg/lb

Daily exposure: mg/kg/day:

assumptions: 1.1 hr/100 acre; 400 lb ai/100 acre; 100 acre/day

0.017 mg/kg/hr x 1.1 hr/day = 0.018 mg/kg/day

Annual exposure: mg/kg/yr:

-Private: assumptions: 1 - 6 day/yr; 100 - 600 acre/yr; 400 - 2400 lb/yr; 4 lb/acre

$$0.018 \text{ mg/kg/day} \times 1 \text{ day/yr} = 0.018 \text{ mg/kg/yr}$$

$$0.018 \text{ mg/kg/day} \times 6 \text{ day/yr} = 0.11 \text{ mg/kg/yr}$$

-Custom: assumptions: 5 - 30 day/yr; 500 - 3000 acre/yr; 2000 - 12000 lb/yr; 4lb/acre

$$0.018 \text{ mg/kg/day} \times 5 \text{ day/yr} = 0.09 \text{ mg/kg/yr}$$

$$0.018 \text{ mg/kg/day} \times 30 \text{ day/yr} = 0.54 \text{ mg/kg/yr}$$

Lifetime daily exposure: mg/kg/day

-Private: assumptions: 40 yr working lifetime; 70 yr lifespan [40/70x365 = 0.0016]

$$0.018 \text{ mg/kg/yr} \times \frac{40 \text{ yr}}{70 \text{ yr} \times 365 \text{ day/yr}} = 2.9 \times 10^{-5} \text{ mg/kg/day}$$

$$0.11 \text{ mg/kg/yr} \times 0.0016 = 18 \times 10^{-5} \text{ mg/kg/day}$$

-Custom: assumptions: 40 yr working lifetime; 70 yr lifespan

$$0.09 \text{ mg/kg/yr} \times 0.0016 = 14 \times 10^{-5} \text{ mg/kg/day}$$

$$0.54 \text{ mg/kg/yr} \times 0.0016 = 86 \times 10^{-5} \text{ mg/kg/day}$$

B. With Protection (80%)

Assume 80% reduction in exposure to body only. (20% of exposure with no protection.)

$$1101.6 \text{ ug} \times 20\% = 220.3 \text{ ug}$$

$$\text{Total Hands} + \text{Body} = 299.2 \text{ ug} + 220.3 \text{ ug} = 519.5 \text{ ug}$$

$$\text{mg/kg: } 519.5 \text{ ug}/70 \text{ kg} = 7.4 \text{ ug/kg}$$

$$\text{mg/hr: } 519.5 \text{ ug}/1.1 \text{ hr} = 472.3 \text{ ug/hr}$$

$$\text{mg/kg/hr: } 519.5 \text{ ug}/70 \text{ kg}/1.1 \text{ hr} = 6.7 \text{ ug/kg/hr}$$

$$\text{mg/kg/lb: } 0.0067 \text{ mg/kg/hr} \times 1.1 \text{ hr}/100 \text{ acre} \times 100 \text{ acre}/400 \text{ lb} = 0.000018 \text{ mg/kg/lb}$$

$$\text{Daily exposure: mg/kg/day: } 6.7 \text{ ug/kg/hr} \times 1.1 \text{ hr/day} = 0.0073 \text{ mg/kg/day}$$

Annual exposure: mg/kg/yr

Private

$$\begin{aligned} 0.0073 \text{ mg/kg/day} \times 1 \text{ day/yr} &= 0.0073 \text{ mg/kg/yr} \\ 0.0073 \text{ mg/kg/day} \times 6 \text{ day/yr} &= 0.044 \text{ mg/kg/yr} \end{aligned}$$

Custom

$$\begin{aligned} 0.0073 \text{ mg/kg/day} \times 5 \text{ day/yr} &= 0.036 \text{ mg/kg/yr} \\ 0.0073 \text{ mg/kg/day} \times 30 \text{ day/yr} &= 0.22 \text{ mg/kg/yr} \end{aligned}$$

Lifetime daily exposure: mg/kg/day

Private

$$\begin{aligned} 0.0073 \text{ mg/kg/yr} \times 0.0016 &= 1.2 \times 10^{-5} \text{ mg/kg/day} \\ 0.044 \text{ mg/kg/yr} \times 0.0016 &= 7. \times 10^{-5} \text{ mg/kg/day} \end{aligned}$$

Custom

$$\begin{aligned} 0.036 \text{ mg/kg/yr} \times 0.0016 &= 5.8 \times 10^{-5} \text{ mg/kg/day} \\ 0.22 \text{ mg/kg/yr} \times 0.0016 &= 35. \times 10^{-5} \text{ mg/kg/day} \end{aligned}$$

C. With Protection (100%)

Assume 100% protection to covered body parts. (Hand exposure determined by cotton gloves.) The following body areas are the only exposure parts to alachlor: face, front of neck ("V" of chest), back of neck, and hands as determined by residues in cotton gloves.

Residue values average of two reps:

Hands (cotton gloves)	0.27 ug/cm ²	x	820 cm ²	=	221.4 ug
Chest	0.146	x	150	=	21.9
Back	0.007	x	110	=	0.77
Forehead	0.011	x	650	=	7.15

Total Hands + Body	251.22 ug
Total Body only	29.82 ug

Corrected for recovery:

$$\begin{aligned} \text{Hands} \quad 221.4 \text{ ug}/0.74 &= 299.2 \text{ ug} \\ \text{Body} \quad 29.82 \text{ ug}/0.89 &= 33.5 \text{ ug} \end{aligned}$$

Total Hands + Body 332.7 ug

$$\text{mg/kg: } 332.7 \text{ ug}/70 \text{ kg} = 4.75 \text{ ug/kg}$$

$$\text{mg/hr: } 332.7 \text{ ug}/1.1 \text{ hr} = 302.4 \text{ ug/hr}$$

$$\text{mg/kg/hr: } 332.7 \text{ ug}/70 \text{ kg}/1.1 \text{ hr} = 4.3 \text{ ug/kg/hr}$$

$$\text{mg/kg/lb: } 4.3 \text{ ug/kg/hr} \times 1.1 \text{ hr/100 acre} \times 100 \text{ acre/400 lb} = 0.012 \text{ ug/kg/lb}$$

$$\text{Daily exposure: mg/kg/day: } 4.3 \text{ ug/kg/day} \times 1.1 \text{ hr/day} = 0.0047 \text{ mg/kg/day}$$

Annual exposure: mg/kg/yr:

Private

$$0.0047 \text{ mg/kg/yr} \times 1 \text{ day/yr} = 0.0047 \text{ mg/kg/yr}$$

$$0.0047 \text{ mg/kg/yr} \times 6 \text{ day/yr} = 0.028 \text{ mg/kg/yr}$$

Custom

$$0.0047 \text{ mg/kg/yr} \times 5 \text{ day/yr} = 0.024 \text{ mg/kg/yr}$$

$$0.0047 \text{ mg/kg/yr} \times 30 \text{ day/yr} = 0.14 \text{ mg/kg/yr}$$

Lifetime daily exposure: mg/kg/day:

Private

$$0.0047 \text{ mg/kg/yr} \times 0.0016 = 0.75 \times 10^{-5} \text{ mg/kg/day}$$

$$0.028 \text{ mg/kg/yr} \times 0.0016 = 4.5 \times 10^{-5} \text{ mg/kg/day}$$

Custom

$$0.024 \text{ mg/kg/yr} \times 0.0016 = 3.8 \times 10^{-5} \text{ mg/kg/day}$$

$$0.14 \text{ mg/kg/yr} \times 0.0016 = 22. \times 10^{-5} \text{ mg/kg/day}$$

D. Respiratory Exposure

Respiratory exposure was abstracted from the current submission without change and multiplied by appropriate factors to obtain specified units. The highest residue value was chosen to determine exposure (1.45 ug/m^3).

$$\text{mg/hr: } 1.45 \text{ ug/m}^3 \times 1.5 \text{ m}^3/\text{hr} = 2.2 \text{ ug/hr}$$

$$\text{mg/kg/hr: } 2.2 \text{ ug/hr/70 kg} = 0.031 \text{ ug/kg/hr}$$

Daily exposure: mg/kg/day:

$$1.45 \text{ ug/m}^3 \times 1.5 \text{ m}^3/\text{hr} \times 1.1 \text{ hr/day} = 2.4 \text{ ug/day}$$

$$2.4 \text{ ug/day/70 kg} = 0.034 \text{ ug/kg/day}$$

$$\text{mg/kg/lb: } 0.0000004 \text{ mg/kg/lb (from current submission)}$$

Canadian Calculation

Open Tank Fill: 28.3 ug/lb/70 kg = 0.0004 mg/kg/lb

Application: 8.78 ug/lb/70 kg = 0.00012 mg/kg/lb

Other Scenarios

Similar calculations were performed for the remaining exposure scenarios (aerial, probe transfer). Original study provides information concerning hours applied, lb/acre used, etc.

TABLES

Abstracted from original study
and current submission and from
Canadian response.

Tables from original study.

EPA Accession No.: 070591

Table VII
AERIAL APPLICATOR EXPOSURE
RECOVERIES OF ALACHLOR
FROM
FIELD FORTIFIED PUF PLUGS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.62	--
check	0.24	--
200.0 ug	180.04	90
200.0 ug	198.01	99
		95 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table VIII

LASSO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

FIELD FORTIFIED PUF PLUGS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.78	--
check	0.63	--
check	0.06	--
check	0.70	--
5.0 ug	5.73	98
5.0 ug	5.64	97
10.0 ug	10.87	101
10.0 ug	10.65	98
15.0 ug	16.23	103
15.0 ug	15.77	100
20.0 ug	22.67	102
20.0 ug	20.70	99
check	63.63	--
check	29.19	--
check	47.56	--
check	54.64	--
25.0 ug	64.47	61
25.0 ug	63.29	76
50.0 ug	88.48	78
50.0 ug	65.45	72
100.0 ug	123.26	71
100.0 ug	116.92	68
200.0 ug	230.54	91
200.0 ug	214.88	83
		85 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table IX
LIASSO II GRANULES
 GROUND APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 FIELD FORTIFIED PTF PLUGS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	1.32	--
check	1.45	--
check	1.57	--
check	1.57	--
10.0 ug	12.32	107
10.0 ug	11.63	100
20.0 ug	22.06	102
20.0 ug	21.94	102
50.0 ug	55.63	108
50.0 ug	53.10	103
200.0 ug	185.66	93
200.0 ug	192.24	95
		101 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OF
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table X
AERIAL APPLICATOR EXPOSURE
RECOVERIES OF ALAMINOR
FROM
LAB FORTIFIED PUF PLUGS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.58	--
check	0.22	--
10.0 ug	9.06	91
10.0 ug	8.74	87
20.0 ug	17.77	89
20.0 ug	17.51	88
50.0 ug	44.04	88
50.0 ug	44.20	88
200.0 ug	175.32	88
200.0 ug	176.86	88
		88 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table XI

20.

LASSO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

LAB FORTIFIED PTF PLUGS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	7.25	--
check	7.17	--
5.0 ug	5.27	171
5.0 ug	5.43	174
10.0 ug	10.47	104
10.0 ug	10.23	100
check	7.23	--
check	6.20	--
15.0 ug	16.45	108
15.0 ug	15.73	105
20.0 ug	21.82	108
20.0 ug	21.80	108
check	6.21	--
check	6.20	--
25.0 ug	28.65	114
25.0 ug	29.15	116
50.0 ug	54.82	109
50.0 ug	56.83	113
check	7.26	--
check	6.26	--
100.0 ug	87.37	87
100.0 ug	89.75	89
200.0 ug	195.72	98
200.0 ug	192.69	96
check	7.64	--
check	6.55	--
500.0 ug	498.96	100
500.0 ug	511.53	102
1000.0 ug	721.70	72
1000.0 ug	782.67	98
		103 % average

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF AGRICULTURE
C-100-1012

Recoveries are corrected for their respective backgrounds.

21

Table XII
 AERIAL APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 FIELD FORTIFIED SILICA GEL TUBES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.01	--
check	0.01	--
0.1 ug	0.12	111
0.1 ug	0.11	100
check	0.01	--
check	0.02	--
2.5 ug	2.56	94
2.5 ug	2.63	105
check	0.02	--
check	0.02	--
20.0 ug	19.06	95
20.0 ug	18.95	95
		100 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XIII

LA550 EC
GROUND APPLICATOR EXPOSURE
RECOVERIES OF ALACHLOR
FROM
FIELD FORTIFIED SILICA GEL

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.03	--
check	0.04	--
check	0.03	--
check	0.03	--
0.1 ug	0.12	93
0.1 ug	0.12	93
0.5 ug	0.55	104
0.5 ug	0.56	106
1.0 ug	1.09	106
1.0 ug	1.04	101
2.5 ug	2.72	108
2.5 ug	2.43	96
check	0.02	--
check	0.02	--
check	0.14	--
check	0.22	--
5.0 ug	4.74	93
5.0 ug	4.64	--
7.5 ug	6.62	87
7.5 ug	6.35	83
10.0 ug	9.47	94
10.0 ug	10.06	100
20.0 ug	16.39	81
20.0 ug	16.07	80
		94 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY.

Table XIV
LAISO - GRANULES
 GROUND APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 FIELD FORTIFIED SILICA GEL

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.05	--
check	0.05	--
check	0.03	--
check	0.02	--
0.5 ug	0.41	74
0.5 ug	0.41	74
2.5 ug	1.76	69
2.5 ug	1.72	67
7.5 ug	6.79	90
7.5 ug	5.39	71
20.0 ug	15.92	79
20.0 ug	13.65	69
		<u>74 % average</u>

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XV
AERIAL APPLICATOR EXPOSURE
RECOVERIES OF ALACHLOR
FROM
LAB FORTIFIED SILICA GEL TUBES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.03	--
check	0.04	--
0.1 ug	0.13	92
0.1 ug	0.15	114
1.0 ug	0.96	92
1.0 ug	0.99	95
check	0.12	--
check	0.17	--
5.0 ug	5.69	111
5.0 ug	5.16	100
10.0 ug	10.17	100
10.0 ug	9.34	92
		100 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table XVI

ASSO 22

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

LAB FORTIFIED SILICA GEL

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.01	--
check	0.01	--
0.1 ug	0.11	100
0.1 ug	0.10	90
0.5 ug	0.61	120
0.5 ug	0.57	112
check	0.02	--
check	0.01	--
1.0 ug	0.94	92
1.0 ug	1.19	117
2.5 ug	2.29	91
2.5 ug	2.50	99
check	0.01	--
check	0.01	--
5.0 ug	4.55	91
5.0 ug	4.19	84
7.5 ug	6.50	87
7.5 ug	6.43	85
check	0.01	--
check	0.01	--
10.0 ug	9.27	83
10.0 ug	8.46	85
20.0 ug	17.42	87
20.0 ug	16.34	82
check	0.00	--
check	0.01	--
25.0 ug	23.87	95
25.0 ug	23.30	93
50.0 ug	45.58	91
50.0 ug	45.46	91

CONTAINS TRADE SECRET
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

(continued)

Table XVI (continued)

LISSO EC

LAB FORTIFIED SILICA GEL

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.01	--
check	0.00	--
100.0 ug	74.67	75
100.0 ug	63.24	63
250.0 ug	217.07	87
250.0 ug	224.79	90
check	0.01	--
check	0.01	--
500.0 ug	403.06	81
500.0 ug	407.86	82
		91 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table XXI
 AERIAL APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 FIELD FORTIFIED GAUZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.01	--
check	----	--
check	0.01	--
check	0.01	--
0.5 ug	0.49	96
0.5 ug	0.43	85
0.5 ug	0.42	82
0.5 ug	0.48	94
check	----	--
check	----	--
check	----	--
check	----	--
10.0 ug	8.15	82
10.0 ug	7.60	76
10.0 ug	8.25	83
10.0 ug	7.95	80
check	----	--
check	----	--
check	----	--
check	----	--
200.0 ug	144.31	72
200.0 ug	141.67	71
200.0 ug	153.68	77
200.0 ug	166.66	83
		82 % average

σ_r = 7.20

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XXII

LASSO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

FIELD FORTIFIED GAUZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.06	--
check	0.05	--
check	0.04	--
check	0.03	--
0.5 ug	0.35	60
0.5 ug	0.34	58
1.0 ug	0.77	72
1.0 ug	0.72	67
5.0 ug	3.96	78
5.0 ug	4.33	86
10.0 ug	8.91	89
10.0 ug	8.97	89
check	0.34	--
check	0.24	--
check	0.30	--
check	0.29	--
25.0 ug	23.53	93
25.0 ug	23.37	92
50.0 ug	44.20	88
50.0 ug	44.80	89
100.0 ug	82.80	83
100.0 ug	84.03	84
200.0 ug	180.03	90
200.0 ug	168.63	84
		81 % average

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF INDUSTRY
GOVERNMENT

82-10.86

Recoveries are corrected for their respective backgrounds.

Table XXIII
ASSO II GRATULES
 GROUND APPLICATOR EXPOSURE
 RECOVERIES OF ALATHLOR
 FROM
 FIELD FORTIFIED GAUZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	17.51	--
check	0.05	--
check	0.05	--
check	0.05	--
1.0 ug	0.99	94
1.0 ug	0.89	84
10.0 ug	9.59	95
10.0 ug	9.66	96
50.0 ug	46.87	94
50.0 ug	47.66	95
200.0 ug	199.07	100
200.0 ug	195.76	98
		95 % average

Recoveries are corrected for their respective backgrounds.

¹Not used for background correction.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XXIV
AERIAL APPLICATOR EXPOSURE
RECOVERIES OF ALACHLOR
FROM
LAS FORTIFIED GAUZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.08	--
check	0.05	--
0.5 ug	0.58	102
0.5 ug	0.74	134
check	0.07	--
check	0.12	--
1.0 ug	0.90	81
1.0 ug	1.08	99
check	0.07	--
check	0.05	--
5.0 ug	4.75	94
5.0 ug	4.65	92
check	0.10	--
check	0.06	--
10.0 ug	8.94	89
10.0 ug	8.89	88
check	0.13	--
check	0.04	--
25.0 ug	20.96	84
25.0 ug	21.85	87
check	0.04	--
check	0.03	--
50.0 ug	37.46	75
50.0 ug	39.95	80
check	0.04	--
check	0.04	--
100.0 ug	84.73	85
100.0 ug	95.91	96
200.0 ug	229.43	109
200.0 ug	201.83	101
		51% average

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MUNSJIO
COMB-314

$\sigma = 13.4$

93%

Recoveries are corrected for their respective backgrounds.

Table XXV

LASSO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

LAB FORTIFIED GAUZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.05	--
check	0.04	--
0.5 ug	0.35	60
0.5 ug	0.35	50
check	0.07	--
check	0.08	--
1.0 ug	0.79	71
1.0 ug	0.91	73
check	0.07	--
check	0.07	--
5.0 ug	4.32	85
5.0 ug	4.06	80
check	0.07	--
check	0.06	--
10.0 ug	9.19	91
10.0 ug	9.05	90
check	0.06	--
check	0.06	--
25.0 ug	19.92	79
25.0 ug	19.10	76
check	0.08	--
check	0.10	--
50.0 ug	41.25	82
50.0 ug	43.55	87
check	0.06	--
check	0.03	--
100.0 ug	81.50	81
100.0 ug	84.14	84

CONTAINS 17
OTHERWISE
INFORMATION
COMPANY

DE 51

DE 51

LASSO EC

37.

LAB FORTIFIED LAUZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.02	--
check	0.02	--
3000.0 ug	2670.32	89
3000.0 ug	2725.64	91
check	0.15	--
check	0.11	--
4000.0 ug	4122.03	103
4000.0 ug	3933.79	96
check	0.25	--
check	0.16	--
5000.0 ug	4938.33	100
5000.0 ug	4999.73	100
check	0.30	--
check	0.18	--
0.5 ug	0.60	72
0.5 ug	0.58	68
check	0.15	--
check	0.13	--
1.0 ug	0.98	84
1.0 ug	1.01	87
check	0.02	--
check	0.02	--
5.0 ug	3.97	79
5.0 ug	3.76	75
check	0.04	--
check	0.18	--
10.0 ug	10.60	105
10.0 ug	10.53	104
check	0.11	--
check	0.09	--
25.0 ug	24.48	98
25.0 ug	25.52	102
check	0.04	--
check	0.05	--
50.0 ug	47.75	95
50.0 ug	48.53	97

CONTAINS TRADE SECRET &
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

44
(continued)

(continued)

LASSO EC

LAE FORTIFIED CANZE PADS

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.25	--
check	0.17	--
100.0 ug	106.54	106
100.0 ug	116.52	116
200.0 ug	122.09	121
200.0 ug	257.64	134
		91 % average

Recoveries are corrected for their respective background.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table XXVI
 AERIAL APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 FIELD FORTIFIED COTTON GLOVES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.87	--
check	1.14	--
10.0 ug	9.04	81
10.0 ug	8.89	79
check	1.03	--
check	1.14	--
100.0 ug	68.50	67
100.0 ug	76.02	75
check	0.93	--
check	0.65	--
1000.0 ug	954.91	95
1000.0 ug	997.62	100
		<u>83 % average</u>

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XXVII

LASSO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

FIELD FORTIFIED COTTON GLOVES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	1.23	--
check	0.80	--
10.0 ug	7.01	69
10.0 ug	6.72	57
25.0 ug	17.15	65
25.0 ug	14.26	53
50.0 ug	39.91	78
50.0 ug	32.37	63
100.0 ug	65.16	64
100.0 ug	62.08	61
check	3.43	--
check	2.07	--
250.0 ug	191.21	75
250.0 ug	198.64	78
500.0 ug	356.53	71
500.0 ug	373.95	74
750.0 ug	510.83	69
750.0 ug	555.93	74
1000.0 ug	764.39	76
1000.0 ug	783.42	78
		<u>68</u> % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

13

Table XXVIII
LAISO II GRANULES
 GROUND APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 FIELD FORTIFIED COTTON GLOVES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.49	--
check	0.46	--
25.0 ug	19.07	74
25.0 ug	20.44	80
100.0 ug	64.50	64
100.0 ug	61.12	61
500.0 ug	356.12	71
500.0 ug	376.77	75
1000.0 ug	664.36	66
1000.0 ug	752.33	75
		<u>71 % average</u>

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XXIX
 AERIAL APPLICATOR EXPOSURE
 RECOVERIES OF ALACHLOR
 FROM
 LAS FORTIFIED COTTON GLOVES

<u>fort level</u>	<u>ug found</u>	<u>% rec.</u>
check	1.53	--
check	1.27	--
10.0 ug	9.13	77
10.0 ug	9.05	76
25.0 ug	22.22	83
25.0 ug	22.87	86
check	0.74	--
check	0.65	--
50.0 ug	39.34	77
50.0 ug	40.84	80
100.0 ug	82.64	82
100.0 ug	82.52	82
500.0 ug	509.98	102
500.0 ug	466.60	93
		84 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table AAX

43.

LAISO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

LAB FORTIFIED COTTON GLOVES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	2.78	--
check	1.38	--
10.0 ug	6.87	48
10.0 ug	7.34	50
check	0.93	--
check	0.35	--
10.0 ug	8.36	77
10.0 ug	6.43	58
check	0.56	--
check	0.00	--
25.0 ug	14.56	57
25.0 ug	17.57	69
check	0.37	--
check	0.21	--
50.0 ug	33.70	67
50.0 ug	34.50	68
check	0.48	--
check	0.33	--
100.0 ug	63.93	64
100.0 ug	56.60	56
500.0 ug	356.18	71
500.0 ug	378.15	76
1000.0 ug	682.16	68
1000.0 ug	651.75	66
2500.0 ug	1706.0	68
2500.0 ug	1614.97	65
5000.0 ug	2965.78	59
5000.0 ug	2572.82	57
10000.0 ug	6260.37	63
10000.0 ug	6312.35	63
		64 % average

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSIEUR
COMPTON

Recoveries are corrected for their respective backgrounds.

Table XXXI

LASSO EC

GROUND APPLICATOR EXPOSURE

RECOVERIES OF ALACHLOR

FROM

LAB FORTIFIED SOIL

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.40	--
check	0.66	--
5.0 ug	5.18	93
5.0 ug	5.03	90
check	3.28	--
50.0 ug	47.56	89
50.0 ug	45.32	84
check	1.77	--
check	0.86	--
100.0 ug	79.53	78
100.0 ug	97.15	96
check	11.22	--
check	4.27	--
200.0 ug	158.13	75
200.0 ug	159.22	76
check	3.08	--
check	3.22	--
300.0 ug	330.55	109
300.0 ug	352.29	116
400.0 ug	326.13	81
400.0 ug	329.03	81
500.0 ug	452.72	80
500.0 ug	541.90	108
600.0 ug	623.42	103
600.0 ug	620.66	101
		92 % average

Recoveries are corrected for their respective backgrounds.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

45.

Table XXXII
GROUND APPLICATOR EXPOSURE
RECOVERIES OF ALACHLOR
FROM
LAS FORTIFIED RUBBER GLOVES

<u>fort. level</u>	<u>ug found</u>	<u>% rec.</u>
check	0.13	--
check	0.07	--
10,000.0 ug	8,775.54	88
10,000.0 ug	8,606.37	86
		<u>87 % average</u>

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

5.1

Table XXXIII

LASSO EC AERIAL APPLICATOR EXPOSURE

TANK FILLSAMPLING CONDITIONS

Wind (mph-direction):
Humidity:
Air Temperature (°C):
Sampling Time (min):

REP NO. 1REP NO. 2

Measured Concentration of
Alachlor in Air (ug/m³)

(polyurethane foam)

6.04
7.0310.52
4.69

(silica gel)

14.11
14.7239.08
35.69

Measured Concentration of
[redacted] (ug/m³)

(charcoal)

9,154.5
12,424.28,033.3
9,151.5

Dermal Deposition of
Alachlor (ug/cm²)

Hands:

0.051

1.409

Head:

0.036

0.065

Forehead:

0.026

0.777

Shoulder:

0.040

0.075

Chest:

7.040

2.515

Back:

0.031

0.070

Thigh:

16.050

50.676

Forearm (under):

0.075

0.181

Forearm (outer):

0.423

0.286

Bicep (under):

0.007

0.051

Bicep (outer):

0.151

0.234

Ankle (under):

0.075

270.90

Ankle (outer):

15.955

0.166

37%

76%

outlier

78%

average
reduction
normal clothes

49

Table XX.:V

LASSO EC AERIAL APPLICATOR EXPOSURE

APPLICATION (PILOT)

		REP NO. 1	REP NO. 2
<u>SAMPLING CONDITIONS</u>	Wind (mph-direction):	7 SW	3 NE
	Humidity:	62%	62%
	Air Temperature (°C):	15	10
	Sampling Time (min):	35	47
Measured Concentration of Alachlor in Air ($\mu\text{g}/\text{m}^3$)	(polyurethane foam)		
	(silica gel)	15.20 17.79	7.53 12.72
	(charcoal)	<2,272.1 ¹ <2,272.7 ¹	<1,694.9 ¹ <1,694.9 ¹
Measured Concentration of [REDACTED] ($\mu\text{g}/\text{m}^3$)			
Dermal Deposition of Alachlor ($\mu\text{g}/\text{cm}^2$)	Hands:	0.177	0.270
	Head:	0.057	0.037
	Forehead:	0.045	0.045
	Shoulder:	0.060	0.049
	Chest:	0.521	0.034
	Back:	0.640	0.206
	Thigh:	0.155	0.105
	Forearm (under):	0.038	0.052
	Forearm (outer):	1.844	0.036
	Bicep (under):	0.065	0.032
	Bicep (outer):	0.043	0.029
	Ankle (under):	0.067	0.079
	Ankle (outer):	0.645	0.061

¹ Based on 200 μg sensitivity and volume of air sampled.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

47% protection
factor
↑
mixed average of 1
and 2
for under + out

50.

Table XXXV
LASSO EC AERIAL APPLICATOR EXPOSURE

<u>APPLICATION (FLAG-MAN)</u>		REP NO. 1	REP NO. 2
<u>SAMPLING CONDITIONS</u>	Wind (mph-direction):	7 SW	3 NE
	Humidity:	62%	62%
	Air Temperature (°C):	15	10
	Sampling Time (min):	25	24
Measured Concentration of Alachlor in Air (ug/m ³)	(polyurethane foam)		
	(silica gel)	168.88 96.48	152.75 103.67
Measured Concentration of [REDACTED] (ug/m ³)	(charcoal)	<3,174.6 ¹ <3,174.6 ¹	<3,333.3 ¹ <3,333.3 ¹
Dermal Deposition of Alachlor (ug/cm ²)	Hands:	6.603	2.273
	Head:	9.904	7.232
	Forehead:	18.637	2.228
	Shoulder:	4.266	12.948
	Chest:	9.039	2.351
	Back:	3.110	29.561
	Thigh:	13.258	2.462
	Forearm (under):	9970 0.045	<1.006 73%
	Forearm (outer):	21.377	3.723
	Bicep (under):	99 0.038	<0.048 96%
	Bicep (outer):	12.650	3.185
	Ankle (under):	99 0.134	<0.133 92%
	Ankle (outer):	16.789	1.623

¹ Based on 200 ug sensitivity and volume of air sampled.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table XXXVI
LASEO EC GROUND APPLICATOR EXPOSURE
OPEN-POUR TANK FILL

		REP NO. 1	REP NO. 2	
<u>SAMPLING CONDITIONS</u>	Wind (mph-direction):	9 S	7 NE	
	Humidity:	64%	62%	
	Air Temperature (°C):	20	15	
	Sampling Time (min):	12	17	
Measured Concentration of Alachlor in Air (ug/m ³)	(polyurethane foam)	1.89	2.33	
		20.39	2.57	
	(silica gel)	6.67	2.94	
		7.50	10.12	
Measured Concentration of [REDACTED] (ug/m ³)	(charcoal)	9,177.0	<4,651.0 ¹	
		7,153.0	<4,651.0 ¹	
Dermal Deposition of Alachlor (ug/cm ²)	Hands:	1.545	0.427	
	Head:	0.012	0.011	
	Forehead:	0.026	0.006	
	Shoulder:	0.037	0.030	
	Chest:	4.958	1.415	
	Back:	0.004	0.079	
	Thigh:	7.794	0.089	
	Forearm (under):	0.007	0.014	52%
	Forearm (outer):	0.003	0.023	
	Bicep (under):	0.002	0.008	95%
	Bicep (outer):	2.267	0.187	
	Ankle (under):	0.008	0.009	98%
	Ankle (outer):	0.111	0.540	
		90% <		
		99% <		
		92% <		

¹ Based on 200 ug sensitivity and volume of air sampled.

Ave 88%

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

53.

Table XA .1
LASSO EC GROUND APPLICATOR EXPOSURE
APPLICATION/INCORPORATION (INSIDE)

<u>SAMPLING CONDITIONS</u>		
	<u>REP NO. 1</u>	<u>REP NO. 2</u>
Wind (spt-direction):	9 S	7 NE
Humidity:	64%	62%
Air Temperature (°C):	20	15
Sampling Time (min):	69	63
Measured Concentration of Alachlor in Air (ug/m ³)		
(polyurethane foam)		
(silica gel)	1.19 1.15	1.33 1.40
Measured Concentration of [REDACTED] (ug/m ³)		
(charcoal)	1,255.0 1,259.0	1,525.0 1,479.0
Dermal Deposition of Alachlor (ug/cm ²)		
Hands:	0.497	0.041
Head:	0.008	0.007
Forehead:	0.011	0.011
Shoulder:	0.058	0.019
Chest:	0.273	0.019
Back:	0.005	0.010
Thigh:	0.258	0.018
Forearm (under):	0.069	0.007
Forearm (outer):	0.015	0.008
Bicep (under):	0.018	0.010
Bicep (outer):	0.050	0.013
Ankle (under):	0.115	0.065
Ankle (outer):	0.017	0.036

64% 12% 23%

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table A.IV
 LASSO EC GROUND APPLICATOR EXPOSURE
PROBE TRANSFER TANK FILL (5 GALLON CAN)

<u>SAMPLING CONDITIONS</u>		<u>REP NO. 1</u>	<u>REP NO. 2</u>
	Wind (mph-direction):	12 NE	10 NE
	Humidity:	50%	47%
	Air Temperature (°C):	17	18
	Sampling Time (min):	15	18
Measured Concentration of Alachlor in Air (ug/m ³)	(polyurethane foam)	3.86 2.46	1.23 3.54
	(silica gel)	9.08 5.69	2.67 2.11
Measured Concentration of [REDACTED] (ug/m ³)	(charcoal)	<6,061.0 ¹ <6,061.0 ¹	<4,444.0 ¹ <4,444.0 ¹
Dermal Deposition of Alachlor (ug/cm ²)	Hands:	0.010	0.012
	Head:	0.156	0.005
	Forehead:	0.992	0.034
	Shoulder:	0.014	0.005
	Chest:	0.014	0.005
	Back:	0.025	0.007
	Thigh:	0.031	0.055
	Forearm (under):	0.010	0.036
	Forearm (outer):	0.015	0.042
	Bicep (under):	0.005	0.005
	Bicep (outer):	0.016	0.005
	Ankle (under):	0.011	0.019
	Ankle (outer):	0.028	0.532

¹ Based on 200 ug sensitivity and volume of air sampled.

CONTAINS TRADE SECRET OR
 OTHERWISE CONFIDENTIAL
 INFORMATION OF MONSANTO
 COMPANY

Table XLVII

LASTO EC GROUND APPLICATION EXPOSURE

PROBE TRANSFER TANK FILL (55 GALLON DRUM)

<u>SAMPLING CONDITIONS</u>		<u>REP NO. 1</u>	<u>REP NO. 2</u>
Wind (mph-direction):		12 SSE	13 SSE
Humidity:		77%	74%
Air Temperature (°C):		21	21
Sampling Time (min):		11	13
Measured Concentration of Alachlor in Air (ug/m ³)	(polyurethane foam)	5.99	7.64
		3.01	5.66
	(silica gel)	9.82	8.31
		5.82	28.46
Measured Concentration of [REDACTED] (ug/m ³)	(charcoal)	(7,142.0) ¹	13,474.0
		(7,142.0) ¹	16,650.0
Dermal Deposition of Alachlor (ug/cm ²)	Hands:	0.717	0.452
	Head:	0.304	0.023
	Forehead:	0.069	0.016
	Shoulder:	0.073	0.024
	Chest:	1.462	0.111
	Back:	1.575	0.015
	Thigh:	4.504	0.222
	Forearm (under):	0.005	0.034
	Forearm (outer):	0.610	0.233
	Bicep (under):	0.046	0.022
	Bicep (outer):	0.366	0.045
	Ankle (under):	0.043	0.013
	Ankle (outer):	0.013	0.012

¹ Based on 200 ug sensitivity and volume of air sampled.CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table XLVIII

Total Body Dose Estimates

Combined Inhalation & Dermal Exposure

	1	2	3	4	5	6	7	8
Exposure	Unit Inhalation Exposure ^{1,2}	Total Inhalation Exposure ^{1,2}	Total Daily Inhalation Exposure ^{1,3} 100% Absorption Assumed	Unit Dermal Exposure ^{1,4}	Total Dermal Exposure ^{1,5}	Total Daily Dermal Exposure ^{1,6} 10% Absorption Assumed	Total Body Dose Dermal & Inhalation (ug/kg bw/100 acres) ⁷	Total Body Dose Dermal & Inhalation (ug/kg bw/pound applied)
Fill (one)	0.08			20.71				
cation (1)	0.23	0.31	0.31	5.38	26.09	2.61	2.92	0.010
cation (men)	1.31	1.31	1.31	211.47	211.47	21.15	22.46	0.075
Fill (tor)	0.19			111.61				
ication/ine. (de cab)	0.19	0.36	0.36	21.00	132.61	13.26	13.64	0.034
er Fill (ter)	2.11			21.41				
ication (de cab)	0.20	2.31	2.31	N.B.	21.41	2.14	4.45	0.011

-11-

3/4 lb/acre treated (refer to footnotes in tables 49 and 50 for explanation of assumptions used).

Unit Inhalation values are from table 49, column 5.

Unitive total of tankfill values and application values.

Column 2 multiplied by absorption rate.

Unit dermal exposure values are from table 50, column 5.

Column 5 multiplied by absorption rate.

Unitive total of columns 3 and 6.

erial application calculated at 970 lbs/100 acres; ground application calculated at 400 lbs/100 acres.

Table XLIX

Applicator Exposure to Alachlor Through Inhalation

1 Operation	2 Average Exposure Level ($\mu\text{g}/\text{M}^3$)	3 Range ($\mu\text{g}/\text{M}^3$)	4 Number of Samples	5 Time (in hours) for 100 acre application	6 Unit Inhalation Exposure ($\mu\text{g}/\text{kg bw}/100$ acres treated)
Tank Fill (airplane)	16.53	4.7-39.1	8	0.2	0.08
Application (pilot)	13.14	7.5-17.1	4	0.7	0.23
Application (flag-man)	130.50	96.5-160.9	4	0.4	1.31
Tank Fill (tractor)	6.00	1.9-20.4	8	1.1	0.19
Application/Ine. (inside cab)	1.30	1.1-1.40	4	5.7	0.19
Hopper Fill (planter)	64.85	1.4-193.0	8	1.3	2.11
Application (inside cab)	0.00	0.6-1.0	4	10.0	0.20

1 Inside tractor cab air concentration values were used for application/incorporation operations.

2 A 60 kg female body weight is assumed; a $1.5 \text{ M}^3/\text{hr}$ breathing rate is assumed.

3 Multiply the value in column 1 by column 4, then multiply by $1.5 \text{ M}^3/\text{hr}$, and then divide by 60 kg.

Table 1.

Applicator Exposure via Dermal Deposition

Operation	1 Average Exposure Level	2 Range	3 Number of Samples	4 Time (in hours) for 100 acre application	5 Unit Dermal Exposure (ug/kg bw/100 acres treated)
Tank Fill (airplane)	6,212	1,889-10,535	2	0.2	20.71 $\frac{60}{70}$
Application (pilot)	461	325-591	2	0.7	5.38 $\frac{60}{70}$
Application (flag-man)	31,721	9,985-53,552	2	0.4	211.47 $\frac{60}{70}$
Tank Fill (tractor)	6,088	2,004-10,173	2	1.1	111.61 $\frac{60}{70}$
Application/Inc. (inside cab)	221	44-398	2	5.7	21.00
Hopper Fill (planter)	988	264-1,711	2	1.3	21.41
Application (inside cab)	N.D.	N.D.	2	10.0	N.D.

1. ug/exposed area.
2. The pads located on the forehead and back of neck, and the gloves were considered representative of the exposed skin on the face, back of neck, "v" of the chest and hands.
3. It is assumed that a long-sleeved shirt is worn. front of neck and "v" of chest.
4. Exposed areas: face = 650 cm²; back of neck = 110 cm²; front of neck and "v" of chest = 150 cm²; hands = 820 cm². Based on W. J. Hayse, Toxicology of Pesticides and Durham and Wolfe, Bull. Who 26, 75-91 (1962).
5. Range: lowest and highest values from forehead, back, chest pads and hand gloves, divided by exposure time (minutes), multiplied by 60 min/hr, multiplied by the appropriate surface areas listed in footnote 4.
6. A 60 kg body weight is assumed.
7. Multiply column 1 by column 4, then divide by 60 kg body weight.

Table LI

Applicator Exposure Through Inhalation

1 Operation	2 Average Exposure Level ($\mu\text{g}/\text{M}^3$) ¹	3 Range ($\mu\text{g}/\text{M}^3$)	4 Number of Samples	5 Time (in hours) for 100 acre application	Unit Inhalation Exposure ($\mu\text{g}/\text{kg}$ bw/100 acres treated) ^{2,3}
Tank Fill (airplane)	9,691	0,033-12,424	4	0.2	40.46
Application (pilot)	N.D.	N.D.	4	0.7	N.D.
Application (flag-man)	N.D.	N.D.	4	0.4	N.D.
Tank Fill (open pour)	6,408	4,651-9,177	4	1.1	117.48
Tank Fill (5 gal. probe)	N.D.	N.D.	4	1.1	N.D.
Tank Fill (55 gal. probe)	11,102	7,142-16,650	4	0.9	166.53
Application/Inc. (inside cab)	1,375	1,239-1,525	4	5.7	195.87

¹ Inside tractor cab air concentration values were used for application/incorporation operations.

² A 60 kg female body weight is assumed; a 1.5 M³/hr breathing rate is assumed.

³ Multiply the value in column 1 by column 4, then multiply by 1.5 M³/hr, and then divide by 60 kg.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL

Table LII
Total Body Dose Estimates
Combined Inhalation & Dermal Exposure

Operation	1 Unit Inhalation Exposure ^{1,2}	2 Total Inhalation Exposure ^{1,3}	3 Total Daily Inhalation Exposure ^{1,4} 10% Absorption Assumed	4 Unit Dermal Exposure ^{1,5}	5 Total Dermal Exposure ^{1,5}	6 Total Daily Dermal Exposure ^{1,6} 10% Absorption Assumed	7 Total Body Dose Dermal & Inhalation (ug/kg bw/100 acres)	8 Total Body Dose Dermal & Inhalation (ug/kg bw/pound applied)
Tank Fill (5 gal. probe)	0.11			26.93				
Application/Inc. (inside cab)	0.19	0.30	0.30	21.00	49.93	4.99	5.29	0.013
Tank Fill (55 gal. probe)	0.21			55.83				
Application/Inc. (inside cab)	0.19	0.40	0.40	21.00	76.83	7.68	8.08	0.020

1. ug/kg bw/area treated (refer to footnotes in tables S1 and S4 for explanation of assumptions used).
2. Unit inhalation values are from table S1, column 5.
3. Additive total of tankfill values and application values.
4. Column 2 multiplied by absorption rate.
5. Unit dermal exposure values are from table S4, column 5.
6. Column 5 multiplied by absorption rate.
7. Additive total of columns 3 and 6.
8. Application calculated at 470 lbs/100 acres.

CONTAINS TRADE SECRET OF
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table LIII

Applicator Exposure to Alachlor Through Inhalation

Operation	1 Average Exposure Level ($\mu\text{g}/\text{M}^3$)	2 Range ($\mu\text{g}/\text{M}^3$)	3 Number of Samples	4 Time (in hours) for 100 acre application	5 Unit Inhalation Exposure ($\mu\text{g}/\text{kg bw}/100$ acres treated) ^{1,2}
Tank Fill (5 gal. probe)	3.83	1.2-9.1	8	1.1	0.11
Tank Fill (55 gal. probe)	9.34	3.0-28.5	8	0.9	0.21

¹ A 60 kg female body weight is assumed; a 1.5 M³/hr breathing rate is assumed.
² Multiply the value in column 1 by column 4, then multiply by 1.5 M³/hr, and then divide by 60 kg.

CONTAINS TRADE SECRET
 OTHERWISE CONFIDENTIAL
 INFORMATION OF Monsanto
 COMPANY

Table LIV
Applicator Exposure to Alachlor Via Dermal Deposition

Operation	Average ¹ Exposure Level ^{1,2,3,4}	Range ² 1,5	Number of Samples ³	Time (in hours) for 100 acre application ⁴	Unit Dermal Exposure ⁵ (ug/kg bw/100 acres treated) ^{6,7}
Tank Fill (5 gal. probe)	1,578	114-3,042	2	1.1	28.93
Tank Fill (55 gal. probe)	3,722	1,843-5,595	2	0.9	55.83

1. ug/exposed area.
2. The pads located on the forehead and back of neck and the gloves were considered representative of the exposed skin on the face, back of neck and "V" of the chest and hands.
3. It is assumed that a long-sleeved shirt is worn.
4. Exposed areas: face = 650 cm²; back of neck = 110 cm²; front of neck and "V" of chest = 150 cm²; hands = 820 cm². Based on W. J. Hayse, Toxicology of Pesticides and Durham and Wolfe, Bull. WHO 26, 75-91 (1962).
5. Range: lowest and highest values from forehead, back, chest pads and hand gloves, divided by exposure time (minutes), multiplied by 60 min/hr, multiplied by the appropriate surface areas listed in footnote 4.
6. A 60 kg body weight is assumed.
7. Multiply column 1 by column 4, then divide by 60 kg body weight.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSIEUR
COMPANY

Tables from current submission.

EPA Accession No.: 252498

Table V

LASSO EC - GROUND - OPEN POUR

Reps are calculated separately then are averaged,
Lbs alachlor: Rep 1 = 80, Rep 2 = 108

I. DERMAL

I. DERMAL											
Operation	Body Site	Deposition		Ala- chlor lbs ³	Deposition		Site Area cm ²	Deposition Total µg/lb AL	Body (70 kg) "Exposure" µg/kg bw/lb AL	Body Load 10% Absor µg/kg bw/lb	
		Rec %	µg/cm ²		µg/cm ² /lb AL	Avg					
			Raw ²		Corr						
Mix/Load	Hands	68	1.545 0.427	2.27 0.628	80 108	0.0284 0.0058	0.0171	820	14.0	0.2000	0.0200
	Forehead	81	0.026 0.006	0.032 0.007	80 108	0.0004 0.0001	0.0003	650	0.19	0.0030	0.0003
	Back (Neck)	81	0.008 0.079	0.010 0.098	80 108	0.0001 0.0009	0.0005	110	0.06	0.0008	0.0001
	Chest (Vee)	81	4.958 1.415	6.127 1.747	80 108	0.0765 0.0162	0.0463	150	6.95 21.2	0.0993 0.3031	0.0099 0.0303
						DERMAL TOTALS					
	Hands	68	0.499 0.041	0.734 0.060	80 108	0.0092 0.0006	0.0049	820	4.02	0.0574	0.0057
	Forehead	81	0.011 0.011	0.014 0.014	80 108	0.0002 0.0001	0.0002	650	0.13	0.0014 0.0002	0.0002 0.0000
	Back (Neck)	81	<0.005 0.010	0.006 0.012	80 108	0.0000 0.0001	0.0001	110	0.013	0.0002	0.0000
	Chest (Vee)	81	0.273 0.019	0.337 0.023	80 108	0.0042 0.0002	0.0022	150	0.33 4.49	0.0047 0.0025	0.0005 0.0062
						DERMAL TOTALS					

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

03/20/84
2-20-84

Table VI

LASSO EC - GROUND - OPEN POUR

Reps are calculated separately then averaged,
Lbs alachlor Rep 1 = 80; Rep 2 = 108

II. INHALATION		2-28-84													
Operation	Type Sample	Rep	③ Raw Data		Rec'd %	Conc'n µg/m³	Corr µg/m³	Avg µg/m³	µg/m³ Avg/Rep	Exposure µg/m³		Exposure, Total @ 100%			
			per Rep	per Rep						x Hr	x 1.5 m³/Hr	Per 80 lb Ala	Per 108 lb Ala BW (70)		
Mix/Load	PUF	1	1.89	20.39	88	2.15	23.2	12.7	10.1	0.20	3.03	0.038	0.0005		
	SiO2		6.67	7.50	94	7.09	7.98	7.53							
	PUF	2	2.33	2.57	88	2.65	2.92	2.79	4.87	0.28	2.05	0.019	0.0003		
	SiO2		2.94	10.12	94	3.13	10.8	6.95							
Applic/Incor	SiO2	1	1.19	1.13	94	1.27		1.24	1.24	1.15	2.14	0.027	0.0004		
					1.20										
	SiO2	2	1.33	1.40	94	1.41		1.45	1.45	1.05	2.28	0.021	0.0003		
					1.49										
													Mix/Load Avg	0.0004	
													Appl Avg	0.0004	

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

µg/kg bw/lb
alachlor

-16-
Table VI Footnotes: References: R.D. No. 396, Special Report MSL-1983, Vol. 7, EPA Accession No. 70591.

¹ Raw data, for Mix/Load: Polyurethane Foam, PUF, Table XXXVI, p. 52

Silica Gel, SiO₂, Table XXXVI, p. 52

For Applic/Incorp, Silica Gel, SiO₂, Table XXXVII, p. 53 (PUF not used).

² Recovery %: PUF, Table VIII, p. 17; 88%

SiO₂, Table XIII, p. 22; 94%

³ Replicate descriptions, Rep. 1, p. 79-80 (20 gals Lasso EC, 80 lbs)

Rep 2, p. 80 (27 gals Lasso EC, 108 lbs).

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Table VII

LASSO EC - GROUND APPLICATION - COMBINED OPERATIONS

<u>Operation</u>	<u>Body Load (70 kg)</u>		
	<u>µg/kg bw/lb Alachlor</u>		
	<u>Dermal¹</u>	<u>Inhal.²</u>	<u>Total</u>
Mix/Load	0.0303	0.0004	0.0307
	0.0062		0.0369
Applic/Incorp	<u>0.0062</u>	<u>0.0004</u>	<u>0.0066</u>
	0.0367		0.0375
Combined Totals	<u>0.0365</u>	<u>0.0008</u>	<u>0.0373</u>

¹ Dermal absorption at 10%

² Lung absorption at 100%

From Table VII, the total body load exposure through the mix/load and application/incorporation operations is 0.0373⁵ µg/kg bw/lb alachlor applied. It is to be noted that of this total body load exposure only 2% or 0.0008 µg/kg bw/lb alachlor is by inhalation, attesting to the very low volatility of alachlor.

Using this body load exposure value a typical grower-applicator lifetime exposure can be calculated. The parameters for the calculation for this typical farmer using Lasso are listed below. It is to be noted that a farmer uses Lasso, a pre-emergent herbicide, only once each year. From Monsanto marketing information for 1981, the average or typical grower-applicator grows 130 acres of corn and 120 acres of soybeans, both treated with Lasso at an average rate of about 2 lbs/acre.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Lasso usage, number of years	40 years
Lifetime	70 years
Area treated per year	250 acres
Average application rate	2.0 lbs alachlor/acre
Total body load dose per pound alachlor applied	0.037 µg/kg bw/lb alachlor

$$\frac{0.037 \text{ µg/kg bw/lb} \times 2.0 \text{ lb/acre} \times 250 \text{ acres} \times 40 \text{ years}}{70 \text{ years} \times 365 \text{ days/year}}$$

$$0.029 \text{ µg/kg bw/day} =$$

$$2.9 \times 10^{-5} \text{ mg/kg bw/day}$$

The result at 2.9×10^{-5} mg/kg bw/day is incrementally higher than the 2.0×10^{-5} mg/kg bw/day value for a grower-applicator reported in 1982 (R.D. No. 396, Special Report MSL-1983, Vol. 1 (of 9), Section I, Part 3, pp. 15-16). Factors contributing to the difference in value are (1) the aforementioned correction for recoveries from field-spiked sampling devices; (2) the shift to a 70 kg body weight from a 60 kg body weight; and (3) an increase of years for using alachlor to 40 from 30. As mentioned, the last two factors represent values used by the Agency in recent risk assessment notices for several other pesticides. The 2.9×10^{-5} mg/kg bw/day value is used for exposure risk assessment calculations.

CONTAINS TRADE SECRET OR
OTHERWISE CONFIDENTIAL
INFORMATION OF MONSANTO
COMPANY

Canadian Calculation.

CANADIAN CALCULATION

TABLE 3 Total Dermal Exposure Field Application (EC)

patch represent- ing area	body area (cm ²)	corrected (1) patch values (ug/cm ² /lb handled) (2)	regional exposure (3) (ug/body area/lb handled) (2 applicators mixing spraying 1 trial)
cotton gloves rubber gloves	820	1.7×10^{-2} 4.64×10^{-1}	13.9 (380.6)*
forehead	650	2.33×10^{-4}	0.15
back (over)	110	5.2×10^{-4}	0.057
chest (over)	150	6.7×10^{-2}	7.05
lower arm (over)	1286	8.6×10^{-4}	1.1
upper arm (under)	5240	6.68×10^{-5}	0.42
upper leg (under)	6028	1.11×10^{-4}	0.67
			23.3
			8.78

Exposed Area

hands

face

back of neck

of chest

lower arms

Covered Area

chest, back
upper arms

upper and
lower leg

The total body exposure for each pound of
pesticide mixed and sprayed is 32.1 ug

32.1 ug/lb used $\times 60$ lb/day used

= 214 ug/kg b.w./day for applicator
wearing rubber gloves during mixing

or

(32.1 ug/lb used $\times 310.6$ ug/lb used $\times 400$ lb)

= 2751.33 ug/kg b.w./day for an applicator
not wearing rubber gloves during mixing

- 1) Original analytical results, corrected for field recovery
- 2) Patch values for each applicator normalized with respect to the number of pounds sprayed, then results of
of two trials averaged.
- 3) Corrected patch value $\times$ body area = regional contact exposure
- 4) Total dermal contact exposure during mixing and spraying
- 5) residue on rubber gloves representing "unprotected hands"