# 

### EEE BRANCH REVIEW

DATE:	IN	OUT 6/3 /25	INOUT ENVIRONMENTAL CHEMISTRY			IN_	OUT		<del></del>
							EFFICACY		
			•	-					
		•							
ETTE OF	PEC NO.			• • • •			J	1 14. 14.	
					EC1624	<del></del>	; <u>, , , , , , , , , , , , , , , , , , ,</u>	***************************************	
PETITIO	ON OR EXP.	PERMIT NO	239-EXP	).•	561624			<u>,</u>	•
DATE D	V. RECEIVE	D April 28, 1	975					•	<del></del>
DATE OF	SUBMISSIO	N_April 10,	1975			·,		<del>lysians and the state</del>	<del></del>
DATE SU	JEMISSION A	ACCEPTED			•				
TYPE P	RODUCT (S):	(I, D, H, F,	N, R, S	<u>Insecti</u>	cide		· · · · · · · · · · · · · · · · · · ·		
PRODUC	r MGR. NO.		<del>, ,</del>						
PRODUC	r name (s)_	Monitor 4 Sp	pary				•		<del></del>
COMPAN	y Name_	Chevron Cher	nical Com	npany	·	,			
SUBMIS	SION PURPO	SE Obtain	Exp. Perr	nit for us	e of Moni	tor in	almo	nds	
CHEMIC	AL & FORMU	LATION 0.S -	Dimethy'	l phosphor	amidothio	ate -	40%		

#### 100.0 Pesticidal Use

#### 100.1..2

(a) READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Keep Pesticide in original container. Do not put concentrate or dilute into food or drink containers.

Not for use or storage in or around the home.

FOR EXPERIMENTAL USE ONLY

#### Directions

ALMONDS: Navel Orangeworm - Apply 2 to 8 qts. (2 to 8 lbs. active) per acre as a single application 2 weeks before to 30 weeks after hull split, or 2 to 4 qts. (2 to 4 lbs. active) per acre as a 2 application program with the first application within 2 weeks before hull split and the second application within 3 weeks after hull split. Apply 400 to 500 gals. spray mix per acre using conventional ground equipment. Do not apply within 8 weeks of harvest. Do not feed hulls to live-stock.

## (b) Fish & Wildlife Cautions on Label:

This product is toxic to birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

# (c) Purpose of Program

To develop more information on effective rates, application timing, yields and quality, and residue data under large-scale field trials.

# (d) Area of Use

California only.

# (e) Amount to be Shipped

88 gals. of Monitor 4 spray.

# (f) Acreage

24 - 86.

### 101.0 Chemical and Physical Properties

101.1 O.S - Dimethyl phosphoramidothioate

101.2 Monitor (Ortho 9006)

$$\begin{array}{c} \text{CH}_{3^0} > \stackrel{0}{\underset{P}{\mid}} - \text{NH}_2 \\ \text{CH}_{3^S} \end{array}$$

- 102.0 <u>Behavior in the Environment</u>
  (Previously submitted)
- 103.0 <u>Toxicological Properties</u> (Previously submitted)

#### 104.0 Hazard Assessment

#### 104.1 Discussion

- 104.1.1 The toxicological studies submitted so far appear adequate.
- 104.1.2 An expansion of MONITOR's use to any new crops with large acreages (other than almonds) will necessitate dietary LC<sub>50</sub> studies with mallard duck and avian reproduction studies.
- The impact of MONITOR on non-target organisms should be minimal. MONITOR is not highly toxic to fish and degrades rapidly in soil (1/2-life = 2-6 days). It is noted that MONITOR is acutely toxic to mammals and birds and that chronic studies show adverse effects on mammals at low levels (3 ppm). However, the short 1/2-life of MONITOR lessens concerns about chronic toxicity. The label precautions and limitations concerning wildlife are presently adequate. Further, one application of a maximum of 8# a.i./A (or two applications of 4# + 4# a.i./A) by ground equipment in an area not considered to be prime wildlife habitat greatly reduces hazard to non-target organisms.

#### 105.0 Conclusions

105.1 The recommended rates of application are confusing. The label recommends 2#-8# a.i./A (single application) or 2#-4# a.i./A (two applications) in 400-500 gals. spray mix/A. The proposed experimental program, however, recommends 0.5-20# a.i./100 gals. of spray mix in 400-500 gpa. This implies a maximum application of 10# a.i./A. Which do you intend? Please clarify or amend where appropriate.

105.2 An expansion of MONITOR"s use to any new crops with large acreages (other than almonds) will necessitate dietary LC  $_{\rm 50}$  studies with mallard duck and avian reproduction studies.

Norman J. Cook Fish & Wildlife Section

Efficacy & Ecological Effects Branch

6/11/75