

METHAMIDOPHOS

Task 3: Environmental Fate Profile

Contract No. 68-01-5830

Final Report

December 9, 1981

SUBMITTED TO:

Environmental Protection Agency
Arlington, Virginia 22202

SUBMITTED BY:

 Enviro Control, Inc.
The Dynamac Building
11140 Rockville Pike
Rockville, MD 20852


A Subsidiary of the Dynamac Corporation

This document contains commercial and/or financial information of Enviro Control, Inc. that is privileged or confidential and is therefore subject to the restrictions on disclosure outside the agency to which such document has been submitted, as provided in the Freedom of Information Act (PL 89-554, September 6, 1966, as amended).

METHAMIDOPHOS

Task 3

METHAMIDOPHOS, MONITOR, TAMARON



O,S-Dimethyl phosphoramidothioate

Environmental Fate Profile

Available data are insufficient to fully assess the environmental fate of methamidophos.

The following products were identified in alkaline and acid hydrolysates (10% NaOH and 10% HCl) of technical methamidophos (~70% purity) at 80 C: O-methyl phosphoric acid, S-methyl phosphorothioate, methanol, methyl mercaptan, and ammonia (Magee, 00014039). The mechanisms by which these compounds originated cannot be determined due to deficiencies in the protocols of the study.

Methamidophos at 1 ppm had half-lives of 2-6 days in silt, loam, and sandy soils at 21 C (Leary and Tutass, 00014076). Methamidophos at 20 ppm had a half-life of 10-12 days in a sandy loam soil at 24 C (Tucker, 00014991). In another study, methamidophos levels declined ~50% (from 1.3 ppm) within 9 days after the last of three applications to a sandy loam soil over a 33-day period at room temperature (Tucker, 00014497). Lubkowitz (05017379) identified the following products in three types of soils incubated with methamidophos at 22 C: O,S-dimethyl phosphorothioate, O-methyl phosphoric acid, S-methyl phosphoroamidothioate, O-methyl phosphoroamidate, and phosphate ion.

More than 90% of the ^{14}C had dissipated from a silt soil 1 week after application of [S-methyl- ^{14}C]methamidophos at 0.2 ppm and incubation at

37 C (Leary and Tutass, 00014076). ^{14}C was detected in the form of amino acids and carbohydrates, indicating that microorganisms metabolized methamidophos. Microbial metabolism was also demonstrated by the fact that only ~10% of the applied ^{14}C dissipated after 1 week in sterile soil at 37 C. Only 8% of the applied ^{14}C dissipated in a volatile form after 3 days in anaerobic soil, versus 70% in aerobic soil.

Studies by Focht and Joseph (00015233 and 05017226), Ramadan and Zidan (05019841 and 05019842), and Zidan and Ramadan (05017741) indicate that methamidophos at actual use levels in soil will not have a long-term adverse effect on populations of bacteria (including nitrogen fixers) and fungi, or on microbial soil respiration, cellulose decomposition, ammonification, sulfur oxidation, and nitrification. Observed inhibitory effects were not severe, and generally lasted less than 1 month.

In soil thin-layer chromatography studies (Thornton et al., 00029887; Tucker, 00014992), [^{14}C]methamidophos was moderately mobile to very mobile in nine soil types (sand, loamy sand, sandy loam, sandy clay loam, silt loam, silty clay, silty clay loam, loam, and clay).

Methamidophos exhibited a bioaccumulation factor of <2 on the 7th day of exposure of the marine diatom Cylindrotheca fusiformis at 1-10 ppm (Tucker, 00014496), on the 8th day of exposure of bass at 0.01 ppm (Stanley, 00014014 and 00014018; Chemagro Corporation, 00014017 and 00014019), and over a 28-day period of exposure of bass at 0.8-1.5 ppm (Baychem Corporation, 00014015 and 00014016). In the bass, depuration to nonquantifiable levels (<0.014 ppm) occurred on the 1st day of depuration. ^{14}C residues had a bioaccumulation factor of ~2 in Daphnia magna on the 3rd day of exposure to [^{14}C]methamidophos at 0.1 ppm (Tucker, 00015242). Methamidophos is very soluble in water (Farm Chemicals Handbook, 1981, Meister Publishing Co., Willoughby, OH) and therefore should have a low octanol/water partition coefficient.

In summary, methamidophos had a half-life of <2 weeks in soil under aerobic conditions. The S-methyl group is metabolized and incorporated into amino

acids and carbohydrates. Products formed in soil and water are O,S-dimethyl phosphorothioate, O-methyl phosphoric acid, S-methyl phosphorothioate, S-methyl phosphoroamidothioate, O-methyl phosphoroamidate, methanol, methyl mercaptan, ammonia, and phosphate ion. Some temporary inhibitory effects on soil microbes and their functions may occur in some soils, but long-term adverse effects are not expected as a result of methamidophos use. Methamidophos will be moderately mobile to very mobile in most types of soils, but degradate mobility is unknown. If methamidophos enters an aquatic system, it will be transformed and residues are not expected to accumulate in aquatic non-target organisms.

Summary of Major Data Gaps

The major data gaps for this chemical are: hydrolysis studies; photodegradation studies in water, on soil, and in air; aerobic and anaerobic soil metabolism studies; a degradate mobility study; laboratory and field volatility studies; and terrestrial field dissipation studies.

Label Restrictions

At present, there are no label restrictions regarding the environmental chemistry of methamidophos.

References

Baychem Corporation. 1972. Chemagro, Division of Baychem Corporation. Residue experiment: Report No. 31933. (Unpublished study prepared by Baychem Corp.) (00014015)

Baychem Corporation. 1972. Chemagro, Division of Baychem Corporation. Residue experiment: Report No. 31938. (Unpublished study received on unknown data under OF0956, submitted by Chevron Chemical Co., Richmond, CA; CDL:093266-H) (00014016)

Chemagro Corporation. 1971. Recovery of Monitor from bass and rainbow trout: Report No. 30,796. Dated Sept. 30, 1971. (Unpublished study prepared by Baychem Corp.) (00014017)

Chemagro Corporation. 1971. Recovery of Monitor from bass and rainbow trout: Report No. 30,977. Dated Sept. 30, 1971. (Unpublished study prepared by Baychem Corp.) (00014019)

Focht, D.D., and H.A. Joseph. 1969? Microbial activity in soils treated with acephate and its major degradation product. (Unpublished study received Mar. 27, 1973 under 239-EX-60; prepared by Univ. of California at Riverside, Dept. of Soil Science and Agricultural Engineering, submitted by Chevron Chemical Co., Richmond, CA; CDL:223489-Q) (00015233)

Focht, D.D., and H. Joseph. 1974. Microbial activity in soils treated with acephate and Monitor. J. Environ. Qual. 3(4):327-328. (05017226)

Leary, J.B., and H.O. Tutass. 1968. Degradation of Monitor insecticide in soil (Unpublished study received Mar. 5, 1970 under OF0956; submitted by Chevron Chemical Co., Richmond, CA; CDL:093264-AN) (00014076)

Lubkowitz, J.A. 1975. Uptake and degradation of methamidophos by tomato plants and soils. in Origin and fate of chemical residues in food, agriculture and fisheries. p. 157-163. Proceedings and report of two research coordination meetings; 1973, 1974, Vienna, Austria. (05017379)

Magee, P.S. 1966. Hydrolysis of Monitor insecticide. (Unpublished study received Mar. 5, 1970 under OF0956; submitted by Chevron Chemical Co., Richmond, CA; CDL:093263-C) (00014039)

Ramadan, E.M., and Z.H. Zidan. 1977. Influence of certain organophosphorus insecticides on soil microflora. 1. Total microbial flora, actinomycetes, fungi, yeasts, and cellulose decomposers. Ann. Agric. Sci. 20(2):57-63. (05019842)

Ramadan, E.M., and Z.H. Zidan. 1977. Influence of certain organophosphorus insecticides on soil microflora. 2. Nonsymbiotic nitrogen fixers and nitrifying bacteria. Ann. Agric. Sci. 20(2):65-71. (05019841)

Stanley, C.W. 1971. A gas chromatographic method for the determination of Monitor in fish and water: Report No. 30975. Dated Sept. 30, 1971. (Unpublished study received on unknown date under OF0956; prepared by Baychem Corp., submitted by Chevron Chemical Co., Richmond, CA; CDL: 093266-J) (00014018)

Stanley, C.W. 1971. Analysis of bass and water for Monitor: Report No. 30979. (Unpublished study received on unknown date under OF0956; prepared by Baychem Corp.; submitted by Chevron Chemical Co., Richmond, CA; CDL: 093266-F) (00014014)

Thornton, J.S., J.B. Hurley, and J.J. Obrist. 1976. Soil thin-layer mobility of twenty-four pesticide chemicals: Report No. 51016. (Unpublished study received Jan. 28, 1980 under 5F1547; submitted by Mobay Chemical Corp., Pittsburgh, PA; CDL:099216-I) (00029887)

Tucker, B.V. 1973. Orthene and Ortho 9006 in Daphnia magna living in treated water. (Unpublished study prepared by Chevron Chemical Co., Richmond, CA) (00015242)

Tucker, B.V. 1972. Orthene leaching in soil (Unpublished study including supplementary report, received Feb. 23, 1972 under 2G1248; submitted by Chevron Chemical Co., Richmond, CA; CDL:091774-AA) (00014992)

Tucker, B.V. 1972. Orthene soil metabolism-laboratory studies. (Unpublished study including supplement, received Feb. 23, 1972 under 2G1248; submitted by Chevron Chemical Co., Richmond, CA; CDL:091774-Z) (00014991)

Tucker, B.V. 1972. Residues in earthworms in Orthene and Ortho 9006 treated soil. (Unpublished study prepared by Chevron Chemical Co., Richmond, CA) (00014497)

Tucker, B.V. 1972. Residues of Orthene and Ortho 9006 in a marine diatom growing in treated water. (Unpublished study received Aug. 7, 1972 under 239-2406; submitted by Chevron Chemical Co., Richmond, CA; CDL:001571-U) (00014496)

Zidan, Z.H., and E.M. Ramadan. 1976. Degradation of some organophosphorus insecticides by fungi. *Egyp. J. Microbiol.* 11(1/2):93-98. (05017741)