

13000  
5  
OPP OFFICIAL RECORD  
HEALTH EFFECTS DIVISION  
SCIENTIFIC DATA REVIEWS  
EPA SERIES 361

EPA Reviewer: William B. Greear, M.P.H. *William B. Greear, 5/19/95*  
Review Section IV, Toxicology Branch I (7509C)  
EPA Secondary: Marion P. Copley, D.V.M.  
Review Section, Toxicology Branch I (7509C) *Marion Copley 5/19/95*

## DATA EVALUATION REPORT

Study Type: 14-Day Intrathyroidal Metabolism Study  
Guideline Series 85-1

EPA ID No.: MRID No.: 431350-03  
Pesticide Chemical Code: 108501  
Tox Chemical No.: 454BB  
DP Barcode: D201875  
Submission No.: S463228

Test Material: AC 92,533

Synonyms: Pendimethalin, Prowl, N-(-1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzeneamine

Sponsor: American Cyanamid Co.  
Princeton, NJ 08543-0400

Testing Facility: Division of Endocrinology, University of  
Massachusetts Medical School, Worcester, MA  
01655

Title of Report: A 14-Day Intrathyroidal Metabolism Study in Male  
Rats with AC 92,553

Study Number: UM-91-06-01

Authors: William J. DeVito and Lewis E. Braverman

Report Issued: April 16, 1993

Executive Summary: Charles River rats were administered 0, 100 or 5000 ppm of pendimethalin in the diet for 14 days (approximate dosages of 0, 5 or 250 mg/kg/day).

At 5000 ppm AC 92,533 for 14 days, TSH was increased and  $T_4$  and  $T_3$  were decreased. No treatment related effects were observed for  $rT_3$  levels, thyroid weight,  $^{131}I$  uptake in MIT, DIT or  $T_4$ . There was a significant increase of  $^{131}I$  uptake by the thyroid of rats in the 5000 ppm group and an increase in incorporation of  $^{131}I$  in  $T_3$ . Total  $T_3$  and  $T_4$  levels in the thyroid were not affected by treatment at 5000 ppm. No treatment related effects were noted in the 100 ppm group.

The study is core-supplementary and does not by itself satisfy the guideline requirement for a series 85-1 metabolism study.

A. Materials

1. Test compound: AC 92,533; Description: orange solid; Lot No.: AC 6539-77A; Purity: 92.98%, Contaminants: not reported.
2. Test animals: Species: rat; Strain: Crl: CD(SD); Age: 13 weeks, Weight: not reported; Source: Charles River Laboratories, Inc.

B. Study Design:14-DAY INTRATHYROIDAL METABOLISM STUDY

Test Group	Dose in Diet (ppm)	Number of Rats
Control	0	10
Low	100	10
High	5000	10

Rats were maintained under "controlled" conditions with a 12 hour on/off light cycle. Rats were maintained on their respective diets for 14 days after which blood samples were obtained by retroorbital bleeding and the serum was frozen at -20 degrees C and later analyzed for serum TSH,  $T_3$ ,  $T_4$ , and  $rT_3$ . Each rat then received an ip injection of 25-50  $\mu\text{Ci}$   $^{131}\text{I}(\text{NaI})$ . Two hours after injection, the rats were sacrificed and blood was collected. One 50  $\mu\text{l}$  aliquot was counted. A 10-20  $\mu\text{l}$  aliquot was subjected to paper electrophoresis. The percentage of  $^{131}\text{I}$  present in the organic forms was determined by counting the appropriate zones by autoradiography. An aliquot of homogenate from each thyroid was hydrolyzed with 20 mg of pancreatin at 37 degrees C. Then the homogenate was subjected to ascending paper chromatography in butanol-ethanol-0.5 N ammonia (5:1:2) solvent system. Monoiodotyrosine (MIT), diiodotyrosine (DIT),  $T_3$ , and  $T_4$  were localized and counted. At termination, body weight and thyroid weight were determined.

C. Results

There was a significant increase in TSH (60%) in the 5000 ppm group when compared to the control and 100 ppm groups. In the 5000 ppm group there were significant decreases in  $T_4$  (80%) and  $T_3$  (39%). No treatment related effects were noted on  $rT_3$  or

thyroid weight. There was an increase in  $^{131}\text{I}$  uptake in rats in the 5000 ppm group (73%) when compared to controls (see Table 1). There were no significant increases of  $^{131}\text{I}$  uptake in MIT, DIT or  $\text{T}_4$ . There was a significant percentage of  $^{131}\text{I}$  incorporation into  $\text{T}_3$  in rats fed 5000 ppm AC 92,553 (see Table 2-taken from p.17 of the study report). Ingestion of 5000 ppm did not significantly affect the total concentration of  $\text{T}_3$  or  $\text{T}_4$  in the thyroid.

Table 1. Thyroid Function at 14 Days

Parameter	Dose Level (ppm)		
	0	100	5000
TSH ( $\mu\text{U}/\text{ml}$ )	46.04 $\pm 21.39$	41.07 $\pm 18.08$	73.82 $\pm 33.90$
$\text{T}_4$ ( $\mu\text{g}/\text{dl}$ )	5.23 $\pm 1.02$	4.12 $\pm 1.00$	1.02 $\pm 0.28$
$\text{T}_3$ (ng/ml)	96.42 $\pm 20.85$	76.66 $\pm 19.07$	58.40 $\pm 13.15$
$\text{rT}_3$ (pg/ml)	70.92 $\pm 17.83$	63.01 $\pm 16.72$	48.52 $\pm 9.76$
$^{131}\text{I}$ uptake % dose	2.75 $\pm 1.04$	2.85 $\pm 1.06$	4.75 $\pm 0.63$
Thyroid Wt. (mg)	27.85 $\pm 7.66$	26.67 $\pm 3.67$	26.27 $\pm 4.24$

#### D. Discussion

Oral exposure of rats to 5000 ppm AC 92,533 increased TSH, and decreased  $\text{T}_4$  and  $\text{T}_3$  levels. No treatment related effects were observed for  $\text{rT}_3$  levels, thyroid weight,  $^{131}\text{I}$  uptake in MIT, DIT or  $\text{T}_4$ . There was a significant increase in incorporation of  $^{131}\text{I}$  in  $\text{T}_3$ . Total  $\text{T}_3$  and  $\text{T}_4$  levels in the thyroid were not affected by treatment at 500 ppm. No treatment related effects were noted in the 100 ppm group.

The study is core-supplementary. It does not by itself satisfy the guideline requirement for a series 85-1 metabolism study.

U.S. ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES/HED/TOX  
ONELINERS

## Study Number UN-91-06-01

Table 2. Percentage  $^{131}\text{I}$  incorporated into organic  $^{131}\text{I}$ , and  $^{131}\text{I}$  labeled MIT, DIT,  $\text{T}_1$  and  $\text{T}_2$  in rats fed either control diet or diet containing 100 or 5000 ppm AC 92,553 for 14 days.

	Percentage of intrathyroidal $^{131}\text{I}$		
	Control	AC 92,553 100 ppm	AC 92,553 5000 ppm
Organic $^{131}\text{I}$	92.8 $\pm$ 0.9	92.8 $\pm$ 1.5	92.4 $\pm$ 1.1
MIT	33.9 $\pm$ 1.1	31.4 $\pm$ 1.1	29.8 $\pm$ 0.8
DIT	43.6 $\pm$ 1.4	44.3 $\pm$ 1.7	46.4 $\pm$ 1.5
$\text{T}_1$	0.37 $\pm$ 0.09	0.32 $\pm$ 0.07	1.1 $\pm$ 0.1*
$\text{T}_2$	1.5 $\pm$ 0.3	1.4 $\pm$ 0.2	1.2 $\pm$ 0.1
Mean $\pm$ SEM			
*p < 0.05 vs control and 100 ppm			

Taken from p.17 of the study report.

Chemical Name: Pendimethalin

TOXCHEM NO.: 454BB

P. C. No: 108501

CITATION	MATERIAL	NRID NUMBER	RESULTS	TOX CAT	CORE GRADE DOC. #
(82-SS) Special Study Species: rat Lab. Name: Univ of Mass. Study No: UM-91-06-01 Date: 4/16/93	Pendimethalin	31350-03	<p>Ten male Charles River rats were administered 0, 100 or 5000 ppm of pendimethalin in the diet for 14 days to evaluate intrathyroidal metabolism.</p> <p>(150 mg/kg/day)</p> <p>At 5000 ppm AC 92,533 for 14 days, TSH was increased and <math>T_4</math> and <math>T_3</math> were decreased. No treatment related effects were observed for <math>rT_4</math> levels, thyroid weight, <math>^{125}I</math> uptake in MIT, DIT or <math>T_4</math>. There was a significant increase of <math>^{125}I</math> uptake by the thyroid of rats in the 5000 ppm group and an increase in incorporation of <math>^{125}I</math> in <math>T_3</math>. Total <math>T_3</math> and <math>T_4</math> levels in the thyroid were not affected by treatment at 5000 ppm. No treatment related effects were noted in the 100 ppm group.</p> <p>(52/8/93)</p>		Supplementary



13544

R059062

<b>Chemical:</b>	<b>Pendimethalin</b>
<b>PC Code:</b>	<b>108501</b>
<b>HED File Code</b>	<b>13000 Tox Reviews</b>
<b>Memo Date:</b>	<b>05/19/95 12:00:00 AM</b>
<b>File ID:</b>	<b>DPD201875</b>
<b>Accession Number:</b>	<b>412-04-0140</b>

**HED Records Reference Center**  
**04/02/2004**