



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

2-25-82

FEB 25 1982

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

SUBJECT: 82-NJ-02,-03,-04 and -05. Proposed Section 18
exemptions for the use of Bayleton on cucurbits

FROM: Edward Zager, Chemist
Residue Chemistry Branch
Hazard Evaluation Division (TS-769)

Edward Zager

THRU: Charles L. Trichilo, Chief
Residue Chemistry Branch
Hazard Evaluation Division (TS-769)

CT

TO: Emergency Response Section
Registration Division (TS-767)

and

Toxicology Branch
Hazard Evaluation Division (TS-769)

The New Jersey Department of Environmental Protection requests
Section 18 exemptions for the use of Bayleton 50 WP (50% active
ingredient) on cucurbits (including cucumbers, melons, squash and
pumpkins).

PP#OE2393 and PP#OF2349 proposing tolerances for residues of
Bayleton (1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-
yl)-2-butanone and its metabolite KWG 0519 (β -(4-chlorophenoxy)
- α -(1,1-dimethylethyl)-1H-1,2,4-triazol-1-ethanol in or on
cucumbers at 0.1 ppm and melons at 0.2 ppm are currently in reject
status due to questions concerning analytical methodology.

The proposed use would permit up to six air or ground applications
at the rate of 0.5-1 oz act/A to cucurbits (including cucumbers,
melons squash and pumpkins) with a 7 day PHI.

The metabolism of Bayleton in cucumbers was discussed in our
review of PP#OE2393 (A. Smith, 12/2/80). For the purpose of this
emergency use, we consider the residue of concern in cucurbits to
be Bayleton and its metabolite KWG 0519. We note that the GC
method used to generate the residue data appears to determine only
the free residues of Bayleton and KWG 0519. However, conjugated
residues of Bayleton and KWG 0519 are not likely to exceed 20%
of the total residue in cucurbits.

Residue data reflecting applications to cucumbers grown in Mexico
were submitted with PP#OE2393. Following 3 foliar applications
at the rate of 1.8 oz act/A residues of Bayleton and KWG 0519
were <0.01-0.04 ppm at 5 days, <0.01-0.02 ppm at 15 days.

327

1/2

Residue data reflecting applications to cantaloupes also grown in Mexico were submitted with PP#0F2349. Following 3 foliar applications at the rate of 1.75 oz act/A residues of Bayleton and KWG 0519 were 0.03-0.13 ppm, at 0 days; 0.05-0.08 ppm at 5 days and 0.03-0.11 ppm at 15 days.

Based on the above data we estimate that residues of Bayleton and KWG 0519 will not exceed 0.2 ppm in or on cucurbits as a result of the proposed use.

Meat, Milk, Poultry and Eggs

There are no feed items involved in this use. Consequently, there will be no problem of secondary residues in meat, milk, poultry and eggs.

Conclusions

1. Residues of Bayleton and its metabolite KWG 0519 will not exceed 0.2 ppm in or on cucurbits as a result of the proposed use.
2. The proposed use will not lead to secondary residues in meat, milk, poultry and eggs.

Recommendation

TOX considerations permitting, we have no objections to the proposed Section 18 exemptions. An agreement should be made with FDA regarding the legal status of treated cucurbits in commerce.

cc: Section 18 S.F.
R.F.,
Circu
Reviewer
Bayleton S.F.

TS-769:RCB:Reviewer:EZager:LOT:X77324:CM#2:RM:810:Date:2/23/82
RDI:Section Head:RJH:Date:2/23/82:RDS:Date:2/23/82