- 1. Chemical: Baquacil
- 2. Formulation: Technical (20% a.i.)
- 3. Citation: Fink, R. (1979) Eight-Day Dietary LC50-Bobwhite Quail Final Report, Sample #5889, EPA Registration No. 10182-RO; Product Name, wildlife International LTD for ICI american, Inc. Asldsboro, N.C. 27530
- Reviewed By: C.E. Laird 4.
- 5. Date Reviewed: April 8, 1980
- 6. Test Type: Avian Eight-Day Dietary LC50
  - Test Species: Bobwhite Quail
- 7. Reported Results:

There were three mortalities in the negative control groups, each the result of toe picking, and three mortalities occurred at the 3160 ppm dose level on day 4 due to nostril and toe picking.

Reviewer's Conclusion:

The study is scientifically sound and indicates the Baquacil is practically non-toxic to bobwhile quail. The study does fulfill the requirements for an avian eight-day dietary LC50.

#### Material/Methods

#### Test Procedure

Not given

#### Statistical Analysis

Finney Probit

### Discussion / Results

The LC50 value was greater than 5620 ppm. There were mortalities in the negative control and three mortalities at the 3160 ppm dosage level.

### Reviewer's Evaluation

### A. <u>Test Procedure</u>

Test procedure complies with the recommended EPA protocol of 1978.

## B. Statistical Analysis

No statistical analysis was performed, because no mortalities occurred at any dosage level due to test chemical.

# C. Conclusion

- 1. Category: Core
- 2. Rationale: N/A
- 3. Repairability: N/A