

DATA EVALUATION RECORD
ACUTE LC₅₀ TEST WITH AN ESTUARINE/MARINE ORGANISM
§72-3(C) - SHRIMP

1. **CHEMICAL:** JAU6476-desthio PC Code No.: 113961

2. **TEST MATERIAL:** JAU 6476-Desthio Purity: 96.5%

Common name: JAU6476-desthio

Chemical:

IUPAC name: 2-(1-chlorocyclopropyl)-1-(2-chlorophenyl)-3-(1,2,4-triazol-1-yl)-propan-2-ol

CAS name: 2-(1-Chlorocyclopropyl)-1-(2-chlorophenyl)-3-(1, 2, 4-triazol-1-yl)-
propan-2-ol

CAS No.: 120983-64-4

Synonyms: SXX0665

3. **CITATION:**

Author: Drottar, K.R., *et al.*

Title: Desthio JAU 6476: A 96-Hour Flow Through Acute
Toxicity Test with the Saltwater Mysid (*Mysidopsis bahia*).

Study Completion Date: June 7, 2002

Laboratory: Wildlife International Ltd.
8598 Commerce Drive
Easton, MD 21601

Sponsor: Bayer Corporation
Agriculture Division
17745 South Metcalf
Stilwell, Kansas 66085-9104

Laboratory Report ID: 149A-129

MRID No.: 46246017

DP Barcode: D303488



PMRA Submission Number 2004-0843

4. REVIEWED BY: Rebecca Bryan, Staff Scientist, Dynamac Corporation

Signature:

Date: 8/30/2004

APPROVED BY: Gregory Hess, Staff Scientist, Dynamac Corporation

Signature:

Date: 9/8/2004


5. APPROVED BY: Kevin Costello, ERB-IV

Signature:

Date:

6. SECONDARY REVIEW BY:


Émilie Larivière, EAD, PMRA, HC

Signature: 

9/8/05

Date: 9/8/2005

Christopher J. Salice, ERB-IV, EPA, EFED

Signature: 

Date: 8-15-05

7. STUDY PARAMETERS:

Scientific Name of Test Organism: *Mysidopsis bahia*

Age or Size of Test Organism: <24 hours old

Definitive Test Duration: 96 hours

Study Method: Flow-through

Type of Concentration: Mean-measured

8. CONCLUSIONS:

The 96-hour acute toxicity of JAU 6476-Desthio to the saltwater mysid, *Mysidopsis bahia*, was studied under flow-through conditions. Mysids were exposed to the test material at nominal concentrations of 0 (negative control), 0 (0.1 mL/L DMF control), 0.013, 0.025, 0.050, 0.10, and 0.20 ppm a.i. Mean-measured concentrations were <0.0025 (LOQ; controls), 0.013, 0.026, 0.050, 0.099, and 0.20 ppm a.i.

After 96 hours, mortality was 10, 5, 15, 95, and 100% in the 0.013, 0.026, 0.050, 0.099, and 0.20 ppm a.i. treatment levels, respectively. No mortalities were observed in the controls. Erratic swimming and/or lethargy were observed in surviving mysids from the

DATA EVALUATION RECORD
ACUTE LC₅₀ TEST WITH AN ESTUARINE/MARINE ORGANISM
§72-3(C) - SHRIMP

1. **CHEMICAL:** JAU6476-desthio PC Code No.: 113961

2. **TEST MATERIAL:** JAU 6476-Desthio Purity: 96.5%

Common name: JAU6476-desthio

Chemical:

IUPAC name: 2-(1-chlorocyclopropyl)-1-(2-chlorophenyl)-3-(1,2,4-triazol-1-yl)-propan-2-ol

CAS name: 2-(1-Chlorocyclopropyl)-1-(2-chlorophenyl)-3-(1, 2, 4-triazol-1-yl)-
propan-2-ol

CAS No.: 120983-64-4

Synonyms: SXX0665

3. **CITATION:**

Author: Drottar, K.R., *et al.*

Title: Desthio JAU 6476: A 96-Hour Flow Through Acute
Toxicity Test with the Saltwater Mysid (*Mysidopsis bahia*).

Study Completion Date: June 7, 2002

Laboratory: Wildlife International Ltd.
8598 Commerce Drive
Easton, MD 21601

Sponsor: Bayer Corporation
Agriculture Division
17745 South Metcalf
Stilwell, Kansas 66085-9104

Laboratory Report ID: 149A-129

MRID No.: 46246017

DP Barcode: D303488

4. **REVIEWED BY:** Rebecca Bryan, Staff Scientist, Dynamac Corporation

Signature:

Date: 8/30/2004

APPROVED BY: Gregory Hess, Staff Scientist, Dynamac Corporation

PMRA Submission Number 2004-0843

Signature:

Date: 9/8/2004

5. **APPROVED BY:** Kevin Costello, ERB-IV

Signature:

Date:

6. **SECONDARY REVIEW BY:**

Émilie Larivière, EAD, PMRA, HC

Signature:

Date: 9/8/2005

Christopher J. Salice, ERB-IV, EPA, EFED

Signature:

Date:

7. **STUDY PARAMETERS:**

Scientific Name of Test Organism: *Mysidopsis bahia*

Age or Size of Test Organism: <24 hours old

Definitive Test Duration: 96 hours

Study Method: Flow-through

Type of Concentration: Mean-measured

8. **CONCLUSIONS:**

The 96-hour acute toxicity of JAU 6476-Desthio to the saltwater mysid, *Mysidopsis bahia*, was studied under flow-through conditions. Mysids were exposed to the test material at nominal concentrations of 0 (negative control), 0 (0.1 mL/L DMF control), 0.013, 0.025, 0.050, 0.10, and 0.20 ppm a.i. Mean-measured concentrations were <0.0025 (LOQ; controls), 0.013, 0.026, 0.050, 0.099, and 0.20 ppm a.i.

After 96 hours, mortality was 10, 5, 15, 95, and 100% in the 0.013, 0.026, 0.050, 0.099, and 0.20 ppm a.i. treatment levels, respectively. No mortalities were observed in the controls. Erratic swimming and/or lethargy were observed in surviving mysids from the solvent control (1 mysid; only at 72-hours), and the 0.013 (1 mysid; 24-48 hours), 0.050 (1-3 mysids; 24-72 hours), 0.099, and 0.20 ppm a.i. treatment levels during the test. No sub-lethal effects were observed in the negative control and 0.026 ppm a.i. treatment level. The **96-hour LC₅₀ value (with 95% C.I.) was 0.060 (0.046-0.079) ppm a.i.**, which categorizes JAU 6476-Desthio (Prothioconazole) as **very highly toxic** to the saltwater mysid, *Mysidopsis bahia*, on an acute toxicity basis. Based on mortality and sub-lethal effects, the

PMRA Submission Number 2004-0843

NOAEC and LOAEC values were 0.026 and 0.050 ppm a.i., respectively.

This study is scientifically valid and fulfills the requirements of an acute LC₅₀ test with an estuarine/marine organism (Subdivision E, §72-3(c) [mysid shrimp]). This study is classified as **ACCEPTABLE**.

Results Synopsis

96-Hour:

LC₅₀: 0.060 ppm a.i.

95% C.I.: 0.046-0.079 ppm a.i.

NOAEC: 0.026 ppm a.i.

LOAEC: 0.050 ppm a.i.

Endpoints affected: Mortality and sub-lethal effects (same conclusions)

9. ADEQUACY OF THE STUDY:

A. Classification: ACCEPTABLE

B. Rationale: This study is scientifically sound and satisfies the guideline requirements for an acute toxicity study with mysid (§72-3(c)). Missing information should be provided to the U.S. EPA.

C. Repairability: N/A

10. BACKGROUND:

11. GUIDELINE DEVIATION:

1. The pretest mortality of the mysid culture was not reported.
2. The total organic carbon concentration found in the dilution water was not reported.
3. The water temperature of 24.4-25.5°C was higher than recommended (approximately 22°C).

12. SUBMISSION PURPOSE: This study was submitted to provide data on the toxicity of JAU 6476-Desthio to mysids for the purpose of chemical registration.

PMRA Submission Number 2004-0843

13. MATERIALS AND METHODS:

A. Test Organisms

Guideline Criteria	Reported Information
<p><u>Species</u> Preferred species are <i>Americamysis bahia</i>, <i>Penaeus setiferus</i>, <i>P. duorarun</i>, <i>P. aztecus</i> and <i>Palaemonetes sp.</i></p>	<p><i>Mysidopsis bahia</i> (same as <i>Americamysis bahia</i>)</p>
<p><u>Age</u> Juvenile (\leq 24 hours old) mysids should be used</p>	<p><24 hours old</p>
<p><u>Supplier</u></p>	<p>Juveniles were collected from in-house laboratory cultures.</p>
<p>All shrimp are from same source?</p>	<p>Yes</p>
<p>All shrimp are from the same year class?</p>	<p>Yes</p>

B. Source/Acclimation

Guideline Criteria	Reported Information
<p><u>Acclimation Period</u> Minimum 10 days</p>	<p>Continuous; held for 14 days prior to test initiation.</p>
<p>Wild caught organisms were quarantined for 7 days?</p>	<p>N/A</p>
<p>Were there signs of disease or injury?</p>	<p>No signs of disease or stress at beginning of test.</p>
<p>If treated for disease, was there no sign of the disease remaining during the 48 hours prior to testing?</p>	<p>N/A</p>

PMRA Submission Number 2004-0843

Guideline Criteria	Reported Information
<p><u>Feeding</u> No feeding during the study and no feeding for 24 hours before the beginning of the test if organisms are over 0.5 g each. Mysids should be fed throughout the study.</p>	Fed live brine shrimp (<i>Artemia</i> sp.) nauplii daily during testing.
<p><u>Pretest Mortality</u> <3% mortality 48 hours prior to testing</p>	Not reported

C. Test System

Guideline Criteria	Reported Information
<p><u>Source of dilution water</u> Soft reconstituted water or water from a natural source, not dechlorinated tap water</p>	Natural seawater collected at Indian River Inlet, Delaware was filtered and diluted (to a salinity of approximately 20‰) with well water. Diluted seawater was then aerated, filtered (0.45 µm) to remove microorganisms and fine particles and UV sterilized.
<p>Does water support test animals without observable signs of stress?</p>	Yes
<p><u>Salinity</u> 30-34 ‰ (parts per thousand) for marine (stenohaline) shrimp and 10-17 ‰ for estuarine (euryhaline) shrimp, weekly range <6 ‰</p>	21‰
<p><u>Water Temperature</u> Approx. 22 ± 1 °C</p>	24.4-25.5°C

PMRA Submission Number 2004-0843

Guideline Criteria	Reported Information
<p><u>pH</u> 8.0-8.3 for marine (stenohaline) shrimp, 7.7-8.0 for estuarine (euryhaline) shrimp, monthly range < 0.8</p>	8.2-8.3
<p><u>Dissolved Oxygen</u> Between 60 and 105% saturation. If needed, aerate prior to introduction of chemical.</p>	6.0-6.8 mg/L ($\geq 82\%$ saturation)
<p><u>Total Organic Carbon</u> Should be <5 mg/L in reconstituted seawater</p>	Not reported
<p><u>Test Aquaria</u> 1. <u>Material</u>: Glass or stainless steel 2. <u>Size</u>: 19.6 L is acceptable for organisms \geq 0.5 g (e.g. pink shrimp, white shrimp, and brown shrimp), 3.9 L is acceptable for smaller organisms (e.g. mysids and grass shrimp). 3. <u>Fill volume</u>: 15 L is acceptable for organisms \geq 0.5 g, 2-3 L is acceptable for smaller organisms.</p>	The 2-L glass beakers (12 cm diameter and 19 cm height) had two nylon mesh- covered holes on each side and were suspended in 9-L glass aquaria. The aquaria were filled with approximately 5- L of test water and the test compartment (beaker) depth was 7.5 cm.
<p><u>Type of Dilution System</u> Must provide reproducible supply of toxicant</p>	Continuous-flow diluter
<p><u>Flow Rate</u> Consistent flow rate of 5-10 vol/24 hours, meter systems calibrated before study and checked twice daily during test period</p>	18 volume additions/24 hours System was calibrated prior to test initiation and checked twice daily during the test.

PMRA Submission Number 2004-0843

Guideline Criteria	Reported Information
<u>Biomass Loading Rate</u> Static: ≤ 0.8 g/L at $\leq 17^\circ\text{C}$, ≤ 0.5 g/L at $> 17^\circ\text{C}$; flow through: ≤ 1 g/L/day (N/A for mysids)	N/A
<u>Photoperiod</u> 16 hours light, 8 hours dark	16 hours light, 8 hours dark, with a 30-minute transition period.
<u>Solvents</u> Not to exceed 0.5 mL/L for static tests or 0.1 mL/L for flow-through tests	Dimethylformamide (DMF), 0.1 mL/L

D. Test Design

Guideline Criteria	Reported Information
<u>Range Finding Test</u> If $\text{LC}_{50} > 100$ mg/L with 30 shrimp, then no definitive test is required.	A range-finding study was conducted, but the results were not reported.
<u>Nominal Concentrations of Definitive Test</u> Control & 5 treatment levels; a geometric series in which each concentration is at least 60% of the next higher one.	0 (negative and solvent controls), 0.013, 0.025, 0.050, 0.10, and 0.20 ppm a.i.
<u>Number of Test Organisms</u> Minimum 20/level, may be divided among containers	20 mysids/level, divided into two replicates of 10 mysids each.
Test organisms randomly or impartially assigned to test vessels?	Yes
Biological observations made every 24 hours?	Yes

PMRA Submission Number 2004-0843

Guideline Criteria	Reported Information
<p><u>Water Parameter Measurements</u></p> <p>1. <u>Temperature</u> Measured constantly or, if water baths are used, every 6 hrs, may not vary >1°C</p> <p>2. <u>DO and pH</u> Measured at beginning of test and ever 48 h in the high, medium, and low doses and in the control</p>	<p>1. Measured in each aquarium at beginning and end of the test and continuously in one negative control vessel.</p> <p>2. Measured daily in alternating replicate aquariums.</p>
<p><u>Chemical Analysis</u> needed if solutions were aerated, if chemical was volatile, insoluble, or known to absorb, if precipitate formed, if containers were not steel or glass, or if flow-through system was used</p>	<p>Analytical determination of test substance was performed on samples collected from alternating replicate test vessels at the beginning of the test, after 48 hours, and at the end of the test.</p>

14. REPORTED RESULTS:

A. General Results

Guideline Criteria	Reported Information
<p>Quality assurance and GLP compliance statements were included in the report?</p>	<p>Yes</p>
<p><u>Recovery of Chemical</u></p>	<p>101-138%, based on matrix blanks fortified at 0.00500, 0.0500, and 0.250 ppm a.i. and extracted and analyzed concurrently with the dilution water (Appendix 3.5, p. 31).</p>
<p><u>Control Mortality</u> Not more than 10% of control organisms may die or show abnormal behavior.</p>	<p>0% negative and solvent control mortality were observed.</p>

PMRA Submission Number 2004-0843

Guideline Criteria	Reported Information
Raw data included?	Yes
Signs of toxicity (if any) were described?	Yes (lethargic and erratic swimming).

Mortality

Concentration (ppm a.i.)		Number of Shrimp	Mean cumulative mortality (%)			
Nominal	Mean Measured		Hours of Study			
			24	48	72	96
Negative Control	<LOQ	20	0	0	0	0
Solvent Control	<LOQ	20	0	0	0	0
0.013	0.013	20	0	5	5	10
0.025	0.026	20	0	5	5	5
0.050	0.050	20	0	0	5	15
0.10	0.099	20	75	95	95	95
0.20	0.20	20	95	100	100	100

LOQ = 0.00250 ppm a.i.

After 96 hours, mortality was 10, 5, 15, 95, and 100% in the mean-measured 0.013, 0.026, 0.050, 0.099, and 0.20 ppm a.i. treatment groups, respectively (Table 3, p. 21). No mortalities were observed in the controls.

Erratic swimming and/or lethargy were observed in surviving mysids from the solvent control (1 mysid; only at 72-hours), and the 0.013 (1 mysid; 24-48 hours), 0.050 (1-3 mysids; 24-72 hours), 0.099, and 0.20 ppm a.i. treatment levels during the test. No sub-lethal effects were observed in the negative control and 0.026 ppm a.i. treatment level.

B. Statistical Results

The 96-hour LC_{50} value (with 95% C.I.) was calculated using the moving average method via the computer program of C.E. Stephan. The 96-hour NOAEC was estimated

PMRA Submission Number 2004-0843

empirically based on mortality and clinical observation data. All toxicity values were determined in terms of the mean measured treatment concentrations.

96-Hour:LC₅₀: 0.060 ppm a.i.

95% C.I.: 0.046-0.079 ppm a.i.

NOAEC: 0.026 ppm a.i.

LOAEC: 0.050 ppm a.i.

Endpoints affected: Mortality and sub-lethal effects

15. VERIFICATION OF STATISTICAL RESULTS:

The 96-hour LC₅₀ was determined using the probit method, which did not provide a reliable 95% confidence interval; this analysis was conducted using TOXANAL statistical software. Consequently, the study author determined 96-hour LC₅₀ value and associated 95% confidence interval are reported in the CONCLUSION section of this DER. NOAEC and LOAEC values were determined visually based on mortality (15% in the 0.050 ppm a.i. treatment) and the reported sub-lethal effects data; >10% effects were considered significant. All toxicity values were determined using the mean-measured treatment concentrations.

96-Hour:LC₅₀: 0.057 ppm a.i.

95% C.I.: 0 to infinity ppm a.i.

Mortality:

NOAEC: 0.026 ppm a.i.

LOAEC: 0.050 ppm a.i.

Sub-lethal:

NOAEC: 0.026 ppm a.i.

LOAEC: 0.050 ppm a.i.

Endpoints affected: Sub-lethal effects (most sensitive)

16. REVIEWER'S COMMENTS:

The reviewer's conclusions were nearly identical to the study authors'. The reviewer's LC₅₀ estimate was associated with an unreliable 95% confidence interval (0 to infinity) because the probit method was insufficient to estimate the 96-hour LC₅₀, given the non-linear distribution of the data. Consequently, the study author determined 96-hour LC₅₀ value and associated 95% confidence interval are reported in the CONCLUSION section of this DER. The NOAEC and LOAEC values determined by the reviewer and study authors

PMRA Submission Number 2004-0843

were identical. Based on the LC₅₀, JAU 6476-Desthio is categorized as very highly toxic to saltwater mysids on an acute toxicity basis.

This study was conducted in accordance with USEPA (40 CFR Part 160), OECD, and JMAFF Good Laboratory Practice Regulations. A Quality Assurance and No Data Confidentiality Statements were included. This study is scientifically valid and fulfills the requirements of an acute LC₅₀ test with an estuarine/marine organism (Subdivision E, §72-3(c) [mysid shrimp]). This study is classified as **ACCEPTABLE**.

17. REFERENCES

- U.S. Environmental Protection Agency. 1996. Series 850 - Ecological Effects Test Guidelines (*draft*), OPPTS Number 850.1035: *Mysid Acute Toxicity Test*.
- U.S. Environmental Protection Agency. 1985. *Standard Evaluation Procedure, Acute Toxicity Test for Estuarine and Marine Organisms (Shrimp 96-Hour Acute Toxicity Test)*. Hazard Evaluation Division. Office of Pesticide Programs. EPA-540/9-85-010. Washington, DC.
- ASTM Standard E 729-88a. 1994. *Standard Guide for Conducting Acute Toxicity Tests with Fishes, Macroinvertebrates, and Amphibians*. American Society for Testing and Materials.
- Stephan, C.E. 1978. U.S. EPA, Environmental Research Laboratory, Duluth, Minnesota. Personal communication.
- Finney, D.J. 1971. *Statistical Methods in Biological Assay*. Second edition. Griffin Press, London.
- Thompson, W.R. 1947. *Bacteriological Reviews*. Vol. II, No. 2. Pp. 115-145.
- Stephan, C.E. 1977. "Methods for Calculating and LC₅₀", *Aquatic Toxicology and Hazard Evaluation*. American Society for Testing and Materials. Publication Number STP 634, pp 65-84.

PMRA Submission Number 2004-0843

APPENDIX 1. OUTPUT OF REVIEWER'S STATISTICAL VERIFICATION:

TOXANAL Results:

RESULTS CALCULATED USING THE PROBIT METHOD

ITERATIONS	G	H	GOODNESS OF FIT PROBABILITY
5	2.347277	8.593819	0

A PROBABILITY OF 0 MEANS THAT IT IS LESS THAN 0.001

SINCE THE PROBABILITY IS LESS THAN 0.05, RESULTS CALCULATED USING THE PROBIT METHOD PROBABLY SHOULD NOT BE USED.

SLOPE = 3.620131
95 PERCENT CONFIDENCE LIMITS = -1.926209 AND 9.166469

LC50 = 5.682581E-02
95 PERCENT CONFIDENCE LIMITS = 0 AND +INFINITY

LC10 = .0253355
95 PERCENT CONFIDENCE LIMITS = 0 AND .0858452

DP Barcode: D303488, D303495

MRID No.: 46246017

PMRA Submission Number 2004-0843

Data Evaluation Report on the Acute Toxicity of the transformation product JAU6476-desthio to Marine Invertebrates- Acute (Crustacean)

PMRA Submission Number 2004-0843

EPA MRID Number 46246017

EAD Assessment of USEPA DER

Reviewer: Émilie Larivière (#1269); PMRA

Date: September 8, 2005

PMRA Submission Number: 2004-0843

Study Type: Non-Target Marine Invertebrates - Acute (Crustacean)

Drottar, K.R., S.J. Palmer, T.Z. Kendall and H.O. Krueger. 2002. Desthio-JAU6476: A 96-hour flow-through acute toxicity test with the saltwater mysid (*Mysidopsis bahia*). Unpublished study performed by Wildlife International, Ltd., Easton, MD. Laboratory ID 149A-129. Study submitted by Bayer Corporation, Stilwell, KS. Bayer Report 110979. June 7, 2002.

PMRA DATA CODE: 9.4.2

EPA DP Barcode: D303488

OECD Data Point: IIA 8.11.1

EPA MRID: 46246017

EPA Guideline: §72-3(B); 850.1035

Company Code: BCZ

Active Code: PRB

Use Site Category: 7, 13, 14

EPA PC Code: 113961

EAD Executive Summary:

The 96-hour acute toxicity of the transformation product JAU 6476-desthio (purity: 96.5%) to the saltwater mysid, *Mysidopsis bahia*, was studied under flow-through conditions. Mysids were exposed to the test material at nominal concentrations of 0 (negative control), 0 (0.1 mL/L dimethylformamide control), 0.013, 0.025, 0.050, 0.10, and 0.20 mg JAU6476-desthio/L. Mean measured concentrations were <0.0025 (LOQ; controls), 0.013, 0.026, 0.050, 0.099, and 0.20 mg JAU6476-desthio/L. This study was conducted following U.S. EPA OPPTS Guideline 850.1035 and ASTM Standard E729-88a, and was in compliance with U.S. EPA (40 CFR Part 160), OECD, and Japan MAFF Good Laboratory Practice Regulations.

PMRA Submission Number 2004-0843

After 96 hours, mortality was 10, 5, 15, 95, and 100% in the 0.013, 0.026, 0.050, 0.099, and 0.20 mg JAU6476-desthio/L treatment levels, respectively. No mortalities were observed in the controls. Erratic swimming and/or lethargy were observed in surviving mysids from the solvent control (1 mysid; only at 72-hours), and the 0.013 (1 mysid; 24-48 hours), 0.050 (1-3 mysids; 24-72 hours), 0.099, and 0.20 mg JAU6476-desthio/L treatment levels during the test. No sub-lethal effects were observed in the negative control and 0.026 mg JAU6476-desthio/L treatment level. The 96-hour LC₅₀ value (with 95% C.I.) was 0.060 (0.046-0.079) mg JAU6476-desthio/L, which categorizes JAU 6476-desthio as very highly toxic to the saltwater mysid, *Mysidopsis bahia*, on an acute toxicity basis, according to the classification scheme of the U.S. EPA (1985). Based on mortality and sub-lethal effects, the NOEC and LOEC values were 0.026 and 0.050 mg JAU6476-desthio/L, respectively.

Results Synopsis**96-Hour:**LC₅₀: 0.060 mg JAU6476-desthio/L 95% C.I.: 0.046-0.079 mg JAU6476-desthio/L

NOEC: 0.026 mg JAU6476-desthio/L

LOEC: 0.050 mg JAU6476-desthio/L

Endpoints affected: Mortality and sub-lethal effects (same conclusions)

Evaluator Comments:

1. The appropriate PMRA information (PMRA Submission Number, PMRA Data Code, PMRA company code, PMRA active ingredient code, PMRA use site category, OECD data point) was added to the PMRA review portion of the DER. The PMRA Submission Number was added to the Header of the DER. Information on the chemical name (IUPAC name, CAS name and synonym) available from the study report, the PMRA Chemistry review and other sources of information was added at the beginning of the DER. The name of the EAD secondary reviewer was added to the front portion of the DER.
2. The name Prothioconazole should be removed from the Conclusions section (section 8), the section on Submission Purpose, Reviewer's Comments of the DER because the study was conducted with the transformation product JAU6476-desthio (SXX 0665) and not the parent compound prothioconazole. Changes are suggested in strikeout font and yellow highlighting.
3. The EAD reviewer verified the LC₅₀ value using probit analysis and obtained identical results to those of the EPA reviewer. The EAD reviewer agrees that the LC₅₀ value and 95% confidence intervals reported by the study authors (0.060 mg a.i./L; 95% C.I.: 0.046-0.079 mg a.i./L) are acceptable and should be the ones reported in the Executive Summary and the Conclusions.

PMRA Submission Number 2004-0843

4. No statistical analysis was done to determine the NOEC. Upon inspection of the data, the 0.05 and 0.099 mg a.i./L treatments showed 15 and 95% mortality, which the EAD reviewer considers to be greater than natural mortality. The EAD reviewer therefore agrees with the study author that the NOEC for mortality is 0.026 mg a.i./L. The EAD also agrees that the NOEC for sub-lethal effects is 0.026 mg a.i./L, based on visual observation of the data.

Study Acceptability: This study is scientifically valid and fulfills the requirements of an acute toxicity test of the transformation product JAU6476-desthio with an estuarine/marine invertebrate (acute - crustacean). This study is classified as **ACCEPTABLE**.